Dynamics of the cell-free DNA methylome of metastatic prostate cancer during androgen-

targeting treatment

Running Head: cfDNA methylome in advanced prostate cancer

Authors: Madonna R. Peter[1,2], Misha Bilenky[3], Ruth Isserlin[4], Gary D. Bader[4], Shu Yi Shen[5],

Daniel D. De Carvalho[5,6], Aaron R. Hansen[7], Pingzhao Hu[8], Neil E. Fleshner[9], Anthony M.

Joshua[7,10], Martin Hirst[3,11], Bharati Bapat[1,2]*

Affiliations:

1. Lunenfeld-Tanenbaum Research Institute, Sinai Health System, Toronto, Canada.

2. Department of Laboratory Medicine & Pathobiology, University of Toronto, Toronto, Canada.

3. Canada's Michael Smith Genome Science Center, BC Cancer Agency, Vancouver, Canada.

4. Terrence Donnelly Centre for Cellular and Biomolecular Research, University of Toronto,

Toronto, Canada.

5. Princess Margaret Cancer Centre, University Health Network, Toronto, Canada

6. Department of Medical Biophysics, University of Toronto. Toronto, Ontario, Canada

7. Division of Medical Oncology and Hematology, Princess Margaret Cancer Centre, Toronto,

Canada.

8. Department of Biochemistry and Medical Genetics, University of Manitoba, Winnipeg,

Canada.

9. Departments of Surgery and Surgical Oncology, Division of Urology, University Health

Network, Toronto, Canada.

10. Department of Medical Oncology, Kinghorn Cancer Centre, Australia.

11. Department of Microbiology and Immunology and Michael Smith Laboratories, University

of British Columbia, Vancouver, Canada.

*Corresponding author: Dr. Bharati Bapat. 60 Murray Street, Toronto, ON, Canada, M1B 1Y5.

Telephone: 416-586-4800, Ext: 5175. Email: bapat@lunenfeld.ca

Abstract

Aim: We examined methylation changes in cell-free DNA (cfDNA) in metastatic castration

resistant prostate cancer (mCRPC) during treatment. **Materials and Methods:** Genome-wide

methylation analysis of sequentially collected cfDNA samples derived from mCRPC patients

undergoing androgen-targeting therapy was performed. **Results:** Alterations in methylation

states previously implicated in prostate cancer progression were observed and patients that

maintained methylation changes throughout therapy tended to have a longer time to clinical

progression (TTP). Importantly, we also report that markers associated with a highly aggressive

form of the disease, Neuroendocrine-CRPC, were associated with a faster TTP. **Conclusion:** Our

findings highlight the potential of monitoring cfDNA methylome during therapy in mCRPC,

which may serve as predictive markers of response to androgen-targeting agents.

Introduction:

Prostate cancer (PCa) is the most common cancer and the second leading cause of cancer-related deaths among males (1, 2). Androgen deprivation therapy (ADT) to reduce systemic androgen levels continues to be the major treatment for aggressive PCa (3). While ADT is initially beneficial, treatment resistance leads to the most lethal form of PCa, metastatic castration resistant prostate cancer (mCRPC) (4). Despite castrate levels of systemic androgens, these tumors can continue to rely on the androgen receptor (AR) pathway to promote tumor growth and metastasis (5). Indeed, androgen-targeting agents, Enzalutamide and Abiraterone acetate, can improve overall and progression-free survival in both the pre- and post-chemotherapy settings (6-9).

Enzalutamide is an antiandrogen that directly inhibits full-length AR. Abiraterone inhibits CYP17A1, which catalyzes extragonadal androgen production (10, 11). Both treatments perform similarly to suppress the androgen pathway, and choice of treatment is often dependent on comorbidities (12). However, primary and eventual secondary resistance to Abiraterone or Enzalutamide remains an ongoing challenge. As a result, the treatment landscape of mCRPC is complex and continues to focus primarily on the order/sequencing of therapies. For instance, recent trials demonstrated early treatment with Abiraterone or Enzalutamide in the hormone-sensitive state could be beneficial (13-15). However, there is a diverse array of molecular drivers contributing to disease heterogeneity amongst mCRPC patients (16, 17), leading to variable response to current treatment strategies. Therefore, reliable biomarkers are needed to facilitate optimal therapy sequences for each patient prior to starting treatments.

Liquid biopsies are emerging as a minimally invasive source of biomarkers, reflecting

tumor turnover (18). Numerous studies have explored the potential of circulating tumor cell

(CTC) or circulating nucleic acid markers in mCRPC (19). Counting the number of CTCs and/or

characterization of molecular alterations (i.e. genomic and transcriptomic) are promising markers

that capture tumor heterogeneity (20). For instance, increased CTC number is associated with

poor prognosis, and expression of the ligand-independent AR-V7 splice variant by CTCs is

associated with resistance to Enzalutamide and Abiraterone (21-23). However, low CTC counts

prior to first-line treatment and the need for robust CTC surface markers for their optimal

detection remain an ongoing challenge (24, 25). Furthermore, some AR-V7 positive patients may

still benefit from androgen therapies, while certain AR-V7 negative patients show variability in

treatment response (26, 27). In addition, highly aggressive androgen-independent

Neuroendocrine-CRPC (NE-CRPC) tumors express lower levels of AR-V7 than

Adenocarcinoma-CRPC patients (28). Therefore, additional markers are needed to identify this

subset of patients.

Circulating cell free DNA (cfDNA) can harbor tumor-specific somatic mutations and

capture tumor heterogeneity (29). There is a high concordance (>90%) of tumor mutations (i.e.

AR, BRCA2) between matched cfDNA and mCRPC biopsies (18). Although the proportion of

circulating tumor DNA (ctDNA) can vary from 1-2% to ~30% of cfDNA in mCRPC, the

sequencing throughput of massively parallel sequencing platforms can sample cfDNA with

sufficient coverage (30). For instance, targeted sequencing of *AR* in cfDNA can detect mutations

associated with resistance to Enzalutamide or Abiraterone (31, 32). However, genomic

aberrations are not the only contributors to molecular and phenotypic heterogeneity in mCRPC.

Aberrations in the epigenome are a hallmark of all stages of PCa, including DNA methylation

alterations (33, 34). Hypermethylation in promoter regions of several genes, such as *GSTP1*,

APC, *HOXD3* and *TBX15*, are currently being investigated as diagnostic and prognostic markers

in earlier stages of PCa (34, 35). Furthermore, in biopsy tissue from metastatic lesions, distinct

DNA methylation patterns were observed between NE-CRPC and Adenocarcinoma-CRPC (28).

CpG methylation can be detected from DNA extracted from tissue and various biofluids

(36-38). Presence of promoter methylation in *GSTP1* and *APC* in cfDNA from mCRPC patients

is prognostic of overall survival (36, 39). Using an array-based platform, differential cfDNA

methylation patterns were observed between Abiraterone responsive versus resistant patients

(37). Recently, NE-CRPC tissue-derived methylation signals were shown to be detectable in

matched plasma samples (40). While these observations are promising, extensive genome wide

analysis of the mCRPC cfDNA methylome during Enzalutamide or Abiraterone treatment has

not been performed. To identify circulating DNA methylation changes associated with response

to either Enzalutamide or Abiraterone treatment, we monitored sequentially collected cfDNA

samples from mCRPC patients, starting from prior to treatment initiation to eventual clinical

progression, and performed genome-wide methylome assessment.

Given that biopsy tissue samples are not easily obtainable in the clinical setting,

especially bone metastases, we addressed whether analysis of intra-patient cfDNA methylation

changes without a priori knowledge of matched tumour methylation patterns, and/or integration

with tissue methylation datasets is required to correlate with response to treatment. We identified

key genes known to have altered methylation states in PCa and found genomic regions

associated with clinical progression. Furthermore, as these changes could be from tumour and

non-tumour derived cfDNA, we combined our findings with published data and revealed cfDNA

methylation patterns associated with worse clinical prognosis during androgen-targeting

treatment.

Patients and Methods

Patient cohort

All patients in this study were recruited from the Princess Margaret Cancer Centre

Genitourinary Clinic, and informed written consent was obtained in accordance with approved

institutional Research Ethics Board protocols from University Health Network (UHN) and Sinai

Health System (SHS). The UHN Genitourinary Biobank performed patient recruitment, blood

collection and clinical follow-up. All patients were chemotherapy naïve and developed resistance

following initial PCa treatments (local therapies and ADT) (**Supplementary Table 1**). A total of

12 Enzalutamide-treated (160 mg daily) and 4 Abiraterone-treated patients (1000 mg daily +

Prednisone/Dexamethasone) completed study visits for cfDNA methylome analysis. Blood was

collected at three time points: prior to starting treatment (**Visit A**), at 12-weeks during treatment

(**Visit B**) and upon clinical progression/treatment change (**Visit C**). While we aimed to collect

Visit B around week-12 (+/- 2 weeks), average time from baseline for Visit B was 12.7 weeks

(ranging 9 - 17 weeks). Serum PSA levels were obtained throughout treatment, and the lowest

PSA level (nadir) was used to determine PSA progression. Clinical progression was determined

by several factors, including radiological evidence of additional/worsening metastases and

symptomatic changes as assessed by the treating physician.

Blood sample processing and cfDNA isolation

Blood was collected at each time point in citrate cell preparation tubes (BD Biosciences,

San Jose, CA, USA) and samples were processed within 2 hours from collection. Plasma was

separated from peripheral blood mononuclear cells (PBMCs) by centrifugation at 1800 x g for 20

minutes. Isolated plasma was divided into 1 mL aliquots and stored at -80°C. Prior to cfDNA

isolation, frozen plasma samples were rapidly thawed at 37°C and spun at 16 000 x g for 5

minutes. For each visit, we utilized 4 mL of plasma and cfDNA was isolated using the QIAamp

Circulating Nucleic Acid Kit (Qiagen, Hilden, Germany). The concentration of cfDNA was

determined using the Qubit dsDNA High Sensitivity Assay Kit (ThermoFisher Scientific,

Waltham, MA, USA) and cfDNA fragment distibrution for subset of samples was obtained using

the Agilent 2100 Bioanalyzer System. To avoid within-patient batch effects, we isolated plasma

cfDNA and performed methylation analysis for all visits from each patient together.

cfMeDIP-seq protocol

Genome-wide methylation profiling was performed using a published protocol, cell-free

methylated DNA immunoprecipitation and high-throughput sequencing (cfMeDIP-seq) (41).

Briefly, isolated cfDNA was first end-repaired and A-tailed using the KAPA HyperPrep kit

(Roche, Mannheim, Germany), followed by adapter ligation with index adapters (Integrated

DNA technologies). Each adapter-ligated cfDNA sample was then spiked with λ phage DNA

(ThermoFisher Scientific) as filler DNA, as well as methylated and unmethylated control

Arabidopsis thaliana (AT) DNA (Diagenode, Denville, NJ, USA) to bring the total DNA amount

to 100ng. Prior to immunoprecipitation, 10% of this DNA is saved as input control. The

Diagenode MagMeDIP and IPure v2 kits were utilized to enrich and purify methylated cfDNA.

Prior to library amplification, qPCR quality checks were performed to ensure: (1) high

specificity (>95%) for enrichment of methylated AT DNA over unmethylated AT DNA, and (2)

appropriate adapter-ligation. This was followed by adapter-specific amplification of each MeDIP

and input control library and gel size selection. All sequencing was performed on the Illumina

HiSeq 2500 platform (50-bp single end reads).

Sequencing data pre-processing and differential methylation analysis

Raw sequencing data was aligned to hg19 reference genome using Burrows-Wheeler

alignment (BWA version 0.7.6a, using 'mem' option with default settings, except using –M

flag). Duplicated reads were marked by Picard tools (https://broadinstitute.github.io/picard/) and

collapsed to allow only one copy of the reads from reads that have the same alignment position.

Reads mapped to the multiple locations were removed by applying alignment quality threshold

(QA>5). As the sequencing data was single-end, the average DNA fragment length was

evaluated for both cfMeDIP and input controls using self-correlation technique (42). For each

library, the fragment coverage genomic profile was calculated using directional extension of all

aligned reads with estimated average fragment length (175bp in our case) using an in-house

BAM2WIG custom tool (M. Bilenky, unpublished, see http://www.epigenomes.ca/tools-and-

software). Within-patient differential methylation analysis for all comparisons was performed

using DMRHunter pipeline developed for cfDNA samples (**Supplemental Methods Section**).

Statistical analysis of mCRPC-associated DMRs

Spearman correlation analysis was performed using the base package of R (v3.5.3).

Receiver operating characteristic (ROC) curve analysis was performed using the pROC package

(1.14.0). All data was visualized using the ggplots2 package (v 3.1) and heatmaps from the

package ComplexHeatmap (1.2). Pathway analysis is described in **Supplemental Methods**

Section.

Results:

Study design and patient cohort

Tumor methylation signatures can be identified through comparison of cancer patients'

cfDNA with either matched tumor tissue or healthy control cfDNA (41). In the case of mCRPC,

biopsy of metastatic lesions in bone is not routinely feasible. Therefore, to identify methylation

markers associated with response to therapy, we opted for within/intra-patient differential

methylation analysis. We hypothesized that sequential collection of cfDNA from visit A

(baseline/pre-treatment) to visit B (week-12) to visit C (clinical progression) would reveal

changes in the cfDNA methylome that reflect responses to treatment, with each patient serving

as their own internal control. As outlined in **Figure 1**, intra-patient comparison could potentially

detect changes in abundance of methylated DNA fragments related to tumor response. For

instance, loss of methylated fragments may reflect loss/reduction of tumor cells sensitive to

treatment. In contrast, gains in methylated fragments could be associated with gains in treatment

resistant tumor cells, since changes in methylome could occur due to non-tumor/treatment

related changes. Therefore, we first examined intra-patient differentially methylated regions

(**DMRs**) to assess overall changes in cfDNA during treatment. We then integrated these DMRs

with tumour-derived methylation datasets.

We prospectively collected plasma from mCRPC patients receiving either Enzalutamide

(n=12) or Abiraterone acetate (n=4) treatment. In total, 45 blood samples were collected and

summarized in **Figure 2a**. Two Enzalutamide-treated patients were unable to provide Visit C

samples (P2 and P26) and one Abiraterone-treated patient progressed prior to Visit B (P36). The

majority of patients (14/16) presented with bone metastatic lesions upon mCRPC diagnosis, with

a subset showing lymph node and soft tissue metastases (**Figure 2b**). Twelve patients

demonstrated an initial favorable prostate-specific antigen (PSA) response ($\geq$ 50% decline from

baseline, **Figure 2c**), and most patients exhibited maximal PSA response (favorable or plateau)

around Visit B (**Supplementary Figure 1**). Overall time to clinical progression (**TTP**) varied for

each patient, ranging from rapid/primary resistance to therapy to treatment-driven resistance

following initial favorable response (**Figure 2d**).

Performance of cfMeDIP-seq strategy with PCa cell line DNA and cfDNA samples

We utilized an established cfMeDIP-seq approach, which was developed for low amounts

of cfDNA and capable of ctDNA fractions as low as 0.001% (41). This technique was able to

distinguish methylation patterns between various cancer types, including pancreatic, breast and

colorectal cancer. Briefly, methylated cfDNA is enriched using an antibody specific for

methylated CpG sites, followed by amplification of adapter-ligated libraries. For each sample,

10% of the DNA was reserved as an input control (without immunoprecipitation).

To benchmark this methodology for PCa samples, we generated a control from genomic

DNA derived from PCa cell line, 22Rv1, sheared to the same size as cfDNA (**Supplementary**

Figure 2). As an additional control, we spiked all samples with methylated and unmethylated

Arabidopsis thaliana (AT) DNA, and confirmed increased recovery of methylated AT DNA

(~80%) compared to unmethylated DNA (<0.06%) (**Supplementary Figure 2a**). Spiked AT

DNA was added following sequencing adapter ligation to cfDNA and not detectable during

sequencing. Increased enrichment of known methylated genes/regions in the genome of 22Rv1

cells, (43) including the *CDKN2A* gene body and the first exon of *TGFB2,* compared to the

unmethylated promoter of *HOXD8* was confirmed (**Supplementary Figure 2b**).

We performed cfMeDIP-seq on all 45 samples collected. The average starting amount of

dsDNA was 27 ng, and ranged from 9 to 50 ng, with no significant differences in cfDNA

amounts across visits and treatments (**Supplementary Figure 2c**). Following alignment and base

quality filtering, 48-62% of the cfMeDIP reads and 80-85% of input control reads remained for

DMR analysis (**Supplementary Figure 2d-e**). A majority of cfMeDIP-seq sequence reads

aligned within gene bodies (major transcript starting from transcriptional start site/TSS to

transcription termination site/TTS) and intergenic regions, as well as CpG Island (CGI), CGI

shores, and TSS/promoter regions (TSS +/-1.5Kb), together representing the entire genome

(**Supplementary Figure 2f**). As expected, enrichment of methylated cfDNA in the MeDIP

fraction over input control fraction was observed, especially within CpG rich regions. Similar

distribution of regions was observed between Enzalutamide and Abiraterone treated patients

(**Supplementary Figure 2 g-h**).

Intra-patient differential methylation analysis

To identify treatment-related differentially methylated regions (DMRs), we developed a

DMR calling strategy for intra-patient analysis termed "cfDNA DMRHunter" (See

Supplemental methods and **Supplementary Figure 3**). As location-specific cfDNA recovery

sensitivity is unknown *a priori*, we used input control libraries to ensure uniform cfDNA read

coverage across the whole genome. DMRHunter also uses the input controls to remove

alignment artifacts from analysis. After filtering low quality and PCR duplicated reads,

DMRHunter builds a background model and identifies differentially methylated CpGs (DMCs)

across the genome. Adjacent DMCs with the same differential methylation state (UP or DOWN)

were merged to identify DMRs with stringent z-score (>1.5), average read count (≥ 10) and false

discovery rate (FDR = 0.01) cut-offs. There were a large number of DMRs across all

comparisons and treatments, ranging from 93 000 to 154 000 DMRs (**Figure 3a**) with no

significant difference in median number of DMRs between comparisons and treatment (**Figure**

3b-c). The overall median DMR size was 74 bp, ranging from 2bp (single CpG sites) up to 4163

bp.

We next examined the directionality of these changes across comparisons.

Conventionally, hypermethylation or hypomethylation is the typical terminology for methylation

changes. However, for cfDNA, it is unclear whether these changes reflect a gain/loss of

methylated CpGs at these specific genomic sites or changes in DNA fragment abundance,

reflecting an altered cellular abundance contributing to the cfDNA pool. For our analysis we

opted to define these DMRs as gains (UP) or loss (DOWN) in methylated fragments. There was

no significant difference in the overall proportion of regions with increased or decreased

methylation in all visit comparisons (**Figures 3d-f**). The majority of DMRs were found within

gene bodies (protein coding and pseudogenes) and intergenic regions (**Figure 3g**). DMRs

near/within transcription start sites (TSS)/promoters and CpG Islands (CGI), as well as non-

coding RNA (ncRNA), were found, but with lower abundance. We chose to focus on DMRs

within promoter and gene body regions, as CpG methylation changes in these locations are

known to be associated with gene expression changes (44).

Methylation changes within promoter and gene body regions

The majority of DMRs within promoters (TSS +/- 1.5 Kb) were associated with protein

coding (PC) genes, followed by long non-coding RNA (lncRNA), pseudogenes (Pseudo) and

short non-coding RNA (sncRNA) regions (**Supplementary Figure 4a**). We observed minimal

overlap among promoters of different gene/ncRNA categories, and overall distribution of DMRs

in these regions did not vary between patients and visits (**Supplementary Figure 4b**). We

focused on protein coding genes with at least one promoter DMR. Analyzing recurrence of these

genes, we found that many of these genes were found in less than 5 patients for each comparison

(**Supplementary Figure 4c-e**). We examined the most common protein coding genes (with

DMRs in ≥ 5 patients) and found many genes overlapping across visit comparisons, with the

majority shared between Abiraterone and Enzalutamide groups (**Figure 3h**). The most common

genes showing either increases or decreases in promoter methylation are in **Supplementary**

Figure 4f, with certain genes implicated in PCa or other cancers. For instance, pyrroline-5-

carboxylate reductase 1 (*PYCR1*), which is involved in proline biosynthesis and can promote

tumor cell growth (45), often demonstrated gain of methylation at visit C. Pathway enrichment

analysis of promoter DMRs (**Supplementary Figures 5 and 6**) showed cancer-related pathways

such as Wnt and PI3K signaling were observed, with neuronal pathways being the most frequent.

Several promoter associated DMRs were only found among Enzalutamide-treated patients

(**Figure 3h and Supplementary table 2)**. The most frequently differentially methylated

promoter region (6-7 patients, 20 comparisons) within the Enzalutamide cohort was *FGFR1*,

which is overexpressed in PCa and implicated in metastasis (46).

We also examined the distribution of DMRs within gene bodies/known transcripts. The

majority of these DMRs were associated with protein coding gene bodies across

(**Supplementary Figure 7a-b**). In contrast to promoter DMRs, many genes with body DMRs

were shared across patients (**Supplementary Figure 7c-e**). Similarly, we focused on commonly

altered gene bodies and found extensive overlap between visit comparisons and treatments

(**Figure 3i**). We further examined net methylation change across these genes by scoring the ratio

of increased to decreased methylation within each gene body, excluding genes that had no net

change (i.e. equal number of UP and DOWN trends). We found 91 genes that were commonly

altered across all comparisons (**Supplementary Figure 7f**). In most cases, regardless of visit

comparison, the methylation pattern for these genes clustered closely for the same patients. One

such group of genes with increases in methylation at Visit C included *NFATC1*, which can

promote tumorigenesis (47), and a histone deacetylase, *HDAC4*. While several Enzalutamide-

specific genes were found (**Supplementary table 3**), common pathways including several

differentiation-related and neuronal pathways were found among gene-body associated DMRs

(**Supplementary Figures 8 & 9**).

Overall, examining the most frequently altered promoters or gene bodies revealed a few

cancer-related genes, but changes across all visit comparisons were variable. We assessed the

overall promoter methylation status of genes that have been established as differentially

methylated in PCa (**Figure 4a**), including *GSTP1*, *TBX15*, *AOX1* and members of the HOX

family of transcription factors (34). Among the most common differentially methylated gene

promoters were genes involved in tumorigenic processes, such as *RUNX3*, *RGS12* and *FBP1*,

with several differentially expressed in NE-CRPC (28), including the neuronal transcription

factor *PHOX2A* (48) and a regulator of apoptosis, *CTBP2* (49). DMRs in genes associated with

diagnosis and prognosis could reflect tumor cell response to treatments. While differential

promoter methylation events in the well-established PCa-specific hypermethylated gene, *GSTP1*,

were infrequent (**Figure 4b**), decreased gene body methylation coinciding with treatment

response (i.e. PSA reduction) was shown in some patients (i.e. P3). Similar methylation regions

were also observed in the PCa cell line LNCaP (50). Similarly, changes in *TBX15* methylation

(1^{st} intron) were associated with response to treatment (i.e. PSA) in certain patients (**Figure 4c**).

While promising, there were no consistent DMRs in these genes that could classify patients by

androgen-targeting treatment response. To further refine potential treatment associated

methylation changes, locations within the genome that experienced fluctuations in methylation

levels across all visits were next explored.

Common CpG sites with differential methylation across treatment visits

To find potential "treatment sensitive" CpG sites in the genome, we next examined

differentially methylated CpG sites (DMCs) among patients that completed all study visits,

specifically CpG sites that fluctuated in ≥2 visit comparisons. DMCs were analyzed as DMR size

for similar genomic regions varied within and across patients. We stratified the proportion of

DMCs that were uniquely found in single visit comparisons and those shared across all visits

(Scenarios 1-6 in **Figure 5a**). We identified DMCs present in two or more visit comparisons for

each patient (Scenarios 7-14) and associated them with gene promoters and bodies. Analysis of

gene promoters containing DMCs revealed minimal overlap amongst patients, with P4, P31, P38

and P3 showing the highest overlap of 8 shared genes: *NPBWR1*, *ZSCAN12*, *PCDHGA11*,

PHOX2A, *TBX10*, *TEX28*, *TKTL1* and *TSPAN32* (**Supplementary Figure 10a**). In contrast, we

observed a high degree of overlap amongst patients when looking at DMCs within gene bodies,

with the exception of P3 (**Supplementary Figure 10b**). Interestingly, we also noted a positive

correlation between the proportion of DMCs shared across all visits and TTP (**Figure 5a**).

We examined 3 major combined scenarios observed among 14 total: (1)

increase/decrease in methylation in visit C vs B and A (scenarios 7 and 8) (**Figure 5a**); (2)

increase/decrease in B and C vs A (scenarios 9 and 10,); (3) those that increase /decrease in B vs

A and then return to same levels in C (13 &14). The proportion of these three combined

scenarios was correlated with time to progression (TTP), and we found that those patients with

more DMCs that increased/decreased at visit B and maintained at visit C (scenarios 9 and 10)

had a longer TTP (**Figure 5b**). Although scenarios 7 & 8 did not correlate with TTP, those

patients with a higher proportion of scenarios 13 & 14 had a possible trend towards faster TTP

(**Supplementary figure 10 c-d**, $p=0.05$). This suggests that genome-wide cfDNA methylation

dynamics can reflect response to treatment in mCRPC patients. Importantly, changes in

methylation levels of these specific regions/genes could potentially be used as a monitoring tool

during androgen-target treatment.

Neuroendocrine-CRPC related cfDNA DMRs

Thus far, fluctuations in DMRs were analyzed for all cfDNA signals, which could include

tumour and non-tumour derived DNA. As many of the DMRs were associated with genes

implicated in several neuronal pathways and differentiation, we integrated cfDNA DMRs with

those identified from biopsy samples from NE-CRPC versus Adenocarcinoma-CRPC in an

independent mCRPC patient series (28). While this published dataset utilized reduced

representation bisulfite sequencing (RRBS), which primarily covers CpG rich regions (i.e.

CGIs), several of these DMCs were found within cfDNA DMRs (29 355 CpG sites out of 84 930

in NE-CRPC). In order to assess whether changes in these regions at earlier time points are

associated with overall response to androgen-targeting agents, we examined the overlap between

DMRs from the A vs B comparison and NE-associated DMRs. As P36 progressed prior to visit

B (**Figure 2d**), we included this patient in the analysis (examining A vs C). In this case, we

examined the changes in methylation from the perspective of visit A. That is, increased

methylation in visit A vs B/C (UP in A) or decreased methylation in visit A (DOWN in A). We

proposed that the abundance of methylated cfDNA fragments from certain regions may reflect

key genes/regions to monitor prior to initiating therapy. The overall proportion of these DMRs

that were increased or decreased in methylation at A vs B (or C for ID 36) varied across the

patients (**Figure 6a**). Interestingly, we noted that these DMRs (A vs B) appeared to be related to

TTP. We examined the ratio of DMRs containing NE-CRPC associated CpG sites with more

methylated fragments in Visit A to those with less methylation. In order to further delineate key

regions, we applied various FDR thresholds to these DMRs to examine the effect on correlation

with TTP (**Figure 6b**). We found an optimal set of regions that were negatively correlated with

TTP: that is, the more methylated cfDNA fragments containing these NE-CRPC associated

CpGs, the faster/shorter the TTP (**Figure 6c**). In a follow-up study analyzing matched biopsy and

plasma methylation signals from 5 Adenocarcinoma-CRPC and 6 NE-CRPC patients, several

prostate cancer and Neuroendocrine related methylation regions were found in cfDNA (40). We

also integrated our A vs B DMRs with this dataset and found similar correlation with TTP

(Spearman R = -0.56, p = 0.024). To determine whether the amount of NE-CRPC related

methylation patterns at visit A was associated with a faster TTP, we performed ROC analysis for

various TTP cut-offs (ranging from ≤20 to 35 weeks). We found that an increased ratio of NE-

CRPC DMRs UP in visit A can stratify patients that progressed as early as 25-weeks (area under

the curve, AUC: 0.927, CI: 0.773-1) (**Figure 6d**). These findings confirm the detection of NE-

CRPC related methylation signatures in cfDNA, but also show for the first time that these sites

may serve as potential predictive markers of resistance to androgen-targeting agents.

<u>Discussion</u>

In this study, we showed that methylome analysis of the cfDNA from mCRPC patients

can capture treatment-related epigenetic alterations, with and without integration with tissue

derived methylation data. We found changes in well-established PCa methylation markers

associated with response to treatment. Importantly, we demonstrated that cfDNA contains NE-

CRPC related methylation signals, which have the potential to be utilized as predictive markers

of treatment response. Methylation of several oncogenic driving pathways were detected in

cfDNA across time points, suggesting that methylome analysis of patients' liquid biopsies can

serve as a monitoring tool during treatment, as well as highlight potential targets for future

therapies.

A key advantage of liquid biopsy-based versus tissue-biopsy approaches is the ability to

capture tumor heterogeneity. To date, much focus has been placed on genomic alterations, which

can help identify *AR* mutations related to resistance to Abiraterone/Enzalutamide (31, 32).

However, few studies have highlighted that cfDNA methylation markers may help further

stratify patients (36, 37). In a recent study of 600 mCRPC patients receiving docetaxel, increased

methylated *GSTP1* at baseline was associated with longer overall survival, and loss of

methylation after two cycles of treatment corresponded with longer time to PSA progression

(51). Furthermore, patients that exhibited loss in methylated *GSTP1* after two cycles of treatment

had better overall survival. In this study, we observed some cases in which there were losses of

GSTP1 methylation in certain treatment responsive patients (PSA reduction). We also examined

other tissue-based methylation markers (promoter and gene body), including *TBX15* and *RUNX3*

(34), and similar heterogeneity amongst patients was observed. This in part could be explained

by differing mechanisms between docetaxel and androgen-targeting agents. Due to intra- and

inter-tumor heterogeneity (52, 53), androgen-targeting agents may not target all androgen-

independent tumor cells, which could also harbor these methylation markers. Furthermore, most

well-studied methylation markers were derived from earlier stages of PCa, whereas further

epigenomic instability is known to occur with the mCRPC state (54).

Through sequential analysis using a genome-wide approach, we highlighted progressive

changes in genes involved in tumorigenic processes in PCa and other cancers. Commonly altered

genes in the promoter region included *PYCR1* and *SHARPIN,* which are known to be upregulated

in PCa tissue and involved in processes such as proliferation and angiogenesis (55). Gene body

methylation changes were observed in genes that regulate various tumorigenic processes,

including *NFATC1* (47)*, PRDM16* (56)*,* and *OPCML* (57). Pathway analysis also further

highlighted key mechanisms that have been established in prostate tumor tissue studies,

including the wnt pathway (16). However, the methylation patterns amongst these genes

demonstrated extensive variability across patients and visits.

In order to further delineate key regions, we further assessed common regions that

changed during treatment. These changes appeared to reflect dynamics of treatment response.

For instance, patients that demonstrated CpG changes (up or down) in visit B and sustained the

changes by visit C appeared to have a longer TTP. In contrast, there was a trend towards shorter

TTP amongst patients that did not demonstrate sustained methylation changes. Among the genes

included was Neuropeptides B/W receptor 1 (*NPBWR1*), which was shown to be upregulated in

basal cells from prostate tissue (58), thus highlighting that genome-wide cfDNA methylation

analysis is able to capture tumor dynamics and may further have additive potential to mutation

analysis.

We also observed that many of the differentially methylated genes overlapped between

Enzalutamide- and Abiraterone-treated patients. Although there are ongoing studies to compare

efficacy of either treatment (59), the current consensus is that both perform similarly, and

deciding between treatments in a patient-specific manner remains an ongoing challenge. We

identified sets of genes altered in the promoter and gene body locations, including one large set

amongst Enzalutamide treated patients. We were limited by the number of Abiraterone-treated

patients, which may have limited heterogeneity, as seen in the larger Enzalutamide cohort.

Further studies are needed to examine if the methylome is different between these treatment

types. However, similar pathways were observed between the treatments, with neuronal

pathways being the most prominent.

458   Given these neuronal-related genes/pathways, we wanted to examine the utility of these

459 methylation changes in helping to predict treatment outcome, especially in identifying

460 Neuroendocrine-CRPC. With our unique study design of sequential sampling, we could detect

461 these methylation trends at earlier time points, demonstrating that the proportion of

462 Neuroendocrine-like DMRs correlated with faster/shorter time to progression. In particular, the

463 higher the proportion of certain DMRs with these NE-CRPC associated methylation patterns, the

464 faster the time to progression. While we did not have confirmatory biopsies or other markers of

465 NE disease (i.e. chromogranin A), increased methylation at Visit A could be utilized as

466 predictive markers of response to androgen-targeting agents. Although we are currently limited

467 to a single cohort, we now have a set of candidate regions that could be validated in additional

468 cohorts.

470 **<u>Conclusions</u>**

471   Our genome-wide sequencing protocol and analysis strategy has demonstrated the utility

472 of intra-patient monitoring of cfDNA methylation. If changes in the methylome during treatment

473 were sustained until clinical progression, these patients appeared to have a better prognosis with

474 androgen-targeting treatment. Interestingly, methylation patterns associated with a highly

475 aggressive form of the disease may serve as potential predictive markers. Further independent

476 validation of these markers is needed to assess this predictive potential.

480   **Future Perspectives**

481     Liquid biopsy based biomarkers will be playing a major role in how we treat patients

with highly aggressive prostate cancer. Many studies focused primarily on genomic alterations

associated with treatment resistance, but epigenomic markers could further explain the molecular

heterogeneous mechanisms that contribute to disease progression. Ultimately, we anticipate that

a combination of genomic and epigenomic markers will help facilitate optimal therapy decisions.

Summary points

• We performed genome-wide methylation analysis of sequentially collected cfDNA

489     samples from mCRPC patients receiving androgen-targeting therapy

490   • The DMRHunter pipeline was developed to identify differentially methylation regions

491     between study visits for each patient

492   • Changes in promoter and gene body methylation was identified in key novel genes as

493     well as previously established genes associated with tumourigenesis and the highly

494     aggressive form of the disease, Neuroendocrine-CRPC

495   • Patients that were able to sustain methylation changes throughout treatment tended to

496     present a longer time to clinical progression, suggesting that fluctuations in these CpG

497     sites could help monitor response to treatment

498   • Elevated levels of Neuroendocrine-CRPC related methylation signals could serve

499     potential predictive markers of resistance to androgen

500   • These findings could help with improving the clinical management of mCRPC using a

501     minimally invasive approach

Authors' contributions: B.B. designed the study for intra-patient comparison. A.M.J. designed

the clinical study for patient recruitment and sample collection. P.H. provided sample size

calculations required for recruitment. M.R.P. performed sample processing, cfMeDIP-seq library

preparation, and data analysis. M.B. developed and executed the DMRHunter pipeline and along

with M.H., assisted with analysis to interpret cfMeDIP-seq datasets. R.I. and G.D.B. performed

pathway analysis. S.Y.S. and D.D.D.C. designed the cfMeDIP–seq protocol and assisted with

optimization for this study design. A.M.J., N.E.F. and A.R.H. were involved in patient follow-up

and interpretation of data in the context of clinicopathological parameters. M.R.P., M.B. and

B.B. composed the manuscript with feedback from all co-authors involved.

Acknowledgements: We acknowledge the UHN Genitourinary clinic and Genitourinary

Biobank for recruiting patients, collecting blood samples and for patient follow-up/data

collection. All sequencing and initial data pre-processing was performed by the Princess

Margaret Genomics Centre and the UHN Bioinformatics and HPC Core, Princess Margaret

Cancer Centre, respectively. Differential methylation analysis was performed using the facilities

provided by the Canada's Michael Smith Genome Sciences Centre (British Columbia Cancer

Agency). We also acknowledge Shivani Kamdar (SHS/LTRI) for assisting with sample

processing and for helping to edit this manuscript.

Disclosures: There are no financial or non-financial interests to be declared.

Ethics approval and consent to participate: All patients in this study provided informed

written consent in accordance with approved institutional Research Ethics Board protocols from

University Health Network (UHN) and Sinai Health System (SHS). Progress updates were

submitted and approved annually by respective Research Ethics Boards. The study was

performed in accordance with the Declaration of Helsinki. No personal information for any

patient is presented in this manuscript. According to REB approvals, each patient was assigned

unique study identifiers, which are not linked to personal health information.

Availability of data and materials: All FASTQ and QC files will be made available in a public

repository upon publication of this manuscript. Additional supplementary data files have also

been included.

Funding: This study was supported by a Prostate Cancer Canada Movember Discovery Grant

(D2014-10) and an Astellas Prostate Cancer Innovation Fund (2017) to B. Bapat.

References

1.   Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, and Jemal A. Global cancer
statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36
cancers in 185 countries. *CA Cancer J Clin.* 68(6), 394-424 (2018).
2.   Global Burden of Disease Cancer C, Fitzmaurice C, Allen C, Barber RM, Barregard L,
Bhutta ZA, et al. Global, Regional, and National Cancer Incidence, Mortality, Years of
Life Lost, Years Lived With Disability, and Disability-Adjusted Life-years for 32 Cancer
Groups, 1990 to 2015: A Systematic Analysis for the Global Burden of Disease Study.
JAMA Oncology. 3(4), 524-48 (2017).
3.   Wong YN, Ferraldeschi R, Attard G, and de Bono J. Evolution of androgen receptor
targeted therapy for advanced prostate cancer. *Nat Rev Clin Oncol.* 11(6), 365-76 (2014).
4.   Sridhar SS, Freedland SJ, Gleave ME, Higano C, Mulders P, Parker C, et al. Castration-
resistant prostate cancer: from new pathophysiology to new treatment. *Eur Urol.* 65(2),
289-99 (2014).
5.   Crawford ED, Petrylak D, and Sartor O. Navigating the evolving therapeutic landscape in
advanced prostate cancer. *Urol Oncol.* 35S, S1-S13 (2017).
6.   Scher HI, Fizazi K, Saad F, Taplin ME, Sternberg CN, Miller K, et al. Increased survival
with enzalutamide in prostate cancer after chemotherapy. *N Engl J Med.* 367(13), 1187-
97 (2012).
7.   Loriot Y, Miller K, Sternberg CN, Fizazi K, De Bono JS, Chowdhury S, et al. Effect of
enzalutamide on health-related quality of life, pain, and skeletal-related events in
asymptomatic and minimally symptomatic, chemotherapy-naive patients with metastatic
castration-resistant prostate cancer (PREVAIL): results from a randomised, phase 3 trial.
Lancet Oncol. 16(5), 509-21 (2015).
8.   de Bono JS, Logothetis CJ, Molina A, Fizazi K, North S, Chu L, et al. Abiraterone and
increased survival in metastatic prostate cancer. *N Engl J Med.* 364(21), 1995-2005
(2011).
9.   Ryan CJ, Smith MR, Fizazi K, Saad F, Mulders PF, Sternberg CN, et al. Abiraterone
acetate plus prednisone versus placebo plus prednisone in chemotherapy-naive men with
metastatic castration-resistant prostate cancer (COU-AA-302): final overall survival
analysis of a randomised, double-blind, placebo-controlled phase 3 study. *Lancet Oncol.*
16(2), 152-60 (2015).
10.   Bambury RM, and Scher HI. Enzalutamide: The story of its development from bench to
bedside. *Urol Oncol.* (2015).
11.   Grist E, de Bono JS, and Attard G. Targeting extra-gonadal androgens in castration-
resistant prostate cancer. *J Steroid Biochem Mol Biol.* 145(157-63 (2015).
12.   Sumanasuriya S, and De Bono J. Treatment of Advanced Prostate Cancer-A Review of
Current Therapies and Future Promise. *Cold Spring Harb Perspect Med.* 8(6), (2018).
13.   Fizazi K, Tran N, Fein L, Matsubara N, Rodriguez-Antolin A, Alekseev BY, et al.
Abiraterone plus Prednisone in Metastatic, Castration-Sensitive Prostate Cancer. *N Engl J*
Med. 377(4), 352-60 (2017).
14.   James ND, de Bono JS, Spears MR, Clarke NW, Mason MD, Dearnaley DP, et al.
Abiraterone for Prostate Cancer Not Previously Treated with Hormone Therapy. *N Engl J*
Med. 377(4), 338-51 (2017).

15.   Armstrong AJ, Szmulewitz RZ, Petrylak DP, Holzbeierlein J, Villers A, Azad A, et al.
ARCHES: A Randomized, Phase III Study of Androgen Deprivation Therapy With
Enzalutamide or Placebo in Men With Metastatic Hormone-Sensitive Prostate Cancer. *J
Clin Oncol.* JCO1900799 (2019).

16.   Robinson D, Van Allen EM, Wu YM, Schultz N, Lonigro RJ, Mosquera JM, et al.
Integrative clinical genomics of advanced prostate cancer. *Cell.* 161(5), 1215-28 (2015).

17.   Armenia J, Wankowicz SAM, Liu D, Gao J, Kundra R, Reznik E, et al. The long tail of
oncogenic drivers in prostate cancer. *Nat Genet.* 50(5), 645-51 (2018).

18.   Wyatt AW, Annala M, Aggarwal R, Beja K, Feng F, Youngren J, et al. Concordance of
Circulating Tumor DNA and Matched Metastatic Tissue Biopsy in Prostate Cancer. *J
Natl Cancer Inst.* 109(12), (2017).

19.   Morrison GJ, and Goldkorn A. Development and Application of Liquid Biopsies in
Metastatic Prostate Cancer. *Curr Oncol Rep.* 20(4), 35 (2018).

20.   Liu W, Yin B, Wang X, Yu P, Duan X, Liu C, et al. Circulating tumor cells in prostate
cancer: Precision diagnosis and therapy. *Oncol Lett.* 14(2), 1223-32 (2017).

21.   Heller G, McCormack R, Kheoh T, Molina A, Smith MR, Dreicer R, et al. Circulating
Tumor Cell Number as a Response Measure of Prolonged Survival for Metastatic
Castration-Resistant Prostate Cancer: A Comparison With Prostate-Specific Antigen
Across Five Randomized Phase III Clinical Trials. *J Clin Oncol.* 36(6), 572-80 (2018).

22.   Bastos DA, and Antonarakis ES. CTC-derived AR-V7 detection as a prognostic and
predictive biomarker in advanced prostate cancer. *Expert Rev Mol Diagn.* 18(2), 155-63
(2018).

23.   Scher HI, Graf RP, Schreiber NA, Jayaram A, Winquist E, McLaughlin B, et al.
Assessment of the Validity of Nuclear-Localized Androgen Receptor Splice Variant 7 in
Circulating Tumor Cells as a Predictive Biomarker for Castration-Resistant Prostate
Cancer. *JAMA Oncol.* (2018).

24.   Antonarakis ES, Lu C, Luber B, Wang H, Chen Y, Zhu Y, et al. Clinical Significance of
Androgen Receptor Splice Variant-7 mRNA Detection in Circulating Tumor Cells of
Men With Metastatic Castration-Resistant Prostate Cancer Treated With First- and
Second-Line Abiraterone and Enzalutamide. *J Clin Oncol.* 35(19), 2149-56 (2017).

25.   Millner LM, Linder MW, and Valdes R, Jr. Circulating tumor cells: a review of present
methods and the need to identify heterogeneous phenotypes. *Ann Clin Lab Sci.* 43(3),
295-304 (2013).

26.   Bernemann C, Schnoeller TJ, Luedeke M, Steinestel K, Boegemann M, Schrader AJ, et
al. Expression of AR-V7 in Circulating Tumour Cells Does Not Preclude Response to
Next Generation Androgen Deprivation Therapy in Patients with Castration Resistant
Prostate Cancer. *Eur Urol.* (2016).

27.   Takeuchi T, Okuno Y, Hattori-Kato M, Zaitsu M, and Mikami K. Detection of AR-V7
mRNA in whole blood may not predict the effectiveness of novel endocrine drugs for
castration-resistant prostate cancer. *Res Rep Urol.* 8(21-5 (2016).

28.   Beltran H, Prandi D, Mosquera JM, Benelli M, Puca L, Cyrta J, et al. Divergent clonal
evolution of castration-resistant neuroendocrine prostate cancer. *Nat Med.* 22(3), 298-305
(2016).

29.   Corcoran RB, and Chabner BA. Application of Cell-free DNA Analysis to Cancer
Treatment. *N Engl J Med.* 379(18), 1754-65 (2018).

30. Ritch E, and Wyatt AW. Predicting therapy response and resistance in metastatic prostate
cancer with circulating tumor DNA. *Urol Oncol.* (2017).
31. Romanel A, Tandefelt DG, Conteduca V, Jayaram A, Casiraghi N, Wetterskog D, et al.
Plasma AR and abiraterone-resistant prostate cancer. *Sci Transl Med.* 7(312), 312re10
(2015).
32. Azad AA, Volik SV, Wyatt AW, Haegert A, Le Bihan S, Bell RH, et al. Androgen
Receptor Gene Aberrations in Circulating Cell-Free DNA: Biomarkers of Therapeutic
Resistance in Castration-Resistant Prostate Cancer. *Clin Cancer Res.* 21(10), 2315-24
(2015).
33. Yegnasubramanian S. Prostate cancer epigenetics and its clinical implications. *Asian
Journal of Andrology.* 18(4), 549-58 (2016).
34. Massie CE, Mills IG, and Lynch AG. The importance of DNA methylation in prostate
cancer development. *J Steroid Biochem Mol Biol.* 166(1-15 (2017).
35. Kron K, Liu L, Trudel D, Pethe V, Trachtenberg J, Fleshner N, et al. Correlation of ERG
expression and DNA methylation biomarkers with adverse clinicopathologic features of
prostate cancer. *Clin Cancer Res.* 18(10), 2896-904 (2012).
36. Mahon KL, Qu W, Devaney J, Paul C, Castillo L, Wykes RJ, et al. Methylated
Glutathione S-transferase 1 (mGSTP1) is a potential plasma free DNA epigenetic marker
of prognosis and response to chemotherapy in castrate-resistant prostate cancer. *Br J
Cancer.* 111(9), 1802-9 (2014).
37. Gordevicius J, Krisciunas A, Groot DE, Yip SM, Susic M, Kwan A, et al. Cell-Free DNA
Modification Dynamics in Abiraterone Acetate-Treated Prostate Cancer Patients. *Clin
Cancer Res.* 24(14), 3317-24 (2018).
38. Zhao F, Olkhov-Mitsel E, van der Kwast T, Sykes J, Zdravic D, Venkateswaran V, et al.
Urinary DNA Methylation Biomarkers for Noninvasive Prediction of Aggressive Disease
in Patients with Prostate Cancer on Active Surveillance. *J Urol.* 197(2), 335-41 (2017).
39. Hendriks RJ, Dijkstra S, Smit FP, Vandersmissen J, Van de Voorde H, Mulders PFA, et
al. Epigenetic markers in circulating cell-free DNA as prognostic markers for survival of
castration-resistant prostate cancer patients. *Prostate.* (2018).
40. Beltran H, Romanel A, Conteduca V, Casiraghi N, Sigouros M, Franceschini GM, et al.
Circulating tumor DNA profile recognizes transformation to castration-resistant
neuroendocrine prostate cancer. *J Clin Invest.* (2020).
41. Shen SY, Singhania R, Fehringer G, Chakravarthy A, Roehrl MHA, Chadwick D, et al.
Sensitive tumour detection and classification using plasma cell-free DNA methylomes.
Nature. 563(7732), 579-83 (2018).
42. Ramachandran P, Palidwor GA, Porter CJ, and Perkins TJ. MaSC: mappability-sensitive
cross-correlation for estimating mean fragment length of single-end short-read
sequencing data. *Bioinformatics.* 29(4), 444-50 (2013).
43. Kamdar SN, Ho LT, Kron KJ, Isserlin R, van der Kwast T, Zlotta AR, et al. Dynamic
interplay between locus-specific DNA methylation and hydroxymethylation regulates
distinct biological pathways in prostate carcinogenesis. *Clin Epigenetics.* 8(32 (2016).
44. Baylin SB, and Jones PA. Epigenetic Determinants of Cancer. *Cold Spring Harb
Perspect Biol.* 8(9), (2016).
45. Zeng T, Zhu L, Liao M, Zhuo W, Yang S, Wu W, et al. Knockdown of PYCR1 inhibits
cell proliferation and colony formation via cell cycle arrest and apoptosis in prostate
cancer. *Med Oncol.* 34(2), 27 (2017).

46. Yang F, Zhang Y, Ressler SJ, Ittmann MM, Ayala GE, Dang TD, et al. FGFR1 is
essential for prostate cancer progression and metastasis. *Cancer Res.* 73(12), 3716-24
(2013).
47. Manda KR, Tripathi P, Hsi AC, Ning J, Ruzinova MB, Liapis H, et al. NFATc1 promotes
prostate tumorigenesis and overcomes PTEN loss-induced senescence. *Oncogene.*
35(25), 3282-92 (2016).
48. Wang R, Chen X, Xu T, Xia R, Han L, Chen W, et al. MiR-326 regulates cell
proliferation and migration in lung cancer by targeting phox2a and is regulated by
HOTAIR. *Am J Cancer Res.* 6(2), 173-86 (2016).
49. Takayama K, Suzuki T, Fujimura T, Urano T, Takahashi S, Homma Y, et al. CtBP2
modulates the androgen receptor to promote prostate cancer progression. *Cancer Res.*
74(22), 6542-53 (2014).
50. Yang YA, Zhao JC, Fong KW, Kim J, Li S, Song C, et al. FOXA1 potentiates lineage-
specific enhancer activation through modulating TET1 expression and function. *Nucleic*
Acids Res. 44(17), 8153-64 (2016).
51. Mahon KL, Qu W, Lin HM, Spielman C, Cain D, Jacobs C, et al. Serum Free Methylated
Glutathione S-transferase 1 DNA Levels, Survival, and Response to Docetaxel in
Metastatic, Castration-resistant Prostate Cancer: Post Hoc Analyses of Data from a Phase
3 Trial. *Eur Urol.* (2018).
52. Brocks D, Assenov Y, Minner S, Bogatyrova O, Simon R, Koop C, et al. Intratumor
DNA methylation heterogeneity reflects clonal evolution in aggressive prostate cancer.
Cell Reports. 8(3), 798-806 (2014).
53. Aryee MJ, Liu W, Engelmann JC, Nuhn P, Gurel M, Haffner MC, et al. DNA
methylation alterations exhibit intraindividual stability and interindividual heterogeneity
in prostate cancer metastases. *Sci Transl Med.* 5(169), 169ra10 (2013).
54. Friedlander TW, Roy R, Tomlins SA, Ngo VT, Kobayashi Y, Azameera A, et al.
Common structural and epigenetic changes in the genome of castration-resistant prostate
cancer. *Cancer Res.* 72(3), 616-25 (2012).
55. Huang H, Du T, Zhang Y, Lai Y, Li K, Fan X, et al. Elevation of SHARPIN Protein
Levels in Prostate Adenocarcinomas Promotes Metastasis and Impairs Patient Survivals.
Prostate. 77(7), 718-28 (2017).
56. Zhu S, Xu Y, Song M, Chen G, Wang H, Zhao Y, et al. PRDM16 is associated with
evasion of apoptosis by prostatic cancer cells according to RNA interference screening.
Mol Med Rep. 14(4), 3357-61 (2016).
57. Simovic I, Castano-Rodriguez N, and Kaakoush NO. OPCML: A Promising Biomarker
and Therapeutic Avenue. *Trends Cancer.* 5(8), 463-6 (2019).
58. Zhang D, Park D, Zhong Y, Lu Y, Rycaj K, Gong S, et al. Stem cell and neurogenic
gene-expression profiles link prostate basal cells to aggressive prostate cancer. *Nat*
Commun. 7(10798 (2016).
59. Hara I, Yamashita S, Nishizawa S, Kikkawa K, Shimokawa T, and Kohjimoto Y.
Enzalutamide Versus Abiraterone as a First-Line Endocrine Therapy for Castration-
Resistant Prostate Cancer: Protocol for a Multicenter Randomized Phase 3 Trial. *JMIR*
Res Protoc. 7(7), e11191 (2018).
727
728

 Figure Legends

Figure 1. Within patient differential methylation analysis strategy to monitor temporal

changes in cfDNA.

In order to help identify methylation changes associated with treatment in cfDNA, we opted to

perform within-patient methylation analysis to identify differentially methylated regions (DMRs)

associated with tumor response and/or resistance to current treatments. Each patient in this study

received either Enzalutamide or Abiraterone acetate and blood was collected prior to initiating

therapy (**Visit A**), at 12-weeks during treatment (**Visit B**) and upon clinical

progression/treatment change (**Visit C**). We applied an established genome-wide method to

detect methylated cfDNA fragments followed by extensive analysis to identify DMRs associated

with treatment response. We compared all study visits available (Visit B vs A, C vs B, and C vs

A) to find losses or gains in methylated cfDNA fragments. For instance, loss of treatment

responsive tumor cells (green methylated DNA fragments at visit B) or gains in treatment

insensitive/resistant tumor cells (blue DNA fragments at visit C) could be detected with this

strategy.

Figure 2. Sample collection and patient clinical follow-up overview.

(**a**) Plasma samples from mCRPC patients receiving Enzalutamide (12 patients) or Abiraterone

acetate (4 patients) were collected at baseline (Visit A), at week-12 (+/- 2 weeks, Visit B) and

upon clinical progression (Visit C). The total number of samples collected at each visit is shown.

2 Enzalutamide-treated patients (P2 and P26) were unable to provide samples for Visit C and 1

Abiraterone-treated patient (P36) progressed prior to Visit B. For these patients, the date of

treatment change/clinical progression was recorded for data analysis. (**b**) Pie chart summarizes

the distribution of metastasis locations at mCRPC diagnosis. (**c**) Bar plot shows best PSA

response for each patient during treatment (nadir), expressed as a percentage of Baseline/Visit A

PSA. Dotted line indicates $\geq$ 50% decline in PSA from Visit A. (**d**) Overall timeline of study

follow-up starting from Visit A to Visit C/Final visit is shown for all patients. Green dots

represent Visit B, red dots are Visit C or final study visit (if Visit C was not collected), yellow

indicates lowest PSA level (nadir) and blue dots show first PSA rise during treatment.

Figure 3. Summary of intra-patient DMR analysis.

(**a**) The total number of DMRs for all patients, visit comparison and treatment. (**b**) Boxplot

summarizing median, first and third quartile of all comparison DMRs. Whiskers represent the

highest and lowest number of DMRs. (**c**) DMR totals were further stratified by treatment type.

Boxplots illustrate percentage of DMRs that increased or decreased in methylation in (**d**) B vs A,

(**e**) C vs B and (**f**) C vs A. (**g**) The overall proportion of DMRs that were identified near/within

CGI, shore, TSS/promoter, ncRNA, intergenic and gene body regions is shown for all visit

comparisons (In order of B vs A, C vs B and C vs A). (**h**) Venn diagrams showing the overlap of

protein coding genes with DMRs near their promoters is shown for all visit comparisons and

between Enzalutamide- and Abiraterone-treated patients. (**i**) Similarly, the overlap of protein

coding regions with DMRs within gene bodies is shown.

Figure 4. Key prostate cancer related methylation changes observed in cfDNA

(**a**) Heatmap shows methylation trends in promoter regions for all patients and comparisons of

the most frequently methylated genes in PCa. For each gene, the proportion of DMRs with

increased (blue) and decreased (red) methylation was calculated and the net change is shown.

Grey indicates no net change in methylation. Representative peaks show methylation levels

(normalized read counts) of (**b**) *GSTP1* (black dotted lines indicating gene body) and (**c**) *TBX15*.

LNCaP cell line MeDIP-seq data is also shown.

Figure 5. Common differentially methylated CpGs (DMCs).

(**a**) For patients that were able to provide samples for all three visits, the proportion of CpG sites

that were differentially methylated in a single comparison and shared across 2 or more

comparisons is shown. 14 possible scenarios are represented by different colors. TTP in weeks

are also indicated for each patient. (**b**) Spearman correlation analysis of the proportion of DMCs

that were found to increase or decrease in B/C vs A (scenarios 9 & 10) against time to

progression (Spearman rho and p values are shown).

Figure 6. NE-CRPC related DMRs in cfDNA.

(**a**) For each patient, the proportion of DMRs that demonstrated methylation differences in NE-

CRPC associated regions in visit A vs B. (**b**) For DMRs that contained NE-CRPC associated

CpGs, various FDR thresholds were applied to assess for optimal regions that correlated with

TTP. (**c**) Using the optimal FDR cut-off, the ratio of NE-associated DMRs with increased

methylated fragments in visit A vs B to decreased fragments was correlated with TTP (Spearman

rho and p value is shown). For P36, A vs C comparison was used as this patient progressed prior

to visit B. (d) ROC analysis shows optimal AUC (0.927, 95% CI: 0.773-1) that stratifies patients

according to TTP (<25 weeks).

Tumor-related cfDNA   **Other cfDNA**

Down: Loss/Reduction of methylated cfDNA fragment
Source: Treatment responsive cells

UP: Gain/Increase of methylated cfDNA fragment
Source: Treatment resistant cells

Visit A: Baseline   **Visit B**: Week-12   **Visit C**: Clinical Progression

a)

	Visit A: Baseline/ Prior to treatment	Visit B: Week-12 (+/- 2 weeks)	Visit C: Clinical Progression/ Change in treatment
Enzalutamide (No. of visits)	12	12	10
Abiraterone (No. of visits)	4	3	4

b) Metastasis sites at mCRPC diagnosis

- Bone
- Bone + Lymph node
- Bone + Lymph node + Lung
- Lymph node

c)

Best PSA response (% of Visit A)

ABI
ENZA

Patient: P3 P7 P1 P2 P4 P8 P9 P15 P31 P12 P6 P38 P26 P37 P36 P28

d)

Weeks from Visit A

- Visit B
- Visit C/Final visit
- PSA Nadir
- First PSA rise