Transcriptome profiling in Indian Cattle revealed novel insights into response to heat stress

Raja Ishaq Nabi Khan [a†], Amit Ranjan Sahu [a†], Waseem Akram Malla [a], Manas Ranjan Praharaj [b], Neelima Hosamani [b†], Shakti Kumar [b], Smita Gupta [a], Shweta Sharma [a], Archana Saxena [a], Anshul Varshney [a], Pragya Singh [a], Vinay Verma [c], Puneet Kumar [c], Gyanendra Singh [c], Aruna Pandey [a], Shikha Saxena [a], Ravi Kumar Gandham [b#], Ashok Kumar Tiwari [d#]

[a] Division of Veterinary Biotechnology, Indian Veterinary Research Institute, Bareilly, India
[b] Computational Biology and Genomics, National Institute of Animal Biotechnology, Hyderabad, India
[c] Division of Physiology and Climatology, Indian Veterinary Research Institute, Bareilly, India
[d] Division of Biological Standardization, Indian Veterinary Research Institute, Bareilly, India

[†]Equal Contributors

[#] Corresponding Authors: Ravi Kumar Gandham ,ravigandham@niab.org.in

Ashok Kumar Tiwari, aktiwari71d@gmail.com

Abstract

Heat stress induced by high environmental temperature and humidity affects livestock production and health. With global warming on the uprise, indigenous cattle known for their heat tolerance are gaining importance than the Crossbreds. However, systems biology behind this phenotype in indigenous cattle is less known. This study, revealed novel insights in response to heat stress in Indian cattle (Tharparkar breed – Indigenous breed vs Vrindhavani – Crossbred).The number of DEGs in Crossbred were found to be more, than in Tharparkar, suggesting a greater dysregulation in systems biology in Crossbred. A contrast in gene expression was observed with 18.5 % of upregulated genes in Crossbred (Vrindavani cattle) were downregulated in Tharparkar and 17.5% upregulated genes in Tharparkar were downregulated in Crossbred. The increased HSPs levels have been found positively correlated with tolerance in many species. Upregulation of *HSF, HSP70, HSP90*, and activation of EIF2 signaling pathway in Tharparkar and vice-versa in Crossbred delineates how Tharparkar, withstands heat stress. Unlike Crossbred, Tharparkar is not only endowed with higher expression of the scavengers (*UBE2G1, UBE2S*, and *UBE2H*) of misfolded proteins but also with protectors (*VCP, Serp1*, and *CALR*) of naïve unfolded proteins. Further, the apoptotic genes that were dysregulated in both genetic groups indicated a relatively higher probability of apoptosis in Crossbred than in Tharparkar. Also, higher expression of the antioxidants in Tharparkar enables it to cope up with higher levels of

37 free radicals generated as a result of heat stress. In this study we found relevant molecules/genes
38 dysregulated in Tharparkar in the direction that can counter heat stress. To best of our knowledge
39 this is a comprehensive comparison between Tharparkar and Crossbred under heat stress using
40 transcriptome analysis.

42 **Introduction**
43

Cattle being homoeothermic, modulate their internal body temperature in sync to
environmental temperature by equilibrating the amount of heat produced within the body and
dissipating it to the ambient environment. The stress that arises due to disproportionate
thermodynamic behavior between cattle and its surrounding environment is termed as heat stress
[1]. Environmental induced hyperthermic stress lowers feed intake, which in turn reduces growth,
milk production and reproductive efficiency, thereby negatively affecting the economics of
livestock keepers [2-4]. Heat stress has been associated with reduced fertility through its deleterious
impact on oocyte maturation and early embryonic development [5]. Increased morbidity and
mortality was observed in animals due to immune depressive effect of heat stress [6].

India has a wide variety of indigenous cattle breeds distributed throughout its agro-
climatic zones. These are known for their natural tolerance to tropical heat [7,8].To meet the
growing demand for milk and to combine the heat tolerance and tick resistance of zebu with the
productivity of temperate dairy breeds [9] several crossbreeding programs were taken up in India.
Every state had its own crossbreeding policy, which is agro-climatic and breed-specific. Though
the zebu crosses with European breeds produced more milk than zebu, they were found not
withstanding heat/solar radiation [10]. Crossbreds are susceptible to tropical diseases and require a
constant input of good management conditions [8]. Antagonism exists between heat tolerance and
milk productivity [11]. The adaptive capacity to heat stress varies between species and genetic
groups within species. Among various adaptive mechanisms, physiological adaptability seems to
be the primary step in cattle. Sahiwal cows better regulate body temperature in response to heat
stress than Karan Fries [8]. It was observed that Ongole cattle rely on the respiration rate to
maintain thermal balance, while, Bali cattle rely on rectal temperature [12]. In Brazil, Sindhi and
Girolando breeds showed better physiological response to thermal stress than Gir cattle [13].
Increase in respiration rate was reported in Nellore breed when exposed to heat load [14].

In India, Tharparkar is one among the best dairy breeds. It is adapted to the Indian states
of Punjab and Haryana. [15,16]. It is considered to be the hardiest, disease resistant, heat tolerant
and tick resistant indigenous cattle breed of the country [17]. This breed has also been used in
several crossbreeding programs. Currently, percentage of purebreds is exceptionally low in India
(Department of Animal Husbandry & Dairying, Govt. of India). Most of the farmer's in India
have Crossbreds and the percentage of exotic inheritance in these Crossbreds is unknown.
Vrindavani, a Crossbred (synthetic), with 27% of Indigenous blood and 73% of exotic
inheritance [18] is a representation of the kind of admixture that prevails in Indian cattle. Therefore,
comparing Tharparkar with Vrindavani may establish the lost importance of Indigenous cattle
and would further emphasize the need of conserving our indigenous purebreds because of
advantageous traits like Heat tolerance.

Studies explaining the difference between the genetic groups (Crossbreds and Indigenous
cattle) have been done mainly to address the physiological responses vis – a – vis heat stress and
very few studies at the genomic level have been taken up [19,20].Also, there is an increasing need to
develop methods by combining the knowledge from -omics technologies to identify heat tolerant
animals [11]. Transcriptome profiling / RNA-Sequencing (RNA-seq) is a high throughput omics
approach to measure relative global changes in the transcripts under specific condition(s) [21-23] to
study the systems biology behind a phenotype [23,24]. RNA - seq allows for analysis of
transcriptome in an unbiased way, with, a tremendous dynamic detection range (>8,000 fold),
and low background signals [25]. It has been used as an investigating tool in understanding disease
pathogenesis [26,27] and differential physiological response to various biotic and abiotic factors
[28,29].

In this study, Tharparkar and Crossbred - Vrindavani cattle were subjected to heat stress
and blood samples were collected on 0^{th} day and 7^{th} day, as it is known that short term
acclimation occurs around 5 - 6 days [30,31]. The transcriptome of 7^{th} day was compared and with
0^{th} day in both the genetic groups to understand their differential response to heat stress.

Results

Physiological Parameters

96   The overview of the analysis is given in Figure 1. Respiration rate (RR), rectal
97   temperature (RT) and T3 level increased significantly (p<0.05) on 7^{th} - day post heat stress in

98 both the genetic groups (n=5) (Figure 2). However, the increase in   RR, RT and T3 level, was

99 found significantly (P<0.05) higher in Crossbred than in Tharparkar.

Comparison of DEGs of Crossbred and Tharparkar under heat stress

The differentially expressed genes for each genetic group were obtained on comparing

the 0 day and 7th day RNA-seq data using EdgeR after obtaining the gene counts from RSEM.

Under heat stress, global expression profiles of Crossbred and Tharparkar were identified with

6042 and 4718 differentially expressed genes (DEGs), respectively (Supplementary Table 1).

Among these, 3481 DEGs were found common between the two genetic groups, while 2561 and

1238 DEGs were uniquely found in Crossbred and Tharparkar, respectively (Figure 3a).

Additionally, 3132 and 2924 genes were upregulated and downregulated in Crossbred,

respectively, while 2367 and 2358 genes were upregulated and downregulated in Tharparkar,

respectively (Figure 3b). On comparison of upregulated and downregulated genes, 724 and 1416

genes were found uniquely upregulated and 514 and 1145 genes were found uniquely

downregulated in Tharparkar and Crossbred, respectively. The comparison also revealed that

17.5% of upregulated genes (1278) in Tharparkar were downregulated in Crossbred and 18.5%

downregulated genes (1344) in Tharparkar were upregulated in Crossbred. However, the number

of common upregulated and downregulated genes in both the genetic groups were 357 (4.9%)

and 498 (6.8%), respectively (Figure 3c).

Analysis of knowledge-based genes

Under heat stress four major physiological processes are found to be usually associated

with heat stress viz. elicitation of unfolded protein response (UPR) in cells; Induction of

apoptosis; Ubiquitination and; Imbalance in production of ROS and antioxidants (refer to

materials and methods for details). Heat shock genes and its associated genes are involved in

elicitation of unfolded protein response (UPR) in cells. Heat shock genes have been found

dysregulated under heat stress in both the genetic groups. Most of the genes encoding Heat

shock proteins (HSPs) - Heat shock 70 kDa protein 4   (*HSPA4),* Heat shock cognate 71 kDa

protein *(HSPB8),* Heat shock 70 kDa protein 1A *(HSPA1A) ,*Heat shock cognate 71 kDa protein

(HSPA8), Heat shock protein HSP 90-beta *(HSP90AB1)* and Heat shock protein HSP 90-alpha

(*HSP90AA1*) and heat shock protein regulating factors- Heat shock factor 1 *(HSF1)* and

Eukaryotic Translation Elongation Factor 1 Alpha 1 *(EEF1A1)* have been found to be

downregulated/not-differentially expressed in Crossbred but upregulated in Tharparkar.
However, Calcium/Calmodulin Dependent Protein Kinase II Delta *(CAMK2D) that* is involved
in the regulation of expression of heat shock genes was upregulated in Crossbred and
downregulated in Tharparkar.

Among the apoptotic genes, genes encoding Bcl-2-like protein 11 *(BCL2L11)*, Tumor
necrosis factor ligand superfamily member 6 *(FASLG)*, TIR domain-containing adapter
molecule 2 *(TICAM2)*, Toll-like receptor 4 *(TLR4)*, Adenomatous polyposis coli protein
(APC), Caspase-3 *(CASP3)*, Mitogen-activated protein kinase 8 *(MAPK8)*, Mixed lineage
kinase domain-like protein *(MLKL)*, Late endosomal/lysosomal adaptor and MAPK and MTOR
activator 5 *(XIP)*, Vimentin *(VIM)*, and High mobility group protein B2 *(HMGB2)* were found to
be upregulated in Crossbred and downregulated in Tharparkar. The number of upregulated genes
involved in achieving the balance of ROS production and antioxidants, were found to be more in
Tharparkar than in Crossbred. Among these, genes encoding Glutathione peroxidase 3
(GPX3), Nudix Hydrolase 2 *(NUDT2)*, Catalase *(CAT)*, Cytochrome c *(CYCS)*, Copper
chaperone for superoxide dismutase *(CCS)*, Peroxiredoxin-5 *(PRDX5)*, Peroxiredoxin-6
(PRDX6), Peroxiredoxin-1 *(PRDX1)*, Superoxide dismutase *(SOD1)*, and Cytochrome b-245
heavy chain *(CYBB)* were found either downregulated/not-differentially expressed in Crossbred
and upregulated in Tharparkar. More number of genes involved in Ubiquitination were
differentially expressed in Crossbred than in the Tharparkar. Genes encoding Ubiquitin-
conjugating enzyme E2 G1 *(UBE2G1)*, Ubiquitin-conjugating enzyme E2 *(UBE2S)*, Ubiquitin-
conjugating enzyme E2 H *(UBE2H)*, Ubiquitin A-52 residue ribosomal protein fusion product 1
(UBA52), and Ubiquitin-activating enzyme E1 *(UBA1)* have been found downregulated/not-
differentially expressed in Crossbred and upregulated in Tharparkar. However, Valosin-
containing protein *(VCP)*, RING finger protein 40 *(RNF40)*, and Ubiquitin-conjugating enzyme
E2 L3 *(UBE2L3)* have been found downregulated in Crossbred but not-differentially expressed
in Tharparkar. Among the genes involved in Unfolded Protein folding response (UPR), genes
encoding Membrane-bound transcription factor site-1 protease *(MBTPS1)*, Cyclic AMP-
responsive element-binding protein 3-like protein 1 *(CREB3L1)*, Stress-associated endoplasmic
reticulum protein 1 *(SERP1)*, Glycogen synthase kinase-3 alpha *(GSK3A)*, Eukaryotic
translation initiation factor 2 subunit 3 *(EIF2S3)*, Calreticulin *(CALR)*, and Stress-associated

endoplasmic reticulum protein 1 (*SERP1)* have been found downregulated in Crossbred and

upregulated in Tharparkar (Figure 4).

Protein - protein interaction (PPI) network revealed functional importance of HSP70
(*HSPA8* and *HSPA1A)* and ubiquitin (UBB, UBA52), in coordinating genes involved in heat
stress

A total of 246 knowledge-based genes were identified from the reactome database. Out of

these 177 and 194 genes were found to be differentially expressed in Tharparkar and Crossbred,

respectively. Among these 126 genes were found to be commonly differentially expressed in

Tharparkar and Crossbred. PPI network for these common knowledge-based genes between

Tharparkar and Crossbred was constructed (Supplementary Figure 1). In PPI networks, hubs

define the functional and structural importance of a network. The genes, which act as hubs in PPI

networks were found to be *UBB, UBA52, HSPA8*, and *HSPA1A* (Supplementary Figure 1).

Among the 4 hubs, *UBB* was downregulated in both genetic groups and the rest were

downregulated in Crossbred and upregulated in Tharparkar.

A change in the expression of the hub protein will have a larger effect than change in

expression of non-hub proteins [32]. Therefore, UBB, UBA52, HSPA8, and HSPA1A are taken to

be critical for coordinating the changes in systems biology under heat stress. The hubs HSPA8

and HSPA1A are connected to genes that are associated with regulation of stress viz.

nucleoporins genes - *NUP188, NUP155, NUP210 & NUP214*; BAG family molecular chaperone

regulators - *BAG1, BAG3 & BAG4*; Heat Shock Protein Family A - *HSPA5, HSPA4, HSPA12B*

& HSPA9; DnaJ Heat Shock Protein Family i.e. HSP40 - *DNAJA1, DNAJC2 & DNAJB6*; Heat

shock factor - *HSF1*; Ubiquitin - *UBB & UBA52* and; Sirtuin - *SIRT1*. The hubs - UBB and

UBA52 are connected to molecules of different proteasome subunits viz. ⍺ type subunits -

PSMA1 & PSMA2; β type subunits - *PSMB4 & PSMB8*; ATPase subunits - *PSMC2 & PSMC5*

and non-ATPase subunits - *PSMD2 & PSMD13*. These hubs were also found connected to

ubiquitin specific peptidases - *USP9X* and *USP7* and Ubiquitin-conjugating enzyme - *UBE2B,*

UBE2G1, UBE2Z, UBE2H, UBE2J2, UBE2S & UBE2D2.

Canonical pathway analysis by Ingenuity Pathway Analysis (IPA) revealed contrast in
signaling pathways in Crossbred and Tharparkar

Canonical pathways associated with Crossbred and Tharparkar are represented in Figure

5a and 5b. In Crossbred, Oncostatin M Signaling, Phospholipase C Signaling, EIF2 Signaling,

Integrin Signaling, IL-3 Signaling, and CXCR4 Signaling were found to be highly inactivated (Z
– score > 2.0) and PTEN signaling was found to be highly activated (Z – score < 2.0). In
Tharparkar, EIF2 Signaling, Androgen Signaling, Oncostatin M Signaling, α-Adrenergic
Signaling, BMP signaling pathway, and UVC-Induced MAPK Signaling were found to be highly
activated and PTEN signaling was found to be inactivated. The canonical pathway Oncostatin M
Signaling and EIF2 Signaling were found to have the highest ratio of genes involved vis-a-vis
the genes in the database in Crossbred and Tharparkar, respectively.

While carrying out comparative analysis through IPA, Calcium-induced T Lymphocyte
Apoptosis, BMP signaling pathway, UVC-Induced MAPK Signaling, Regulation of Cellular
Mechanics by Calpain Protease, fMLP Signaling in Neutrophils, Melatonin Signaling, and
Leukocyte Extravasation Signaling, were found inactivated in Crossbred and activated in
Tharparkar (Supplementary Figure 2). Genes involved in Oncostatin M Signaling-   Growth
factor receptor-bound protein 2 (*GRB2)*, GTPase HRas *(HRAS)*, Janus kinase 1 *(JAK1)*, Janus
kinase 3 *(JAK3)*, Mitogen-activated protein kinase kinase 1 *(MAP2K1)*, Mitogen-activated
protein kinase 1 *(MAPK1)*, Oncostatin-M *(OSM)*, Ras-related protein Rap-1b *(RAP1B)*, Ras-
related protein Rap-2a *(RAP2A)*, Signal transducer and activator of transcription 1-alpha/beta
(STAT1), Signal transducer and activator of transcription 5B *(STAT5B)*, Non-receptor tyrosine-
protein kinase *(TYK2)*, and Ras-related protein *(RRAS)* were found downregulated in Crossbred
and upregulated in Tharparkar (Figure 6a, b). While the key genes involved in PTEN Signaling
pathway – Fas Ligand (*FASLG)*, member of RAS oncogene family *(RAP2A)*, Bcl-2-like protein
11 *(BIM)*,Caspase-3 *(CASP3)* and microspherule protein 1 (*MSP58)* were found upregulated in
Crossbred and downregulated in Tharparkar as well (Figure 7a, b).

Variation in microRNAs and Transcription factors

IPA, on evaluating the differentially expression genes predicts miRNAs and
Transcription Factors (upstream regulators). In Crossbred, 111 miRNAs were found to be
inactivated and 37 activated. In Tharparkar, 205 miRNAs were found to be inactivated and 272
activated. Among them, 52 microRNAs were found common between the two genetic groups.
Most of the common miRNAs were found activated in Crossbred and inactivated in Tharparkar
(Supplementary Figure 3). miR-4779, miR-4651, miR-1207-5p, miR-6967-5p and miR-504-3p
are the top 5 miRNAs that were activated in Crossbred and inactivated in Tharparkar.

Various Transcription factors were found to regulate the expression of the identified
DEGs. Transcription factors, 19 in Tharparkar (11 activated and 8 inactivated) and 26 in
Crossbred (8 activated and 18 inactivated) were identified in IPA that regulate the expression of
DEGs. Among them, *PAX5, MTA3, MYC, PROX1* and *SMAD7* in Crossbred and, *HMGA1, MAF,*
MAX NOTCH22 and *NCOR1* in Tharparkar are the top 5 upregulated and activated TFs. On
comparing the TFs of Tharparkar and Crossbred, it was found that *BHLHE40, HMGA1,*
HMGB1, IKZF1, and *TCF7* were found to be common. *BHLHE40, HMGA1,* and *TCF7* were
found to be activated in Tharparkar and inactivated in Crossbred and it was vice - versa with
HMGB1 and *IKZF1* (Supplementary Figure 4)

Real-time validation.

Six genes (*HSF1, SOD1, CALR, GSK3A, CAT & GPX3*) that were upregulated in
Tharparkar but downregulated/not expressed in crossbred and four genes (*CASP3, FASLG,*
BCL2L11 & APC) that were upregulated in crossbred but downregulated in Tharparkar were
considered for Real time PCR based on their role in heat stress. Briefly, CALR is involved in
unfolded protein response upon Endoplasmic reticulum stress; HSF1 in trimer induces the
expression of HSPs; GSK3A, kinase inhibits trimerization of HSF1 regulating the expression of
HSPs; SOD1 converts free radical into hydrogen peroxide, which is further converted by GPX3A
and/or CAT into water and oxygen molecules in regulatory pathway of ROS scavenging and ;
BCL2L11, FASLG, APC and CASP3 induce apoptosis in stressed cell. The expression of genes
was in concordance with the RNA- Seq results (Supplementary Figure 5 and Supplementary
table 2).

Discussion

Heat stress is a natural phenomenon that affects domestic animals in tropical, sub-tropical
and often in temperate regions of the world during summer months. Heat and humidity during
the summer months combine to make an uncomfortable environment for dairy cattle. Heat stress
negatively impacts a variety of dairy parameters resulting in economic losses [33]. Response to
heat stress varies with species and genetic groups within species [5,34,35]. In this study,
transcriptome of genetic groups – Crossbred and Tharparkar cattle under heat stress was
evaluated to understand their differential response to heat stress.

Animals (n=5) of both the genetic groups were exposed to a temperature of 42 °C for 7
days. Around 5^{th}- 6^{th} day, short term heat acclimation occurs [30,31]. This time point was selected
to understand the differences in systems biology to heat stress in the two genetic groups.
Initially, heat stress indicators - RR, RT, and T3 level were evaluated. RR was found to increase
in both genetic groups under heat treatment and the increase in Crossbred was found to be
significantly ($P<0.05$) different from that in Tharparkar. A positive correlation exists between
RR and heat treatment [36-38]. This increase is an attempt to dissipate excess body heat by
vaporizing more moisture in expired air or response to a greater requirement of oxygen by
tissues under heat stress. Also, the physiological response to heat stress includes reduced heat
production, which is achieved by lowering feed intake and thyroid hormone secretion [39]. T3 level
increases under heat stress [40,41]. A significant increase in T3 level in Crossbred as compared to
Tharparkar indicates an effective regulatory mechanism in modulating T3 levels in Tharparkar in
response to heat stress. The T3 triggered metabolism may be one of the reasons that increases
heat production resulting in high rectal temperature in Crossbred in comparison to Tharparkar as
was found in our study. The significant increase in RR, RT and T3 level in Crossbreed than in
Tharparkar, suggests the inability of Crossbred to cope up with heat stress in comparison to
Tharparkar.

A phenotype is defined by the changes in systems biology. Transcriptome profiling by
RNA-seq is the most common methodology to study the changes in systems biology. RNA
profiling based on next-generation sequencing enables to measure and compare gene expression
patterns[21]. The transcriptome of Tharparkar and Crossbred indicated differential response to heat
stress as evident from the DEGs, which were either distinct to both or have a difference in
expression. The number of DEGs were higher in Crossbred than in Tharparkar, suggesting a
greater dysregulation in systems biology in Crossbred. Among the dysregulated genes, the
number of upregulated genes were more than the downregulated genes in both genetic groups.
However, a contrast in expression was observed with 18.5 % of upregulated genes in Crossbred,
were downregulated in Tharparkar and 17.5% upregulated genes in Tharparkar
weredownregulated in Crossbred. Some of the genes were confirmed for their expression by
Realtime PCR. The genes selected for Realtime PCR were selected based on their role in the
major physiological processes usually associated with heat stress.

IPA revealed activation or inactivation of several pathways in both the genetic groups. It
is known that - EIF2 signalling, helps in initiation of global protein translation [42]; MAPK-
signalling pathway, induces cell proliferation [43]; androgen signalling, enhances pro-survival and
anti-apoptotic activity in cell [44]; a-Adrenergic signalling, maintains immune defence mechanism
[45] and, helps in tissue repair upon stress [46] and increases angiogenesis [47]; integrin pathway, resists
the cell against apoptosis and other environmental insults [48]; IL-3 signalling, aids in cell survival
and haematopoiesis [49]; CXCR4 signalling modulates cell survival and cell motility [50] and ;
Phospholipase C signalling aids in cell survival in stress through protein kinase C dependent
phosphorylation of BCL-2 [51]. Inactivation of these pathways except MAPK-signalling pathway
in Crossbred and activation of α-Adrenergic signalling, Androgen signalling, EIF2 signalling
and MAPK signalling in Tharparkar indicates that the systems biology in Tharparkar is moving
towards countering the effects due to heat stress.

While exploring the DEGs at a functional level we considered a knowledge-based
approach. Under this, four major physiological processes are found to be usually associated with
heat stress viz. elicitation of unfolded protein response (UPR) in cells; Ubiquitination; Induction
of apoptosis and; Imbalance in production of ROS and antioxidants (Figure 13). Heat shock and
its associated genes are involved in elicitation of unfolded protein response (UPR) in cells. Most
of the heat shock genes were found upregulated in Tharparkar and downregulated in Crossbred.
The increased HSP levels have been found positively correlated with tolerance in many
species[52,53]. HSF1, that positively regulates the transcription of *HSP70* and *HSP90* [54,55] was
found upregulated in Tharparkar and downregulated in Crossbred. Upregulation of *HSF1,*
HSP70 and HSP90 in Tharparkar and vice-versa in Crossbred corroborates to state that
Tharparkar is better equipped to counter heat stress than Crossbred. Further, to ensure that the
HSP70 in Tharparkar is maintained at an optimum level, dysregulation of *CAMK2D* and *GSK3A*
seems to act as negative feedback. CAMK2D that induces the transcription of HSP70 via HSF1
[56] was found downregulated in Tharparkar. GSK3A that inhibits the trimerization of HSF1 that is
needed for the induction of HSP70 [57] was found upregulated in Tharparkar. The decreased level
of HSP70 in Crossbred makes it inevitable that such negative feedbacks would further reduce its
level and hence, *GSK3A* was found downregulated and *CAMK*, upregulated (Figure.13).

Ubiquitination is positively correlated with heat tolerance [58,59]. Ubiquitin-Proteasome

System (UPS) regulates the levels of proteins and acts by removing the misfolded or damaged

proteins that may accumulate as a result of exposure to abiotic stress. Malfunctioning of

ubiquitin-proteasome system (UPS) could have negative consequences for protein regulation,

including loss of function [60]. In Tharparkar after heat acclimation, HSP70 tends to activate the

ubiquitination pathway to minimize the accumulation of the unfolded proteins that can't be

refolded by it [61]. This pathway activation is supported by upregulation of E2 ligases - *UBE2G1,*

UBE2S, and *UBE2H* that catalyze covalent attachment of E2 to E3 [62-65] in Tharparkar. USP7

that deubiquitinates target proteins [66,67] was found upregulated in Crossbred and downregulated

in Tharparkar. Further, a group of molecules – *VCP, SERP1*, and *CALR* that ensure the

protection of naïve proteins during their transport within the cell [68-70] were found upregulated in

Tharparkar and downregulated in Crossbred. Unlike Crossbred, Tharparkar is not only endowed

with higher expression of the scavengers of misfolded proteins but also with protectors of naïve

unfolded proteins.

Activation of apoptosis pathway is one of the major physiological processes linked with

heat stress. Among the apoptotic genes, *BCL2L11, FASLG, MLKL, CASP3, MAPK8,* and *VIM*

have been found upregulated in Crossbred and downregulated in Tharparkar under heat stress.

BCL2L11 induces apoptosis by neutralizing key molecules of pro-survival BCL2 sub-family

[71,72], FASLG   transduces the apoptotic signal into cells[73,74], CASP3 activates caspases and

executes apoptosis [75], and MAPK8, MLKL, and VIM also induce apoptosis [76,77]. PTEN

signaling pathway that drives apoptosis [78,79] was found inactivated in Tharparkar and activated in

Crossbred. This indicates a relatively higher probability of apoptosis in Crossbred than in

Tharparkar.

The ability to balance the ROS and antioxidant level, is one of the key factors that would

determine the tolerance of an individual to heat stress. The antioxidant triad of GPX, SOD, and

CAT that forms the first line of defense against reactive oxygen species [80-82], was found

upregulated in Tharparkar and downregulated in Crossbred. Additionally, genes belonging to

Peroxiredoxins - *PRDX3, PRDX5* and *PRDX6* that catalyze the reduction of hydrogen peroxide

and organic hydroperoxides [83-87],were also found upregulated in Tharparkar and were either

downregulated or not-differentially expressed in Crossbred. Higher expression of the

antioxidants in Tharparkar enables it to cope up with higher levels of free radicals generated as a
result of heat stress while Crossbred is unable to do so.

Conclusion

A contrast in expression was observed with 18.5 % of upregulated genes in Crossbred
were downregulated in Tharparkar and 17.5% upregulated genes in Tharparkar were
downregulated in Crossbred. Transcripts of molecules that stimulate heat shock response,
Ubiquitination, unfolded protein response and antioxidant level were found upregulated in
Tharparkar and downregulated in Crossbred. EIF2 Signaling that promotes protein translation
and PTEN signaling that drives apoptosis were found activated and inactivated in Tharparkar,
respectively and vice-versa in Crossbred. We found relevant molecules/genes dysregulated in
Tharparkar in the direction that counters heat stress. A proposed contrasting interplay of
molecules in both the two groups is shown in Figure 8. To the best of our knowledge this is a
comprehensive comparison between Tharparkar and Crossbred at a global level using
transcriptome analysis.

Methods

Experimental condition and Ethical Statement

The animals used for the study were from the Indian Veterinary Research Institute. The
permission to conduct the study was granted by Indian Veterinary Research Institutional Animal
Ethics Committee (IVRI-IAEC) under the Committee for Control and Supervision of
Experiments on Animals (CPCSEA), India, vide letter no 387/CPSCEA. Genetic groups -
Tharparkar (Indigenous breeds) and Vrindavani (synthetic Crossbred) were considered in this
study. Prior to experiment, the animals – 05 Tharparkar and 05 Crossbred (Vrindavani) cattle,
were acclimatized for 15 days outside the Psychometric chamber. The experiment was conducted
during October when the environmental Temperature Humidity Index (THI) was 73.0242. These
animals were exposed in Psychometric chamber at 42 °C for six hours for 7 days (THI
=78.5489). All the animals were fed with wheat straw and concentrate mixture in 60:40 ratios.
Respiration rate (RR) and rectal temperature (RT) of animals from each genetic group were
measured on 0 day (Control, n=5) before exposure to Psychometric chamber and on 7[th] day of
heat exposure (Treated, n=5). Blood samples were collected from the animals at the mentioned

time points and serum concentration of Triiodothyronine (T3) was estimated by RIA technique

using T_3 [125]I (Immunotech) as per the manufacturer's instructions.

RNA sequencing (RNA-seq**)**

PBMCs were collected from the blood samples using Ficol histopaque gradient method**.**

Total RNA from each of the collected samples (PBMCs) was isolated using the RNeasy Mini kit

(Qiagen GmbH, Germany) according to the manufacturer's protocol. The integrity and quantity

of isolated RNA were assessed on a Bioanalyzer 2100 (Agilent Technologies, Inc). The library

was prepared using NEBNext Ultra RNA Library Prep Kit for Illumina (NewEngland Biolabs

Inc.) following the manufacturer's protocol. Approximately, 100ng of RNA from each sample

was used for RNA library preparation. The quality of the libraries was assessed on Bioanalyzer.

Libraries were quantified using a Qubit 2.0 Fluorometer (Life technologies) and by qPCR.

Library (1.3ml, 1.8pM) was denatured, diluted and loaded onto a flow cell for sequencing.

cDNA library preparation and Illumina Sequencing was performed at Sandor Life Sciences Pvt.

(Hyderabad, India). Finally, the RNA-seq data were provided in FASTQ format.

Data processing

The sequenced reads were paired-end and 150bp in length. Quality control checks on raw

sequence data from each sample were performed using FastQC (Babraham Bioinformatics).

Processing of the data was performed using prinseq-lite software [88] to remove reads of low

quality (mean phred score 25) and short length (< 50) for downstream analysis. The data was

submitted to the GEO database with accession number GSE136652.

Identification of Differentially Expressed Genes (DEGs)

Bos taurus reference genome (release 94) and its associated gene transfer file (GTF) were

downloaded from Ensembl FTP genome browser [89]. The reference genome was prepared and

indexed by RNA-Seq by expectation maximization (RSEM) [90] by rsem-prepare-reference

command. Further, the clean reads obtained from filtering of raw data were aligned to the

indexed reference genome by Bowtie2 [91] to estimate gene abundance in counts by rsem-

calculate-expression command. To compare the gene expression levels among different samples,

the aligned reads were used to generate a data matrix by rsem-generate-data-matrix command.

In each genetic group, all the samples of day 7 (treated) were compared with the day 0 (Control)

for the calculation of differential gene expression by edgeR [92] package. The Ensemble IDs of the

differentially expressed genes (DEGs) were converted to the respective gene ID by g: Convert of

g: Profiler [93,94].

Functional Analysis of DEGs

Under heat stress four major physiological processes are found to be usually associated -

Induction of apoptosis [95,96]; Ubiquitination [97,98]; elicitation of unfolded protein response (UPR)

in cells [99] and ; Imbalance in production of ROS and antioxidants [100,101]. The genes involved in

these processes were retrieved from Reactome database [102]. From these genes that are involved

in the four major physiological processes, the genes that were differentially expressed in both the

genetic groups were extracted to compare and contrast their expression between genetic groups.

These genes were referred to as knowledge-based genes. Their protein-protein interaction

network was also studied in both the genetic groups.

Predicted protein-protein interaction of the knowledge-based genes

408   Protein-protein interaction (PPI) network among the knowledge-based DEGs that were

409   found common to both Tharparkar and Crossbred, was retrieved using interactions available in

410   the String database [103]. The degree was calculated using igraph package (https://cran.r-

411   project.org/web/packages/igraph/ index.html). The PPI network was then visualized using

412   Cytoscape software V. 3.7 [104]

Ingenuity Pathway Analysis (IPA) Analysis

QIAGEN's IPA (QIAGEN, Redwood City, USA) [105] is used to quickly visualize and

understand complex omics data and perform insightful data analysis and interpretation by

placing experimental results within the context of biological systems. Here, IPA was used to

analyze the identified DEGs of Crossbred and Tharparkar.The list of DEGs from each genetic

group was used to identify the canonical pathways and the most significant biological processes

against Ingenuity Pathways Knowledge Base (IKB). Core analysis for each dataset was

performed to know activated (Z score > 2) or inactivated (Z score < -2) canonical pathways.

Upstream regulators - Transcription factors and microRNAs were also identified.

Validation of reference genes identified

423   Genes - *BCL2L11, FASLG, CASP3, CAT, SOD1, GSK3A, CALR, HSF1, APC,* and *GPX3*

424 were selected based on their role in heat stress and qRT-PCR was performed on Applied

425 Biosystems 7500 Fast system. *GAPDH* was taken as the internal control. Each of the samples

426 was run in triplicates and relative expression of each gene was calculated using the

427 $2^{-\Delta\Delta CT}$ method with control as the calibrator [106].

Statistical Analysis

429   Respiration rate, Rectal temperature and T3 level were compared using student's *t*-test in

430 JMP9 (SAS Institute Inc., Cary, USA) to test the significance of the difference between the

431 control (0 day) and treated (7th day). This comparison was done within and between genetic

432 groups. Differences within/between groups were considered significant at $P \leq 0.05$.

Declarations

Ethics approval and consent to participate

435 The permission to conduct the study was granted by Indian Veterinary Research Institutional

436 Animal Ethics Committee (IVRI-IAEC) under the Committee for Control and Supervision of

437 Experiments on Animals (CPCSEA), India, vide letter no 387/CPSCEA.

Consent for publication

439 Not applicable.

Availability of data and materials

441 The data was submitted to the GEO database with accession number GSE136652

Competing interests

443 None of the authors had a conflict of interest to declare

Funding

445 This study was supported by "National Innovations in Climate Resilient Agriculture (NICRA) -

446 Identification of unique factors in indigenous livestock making them resilient to climate change

447 in relation to diseases" for the funds to carryout Sampling, Wet Lab experiments and for the
448 procurement of License of IPA.

Authors' contributions

450 AKT and RG conceived and designed the research. SG, SmS, AS,AV, VV , PK , ShS and GS
451 conducted the wet lab work. RINK, ARS, NH, WAM, MRP, SK , AP and RG analyzed the data.
452 RINK, ARS, MRP, RG , AS and GS helped in manuscript drafting and editing. AKT and RG
453 proofread the manuscript

Acknowledgements

455 We also thank Department of Biotechnology, Govt of India for providing fellowship and
456 contingency for RK (DBT Fellow No. DBT/2017/IVRI/768), ARS (DBT Fellow No.
457 DBT/2014/IVRI/170) and WAM (DBT Fellow No. DBT/2017/IVRI/769).

References

Mehla, K. *et al.* Genome-wide analysis of the heat stress response in Zebu (Sahiwal) cattle. *Gene* **533**, 500-507 (2014).

Armstrong, D. V. Heat stress interaction with shade and cooling. *J Dairy Sci* **77**, 2044-2050, doi:10.3168/jds.S0022-0302(94)77149-6 (1994).

Rojas-Downing, M. M., Nejadhashemi, A. P., Harrigan, T. & Woznicki, S. A. Climate change and livestock: Impacts, adaptation, and mitigation. *Climate Risk Management* **16**, 145-163 (2017).

Hahn, G. Dynamic responses of cattle to thermal heat loads. *Journal of animal science* **77**, 10-20 (1999).

Dash, S. *et al.* Effect of heat stress on reproductive performances of dairy cattle and buffaloes: A review. *Veterinary world* **9**, 235 (2016).

Renaudeau, D. *et al.* Adaptation to hot climate and strategies to alleviate heat stress in livestock production. *Animal* **6**, 707-728 (2012).

Berman, A. Invited review: Are adaptations present to support dairy cattle productivity in warm climates? *Journal of Dairy Science* **94**, 2147-2158 (2011).

Rashamol, V. P. *et al.* Physiological adaptability of livestock to heat stress: an updated review. *J Anim Behav Biometeorol* **6**, 62-71 (2018).

Davis, S. R., Spelman, R. J. & Littlejohn, M. D. BREEDING AND GENETICS SYMPOSIUM:Breeding heat tolerant dairy cattle: the case for introgression of the "slick" prolactin receptor variant into dairy breeds. *J Anim Sci* **95**, 1788-1800, doi:10.2527/jas.2016.0956 (2017).

Mcmanus, C. M., Louvandini, H., Paim, T. P., Silva, F. C. P. e. & Bernal, F. E. M. Factors affecting heat tolerance in crossbred cattle in central Brazil. *Ciência Animal Brasileira* **15**, 152-158 (2014).

Carabano, M. J., Ramon, M., Menendez-Buxadera, A., Molina, A. & Diaz, C. Selecting for heat tolerance. *Anim Front* **9**, 62-68, doi:10.1093/af/vfy033 (2019).

Aritonang, S., Yuniati, R., Abinawanto, Imron, M. & Bowolaksono, A. in *AIP Conference Proceedings.* 030098 (AIP Publishing).

13 Cardoso, C. *et al.* Physiological and thermographic response to heat stress in zebu cattle. *Livestock Science* **182**, 83-92 (2015).
14 Valente, É. E. L. *et al.* Intake, physiological parameters and behavior of Angus and Nellore bulls subjected to heat stress. *Semina: Ciências Agrárias* **36**, 4565-4574 (2015).
15 Sodhi, M., Mukesh, M., Prakash, B., Ahlawat, S. & Sobti, R. Microsatellite DNA typing for assessment of genetic variability in Tharparkar breed of Indian zebu (Bos indicus) cattle, a major breed of Rajasthan. *Journal of genetics* **85**, 165-170 (2006).
16 Rachagani, S., Gupta, I. D., Gupta, N. & Gupta, S. Genotyping of β-Lactoglobulin gene by PCR-RFLP in Sahiwal and Tharparkar cattle breeds. *BMC genetics* **7**, 31 (2006).
17 Choudhary, S. *et al.* Tharparkar: The pride of desert. *Journal of entomology and zoology studies* **6**, 1915-1919 (2018).
18 Ahmad, S. F. *et al.* Revelation of genomic breed composition in a crossbred cattle of India with the help of Bovine50K BeadChip. *Genomics* **112**, 1531-1535, doi:10.1016/j.ygeno.2019.08.025 (2020).
19 Salleh, M. *et al.* RNA-Seq transcriptomics and pathway analyses reveal potential regulatory genes and molecular mechanisms in high-and low-residual feed intake in Nordic dairy cattle. *BMC genomics* **18**, 258 (2017).
20 Cánovas, A., Rincon, G., Islas-Trejo, A., Wickramasinghe, S. & Medrano, J. F. SNP discovery in the bovine milk transcriptome using RNA-Seq technology. *Mammalian genome* **21**, 592-598 (2010).
21 Mortazavi, A., Williams, B. A., McCue, K., Schaeffer, L. & Wold, B. Mapping and quantifying mammalian transcriptomes by RNA-Seq. *Nature methods* **5**, 621 (2008).
22 Wang, Z., Gerstein, M. & Snyder, M. RNA-Seq: a revolutionary tool for transcriptomics. *Nature reviews genetics* **10**, 57 (2009).
23 Wani, S. A. *et al.* Contrasting gene expression profiles of monocytes and lymphocytes from Peste-des-petits-ruminants virus infected goats. *Frontiers in Immunology* **10** (2019).
24 Manjunath, S. *et al.* Comparative and temporal transcriptome analysis of peste des petits ruminants virus infected goat peripheral blood mononuclear cells. *Virus research* **229**, 28-40 (2017).
25 Li, P., Piao, Y., Shon, H. S. & Ryu, K. H. Comparing the normalization methods for the differential analysis of Illumina high-throughput RNA-Seq data. *BMC bioinformatics* **16**, 347 (2015).
26 Khanduri, A. *et al.* Dysregulated miRNAome and proteome of PPRV infected goat PBMCs reveal a coordinated immune response. *Frontiers in immunology* **9**, 2631 (2018).
27 Pandey, A. *et al.* Modulation of host miRNAs transcriptome in lung and spleen of Peste des Petits ruminants virus infected sheep and goats. *Frontiers in microbiology* **8**, 1146 (2017).
28 O'Loughlin, A. *et al.* Transcriptomic analysis of the stress response to weaning at housing in bovine leukocytes using RNA-seq technology. *BMC genomics* **13**, 250 (2012).
29 Salama, A. *et al.* Different levels of response to heat stress in dairy goats. *Small Ruminant Research* **121**, 73-79 (2014).
30 Dangi, S. S. *et al.* Modulatory effect of betaine on expression dynamics of HSPs during heat stress acclimation in goat (Capra hircus). *Gene* **575**, 543-550 (2016).
31 Garrett, A. *et al.* Short-term heat acclimation is effective and may be enhanced rather than impaired by dehydration. *American journal of human biology* **26**, 311-320 (2014).
32 He, X. & Zhang, J. Why do hubs tend to be essential in protein networks? *PLoS Genet* **2**, e88, doi:10.1371/journal.pgen.0020088 (2006).
33 Polsky, L. & von Keyserlingk, M. A. Invited review: Effects of heat stress on dairy cattle welfare. *Journal of dairy science* **100**, 8645-8657 (2017).

34 Ross, J. *et al.* Physiological consequences of heat stress in pigs. *Animal Production Science* **55**,
1381-1390 (2015).
35 Mayorga, E. J., Renaudeau, D., Ramirez, B. C., Ross, J. W. & Baumgard, L. H. Heat stress
adaptations in pigs. *Animal Frontiers* **9**, 54-61 (2018).
36 Blackshaw, J. K. & Blackshaw, A. Heat stress in cattle and the effect of shade on production and
behaviour: a review. *Australian Journal of Experimental Agriculture* **34**, 285-295 (1994).
37 Beede, D. & Collier, R. Potential nutritional strategies for intensively managed cattle during
thermal stress. *Journal of Animal Science* **62**, 543-554 (1986).
38 Ganaie, A. *et al.* Biochemical and physiological changes during thermal stress in bovines: A
review. *Iranian Journal of Applied Animal Science* **3**, 423-430 (2013).
39 Hansen, P. Physiological and cellular adaptations of zebu cattle to thermal stress. *Animal
reproduction science* **82**, 349-360 (2004).
40 Magdub, A., Johnson, H. & Belyea, R. Effect of environmental heat and dietary fiber on thyroid
physiology of lactating cows. *Journal of Dairy Science* **65**, 2323-2331 (1982).
41 Williamson, R., Misson, B. & Davison, T. The effect of exposure to 40 on the heat production and
the serum concentrations of triiodothyronine, thyroxine, and corticosterone in immature
domestic fowl. *General and comparative endocrinology* **60**, 178-186 (1985).
42 Rajesh, K. *et al.* Phosphorylation of the translation initiation factor eIF2alpha at serine 51
determines the cell fate decisions of Akt in response to oxidative stress. *Cell Death Dis* **6**, e1591,
doi:10.1038/cddis.2014.554 (2015).
43 Zhang, W. & Liu, H. T. MAPK signal pathways in the regulation of cell proliferation in mammalian
cells. *Cell Res* **12**, 9-18, doi:10.1038/sj.cr.7290105 (2002).
44 Shiota, M., Yokomizo, A. & Naito, S. Pro-survival and anti-apoptotic properties of androgen
receptor signaling by oxidative stress promote treatment resistance in prostate cancer. *Endocr
Relat Cancer* **19**, R243-253, doi:10.1530/ERC-12-0232 (2012).
45 Lacoste, A., De Cian, M. C., Cueff, A. & Poulet, S. A. Noradrenaline and alpha-adrenergic signaling
induce the hsp70 gene promoter in mollusc immune cells. *J Cell Sci* **114**, 3557-3564 (2001).
46 Xian, D. *et al.* Emerging Roles of Redox-Mediated Angiogenesis and Oxidative Stress in
Dermatoses. *Oxid Med Cell Longev* **2019**, 2304018, doi:10.1155/2019/2304018 (2019).
47 Zhao, X. *et al.* Overexpression of Cardiomyocyte alpha1A-Adrenergic Receptors Attenuates
Postinfarct Remodeling by Inducing Angiogenesis Through Heterocellular Signaling. *Arterioscler
Thromb Vasc Biol* **35**, 2451-2459, doi:10.1161/ATVBAHA.115.305919 (2015).
48 Stupack, D. G. & Cheresh, D. A. Get a ligand, get a life: integrins, signaling and cell survival. *J Cell
Sci* **115**, 3729-3738, doi:10.1242/jcs.00071 (2002).
49 Reddy, E. P., Korapati, A., Chaturvedi, P. & Rane, S. IL-3 signaling and the role of Src kinases, JAKs
and STATs: a covert liaison unveiled. *Oncogene* **19**, 2532-2547, doi:10.1038/sj.onc.1203594
(2000).
50 Young, B., Purcell, C., Kuang, Y. Q., Charette, N. & Dupre, D. J. Superoxide Dismutase 1
Regulation of CXCR4-Mediated Signaling in Prostate Cancer Cells is Dependent on Cellular
Oxidative State. *Cell Physiol Biochem* **37**, 2071-2084, doi:10.1159/000438566 (2015).
51 Bai, X. C. *et al.* Phospholipase C-gamma1 is required for survival in heat stress: involvement of
protein kinase C-dependent Bcl-2 phosphorylation. *J Biochem* **131**, 207-212,
doi:10.1093/oxfordjournals.jbchem.a003089 (2002).
52 Parsell, D. & Lindquist, S. The function of heat-shock proteins in stress tolerance: degradation
and reactivation of damaged proteins. *Annual review of genetics* **27**, 437-496 (1993).
53 Rout, P., Kaushik, R. & Ramachandran, N. Differential expression pattern of heat shock protein
70 gene in tissues and heat stress phenotypes in goats during peak heat stress period. *Cell Stress
and Chaperones* **21**, 645-651 (2016).

54 Westerheide, S. D., Anckar, J., Stevens, S. M., Sistonen, L. & Morimoto, R. I. Stress-inducible
regulation of heat shock factor 1 by the deacetylase SIRT1. *Science* **323**, 1063-1066 (2009).

55 Prodromou, C. Mechanisms of Hsp90 regulation. *Biochemical Journal* **473**, 2439-2452 (2016).

56 Holmberg, C. I. *et al.* Phosphorylation of serine 230 promotes inducible transcriptional activity of
heat shock factor 1. *The EMBO journal* **20**, 3800-3810 (2001).

57 Wang, X., Grammatikakis, N., Siganou, A. & Calderwood, S. K. Regulation of molecular
chaperone gene transcription involves the serine phosphorylation, 14-3-3ε binding, and
cytoplasmic sequestration of heat shock factor 1. *Molecular and cellular biology* **23**, 6013-6026
(2003).

58 Ryu, K. Y. *et al.* The mouse polyubiquitin gene UbC is essential for fetal liver development, cell-
cycle progression and stress tolerance. *The EMBO journal* **26**, 2693-2706 (2007).

59 Lyzenga, W. J. & Stone, S. L. Abiotic stress tolerance mediated by protein ubiquitination. *Journal*
of experimental botany **63**, 599-616 (2011).

60 Sulistio, Y. A. & Heese, K. The ubiquitin-proteasome system and molecular chaperone
deregulation in Alzheimer's disease. *Molecular neurobiology* **53**, 905-931 (2016).

61 Fernández-Fernández, M. R., Gragera, M., Ochoa-Ibarrola, L., Quintana-Gallardo, L. & Valpuesta,
J. M. Hsp70–a master regulator in protein degradation. *FEBS letters* **591**, 2648-2660 (2017).

62 Pabarcus, M. K. *et al.* CYP3A4 ubiquitination by gp78 (the tumor autocrine motility factor
receptor, AMFR) and CHIP E3 ligases. *Archives of biochemistry and biophysics* **483**, 66-74 (2009).

63 David, Y., Ziv, T., Admon, A. & Navon, A. The E2 ubiquitin-conjugating enzymes direct
polyubiquitination to preferred lysines. *Journal of Biological Chemistry* **285**, 8595-8604 (2010).

64 Sheng, Y. *et al.* A human ubiquitin conjugating enzyme (E2)-HECT E3 ligase structure-function
screen. *Molecular & Cellular Proteomics* **11**, 329-341 (2012).

65 Kaiser, P. *et al.* A human ubiquitin-conjugating enzyme homologous to yeast UBC8. *Journal of*
Biological Chemistry **269**, 8797-8802 (1994).

66 Li, M. *et al.* Deubiquitination of p53 by HAUSP is an important pathway for p53 stabilization.
Nature **416**, 648 (2002).

67 Zhao, Y. *et al.* Noncanonical regulation of alkylation damage resistance by the OTUD4
deubiquitinase. *The EMBO journal* **34**, 1687-1703 (2015).

68 Kadowaki, H. *et al.* Pre-emptive quality control protects the ER from protein overload via the
proximity of ERAD components and SRP. *Cell reports* **13**, 944-956 (2015).

69 Hori, O. *et al.* Deletion of SERP1/RAMP4, a component of the endoplasmic reticulum (ER)
translocation sites, leads to ER stress. *Molecular and cellular biology* **26**, 4257-4267 (2006).

70 Nauseef, W. M., McCormick, S. J. & Clark, R. A. Calreticulin functions as a molecular chaperone
in the biosynthesis of myeloperoxidase. *Journal of Biological Chemistry* **270**, 4741-4747 (1995).

71 O'Connor, L. *et al.* Bim: a novel member of the Bcl-2 family that promotes apoptosis. *The EMBO*
journal **17**, 384-395 (1998).

72 Marani, M., Tenev, T., Hancock, D., Downward, J. & Lemoine, N. R. Identification of novel
isoforms of the BH3 domain protein Bim which directly activate Bax to trigger apoptosis.
Molecular and cellular biology **22**, 3577-3589 (2002).

73 Muneta, Y., Shimoji, Y., Inumaru, S. & Mori, Y. Molecular cloning, characterization, and
expression of porcine Fas ligand (CD95 Ligand). *Journal of Interferon & Cytokine Research* **21**,
305-312 (2001).

74 Ruiz-García, R. *et al.* Decreased activation-induced cell death by EBV-transformed B-cells from a
patient with autoimmune lymphoproliferative syndrome caused by a novel FASLG mutation.
Pediatric research **78**, 603 (2015).

75 Nicholson, D. W. *et al.* Identification and inhibition of the ICE/CED-3 protease necessary for
mammalian apoptosis. *Nature* **376**, 37 (1995).
76 Dai, R., Frejtag, W., He, B., Zhang, Y. & Mivechi, N. F. c-Jun NH2-terminal kinase targeting and
phosphorylation of heat shock factor-1 suppress its transcriptional activity. *Journal of Biological*
Chemistry **275**, 18210-18218 (2000).
77 Cao, W.-X. *et al.* MLKL mediates apoptosis via a mutual regulation with PERK/eIF2α pathway in
response to reactive oxygen species generation. *Apoptosis* **23**, 521-531 (2018).
78 Lu, X.-X. *et al.* PTEN inhibits cell proliferation, promotes cell apoptosis, and induces cell cycle
arrest via downregulating the PI3K/AKT/hTERT pathway in lung adenocarcinoma A549 cells.
BioMed research international **2016** (2016).
79 Chen, C.-Y., Chen, J., He, L. & Stiles, B. L. PTEN: tumor suppressor and metabolic regulator.
Frontiers in endocrinology **9** (2018).
80 Lee, Y. S. *et al.* Dysregulation of adipose glutathione peroxidase 3 in obesity contributes to local
and systemic oxidative stress. *Molecular Endocrinology* **22**, 2176-2189 (2008).
81 Bierl, C., Voetsch, B., Jin, R. C., Handy, D. E. & Loscalzo, J. Determinants of human plasma
glutathione peroxidase (GPx-3) expression. *Journal of Biological Chemistry* **279**, 26839-26845
(2004).
82 Ighodaro, O. & Akinloye, O. First line defence antioxidants-superoxide dismutase (SOD), catalase
(CAT) and glutathione peroxidase (GPX): Their fundamental role in the entire antioxidant
defence grid. *Alexandria Journal of Medicine* **54**, 287-293 (2018).
83 Tsuji, K., Copeland, N., Jenkins, N. & Obinata, M. Mammalian antioxidant protein complements
alkylhydroperoxide reductase (ahpC) mutation in Escherichia coli. *Biochemical Journal* **307**, 377-
381 (1995).
84 Yamashita, H. *et al.* Characterization of human and murine PMP20 peroxisomal proteins that
exhibit antioxidant activity in vitro. *Journal of Biological Chemistry* **274**, 29897-29904 (1999).
85 Knoops, B. *et al.* Cloning and characterization of AOEB166, a novel mammalian antioxidant
enzyme of the peroxiredoxin family. *Journal of Biological Chemistry* **274**, 30451-30458 (1999).
86 Kang, S. W., Baines, I. C. & Rhee, S. G. Characterization of a mammalian peroxiredoxin that
contains one conserved cysteine. *Journal of Biological Chemistry* **273**, 6303-6311 (1998).
87 Chen, J.-W., Dodia, C., Feinstein, S. I., Jain, M. K. & Fisher, A. B. 1-Cys peroxiredoxin, a
bifunctional enzyme with glutathione peroxidase and phospholipase A2 activities. *Journal of*
Biological Chemistry **275**, 28421-28427 (2000).
88 Schmieder, R. & Edwards, R. Quality control and preprocessing of metagenomic datasets.
Bioinformatics **27**, 863-864 (2011).
89 Cunningham, F. *et al.* Ensembl 2015. *Nucleic acids research* **43**, D662-D669 (2014).
90 Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. *BMC bioinformatics* **12**, 323 (2011).
91 Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nature methods* **9**,
357 (2012).
92 Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. *Bioinformatics* **26**, 139-140 (2010).
93 Reimand, J., Kull, M., Peterson, H., Hansen, J. & Vilo, J. g: Profiler—a web-based toolset for
functional profiling of gene lists from large-scale experiments. *Nucleic acids research* **35**, W193-
W200 (2007).
94 Reimand, J. *et al.* g: Profiler—a web server for functional interpretation of gene lists (2016
update). *Nucleic acids research* **44**, W83-W89 (2016).

95   Gu, Z. *et al.* Heat stress induced apoptosis is triggered by transcription-independent p53, Ca 2+
dyshomeostasis and the subsequent Bax mitochondrial translocation. *Scientific reports* **5**, 11497
(2015).
96   Li, L. *et al.* Heat stress induces apoptosis through a Ca2+-mediated mitochondrial apoptotic
pathway in human umbilical vein endothelial cells. *PloS one* **9**, e111083 (2014).
97   Kelly, S. M., VanSlyke, J. K. & Musil, L. S. Regulation of Ubiquitin-Proteasome System–mediated
Degradation by Cytosolic Stress. *Molecular biology of the cell* **18**, 4279-4291 (2007).
98   Flick, K. & Kaiser, P. in *Seminars in cell & developmental biology.* 515-522 (Elsevier).
99   Homma, T. & Fujii, J. Heat stress promotes the down-regulation of IRE1α in cells: An atypical
modulation of the UPR pathway. *Experimental cell research* **349**, 128-138 (2016).
100   Belhadj Slimen, I. *et al.* Reactive oxygen species, heat stress and oxidative-induced
mitochondrial damage. A review. *International journal of hyperthermia* **30**, 513-523 (2014).
101   Yang, L., Tan, G.-Y., Fu, Y.-Q., Feng, J.-H. & Zhang, M.-H. Effects of acute heat stress and
subsequent stress removal on function of hepatic mitochondrial respiration, ROS production
and lipid peroxidation in broiler chickens. *Comparative Biochemistry and Physiology Part C:*
Toxicology & Pharmacology **151**, 204-208 (2010).
102   Fabregat, A. *et al.* The reactome pathway knowledgebase. *Nucleic acids research* **46**, D649-D655
(2017).
103   Szklarczyk, D. *et al.* STRING v11: protein–protein association networks with increased coverage,
supporting functional discovery in genome-wide experimental datasets. *Nucleic acids research*
47, D607-D613 (2018).
104   Shannon, P. *et al.* Cytoscape: a software environment for integrated models of biomolecular
interaction networks. *Genome research* **13**, 2498-2504 (2003).
105   Krämer, A., Green, J., Pollard Jr, J. & Tugendreich, S. Causal analysis approaches in ingenuity
pathway analysis. *Bioinformatics* **30**, 523-530 (2013).
106   Schmittgen, T. D. & Livak, K. J. Analyzing real-time PCR data by the comparative C T method.
Nature protocols **3**, 1101 (2008).

Legends

Figure 1: Overview of the work done : Two genetic groups (Tharparkar and Crossbred) of cattle

were exposed to a temperature of 42 °C for 7 days. Heat stress indicators - Respiration rate (RR),

Rectal temperature and T3 level before exposure to heat (0day – control group) and at 7[th] day of

exposure (treated) were measured to evaluate heat stress. At these time points, RNA was isolated

from PBMCs for high throughput sequencing. Transcriptome analysis was done to identify

differentially expressed genes (DEGs) under heat treatment in both genetic groups. Genes

involved in physiological processes (heat stress response, apoptosis, ubiquitination, unfolded

protein response and antioxidant level) that are commonly associated with heat stress were

compared between the two genetic groups. Further, functional annotation of DEGs was done
using IPA.

Figure 2: Respiration rate, Rectal Temperature and T3 level measured at 0 day (control) and 7
day post-heat exposure (treated) in Crossbred and Tharparkar (n=5) Levels sharing the same
superscript are not significantly ($P > 0.05$) different from each other.

Figure 3: Expression of DEGs in Crossbred and Tharparkar under heat stress: (a) Venn diagrams
showing unique/common DEGs between Crossbred and Tharparkar (b) Number of upregulated
and downregulated in both genetic groups (c) Contrast in the expression of common DEGs

Figure 4: Contrast in the expression of genes involved in heat stress response, apoptosis,
ubiquitination, unfolded protein response and balance in the production of ROS and antioxidants
between two genetic groups.

Figure 5 : Canonical pathways activated/inactivated in (a) Crossbred (b) Tharparkar under heat
stress generated in the core analysis of Ingenuity pathway analysis tool. Orange color pathways
are activated ($Z > 2$) and blue color pathways are inactivated ($Z < -2$). Height of the bar graphs
indicates -log (p-value) and line graph showing the ratio of list genes found in each pathway over
the total number of genes in that pathway.

Figure 6: Canonical pathways generated in Ingenuity Pathway Analysis of Oncostatin M
signaling pathway of DEGs in (A) Crossbred, (B) Tharparkar. Genes that were upregulated are
shown in red and downregulated in green. The intensity of red and green corresponds to an
increase and decrease, respectively, in Log2 fold change. Genes in grey were not significantly
dysregulated and those in white are not present in the dataset but have been incorporated in the
network through the relationship with other molecules by IPA.

Figure 7: Canonical pathways generated in Ingenuity Pathway Analysis of PTEN signaling
pathway of DEGs in (A) Crossbred, (B) Tharparkar. Genes that were upregulated are shown in
red and downregulated in green. The intensity of red and green corresponds to an increase and
decrease, respectively, in Log2 fold change. Genes in grey were not significantly dysregulated
and those in white are not present in the dataset but have been incorporated in the network
through the relationship with other molecules by IPA.

Figure 8: Predicted interplay of molecules that is underway during heat stress in both

groups : Heat stress causes unfolding of native proteins. HSP70 acts as a chaperone to facilitate

refolding to restore the structure of unfolded proteins. Under normal condition, HSP70 is bound

to HSF1 thereby preventing HSF1 to promote transcription of HSP70. Under heat stress ATP

binds to the HSP70 and HSF1 complex to release HSF1, promoting the binding of the unfolded

protein to HSP70 and ATP. CAMK2D that induces the transcription of HSP70 via HSF1 was

found downregulated in Tharparkar. GSK3A that inhibits the trimerization of HSF1 that is

needed for the induction of HSP70 expression was found upregulated in Tharparkar. The

decreased level of HSP70 in Crossbred makes it inevitable that such negative feedbacks would

further reduce its level and GSK3A was found downregulated and CAMK2D, upregulated.

Further, in Tharparkar, HSP70 tends to activate ubiquitination pathway to decrease the

accumulation of unfolded proteins that can't be refolded by it. This pathway activation is

supported by upregulation of E3 ligases (UBE2G1, UBE2S, and UBE2H) in Tharparkar.

However, the E3 ligase in Crossbred was found downregulated. With HSP70 being upregulated

and having cytoprotection activity, Tharparkar shows the decline in apoptosis as compared to

Crossbred. This is supported by downregulation of BCL2L11, FASLG, MLKL, CASP3,

MAPK8 and VIM in Tharparkar and vice-versa. Besides, higher expression of the antioxidants

(SOD, CAT, GPX) in Tharparkar enables it to cope up with higher levels of free radicals

generated as a result of heat stress while Crossbred is unable to do so. Green arrow indicates

downregulation and Maroon arrow indicates upregulation.

Crossbred_DEGs　　　　　　　　　**Tharparkar_DEGs**

(a)

(c)

(b)

(a)

(b)

(a)　　　　　　　　　　**(b)**

(a)

(b)