Systems Biology under heat stress in Indian Cattle

Raja Ishaq Nabi Khan a†, Amit Ranjan Sahu a†, Waseem Akram Malla a, Manas Ranjan Praharaj b, Neelima Hosamani b† , Shakti Kumar b, Smita Gupta a, Shweta Sharma a , Archana Saxena a , Anshul Varshney a , Pragya Singh a , Vinay Verma c , Puneet Kumarc, Gyanendra Singh c , Aruna Pandey a, Shikha Saxena a, Ravi Kumar Gandham b# , Ashok Kumar Tiwari d#

a Division of Veterinary Biotechnology, Indian Veterinary Research Institute, Bareilly, India
b Computational Biology and Genomics, National Institute of Animal Biotechnology, Hyderabad, India
c Division of Physiology and Climatology,   Indian Veterinary Research Institute, Bareilly, India
d Division of Biological Standardization,   Indian Veterinary Research Institute, Bareilly, India

†Equal Contributors

Corresponding Authors: Ravi Kumar Gandham, ravigandham@niab.org.in
                          Ashok Kumar Tiwari, aktiwari71d@gmail.com

Abstract

Transcriptome profiling of Vrindavani and Tharparkar cattle revealed that more numbers of genes dysregulated in Vrindavani than in Tharparkar. A contrast in gene expression was observed with 18.5 % of upregulated genes in Vrindavani were downregulated in Tharparkar and 17.5% upregulated genes in Tharparkar were downregulated in Vrindavani. Functional annotation of genes differentially expressed in Tharparkar and Vrindavani revealed that the systems biology in Tharparkar is moving towards counteracting the effects due to heat stress. Unlike Vrindavani, Tharparkar is not only endowed with higher expression of the scavengers (*UBE2G1, UBE2S*, and *UBE2H*) of misfolded proteins but also with protectors (*VCP, Serp1*, and *CALR*) of naïve unfolded proteins. Further, higher expression of the antioxidants in Tharparkar enables it to cope up with higher levels of free radicals generated as a result of heat stress. In this study we found relevant genes dysregulated in Tharparkar in the direction that can counter heat stress.

Keywords: Heat Stress; Transcriptome; Tharparkar; Vrindavani

Introduction

Cattle being homoeothermic modulate their internal body temperature in sync to environmental temperature by equilibrating the amount of heat produced within the body and dissipating it to the ambient environment. The stress that arises due to disproportionate

thermodynamic behavior between cattle and its surrounding environment is termed as heat stress [1]. Environmental induced hyperthermic stress lowers feed intake, which in turn reduces growth, milk production and reproductive efficiency, thereby negatively affecting the economics of livestock keepers [2-4]. Heat stress has been associated with reduced fertility through its deleterious impact on oocyte maturation and early embryonic development [5]. Increased morbidity and mortality was observed in animals due to immune depressive effect of heat stress [6].

India has a wide variety of indigenous cattle breeds distributed throughout its agro-climatic zones. These are known for their natural tolerance to tropical heat [7,8]. To meet the growing demand for milk and to combine the heat tolerance and tick resistance of zebu with the productivity of temperate dairy breeds [9] several crossbreeding programs were taken up in India. Every state had its own crossbreeding policy, which is agro-climatic and breed-specific. Though the zebu crosses with European breeds produced more milk than zebu, they were found not withstanding heat/solar radiation [10]. Crossbreds are susceptible to tropical diseases and require a constant input of good management conditions [8]. Antagonism exists between heat tolerance and milk productivity [11]. The adaptive capacity to heat stress varies between species and genetic groups within species. Among various adaptive mechanisms, physiological adaptability seems to be the primary step in cattle. Sahiwal cows better regulate body temperature in response to heat stress than Karan Fries [8]. It was observed that Ongole cattle rely on the respiration rate to maintain thermal balance, while, Bali cattle rely on rectal temperature [12]. In Brazil, Sindhi and Girolando breeds showed better physiological response to thermal stress than Gir cattle [13]. Increase in respiration rate was reported in Nellore breed when exposed to heat load [14].

In India, Tharparkar is one among the best dairy breeds. It is adapted to the Indian states of Punjab and Haryana. [15,16]. It is considered to be the hardiest, disease resistant, heat tolerant and tick resistant indigenous cattle breed of the country [17]. This breed has also been used in several crossbreeding programs. Currently, percentage of purebreds is exceptionally low in India (Department of Animal Husbandry & Dairying, Govt. of India). Most of the farmer's in India have Crossbreds and the percentage of exotic inheritance in these Crossbreds is unknown. Vrindavani, a Crossbred (synthetic), with 27% of Indigenous blood and 73% of exotic inheritance [18] is a representation of the kind of admixture that prevails in Indian cattle. Therefore, comparing Tharparkar with Vrindavani may establish the lost importance of Indigenous cattle

and would further emphasize the need of conserving our indigenous purebreds because of
advantageous traits like Heat tolerance.

Studies explaining the difference between the genetic groups (Crossbreds and Indigenous
cattle) have been done mainly to address the physiological responses vis – a – vis heat stress and
very few studies at the genomic level have been taken up [19,20].Also, there is an increasing need to
develop methods by combining the knowledge from -omics technologies to identify heat tolerant
animals [11]. Transcriptome profiling / RNA-Sequencing (RNA-seq) is a high throughput omics
approach to measure relative global changes in the transcripts under specific condition(s) [21-23] to
study the systems biology behind a phenotype [23,24]. RNA - seq allows for analysis of
transcriptome in an unbiased way, with, a tremendous dynamic detection range (>8,000 fold),
and low background signals [25]. It has been used as an investigating tool in understanding disease
pathogenesis [26,27] and differential physiological response to various biotic and abiotic factors
[28,29].

In this study, Tharparkar and Vrindavani cattle were subjected to heat stress and blood
samples were collected on 0^{th} day and 7^{th} day, as it is known that short term acclimation occurs
around 5 - 6 days [30,31]. The transcriptome of 7^{th} day was compared and with 0^{th} day in both the
genetic groups to understand their differential response to heat stress.

Results

Physiological Parameters

86   The overview of the analysis is given in Figure 1. Respiration rate (RR), rectal
87 temperature (RT) and T3 level increased significantly (p<0.05) on 7^{th} - day post heat stress in
88 both the genetic groups (n=5) (Figure 2). However, the increase in RR, RT and T3 level, was
89 found significantly (P<0.05) higher in Vrindavani than in Tharparkar.

Comparison of DEGs of Vrindavani and Tharparkar under heat stress

The differentially expressed genes for each genetic group were obtained on comparing
the 0 day and 7^{th} day RNA-seq data using EdgeR after obtaining the gene counts from RSEM.
Under heat stress, global expression profiles of Vrindavani and Tharparkar were identified with
6042 and 4718 differentially expressed genes (DEGs), respectively (Supplementary Table 1).
Among these, 3481 DEGs were found common between the two genetic groups, while 2561 and

1238 DEGs were uniquely found in Vrindavani and Tharparkar, respectively (Figure 3a).

Additionally, 3132 and 2924 genes were upregulated and downregulated in Vrindavani,

respectively, while 2367 and 2358 genes were upregulated and downregulated in Tharparkar,

respectively (Figure 3b). On comparison of upregulated and downregulated genes, 724 and 1416

genes were found uniquely upregulated and 514 and 1145 genes were found uniquely

downregulated in Tharparkar and Vrindavani, respectively. The comparison also revealed that

17.5% of upregulated genes (1278) in Tharparkar were downregulated in Vrindavani and 18.5%

downregulated genes (1344) in Tharparkar were upregulated in Vrindavani. However, the

number of common upregulated and downregulated genes in both the genetic groups were 357

(4.9%) and 498 (6.8%), respectively (Figure 3c).

Functional annotation of DEGs in Reactome

DEGs in Tharparkar and Vrindavani

On functional annotation of all the DEGs (6042 and 4718 in Vrindavani and Tharparkar,

respectively), the commonly enriched significant pathways in both genetic groups included,

neutrophil degranulation, L13a-mediated translational silencing of ceruloplasmin expression,

Formation of a pool of free 40S subunits, ubiquitination and proteasome degradation, SRP-

dependent cotranslational protein targeting to membrane and nonsense-mediated decay and

selenocysteine synthesis. Among the most enriched pathways, peptide chain elongation, gap-

filling DNA repair synthesis and ligation in TC-NER were the pathways significantly enriched in

Tharparkar and not in Vrindavani. The pathways - SCF-beta-TrCP mediated degradation of

Emi1, Autodegradation of Cdh1 by Cdh1:APC/C and Regulation of RAS by GAPs were the

pathways significantly enriched in Vrindavani and not in Tharparkar.

Common DEGs in Tharparkar and Vrindavani

On functional annotation of common the DEGs (3481), the enriched significant pathways

included, neutrophil degranulation, L13a-mediated translational silencing of ceruloplasmin

expression, formation of a pool of free 40S subunits, ubiquitination and proteasome degradation,

SRP-dependent cotranslational protein targeting to membrane, RNA Polymerase II Transcription

Termination, Peptide chain elongation, nonsense-mediated decay and selenocysteine synthesis.

On functionally annotation of 1278 common genes that were upregulated in Tharparkar and

downregulated in Vrindavani, the enriched pathways included all the pathways as mentioned for

the common except for ubiquitination & proteasome degradation. In addition, cellular response
to stress was found significantly enriched these genes upregulated in Tharparkar and
downregulated in Vrindavani. Further, on assessing the 1344 genes that were upregulated in
Vrindavani and downregulated in Tharparkar, RUNX1 regulates transcription of genes involved
in BCR signaling, p75NTR recruits signaling complexes, chromatin modifying enzymes,
chromatin organization, death receptor signaling, resolution of D-loop structures through
Holliday junction intermediates and TP53 regulates transcription of DNA repair genes pathways
were significantly enriched.

Unique DEGs in Tharparkar

On functional annotation of unique DEGs (1238) expressed only in Tharparkar, the enriched
significant pathways included, nucleotide excision repair, mitotic G1 phase and G1/S transition,
DNA replication pre-initiation, cell cycle, synthesis of DNA, translation, DNA repair, activation
of the pre-replicative complex, cell cycle checkpoints and inhibition of replication initiation of
damaged DNA by RB1/E2F.

Unique DEGs in Vrindavani

On functional annotation of unique DEGs (2561) expressed only in Vrindavani, the enriched
significant pathways included, unfolded protein response (UPR), loss of proteins required for
interphase microtubule organization from the centrosome, amplification of signal from
unattached kinetochores via a MAD2 inhibitory signal, mitotic spindle checkpoint, organelle
biogenesis and maintenance and loss of Nlp from mitotic centrosomes.

Functional annotation of DEGs in IPA

Canonical pathway analysis

Canonical pathway analysis by Ingenuity Pathway Analysis (IPA) revealed contrast in signaling
pathways in Vrindavani and Tharparkar. Canonical pathways associated with Vrindavani and
Tharparkar are represented in Figure 4a and 4b. In Vrindavani, Oncostatin M Signaling,
Phospholipase C Signaling, EIF2 Signaling, Integrin Signaling, IL-3 Signaling, and CXCR4
Signaling were found to be highly inactivated (Z – score > 2.0) and PTEN signaling was found
to be highly activated (Z – score < 2.0). In Tharparkar, EIF2 Signaling, Androgen Signaling,
Oncostatin M Signaling, α-Adrenergic Signaling, BMP signaling pathway, and UVC-Induced

MAPK Signaling were found to be highly activated and PTEN signaling was found to be
inactivated. The canonical pathway Oncostatin M Signaling and EIF2 Signaling were found to
have the highest ratio of genes involved vis-a-vis the genes in the database in Vrindavani and
Tharparkar, respectively.

While carrying out comparative analysis through IPA, Calcium-induced T Lymphocyte
Apoptosis, BMP signaling pathway, UVC-Induced MAPK Signaling, Regulation of Cellular
Mechanics by Calpain Protease, fMLP Signaling in Neutrophils, Melatonin Signaling, and
Leukocyte Extravasation Signaling, were found inactivated in Vrindavani and activated in
Tharparkar (Supplementary Figure 1). Genes involved in Oncostatin M Signaling- Growth
factor receptor-bound protein 2 (*GRB2*), GTPase HRas *(HRAS)*, Janus kinase 1 *(JAK1)*, Janus
kinase 3 *(JAK3)*, Mitogen-activated protein kinase kinase 1 *(MAP2K1)*, Mitogen-activated
protein kinase 1 *(MAPK1)*, Oncostatin-M *(OSM)*, Ras-related protein Rap-1b *(RAP1B)*, Ras-
related protein Rap-2a *(RAP2A)*, Signal transducer and activator of transcription 1-alpha/beta
(STAT1), Signal transducer and activator of transcription 5B *(STAT5B)*, Non-receptor tyrosine-
protein kinase *(TYK2)*, and Ras-related protein *(RRAS)* were found downregulated in Vrindavani
and upregulated in Tharparkar (Figure 5a, b). While the key genes involved in PTEN Signaling
pathway – Fas Ligand (*FASLG*), member of RAS oncogene family *(RAP2A)*, Bcl-2-like protein
11 *(BIM)*,Caspase-3 *(CASP3)* and microspherule protein 1 (*MSP58)* were found upregulated in
Vrindavani and downregulated in Tharparkar as well (Figure 6a, b).

Variation in microRNAs and Transcription factors

IPA, on evaluating the differentially expression genes predicts miRNAs and
Transcription Factors (upstream regulators). In Vrindavani, 111 miRNAs were found to be
inactivated and 37 activated. In Tharparkar, 205 miRNAs were found to be inactivated and 272
activated. Among them, 52 microRNAs were found common between the two genetic groups.
Most of the common miRNAs were found activated in Vrindavani and inactivated in Tharparkar
(Supplementary Figure 2). miR-4779, miR-4651, miR-1207-5p, miR-6967-5p and miR-504-3p
are the top 5 miRNAs that were activated in Vrindavani and inactivated in Tharparkar.

Various Transcription factors were found to regulate the expression of the identified
DEGs. Transcription factors, 19 in Tharparkar (11 activated and 8 inactivated) and 26 in
Vrindavani (8 activated and 18 inactivated) were identified in IPA that regulate the expression of

DEGs. Among them, *PAX5, MTA3, MYC, PROX1* and *SMAD7* in Vrindavani and, *HMGA1,*

MAF, MAX NOTCH22 and *NCOR1* in Tharparkar are the top 5 upregulated and activated TFs.

On comparing the TFs of Tharparkar and Vrindavani, it was found that *BHLHE40, HMGA1,*

HMGB1, IKZF1, and *TCF7* were found to be common. *BHLHE40, HMGA1,* and *TCF7* were

found to be activated in Tharparkar and inactivated in Vrindavani and it was vice - versa with

HMGB1 and *IKZF1* (Supplementary Figure 3)

Disease and Functions

In the diseases and functions category, on evaluating all the DEGs in both the genetic

groups, survival of the organism, ubiquitination, ubiquitination of protein and repair of DNA

were found activated in Vrindavani and were found either inactivated/ not-activated in

Tharparkar. Similarly, homeostasis, development of hematopoietic cells, leukopoiesis was found

relatively activated in Tharparkar in comparison to Vrindavani. Further, translation, Expression

of protein, translation of protein and degranulation were found inactivated in Vrindavani.

On correlating the results of disease and function with the results of reactome four major

processes that were considered to be associated with heat stress viz. elicitation of unfolded

protein response (UPR) in cells; Induction of apoptosis; Ubiquitination and; Imbalance in

production of ROS and antioxidants. Heat shock genes and its associated genes are involved in

elicitation of unfolded protein response (UPR) in cells. Heat shock genes have been found

dysregulated under heat stress in both the genetic groups. Most of the genes encoding Heat

shock proteins (HSPs) - Heat shock 70 kDa protein 4   *(HSPA4),* Heat shock cognate 71 kDa

protein *(HSPB8),* Heat shock 70 kDa protein 1A *(HSPA1A) ,*Heat shock cognate 71 kDa protein

(HSPA8), Heat shock protein HSP 90-beta *(HSP90AB1)* and Heat shock protein HSP 90-alpha

(*HSP90AA1*) and heat shock protein regulating factors- Heat shock factor 1 *(HSF1)* and

Eukaryotic Translation Elongation Factor 1 Alpha 1 *(EEF1A1)* have been found to be

downregulated/not-differentially expressed in Vrindavani but upregulated in Tharparkar.

However, Calcium/Calmodulin Dependent Protein Kinase II Delta *(CAMK2D) that* is involved

in the regulation of expression of heat shock genes was upregulated in Vrindavani and

downregulated in Tharparkar.

Among the apoptotic genes, genes encoding Bcl-2-like protein 11 *(BCL2L11) ,* Tumor

necrosis factor ligand superfamily member 6   *(FASLG) ,*   TIR domain-containing adapter

molecule 2 *(TICAM2)* , Toll-like receptor 4 *(TLR4),* Adenomatous polyposis coli protein *(APC),* Caspase-3 *(CASP3),* Mitogen-activated protein kinase 8 *(MAPK8),* Mixed lineage kinase domain-like protein *(MLKL),* Late endosomal/lysosomal adaptor and MAPK and MTOR activator 5 *(XIP),* Vimentin *(VIM),* and High mobility group protein B2 *(HMGB2)* were found to be upregulated in Vrindavani and downregulated in Tharparkar. The number of upregulated genes involved in achieving the balance of ROS production and antioxidants, were found to be more in Tharparkar than in Vrindavani. Among these, genes encoding Glutathione peroxidase 3 (*GPX3),* Nudix Hydrolase 2 *(NUDT2),* Catalase *(CAT),* Cytochrome c *(CYCS),* Copper chaperone for superoxide dismutase *(CCS),* Peroxiredoxin-5 *(PRDX5),* Peroxiredoxin-6 *(PRDX6),* Peroxiredoxin-1 *(PRDX1),* Superoxide dismutase *(SOD1),* and Cytochrome b-245 heavy chain *(CYBB*) were found either downregulated/not-differentially expressed in Vrindavani and upregulated in Tharparkar.   More number of genes involved in Ubiquitination were differentially expressed in Vrindavani than in the Tharparkar. Genes encoding Ubiquitin-conjugating enzyme E2 G1 *(UBE2G1),* Ubiquitin-conjugating enzyme E2 *(UBE2S),* Ubiquitin-conjugating enzyme E2 H *(UBE2H),* Ubiquitin A-52 residue ribosomal protein fusion product 1 *(UBA52),* and Ubiquitin-activating enzyme E1 *(UBA1)* have been found downregulated/not-differentially expressed in Vrindavani and upregulated in Tharparkar. However, Valosin-containing protein *(VCP),* RING finger protein 40 *(RNF40)*, and Ubiquitin-conjugating enzyme E2 L3 *(UBE2L3)* have been found downregulated in Vrindavani but not-differentially expressed in Tharparkar. Among the genes involved in Unfolded Protein folding response (UPR), genes encoding Membrane-bound transcription factor site-1 protease *(MBTPS1),* Cyclic AMP-responsive element-binding protein 3-like protein 1 *(CREB3L1),* Stress-associated endoplasmic reticulum protein 1 *(SERP1),* Glycogen synthase kinase-3 alpha *(GSK3A),* Eukaryotic translation initiation factor 2 subunit 3 *(EIF2S3),* Calreticulin *(CALR),* and Stress-associated endoplasmic reticulum protein 1 (*SERP1)* have been found downregulated in Vrindavani and upregulated in Tharparkar (Figure 7).

Protein - protein interaction (PPI) network

Among the genes involved in the 4 major processes considered, protein - protein interaction (PPI) network revealed functional importance of HSP70 (HSPA8 and HSPA1A) and ubiquitin (UBB, UBA52), in coordinating genes involved in heat stress. Out of the total 246 genes

identified from the reactome database among these processes, 177 and 194 genes were found to

be differentially expressed in Tharparkar and Vrindavani, respectively. Among these 126 genes

were found to be commonly differentially expressed in Tharparkar and Vrindavani. PPI network

for these common genes between Tharparkar and Vrindavani was constructed (Supplementary

Figure 4). In PPI networks, hubs define the functional and structural importance of a network.

The genes, which act as hubs in PPI networks were found to be *UBB, UBA52, HSPA8*, and

HSPA1A (Supplementary Figure 4). Among the 4 hubs, *UBB* was downregulated in both genetic

groups and the rest were downregulated in Vrindavani and upregulated in Tharparkar.

A change in the expression of the hub protein will have a larger effect than change in

expression of non-hub proteins [32]. Therefore, UBB, UBA52, HSPA8, and HSPA1A are taken to

be critical for coordinating the changes in systems biology under heat stress. The hubs HSPA8

and HSPA1A are connected to genes that are associated with regulation of stress viz.

nucleoporins genes - *NUP188, NUP155, NUP210 & NUP214*; BAG family molecular chaperone

regulators - *BAG1, BAG3 & BAG4*; Heat Shock Protein Family A - *HSPA5, HSPA4, HSPA12B*

& HSPA9; DnaJ Heat Shock Protein Family i.e. HSP40 - *DNAJA1, DNAJC2 & DNAJB6*; Heat

shock factor - *HSF1*; Ubiquitin - *UBB & UBA52* and; Sirtuin - *SIRT1*. The hubs - UBB and

UBA52 are connected to molecules of different proteasome subunits viz. ▯ type subunits -

PSMA1 & PSMA2; β type subunits - *PSMB4 & PSMB8*; ATPase subunits - *PSMC2 & PSMC5*

and non-ATPase subunits - *PSMD2 & PSMD13*. These hubs were also found connected to

ubiquitin specific peptidases - *USP9X* and *USP7* and Ubiquitin-conjugating enzyme - *UBE2B,*

UBE2G1, UBE2Z, UBE2H, UBE2J2, UBE2S & UBE2D2.

Real-time validation

Six genes (*HSF1, SOD1, CALR, GSK3A, CAT & GPX3*) that were upregulated in

Tharparkar but downregulated/not expressed in Vrindavani and four genes (*CASP3, FASLG,*

BCL2L11 & APC) that were upregulated in Vrindavani but downregulated in Tharparkar were

considered for Real time PCR based on their role in heat stress. The expression of genes was in

concordance with the RNA- Seq results (Supplementary Figure 5 and Supplementary table 2).

Discussion

Heat stress is a natural phenomenon that affects domestic animals in tropical, sub-tropical

and often in temperate regions of the world during summer months. Heat and humidity during

the summer months combine to make an uncomfortable environment for dairy cattle. Heat stress

negatively impacts a variety of dairy parameters resulting in economic losses [33]. Response to

heat stress varies with species and genetic groups within species [5,34,35]. In this study,

transcriptome of genetic groups – Vrindavani and Tharparkar cattle under heat stress was

evaluated to understand their differential response to heat stress.

Animals (n=5) of both the genetic groups were exposed to a temperature of 42 °C for 7

days. Around 5[th]- 6[th] day, short term heat acclimation occurs [30,31]. This time point was selected

to understand the differences in systems biology to heat stress in the two genetic groups.

Initially, heat stress indicators - RR, RT, and T3 level were evaluated. RR was found to increase

in both genetic groups under heat treatment and the increase in Vrindavani was found to be

significantly (P<0.05) different from that in Tharparkar. A positive correlation exists between

RR and heat treatment [36-38]. This increase is an attempt to dissipate excess body heat by

vaporizing more moisture in expired air or response to a greater requirement of oxygen by

tissues under heat stress. Also, the physiological response to heat stress includes reduced heat

production, which is achieved by lowering feed intake and thyroid hormone secretion [39]. T3 level

increases under heat stress [40,41]. A significant increase in T3 level in Vrindavani as compared to

Tharparkar indicates an effective regulatory mechanism in modulating T3 levels in Tharparkar in

response to heat stress. The T3 triggered metabolism may be one of the reasons that increases

heat production resulting in high rectal temperature in Vrindavani in comparison to Tharparkar

as was found in our study. The significant increase in RR, RT and T3 level in Crossbreed than in

Tharparkar, suggests the inability of Vrindavani to cope up with heat stress in comparison to

Tharparkar.

A phenotype is defined by the changes in systems biology. Transcriptome profiling by

RNA-seq is the most common methodology to study the changes in systems biology. RNA

profiling based on next-generation sequencing enables to measure and compare gene expression

patterns[21]. The transcriptome of Tharparkar and Vrindavani indicated differential response to

heat stress as evident from the DEGs, which were either distinct to both or have a difference in

expression. The number of DEGs were higher in Vrindavani than in Tharparkar, suggesting a

greater dysregulation in systems biology in Vrindavani. Among the dysregulated genes, the

number of upregulated genes were more than the downregulated genes in both genetic groups.

However, a contrast in expression was observed with 18.5 % of upregulated genes in Vrindavani,

were downregulated in Tharparkar and 17.5% upregulated genes in Tharparkar were downregulated in Vrindavani.

The differentially expressed genes in each genetic group were functionally annotated using both reactome and IPA. In the reactome, the enriched pathways were identified and in IPA, based on the expression of the DEGs the activated/inactivated/not-activated pathways were identified. In reactome, on functionally annotating the DEGs of each of the genetic groups, key pathways related to stress – neutrophil degranulation [42], L13a-mediated translational silencing of ceruloplasmin expression [43], ubiquitination and proteasome degradation [44], SRP-dependent cotranslational protein targeting to membrane [45], Nonsense-Mediated Decay [46] and selenocysteine synthesis [47] were enriched in both the genetic groups. Among the common genes, the genes upregulated in Tharparkar but downregulated in Vrindavani showed enrichment towards cellular response to stress besides the above-mentioned pathways. This indicated that Tharparkar must be responding to withstand heat stress. Further, on evaluating the common genes that were upregulated in Vrindavani and downregulated in Tharparkar, pathways - chromatin modifying enzymes, chromatin organization and death receptor signaling indicated towards apoptosis and DNA repair in Vrindavani. These findings were further corroborated with the pathway enrichment analysis of unique genes in both the genetic groups. IPA analysis of disease and biological function showed repair of DNA and ubiquitination, activated in Vrindavani and senescence and degranulation of cells inactivated in Tharparkar. The results from reactome and IPA suggested Tharparkar may be more resilient than Vrindavani.

IPA revealed activation or inactivation of several pathways in both the genetic groups. It is known that - EIF2 signalling, helps in initiation of global protein translation [48]; MAPK-signalling pathway, induces cell proliferation [49]; androgen signalling, enhances pro-survival and anti-apoptotic activity in cell [50]; a-Adrenergic signalling, maintains immune defence mechanism [51] and, helps in tissue repair upon stress [52] and increases angiogenesis [53]; integrin pathway, resists the cell against apoptosis and other environmental insults [54]; IL-3 signalling, aids in cell survival and haematopoiesis [55]; CXCR4 signalling modulates cell survival and cell motility [56] and ; Phospholipase C signalling aids in cell survival in stress through protein kinase C dependent phosphorylation of BCL-2 [57]. Inactivation of these pathways except MAPK-signalling pathway in Vrindavani and activation of α-Adrenergic signalling, Androgen signalling, EIF2 signalling

and MAPK signalling in Tharparkar indicates that the systems biology in Tharparkar is moving
towards countering the effects due to heat stress.

On correlating the reactome data with the IPA four major physiological processes -
elicitation of unfolded protein response (UPR) in cells; Ubiquitination; Induction of apoptosis
and; Imbalance in production of ROS and antioxidants were considered for further evaluation
(Figure 8). Heat shock and its associated genes are involved in elicitation of unfolded protein
response (UPR) in cells. Most of the heat shock genes were found upregulated in Tharparkar and
downregulated in Vrindavani. The increased HSP levels have been found positively correlated
with tolerance in many species[58,59]. HSF1, that positively regulates the transcription of *HSP70*
and *HSP90* [60,61] was found upregulated in Tharparkar and downregulated in Vrindavani.
Upregulation of *HSF1, HSP70 and HSP90* in Tharparkar and vice-versa in Vrindavani
corroborates to state that Tharparkar is better equipped to counter heat stress than Vrindavani.
Further, to ensure that the HSP70 in Tharparkar is maintained at an optimum level, dysregulation
of *CAMK2D* and *GSK3A* seems to act as negative feedback. CAMK2D that induces the
transcription of HSP70 via HSF1 [62] was found downregulated in Tharparkar. GSK3A that
inhibits the trimerization of HSF1 that is needed for the induction of HSP70 [63] was found
upregulated in Tharparkar. The decreased level of HSP70 in Vrindavani makes it inevitable that
such negative feedbacks would further reduce its level and hence, *GSK3A* was found
downregulated and *CAMK*, upregulated (Figure 8).

Ubiquitination is positively correlated with heat tolerance [64,65]. Ubiquitin-Proteasome
System (UPS) regulates the levels of proteins and acts by removing the misfolded or damaged
proteins that may accumulate as a result of exposure to abiotic stress. Malfunctioning of
ubiquitin-proteasome system (UPS) could have negative consequences for protein regulation,
including loss of function [66]. In Tharparkar after heat acclimation, HSP70 tends to activate the
ubiquitination pathway to minimize the accumulation of the unfolded proteins that can't be
refolded by it [67]. This pathway activation is supported by upregulation of E2 ligases - *UBE2G1,*
UBE2S, and *UBE2H* that catalyze covalent attachment of E2 to E3 [68-71] in Tharparkar. USP7
that deubiquitinates target proteins [72,73] was found upregulated in Vrindavani and downregulated
in Tharparkar. Further, a group of molecules – *VCP, SERP1*, and *CALR* that ensure the
protection of naïve proteins during their transport within the cell [74-76] were found upregulated in

Tharparkar and downregulated in Vrindavani. Unlike Vrindavani, Tharparkar is not only endowed with higher expression of the scavengers of misfolded proteins but also with protectors of naïve unfolded proteins.

Activation of apoptosis pathway is one of the major physiological processes linked with heat stress. Among the apoptotic genes, *BCL2L11, FASLG, MLKL, CASP3, MAPK8,* and *VIM* have been found upregulated in Vrindavani and downregulated in Tharparkar under heat stress. BCL2L11 induces apoptosis by neutralizing key molecules of pro-survival BCL2 sub-family [77,78], FASLG transduces the apoptotic signal into cells[79,80], CASP3 activates caspases and executes apoptosis [81], and MAPK8, MLKL, and VIM also induce apoptosis [82,83]. PTEN signaling pathway that drives apoptosis [84,85] was found inactivated in Tharparkar and activated in Vrindavani. This indicates a relatively higher probability of apoptosis in Vrindavani than in Tharparkar.

The ability to balance the ROS and antioxidant level, is one of the key factors that would determine the tolerance of an individual to heat stress. The antioxidant triad of GPX, SOD, and CAT that forms the first line of defense against reactive oxygen species [86-88], was found upregulated in Tharparkar and downregulated in Vrindavani. Additionally, genes belonging to Peroxiredoxins - *PRDX3, PRDX5* and *PRDX6* that catalyze the reduction of hydrogen peroxide and organic hydroperoxides [89-93], were also found upregulated in Tharparkar and were either downregulated or not-differentially expressed in Vrindavani. Higher expression of the antioxidants in Tharparkar enables it to cope up with higher levels of free radicals generated as a result of heat stress while Vrindavani is unable to do so.

Conclusion

A contrast in expression was observed with 18.5 % of upregulated genes in Vrindavani were downregulated in Tharparkar and 17.5% upregulated genes in Tharparkar were downregulated in Vrindavani. Transcripts of molecules that stimulate heat shock response, Ubiquitination, unfolded protein response and antioxidant level were found upregulated in Tharparkar and downregulated in Vrindavani. EIF2 Signaling that promotes protein translation and PTEN signaling that drives apoptosis were found activated and inactivated in Tharparkar, respectively and vice-versa in Vrindavani. We found relevant molecules/genes dysregulated in Tharparkar in the direction that counters heat stress. A proposed contrasting interplay of

molecules in both the two groups is shown in Figure 8. To the best of our knowledge this is a

comprehensive comparison between Tharparkar and Vrindavani at a global level using

transcriptome analysis.

Methods

Experimental condition and Ethical Statement

The animals used for the study were from the Indian Veterinary Research Institute. The

permission to conduct the study was granted by Indian Veterinary Research Institutional Animal

Ethics Committee (IVRI-IAEC) under the Committee for Control and Supervision of

Experiments on Animals (CPCSEA), India, vide letter no 387/CPSCEA. Genetic groups -

Tharparkar (Indigenous breeds) and Vrindavani (synthetic Crossbred) were considered in this

study. Prior to experiment, the animals – 05 Tharparkar and 05 Vrindavani cattle, were

acclimatized for 15 days outside the Psychometric chamber. The experiment was conducted

during October when the environmental Temperature Humidity Index (THI) was 73.0242. These

animals were exposed in Psychometric chamber at 42 °C for six hours for 7 days (THI

=78.5489). All the animals were fed with wheat straw and concentrate mixture in 60:40 ratios.

Respiration rate (RR) and rectal temperature (RT) of animals from each genetic group were

measured on 0 day (Control, n=5) before exposure to Psychometric chamber and on 7[th] day of

heat exposure (Treated, n=5). Blood samples were collected from the animals at the mentioned

time points and serum concentration of Triiodothyronine (T3) was estimated by RIA technique

using T_3 [125]I (Immunotech) as per the manufacturer's instructions.

RNA sequencing (RNA-seq**)**

PBMCs were collected from the blood samples using Ficol histopaque gradient method**.**

Total RNA from each of the collected samples (PBMCs) was isolated using the RNeasy Mini kit

(Qiagen GmbH, Germany) according to the manufacturer's protocol. The integrity and quantity

of isolated RNA were assessed on a Bioanalyzer 2100 (Agilent Technologies, Inc). The library

was prepared using NEBNext Ultra RNA Library Prep Kit for Illumina (NewEngland Biolabs

Inc.) following the manufacturer's protocol. Approximately, 100ng of RNA from each sample

was used for RNA library preparation. The quality of the libraries was assessed on Bioanalyzer.

Libraries were quantified using a Qubit 2.0 Fluorometer (Life technologies) and by qPCR.

Library (1.3ml, 1.8pM) was denatured, diluted and loaded onto a flow cell for sequencing.

cDNA library preparation and Illumina Sequencing was performed at Sandor Life Sciences Pvt.
(Hyderabad, India). Finally, the RNA-seq data were provided in FASTQ format.

Data processing

The sequenced reads were paired-end and 150bp in length. Quality control checks on raw
sequence data from each sample were performed using FastQC (Babraham Bioinformatics).
Processing of the data was performed using prinseq-lite software [94] to remove reads of low
quality (mean phred score 25) and short length (< 50) for downstream analysis. The data was
submitted to the GEO database with accession number GSE136652.

Identification of Differentially Expressed Genes (DEGs)

Bos taurus reference genome (release 94) and its associated gene transfer file (GTF) were
downloaded from Ensembl FTP genome browser [95]. The reference genome was prepared and
indexed by RNA-Seq by expectation maximization (RSEM) [96] by rsem-prepare-reference
command. Further, the clean reads obtained from filtering of raw data were aligned to the
indexed reference genome by Bowtie2 [97] to estimate gene abundance in counts by rsem-
calculate-expression command. To compare the gene expression levels among different samples,
the aligned reads were used to generate a data matrix by rsem-generate-data-matrix command.
In each genetic group, all the samples of day 7 (treated) were compared with the day 0 (Control)
for the calculation of differential gene expression by edgeR [98] package. The Ensemble IDs of the
differentially expressed genes (DEGs) were converted to the respective gene ID by g: Convert of
g: Profiler [99,100].

Functional Analysis of DEGs - Reactome and Ingenuity pathway analysis (IPA)

The differentially expressed genes in both the genetic groups were functionally annotated
in Reactome [101] and IPA (Ingenuity Pathway Analysis). Reactome provides bioinformatics tools
for visualisation, interpretation and analysis of pathway knowledge to support basic research.
Reactome analysis was done for all DEGs in both genetic groups, common DEGs and for unique
DEGs for each genetic group. For common DEGs, analysis was also done for contrasting genes
(genes upregulated in Tharparkar but downregulated in Vrindavani and the vice-versa).

QIAGEN's IPA (QIAGEN, Redwood City, USA) [102] is used to quickly visualize and
understand complex omics data and perform insightful data analysis and interpretation by

placing experimental results within the context of biological systems. Here, IPA was used to

analyze the identified DEGs of Vrindavani and Tharparkar. The list of DEGs from each genetic

group was used to identify the canonical pathways and the most significant biological processes

against Ingenuity Pathways Knowledge Base (IKB). Core analysis for each dataset was

performed to know activated (Z score > 2) or inactivated (Z score < −2) canonical pathways.

Upstream regulators - Transcription factors and microRNAs were identified. Significant

Diseases and functions were also identified.

The results obtained from Reactome and IPA were correlated to narrow down to

processes - Induction of apoptosis, Ubiquitination, elicitation of unfolded protein response

(UPR) in cells and Imbalance in production of ROS and antioxidants. The unique genes involved

in all related pathways to the processes mentioned above were extracted from the Reactome

analysis and were found to be 246 in number. From among these genes the number of

differentially expressed genes in Tharparkar and Vrindavani, and the common genes between the

genetic groups were extracted.

Predicted protein-protein interaction network

Protein interaction network (interactome) analysis provides an effective way to

understand the interrelationships between genes [103]. Protein-protein interactions (PPI) among the

common genes involved in the processes mentioned, were retrieved using String database [104].

The degree was calculated using igraph package (https://cran.r-project.org/web/packages/igraph/

index.html). The PPI network was then visualized using Cytoscape software V. 3.7 [105] using

DEGs and degree value from igraph.

Validation of reference genes identified

The quantity of data generated from RNA sequencing is large and therefore it is

important to validate differential expression by real-time RT-PCR. Genes - *BCL2L11, FASLG,*

CASP3, CAT, SOD1, GSK3A, CALR, HSF1, APC, and *GPX3* were selected based on their role in

heat stress and qRT-PCR was performed on Applied Biosystems 7500 Fast system. *GAPDH* was

taken as the internal control. Each of the samples was run in triplicates and relative expression of

each gene was calculated using the $2^{-\Delta\Delta CT}$ method with control as the calibrator [106].

Statistical Analysis

Respiration rate, Rectal temperature and T3 level were compared using student's *t*-test in
JMP9 (SAS Institute Inc., Cary, USA) to test the significance of the difference between the
control (0 day) and treated (7th day). This comparison was done within and between genetic
groups. Differences within/between groups were considered significant at $P \leq 0.05$.

Declarations

Ethics approval and consent to participate

490   The permission to conduct the study was granted by Indian Veterinary Research Institutional
491   Animal Ethics Committee (IVRI-IAEC) under the Committee for Control and Supervision of
492   Experiments on Animals (CPCSEA), India, vide letter no 387/CPSCEA.

Consent for publication

494   Not applicable.

Availability of data and materials

496   The data was submitted to the GEO database with accession number GSE136652

Competing interests

498   None of the authors had a conflict of interest to declare

Funding

500   This study was supported by "National Innovations in Climate Resilient Agriculture (NICRA) -
501   Identification of unique factors in indigenous livestock making them resilient to climate change
502   in relation to diseases" for the funds to carryout Sampling, Wet Lab experiments and for the
503   procurement of License of IPA.

Authors' contributions

505   AKT and RG conceived and designed the research. SG, SmS, AS,AV, VV , PK , ShS and GS
506   conducted the wet lab work. RINK, ARS, NH, WAM, MRP, SK , AP and RG analyzed the data.

507 RINK, ARS, MRP, RG , AS and GS helped in manuscript drafting and editing. AKT and RG

508 proofread the manuscript

Acknowledgements

510 We also thank Department of Biotechnology, Govt of India for providing fellowship and

511 contingency for RK (DBT Fellow No. DBT/2017/IVRI/768), ARS (DBT Fellow No.

512 DBT/2014/IVRI/170) and WAM (DBT Fellow No. DBT/2017/IVRI/769).

References

1   Mehla, K. *et al.* Genome-wide analysis of the heat stress response in Zebu (Sahiwal) cattle. *Gene*
533, 500-507 (2014).
2   Armstrong, D. V. Heat stress interaction with shade and cooling. *J. Dairy Sci.* **77**, 2044-2050,
doi:10.3168/jds.S0022-0302(94)77149-6 (1994).
3   Rojas-Downing, M. M., Nejadhashemi, A. P., Harrigan, T. & Woznicki, S. A. Climate change and
livestock: Impacts, adaptation, and mitigation. *Climate Risk Management* **16**, 145-163 (2017).
4   Hahn, G. Dynamic responses of cattle to thermal heat loads. *J. Anim. Sci.* **77**, 10-20 (1999).
5   Dash, S. *et al.* Effect of heat stress on reproductive performances of dairy cattle and buffaloes: A
review. *Veterinary world* **9**, 235 (2016).
6   Renaudeau, D. *et al.* Adaptation to hot climate and strategies to alleviate heat stress in livestock
production. *Animal* **6**, 707-728 (2012).
7   Berman, A. Invited review: Are adaptations present to support dairy cattle productivity in warm
climates? *J. Dairy Sci.* **94**, 2147-2158 (2011).
8   Rashamol, V. P. *et al.* Physiological adaptability of livestock to heat stress: an updated review. *J*
Anim Behav Biometeorol **6**, 62-71 (2018).
9   Davis, S. R., Spelman, R. J. & Littlejohn, M. D. BREEDING AND GENETICS SYMPOSIUM:Breeding
heat tolerant dairy cattle: the case for introgression of the "slick" prolactin receptor variant into
dairy breeds. *J Anim Sci* **95**, 1788-1800, doi:10.2527/jas.2016.0956 (2017).
10   Mcmanus, C. M., Louvandini, H., Paim, T. P., Silva, F. C. P. e. & Bernal, F. E. M. Factors affecting
heat tolerance in crossbred cattle in central Brazil. *Ciência Animal Brasileira* **15**, 152-158 (2014).
11   Carabano, M. J., Ramon, M., Menendez-Buxadera, A., Molina, A. & Diaz, C. Selecting for heat
tolerance. *Anim Front* **9**, 62-68, doi:10.1093/af/vfy033 (2019).
12   Aritonang, S., Yuniati, R., Abinawanto, Imron, M. & Bowolaksono, A. in *AIP Conference*
Proceedings. 030098 (AIP Publishing).
13   Cardoso, C. *et al.* Physiological and thermographic response to heat stress in zebu cattle.
Livestock Science **182**, 83-92 (2015).
14   Valente, É. E. L. *et al.* Intake, physiological parameters and behavior of Angus and Nellore bulls
subjected to heat stress. *Semina: Ciências Agrárias* **36**, 4565-4574 (2015).
15   Sodhi, M., Mukesh, M., Prakash, B., Ahlawat, S. & Sobti, R. Microsatellite DNA typing for
assessment of genetic variability in Tharparkar breed of Indian zebu (Bos indicus) cattle, a major
breed of Rajasthan. *Journal of genetics* **85**, 165-170 (2006).
16   Rachagani, S., Gupta, I. D., Gupta, N. & Gupta, S. Genotyping of β-Lactoglobulin gene by PCR-
RFLP in Sahiwal and Tharparkar cattle breeds. *BMC Genet.* **7**, 31 (2006).

17 Choudhary, S. *et al.* Tharparkar: The pride of desert. *Journal of entomology and zoology studies*
6, 1915-1919 (2018).

18 Ahmad, S. F. *et al.* Revelation of genomic breed composition in a crossbred cattle of India with
the help of Bovine50K BeadChip. *Genomics* **112**, 1531-1535, doi:10.1016/j.ygeno.2019.08.025
(2020).

19 Salleh, M. *et al.* RNA-Seq transcriptomics and pathway analyses reveal potential regulatory
genes and molecular mechanisms in high-and low-residual feed intake in Nordic dairy cattle.
BMC Genomics **18**, 258 (2017).

20 Cánovas, A., Rincon, G., Islas-Trejo, A., Wickramasinghe, S. & Medrano, J. F. SNP discovery in the
bovine milk transcriptome using RNA-Seq technology. *Mamm. Genome* **21**, 592-598 (2010).

21 Mortazavi, A., Williams, B. A., McCue, K., Schaeffer, L. & Wold, B. Mapping and quantifying
mammalian transcriptomes by RNA-Seq. *Nat. Methods* **5**, 621 (2008).

22 Wang, Z., Gerstein, M. & Snyder, M. RNA-Seq: a revolutionary tool for transcriptomics. *Nature*
reviews genetics **10**, 57 (2009).

23 Wani, S. A. *et al.* Contrasting gene expression profiles of monocytes and lymphocytes from
Peste-des-petits-ruminants virus infected goats. *Front. Immunol.* **10** (2019).

24 Manjunath, S. *et al.* Comparative and temporal transcriptome analysis of peste des petits
ruminants virus infected goat peripheral blood mononuclear cells. *Virus Res.* **229**, 28-40 (2017).

25 Li, P., Piao, Y., Shon, H. S. & Ryu, K. H. Comparing the normalization methods for the differential
analysis of Illumina high-throughput RNA-Seq data. *BMC Bioinformatics* **16**, 347 (2015).

26 Khanduri, A. *et al.* Dysregulated miRNAome and proteome of PPRV infected goat PBMCs reveal a
coordinated immune response. *Front. Immunol.* **9**, 2631 (2018).

27 Pandey, A. *et al.* Modulation of host miRNAs transcriptome in lung and spleen of Peste des
Petits ruminants virus infected sheep and goats. *Front. Microbiol.* **8**, 1146 (2017).

28 O'Loughlin, A. *et al.* Transcriptomic analysis of the stress response to weaning at housing in
bovine leukocytes using RNA-seq technology. *BMC Genomics* **13**, 250 (2012).

29 Salama, A. *et al.* Different levels of response to heat stress in dairy goats. *Small Rumin. Res.* **121**,
73-79 (2014).

30 Dangi, S. S. *et al.* Modulatory effect of betaine on expression dynamics of HSPs during heat
stress acclimation in goat (Capra hircus). *Gene* **575**, 543-550 (2016).

31 Garrett, A. *et al.* Short-term heat acclimation is effective and may be enhanced rather than
impaired by dehydration. *Am. J. Hum. Biol.* **26**, 311-320 (2014).

32 He, X. & Zhang, J. Why do hubs tend to be essential in protein networks? *PLoS Genet* **2**, e88,
doi:10.1371/journal.pgen.0020088 (2006).

33 Polsky, L. & von Keyserlingk, M. A. Invited review: Effects of heat stress on dairy cattle welfare. *J.*
Dairy Sci. **100**, 8645-8657 (2017).

34 Ross, J. *et al.* Physiological consequences of heat stress in pigs. *Animal Production Science* **55**,
1381-1390 (2015).

35 Mayorga, E. J., Renaudeau, D., Ramirez, B. C., Ross, J. W. & Baumgard, L. H. Heat stress
adaptations in pigs. *Animal Frontiers* **9**, 54-61 (2018).

36 Blackshaw, J. K. & Blackshaw, A. Heat stress in cattle and the effect of shade on production and
behaviour: a review. *Australian Journal of Experimental Agriculture* **34**, 285-295 (1994).

37 Beede, D. & Collier, R. Potential nutritional strategies for intensively managed cattle during
thermal stress. *J. Anim. Sci.* **62**, 543-554 (1986).

38 Ganaie, A. *et al.* Biochemical and physiological changes during thermal stress in bovines: A
review. *Iranian Journal of Applied Animal Science* **3**, 423-430 (2013).

39 Hansen, P. Physiological and cellular adaptations of zebu cattle to thermal stress. *Anim. Reprod.*
Sci. **82**, 349-360 (2004).

40   Magdub, A., Johnson, H. & Belyea, R. Effect of environmental heat and dietary fiber on thyroid
596   physiology of lactating cows. *J. Dairy Sci.* **65**, 2323-2331 (1982).

597   41   Williamson, R., Misson, B. & Davison, T. The effect of exposure to 40 on the heat production and
598   the serum concentrations of triiodothyronine, thyroxine, and corticosterone in immature
599   domestic fowl. *Gen. Comp. Endocrinol.* **60**, 178-186 (1985).

600   42   Rodeberg, D. A., Meyer, J. G. & Babcock, G. F. Heat shock response: presence and effects in burn
601   patient neutrophils. *Journal of leukocyte biology* **66**, 773-780, doi:10.1002/jlb.66.5.773 (1999).

602   43   Orzheshkovskyi, V. V. & Trishchynska, M. A. Ceruloplasmin: Its Role in the Physiological and
603   Pathological Processes. *Neurophysiology* **51**, 141-149, doi:10.1007/s11062-019-09805-9 (2019).

604   44   Flick, K. & Kaiser, P. Protein degradation and the stress response. *Seminars in cell &
605   developmental biology* **23**, 515-522, doi:10.1016/j.semcdb.2012.01.019 (2012).

606   45   Zhang, D., Sweredoski, M. J., Graham, R. L., Hess, S. & Shan, S. O. Novel proteomic tools reveal
607   essential roles of SRP and importance of proper membrane protein biogenesis. *Molecular &
608   cellular proteomics : MCP* **11**, M111 011585, doi:10.1074/mcp.M111.011585 (2012).

609   46   Goetz, A. E. & Wilkinson, M. Stress and the nonsense-mediated RNA decay pathway. *Cellular
610   and molecular life sciences : CMLS* **74**, 3509-3531, doi:10.1007/s00018-017-2537-6 (2017).

611   47   Hariharan, S. & Dharmaraj, S. Selenium and selenoproteins: it's role in regulation of
612   inflammation. *Inflammopharmacology* **28**, 667-695, doi:10.1007/s10787-020-00690-x (2020).

613   48   Rajesh, K. *et al.* Phosphorylation of the translation initiation factor eIF2alpha at serine 51
614   determines the cell fate decisions of Akt in response to oxidative stress. *Cell Death Dis* **6**, e1591,
615   doi:10.1038/cddis.2014.554 (2015).

616   49   Zhang, W. & Liu, H. T. MAPK signal pathways in the regulation of cell proliferation in mammalian
617   cells. *Cell Res* **12**, 9-18, doi:10.1038/sj.cr.7290105 (2002).

618   50   Shiota, M., Yokomizo, A. & Naito, S. Pro-survival and anti-apoptotic properties of androgen
619   receptor signaling by oxidative stress promote treatment resistance in prostate cancer. *Endocr
620   Relat Cancer* **19**, R243-253, doi:10.1530/ERC-12-0232 (2012).

621   51   Lacoste, A., De Cian, M. C., Cueff, A. & Poulet, S. A. Noradrenaline and alpha-adrenergic signaling
622   induce the hsp70 gene promoter in mollusc immune cells. *J Cell Sci* **114**, 3557-3564 (2001).

623   52   Xian, D. *et al.* Emerging Roles of Redox-Mediated Angiogenesis and Oxidative Stress in
624   Dermatoses. *Oxid Med Cell Longev* **2019**, 2304018, doi:10.1155/2019/2304018 (2019).

625   53   Zhao, X. *et al.* Overexpression of Cardiomyocyte alpha1A-Adrenergic Receptors Attenuates
626   Postinfarct Remodeling by Inducing Angiogenesis Through Heterocellular Signaling. *Arterioscler
627   Thromb Vasc Biol* **35**, 2451-2459, doi:10.1161/ATVBAHA.115.305919 (2015).

628   54   Stupack, D. G. & Cheresh, D. A. Get a ligand, get a life: integrins, signaling and cell survival. *J Cell
629   Sci* **115**, 3729-3738, doi:10.1242/jcs.00071 (2002).

630   55   Reddy, E. P., Korapati, A., Chaturvedi, P. & Rane, S. IL-3 signaling and the role of Src kinases, JAKs
631   and STATs: a covert liaison unveiled. *Oncogene* **19**, 2532-2547, doi:10.1038/sj.onc.1203594
632   (2000).

633   56   Young, B., Purcell, C., Kuang, Y. Q., Charette, N. & Dupre, D. J. Superoxide Dismutase 1
634   Regulation of CXCR4-Mediated Signaling in Prostate Cancer Cells is Dependent on Cellular
635   Oxidative State. *Cell Physiol Biochem* **37**, 2071-2084, doi:10.1159/000438566 (2015).

636   57   Bai, X. C. *et al.* Phospholipase C-gamma1 is required for survival in heat stress: involvement of
637   protein kinase C-dependent Bcl-2 phosphorylation. *J Biochem* **131**, 207-212,
638   doi:10.1093/oxfordjournals.jbchem.a003089 (2002).

639   58   Parsell, D. & Lindquist, S. The function of heat-shock proteins in stress tolerance: degradation
640   and reactivation of damaged proteins. *Annu. Rev. Genet.* **27**, 437-496 (1993).

59 Rout, P., Kaushik, R. & Ramachandran, N. Differential expression pattern of heat shock protein
70 gene in tissues and heat stress phenotypes in goats during peak heat stress period. *Cell Stress*
Chaperones **21**, 645-651 (2016).
60 Westerheide, S. D., Anckar, J., Stevens, S. M., Sistonen, L. & Morimoto, R. I. Stress-inducible
regulation of heat shock factor 1 by the deacetylase SIRT1. *Science* **323**, 1063-1066 (2009).
61 Prodromou, C. Mechanisms of Hsp90 regulation. *Biochem. J.* **473**, 2439-2452 (2016).
62 Holmberg, C. I. *et al.* Phosphorylation of serine 230 promotes inducible transcriptional activity of
heat shock factor 1. *The EMBO journal* **20**, 3800-3810 (2001).
63 Wang, X., Grammatikakis, N., Siganou, A. & Calderwood, S. K. Regulation of molecular
chaperone gene transcription involves the serine phosphorylation, 14-3-3ε binding, and
cytoplasmic sequestration of heat shock factor 1. *Mol. Cell. Biol.* **23**, 6013-6026 (2003).
64 Ryu, K. Y. *et al.* The mouse polyubiquitin gene UbC is essential for fetal liver development,
cell-cycle progression and stress tolerance. *The EMBO journal* **26**, 2693-2706 (2007).
65 Lyzenga, W. J. & Stone, S. L. Abiotic stress tolerance mediated by protein ubiquitination. *J. Exp.*
Bot. **63**, 599-616 (2011).
66 Sulistio, Y. A. & Heese, K. The ubiquitin-proteasome system and molecular chaperone
deregulation in Alzheimer's disease. *Mol. Neurobiol.* **53**, 905-931 (2016).
67 Fernández-Fernández, M. R., Gragera, M., Ochoa-Ibarrola, L., Quintana-Gallardo, L. & Valpuesta,
J. M. Hsp70–a master regulator in protein degradation. *FEBS Lett.* **591**, 2648-2660 (2017).
68 Pabarcus, M. K. *et al.* CYP3A4 ubiquitination by gp78 (the tumor autocrine motility factor
receptor, AMFR) and CHIP E3 ligases. *Arch. Biochem. Biophys.* **483**, 66-74 (2009).
69 David, Y., Ziv, T., Admon, A. & Navon, A. The E2 ubiquitin-conjugating enzymes direct
polyubiquitination to preferred lysines. *J. Biol. Chem.* **285**, 8595-8604 (2010).
70 Sheng, Y. *et al.* A human ubiquitin conjugating enzyme (E2)-HECT E3 ligase structure-function
screen. *Mol. Cell. Proteomics* **11**, 329-341 (2012).
71 Kaiser, P. *et al.* A human ubiquitin-conjugating enzyme homologous to yeast UBC8. *J. Biol. Chem.*
269, 8797-8802 (1994).
72 Li, M. *et al.* Deubiquitination of p53 by HAUSP is an important pathway for p53 stabilization.
Nature **416**, 648 (2002).
73 Zhao, Y. *et al.* Noncanonical regulation of alkylation damage resistance by the OTUD4
deubiquitinase. *The EMBO journal* **34**, 1687-1703 (2015).
74 Kadowaki, H. *et al.* Pre-emptive quality control protects the ER from protein overload via the
proximity of ERAD components and SRP. *Cell Rep.* **13**, 944-956 (2015).
75 Hori, O. *et al.* Deletion of SERP1/RAMP4, a component of the endoplasmic reticulum (ER)
translocation sites, leads to ER stress. *Mol. Cell. Biol.* **26**, 4257-4267 (2006).
76 Nauseef, W. M., McCormick, S. J. & Clark, R. A. Calreticulin functions as a molecular chaperone
in the biosynthesis of myeloperoxidase. *J. Biol. Chem.* **270**, 4741-4747 (1995).
77 O'Connor, L. *et al.* Bim: a novel member of the Bcl-2 family that promotes apoptosis. *The EMBO*
journal **17**, 384-395 (1998).
78 Marani, M., Tenev, T., Hancock, D., Downward, J. & Lemoine, N. R. Identification of novel
isoforms of the BH3 domain protein Bim which directly activate Bax to trigger apoptosis. *Mol.*
Cell. Biol. **22**, 3577-3589 (2002).
79 Muneta, Y., Shimoji, Y., Inumaru, S. & Mori, Y. Molecular cloning, characterization, and
expression of porcine Fas ligand (CD95 Ligand). *J. Interferon Cytokine Res.* **21**, 305-312 (2001).
80 Ruiz-García, R. *et al.* Decreased activation-induced cell death by EBV-transformed B-cells from a
patient with autoimmune lymphoproliferative syndrome caused by a novel FASLG mutation.
Pediatr. Res. **78**, 603 (2015).

81   Nicholson, D. W. *et al.* Identification and inhibition of the ICE/CED-3 protease necessary for
mammalian apoptosis. *Nature* **376**, 37 (1995).

82   Dai, R., Frejtag, W., He, B., Zhang, Y. & Mivechi, N. F. c-Jun NH2-terminal kinase targeting and
phosphorylation of heat shock factor-1 suppress its transcriptional activity. *J. Biol. Chem.* **275**,
18210-18218 (2000).

83   Cao, W.-X. *et al.* MLKL mediates apoptosis via a mutual regulation with PERK/eIF2α pathway in
response to reactive oxygen species generation. *Apoptosis* **23**, 521-531 (2018).

84   Lu, X.-X. *et al.* PTEN inhibits cell proliferation, promotes cell apoptosis, and induces cell cycle
arrest via downregulating the PI3K/AKT/hTERT pathway in lung adenocarcinoma A549 cells.
BioMed research international **2016** (2016).

85   Chen, C.-Y., Chen, J., He, L. & Stiles, B. L. PTEN: tumor suppressor and metabolic regulator. *Front.*
Endocrinol. (Lausanne) **9** (2018).

86   Lee, Y. S. *et al.* Dysregulation of adipose glutathione peroxidase 3 in obesity contributes to local
and systemic oxidative stress. *Mol. Endocrinol.* **22**, 2176-2189 (2008).

87   Bierl, C., Voetsch, B., Jin, R. C., Handy, D. E. & Loscalzo, J. Determinants of human plasma
glutathione peroxidase (GPx-3) expression. *J. Biol. Chem.* **279**, 26839-26845 (2004).

88   Ighodaro, O. & Akinloye, O. First line defence antioxidants-superoxide dismutase (SOD), catalase
(CAT) and glutathione peroxidase (GPX): Their fundamental role in the entire antioxidant
defence grid. *Alexandria Journal of Medicine* **54**, 287-293 (2018).

89   Tsuji, K., Copeland, N., Jenkins, N. & Obinata, M. Mammalian antioxidant protein complements
alkylhydroperoxide reductase (ahpC) mutation in Escherichia coli. *Biochem. J.* **307**, 377-381
(1995).

90   Yamashita, H. *et al.* Characterization of human and murine PMP20 peroxisomal proteins that
exhibit antioxidant activity in vitro. *J. Biol. Chem.* **274**, 29897-29904 (1999).

91   Knoops, B. *et al.* Cloning and characterization of AOEB166, a novel mammalian antioxidant
enzyme of the peroxiredoxin family. *J. Biol. Chem.* **274**, 30451-30458 (1999).

92   Kang, S. W., Baines, I. C. & Rhee, S. G. Characterization of a mammalian peroxiredoxin that
contains one conserved cysteine. *J. Biol. Chem.* **273**, 6303-6311 (1998).

93   Chen, J.-W., Dodia, C., Feinstein, S. I., Jain, M. K. & Fisher, A. B. 1-Cys peroxiredoxin, a
bifunctional enzyme with glutathione peroxidase and phospholipase A2 activities. *J. Biol. Chem.*
275, 28421-28427 (2000).

94   Schmieder, R. & Edwards, R. Quality control and preprocessing of metagenomic datasets.
Bioinformatics **27**, 863-864 (2011).

95   Cunningham, F. *et al.* Ensembl 2015. *Nucleic Acids Res.* **43**, D662-D669 (2014).

96   Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

97   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**, 357
(2012).

98   Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. *Bioinformatics* **26**, 139-140 (2010).

99   Reimand, J., Kull, M., Peterson, H., Hansen, J. & Vilo, J. g: Profiler—a web-based toolset for
functional profiling of gene lists from large-scale experiments. *Nucleic Acids Res.* **35**, W193-
W200 (2007).

100   Reimand, J. *et al.* g: Profiler—a web server for functional interpretation of gene lists (2016
update). *Nucleic Acids Res.* **44**, W83-W89 (2016).

101   Jassal, B. *et al.* The reactome pathway knowledgebase. *Nucleic acids research* **48**, D498-D503,
doi:10.1093/nar/gkz1031 (2020).

102 Krämer, A., Green, J., Pollard Jr, J. & Tugendreich, S. Causal analysis approaches in ingenuity
pathway analysis. *Bioinformatics* **30**, 523-530 (2013).
103 Manjunath, S. *et al.* Genomic analysis of host - Peste des petits ruminants vaccine viral
transcriptome uncovers transcription factors modulating immune regulatory pathways. *Vet. Res.*
46, 15, doi:10.1186/s13567-015-0153-8 (2015).
104 Szklarczyk, D. *et al.* STRING v11: protein–protein association networks with increased coverage,
supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res.* **47**,
D607-D613 (2018).
105 Shannon, P. *et al.* Cytoscape: a software environment for integrated models of biomolecular
interaction networks. *Genome Res.* **13**, 2498-2504 (2003).
106 Schmittgen, T. D. & Livak, K. J. Analyzing real-time PCR data by the comparative C T method.
Nat. Protoc. **3**, 1101 (2008).

Legends

Figure 1: Overview of the work done : Two genetic groups (Tharparkar and Vrindavani) of
cattle were exposed to a temperature of 42 °C for 7 days. Heat stress indicators - Respiration rate
(RR), Rectal temperature and T3 level before exposure to heat (0day – control group) and at 7[th]
day of exposure (treated) were measured to evaluate heat stress. At these time points, RNA was
isolated from PBMCs for high throughput sequencing. Transcriptome analysis was done to
identify differentially expressed genes (DEGs) under heat treatment in both genetic groups.
Genes involved in physiological processes (heat stress response, apoptosis, ubiquitination,
unfolded protein response and antioxidant level) that are commonly associated with heat stress
were compared between the two genetic groups. Further, functional annotation of DEGs was
done using IPA.

Figure 2: Respiration rate, Rectal Temperature and T3 level measured at 0 day (control) and 7
day post-heat exposure (treated) in Vrindavani and Tharparkar (n=5) Levels sharing the same
superscript are not significantly (P > 0.05) different from each other.

Figure 3: Expression of DEGs in Vrindavani and Tharparkar under heat stress: (a) Venn
diagrams showing unique/common DEGs between Vrindavani and Tharparkar (b) Number of
upregulated and downregulated in both genetic groups (c) Contrast in the expression of common
DEGs

Figure 4: Canonical pathways activated/inactivated in (a) Vrindavani (b) Tharparkar under heat stress generated in the core analysis of Ingenuity pathway analysis tool. Orange color pathways are activated ($Z > 2$) and blue color pathways are inactivated ($Z < -2$). Height of the bar graphs indicates -log (p-value) and line graph showing the ratio of list genes found in each pathway over the total number of genes in that pathway.

Figure 5: Canonical pathways generated in Ingenuity Pathway Analysis of Oncostatin M signaling pathway of DEGs in (A) Vrindavani, (B) Tharparkar. Genes that were upregulated are shown in red and downregulated in green. The intensity of red and green corresponds to an increase and decrease, respectively, in Log2 fold change. Genes in grey were not significantly dysregulated and those in white are not present in the dataset but have been incorporated in the network through the relationship with other molecules by IPA.

Figure 6: Canonical pathways generated in Ingenuity Pathway Analysis of PTEN signaling pathway of DEGs in (A) Vrindavani, (B) Tharparkar. Genes that were upregulated are shown in red and downregulated in green. The intensity of red and green corresponds to an increase and decrease, respectively, in Log2 fold change. Genes in grey were not significantly dysregulated and those in white are not present in the dataset but have been incorporated in the network through the relationship with other molecules by IPA.

Figure 7: Contrast in the expression of genes involved in heat stress response, apoptosis, ubiquitination, unfolded protein response and balance in the production of ROS and antioxidants between two genetic groups.

Figure 8: Predicted interplay of molecules that is underway during heat stress in both groups : Heat stress causes unfolding of native proteins. HSP70 acts as a chaperone to facilitate refolding to restore the structure of unfolded proteins. Under normal condition, HSP70 is bound to HSF1 thereby preventing HSF1 to promote transcription of HSP70. Under heat stress ATP binds to the HSP70 and HSF1 complex to release HSF1, promoting the binding of the unfolded protein to HSP70 and ATP. CAMK2D that induces the transcription of HSP70 via HSF1 was found downregulated in Tharparkar. GSK3A that inhibits the trimerization of HSF1 that is needed for the induction of HSP70 expression was found upregulated in Tharparkar. The decreased level of HSP70 in Vrindavani makes it inevitable that such negative feedbacks would further reduce its level and GSK3A was found downregulated and CAMK2D, upregulated.

Further, in Tharparkar, HSP70 tends to activate ubiquitination pathway to decrease the
accumulation of unfolded proteins that can't be refolded by it. This pathway activation is
supported by upregulation of E3 ligases (UBE2G1, UBE2S, and UBE2H) in Tharparkar.
However, the E3 ligase in Vrindavani was found downregulated. With HSP70 being upregulated
and having cytoprotection activity, Tharparkar shows the decline in apoptosis as compared to
Vrindavani. This is supported by downregulation of BCL2L11, FASLG, MLKL, CASP3,
MAPK8 and VIM in Tharparkar and vice-versa. Besides, higher expression of the antioxidants
(SOD, CAT, GPX) in Tharparkar enables it to cope up with higher levels of free radicals
generated as a result of heat stress while Vrindavani is unable to do so. Green arrow indicates
downregulation and Maroon arrow indicates upregulation.

Supplementary.Figure 1: Comparison of activated/inactivated pathways in Vrindavani and
Tharparkar. Activated pathways have Z score > 2 and indicated by red colour while inactivated
pathways are having Z score < - 2 and indicated by green colour.

Supplementary Figure 2: Comparison of activated/inactivated miRNAs in Vrindavani and
Tharparkar as predicted by IPA upstream analysis. Activated pathways have Z score > 2 and
indicated by red colour while inactivated pathways are having Z score < - 2 and indicated by
green colour.

Supplementary Figure 3: Comparison of activated/inactivated Transcription factors as
predicted by IPA upstream analysis (Transcription factors of Vrindavani are red-coloured and
Tharparkar are blue-coloured) vis-à-vis their Log2FC in both genetic groups. The encircled ones
are common to both groups.

Supplementary Figure 4: Predicted Protein-protein interaction network of expressed genes
common to Tharparkar and Vrindavani. The diameter of the node represents the
connectivity/degree of the node among the genes.

Supplementary Figure 5: Validation of RNA sequencing data by Real-Time data in Vrindavani
(a) and Tharparkar (b). The expression of 10 selected genes was found in concordance with RNA
Sequencing data. The correlation (r2 = 0.9942 in (a) and 0.9972 in (b)) was found to be
significant (P< .01) in both cases.

Crossbred_DEGs　　　　　　　　　　　　**Tharparkar_DEGs**

(a)

(c)

(b)

(a)

(b)

(a)　　　　　　　　　　　　　　　　　**(b)**

(a)

(b)