Quantify the Requirements to Achieve Grain Zn Biofortification of High-yield Wheat on Calcareous Soils

Sen Wang[1,2,3], Zhaohui Wang[1,2*], Shasha Li[1,2], Chaopeng Diao[1,2], Lu Liu[1,2], Ning Huang[1,2],
Ming Huang[1,2], Xiaoli Hui[1,2], Laichao Luo[1,2], Gang He[1,2], Hanbing Cao[1,2]

[1]State Key Laboratory of Crop Stress Biology for Arid Areas, Northwest A&F University, Yangling,
Shaanxi, China

[2]Key Laboratory of Plant Nutrition and Agri-environment in Northwest China, Ministry of
Agriculture and Rural Affairs, College of Natural Resources and Environment, Northwest A&F
University, Yangling, Shaanxi, China

[3]Shenzhen Branch, Guangdong Laboratory for Lingnan Modern Agriculture, Genome Analysis
Laboratory of the Ministry of Agriculture and Rural Affairs, Agricultural Genomics Institute at
Shenzhen, Chinese Academy of Agricultural Sciences, Shenzhen, Guangdong, China

* Correspondence:
Zhaohui Wang
w-zhaohui@263.net

Keywords: wheat, grain Zn biofortification, Zn distribution, Zn uptake, quantification.

Abstract

The solution to address global human Zn deficiency is Zn biofortification of staple food crops, aimed
at high grain Zn concentration as well as high yield. However, the desired high grain Zn
concentration above 40 mg kg^{-1} is rarely observed for high-yield wheat on worldwide calcareous
soils, due to inadequate Zn uptake or Zn distribution to grain. The present study aims to investigate
how much Zn uptake or distribution is adequate to achieve the Zn.t of high-yield wheat on calcareous
soils with low available Zn (~ 0.5 mg kg^{-1}). Of the 123 cultivars tested in a three-year field
experiment, 19 high-yield cultivars were identified with similar yields around 7.0 t ha^{-1} and various
grain Zn concentrations from 9.3 to 26.7 mg kg^{-1}. The adequate Zn distribution to grain was defined
from the view of Zn biofortification, as the situation where the Zn distribution to grain (Zn harvest
index) increased to the observed maximum of ~ 91.0% and the Zn concentration of vegetative parts
(straw Zn concentration) decreased to the observed minimum of ~ 1.5 mg kg^{-1} (Zn.m). Under the
assumed condition of adequate Zn distribution to grain (~ 91.0%), all the extra Zn above Zn.m was
remobilized from straw to grain and the grain Zn concentration would be increased to its highest
attainable level, which was 14.5 ~ 31.3 mg kg^{-1} for the 19 high-yield cultivars but still lower than 40
mg kg^{-1}. Thus, even with the adequate Zn distribution to grain, the current Zn uptake is still not
adequate and needs to be increased to 308 g ha^{-1} or higher to achieve Zn.t for high-yield wheat (7.0 t
ha^{-1}) on low-Zn calcareous soils. Besides, the established method here can also provide the priority
measures and quantitative guidelines to achieve Zn biofortification in other wheat production
regions.

1   Introduction

Although over 50 years have passed since the first case of human zinc (Zn) deficiency (Prasad,
2013), now this problem is still afflicting over 16% of the world population, mostly in Africa and
South Asia (Kumssa et al., 2015). In these areas, residents cannot get sufficient Zn intake from daily
diets which are mainly low-Zn cereals like wheat, maize, and rice (Wessells and Brown, 2012).
Increasing the Zn concentration of cereal grains, known as Zn biofortification, is the main solution to
alleviate human Zn deficiency, and the target of Zn fortified wheat grain is 40-50 mg kg^{-1} (Cakmak
2018), much higher than the current worldwide grain Zn concentration of 20-30 mg kg^{-1} (Liu et al.,
2014; Chen et al., 2017). To close the gap between the current and target level, it is necessary to fully
understand the factors affecting grain Zn concentration.

Grain yield, as an indicator of sink capacity, has essential influence on grain Zn concentration. For
instance, 45% decrease in yield of field-grown wheat generated 60% increase in grain Zn
concentration (Zhang et al., 2012). The negative correlation between grain Zn concentration and
yield has been widely reported in different locations (Liu et al., 2014) and wheat germplasms
(Guttieri et al., 2015), and the lowered grain Zn concentration in modern cultivars is mainly caused
by the wide adoption of high-yield cultivars after the Green Revolution (Fan et al., 2008; Velu et al.,
2017). To feed 9.7 billion population by 2050, the staple crop production has to increase by 60%
(FAO, 2014), and thus crop Zn biofortification should increase yield and grain Zn concentration
simultaneously (Bouis and Welch, 2010). To achieve high yield and high grain Zn concentration for
wheat, it is necessary to enhance both shoot Zn uptake and its distribution to grain (Wang et al.,
2018), which are the two sources of Zn accumulated in grain (Kutman et al., 2012; Xue et al., 2012).
For example, wheat grain Zn concentration was increased by enhanced nitrogen (N) supply (Kutman
et al., 2011; Hui et al., 2017) or ectopic-expressed rice *NICOTIANAMINE SYNTHASE 2* gene (Singh
et al., 2017b), because abundant N-containing ligands like nicotianamine (NA) in the phloem
enhanced the Zn transportation to grain (Barunawati et al., 2013). Also, the enhanced root Zn uptake
increased wheat grain Zn concentration by enlarged root contact area with soil (Ercoli et al., 2017;
Singh et al., 2017a), external Zn fertilization (Saha et al., 2017; Liu et al., 2017) or the mobilization
of soil intrinsic Zn (Wang et al., 2017). However, most of existing Zn biofortification studies pay
much less attention to grain yield, and thus cannot fully achieve the target of Zn biofortification.

The most cost-effective approach to realize crop Zn biofortification is developing cultivars with high
yield and high grain Zn concentration (Gregory et al., 2017). For example, a few breeding lines or
cultivars of spring wheat exhibited high grain Zn concentration ~ 40 mg kg^{-1} and high yield ~ 4 t ha^{-1},
under soil available Zn of 1.9 mg kg^{-1} in Canada, India, Pakistan, and Mexico (Gao et al., 2011; Velu
et al., 2012). Similarly, under soil available Zn of 1.2 mg kg^{-1} in Iran, several bread wheat genotypes
showed high grain Zn concentration > 50 mg kg^{-1} and high yield > 6 t ha^{-1} (Amiri et al., 2015). Thus,
under relatively high soil available Zn > 1.0 mg kg^{-1}, breeding new wheat cultivars can achieve high
grain Zn concentration and high yield simultaneously. But on calcareous soils with low available Zn
~ 0.5 mg kg^{-1}, the success of Zn biofortification of high-yield wheat is rarely achieved. For instance,
the highest grain Zn concentration of spring wheat and bread wheat lines was only 25 mg kg^{-1} under
soil available Zn of 0.3 ~ 0.6 mg kg^{-1} in Portugal and Iran (Gomez-Coronado et al., 2016;
Khoshgoftarmanesh et al., 2013). In the present study on calcareous soils with available Zn of 0.4 mg
kg^{-1} in northwest China, the grain Zn concentrations of 123 wheat cultivars were all below 30 mg kg^{-1}.
Does this mean that high grain Zn concentration of 40 mg kg^{-1} cannot be achieved by breeding for
high-yield wheat on calcareous soils?

The increase of grain Zn concentration for high-yield wheat relies on both high Zn uptake and high
Zn distribution to grain (Wang et al., 2018). In the present work, the Zn harvest index varied from
45.5% to 94.0% and the straw Zn concentration at maturity varied from 1.2 to 10.5 mg kg^{-1}, while the

minimum Zn concentration in wheat vegetative parts was ~ 5.0 mg kg^{-1} in hydroponics (Kutman et
al. 2012), indicating that some Zn in straw might be still transported to grain. Accordingly,
inadequate Zn distribution to grain can be the reason for the low grain Zn concentration observed for
high-yield wheat. Then, from the view of Zn biofortification, how much Zn distribution to grain is
adequate for field-grown high-yield wheat? With adequate Zn distribution, whether the Zn uptake is
adequate to achieve high grain Zn concentration (> 40 mg kg^{-1}) for high-yield wheat? To test these
hypotheses, the Zn concentration, uptake, and distribution of 123 wheat cultivars over three years
were analyzed in the present study.

2   Materials and methods

2.1   Plant materials and field experiment

The field experiment was conducted in Yongshou (108°12′E, 34°44′N) located on the southern Loess
Plateau, where the climate is the temperate continental monsoon climate and over half of the annual
precipitation (~ 550 mm) occurs in the summer fallow period from July to September. The top 20 cm
soil properties were determined according to Bao (2000) as follows: pH 8.4 (CO_2-free water), cation
exchange capacity 20.3 cmol kg^{-1}, soil organic matter 12.9 g kg^{-1}, total N 0.9 g kg^{-1}, mineral N 26.6
mg kg^{-1}, available phosphorus (P) 16.9 mg kg^{-1}, available potassium (K) 123.4 mg kg^{-1}, available iron
(Fe) 7.5 mg kg^{-1}, available manganese (Mn) 18.1 mg kg^{-1}, available copper (Cu) 1.3 mg kg^{-1}, and
available Zn 0.4 mg kg^{-1}.

One hundred and twenty-three wheat cultivars were collected (Supp Table 1) and tested in the three-
year field experiment with a randomized complete block design. Each cultivar had four replications
and each plot consisted of four 200 cm long rows with 2.5 cm seed spacing and 20 cm row spacing,
under the nutrient supply of 150 kg N ha^{-1} and 100 kg P_2O_5 ha^{-1}. All cultivars were sown during 28th
to 30th September in 2013, 2nd to 3rd October in 2014, and 26th to 28th September in 2015, and
harvested during 14th to 17th June in 2014, 18th to 20th June in 2015, and 18th to 20th June in 2016.
The rainfalls during summer fallow periods and growing seasons were 271 and 267 mm in 2013-
2014, 317 and 314 mm in 2014-2015, and 228 and 186 mm in 2015-2016, respectively. Throughout
the experimental period, no irrigation was conducted, and herbicide and pesticide were used when
necessary.

2.2   Sampling and chemical analyses

At maturity, the plants of 30 ears were randomly sampled from the central two rows of each cultivar
plot, and the roots were cut off at the stem-root joint part. Then, the shoots were air-dried and
separated into stems, glumes, and grains, which were washed with deionized water and oven-dried at
65℃ to determine dry weight. After that, the oven-dried samples were ground by a ball mill (Retsch
MM400, Germany) for chemical analyses. The remaining ears in the central two rows of each plot
were also harvested and weighted, plus the above grain weight, to estimate the grain yield, glume
biomass, and stem biomass in oven-dried weight.

Plant Zn concentration was determined as Ozbek and Akman (2016) with some modifications.
Briefly, 0.2 g sample was mixed with 5.0 ml HNO_3 (65%) in a 50 ml Teflon tube and predigested at
120℃ for 0.5 h, and then 1.0 ml H_2O_2 (30%) was added before microwave digestion (MW Pro,
Anton Paar, Austria). Each sample was digested in two technical duplicates and the standard wheat
flour (GBW10011 GSB-2) was used for quality control. The digestion solution was diluted with
ultrapure water (18.25 MΩ cm^{-1}), and Zn concentration was determined by ICP-MS (iCAP Qc,

Thermo Fisher Scientific, USA). The measured Zn isotope was ^{66}Zn, and ^{73}Ge was added as internal
standard to calibrate the signal fluctuation.

2.3   Quantification of the potential increase of grain Zn concentration by adequate Zn
126
distribution

From the perspective of Zn biofortification, the adequate Zn distribution to grain refers to the
situation where nearly all the mobilizable Zn in shoot vegetative parts was transported or remobilized
to grain, and correspondingly, the ratio of grain Zn uptake to shoot Zn uptake (Zn harvest index)
increased to the maximum and the Zn concentration of vegetative parts (straw Zn concentration,
Zn.s) decreased to the minimum. To determine the adequate Zn distribution to grain for high-yield
wheat, we selected the cultivars with yields higher than the corresponding median of all cultivars in
each year, calculated the Zn uptake and Zn harvest index, and combined stem and glume Zn
concentrations into Zn.s by the weighted mean method. In each year, the observed maximum Zn
harvest index or minimum straw Zn concentration (Zn.m) was used to indicate the level of adequate
Zn distribution to grain. Since Zn concentration was easier to measure than Zn harvest index, we
used Zn.m to calculate the potential increase of grain Zn concentration (Zn.p) caused by the extra Zn
remobilized from straw to grain under the condition of adequate Zn distribution to grain.

$$\text{Zn uptake (g ha}^{-1}) = \frac{\text{Zn concentration (mg kg}^{-1}) \times \text{yield or biomass(kg ha}^{-1})}{1000}$$

$$\text{Zn. s (mg kg}^{-1}) = \frac{\text{stem Zn uptake(g ha}^{-1}) + \text{glume Zn uptake(g ha}^{-1})}{\text{stem biomass(kg ha}^{-1}) + \text{glume biomass(kg ha}^{-1})} \times 1000$$

$$\text{Zn. p (mg kg}^{-1}) = \frac{[\text{Zn. s (mg kg}^{-1}) - \text{Zn. m (mg kg}^{-1})] \times \text{Bm. s (kg ha}^{-1})}{\text{GrY (kg ha}^{-1})}$$

140   Here Bm.s and GrY refer to straw biomass and grain yield. The sum of Zn.p and the current grain Zn
141   concentration (Zn.c) was defined as the attainable grain Zn concentration (Zn.a).

142   Shoot Zn uptake and Zn harvest index (ZnHI) were calculated as described by Kutman et al. (2011)
143   and Xue et al. (2012).

$$\text{Shoot Zn uptake} = \text{grain Zn uptake} + \text{glume Zn uptake} + \text{stem Zn uptake}$$

$$\text{Zn harvest index (\%)} = \frac{\text{Grain Zn uptake (g ha}^{-1})}{\text{Shoot Zn uptake (g ha}^{-1})} \times 100$$

2.4   Statistical analyses and graphing

Analysis of variance (ANOVA) was used to test the effects of year, cultivar, and year × cultivar
interaction on yield or biomass, Zn concentrations, and Zn uptakes. The relationships of grain Zn
concentration with other traits were tested by Pearson correlation or linear regression. All the
statistical analyses were completed by PROC GLM, PROC CORR or PROC REG in SAS v 8.01,
with 0.05 set as the significance level. Scatter and bar plots were created by Sigmaplot 12.5, and the
concept chart illustrating Zn biofortification guidelines was created in Adobe Illustrator CC.

3   Results

3.1 Relationship of grain Zn concentration with other traits for high-yield wheat

Of the 123 tested wheat cultivars ordered by yield from low to high, 19 cultivars consistently
exhibited higher yields than the median in each year, and were identified as high-yield cultivars
(Figure 1). Although yield and biomasses showed significant differences among three years, the 19
high-yield cultivars exhibited similar yields (~ 7 t ha^{-1}) and varied greatly in grain Zn concentration,
from 14.7 to 26.7 mg kg^{-1} in 2014, 9.3 to 22.2 mg kg^{-1} in 2015, and 13.5 to 17.0 mg kg^{-1} in 2016
(Supp Table 2 and Figure 2A). Besides, grain Zn concentration showed no relation with yield, the
biomass and Zn concentration of stem and glume (Figure 2A and Supp Figure 1). However, shoot Zn
uptake was positively correlated with grain Zn concentration in each year and Zn harvest index was
positively correlated with grain Zn concentration in 2015 (Figure 2C and 2D).

3.2 Minimum straw Zn concentration and maximum Zn harvest index

Shoot Zn uptake and Zn harvest index showed large variations among the 19 high-yield cultivars or
three experimental years (Supp Table 2, Figure 3A and 3B). The observed maximum Zn harvest
index was 91.8% in 2014, 87.7% in 2015, and 94.0% in 2016 (Figure 3B), while the observed
minimum straw Zn concentration (Zn.m) was 1.3 mg kg^{-1} in 2014, 1.9 mg kg^{-1} in 2015, and 1.2 mg
kg^{-1} in 2016 (Figure 3C). With Zn.m used to indicate the level of adequate Zn distribution to grain,
the gap between current straw Zn concentration and Zn.m was used to calculate the amount of extra
Zn which could be remobilized into grain.

3.3 Potential increase of grain Zn concentration and attainable grain Zn concentration

The potential increase of grain Zn concentration (Zn.p) due to extra Zn remobilized from straw to
grain of the 19 high-yield cultivars ranged from 2.1 to 5.5 mg kg$_{-1}$ in 2014, 3.0 to 6.5 mg kg^{-1} in
2015, and 3.5 to 7.5 mg kg^{-1} in 2016 (Figure 4). When Zn.p was added to the current grain Zn
concentration (Zn.c), the attainable grain Zn concentration (Zn.a) varied from 17.0 to 31.3 mg kg^{-1} in
2014, 14.5 to 27.5 mg kg^{-1} in 2015, and 17.6 to 22.8 mg kg^{-1} in 2016 (Figure 4), lower than the Zn
biofortification target (Zn.t) of 40 mg kg^{-1}. Thus, even with the adequate Zn distribution to grain, the
current shoot Zn uptake was not adequate to ahieve Zn.t of high-yield wheat. For an assumed high-
yield (~ 7 t ha^{-1}) wheat cultivar with adequate Zn distribution to grain (~ 91.0%), the required shoot
Zn uptake should be at least 308 g ha^{-1} to achieve the Zn.t.

4 Discussion

**4.1 Using the minimum straw Zn concentration to estimate the adequate Zn distribution to
grain**

In the present work, the adequate Zn distribution to grain is defined from the view of Zn
biofortification, and it refers to the situation where all the mobilizable Zn in shoot vegetative parts
was transported or remobilized to grain, meaning that Zn harvest index increased to its maximum and
straw Zn concentration decreased to its minimum (Zn.m). Then, the gap between current straw Zn
concentration and Zn.m can be used to calculate the amount of extra remobilized Zn from straw to
grain, which results in the potential increase and the attainable highest grain Zn concentration under
specific conditions. For high-yield bread wheat grown on low-Zn calcareous soils, the observed
minimum straw Zn concentration is used as Zn.m which is ~ 1.5 mg kg^{-1} and lower than the reported
5.0 mg kg^{-1} for durum wheat in hydroponics (Kutman et al., 2012). Presumably, the senescent
vegetative parts of field-grown wheat need less Zn to constitute structural components and more Zn
can be remobilized into grain. Under uniform environmental conditions, straw Zn concentration and
Zn.m can be used to determine whether the Zn distribution to grain is adequate or not to promote Zn

biofortification. Since the Zn biofortification of high-yield wheat needs both high Zn uptake and high
Zn distribution to grain (Wang et al., 2018), the Zn concentration of vegetative parts, which is easier
to measure than the Zn harvest index, deserves to be considered in future studies on wheat Zn
biofortification.

4.2   Quantify the requirements to achieve wheat Zn biofortification in different environments

Based on the adequate Zn distribution to grain (~ 91.0%) as indicated by Zn.m, we can get the
attainable highest grain Zn concentration (Zn.a) and compare it with the grain Zn biofortification
target (Zn.t, 40 mg kg^{-1}), which can tell us whether the current Zn uptake is adequate to achieve Zn.t
for high-yield wheat (~ 7 t ha^{-1}). Besides, the established quantitative analysis method here are also
suitable for other wheat production regions with the aim of increasing the current grain Zn
concentration (Zn.c) and grain yield. The priority measures to achieve the Zn.t of high-yield wheat
are different under different conditions (Figure 5). In wheat planting areas where Zn.a is close to or
higher than Zn.t and the soil available Zn is often relatively high (> 1.0 mg kg^{-1}), the priority to
realize Zn.t is to adopt or develop the cultivars with high Zn distribution to grain. For example, in
North China Plains with soil available Zn of 3.4 mg kg^{-1}, optimal N application generated high yield
of 7.5 t ha^{-1} and shoot Zn uptake of 370 g ha^{-1} (Xue et al., 2014), and the Zn.a would be 44.9 mg kg^{-1}
and higher than Zn.t, indicating the necessity to introduce high-Zn-distribution wheat cultivars. In
wheat production regions where Zn.a is lower than Zn.t and higher than Zn.c and the soil available
Zn is often relatively low (< 0.5 mg kg^{-1}), developing high-Zn-distribution cultivars should be
combined with agronomic measures like Zn fertilization to promote Zn uptake. For instance, in Iran
rainfed drylands with soil available Zn of 0.2 mg kg^{-1} (Norouzi et al., 2014), the tested wheat
cultivars exhibited high yield of 6.6 t ha^{-1} and shoot Zn uptake of 174 g ha^{-1}. Even if the Zn.c of 17.0
mg kg^{-1} was increased to the Zn.a of 26.4 mg kg^{-1} by introducing new cultivars, there was still a large
gap to Zn.t which needed to be closed by increasing Zn uptake. In wheat production regions where
Zn.a is close to Zn.c and lower than Zn.t, the Zn distribution to grain has reached its maximum and
the priority measure of Zn biofortification is to increase Zn uptake by agronomic practices. However,
this case is rarely observed up to now because researchers have not paid enough attention to the Zn in
vegetative parts in wheat Zn biofortification studies.

On calcareous soils with low available Zn < 0.5 mg kg^{-1}, the current wheat grain Zn concentration is
around 20 mg kg^{-1}, as reported in China, Kazakhstan, Mexico, Turkey, and Zambia, etc. (Zou et al.,
2012; Liu et al., 2014). In the present study on calcareous soils with available Zn of 0.4 mg kg^{-1}, the
wheat cultivar Zhengmai 7698 had the highest Zn.a of 31.3 mg kg^{-1} and the yield of 6.6 t ha^{-1} in
2014, which was still lower than the Zn.t. To achieve high grain Zn concentration of 40 mg kg^{-1} and
high yield of 7.0 t ha^{-1} simultaneously, the current shoot Zn uptake should be increased to at least 308
g ha^{-1} by agronomic measures like soil Zn fertilization with the wheat cultivars with high Zn
distribution to grain (~ 91%). These guidelines deserve to be tested in field studies on Zn
biofortification.

5   Conclusion

From the view of crop Zn biofortification, the adequate Zn distribution to grain can be defined as the
case where Zn harvest index increased to its maximum and straw Zn concentration decreased to its
minimum. For the high-yield (~ 7 t ha^{-1}) wheat grown on low-Zn (~ 0.5 mg kg^{-1}) calcareous soils, the
maximum Zn harvest index was ~ 91.0% and the minimum straw Zn concentration was ~ 1.5 mg kg^{-1}.
Under the condition of adequate Zn distribution to grain, the gap between straw Zn concentration and
its minimum could be used to determine the extra Zn remobilized to grain and the highest attainable

grain Zn concentration, which was 14.5 ~ 31.3 mg kg^{-1} for the high-yield wheat cultivars and lower
than the grain Zn biofortification target (Zn.t) of 40 mg kg^{-1}. Therefore, even with the adequate Zn
distribution to grain (~ 91%), the current shoot Zn uptake is still not adequate to achieve the Zn.t of
high-yield wheat grown on low-Zn calcareous soil, and the priority measure of Zn biofortification is
to increase Zn uptake to 308 g ha^{-1} or higher by agronomic practices like Zn fertilization. The
established method here can also provide the most suitable guidelines and quantitative requirements
to achieve grain Zn biofortification in other wheat production regions.

6   Abbreviations

247   Zn.s, straw Zn concentration; Zn.m, minimum straw Zn concentration; Zn.p, potential increase of
248   grain Zn concentration; Zn.c, current grain Zn concentration; Zn.a, attainable grain Zn concentration;
249   ZnHI, Zn harvest index; Zn.t, grain Zn biofortification target.

7   Conflict of Interest

251   The authors declare that the research was conducted in the absence of any commercial or financial
252   relationships that could be construed as a potential conflict of interest.

8   Author Contributions

254   SW, ZHW, SSL, CPD, and LL contributed conception and design of the study. SW, SSL, CPD, LL,
255   NH, MH, XLH, LCL, GH, and HBC conducted experiment, sampling, chemical and statistical
256   analyses. SW wrote the first draft of the manuscript and ZHW, MH, XLH, LCL, GH, and HBC read
257   and revised the manuscript. All authors contributed to manuscript revision, read and approved the
258   submitted version.

9   Funding

260   The authors thank for the financial supports from the China Agricultural Research System (CARS-3),
261   the Special Fund for Agro-scientific Research in the Public Interest under Grant (201303104), the
262   Agricultural Scientific Research Talent and Team Program, the National Natural Science Foundation
263   of China (NSFC) (41571282, 41401330), and the China Postdoctoral Science Foundation
264   (2019M650919).

10   Acknowledgments

266   During the wheat germplasm collection, we thank for the selfless help from the staff in different
267   research laboratories and experimental stations of the China Agricultural Research System. We also
268   thank the professors and doctors of the Soil Fertility and Plant Nutrition Team of Northwest A&F
269   University, for their critical comments and constructive suggestions.

11   References

Amiri, R., Bahraminejad, S., Sasani, S., Jalali-Honarmand, S., and Fakhri, R. (2015). Bread wheat
genetic variation for grain's protein, iron and zinc concentrations as uptake by their genetic ability.
Eur. J. Agron. 67, 20-26. doi: 10.1016/j.eja.2015.03.004.

Bao, S. D. (2000). *Soil and agricultural chemistry analysis (in Chinese)*. Beijing: China Agriculture
Press.

Barunawati, N., Giehl, R. F. H., Bauer, B., and von Wiren, N. (2013). The influence of inorganic
nitrogen fertilizer forms on micronutrient retranslocation and accumulation in grains of winter wheat.
Front. Plant Sci. 4, 320. doi: 10.3389/fpls.2013.00320.

Bouis, H. E., and Welch, R. M. (2010). Biofortification-A Sustainable Agricultural Strategy for
Reducing Micronutrient Malnutrition in the Global South. *Crop Sci.* 50(2), S20-S32. doi:
10.2135/cropsci2009.09.0531.

Cakmak, I., and Kutman, U.B. (2018). Agronomic biofortification of cereals with zinc: a review.
Eur. J. Soil Sci. 69(1), 172-180. doi: doi:10.1111/ejss.12437.

Chen, X. P., Zhang, Y. Q., Tong, Y. P., Xue, Y. F., Liu, D. Y., Zhang, W., et al. (2017). Harvesting
more grain zinc of wheat for human health. *Sci. Rep.* 7(1), 7016. doi: 10.1038/s41598-017-07484-2.

Ercoli, L., Schüßler, A., Arduini, I., and Pellegrino, E. (2017). Strong increase of durum wheat iron
and zinc content by field-inoculation with arbuscular mycorrhizal fungi at different soil nitrogen
availabilities. *Plant Soil* 419(1), 153-167. doi: 10.1007/s11104-017-3319-5.

Fan, M. S., Zhao, F. J., Fairweather-Tait, S. J., Poulton, P. R., Dunham, S. J., and Mcgrath, S. P.
(2008). Evidence of decreasing mineral density in wheat grain over the last 160 years. *J. Trace Elem.*
Med. Bio. 22(4), 315–324. doi: 10.1016/j.jtemb.2008.07.002.

FAO (2014). "*Building a common vision for sustainable food and agriculture: principles and*
approaches". (Rome: FAO).

Gao, X., Mohr, R. M., McLaren, D. L., and Grant, C. A. (2011). Grain cadmium and zinc
concentrations in wheat as affected by genotypic variation and potassium chloride fertilization. *Field*
Crop Res. 122(2), 95-103. doi: 10.1016/j.fcr.2011.03.005.

Gomez-Coronado, F., Poblaciones, M. J., Almeida, A. S., and Cakmak, I. (2016). Zinc (Zn)
concentration of bread wheat grown under Mediterranean conditions as affected by genotype and
soil/foliar Zn application. *Plant Soil* 401(1), 331-346. doi: 10.1007/s11104-015-2758-0.

Gregory, P. J., Wahbi, A., Adu-Gyamfi, J., Heiling, M., Gruber, R., Joy, E. J. M., et al. (2017).
Approaches to reduce zinc and iron deficits in food systems. *Glob. Food Secur.* 15, 1-10. doi:
10.1016/j.gfs.2017.03.003.

Guttieri, M. J., Baenziger, P. S., Frels, K., Carver, B., Arnall, B., and Waters, B. M. (2015). Variation
for Grain Mineral Concentration in a Diversity Panel of Current and Historical Great Plains Hard
Winter Wheat Germplasm. *Crop Sci.* 55(3), 1035-1052. doi: 10.2135/cropsci2014.07.0506.

Hui, X. L., Wang, Z. H., Luo, L. C., Ma, Q. X., Wang, S., Dai, J., et al. (2017). Winter Wheat Grain
Yield and Zn Concentration Affected by Long-Term N and P Application in Dryland (in Chinese).
Sci. Agric. Sin. 50(16), 3175-3185. doi: 10.3864/j.issn.0578-1752.2017.16.012.

Khoshgoftarmanesh, A. H., SanaeiOstovar, A., Sadrarhami, A., and Chaney, R. (2013). Effect of tire
rubber ash and zinc sulfate on yield and grain zinc and cadmium concentrations of different zinc-
deficiency tolerance wheat cultivars under field conditions. *Eur. J. Agron.* 49, 42-49. doi:
10.1016/j.eja.2013.02.013.

Kumssa, D. B., Joy, E. J., Ander, E. L., Watts, M. J., Young, S. D., Walker, S., et al. (2015). Dietary
calcium and zinc deficiency risks are decreasing but remain prevalent. *Sci. Rep.* 5, 10974. doi:
10.1038/srep10974.

Kutman, U. B., Kutman, B.Y., Ceylan, Y., Ova, E. A., and Cakmak, I. (2012). Contributions of root
uptake and remobilization to grain zinc accumulation in wheat depending on post-anthesis zinc
availability and nitrogen nutrition. *Plant Soil* 361(1-2), 177-187. doi: 10.1007/s11104-012-1300-x.

Kutman, U. B., Yildiz, B., and Cakmak, I. (2011). Effect of nitrogen on uptake, remobilization and
partitioning of zinc and iron throughout the development of durum wheat. *Plant Soil* 342(1-2), 149-
164. doi: 10.1007/s11104-010-0679-5.

Liu, D. Y., Zhang, W., Pang, L. L., Zhang, Y. Q., Wang, X. Z., Liu, Y. M., et al. (2017). Effects of
zinc application rate and zinc distribution relative to root distribution on grain yield and grain Zn
concentration in wheat. *Plant Soil* 411(1), 167-178. doi: 10.1007/s11104-016-2953-7.

Liu, H., Wang, Z. H., Li, F. C., Li, K. Y., Yang, N., Yang, Y. E., et al. (2014). Grain iron and zinc
concentrations of wheat and their relationships to yield in major wheat production areas in China.
Field Crop Res. 156, 151-160. doi: 10.1016/j.fcr.2013.11.011.

Norouzi, M., Khoshgoftarmanesh, A. H., and Afyuni, M. (2014). Zinc fractions in soil and uptake by
wheat as affected by different preceding crops. *Soil Sci. Plant Nutr.* 60(5), 670-678. doi:
10.1080/00380768.2014.937674.

Ozbek, N., and Akman, S. (2016). Method development for the determination of calcium, copper,
magnesium, manganese, iron, potassium, phosphorus and zinc in different types of breads by
microwave induced plasma-atomic emission spectrometry. *Food Chem.* 200, 245-248. doi:
10.1016/j.foodchem.2016.01.043.

Prasad, A. S. (2013). Discovery of Human Zinc Deficiency: Its Impact on Human Health and
Disease. *Adv. Nutr.* 4(2), 176-190. doi: 10.3945/an.112.003210.

Saha, S., Chakraborty, M., Sarkar, D., Batabyal, K., Mandal, B., Murmu, S., et al. (2017).
Rescheduling zinc fertilization and cultivar choice improve zinc sequestration and its bioavailability
in wheat grains and flour. *Field Crop Res.* 200, 10-17. doi: 10.1016/j.fcr.2016.09.006.

Singh, D., Rajawat, M. V. S., Kaushik, R., Prasanna, R., and Saxena, A. K. (2017a). Beneficial role
of endophytes in biofortification of Zn in wheat genotypes varying in nutrient use efficiency grown
in soils sufficient and deficient in Zn. *Plant Soil* 416(1), 107-116. doi: 10.1007/s11104-017-3189-x.

Singh, S. P., Keller, B., Gruissem, W., and Bhullar, N. K. (2017b). Rice NICOTIANAMINE
SYNTHASE 2 expression improves dietary iron and zinc levels in wheat. *Theor. Appl. Genet.*
130(2), 283-292. doi: 10.1007/s00122-016-2808-x.

Velu, G., Singh, R. P., Huerta-Espino, J., Peña, R.J., Arun, B., Mahendru-Singh, A., et al. (2012).
Performance of biofortified spring wheat genotypes in target environments for grain zinc and iron
concentrations. *Field Crop Res.* 137, 261-267. doi: 10.1016/j.fcr.2012.07.018.

Velu, G., Singh, R. P., Huerta, J., and Guzmán, C. (2017). Genetic impact of Rht dwarfing genes on
grain micronutrients concentration in wheat. *Field Crop Res.* 214(Supplement C), 373-377. doi:
10.1016/j.fcr.2017.09.030.

Wang, S., Wang, Z. H., Gao, Y. J., Liu, L., Yu, R., Jin, J. J., et al. (2017). EDTA alone enhanced soil
zinc availability and winter wheat grain Zn concentration on calcareous soil. *Environ. Exp. Bot.* 141,
19-27. doi: 10.1016/j.envexpbot.2017.06.008.

Wang S., Wang Z. H., Li S. S., Diao C. P., Liu L., Hui X. L., et al. (2018). Identification of high-
yield and high-Zn wheat cultivars for overcoming "yield dilution" in dryland cultivation. *Eur. J.
Agron.* 101, 57-62. doi: 10.1016/j.eja.2018.08.008.

Wessells, K. R., and Brown, K. H. (2012). Estimating the Global Prevalence of Zinc Deficiency:
Results Based on Zinc Availability in National Food Supplies and the Prevalence of Stunting. *Plos
One* 7(11), e50568-e50568. doi: 10.1371/journal.pone.0050568.

Xue, Y. F., Yue, S. C., Zhang, Y. Q., Cui, Z. L., Chen, X. P., Yang, F. C., et al. (2012). Grain and
shoot zinc accumulation in winter wheat affected by nitrogen management. *Plant Soil* 361(1-2), 153-
163. doi: 10.1007/s11104-012-1510-2.

Xue, Y. F., Zhang, W., Liu, D. Y., Yue, S. C., Cui, Z. L., Chen, X. P., et al. (2014). Effects of
nitrogen management on root morphology and zinc translocation from root to shoot of winter wheat
in the field. *Field Crop Res.* 161(0), 38-45. doi: 10.1016/j.fcr.2014.01.009.

Zhang, Y. H., Zhang, Y. P., Liu, N., Su, D., Xue, Q. W., Stewart, B. A., et al. (2012). Effect of
source-sink manipulation on accumulation of micronutrients and protein in wheat grains. *J. Plant
Nutr. Soil Sci.* 175(4), 622-629. doi: 10.1002/jpln.201100224.

Zou, C. Q., Zhang, Y. Q., Rashid, A., Ram, H., Savasli, E., Arisoy, R. Z., et al. (2012).
Biofortification of wheat with zinc through zinc fertilization in seven countries. *Plant Soil* 361(1-2),
119-130. doi: 10.1007/s11104-012-1369-2.

12   Figure Legends

Figure 1. Yield variations of the 123 tested wheat cultivars in 2014 (A), 2015 (B), and 2016 (C). The
19 high-yield cultivars (red bar, 'H') were identified with their yields consistently higher than the
corresponding median of all cultivars in each year.

Figure 2. Relationships of grain Zn concentration to grain yield (A), straw Zn concentration (C),
shoot Zn uptake (B), and Zn harvest index (D) for the high-yield wheat cultivars over three
experimental years. Determination coefficients (*P*-values) indicate the relationships of grain Zn
concentration to these traits, and the significant regressions at $P < 0.05$ are shown by solid lines.

Figure 3. Variations of shoot Zn uptake (A), Zn harvest index (B), and straw Zn concentration (C) of
the high-yield cultivars over three experimental years. For each box plot, like the straw Zn
concentration in 2014, the five values from top to bottom are the number of original observations (19
cultivars × four replications = 76), the maximum (10.01), the median (4.41, dashed line), the average
(3.97, solid line) and the minimum (1.33), respectively.

Figure 4. Grain Zn concentration (Zn.c, orange bar) and its biofortification potential (Zn.p, green bar)
of the 19 high-yield wheat cultivars in 2014 (A), 2015 (B), and 2016 (C). Error bars stand for
standard error (n=4).

Figure 5. The most suitable Zn biofortification measures in different scenarios of the current grain
Zn concentration (Zn.c), attainable grain Zn concentration (Zn.a), and Zn biofortification target level
(Zn.t). This is proposed as a guiding workflow to realize grain Zn biofortification target for high-
yield wheat.

13   Supplementary Material

Supplementary Figure 1. Relationships of wheat grain Zn concentration to the biomasses of glume
(A) and stem (B) and Zn concentrations of glume (C) and stem (D) for the 19 high-yield cultivars
over the three experimental years. Determination coefficients (*P*-value) indicate the relationship of
grain Zn concentration to each variable.

Supplementary Table 2. Tested wheat cultivars in the three-year field experiment, with the
identified 19 high-yield ones in red color.

Supplementary Table 2. Analysis of variance (ANOVA) for the effects of year, cultivar, and year $\times$
cultivar interaction on the different traits of high-yield cultivars.

Traits	High-yield group		
	Year	**Cultivar**	**Year $\times$ Cultivar**
Grain yield	20.18 (P = 0.002)	0.97 (P = 0.507)	0.72 (P = 0.865)
Glume biomass	19.7 (P = 0.002)	4.41 (P < 0.001)	0.83 (P = 0.734)
Stem biomass	21.07 (P = 0.002)	1.54 (P = 0.114)	0.86 (P = 0.694)
Straw biomass	26.13 (P = 0.001)	1.33 (P = 0.207)	0.82 (P = 0.749)
Glume Zn concentration	0.47 (P = 0.646)	1.70 (P = 0.069)	0.90 (P = 0.628)
Stem Zn concentration	1.93 (P = 0.226)	1.32 (P = 0.216)	0.77 (P = 0.812)
Straw Zn concentration	1.51 (P = 0.294)	1.27 (P = 0.242)	0.72 (P = 0.864)
Shoot Zn uptake	13.9 (P = 0.006)	2.33 (P = 0.009)	1.49 (P = 0.059)
Zn harvest index	5.94 (P = 0.038)	1.07 (P = 0.406)	1.43 (P = 0.084)
Grain Zn concentration (Zn.c)	29.63 (P = 0.001)	3.02 (P = 0.001)	2.89 (P < 0.001)
Grain Zn biofortification potential (Zn.p)	1.49 (P = 0.297)	1.45 (P = 0.148)	0.73 (P = 0.864)
Attainable grain Zn concentration (Zn.a)	12.06 (P = 0.008)	3.47 (P < 0.001)	1.85 (P = 0.008)