Title page

Quantify the requirements to achieve grain Zn biofortification of high-yield wheat on calcareous soils

Sen Wang [1, 2, 3], Zhaohui Wang [1, 2*], Shasha Li [1, 2], Chaopeng Diao [1, 2], Lu Liu [1, 2], Ning Huang [1, 2], Ming Huang [1, 2], Xiaoli Hui [1, 2],

Laichao Luo [1, 2], Gang He [1, 2], Hanbing Cao [1, 2]

[1] State Key Laboratory of Crop Stress Biology for Arid Areas, Northwest A&F University, Yangling 712100, Shaanxi, China

[2] Key Laboratory of Plant Nutrition and Agri-environment in Northwest China, Ministry of Agriculture and Rural Affairs,

College of Natural Resources and Environment, Northwest A&F University, Yangling 712100, Shaanxi, China

[3] Shenzhen Branch, Guangdong Laboratory for Lingnan Modern Agriculture, Genome Analysis Laboratory of the Ministry of

Agriculture and Rural Affairs, Agricultural Genomics Institute at Shenzhen, Chinese Academy of Agricultural Sciences,

Shenzhen, 518120, Guangdong, China

[*] Correspondence:

Prof. Zhaohui Wang,

Tel: +86 29 8708 2234;

Fax: +86 29 8708 2234.

E-mail: w-zhaohui@263.net.

Declarations of interest: none

Abstract: The solution to address global human Zn deficiency is Zn biofortification of staple food crops, aimed at high grain Zn concentration as well as high yield. However, the desired high grain Zn concentration above 40 mg kg^{-1} is rarely observed for high-yield wheat on worldwide calcareous soils, due to inadequate Zn uptake or Zn distribution to grain. The present study aims to investigate how much Zn uptake or distribution is adequate to achieve the Zn.t of high-yield wheat on calcareous soils with low available Zn ($\sim$ 0.5 mg kg^{-1}). Of the 123 cultivars tested in a three-year field experiment, 19 high-yield cultivars were identified with similar yields around 7.0 t ha^{-1} and various grain Zn concentrations from 9.3 to 26.7 mg kg^{-1}. The adequate Zn distribution to grain was defined from the view of Zn biofortification, as the situation where the Zn distribution to grain (Zn harvest index) increased to the observed maximum of $\sim$ 91.0% and the Zn concentration of vegetative parts (straw Zn concentration) decreased to the observed minimum of $\sim$ 1.5 mg kg^{-1} (Zn.m). Under the assumed condition of adequate Zn distribution to grain ($\sim$ 91.0%), all the extra Zn above Zn.m was remobilized from straw to grain and the grain Zn concentration would be increased to its highest attainable level, which was 14.5 $\sim$ 31.3 mg kg^{-1} for the 19 high-yield cultivars but still lower than 40 mg kg^{-1}. Thus, even with the adequate Zn distribution to grain, the current Zn uptake is still not adequate and needs to be increased to 308 g ha^{-1} or higher to achieve Zn.t for high-yield wheat (7.0 t ha^{-1}) on low-Zn calcareous soils. Besides, the established method here can also provide the priority measures and quantitative guidelines to achieve Zn biofortification in other wheat production regions.

Keywords: wheat; grain Zn biofortification; Zn distribution; Zn uptake; quantification.

1. Introduction

Although over 50 years have passed since the first case of human zinc (Zn) deficiency (Prasad, 2013), now this problem is still afflicting over 16% of the world population, mostly in Africa and South Asia (Kumssa et al., 2015). In these areas, residents cannot get sufficient Zn intake from daily diets which are mainly low-Zn cereals like wheat, maize, and rice (Wessells and Brown, 2012). Increasing the Zn concentration of cereal grains, known as Zn biofortification, is the main solution to alleviate human Zn deficiency, and the target of Zn fortified wheat grain is 40-50 mg kg^{-1} (Cakmak 2018), much higher than the current worldwide grain Zn concentration of 20-30 mg kg^{-1} (Liu et al., 2014; Chen et al., 2017). To close the gap between the current and target level, it is necessary to fully understand the factors affecting grain Zn concentration.

Grain yield, as an indicator of sink capacity, has essential influence on grain Zn concentration. For instance, 45% decrease in yield of field-grown wheat generated 60% increase in grain Zn concentration (Zhang et al., 2012). The negative correlation between grain Zn concentration and yield has been widely reported in different locations (Liu et al., 2014) and wheat germplasms (Guttieri et al., 2015), and the lowered grain Zn concentration in modern cultivars is mainly caused by the wide adoption of high-yield cultivars after the Green Revolution (Fan et al., 2008; Velu et al., 2017). To feed 9.7 billion population by 2050, the staple crop production has to increase by 60% (FAO, 2014), and thus crop Zn biofortification should increase yield and grain Zn concentration simultaneously (Bouis and Welch, 2010). To achieve high yield and high grain Zn concentration for wheat, it is necessary to enhance both shoot Zn uptake and its distribution to grain (Wang et al., 2018), which are the two sources of Zn accumulated in grain (Kutman et al., 2012; Xue et al., 2012). For example, wheat grain Zn concentration was increased by enhanced nitrogen (N) supply (Kutman et al., 2011; Hui et al., 2017) or ectopic-expressed rice *NICOTIANAMINE SYNTHASE 2* gene (Singh et al., 2017b), because abundant N-containing ligands like nicotianamine (NA) in the phloem enhanced the Zn transportation to grain (Barunawati et al., 2013). Also, the enhanced root Zn uptake increased wheat grain Zn concentration by enlarged root contact area with soil (Ercoli et al., 2017; Singh et al., 2017a), external Zn fertilization (Saha et al., 2017; Liu et al., 2017) or the mobilization of soil intrinsic Zn (Wang et al., 2017). However, most of existing Zn biofortification studies pay much less attention to grain yield, and thus cannot fully achieve the target of Zn biofortification.

The most cost-effective approach to realize crop Zn biofortification is developing cultivars with high yield and high grain Zn concentration (Gregory et al., 2017). For example, a few breeding lines or cultivars of spring wheat exhibited high grain Zn concentration ~ 40 mg kg^{-1} and high yield ~ 4 t ha^{-1}, under soil available Zn of 1.9 mg kg^{-1} in Canada, India, Pakistan, and Mexico (Gao et al., 2011; Velu et al., 2012). Similarly, under soil available Zn of 1.2 mg kg^{-1} in Iran, several bread wheat genotypes showed high grain Zn concentration > 50 mg kg^{-1} and high yield > 6 t ha^{-1} (Amiri et al., 2015). Thus, under relatively high soil available Zn > 1.0 mg kg^{-1}, breeding new wheat cultivars can achieve high grain Zn concentration and high yield simultaneously. But on calcareous soils with low available Zn ~ 0.5 mg kg^{-1}, the success of Zn biofortification of high-yield wheat is rarely achieved. For instance, the highest grain Zn concentration of spring wheat and bread wheat lines was only 25 mg kg^{-1} under soil available Zn of 0.3 ~ 0.6 mg kg^{-1} in Portugal and Iran (Gomez-Coronado et al., 2016; Khoshgoftarmanesh et al., 2013). In the present study on calcareous soils with available Zn of 0.4 mg kg^{-1} in northwest China, the grain Zn concentrations of 123 wheat cultivars were all below 30 mg kg^{-1}. Does this mean that high grain Zn concentration of 40 mg kg^{-1} cannot be achieved by breeding for high-yield wheat on calcareous soils?

The increase of grain Zn concentration for high-yield wheat relies on both high Zn uptake and high Zn distribution to grain (Wang et al., 2018). In the present work, the Zn harvest index varied from 45.5% to 94.0% and the straw Zn concentration at maturity varied from 1.2 to 10.5 mg kg^{-1}, while the minimum Zn concentration in wheat vegetative parts was ~ 5.0 mg kg^{-1} in hydroponics (Kutman et al. 2012), indicating that some Zn in straw might be still transported to grain. Accordingly, inadequate Zn distribution to grain can be the reason for the low grain Zn concentration observed for high-yield wheat. Then, from the view of Zn biofortification, how much Zn distribution to grain is adequate for field-grown high-yield wheat? With adequate Zn distribution, whether the Zn uptake is adequate to achieve high grain Zn concentration (> 40 mg kg^{-1}) for high-yield wheat? To test these hypotheses, the Zn concentration, uptake, and distribution of 123 wheat cultivars over three years were analyzed in the present study.

2. Materials and methods

2.1 Plant materials and field experiment

The field experiment was conducted in Yongshou (108°12′E, 34°44′N) located on the southern Loess Plateau, where the climate is the temperate continental monsoon climate and over half of the annual precipitation (~ 550 mm) occurs in the summer fallow period from July to September. The top 20 cm soil properties were determined according to Bao (2000) as follows: pH 8.4 (CO_2-free water), cation exchange capacity 20.3 cmol kg^{-1}, soil organic matter 12.9 g kg^{-1}, total N 0.9 g kg^{-1}, mineral N 26.6 mg kg^{-1}, available phosphorus (P) 16.9 mg kg^{-1}, available potassium (K) 123.4 mg kg^{-1}, available iron (Fe) 7.5 mg kg^{-1}, available manganese (Mn) 18.1 mg kg^{-1}, available copper (Cu) 1.3 mg kg^{-1}, and available Zn 0.4 mg kg^{-1}.

One hundred and twenty-three wheat cultivars were collected (Table S1) and tested in the three-year field experiment with a randomized complete block design. Each cultivar had four replications and each plot consisted of four 200 cm long rows with 2.5 cm seed spacing and 20 cm row spacing, under the nutrient supply of 150 kg N ha^{-1} and 100 kg P_2O_5 ha^{-1}. All cultivars were sown during 28th to 30th September in 2013, 2nd to 3rd October in 2014, and 26th to 28th September in 2015, and harvested during 14th to 17th June in 2014, 18th to 20th June in 2015, and 18th to 20th June in 2016. The rainfalls during summer fallow periods and growing seasons were 271 and 267 mm in 2013-2014, 317 and 314 mm in 2014-2015, and 228 and 186 mm in 2015-2016, respectively. Throughout the experimental period, no irrigation was conducted, and herbicide and pesticide were used when necessary.

2.2 Sampling and chemical analyses

At maturity, the plants of 30 ears were randomly sampled from the central two rows of each cultivar plot, and the roots were cut off at the stem-root joint part. Then, the shoots were air-dried and separated into stems, glumes, and grains, which were washed with deionized water and oven-dried at 65℃ to determine dry weight. After that, the oven-dried samples were ground by a ball mill (Retsch MM400, Germany) for chemical analyses. The remaining ears in the central two rows of each plot were also harvested and weighted, plus the above grain weight, to estimate the grain yield, glume biomass, and stem biomass in oven-dried weight.

Plant Zn concentration was determined as Ozbek and Akman (2016) with some modifications. Briefly, 0.2 g sample was mixed with 5.0 ml HNO_3 (65%) in a 50 ml Teflon tube and predigested at 120℃ for 0.5 h, and then 1.0 ml H_2O_2 (30%) was

added before microwave digestion (MW Pro, Anton Paar, Austria). Each sample was digested in two technical duplicates and

the standard wheat flour (GBW10011 GSB-2) was used for quality control. The digestion solution was diluted with ultrapure

water (18.25 MΩ cm^{-1}), and Zn concentration was determined by ICP-MS (iCAP Qc, Thermo Fisher Scientific, USA). The

measured Zn isotope was ^{66}Zn, and ^{73}Ge was added as internal standard to calibrate the signal fluctuation.

2.3 Quantification of the potential increase of grain Zn concentration by adequate Zn distribution

From the perspective of Zn biofortification, the adequate Zn distribution to grain refers to the situation where nearly all the

mobilizable Zn in shoot vegetative parts was transported or remobilized to grain, and correspondingly, the ratio of grain Zn

uptake to shoot Zn uptake (Zn harvest index) increased to the maximum and the Zn concentration of vegetative parts (straw Zn

concentration, Zn.s) decreased to the minimum. To determine the adequate Zn distribution to grain for high-yield wheat, we

selected the cultivars with yields higher than the corresponding median of all cultivars in each year, calculated the Zn uptake and

Zn harvest index, and combined stem and glume Zn concentrations into Zn.s by the weighted mean method. In each year, the

observed maximum Zn harvest index or minimum straw Zn concentration (Zn.m) was used to indicate the level of adequate Zn

distribution to grain. Since Zn concentration was easier to measure than Zn harvest index, we used Zn.m to calculate the potential

increase of grain Zn concentration (Zn.p) caused by the extra Zn remobilized from straw to grain under the condition of adequate

Zn distribution to grain.

$$\text{Zn uptake (g ha}^{-1}) = \frac{\text{Zn concentration (mg kg}^{-1}) \times \text{yield or biomass(kg ha}^{-1})}{1000}$$

$$\text{Zn.s (mg kg}^{-1}) = \frac{\text{stem Zn uptake(g ha}^{-1}) + \text{glume Zn uptake(g ha}^{-1})}{\text{stem biomass(kg ha}^{-1}) + \text{glume biomass(kg ha}^{-1})} \times 1000$$

$$\text{Zn.p (mg kg}^{-1}) = \frac{[\text{Zn.s (mg kg}^{-1}) - \text{Zn.m (mg kg}^{-1})] \times \text{Bm.s (kg ha}^{-1})}{\text{GrY (kg ha}^{-1})}$$

Here Bm.s and GrY refer to straw biomass and grain yield. The sum of Zn.p and the current grain Zn concentration (Zn.c)

was defined as the attainable grain Zn concentration (Zn.a).

Shoot Zn uptake and Zn harvest index (ZnHI) were calculated as described by Kutman et al. (2011) and Xue et al. (2012).

Shoot Zn uptake = grain Zn uptake + glume Zn uptake + stem Zn uptake

$$\text{Zn harvest index (\%)} = \frac{\text{Grain Zn uptake (g ha}^{-1})}{\text{Shoot Zn uptake (g ha}^{-1})} \times 100$$

2.4 Statistical analyses and graphing

Analysis of variance (ANOVA) was used to test the effects of year, cultivar, and year × cultivar interaction on yield or

biomass, Zn concentrations, and Zn uptakes. The relationships of grain Zn concentration with other traits were tested by Pearson

correlation or linear regression. All the statistical analyses were completed by PROC GLM, PROC CORR or PROC REG in

SAS v 8.01, with 0.05 set as the significance level. Scatter and bar plots were created by Sigmaplot 12.5, and the concept chart

illustrating Zn biofortification guidelines was created in Adobe Illustrator CC.

3. Results

3.1 Relationship of grain Zn concentration with other traits for high-yield wheat

Of the 123 tested wheat cultivars ordered by yield from low to high, 19 cultivars consistently exhibited higher yields than

the median in each year, and were identified as high-yield cultivars (Fig. 1). Although yield and biomasses showed significant

differences among three years, the 19 high-yield cultivars exhibited similar yields ($\sim$ 7 t ha^{-1}) and varied greatly in grain Zn

concentration, from 14.7 to 26.7 mg kg^{-1} in 2014, 9.3 to 22.2 mg kg^{-1} in 2015, and 13.5 to 17.0 mg kg^{-1} in 2016 (Table S2 and

Fig. 2a). Besides, grain Zn concentration showed no relation with yield, the biomass and Zn concentration of stem and glume

(Fig. 2a and Fig. S1). However, shoot Zn uptake was positively correlated with grain Zn concentration in each year and Zn

harvest index was positively correlated with grain Zn concentration in 2015 (Fig. 2c and 2d).

3.2 Minimum straw Zn concentration and maximum Zn harvest index

Shoot Zn uptake and Zn harvest index showed large variations among the 19 high-yield cultivars or three experimental

years (Table S2, Fig. 3a and 3b). The observed maximum Zn harvest index was 91.8% in 2014, 87.7% in 2015, and 94.0% in

2016 (Fig. 3b), while the observed minimum straw Zn concentration (Zn.m) was 1.3 mg kg^{-1} in 2014, 1.9 mg kg^{-1} in 2015, and

1.2 mg kg^{-1} in 2016 (Fig. 3c). With Zn.m used to indicate the level of adequate Zn distribution to grain, the gap between current

straw Zn concentration and Zn.m was used to calculate the amount of extra Zn which could be remobilized into grain.

3.2 Potential increase of grain Zn concentration and attainable grain Zn concentration

The potential increase of grain Zn concentration (Zn.p) due to extra Zn remobilized from straw to grain of the 19 high-yield

cultivars ranged from 2.1 to 5.5 mg kg^{-1} in 2014, 3.0 to 6.5 mg kg^{-1} in 2015, and 3.5 to 7.5 mg kg^{-1} in 2016 (Fig. 4). When Zn.p

was added to the current grain Zn concentration (Zn.c), the attainable grain Zn concentration (Zn.a) varied from 17.0 to 31.3 mg

kg^{-1} in 2014, 14.5 to 27.5 mg kg^{-1} in 2015, and 17.6 to 22.8 mg kg^{-1} in 2016 (Fig. 4), lower than the Zn biofortification target

(Zn.t) of 40 mg kg^{-1}. Thus, even with the adequate Zn distribution to grain, the current shoot Zn uptake was not adequate to

ahieve Zn.t of high-yield wheat. For an assumed high-yield ($\sim$ 7 t ha^{-1}) wheat cultivar with adequate Zn distribution to grain ($\sim$

91.0%), the required shoot Zn uptake should be at least 308 g ha^{-1} to achieve the Zn.t.

4.   Discussion

4.1 Using the minimum straw Zn concentration to estimate the adequate Zn distribution to grain

In the present work, the adequate Zn distribution to grain is defined from the view of Zn biofortification, and it refers to the

situation where all the mobilizable Zn in shoot vegetative parts was transported or remobilized to grain, meaning that Zn harvest

index increased to its maximum and straw Zn concentration decreased to its minimum (Zn.m). Then, the gap between current

straw Zn concentration and Zn.m can be used to calculate the amount of extra remobilized Zn from straw to grain, which results

in the potential increase and the attainable highest grain Zn concentration under specific conditions. For high-yield bread wheat

grown on low-Zn calcareous soils, the observed minimum straw Zn concentration is used as Zn.m which is $\sim$ 1.5 mg kg^{-1} and

lower than the reported 5.0 mg kg^{-1} for durum wheat in hydroponics (Kutman et al., 2012). Presumably, the senescent vegetative

parts of field-grown wheat need less Zn to constitute structural components and more Zn can be remobilized into grain. Under

uniform environmental conditions, straw Zn concentration and Zn.m can be used to determine whether the Zn distribution to

grain is adequate or not to promote Zn biofortification. Since the Zn biofortification of high-yield wheat needs both high Zn

uptake and high Zn distribution to grain (Wang et al., 2018), the Zn concentration of vegetative parts, which is easier to measure

than the Zn harvest index, deserves to be considered in future studies on wheat Zn biofortification.

4.2 Quantify the requirements to achieve wheat Zn biofortification in different environments

169   Based on the adequate Zn distribution to grain ($\sim$ 91.0%) as indicated by Zn.m, we can get the attainable highest grain Zn

170   concentration (Zn.a) and compare it with the grain Zn biofortification target (Zn.t, 40 mg kg^{-1}), which can tell us whether the

current Zn uptake is adequate to achieve Zn.t for high-yield wheat ($\sim$ 7 t ha^{-1}). Besides, the established quantitative analysis method here are also suitable for other wheat production regions with the aim of increasing the current grain Zn concentration (Zn.c) and grain yield. The priority measures to achieve the Zn.t of high-yield wheat are different under different conditions (Fig. 5). In wheat planting areas where Zn.a is close to or higher than Zn.t and the soil available Zn is often relatively high (> 1.0 mg kg^{-1}), the priority to realize Zn.t is to adopt or develop the cultivars with high Zn distribution to grain. For example, in North China Plains with soil available Zn of 3.4 mg kg^{-1}, optimal N application generated high yield of 7.5 t ha^{-1} and shoot Zn uptake of 370 g ha^{-1} (Xue et al., 2014), and the Zn.a would be 44.9 mg kg^{-1} and higher than Zn.t, indicating the necessity to introduce high-Zn-distribution wheat cultivars. In wheat production regions where Zn.a is lower than Zn.t and higher than Zn.c and the soil available Zn is often relatively low (< 0.5 mg kg^{-1}), developing high-Zn-distribution cultivars should be combined with agronomic measures like Zn fertilization to promote Zn uptake. For instance, in Iran rainfed drylands with soil available Zn of 0.2 mg kg^{-1} (Norouzi et al., 2014), the tested wheat cultivars exhibited high yield of 6.6 t ha^{-1} and shoot Zn uptake of 174 g ha^{-1}. Even if the Zn.c of 17.0 mg kg^{-1} was increased to the Zn.a of 26.4 mg kg^{-1} by introducing new cultivars, there was still a large gap to Zn.t which needed to be closed by increasing Zn uptake. In wheat production regions where Zn.a is close to Zn.c and lower than Zn.t, the Zn distribution to grain has reached its maximum and the priority measure of Zn biofortification is to increase Zn uptake by agronomic practices. However, this case is rarely observed up to now because researchers have not paid enough attention to the Zn in vegetative parts in wheat Zn biofortification studies.

On calcareous soils with low available Zn < 0.5 mg kg^{-1}, the current wheat grain Zn concentration is around 20 mg kg^{-1}, as reported in China, Kazakhstan, Mexico, Turkey, and Zambia, etc. (Zou et al., 2012; Liu et al., 2014). In the present study on calcareous soils with available Zn of 0.4 mg kg^{-1}, the wheat cultivar *Zhengmai 7698* had the highest Zn.a of 31.3 mg kg^{-1} and the yield of 6.6 t ha^{-1} in 2014, which was still lower than the Zn.t. To achieve high grain Zn concentration of 40 mg kg^{-1} and high yield of 7.0 t ha^{-1} simultaneously, the current shoot Zn uptake should be increased to at least 308 g ha^{-1} by agronomic measures like soil Zn fertilization with the wheat cultivars with high Zn distribution to grain ($\sim$ 91%). These guidelines deserve to be tested in field studies on Zn biofortification.

5. Conclusion

From the view of crop Zn biofortification, the adequate Zn distribution to grain can be defined as the case where Zn harvest index increased to its maximum and straw Zn concentration decreased to its minimum. For the high-yield ($\sim$ 7 t ha^{-1}) wheat grown on low-Zn ($\sim$ 0.5 mg kg^{-1}) calcareous soils, the maximum Zn harvest index was $\sim$ 91.0% and the minimum straw Zn concentration was $\sim$ 1.5 mg kg^{-1}. Under the condition of adequate Zn distribution to grain, the gap between straw Zn concentration and its minimum could be used to determine the extra Zn remobilized to grain and the highest attainable grain Zn concentration, which was 14.5 $\sim$ 31.3 mg kg^{-1} for the high-yield wheat cultivars and lower than the grain Zn biofortification target (Zn.t) of 40 mg kg^{-1}. Therefore, even with the adequate Zn distribution to grain ($\sim$ 91%), the current shoot Zn uptake is still not adequate to achieve the Zn.t of high-yield wheat grown on low-Zn calcareous soil, and the priority measure of Zn biofortification is to increase Zn uptake to 308 g ha^{-1} or higher by agronomic practices like Zn fertilization. The established method here can also provide the most suitable guidelines and quantitative requirements to achieve grain Zn biofortification in other wheat production regions.

Acknowledgements

The authors thank for the financial supports from the China Agricultural Research System (CARS-3), the Special Fund for Agro-scientific Research in the Public Interest under Grant (201303104), the Agricultural Scientific Research Talent and Team Program, the National Natural Science Foundation of China (NSFC) (41571282, 41401330), and the China Postdoctoral Science Foundation (2019M650919). During the wheat germplasm collection, we thank for the selfless help from the staff in different research laboratories and experimental stations of the China Agricultural Research System. We also thank the professors and doctors of the Soil Fertility and Plant Nutrition Team of Northwest A&F University, for their critical comments and constructive suggestions.

References

1. Amiri, R., Bahraminejad, S., Sasani, S., Jalali-Honarmand, S., and Fakhri, R. (2015). Bread wheat genetic variation for grain's protein, iron and zinc concentrations as uptake by their genetic ability. Eur. J. Agron. 67, 20-26. doi:

10.1016/j.eja.2015.03.004.

2.   Bao, S. D. (2000). Soil and agricultural chemistry analysis (in Chinese). Beijing: China Agriculture Press.

3.   Barunawati, N., Giehl, R. F. H., Bauer, B., and von Wiren, N. (2013). The influence of inorganic nitrogen fertilizer forms

on micronutrient retranslocation and accumulation in grains of winter wheat. Front. Plant Sci. 4, 320. doi:

10.3389/fpls.2013.00320.

4.   Bouis, H. E., and Welch, R. M. (2010). Biofortification-A Sustainable Agricultural Strategy for Reducing Micronutrient

Malnutrition in the Global South. Crop Sci. 50(2), S20-S32. doi: 10.2135/cropsci2009.09.0531.

5.   Cakmak, I., and Kutman, U.B. (2018). Agronomic biofortification of cereals with zinc: a review. Eur. J. Soil Sci. 69(1),

172-180. doi: doi:10.1111/ejss.12437.

6.   Chen, X. P., Zhang, Y. Q., Tong, Y. P., Xue, Y. F., Liu, D. Y., Zhang, W., et al. (2017). Harvesting more grain zinc of wheat

for human health. Sci. Rep. 7(1), 7016. doi: 10.1038/s41598-017-07484-2.

7.   Crespo-Herrera, L. A., Govindan, V., Stangoulis, J., Hao, Y., and Singh, R. P. (2017). QTL Mapping of Grain Zn and Fe

Concentrations in Two Hexaploid Wheat RIL Populations with Ample Transgressive Segregation. Front. Plant Sci. 8, 1800.

doi: 10.3389/fpls.2017.01800.

8.   Drahoňovský, J., Száková, J., Mestek, O., Tremlová, J., Kaňa, A., Najmanová, J., et al. (2016). Selenium uptake,

transformation and inter-element interactions by selected wildlife plant species after foliar selenate application. Environ.

Exp. Bot. 125, 12-19. doi: 10.1016/j.envexpbot.2016.01.006.

9.   Ercoli, L., Schüßler, A., Arduini, I., and Pellegrino, E. (2017). Tabbita. Plant Soil 419(1), 153-167. doi: 10.1007/s11104-

017-3319-5.

10.   Erdman, J. (1979). Oilseed phytates: nutritional implications. J. Am. Oil Chem. Soc. 56(8), 736-741.

11.   Fan, M. S., Zhao, F. J., Fairweather-Tait, S. J., Poulton, P. R., Dunham, S. J., and Mcgrath, S. P. (2008). Evidence of

decreasing mineral density in wheat grain over the last 160 years. J. Trace Elem. Med. Bio. 22(4), 315–324. doi:

10.1016/j.jtemb.2008.07.002.

12. FAO (2014). "Building a common vision for sustainable food and agriculture: principles and approaches". (Rome: FAO).

13. Gao, X., Mohr, R. M., McLaren, D. L., and Grant, C. A. (2011). Grain cadmium and zinc concentrations in wheat as affected

by genotypic variation and potassium chloride fertilization. Field Crop Res. 122(2), 95-103. doi: 10.1016/j.fcr.2011.03.005.

14. Gomez-Coronado, F., Poblaciones, M. J., Almeida, A. S., and Cakmak, I. (2016). Zinc (Zn) concentration of bread wheat

grown under Mediterranean conditions as affected by genotype and soil/foliar Zn application. Plant Soil 401(1), 331-346.

doi: 10.1007/s11104-015-2758-0.

15. Gregory, P. J., Wahbi, A., Adu-Gyamfi, J., Heiling, M., Gruber, R., Joy, E. J. M., et al. (2017). Approaches to reduce zinc

and iron deficits in food systems. Glob. Food Secur. 15, 1-10. doi: 10.1016/j.gfs.2017.03.003.

16. Grüter, R., Costerousse, B., Bertoni, A., Mayer, J., Thonar, C., Frossard, E., et al. (2017). Green manure and long-term

fertilization effects on soil zinc and cadmium availability and uptake by wheat (Triticum aestivum L.) at different growth

stages. Sci. Total Environ. 599-600, 1330-1343. doi: 10.1016/j.scitotenv2017.05.070.

17. Guttieri, M. J., Baenziger, P. S., Frels, K., Carver, B., Arnall, B., and Waters, B. M. (2015). Genc. Crop Sci. 55(3), 1035-

1052. doi: 10.2135/cropsci2014.07.0506.

18. Hamnér, K., Weih, M., Eriksson, J., and Kirchmann, H. (2017). Influence of nitrogen supply on macro- and micronutrient

accumulation during growth of winter wheat. Field Crop Res. 213, 118-129. doi: http://dx.doi.org/10.1016/j.fcr.2017.08.002.

19. Hui, X. L., Wang, Z. H., Luo, L. C., Ma, Q. X., Wang, S., Dai, J., et al. (2017). Winter Wheat Grain Yield and Zn

Concentration Affected by Long-Term N and P Application in Dryland (in Chinese). Sci. Agric. Sin. 50(16), 3175-3185.

doi: 10.3864/j.issn0578-1752.2017.16.012.

20. Jin, J. J., Wang, Z. H., Dai, J., Wang, S., Gao, Y. J., Cao, H. B., et al. (2014). Effects of long-term N and P fertilization with

different rates on Zn concentration in grain of winter wheat (in Chinese). J. Plant Nutr. Fertil. 20(06), 1358-1367. doi: 10.

11674/zwyf. 2014. 0605.

21. Khoshgoftarmanesh, A. H., Schulin, R., Chaney, R.L., Daneshbakhsh, B., and Afyuni, M. (2010). Micronutrient-efficient

genotypes for crop yield and nutritional quality in sustainable agriculture. A review. Agron. Sustain. Dev. 30(1), 83-107.

doi: 10.1051/agro/2009017.

22. Khoshgoftarmanesh, A. H., SanaeiOstovar, A., Sadrarhami, A., and Chaney, R. (2013). Effect of tire rubber ash and zinc

sulfate on yield and grain zinc and cadmium concentrations of different zinc-deficiency tolerance wheat cultivars under

field conditions. Eur. J. Agron. 49, 42-49. doi: 10.1016/j.eja.2013.02.013.

23. Kumssa, D. B., Joy, E. J., Ander, E. L., Watts, M. J., Young, S. D., Walker, S., et al. (2015). Dietary calcium and zinc

deficiency risks are decreasing but remain prevalent. Sci. Rep. 5, 10974. doi: 10.1038/srep10974.

24. Kutman, U. B., Kutman, B.Y., Ceylan, Y., Ova, E. A., and Cakmak, I. (2012). Contributions of root uptake and

remobilization to grain zinc accumulation in wheat depending on post-anthesis zinc availability and nitrogen nutrition. Plant

Soil 361(1-2), 177-187. doi: 10.1007/s11104-012-1300-x.

25. Kutman, U. B., Yildiz, B., and Cakmak, I. (2011). Effect of nitrogen on uptake, remobilization and partitioning of zinc and

iron throughout the development of durum wheat. Plant Soil 342(1-2), 149-164. doi: 10.1007/s11104-010-0679-5.

26. Liu, D. Y., Zhang, W., Pang, L. L., Zhang, Y. Q., Wang, X. Z., Liu, Y. M., et al. (2017). Effects of zinc application rate and

zinc distribution relative to root distribution on grain yield and grain Zn concentration in wheat. Plant Soil 411(1), 167-178.

doi: 10.1007/s11104-016-2953-7.

27. Liu, H., Wang, Z. H., Li, F. C., Li, K. Y., Yang, N., Yang, Y. E., et al. (2014). Grain iron and zinc concentrations of wheat

and their relationships to yield in major wheat production areas in China. Field Crop Res. 156, 151-160. doi:

10.1016/j.fcr.2013.11.011.

28. Norouzi, M., Khoshgoftarmanesh, A. H., and Afyuni, M. (2014). Zinc fractions in soil and uptake by wheat as affected by

different preceding crops. Soil Sci. Plant Nutr. 60(5), 670-678. doi: 10.1080/00380768.2014.937674.

29. Ozbek, N., and Akman, S. (2016). Method development for the determination of calcium, copper, magnesium, manganese,

iron, potassium, phosphorus and zinc in different types of breads by microwave induced plasma-atomic emission

spectrometry. Food Chem. 200, 245-248. doi: 10.1016/j.foodchem.2016.01.043.

30. Prasad, A. S. (2013). Discovery of Human Zinc Deficiency: Its Impact on Human Health and Disease. Adv. Nutr. 4(2), 176-

190. doi: 10.3945/an.112.003210.

31.   Saha, S., Chakraborty, M., Sarkar, D., Batabyal, K., Mandal, B., Murmu, S., et al. (2017). Rescheduling zinc fertilization

and cultivar choice improve zinc sequestration and its bioavailability in wheat grains and flour. Field Crop Res. 200**,** 10-17.

doi: 10.1016/j.fcr.2016.09.006.

32.   Singh, D., Rajawat, M. V. S., Kaushik, R., Prasanna, R., and Saxena, A. K. (2017a). Beneficial role of endophytes in

biofortification of Zn in wheat genotypes varying in nutrient use efficiency grown in soils sufficient and deficient in Zn.

Plant Soil 416(1)**,** 107-116. doi: 10.1007/s11104-017-3189-x.

33.   Singh, S. P., Keller, B., Gruissem, W., and Bhullar, N. K. (2017b). Rice NICOTIANAMINE SYNTHASE 2 expression

improves dietary iron and zinc levels in wheat. Theor. Appl. Genet. 130(2)**,** 283-292. doi: 10.1007/s00122-016-2808-x.

34.   Soltani, S., Khoshgoftarmanesh, A. H., Afyuni, M., Shrivani, M., and Schulin, R. (2014). The effect of preceding crop on

wheat grain zinc concentration and its relationship to total amino acids and dissolved organic carbon in rhizosphere soil

solution. Biol. Fert. Soils 50(2)**,** 239-247. doi: 10.1007/s00374-013-0851-1.

35.   Tabbita, F., Pearce, S., and Barneix, A. J. (2017). Breeding for increased grain protein and micronutrient content in wheat:

Ten years of the GPC-B1 gene. J. Cereal Sci. doi: 10.1016/j.jcs.2017.01.003.

36.   Velu, G., Singh, R. P., Huerta-Espino, J., Peña, R.J., Arun, B., Mahendru-Singh, A., et al. (2012). Performance of biofortified

spring wheat genotypes in target environments for grain zinc and iron concentrations. Field Crop Res. 137, 261-267. doi:

10.1016/j.fcr.2012.07.018.

37.   Velu, G., Singh, R. P., Huerta, J., and Guzmán, C. (2017). Genetic impact of Rht dwarfing genes on grain micronutrients

concentration in wheat. Field Crop Res. 214(Supplement C)**,** 373-377. doi: 10.1016/j.fcr.2017.09.030.

38.   Wang, S., Wang, Z. H., Gao, Y. J., Liu, L., Yu, R., Jin, J. J., et al. (2017). EDTA alone enhanced soil zinc availability and

winter wheat grain Zn concentration on calcareous soil. Environ. Exp. Bot. 141**,** 19-27. doi:

10.1016/j.envexpbot.2017.06.008.

39.   Wang S., Wang Z. H., Li S. S., Diao C. P., Liu L., Hui X. L., et al. (2018). Identification of high-yield and high-Zn wheat

cultivars for overcoming "yield dilution" in dryland cultivation. Eur. J. Agron. 101, 57-62. doi: 10.1016/j.eja.2018.08.008.

40. Wang, Z. M., Liu, Q., Pan, F., Yuan, L. X., and Yin, X. B. (2015). Effects of increasing rates of zinc fertilization on phytic

acid and phytic acid/zinc molar ratio in zinc bio-fortified wheat. Field Crop Res. 184, 58-64. doi: 10.1016/j.fcr.2015.09.007.

41. Wessells, K. R., and Brown, K. H. (2012). Estimating the Global Prevalence of Zinc Deficiency: Results Based on Zinc

Availability in National Food Supplies and the Prevalence of Stunting. Plos One 7(11), e50568-e50568. doi:

10.1371/journal.pone.0050568.

42. Xue, Y. F., Yue, S. C., Zhang, Y. Q., Cui, Z. L., Chen, X. P., Yang, F. C., et al. (2012). Grain and shoot zinc accumulation in

winter wheat affected by nitrogen management. Plant Soil 361(1-2), 153-163. doi: 10.1007/s11104-012-1510-2.

43. Xue, Y. F., Zhang, W., Liu, D. Y., Yue, S. C., Cui, Z. L., Chen, X. P., et al. (2014). Effects of nitrogen management on root

morphology and zinc translocation from root to shoot of winter wheat in the field. Field Crop Res. 161(0), 38-45. doi:

10.1016/j.fcr.2014.01.009.

44.   Zhang, W., Liu, D. Y., Liu, Y. M., Cui, Z. L., Chen, X. P., and Zou, C. Q. (2016). Zinc uptake and accumulation in winter

wheat relative to changes in root morphology and mycorrhizal colonization following varying phosphorus application on

calcareous soil. Field Crop Res. 197, 74-82. doi: 10.1016/j.fcr.2016.08.010.

45. Zhang, Y. H., Zhang, Y. P., Liu, N., Su, D., Xue, Q. W., Stewart, B. A., et al. (2012). Effect of source-sink manipulation on

accumulation of micronutrients and protein in wheat grains. J. Plant Nutr. Soil Sci. 175(4), 622-629. doi:

10.1002/jpln.201100224.

46. Zou, C. Q., Zhang, Y. Q., Rashid, A., Ram, H., Savasli, E., Arisoy, R. Z., et al. (2012). Biofortification of wheat with zinc

through zinc fertilization in seven countries. Plant Soil 361(1-2), 119-130. doi: 10.1007/s11104-012-1369-2.

 Figures

330 Fig. 1 Yield variations of the 123 tested wheat cultivars in 2014 (a), 2015 (b), and 2016 (c). The 19 high-yield cultivars (red bar,

331 'H') were identified with their yields consistently higher than the corresponding median of all cultivars in each year.

Fig. 2 Relationships of grain Zn concentration to grain yield (a), straw Zn concentration (c), shoot Zn uptake (b), and Zn harvest index (d) for the high-yield wheat cultivars over three experimental years. Determination coefficients (*P*-values) indicate the relationships of grain Zn concentration to these traits, and the significant regressions at *P* < 0.05 are shown by solid lines.

337 Fig. 3 Variations of shoot Zn uptake (a), Zn harvest index (b), and straw Zn concentration (c) of the high-yield cultivars over

338 three experimental years. For each box plot, like the straw Zn concentration in 2014, the five values from top to bottom are the

339 number of original observations (19 cultivars × four replications = 76), the maximum (10.01), the median (4.41, dashed line),

340 the average (3.97, solid line) and the minimum (1.33), respectively.

Fig. 4 Grain Zn concentration (Zn.c, orange bar) and its biofortification potential (Zn.p, green bar) of the 19 high-yield wheat

cultivars in 2014 (a), 2015 (b), and 2016 (c). Error bars stand for standard error (n=4).

345 Fig. 5 The most suitable Zn biofortification measures in different scenarios of the current grain Zn concentration (Zn.c), attainable

346 grain Zn concentration (Zn.a), and Zn biofortification target level (Zn.t). This is proposed as a guiding workflow to realize grain

347 Zn biofortification target for high-yield wheat.

 Supplementary figures

350 Fig. S1 Relationships of wheat grain Zn concentration to the biomasses of glume (a) and stem (b) and Zn concentrations of glume

351 (c) and stem (d) for the 19 high-yield cultivars over the three experimental years. Determination coefficients (*P*-value) indicate

352 the relationship of grain Zn concentration to each variable.

 Supplementary tables

354 Table S1 Tested wheat cultivars in the three-year field experiment, with the identified 19 high-yield ones in red color.

Code	Cultivar	Released year	Code	Cultivar	Released year	Code	Cultivar	Released year
001	*Heng 4399*	2008	042	*Huaimai 25*	2007	083	*Zhoumai 27*	2011
002	*Hengguan 35*	2004	043	*Wanmai 68*	2010	084	*Wenmai 6*	2000
003	*Heng 136*	2009	044	*Shannong 20*	2010	085	*Aikang 58*	2005
004	*Gaoyou 2018*	2008	045	*Huaimai 22*	2007	086	*Zhoumai 22*	2007
005	*Han 5093*	2010	046	*Yannong 5158*	2007	087	*Pingan 5*	2011
006	*Han 6172*	2001	047	*Yannong 21*	2002	088	*Baofeng 7228*	2000
007	*Shimai 15*	2005	048	*Su 553*	2011	089	*Guomai 0319*	2009
008	*Jimai 32*	1992	049	*Shiyou 20*	2009	090	*Hemai 1*	2006
009	*Jimai 22*	1985	050	*Linmai 4*	2006	091	*Zhengnong 17*	2004
010	*Liangxing 99*	2004	051	*Liangxing 66*	2008	092	*Zhoumai 18*	2004
011	*Jinmai 90*	2011	052	*Hemai 17*	2011	093	*Yangfumai 2*	2003
012	*Yaomai 16*	2011	053	*Liangxing 77*	2010	094	*Emai 580*	2012
013	*Yunhan 20410*	2007	054	*Linmai 2*	2004	095	*Xiangmai 25*	2008
014	*Lin Y8159*	2013	055	*Shannong 15*	2006	096	*Emai 596*	2009
015	*Yunhan 618*	2010	056	*Weimai 8*	2003	097	*Zhengmai 9023*	2001
016	*Yunhan 719*	2009	057	*Yannong 0428*	2008	098	*Chuanmai 104*	2012
017	*Yunhan 805*	2011	058	*Qingnong 2*	2010	099	*Xikemai 2*	2005
018	*Yunhan 21-30*	2003	059	*Yannong 999*	2011	100	*Neimai 11*	2007
019	*Yunhan 22-33*	2005	060	*Jimai 22*	2006	101	*Neimai 836*	2004
020	*Luohan 6*	2006	061	*Shannong 14*	2006	102	*Chuanmai 107*	2000
021	*Luohan 7*	2007	062	*Heima 1*	2004	103	*Linmai 6*	2003
022	*Luohan 9*	2009	063	*Luyuan 502*	2011	104	*Yunmai 56*	2008
023	*Luohan 11*	2008	064	*Taishan 22*	2004	105	*Shaan 509*	2011
024	*Luohan 13*	2009	065	*Jinan 17*	1999	106	*Zhongmai 895*	2012
025	*Changmai 251*	2011	066	*Tainong 18*	2008	107	*Xiaoyan 22*	2003
026	*Changmai 6697*	2008	067	*Luomai 4*	2003	108	*Xinong 3517*	2008
027	*Shimai 19*	2009	068	*Zhoumai 16*	2002	109	*Xinong 223*	2012
028	*Triticale*	1980	069	*Zhongyu 10*	2007	110	*Xiaoyan 15*	1995
029	*Yangmai 13*	2002	070	*Jimai 20*	2003	111	*9418*	2000
030	*Ningmai 13*	2005	071	*Xiangmai 969*	2006	112	*Shaanmai 139*	2011
031	*Ningmai 14*	2006	072	*Yumai 58*	2001	113	*Xinong 9871*	2008
032	*Xinong 979*	2005	073	*Yumai 18-99*	2003	114	*Changhan 58*	2004
033	*Huamai 5*	2010	074	*Yanzhan 4110*	2003	115	*Changwu 521*	2008
034	*Sumai 188*	2012	075	*Rumai 0319*	2009	116	*Chang 134*	1997
035	*Taikong 6*	2003	076	*Pumai 9*	2004	117	*Jinmai 47*	1995
036	*Shenmai 22*	2010	077	*Zhoumai 26*	2012	118	*Chang 6359*	2005
037	*Xinfumai 1*	2007	078	*Xuke 1*	2007	119	*Bei 1*	2013
038	*An 0817*	2016	079	*Xinmai 26*	2010	120	*Bei 4*	2013
039	*Longping 203*	2013	080	*Fanmai 8*	2008	121	*Bei 9*	2013
040	*Yannong 19*	2001	081	*Zhoumai 24*	2009	122	*Xikemai 6*	2008
041	*Xinmai 8*	2011	082	*Zhengmai 7698*	2011	123	*Yumai 13*	2010

355 Table S2 Analysis of variance (ANOVA) for the effects of year, cultivar, and year × cultivar interaction on the different traits of
356 high-yield cultivars.

Traits	High-yield group		
	Year	Cultivar	Year × Cultivar
Grain yield	20.18 (P = 0.002)	0.97 (P = 0.507)	0.72 (P = 0.865)
Glume biomass	19.7 (P = 0.002)	4.41 (P < 0.001)	0.83 (P = 0.734)
Stem biomass	21.07 (P = 0.002)	1.54 (P = 0.114)	0.86 (P = 0.694)
Straw biomass	26.13 (P = 0.001)	1.33 (P = 0.207)	0.82 (P = 0.749)
Glume Zn concentration	0.47 (P = 0.646)	1.70 (P = 0.069)	0.90 (P = 0.628)
Stem Zn concentration	1.93 (P = 0.226)	1.32 (P = 0.216)	0.77 (P = 0.812)
Straw Zn concentration	1.51 (P = 0.294)	1.27 (P = 0.242)	0.72 (P = 0.864)
Shoot Zn uptake	13.9 (P = 0.006)	2.33 (P = 0.009)	1.49 (P = 0.059)
Zn harvest index	5.94 (P = 0.038)	1.07 (P = 0.406)	1.43 (P = 0.084)
Grain Zn concentration (Zn.c)	29.63 (P = 0.001)	3.02 (P = 0.001)	2.89 (P < 0.001)
Grain Zn biofortification potential (Zn.p)	1.49 (P = 0.297)	1.45 (P = 0.148)	0.73 (P = 0.864)
Attainable grain Zn concentration (Zn.a)	12.06 (P = 0.008)	3.47 (P < 0.001)	1.85 (P = 0.008)

357 Data in table are ANOVA F values (Pr > F).