High-Resolution, Multidimensional Phylogenetic Metrics Identify Class I Aminoacyl-tRNA Synthetase Evolutionary Mosaicity and Inter-modular `Coupling

Charles W. Carter, Jr[1], Alex Popinga[2], Remco Bouckaert[2], and Peter R. Wills[3]

[1]Department of Biochemistry and Biophysics, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599-7260

ORCID ID (Carter): /0000-0002-2653-4452

[2]Centre for Computational Evolution, University of Auckland, PB 92019, Auckland 1142, New Zealand

ORCID ID (Bouckaert): /0000-0001-6765-3813

[3]Department of Physics and Te Ao Marama Centre for Fundamental Inquiry, University of Auckland, PB 92019, Auckland 1142, New Zealand

ORCID ID (Wills): /0000-0002-2670-7624

*Corresponding author

Email: carter@med.unc.edu

Running Title: Mosaic Ancestry of Class I Aminoacyl-tRNA Synthetases

Keywords: BEAST2, DensiTree, protein mosaic structure, RNA World hypothesis

Abbreviations: aaRS, aminoacyl-tRNA synthetase; CP1, Connecting peptide 1; MSA, multiple sequence alignment; MCMC, Markov chain monte carlo

23 Abstract

The provenance of the aminoacyl-tRNA synthetases (aaRS) poses unusually challenging
questions because of their role in the emergence and evolution of genetic coding. We investigate
evidence about their ancestry from highly curated structure-based multiple sequence alignments
of a small "scaffold" that is structurally invariant in all 10 canonical Class I aaRS. Statistically
different values of two uncorrelated phylogenetic metrics—residue by residue conservation
derived from Clustal and row-by-row cladistic congruence derived from BEAST2—suggest that
the Class I scaffold is a mosaic assembled from distinct, successive genetic sources. These data are
especially significant in light of: (i) experimental fragmentations of the Class I scaffold into three
partitions that retain catalytic activities in proportion to their length; and (ii) multiple sources of
evidence that two of these partitions arose from an ancestral Class I aaRS gene encoding a Class
II ancestor in frame on the opposite strand. Two additional metrics output by BEAST2 vary in
accordance with the presumed functionality endowed by the various modules. The new evidence
supplements previous aaRS phylogenies. It identifies a previously characterized 46-residue Class
I "protozyme" as preceding the adaptive radiation of the superfamily containing variations of the
Rossmann dinucleotide binding fold related to amino acid discrimination, and thus as root of that
molecular tree. Such a rooting is consistent with near simultaneous emergence of genetic coding
and the origin of the proteome, resolving a conundrum posed by previous inferences that Class I
aaRS evolved long after the genetic code had been implemented in an RNA world. Further, it
establishes a timeline for the growth of coding from a binary amino acid alphabet by pinpointing
discontinuous enhancements of aaRS fidelity.

44 Author Summary

Phylogenetic analysis uncovers evolutionary connections between different protein superfamily
members. We describe complementary, uncorrelated, phylogenetic metrics that support multiple
evolutionary histories for different segments within members of the Class I aminoacyl-tRNA
synthetase superfamily. Using a carefully curated 3D crystal structure superposition as the
primary source of the multiple sequence alignment substantially reduced dependence of these
metrics on empirical amino acid substitution matrices. Two metrics are derived from the amino
acid distribution observed in each successive position. A third depends on how individual
sequences distribute into phylogenetic tree branches for each of the ten amino acids activated by
the superfamily. All metrics confirm that a segment previously identified as an inserted element
is, indeed, a more recent acquisition, despite its structural conservation. The residue-by-residue
conservation metrics reveal significant co-variation of mutational frequencies between a core
segment that forms the amino acid binding site and a neighboring segment derived from the more
recent insertion element. We attribute that covariation to the differentiation of superfamily
members as evolutionary divergence enhanced amino acid specificity. Finally, evidence that the
insertion element is a recent acquisition implies a new branching order for much of the proteome.

Introduction

The emergence of the aminoacyl-tRNA synthetases, aaRS, is a quintessential chicken and egg
puzzle whose solution would demystify the origins of coded protein synthesis. How did aaRS
enzymes gain the reflexive property of collectively being able to use relationships in the universal
genetic code to convert the sequences of base triplets in their own genes into functional amino acid
sequences that make the code work? The detailed trajectory by which genes for the two essential
superfamilies appeared, grew to acquire catalytic proficiencies and radiated to refine their dual
amino acid and cognate tRNA specificities is thus a crucial chapter in the book of life.

Both aaRS Classes have separate catalytic and anticodon-binding domains. Only the catalytic
domains within each superfamily share the same architectures [1]. In this paper, we develop and
analyze high-resolution phylogenetic metrics relevant to the mosaic evolutionary histories of those
shared architectures. Anticodon-binding domains are idiosyncratic and not considered here.

Phylogenetic analyses of the Class I aaRS superfamily [2-8] generally agree that clades for each
amino acid are monophyletic, divide into three subclasses [1, 9], including IleRS, ValRS, LeuRS,
and MetRS (Subclass IA); GluRS and GlnRS (Subclass IB); and TyrRS and TrpRS (Subclass IC).
Restricting analysis to the bacterial aaRS as they point more faithfully to the root of protein
synthesis [10], we exclude the archaeal Subclass IB Class I LysRS. CysRS and ArgRS families,
assigned originally to Subclass IA [9], are difficult to assign, with one or the other appearing
instead with Subclass IB [for example, see 5]. A parallel sub-classification in Class II aaRS
strengthens connections [11-13] between Class I and II phylogenies and the physical chemistry
differentiating the 10 amino acids in each Class.

Three segments within the Class I aaRS catalytic domains suggest a succession of
intermediate states dating from well before the Last Universal Common Ancestor.

Genetic deconstruction of Class I aaRS modules [14] identified a core whose ~130 residues form
a nearly intact active site, and which experiments show retain a major fraction of full-length aaRS
catalytic proficiency—estimated as the transition-state stabilization free energy—for both amino
acid and tRNA substrates [15-17] (Fig 1). Because of their high catalytic proficiency in both
essential reactions, these and similar cores derived from Class II aaRS were called "Urzymes",
from the German prefix "Ur" = original, authentic. Class I and II Urzymes differentiate between
amino acids from the two sets of substrates, with Class I Urzymes activating Class I, in preference
to Class II amino acids by ~1.0 kcal/mole, and conversely [18-20].

Even smaller 46 residue subsets from both Class I and II retain nearly half the full catalytic
proficiency in the critical amino-acid activation reaction [21], whose uncatalyzed rate is rate-
limiting for protein synthesis. These primitive modules were called "protozymes", from the Greek
προτο = first, to distinguish them from Urzymes. The Class I protozyme coincides with the N-
terminal crossover connection of the Rossmann fold, which contains a distinctive 3D packing
motif that recurs in ~25% of the proteome [22] and imposes distinct, functionally relevant
conformational states tightly coupled to catalysis in TrpRS [23-29].

Fully active Class I and II protozymes have been expressed from a single gene designed to encode
their structures on opposite strands. Experimental Michaelis-Menten parameters of all four
protozymes—Class I and II; native and bidirectional—are, remarkably, the same within
experimental error [21]. Experimental [21, 30, 31] and bioinformatic evidence [10] therefore

support the hypothesis of Rodin and Ohno [32, 33] that the two aaRS Classes descended from
opposite strands of a single bidirectional gene.

The progressive biochemical functionality of aaRS protozymes, Urzymes, and catalytic domains;
the fact that their structures are universally conserved within both Classes; and the evidence that
they descended from one bidirectional ancestral gene; all imply that they are legitimate
experimental models for ancestral evolutionary aaRS forms that participated throughout the
emergence of genetic coding and well before LUCA [14]. Moreover, to date, no such sequential
intermediates have been derived for other superfamilies. Thus, more appears to be known about
the modular evolution of the two aaRS superfamilies than for any other ancient superfamily,
making the Class I aaRS a highly relevant subject for this work.

A variable-length insertion interrupts Class I aminoacyl-tRNA synthetase catalytic cores.

Class I aaRS catalytic domains contain a variable-length insertion called "connecting peptide 1"
or CP1 [34, 35], which is detailed in Fig 1. CP1 insertions include the large editing domains of the
aaRS for aliphatic amino acids, and thus account for the variable size of Class I catalytic domains.
CP1 always intersects the Rossmann fold immediately after the Protozyme between structurally
homologous residues that are ~4.5 Å apart and can be replaced by a peptide bond without structural
disruption to produce the Class I Urzyme [30].

The scaffold identifies invariant secondary structures outside the Urzyme.

Evidence that the essential catalytic architectures observed in aaRS Urzymes arose from a
bidirectional gene implies, *ipso facto*, that sequences inconsistent with bidirectional coding
represent accretion of new genetic material. In this paper we address an apparent contradiction
between the Rodin-Ohno hypothesis and extended conservation into the CP1 segment.
Independent 3D superposition of aaRS crystal structures [2, 30, 36] revealed considerable
structural conservation within the catalytic domains of all ten members of each Class—see Fig 1
of [36]. Structural homology across the Class I superfamily extends beyond the Urzyme, into CP1.
We call the secondary structures within these conserved cores "scaffolds".

The CP1 insertions are incompatible with, and their introduction would necessarily have ended,
bidirectional genetic coding of ancestral Class I and II aaRS (Fig 1B). Class I Urzyme boundaries
delimited decisively by potential bidirectional coding of Class II Urzymes constitute only about
80% of the Class I scaffold; the remainder consists of 10-residue α-helical segments near the
beginning and end of the CP1 insertion (Fig 1). Thus, if the Rodin-Ohno hypothesis is correct,
then the CP1 insert must derive from a distinct genetic source.

Data summarized in the three previous sections lead us to expect metrics derived from MSAs for
modules with distinct ancestries to differ quantitatively. For these reasons, we developed new,
accurate ways to quantify the phylogenetic signatures of the protozyme, Urzyme, and CP1 aaRS
segments in order to assess the likelihood that CP1 represents a more recent acquisition by
ancestral Urzymes.

Novel applications of phylogenetics

This work introduces four novelties that, together, reinforce the conclusion that modular
components of the ancestral aaRS have different genetic histories:

(i) We threaded sequences into closely-related crystal structures to assemble multiple sequence
alignments (MSAs) based on three-dimensional structure superposition, to avoid depending on
amino acid substitution matrices to define equivalent positions.

(ii) We develop multi-dimensional phylogenetic metrics from the ensemble of phylogenetic trees
obtained by Markov Chain Monte Carlo (MCMC) simulations to compare segments of the MSA.
We show that these three metrics are uncorrelated, and demonstrate their statistical significance
by multiple regression.

(iii) We increase the effective analytical resolution by applying the metrics to partitions of the
MSA that have been extensively characterized experimentally, providing novel insight into the
functional modularity of the superfamily and molecular phylogenetic support that they are
successive evolutionary intermediates in the generation of the Class I aminoacyl-tRNA synthetase
superfamily.

(iv) We identify key two-way interactions between mutation rates in different MSA segments.
Both involve the amino acid binding site and one is central to the enhancement of amino acid
specificity enabled by the CP1 insertion.

These results suggest important modifications of conventional evolutionary scenarios for major
parts of the proteome containing Rossmann dinucleotide fold domains, and strengthen the proposal
that the genetic code development is coupled intimately to the structural evolution of aaRS.

Results

Phylogenetic metrics derived from trees constructed for different subset MSAs—CP1; the
Urzyme; and its three distinct modules Urz_A (the protozyme), Urz_B (the amino acid binding
site, and Urz_C (the PP_i binding site)—furnish unprecedented insight into the genetic modularity
of the Class I aaRS. In summary, they suggest that CP1 is derived from a substantially more recent
and less cladistically coherent genetic entity, as previously posited [30]. Similarly, they suggest
that the protozyme may be from an older genetic entity.

168 CP1 has lower and more uniform apparent site-to-site mutation rates between aaRS
169 for different amino acids.

The BEAST 2 MCMC algorithm tracks the extent of site-to-site evolutionary of sequence variation
in using the Tree height and Shape metrics described in Methods. CP1 sequences from the scaffold
exhibit much reduced mutation rates and variance, consistent with their being a more recent
evolutionary acquisition. Sequences within the Class I Urzymes exhibit higher and more variable
mutation rates, consistent with greater adaptive radiation, hence deeper ancestry than CP1
sequences.

The Tree height metric is summarized in (Fig 2A, B). Sequences responsible for the elevated
Urzyme Tree height are identified by regression against the independent parameters of the design
matrix (Table 1) in Fig 2C. Tree height $= 2.17 - 1.4 * Segment\ B + 3.2 * Urzyme - 2.7 *$
$protozyme + 1.7 * Segment\ B * protozyme$. All β coefficients are significant with P-values <
0.005. Foremost among the positive contributors is the Urzyme. However, the interaction between
the Protozyme (segment A) and segment B is also quite significant. We note that CP1 sequences
exhibit substantially smaller Tree height and variance (i.e. higher Shape), consistent with it being
a more recent genetic entity.

The negative logarithm of the Shape parameter is highly correlated with the conservation quality,

$\langle Q \rangle$, defined by Clustal [37, 38] (Fig 3A), lending intuitive insight into the physical meaning of

the latter. The designation "Conservation Quality" is apparently misleading in suggesting that the

significantly smaller $\langle Q \rangle$ value for the CP1 MSA (Fig 3B) implies that it is less well conserved,

in apparent conflict with its reduced Tree height. In fact, the colinearity of $-\log_{10}(\text{Shape})$ and $\langle Q \rangle$

led us to pursue the equivalence between the two metrics. Fig 3C, D show that the two metrics

depend in nearly identical fashion on predictor columns from the design matrix in Table 1. It

appears then that the $\langle Q \rangle$ metric does not measure mutational variation itself, but rather the inverse

of its variance.

Moreover, the extended similarity between Fig 3C and Fig 3D suggest that the site-by-site metrics
(Tree height and Shape) can provide quite high resolution evidence on the evolution of modularity.
Regression of Shape on the independent parameters of the design matrix in Table 1 (Fig 4) resulted
in the unique model: $\text{Shape} = 1.37 + 0.4 * Segment\ B - 1.08 * Segment\ C + 1.56 * CP1 -$
$0.57 * Segment\ B * CP1$. All β coefficients are significant and two have P-values < 0.005.

198   Class I Urzyme- and CP1-based trees form distinctly different clades.

Urzyme and CP1 partitions of the MSAs produce substantially different trees (Fig 5A). In
particular, although all ten Class I aaRS clades are monophyletic in the trees for the Urzyme MSAs,
three enzymatic clades in the CP1 trees—MetRS, ValRS, and TyrRS—are polyphyletic.
Moreover, the Urzyme clades are constrained by tight envelopes, whereas the CP1 envelopes are
poorly defined.

Support, S_i, the fraction of all trees for which each aaRS clade, i, is monophyletic, was averaged

over all Class I aaRS types to give the mean support, $\langle S \rangle = \sum S_i / 10$. That operation was repeated

for trees built for the full scaffold MSA, (Full), the Urzyme, CP1, protozyme (segment A in Fig
1), and seven subset MSAs each consisting of twenty amino acids in blocks of five residues as

described in Methods. For the eleven rows of the design matrix (Table 2), we computed $\langle S \rangle$ for

populations of trees constructed using the conventional WAG [39] substitution matrix and repeated
using the more recent LG [40] substitution matrix used in [8].

Contributors to the variance of $\langle S \rangle$ were assessed using stepwise multiple regression, which

resulted in the unique model: $\langle S \rangle = 0.67 + 0.04 * protozyme - 0.33 * CP1 + 0.1 *$

$protozyme * CP1$ summarized in Fig 5. All three coefficients are significant at better than the
99% confidence level, with P-values < 0.005. As suggested by its position in the regression curve
in Fig 5, trees built from CP1 residues have significantly less support than those from the Urzyme
or any of its 20-residue subsets. The A segment coincides with the protozyme MSA; its dominant

impact on the variance of $\langle S \rangle$, contributing positively both via its intrinsic effect and by its two-

way interaction with CP1, is consistent with its being close to the root of the Class I aaRS
superfamily tree.

Thus, although the 3D structures from the Urzyme and CP1 partitions have comparable structural
homology, they have markedly different phylogenetic signatures.

Phylogenetic metrics—Tree height, Shape, and Support—reveal significant, high-resolution genetic mosaicity.

Our quantitative evidence probes far deeper into evolutionary time than previous phylogenetic analyses of Class I aaRS [3, 7, 41-43]. That depth both calls for caution and is a source of great interest. Constructing aaRS trees is fundamentally ambiguous because at each node in any conceivable tree, the coding alphabet and dimension of the operational substitution matrix necessarily both change by integer steps as Class I and II trees branch into multiple families. Trees for the two superfamilies are thus necessarily interdependent, so that the dimensions of all possible substitution matrices start from 2 and end at 20. Thus, it is uncertain what should be inferred from phylogenetic metrics for a single superfamily.

These difficulties are substantially offset by the consistency of site-by-site and row-by-row metrics with the construction and experimental characterization of aaRS protozymes [21] and Urzymes [14, 15, 18, 19, 24, 25, 31] (Fig 1). Such deep evolutionary intermediates are, at present, manifestly unique to the aaRS. The extensive experimental and structural context of that consistency strengthens our conclusions even without comparable analysis of Class II aaRS, now in progress, especially in light of the following observations.

The three metrics are uncorrelated. The CP1 MSA is a substantial outlier for all three metrics, suggesting that the metrics may be correlated. Removing the CP1 entries from the design matrix eliminates any correlation between Tree height, Shape, or $\langle S \rangle$, Table 2. The three types of metrics are therefore essentially uncorrelated and provide independent insights.

Differences between Urzyme and CP1 sequences are statistically meaningful. If the Class I aaRS sequence partitions compared in Table 1 were all drawn from continuously replicated ancient genetic sources, subject to comparable selection history since their emergence, the null hypotheses would produce similarly conserved sequences and comparably congruent trees for the Urzyme and CP1 partitions. The log-worth values (i.e., –log(P)) for the higher mutational frequency (Tree height; 2.7), its variance (Shape; 3.3) and lack of congruence for phylogenetic trees, ($\langle S \rangle$; 3.4), imply with high statistical significance that all metrics for CP1 and Urzyme sequences are drawn from different populations, corroborating the argument—based on bidirectional coding ancestry—that the CP1 sequences represent genetic information acquired more recently by the Urzymes.

The numbers of residues in different segments drawn from the MSA have no detectable impact on any phylogenetic metric. One might suppose that degraded congruence of the CP1 trees results from the fact its MSA has only ~0.25 as many amino acids as the Urzyme MSA. The insignificant impact of NUMB on the regression models for any score and the clustering of the 20-residue $\langle S \rangle$ values with the intact Urzyme MSA (Fig 5) confute that expectation.

Threading does not force any particular comparison between different aaRS types. Threading increased the reliability of structure-based alignments within any aaRS type by adding sequences. Scaffold positions were rigorously defined as structural homologs from the close proximity of their alpha carbon coordinates in multiple PDB structure alignments (i) among aaRS for any single amino acid, drawn from very diverse bacterial species, that only then (ii) produced a grand scaffold MSA across all Class I aaRS types. Additions produced by threading thus have only second-order effects on structural superpositions of aaRS for different amino acids, adding precision to our analysis without overtly influencing choices on which our conclusions depend.

Distinguishing features of CP1 sequences are evident without considering indels. Figs 2-5 together
illustrate a comprehensive, near-optimally quantitative three-dimensional comparison of partitions
in the Full Class I aaRS scaffold MSA. The crux of what the data suggest is that CP1 is more recent
than the Urzyme, its evolutionary divergence and variance—evidenced by its Tree height and
Shape—is much reduced, but its phylogenetic consistency—evidenced by $\langle S \rangle$—is much reduced,
relative to the corresponding metrics for Urzyme segments. This conclusion is evident, from
sequences with strict 1-1 correspondences between 3D crystal structures, excluding the large and
variable-length indels that dominate CP1 insertions in most aaRS types.

Neither convergent evolution nor horizontal gene transfer are likely explanations for the
Urzyme/CP1 distinction. The congruent clade structure of Urzyme-derived sequences from the
scaffold fall into consensus groupings characterizing Class I aaRS as a coherent superfamily and
does not require mechanisms beyond mutation and selection from a single common ancestor. The
aberrant behavior of sequence variations in the CP1 insertion (Figs. 2-5) might suggest appeal to
such processes. Treatments of Class I aaRS evolution based on full MSAs [6, 44, 45] show
evidence of horizontal gene transfer (HGT), genetic transpositions and large scale
insertion/deletion events, of which CP1 is the foremost example.

Wherever full-length bacterial Class I enzymes with a particular amino acid substrate specificity
are represented by more than one canonical structure, as has been described for IleRS and MetRS
[see Fig. 3 in 44], that bipartite distribution in genome space is adequately explained in terms of
early HGT into bacteria from an archaean/eukaryotic ancestor, but not in terms of convergent
evolution. CP1 sequence disparities at homologous sequence positions in TyrRS, MetRS, and
ValRS behave in the opposite manner: the higher variance of their $\langle Q \rangle$ values producing lower
$\langle S \rangle$ values by allowing their evolutionary paths to wander widely, often crossing in sequence
space, instead of forming multiple well-defined clades reasonably distant from one another in
sequence space that would suggest HGT.

Phylogenetic metrics have functionally relevant interpretations.

Regressions in Fig 3 were performed to demonstrate equivalence of the –log10(Shape) and the
Clustal $\langle Q \rangle$ metric. The dominant main effect of the Urzyme accounts for >60% of the variance
of the dependent variable, with the balance coming from the opposing effects of the B and C
subsets (i.e., their β coefficients are of opposite sign). Regression models of the three independent
metrics derived from BEAST 2 tree constructions all depend heavily on significant two-way
interactions between segments of the different MSAs. Functional properties of the Class I aaRS
modules are experimentally sufficiently well-characterized to sustain functional interpretations of
the interaction terms for site-to-site metrics (Figs 2, 4) and an evolutionary interpretation of the
row-by-row Support metric (Fig 5).

The relationship between Shape and segment B changes sign, depending on whether CP1 is
present in the MSA. CP1 forms an annulus around one end of the two halves of the Urzyme (Fig
6A). Molecular dynamic simulations of the TrpRS Urzyme [17] show that in these two segments,
which together form the amino acid binding site in Class I aaRS, exhibit extensive relative motion.
Further, the distance between the two parts of the amino acid binding site decreases in the TrpRS
catalytic conformational transition and that motion is coupled to the relative motion of CP1 and
the anticodon-binding domain. Moreover, steady state kinetic measurements of amino acid
specificity confirm that the relative domain motion is coupled to the relative specificity for
Tryptophan vs Tyrosine [24].

Thus, the interaction between CP1 and the amino acid binding site of the Urzyme constrains the
amino acid binding site dynamically. Structural relationships between CP1 and the amino acid
biniding site in TrpRS are highlighted in Fig 6B. The experimental demonstration of the interaction
between CP1 and the amino acid binding site validates the sign and strength of the B*CP1
contribution to Shape much as a pre-formed space in a partially assembled puzzle validates the
Shape of the missing piece.

The relationship between Tree height and segment B changes sign, depending on whether the
protozyme is present in the MSA. As noted in regard to Fig 2A,B, the Urzyme itself has a dominant
impact increasing the Tree height, and this effect is reduced substantially by the protozyme but
increased by the protozyme*Segment B interaction. Yet, sequences in Set 1 from within Segment
B have the highest Tree height. Another reflection of the same phenomenon is that the protozyme
and Urzyme appear in very much the same place on the regression plots in Figs 2 and 5, yet are
quite well separated in Fig 4, where the protozyme is midway between the Urzyme and CP1.

The strong interaction in the Tree height regression model is between the protozyme—locus of
ATP bnding—and segment B—locus of amino acid binding (Fig 6B, C). It is reasonable that the
interaction term arises because of the contravariant effects of the protozyme, which has retained
significant sequences over the evolution of the synthetases because it has the ATP binding site and
Segment B, which has the amino acid binding site, and would be expected to exhibit the most
significant evolutionary sequence variation between families specific for different amino acids.

The relationship between Support and CP1 changes sign, depending on whether the protozyme is
present in the MSA. Residues from the protozyme contribute even more decisively to the average
support than those derived from elsewhere in the Urzyme. CP1 has the most significant impact on
the regression model for the Support metric. Its coefficient is -0.33, more than three times that of
the next most important predictor. This effect can be seen in the regression plot in Fig 5, in which
the CP1 MSA is more widely separated from the other MSAs than in the models for any other
metric. However, the presence of the protozyme sequences in the Full MSA is sufficient to change
the impact of CP1 from negative to positive, giving the β-coefficient of +0.1 for the
protozyme*CP1 interaction term. This would appear to reinforce the conclusion, based also on the
widespread occurrence of the protozyme packing motif [22] and its functional activation of ATP
[21], that the protozyme was the original root of the entire superfamily and is older than the
Urzyme itself, as previously proposed [46].

Discussion

Our long-range goal is to understand how evolution of aaRS•tRNA cognate pairs effected the
stepwise increases in dimensionality that resulted in the progression of entries into the universal

genetic coding table. To that end we assembled three novel phylogenetic metrics here, to show
that they identify high-resolution mosaicity in carefully curated, structure-based MSAs of
secondary structures shared by all members of the Class I aaRS superfamily. All metrics support
prior arguments that the Class I aaRS evolved by a succession of intermediates—
protozyme=>Urzyme=>Catalytic domain—with increasingly sophisticated catalytic [14, 18, 19,
21, 31] and discriminatory [46-48] capabilities.

Evolutionary refinement of protein catalysis was predicated on expanding the genetic code.

The Tree height, Shape, and and Support metrics identify differences in the genetic origins of
successively acquired contributors to aaRS function: protozyme=>Urzyme=>catalytic domain
with CP1. They correlate inversely with experimentally measured catalytic proficiencies for the
catalysts from which the corresponding sequences are derived (Fig 7A). Those proficiencies
themselves correlate quantitatively with parallel increases in catalytic proficiency with the
concomitant additions of mass in the Class I and II aaRS evolutionary intermediates they entail
[See Fig. 6 in 14]. Both phylogenetic and functional properties of these intermediates thus appear
to probe far deeper into evolutionary time than previous phylogenetic analyses of Class I aaRS [3,
7, 41-43].

The precision with which protein active sites distinguish substrates from one another and transition
states from substrates was the result of the evolutionary process we wish to characterize. Expansion
of genetic coding itself depended critically on developing a system for the placement of a precisely
defined amino acid sidechain at a particular point on a peptide backbone. Phylogenetic analysis of
segmental Class I aaRS MSAs represents a uniquely promising opportunity to test the hypothesis
that contemporary enzymes are mosaic structures rooted in simpler catalytic polypeptides and
assembled from detectably different genetic ancestors.

High-resolution structural modularity implies functionally significant discontinuities in the evolution of genetic coding.

If CP1 insertions were indeed assimilated from one or more similar genetic sources after Class I
aaRS Urzymes had evolved significantly from modest ancestral protozymes, it could have at least
three noteworthy biological implications:

(i) Class I protozymes, whose high catalytic activity mobilizes ATP for biosynthesis, an activity
found in many proteins—may be the root of that major portion of the proteome built from β-α-β
crossover connections—the Rossmannoid protein superfamily [22] and potentially β-barrel
proteins [49, 50], which are central to intermediary metabolism and nucleotide biosynthesis [51].

(ii) AARS protozyme and Urzyme populations would have functioned first as quasispecies in
translation, limiting the sophistication of the early proteome.

(iii) CP1 assimilations would have transformed selection pressures for subsequent aaRS evolution
by facilitating enhanced fidelity.

Primordial aaRS quasispecies covered progressively smaller regions of sequence space, closer to
the "root". Early aaRS evolution cannot have been an ordinary mutation/selection process. All
solutions to the problem normally framed as a "chicken and egg problem" [52] imply historical
context and a continuity principle, which must be defined by the genotype-phenotype mapping
[53]: at any stage during the emergence of coding. Some prior system must have been interpreted
extant aaRS protogenes. A key aspect of such prior systems is that as the alphabet size, diversity,
and consequent fidelity increased, they would, *ipso facto*, have created more narrowly targeted
selection pressures, strongly coupling mutation and selection in early stages of genetic coding (Fig
7B). The subsequent coevolution of CP1 and Urzyme sequences would necessarily have preserved
Urzyme functionality, while the new CP1 could adapt flexibly to its developing role of enhancing
specificity as described in the next section.

Different background colors in Fig 7B denote how branching of the tree to allow introduction of
the n^{th} amino acid into the alphabet enforces a highly cooperative re-optimizing of the $n-1$ aaRS
types already present. As each new, refined amino acid type emerged, all extant gene sequences
adapted to opportunities introduced by progressively finer discrimination between amino acid side
chain physical chemistry. In turn, adaptation to a more diverse alphabet sharpens the new aaRS
fidelities (i.e., the contraction of branch thicknesses in Fig 7B), implicating a bootstrapping
feedback and enhancing the cooperativity of the transition to higher-dimension coding alphabets
[46]. That cooperativity creates a Lamarckian correlation between selection pressure and its
outcome—the result of any mutation being nearly synonymous with the selection pressure it faced,
especially as the code differentiated. Survival would have depended on the relationship between
the shapes of fitness landscapes and error rates of catalysis by the extant quasispecies [54]. The
earliest alphabets, including at least the first two amino acid types, were also tightly constrained
by bidirectional coding (gray shaded background, Fig 7B).

We have argued [46, 48] that a single functional island in sequence space (i.e. quasispecies) would
invariably have been a strong attractor, irrespective of the detailed Shape of the fitness landscape
that stabilized it, because all mutations that moved the system slightly off its optimum phenotype
would be subject to strongly restorative selection pressures. However, if ancestral protozymes
from a bidirectional gene had broad, relatively flat (and necessarily co-dependent) fitness
landscapes, matched to correspondingly high error rates, that could have favored bifurcated
quasispecies that enhanced genetic coding by recruiting new amino acid types and simultaneously
increasing the precision with which child specificities could be encoded.

Insertion of CP1 likely enabled saltatory improvements in fidelity. Structural and biochemical data
suggest that the CP1 insertions created stepwise enhancements in the evolution of genetic coding
by enabling conformationally-driven mechanisms to enhance specificity. The shortest CP1
insertions have ~75 residues in TrpRS and TyrRS that recur essentially intact in the longer CP1
insertions of the remaining eight Class I aaRS [30, 36]. It seems likely that the initial insertion
needed to be that long. CP1 forms an annulus around the Urzyme (Fig 6), constraining relative
movements of the protozyme and specificity helix that form the amino acid binding site [17]. For
that reason, the original hypothesis as to its origin was near-simultaneous insertion of a mobile
genetic element into all extant Class I Urzymes [30], in which case its root sequence would have
been more recent than that of the Urzymes, yet earlier than contemporary full-length aaRS
enzymes.

Amino acid specificities [18, 19] suggest that although capable of 10^9-fold rate accelerations, Class
I and II Urzymes select an amino acid from the correct Class only 80% of the time. However, Wills

& Carter [20] note that their within-Class specificity is consistent with each Class distinguishing
two kinds of amino acids, to operate a four-letter coding alphabet. These modest fidelities suggest
a fundamental limit to the precision of which bidirectional coding was ultimately capable.

The most evident contribution of CP1 to fidelity is that the editing domains present in the larger
subclass IA aaRS for aliphatic amino acids Ile, Val, and Leu are elaborations of the CP1 motif
present in the simplest subclass IC aaRS for Tyr and Trp. It seems likely, however, that CP1
functioned earlier to enhance fidelity by dynamically constraining the size and Shape of the amino
acid binding pocket. Several groups found that comprehensive mutation of side chains in the
immediate vicinity of the amino acid substrate, all within the Urzyme architecture, would not
change specificities of subclass IB GlnRS to Glu [55, 56] or subclass IC TrpRS to Tyr [57].
Changing GlnRS specificity to Glutamate (Bullock, et al. [56]) required wholesale mutations in
the second layer surrounding the amino acid binding site outside the Urzyme, but within in the
GlnRS CP1 domain.

Similarly, a modular thermodynamic cycle comparing specificities of full-length TrpRS, Urzyme
and Urzyme plus either CP1 or the anticodon binding domain (ABD) showed rejection of Tyrosine
by *G. stearothermophilus* TrpRS requires cooperation between CP1 and the ABD [24]. CP1 must
coordinate its movements with those of the ABD to perform the allosteric communication
necessary to enhance side chain selectivity beyond the modest capabilities of the Urzyme [25, 27].
In both cases, fine tuning specific recognition of amino acid substrates required insertion of CP1
and the ABD and, subsequently, coupling between them.

Inserting the CP1 motif would necessarily have disrupted bidirectional coding (Fig 1B), thus
dividing aaRS evolutionary histories decisively into distinct stages (Fig 7, between green and
yellow backgrounds). Selective advantages of CP1 insertion thus appear to have been to (i) end
constraints imposed by bidirectional coding and (ii) transcend the fundamental limitation on
specificity posed by the Urzyme architecture. Either or both would have allowed substantial
increases in fidelity to develop. CP1 therefore likely dramatically transformed the Class I aaRS
fitness landscape, and was likely necessary to expand the coding alphabet.

A revised branching order suggests that Class I aaRS protozymes and Urzymes root the
Rossmannoid superfamily. Putting Class I aaRS protozymes at the root of that superfamily
reconfigures many branching orders within the proteome to reflect that aaRS Urzymes were not a
late-developing branch in the Rossmannoid superfamily radiation, but instead were ancestral to it
(Fig 8). Descent of Class I and II aaRS from a single bidirectional ancestral gene [10, 14, 18, 19,
21, 32, 33] would underscore the likelihood that the aaRS of both families diverged, rather than
converging to similar functions from different sources. Thus, the genetic coding table itself likely
evolved by bifurcating pre-existing aaRS genes into specialized enzymes whose more
discriminating specificities for tRNA and amino acid substrates enabled daughter synthetases to
differentiate groups amino acids that previously had functioned as synonymous [46, 48, 58, 59]. It
would be surprising if other branches of the proteome had not diverged from the ancestral aaRS,
as suggested in Fig 6.

Phylogenetic trees have meaningful fine structure

The coherence of the phylogenetic metrics described in previous sections makes a strong case for
integrating complementary experimental data into a coherent application of Bayesian phylogenetic
analysis. Their functional significance emerges only in the context of partitioning the overall MSA
according to structural and functional criteria established with other disciplines and in the presence

of suitable controls for the effect of sequence length (subsets 0-6 from the Urzyme MSA; Fig 1C,
Table 1). The evidence that emerges for fine structure within protein domains is without precedent.

In turn, the phylogenetic analysis provides novel validation of the experimental work that guided
decisions on partitioning the MSA. Identification of how the Tree height and Shape metrics depend
on intermodular two-way interactions validates the considerable experimental work demonstrating
how intermodular coupling contributes to catalysis and specificity. Further, because these
interactions arise from the coherence of the entire Class I superfamily, they imply that much of
that experimental evidence applies, perforce, to similar interactions between ATP and amino acid
binding sites and between amino acid binding sites and CP1 in all or most Class I aaRS.

These observations suggest that this work points the way to more substantial applications of
software like BEAST2 [60] to a broader range of evolutionary questions.

Genetic coding and the proteome probably emerged from an RNA•polypeptide
partnership.

Previously accepted phylogenies of the Class I aaRS root well before the Last Universal Common
Ancestor (LUCA) [3, 7, 41-43], but appeared also to have radiated after earlier bifurcations in the
immense meta-family [3] containing folds based on the Rossmann dinucleotide-binding fold [61].
This paradoxically late radiation of protein assignment catalysts is the foremost phylogenetic
evidence favoring the RNA World hypothesis. Genetic coding by protein aaRS was therefore
thought, necessarily. to have replaced a prior implementation based on ribozymal assignment
catalysts [62, 63].

Evidence described here that CP1 is a much more recent acquisition by more ancestral aaRS
Urzymes supports a plausible alternative branching order (Fig 8). Attributing the apparently late
adaptive radiation of Class I aaRS to CP1 achieves consistency with an origin of genetic coding
from a bidirectional gene administering a binary coding alphabet in a peptide/RNA world.
Resolving the paradox in this way complements—without necessarily contradicting—the
phylogenetic analyses of Koonin [62, 63].

Methods

Amino acid chemistry underlies the metric form of amino acid substitution matrices (aaSM)
generally required for the logically circular process of creating an amino acid sequence alignment
by optimizing the constraints of some aaSM and then building a phylogenetic tree from the
alignment according to those constraints [8].

We superposed the available bacterial Class I aaRS crystal structures to base the multiple sequence
alignment (MSA) exclusively on precise three-dimensional structural homology, freeing the MSA
from dependence on empirical substitution matrices and estimates of relative rates of mutation
[64]. In the first place, this was done within each aaRS type. The identification of sequence
positions displaying very high conservation of both structure and amino acid occupancy provided
unambiguous anchors for the production of much more extensive sequence alignments. Popinga
[65] performed this task through extensive use of HHpred [66] and MODELLER [67] to thread
amino acid sequences for each aaRS type using experimentally determined 3D structures as
templates. This procedure produced, for each aaRS type, a greatly expanded MSA in which various
well-conserved secondary structural regions were near-perfectly structure- and sequence-aligned
between species known to have diverged not long after the LUCA.

In the final stage of alignment, the structures of the individual aaRS types were brought together
to identify regions of universal structural homology, the Class I aaRS "scaffold". The product was
an MSA across all Class I aaRS types in which each sequence position could be assumed to have
arisen, as far as is reasonably possible, from the same LUCA-ancestral codon. While the validity
of this assumption is by its very nature untestable, the close structural homology of Class I aaRS
of all types and the rigour of our procedure gives us confidence and justification as good as that
underlying any cross-species phylogenetic enterprise. However, we took further steps to restrict
and constrain the data upon which we later built putative phylogenies. First, individual scaffold
elements are more extensive in some extant aaRS types than others and it is impossible to identify
any proper universal homology among the loops, turns and structurally disordered regions that join
them. All of these were excluded from our grand Class I "scaffold" MSA. Second, we included
only eubacterial aaRS. It has been suggested previously [10] that bacterial aaRS sequences are
closer to the ancestral root than those from other domains. While analysis of loop structures
demonstrates that this is contestable in respect of the ages of biology's three main evolutionary
domains (ref Berezovsky? Lupas? Trifonov?) it is generally clear that cytoplasmic proteins of
bacteria have been subject to many fewer complex selection pressures than their archeal and
eukaryotic counterparts. The complex functional and regulatory roles played by some aaRS
proteins and their involvement in numerous genetic syndromes attests to this (ref).

Thus, the final MSA contained roughly bacterial 20 sequences for each of 10 Class I aaRS,
circumventing as many problems as possible in aligning sequences that diverged to produce
different substrate specificities in the pre-LUCA æon. The use of scaffold sequences for
phylogenetic analysis gave the best guarantee that results would reflect relatively neutral evolution
within the context of asymmetric, specialised selection pressures producing variant substrate
specificities by fine-tuning the size, Shape and chemistry of more intricately constructed amino
acid side-chain structures and pockets.

The MSA for the Class I scaffold was output using VMD [68], and is provided in the supplement
(Supplement_MSA_files.txt) together with all subsets used in this work. The scaffold fasta file
was then partitioned along lines of the experimental deconstruction of the Class I aaRS superfamily
[14] into the disjointed segments of the Urzyme, separated by structurally conserved segments
from CP1. Finally, because the Urzyme (83) and CP1 (20) partitions of the scaffold MSAs have
different sequence lengths, seven subsets comprising 20 sequence positions distributed throughout
the Urzyme were selected arbitrarily by a balanced, randomized procedure [69] to test the effect
of sequence length on our phylogenetic signatures.

We compiled a three-dimensional comparison of complementary metrics of the overall scaffold
MSA, the full Class I Urzyme, the protozyme, and CP1 subsets. BEAST 2 accumulates two metrics
from the ensemble of trees generated by Markov Chain Monte Carlo simulation. The Tree height
reflects the mean overall mutation rates, and site-specific mutation rates are fitted to a gamma
distribution [70] by adjusting a Shape parameter, $\alpha = 1/CV^2$, where CV is the coefficient of
variation or the ratio of the standard deviation to the mean and thus a measure of Tree height
variance.

For the third dimension, molecular kinships between rows of the MSA within each partition were
compared by clustering the sequences into clades according to their evolutionary origin.
Phylogenetic trees were computed using BEAST2 [60], allowing the use of multiple amino acid
substitution matrices (primarily WAG and LG). Trees were visualized with DensiTree [71] and
FigTree [72]. For the full Class I scaffold and each partition of interest, DensiTree was used to

extract ~10,000 trees generated by BEAST2 in building trees. From these we calculated values of
a parameter representing the support for clades that each corresponded exclusively to an individual
aaRS type, i. This support, S_i, is the percentage of all trees for which the aaRS type in question
appears to be monophyletic, meaning that the leaves of that aaRS type descend from the same most
recent branch point (common ancestor) with no descendants arising from a different aaRS type.

The mean phylogenetic support, $\langle S \rangle$, is a metric derived from a row-by-row calculation over all
10,000 phylogenetic trees generated using the sequence data for a chosen partition of the scaffold
MSA.

The conservation quality, Q_j, defined by Clustal [37, 38] was computed down each column, j, of
the grand MSA, which included all aaRS types. While the definition of Q seems convoluted, it has
been constructed as a metric whose value reflects the degree of amino acid diversity generated by
typical evolutionary amino acid substitution processes, reflected in the substitution matrix that it
uses as a reference. Different matrices do not give widely divergent Q parameters. We used the
ClustalW default matrix (PAM 250; [73]) and calculated the average, $\langle Q \rangle$, over all positions
within a partition of the MSA a parametric representation of the partition-wide variation in amino
acid occupancy calculated column-by-column over individual sequence positions.

Columns for various predictors were appended to the table of $\langle Q \rangle$ and $\langle S \rangle$ values to form the
design matrix (Table 2) used for multiple regression analysis with the JMP software [74]. Multiple
Regression was used to assess the statistical strength of contributions of the predictors (i.e., the
independent variables, A, B, C, CP1 defined in Fig 1 and the number of amino acids) to $\langle Q \rangle$ and
$\langle S \rangle$ values, using stepwise searches to identify the best set of predictors followed by least squares
estimation of the corresponding coefficients and their Student t–test P values.

Acknowledgments. This work was supported by NIGMS (Grant number R01-78227 to
C.W.C., Jr). We are grateful to Corbin Jones, Greg Fournier, Günter Wachtershäuser, and Bill
Martin for helpful feedback in preparing the manuscript and Loren Williams for suggesting that
ribosomal protein sequences may exhibit a similar genetic mosaicity.

References

1. Carter CW, Jr. Cognition, Mechanism, and Evolutionary Relationships in Aminoacyl-tRNA
Synthetases. Ann Rev Biochem. 1993;62:715-48.

2. O'Donoghue P, Luthey-Schulten Z. On the Evolution of Structure in Aminoacyl-tRNA
Synthetases. Microbiol Mol Biol Rev. 2003; 67(4):550–73.

3. Aravind L, Anantharaman V, Koonin EV. Monophyly of Class I Aminoacyl tRNASynthetase,
USPA, ETFP, Photolyase, and PP-ATPase Nucleotide-Binding Domains: Implication for
Protein Evolution in the RNAWorld. PROTEINS: Struct Funct Gen. 2002;48:1–14.

4. Leipe DD, Wolf YI, Koonin EV, Aravind L. Classification and Evolution of P-loop GTPases
and Related ATPases. J Mol Biol. 2002;317:41-72.

5. Roach JM, Sharma S, Kapustina M, Carter CW, Jr. Structure alignment via Delaunay
tetrahedralization. PROTEINS: Struct Funct Bioinf. 2005;60(1):66-81.

6. Wolf YI, Aravind L, Grishin NV, Koonin EV. Evolution of Aminoacyl-tRNA Synthetases—
Analysis of Unique Domain Architectures and Phylogenetic Trees Reveals a Complex History
of Horizontal Gene Transfer Events. Genome Research. 1999;9:689–710.

7. Caetano-Anollés G, Wang M, Caetano-Anollés D. Structural Phylogenomics Retrodicts the
Origin of the Genetic Code and Uncovers the Evolutionary Impact of Protein Flexibility. Plos
One. 2013;8(8):e72225. doi: 10.1371/journal.pone.0072225.

8. Shore J, Holland BR, Sumner JG, Nieselt K, Wills PR. The Ancient Operational Code is
Embedded in the Amino Acid Substitution Matrix and aaRS Phylogenies. J Mol Evol. 2019;
88:136–50. doi: 10.1007/s00239-019-09918-z.

9. Cusack S. Eleven down and nine to go. Nat Str Biol. 1995;2:824-31.

10. Chandrasekaran SN, Yardimci G, Erdogan O, Roach JM, Carter CW, Jr. Statistical Evaluation
of the Rodin-Ohno Hypothesis: Sense/Antisense Coding of Ancestral Class I and II
Aminoacyl-tRNA Synthetases. Molecular Biology and Evolution. 2013;30(7):1588-604. doi:
10.1093/molbev/mst070.

11. Carter CW, Jr., Wolfenden R. Acceptor-stem and anticodon bases embed amino acid chemistry
into tRNA. RNA Biology. 2016;13(2):145–51. doi: 10.1080/15476286.2015.1112488.

12. Carter CW, Jr., Wolfenden R. tRNA Acceptor-Stem and Anticodon Bases Form Independent
Codes Related to Protein Folding. Proc Nat Acad Sci USA. 2015;112 (24):7489-94. doi:
http://www.pnas.org/cgi/doi/10.1073/pnas.1507569112.

13. Wolfenden R, Lewis CA, Yuan Y, Carter CW, Jr. Temperature dependence of amino acid
hydrophobicities. Proc Nat Acad Sci USA. 2015;112 (24):7484-8. doi:
10.1073/pnas.1507565112.

14. Carter CW, Jr. Coding of Class I and II aminoacyl-tRNA synthetases. Advances in
Experimental Medicine and Biology: Protein Reviews. 2017;18:103-48. doi: DOI
10.1007/5584_2017_93.

15. Li L, Francklyn C, Carter CW, Jr. Aminoacylating Urzymes Challenge the RNA World
Hypothesis. J Biol Chem. 2013;288:26856-63. doi: 10.1074/jbc.M113.496125

16. Li L, Weinreb V, Francklyn C, Carter CW, Jr. Histidyl-tRNA Synthetase Urzymes: Class I
and II Aminoacyl-tRNA Synthetase Urzymes have Comparable Catalytic Activities for
Cognate Amino Acid Activation. J Biol Chem. 2011;286:10387-95. doi:
10.1074/jbc.M110.198929.

17. Pham Y, Kuhlman B, Butterfoss GL, Hu H, Weinreb V, Carter CW, Jr. Tryptophanyl-tRNA
synthetase Urzyme: a model to recapitulate molecular evolution and investigate intramolecular
complementation. J Biol Chem. 2010;285:38590-601. doi: 10.1074/jbc.M110.136911

18. Carter CW, Jr. What RNA World? Why a Peptide/RNA Partnership Merits Renewed
Experimental Attention. Life. 2015;5:294-320. doi: 10.3390/life5010294.

19. Carter CW, Jr., Li L, Weinreb V, Collier M, Gonzales-Rivera K, Jimenez-Rodriguez M, et al. The Rodin-Ohno Hypothesis That Two Enzyme Superfamilies Descended from One Ancestral Gene: An Unlikely Scenario for the Origins of Translation That Will Not Be Dismissed. Biology Direct. 2014;9:11.

20. Wills PR, Carter CW, Jr. Impedance matching and the choice between alternative pathways for the origin of genetic coding. International Journal of Molecular Sciences. 2020;21:7392. doi: 10.3390/ijms21197392.

21. Martinez L, Jimenez-Rodriguez M, Gonzalez-Rivera K, Williams T, Li L, Weinreb V, et al. Functional Class I and II Amino Acid Activating Enzymes Can Be Coded by Opposite Strands of the Same Gene. J Biol Chem. 2015;290(32):19710–25. doi: 10.1074/jbc.M115.642876

22. Cammer S, Carter CW, Jr. Six Rossmannoid Folds, Including the Class I Aminoacyl-tRNA Synthetases, Share a Partial Core with the Anticodon-Binding Domain of a Class II Aminoacyl-tRNA Synthetase. Bioinformatics. 2010;26(6):709-14. doi: 10.1093/bioinformatics/btq039.

23. Carter CW, Jr. Escapement mechanisms: efficient free energy transduction by reciprocally-coupled gating. Proteins: Structure, Function, and Bioinformatics. 2019;88:710–7. doi: 10.1002/prot.25856.

24. Weinreb V, Li L, Chandrasekaran SN, Koehl P, Delarue M, Carter CW, Jr Enhanced Amino Acid Selection in Fully-Evolved Tryptophanyl-tRNA Synthetase, Relative to its Urzyme, Requires Domain Movement Sensed by the D1 Switch, a Remote, Dynamic Packing Motif J Biol Chem 2014;289:4367-76. doi: 10.1074/jbc.M113.538660.

25. Li L, Carter CW, Jr. Full Implementation of the Genetic Code by Tryptophanyl-tRNA Synthetase Requires Intermodular Coupling. J Biol Chem. 2013;288(29 November):34736–45. doi: 10.1074/jbc.M113.510958.

26. Weinreb V, Li L, Carter CW, Jr. A Master Switch Couples Mg^{2+}-Assisted Catalysis to Domain Motion in B. stearothermophilus Tryptophanyl-tRNA Synthetase. Structure. 2012;20:128-38.

27. Chandrasekaran SN, Carter CW, Jr. Adding torsional interaction terms to the Anisotropic Network Model improves the PATH performance, enabling detailed comparison with experimental rate data Structural Dynamics. 2017;4:032103.

28. Chandrasekaran SN, Das J, Dokholyan NV, Carter CW, Jr. A modified PATH algorithm rapidly generates transition states comparable to those found by other well established algorithms. Structural Dynamics. 2016;3:012101. doi: 10.1063/1.4941599. PubMed Central PMCID: PMCPMC4769271.

29. Carter J, Charles W., Chandrasekaran SN, Weinreb V, Li L, Williams T, editors. Combining multi-mutant and modular thermodynamic cycles to measure energetic coupling networks in enzyme catalysis Structural Dynamics; 2016; American Crystallographic Association Annual Meeting: American Crystallographic Association.

30. Pham Y, Li L, Kim A, Erdogan O, Weinreb V, Butterfoss G, et al. A Minimal TrpRS Catalytic Domain Supports Sense/Antisense Ancestry of Class I and II Aminoacyl-tRNA Synthetases. Mol Cell. 2007;25:851-62.

31. Carter CW, Jr. Urzymology: Experimental Access to a Key Transition in the Appearance of
Enzymes. J Biol Chem. 2014;289(44):30213–20. doi: 10.1047/jbcR114.576495.

32. Rodin A, Rodin SN, Carter CW, Jr. On Primordial Sense-Antisense Coding. Journal of
Molecular Evolution. 2009;69:555-67.

33. Rodin SN, Ohno S. Two Types of Aminoacyl-tRNA Synthetases Could be Originally Encoded
by Complementary Strands of the Same Nucleic Acid. Orig Life Evol Biosph. 1995;25:565-
89.

34. Burbaum JJ, Schimmel P. Assembly of a Class I tRNA Synthetase from Products of an
Artificially Split Gene. Biochem. 1991;30:319-24.

35. Burbaum JJ, Starzyk RM, Schimmel P. Understanding Structural Relationships in Proteins of
Unsolved Three-Dimensional Structure. PROTEINS: Struct Funct Gen. 1990;7:99-111.

36. Wills PR. Genetic Information, Physical Interpreters and Thermodynamics; The Material-
Informatic Basis of Biosemiosis Biosemiotics 2014;7:141–65. doi: 10.1007/s12304-013-9196-
2.

37. Thompson JD, Higgins DG, Gibson TJ. Clustal W: improving the sensitivity of progressive
multiple sequence alignment through sequence weighting, positions-specific gap penalties and
weight matrix choice. Nucleic Acids Res. 1994;22:4673-80.

38. Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG. The ClustalX windows
interface: flexible strategies for multiple sequence alignment aided by quality analysis tools.
Nucleic Acids Res. 1997;24:4876-82.

39. Whelan S, Goldman N. A General Empirical Model of Protein Evolution Derived from
Multiple Protein Families Using a Maximum-Likelihood Approach. Molecular Biology and
Evolution. 2001;18(5):691-9.

40. Le SQ, Gascuel O. An Improved, General Amino-Acid Replacement Matrix. Molecular
Biology and Evolution 2008 2008;25((7)):1307-20.

41. Fournier GP, Alm EJ. Ancestral Reconstruction of a Pre-LUCA Aminoacyl-tRNA Synthetase
Ancestor Supports the Late Addition of Trp to the Genetic Code. J Mol Evol. 2015;80:171-85.
doi: 10.1007/s00239-015-9672-1.

42. Fournier GP, Andam CP, Alm EJ, Gogarten JP. Molecular Evolution of Aminoacyl tRNA
Synthetase Proteins in the Early History of Life. Orig Life Evol Biosph. 2011;41 621–32

43. Katsnelson MI, Wolf YI, Koonin EV. Towards physical principles of biological evolution.
arXiv. 2018;orgabs1709.00284.

44. Shiba K. Glycyl-tRNA Synthetases. In: Ibba M, Francklyn C, Cusack S, editors. The
Aminoacyl-tRNA Synthetases Austin: Landes Bioscience; 2005.

45. Soucy SM, Huang J, Gogarten JP. Horizontal gene transfer: building the web of life. Nat Rev
Gen. 2015;16(AUGUST):472

46. Carter CW, Jr, Wills PR. Interdependence, Reflexivity, Fidelity, and Impedance Matching, and
the Evolution of Genetic Coding. Molecular Biology and Evolution. 2018;35(2):269-86. doi:
10.1101/139139. PubMed Central PMCID: PMC29077934.

47. Wills PR. Reflexivity, Coding, and Quantum Biology. BioSystems. 2019;In preparation.

48. Wills PR, Carter CW, Jr. Insuperable problems of an initial genetic code emerging from an
RNA World. BioSystems. 2018;164:155-66. doi: 10.1101/140657

49. Goldman AD, Beatty JT, Landweber LF. The TIM Barrel Architecture Facilitated the Early
Evolution of Protein-Mediated Metabolism. J Mol Evol 2016;82(Jan (1)):17-26. Epub 2016
Jan 5. doi: 10.1007/s00239-015-9722-8. PubMed Central PMCID: PMC26733481.

50. Hemmingsen JM, Gernert KM, Richardson JS, Richardson DC. The tyrosine corner: a feature
of most Greek key beta-barrel proteins. Protein Science. 1994;3(11):1927-37.

51. Smith JL. Enzymes of nucleotide synthesis. Curr Op Struct Biol. 1995;5:752-7.

52. Tyson NdG. "Just to settle it once and for all: Which came first the Chicken or the Egg? The
Egg – laid by a bird that was not a Chicken. 2013-01-28. 2013:Tweet.

53. Fontana W, Schuster P. Continuity in Evolution: On the Nature of Transitions. SCIENCE.
1998;280(29 MAY):1451-5.

54. Lauring AS, Andino R. Quasispecies Theory and the Behavior of RNA Viruses. PLoS
Pathogens 2010;6 (7):e1001005.

55. Perona JJ, Gruic-Sovulj I. Synthetic and Editing Mechanisms of Aminoacyl-tRNA
Synthetases. Topics in Current Chemistry 2013;DOI: 10.1007/128_2013_456. doi:
10.1007/128_2013_456.

56. Bullock T, Uter N, Nissan TA, Perona JJ. Amino Acid Discrimination by a class I aminoacyl-
tRNA synthetase specified by negative determinants. J Mol Biol. 2003;328:395-408.

57. Sever S, Rogers K, Rogers MJ, Carter CW, Jr., Söll D. *Escherichia coli* tryptophanyl-tRNA
synthetase mutants selected for tryptophan auxotrophy implicate the dimer interface in
optimizing amino acid binding. Biochemistry. 1996;35:32-40.

58. Carter CW, Jr., Wills PR. The Roots of Genetic Coding in Aminoacyl-tRNA Synthetase
Duality Annual Review of Biochemistry. 2021;89; In Preparation.

59. Shore J, Holland BR, Sumner JG, Nieselt K, Wills PR. The Ancient Operational Code is
Embedded in the Amino Acid Substitution Matrix and aaRS Phylogenies. Journal of Molecular
Evolution. 2019; 88:136–50. doi: 10.1007/s00239-019-09918-z.

60. Bouckaert R, Vaughan TG, Sottani JB, Duchene S, Fourment M, Gavryushkina A, et al.
BEAST 2.5: An Advanced Software Platform for Bayesian Evolutionary Analysis. PLoS
Comput Biol. 2019;15(4):e1006650.

61. Buehner M, Ford GC, Moras D, Olsen KW, Rossmann MG. D-Glyceraldehyde 3-Phosphate
Dehydrogenase: Three Dimensional Structure and Evolutionary Significance. Proc Nat Acad
Sci USA. 1973;70:3052-4.

62. Koonin EV. The Logic of Chance: The Nature and Origin of Biological Evolution. Upper
Saddle River, NJ: Pearson Education; FT Press Science; 2011.

63. Koonin EV, Novozhilov AS. Origin and Evolution of the Genetic Code: The Universal
Enigma. IUBMB Life, . 2009;61((2) February):99–111. doi: 10.1002/iub.146.

64. Morrison DA. Multiple Sequence Alignment is not a Solved Problem. arXiv.
2018:1808.07717.

65. Popinga A. From the Origins of Life to Epidemics: Bayesian Inference, Stochastic Simulation,
and Dynamics of Bioinformatic Systems. [Doctoral]. Auckland, NZ: University of Auckland:
Supplementary Data. http://github.com/alexpopinga/aaRS-Pipeline, accessed 11 April 2019;
2019.

66. Söding J, Biegert A, Lupas AN. The HHpred interactive server for protein homology detection
and structure prediction. Nucl Acids Res. 2005;33:W244–W8 doi: 10.1093/nar/gki408.

67. Webb B, Sali A. Comparative Protein Structure Modeling Using Modeller. Current Protocols
in Bioinformatics. 54: John Wiley & Sons, Inc.,; 2016. p. 5.6.1-5.6.37.

68. Humphrey W, Dalke A, Schulten K. VMD - Visual Molecular Dynamics. Journal of Molecular
Graphics. 1996;14:33-8.

69. Carter CW, Jr., Carter CW. Protein Crystallization Using Incomplete Factorial Experiments.
Journal of Biological Chemistry. 1979;254:12219-23.

70. Yang Z. Maximum Likelihood Phylogenetic Estimation from DNA Sequences with Variable
Rates over Sites: Approximate Methods. J Mol Evol. 1994;39:306-14.

71. Bouckaert RR. DensiTree: making sense of sets of phylogenetic trees. BIOINFORMATICS.
2010;26 (10):1372–3. doi: 10.1093/bioinformatics/btq110.

72. Rambaut A. FigTree. 1.4.0 ed: University of Edinburgh; 2010.

73. Gonnet GH, Cohen MA, Benner SA. Exhaustive matching of the entire protein sequence
database. Science. 1992;256:1443–5.

74. SAS. JMP: The Statistical Discovery Software. V.13.1 ed. Cary, NC: SAS Institute, Cary NC;
2015.

75. Pymol. The PyMOL Molecular Graphics System. Schrödinger, LLC.

774
775
776   Table 1. Design matrix for regression analysis[1].
777

Subset	$\langle S \rangle_{WAG}$	$\langle S \rangle_{LG}$	$\langle Q \rangle$	Tree height	Shape	A	B	C	CP1	NUMB
Full	0.9	0.86	48.8	3.69	1.17	4	3	1	2	103
Urzyme	0.87	0.82	52.4	3.40	1.60	4	3	1	0	83
Protoz	0.85	0.85	52.3	2.71	1.38	4	0	0	0	46
CP1	0.34	0.35	33.7	2.17	4.49	0	0	0	2	20
Urz_0	0.76	0.73	50.1	3.03	1.46	1	1	0	0	20
Urz_1	0.71	0.68	57.8	3.99	0.91	0	1	0.5	0	20
Urz_2	0.72	0.7	55	2.78	1.13	2	0	0	0	20
Urz_3	0.62	0.62	45.6	2.54	2.26	0	2	0	0	20
Urz_4	0.82	0.77	58.5	2.41	1.21	2	1	0.5	0	20
Urz_5	0.78	0.8	45.6	2.88	1.98	4	0	0	0	20
Urz_6	0.62		56.6	2.65	1.13	0	2	1	0	20

778
779
780

[1] Shaded columns are dependent variables; unshaded columns are independent variables and serve as predictors. Numerals in columns A, B, C, and CP1 are proportional to the total number of amino acids in the respective MSAs. Two additional independent variables, Urzyme and protozyme, constructed in a related fashion, are not shown. NUMB is the number of amino acids in the alignment.

781 Table 2. Cross-correlation R^2 values between phylogenetic metrics
782

	Tree height	Shape
Shape	0.15	
<S>	0.10	0.03

Fig Legends

Fig 1. Structural modules underlying the hypothesis and organization of data adduced as
evidence. **A**. The Class I Urzyme and internal CP1 insertion that together make up the Class I
catalytic domain. Cartoons were prepared with Pymol [75] from coordinates (PDB ID 1I6L) for
the tryptophanyl-tRNA synthetase, the smallest Class I aaRS. The activated aminoacyladenylate
is drawn as sticks. The primary size variation the remaining nine Class I aaRS occurs as further
insertions within CP1 as structures displayed here are essentially unchanged. Anticodon-binding
domains are idiosyncratic, and not shown. **B**. Overlapping portions of Class I and II aaRS as
envisioned in an ancestral bidirectional gene [33] coincide with the respective Urzymes [19, 30].
Vertical lines denote ancestral base-pairing. Grey segments were presumably more recent
additions. In particular, insertion of CP1 is incompatible with bidirectional coding. **C**. Schematic
location of CP1 between two roughly equal fragments of the Urzyme, colored to allow
identification of structural fragments in **A**. Intensely colored segments are highly conserved
secondary structures in all 10 Class I aaRS and compose the Class I scaffold. Subclass I A enzymes
contain one or more additional insertions (red). **D**. Sampled alignments consisting of different
segments totaling 20 amino acids, selected as described in Methods. **C** and **D** amplify the Class I
base-paired portion of **B**. The protozyme (Urz_A) is the amino terminal β-α-β crossover
connection (blue), the amino acid binding pocket (Urz_B) is formed by the protozyme and two
intermediate α-helices (amber). The KMSKS loop (Urz_C) is rose. Previous work [25] established
that the CP1 insertion (green) has little impact on the enzymatic properties of the TrpRS Urzyme
unless the anticodon-binding domain is also present. Colors match those in (**A**).

Fig 2. Histograms of the Tree height (**A**) and Shape (**B**) metrics highlight differences
between sequence variability within CP1 (blue) and Urzyme (red) sequences. The Tree height is
the reciprocal of the estimated mutation rate per site. CP1 has the lowest site-specific mutation rate
and the highest Shape parameter (blue dot implies statistical significance). The Urz_1 subset of
amino acids includes a stretch of 10 amino acids along the specificity determining helix, where the
highest site-specific mutation frequency occurs within the Urzyme. **C**. Regression model showing
the dependence of Tree height on MSA subsets. The horizontal blue line in this and other
regression curves is the average value of the dependent variable. Dots represent the various MSAs
in the design matrix and are colored and distinguished by different symbols for identification. R^2
for this model is 0.87. Numerals refer to the 20 residue subsets defined in Fig 1. The F-ratio is 9.9
with a P-value of 0.008.

Fig 3. Position-specific metrics. **A**. Colinearity of the logarithm of the Shape parameter of
the gamma distribution of the Tree heights estimated by BEAST2 for different MSAs derived from
the Class I scaffold $\langle Q \rangle$, and the mean conservation quality scores, $\langle Q \rangle$ [38] obtained directly
from respective MSAs. **B**. The conservation quality, <Q>, for the four segments of the Class I
scaffold alignment. Class I signature peptides [1] are in parentheses. Error bars show the standard
error of the mean over all positions within the segment. The numbers of amino acids in each
segment are given for each histogram. **C**. and **D**. are regression models showing nearly identical
dependence of $\log_{10}$(Shape) and $\langle Q \rangle$, respectively, on predictors in the design matrix (Table 1).
Dots represent different MSAs. They are labeled and colored as in Fig 2 to emphasize the
extraordinary similarity of the two site-by-site metrics derived two different ways.

Fig 4. Shape parameter dependence. **A.** Plot of the multivariate regression model for Shape.
$R^2 = 0.94$ for the regression and the P value for the F ratio is 0.0006. **B**. Regression coefficients
and their Student t-test probabilities. Points for trees constructed using both WAG and LG
substitution matrices are shown for comparison. Significant predictors of Shape are the amino acid
positions in CP1, those in the B-fragment containing the amino acid specificity-determining helix
(sand; Fig 1A), and their two-way interaction.

Fig 5. Phylogenetic support. **A.** DensiTree representations of the Urzyme and CP1
segments of the Class I structural scaffold. Each aaRS is monophyletic in the Urzyme alignment,
whereas several of the clades in the CP1 alignment, highlighted in color, have multiple ancestries.
B. Regression model for $\langle S \rangle_i$. Dots are colored and labeled as in Fig 3. Values in the Regression
table are only for the WAG matrix, as use of both WAG and LG matrices unduly enhances the P-
values of the β-coefficients. The R^2 and P value of the model's F-ratio are shown under the X-axis.
Coefficients and their statistics for the model are given in Table 2. Individual $\langle S \rangle$ values are
labeled to enable comparison with Fig 1C. R^2 was 0.94 for 20 observations, and the F-ratio for the
regression table was 92.4 with a P-value <0.0001.

Fig 6. A. Structural relationships between the TrpRS Urzyme, CP1, and the anticodon-
binding domain (ABD). Coloring in this Fig differs from that in Figs 1 and **B**. The CP1 motif forms
an annulus that constrains motions of the specificity determining helix (sand) and the Protozyme
(yellow), constraining the effective size of the amino acid binding pocket when the ABD (teal)
changes its orientation. Experimental evidence described in the text confirms that these constraints
enable full-length TrpRS to reject tyrosine in the transition state complex for tryptophan activation.
B. Structural cartoon with details of the interaction illustrated in **A**. Coloring in this Fig is the same
as that in Fig 1. The specificity-determining helix is the site of the Set 1 segment, which is colored
in Blue. The TrpRS CP1 module is green and the scaffold segments are red. Note the close contact
between CP1—especially the red scaffold segment—and Set 1. **C.** Interactions between the TrpRS
protozyme—locus of the ATP binding site, and segment B, locus of the amino acid binding site.
Subsets 2 and 1, which are contained entirely within the respective segments, are highlighted in
dark blue.

Fig 7. Assignment catalysis and code evolution. **A.** Correlations between rate accelerations
and phylogenetic metrics, $\langle Q \rangle$ and $\langle S \rangle$. Parameters for regression against experimental catalytic
proficiencies for corresponding putative evolutionary intermediate constructs are both significant.
Blue, green, and amber triangles represent protozymes, Urzymes, and catalytic domains. **B.**
Timeline for growth of the genetic coding alphabet from a two-letter code. Introducing new aaRS
into the context of the ancestral bidirectional gene (grey background and dashed connecting lines)
simultaneously enhanced specificity and created fundamental changes in selection pressure.
Different colored backgrounds signify altered selection pressures that apply to all extant aaRS at
a given stage of the coding alphabet. Increasing cardinality of the alphabet induces (i) sequence
space inflation as a greater number of distinguishable sequences can be specified; and (ii)
restriction in the diameters of the quasispecies as they approach fully coded sequences in the final
20-letter alphabet. Bold face landmarks (CP1 and ID) denote qualitative changes in aaRS
architecture shown experimentally to enrich specificity as discussed in the text. Note that this
timeline like that in Fig 5 represents evolutionary events before the LUCA.

Fig 8. Modified Class I aaRS phylogeny consistent with bidirectional coding ancestry in a
peptide/RNA world. Adapted with permission from Fig 12-2A of [62]. Green circles represent key
stages in the emergence of genetic coding, beginning with ATP-dependent amino acid activation
by protozymes coded by a bidirectional gene [21]. Magenta background indicates the progression
illustrated in detail in Fig7B.

Fig 1.

Term	β	Std Error	t Ratio	Prob>|t|
Intercept	2.17	0.26	8.35	0.0002
B	-1.39	0.32	-4.38	0.0047
Urzyme	3.21	0.61	5.27	0.0019
Protozyme	-2.70	0.57	-4.75	0.0031
B*Protozyme	1.66	0.33	5.05	0.0023

878
879
880   Fig 2.
881

883
884
885 Fig 3.
886

Term	β	Std Error	t Ratio	Prob>ltl
Intercept	1.37	0.17	8.26	0.0002
B	0.40	0.16	2.45	0.0498
C	-1.08	0.40	-2.73	0.0342
CP1	**1.56**	**0.19**	**8.43**	**0.0002**
B*CP1	**-0.57**	**0.10**	**-5.96**	**0.001**

891
892
893   Fig 5.
894

Fig 5.

899
900
901   Fig 6.
902

903
904
905   Fig 7.
906

907
908
909   Fig 8.
910