1

**Triclosan depletes the membrane potential in *Pseudomonas aeruginosa* biofilms inhibiting**

**aminoglycoside induced adaptive resistance**

Michael M. Maiden and Christopher M. Waters[*]

[1]Departments of Microbiology and Molecular Genetics and The BEACON Center for The Study
of Evolution in Action, Michigan State University, East Lansing, Michigan, United States of
America

*Corresponding Author

E-mail: watersc3@msu.edu (CW)

**Abstract**

Biofilm-based infections are difficult to treat due to their inherent resistance to antibiotic

treatment. Discovering new approaches to enhance antibiotic efficacy in biofilms would be

highly significant in treating many chronic infections. Exposure to aminoglycosides induces

adaptive resistance in *Pseudomonas aeruginosa* biofilms. Adaptive resistance is primarily the

result of active antibiotic export by RND-type efflux pumps, which use the proton motive force

as an energy source. We show that the protonophore uncoupler triclosan depletes the membrane

potential of biofilm growing *P. aeruginosa,* leading to decreased activity of RND-type efflux

pumps. This disruption results in increased intracellular accumulation of tobramycin and

enhanced antimicrobial activity *in vitro*. In addition, we show that triclosan enhances tobramycin

effectiveness *in vivo* using a mouse wound model. Combining triclosan with tobramycin is a new

anti-biofilm strategy that targets bacterial energetics, increasing the susceptibility of *P.*

*aeruginosa* biofilms to aminoglycosides.

**Author summary**

Adaptive resistance is a phenotypic response that allows *P. aeruginosa* to transiently

survive aminoglycosides such as tobramycin. To date, few compounds have been identified that

target adaptive resistance. Here, we show the protonophore uncoupler triclosan disrupts the

membrane potential of *P. aeruginosa*. The depletion of the membrane potential reduces efflux

pump activity, which is essential for adaptive resistance, leading to increased tobramycin

accumulation and a shorter onset of action. Our results demonstrate that in addition to its

canonical mechanism inhibiting membrane biosynthesis, triclosan can exert antibacterial

properties by functioning as a protonophore that targets *P. aeruginosa* energetics.

## Introduction

**Introduction**

*Pseudomonas aeruginosa* is a Gram-negative opportunistic pathogen that can form highly

recalcitrant biofilms in immunocompromised hosts such as those suffering from cystic fibrosis

(CF) and diabetes (1-4). Biofilms are a community of cells enmeshed in a self-made matrix that

is highly tolerant to antibiotic treatment (1-4). In addition to biofilm growth, *P. aeruginosa*

possesses many antibiotic resistance mechanisms including low outer membrane (OM)

permeability, which is estimated to be 1/100[th] that of *Escherichia coli* (5), the expression of

proton motive force (PMF) dependent resistance-nodulation-cell division (RND) family of

multidrug efflux pumps, and a chromosomal encoded AmpC β-lactamase, making it naturally

resistant to β-lactams (4).

*P. aeruginosa* also exhibits a unique form of adaptive resistance in response to

aminoglycosides, cationic antimicrobial peptides, and polymyxins (6, 7). Adaptive resistance is

mainly mediated by the induction of RND-type efflux pumps and by alterations to the OM

within 2-hrs of exposure to subinhibitory concentrations of aminoglycosides (8-11).

Aminoglycosides cause the mistranslation of the PA5471 leader peptide PA5471.1, a peptide

which inhibits the MexZ repressor, thereby triggering the induction of the RND-type efflux

pump MexXY (12). This phenomenon has been observed *in vitro,* in animal models, and in CF

patients (13-17).

Aminoglycoside uptake occurs in three steps (18-20). In the first step, termed "self-

promoted uptake," positively charged aminoglycosides bind negatively charged components of

the OM, triggering permeabilization and diffusion (18-20). In the second step, energy dependent

phase-I (EDP-I), aminoglycosides slowly cross the inner membrane (IM) dependent upon the

membrane potential ($\Delta\psi$) (18-20). In the third step, EDP-II, aminoglycosides bind to the A-site

of the 30S subunit of membrane-associated ribosomes and cause the synthesis of misfolded

proteins that insert into the IM triggering permeabilization of the cytosolic barrier (18-20). This

leads to an irreversible entry of aminoglycosides into the cell, thereby inhibiting translation.

Despite the well-documented phenomenon of adaptive resistance,

aminoglycosides continue to be the standard of care for pseudomonal infections (21). In fact,

pulmonary infections in CF patients are treated with extended doses of the aminoglycoside

tobramycin (300-mg nebulized twice-a-day for 28-days), reaching mean sputum concentrations

of ~737 µg/g per dose (21, 22). As antibacterial resistance continues to be an emerging public

health crisis, there is a crucial need for new therapeutic approaches that decrease resistance (23).

One approach is to target bacterial energetics (24). The feasibility of this approach was recently

demonstrated by the discovery of the protonophore uncoupler bedaquiline, which is the first new

Food and Drug Administration (FDA) approved *Mycobacterium tuberculosis* (*Mtb*) drug in 40-

years (25-27).

We previously performed a high throughput screen (HTS) to identify compounds that

enhanced tobramycin against *P. aeruginosa* biofilms and unexpectedly discovered that the

combination of triclosan and tobramycin (or other aminoglycosides) was 100-times more

effective than either triclosan, tobramycin, or other aminoglycosides alone at killing established

*P. aeruginosa* biofilms (28). Triclosan is a fatty acid synthesis inhibitor, targeting the enoyl-acyl

carrier reductase FabI  (29). However, it is well-known that *P. aeruginosa* is inherently resistant

to triclosan due to the expression of both the RND-type efflux pump TriABC and a triclosan

resistant enoyl-acyl carrier reductase FabV, and our previous results suggested that triclosan does

not enhance tobramycin activity by targeting FabI (28, 30, 31). Therefore, the mechanism of

action of triclosan enhancement of aminoglycoside activity remained unknown.

The PMF is the sum of the membrane potential due to charge separation ($\Delta\psi$,

positive$_{outside}$/negative$_{inside}$) and the proton gradient ($\Delta pH$, acidic$_{outside}$/alkaline$_{inside}$) across the

inner membrane (32, 33). Recently, it has been shown that in addition to fatty acid synthesis

inhibition, triclosan possesses protonophore activity that can disrupt the $\Delta\psi$ of mitochondria

isolated from rats, artificial bilayer lipid membranes, and the Gram-positive bacterium *Bacillus*

*subtilis* (34-38). This led us to hypothesize that triclosan enhances aminoglycoside activity by

targeting the $\Delta\psi$ in *P. aeruginosa* (28). In fact, from the same HTS in which triclosan was

identified, we discovered that the protonophore uncoupler oxyclozanide acted synergistically

with tobramycin to kill *P. aeruginosa* biofilms (39).

Here we demonstrate that triclosan increases *P. aeruginosa* susceptibility to tobramycin

by disrupting adaptive resistance. This occurs because triclosan decreases the $\Delta\psi$, leading to a

reduction in the activity of RND-type efflux pumps, which increases tobramycin accumulation

within cells. Finally, we demonstrate that triclosan in combination with tobramycin embedded in

a hydrogel is effective *in vivo* using a murine wound model. As numerous toxicity studies have

demonstrated that triclosan is safe when used appropriately (40-42), this combination has the

clinical potential to improve the treatment of biofilm-based infections by targeting bacterial

energetics.

**Results**

**The disruption of fatty acid synthesis does not increase *P. aeruginosa* susceptibility to**

**tobramycin.**

115            To determine if the inhibition of fatty acid biosynthesis and membrane biogenesis by

116     triclosan is responsible for increasing *P. aeruginosa* susceptibility to tobramycin, we tested if the

117     specific fatty acid synthesis inhibitor AFN-1252 synergized with tobramycin to kill *P.*

118     *aeruginosa* biofilms. AFN-1252 forms a binary complex with the active site of FabI, which is an

119     enoyl-acyl reductase responsible for the final elongation step in fatty acid synthesis and is

120     inhibited by triclosan (43).

121            AFN-1252 alone and in combination with tobramycin was not effective at killing

biofilms (Fig. S1). At the maximal concentrations of tobramycin and AFN-1252 used we

observed slightly more than 2-fold killing; however, this difference was not statistically

significant and far weaker than the ~100-fold killing observed when triclosan is combined with

tobramycin (28). These findings are in agreement with our previous genetic experiments using a

FabI mutant of *P. aeruginosa* that showed identical resistance to tobramycin treatment as the

parental strain (28). Together, both our genetic and molecular experiments suggest that the

inhibition of FabI by triclosan does not account for increased *P. aeruginosa* susceptibility to

tobramycin.

**Triclosan combined with tobramycin causes synergistic permeabilization of cells within**

**biofilms.**

132            Aminoglycosides corrupt protein synthesis by binding to the A-site of the 30S subunit of

membrane bound ribosomes causing mistranslation of membrane proteins and inner membrane

permeabilization (18-20). We initially hypothesized that triclosan enhanced tobramycin killing

by permeabilizing cells in biofilms. To test this hypothesis, we stained cells with the cell-

impermeant TO-PRO™-3 iodide dye, which selectively emits a fluorescent signal in

permeabilized cells.

Treatment for 2-hrs with triclosan or tobramycin alone resulted in ~15% and ~11% of the

cells within biofilms to become permeabilized, respectively (Fig. 1). Neither of these increases

were statistically significantly compared to untreated controls. Alternatively, the combination of

triclosan and tobramycin permeabilized ~50% of cells after only 2-hrs of treatment. This result

suggests that neither triclosan nor tobramycin treatment alone is sufficient to permeabilize *P.*

*aeruginosa* cells growing in biofilms.

**Mutations in *fusA1* are resistant to the combination.**

To further elucidate the mechanism of action of this combination, we selected for

resistance mutants and then performed whole genome sequencing to identify causative

mutations, using a modified method developed by Lindsey et al. (44). 27 *P. aeruginosa* biofilms

serially passaged in parallel were exposed to sudden, moderate and gradual treatment regimens

consisting of ever-increasing concentrations of tobramycin and triclosan (Table S1). No resistant

mutations were isolated from populations exposed to sudden or moderate treatment regimens.

After 30 passages, two resistant populations from the gradual treatment regimen were isolated

(Table 1), and 96-colonies were isolated from each mutant pool that were found to be resistant to

the combination.

To determine the mutation(s) responsible for the resistance to the combination we

performed whole genome sequencing on two mutants from the F1 and four mutants from the B2

mutant populations (Table 1). These 6 mutants all showed complete resistance to tobramycin and

partial resistance to the combination compared to the ancestral strain (Fig. 2). The low level of

killing generated by the combination may be attributed to triclosan as all six of these mutants

exhibited increased sensitivity to triclosan treatment alone versus the ancestral strain. This

phenomenon, termed "collateral sensitivity," is the process by which resistance to one

antimicrobial simultaneously results in increased sensitivity to unrelated antimicrobials (45).

One common feature of all six mutants was a single nucleotide polymorphism (SNP) in

the *fusA1* gene encoding for EF-G1A, a protein responsible for ribosomal translocation and

recycling (46). Two unique mutations, a SNP at residues L40Q or T64A were identified. These

mutations are near or within the switch I domain, respectively, which is required for hydrolysis

of guanosine triphosphate and the "power-stroke" of EF-G1A. Because all mutants having a

L40Q SNP came from mutant pool F1 and all mutants having a T64A SNP came from mutant

pool B2, we hypothesize that these SNPs occurred early on during selection, allowing for clonal

expansion. However, no two isolates have the exact same set of SNPs, indicating the sequenced

strains are not siblings and each isolate is a distinct mutant as detailed in the discussion

(supplemental spreadsheet). Importantly, mutations in *fusA1* that have been found in clinical

isolates, including L40Q, confer resistance to tobramycin through an unknown mechanism (47).

Together, these results suggest that tobramycin is the component of the combination that is

primarily responsible for killing.

**Initial tobramycin treatment is required for the combination to be effective.**

The corruption of translation is thought to be the first step in aminoglycoside induced

adaptive resistance (12). This led us to hypothesize that tobramycin must first corrupt translation

thereby triggering adaptive resistance and the induction of RND-type efflux pumps for the

combination to be synergistic (8-10, 12). To test if tobramycin induction of adaptive resistance

was important for synergy, we first treated *P. aeruginosa* biofilms with tobramycin alone for 3-

hrs and then washed the biofilms three time in Dulbecco's Phosphate Buffer Solution (DPBS) for

3-mins each followed by subsequent treatment with triclosan alone for 3-hrs. We performed a

similar experiment first treating with triclosan followed by tobramycin treatment.

Sequential treatment with tobramycin followed by triclosan addition caused synergy,

resulting in ~2-$\log_{10}$ reduction in cells within biofilms compared to untreated controls (Fig. S2).

Intriguingly, despite no longer having tobramycin in the media, the cells within the biofilms were

killed when treated with triclosan. We speculate that enough tobramycin remains after washing

due to association of aminoglycosides with the extracellular polymeric substances (EPSs)

making up the biofilm matrix and cell itself, which has been demonstrated to be resistant to

removal by washing (19, 48). Alternatively, initial treatment with triclosan followed by treatment

with tobramycin did not cause synergy or result in significant biofilm killing. These data suggest

that a phenotypic switch induced by aminoglycoside exposure is important for synergistic

activity.

In support of these experiments, we previously showed that tobramycin induces classic

adaptive resistance, rendering *P. aeruginosa* refractory to killing by tobramycin until 6-hours,

with the greatest resistance observed at 2-hrs (28). Furthermore, we found that only antibiotics

that corrupt translation (aminoglycosides and tetracycline) act synergistically or are enhanced

when combined with triclosan (28). Our past and current results lead us to hypothesize that

tobramycin triggers adaptive resistance, which is then disrupted by triclosan.

**Triclosan reduces RND-type efflux pump activity.**

204             Because RND-type efflux pumps are a major component of aminoglycoside induced

adaptive resistance (8-10, 12), we measured their activity in biofilms after treatment with

tobramycin alone or in combination with triclosan by measuring cellular accumulation of

ethidium bromide. Ethidium bromide is a substrate of RND-type efflux pumps, and its

accumulation within cells can be used as a proxy for their activity (49). In this assay, the

protonophore carbonyl cyanide 3-chlorophenylhydrazone (CCCP), which is known to reduce

efflux pump activity, was used as a positive control (49). CCCP treatment led to increased

ethidium bromide accumulation in *P. aeruginosa* biofilms as expected (Fig. 3). Similarly,

triclosan reduced efflux pump activity as indicated by increased ethidium bromide accumulation,

but tobramycin treatment alone had no effect. The combination of triclosan and tobramycin was

similar to triclosan alone. These data suggest that triclosan reduces the activity of RND-type

efflux pumps.

**Triclosan increased intercellular accumulation of tobramycin.**

218             Triclosan inhibition of RND-type efflux pumps should increase the accumulation of

219     tobramycin within cells. To test this, we conjugated tobramycin with the fluorescent Texas Red

220     dye (TbTR)  as previously described (50), and measured its accumulation within cells growing as

221     biofilms. Conversely, we also measured TbTR extrusion from cells into treatment-free recovery

222     media.

223             For accumulation assays, biofilms were treated with TbTR with and without triclosan for

30-mins. Based on our previously published time-killing studies, this time point is before the

combination causes cell death (28). Prior to reading fluorescence, biofilms were washed,

mechanically disrupted with a wooden stick, and cells were lysed to release TbTR. Triclosan in

combination with TbTR resulted in a statistically significant increase in TbTR accumulation

whereas CCCP did not significantly increase TbTR accumulation (Fig. 4A). This is in agreement

with CCCPs reduced potency compared to triclosan at disrupting the membrane potentials of

bacteria due to differences in lipophilicity and reduced activity against RND-type efflux pumps

in *P. aeruginosa* biofilms (36).

To confirm that co-treatment with triclosan increased tobramycin accumulation within

cells, biofilms were treated with TbTR with and without triclosan for 30-mins. Then biofilms

were washed three-times in DPBS, recovered in treatment-free media for 30-mins and the

supernatant was measured. Triclosan in combination with TbTR resulted in greater overall

extrusion into treatment-free media, indicating more TbTR had accumulated in cells prior to

recovery (Fig. 4B). These results support the conclusion that triclosan reduces the activity of

RND-type efflux pumps, leading to increased accumulation of TbTR within cells.

**Triclosan inhibits tobramycin induced $\Delta\psi$.**

Triclosan has a hydroxyl group with a dissociable proton that can disrupt the $\Delta\psi$ in

eukaryotic cells, mitochondria, and bacteria (34-38). Because adaptive resistance is primarily the

result of PMF driven RND-type efflux pumps (8-10, 12), we speculated that the depletion of the

$\Delta\psi$ could lead to their inhibition and the abolishment of adaptive resistance. In these

experiments, we measured changes in the $\Delta\psi$ rather than the $\Delta pH$ because at neutral pH the PMF

in bacteria is primarily composed of the $\Delta\psi$ due to homeostatic buffering (51, 52), and RND-

type efflux pump activity is primarily driven by the $\Delta\psi$ which is generated more rapidly than

ΔpH (53, 54). To measure changes in the Δψ of *P. aeruginosa* growing as biofilms, cells were

stained with the fluorescent Δψ indicator DiOC2(3) dye and analyzed by flow cytometry. Cells

were also co-stained with the membrane impermeant TO-PRO™-3 iodide dye to exclude

permeabilized cells from analysis of Δψ.

Triclosan treatment for 2-hrs reduced the population of cells maintaining a Δψ 4-fold,

from ~20% to ~5% (Fig. 5). Tobramycin treatment resulted in an increase in the population of

cells with a Δψ almost 2-fold, from ~20% to ~37%. This increase is indicative of adaptive

resistance (8-10, 12). Triclosan in addition to tobramycin reduced the population of cells

maintaining a Δψ almost 4-fold from tobramycin treatment alone to levels comparable to

untreated biofilms. These data suggest that triclosan depletes the Δψ at an essential time in

aminoglycoside induced adaptive resistance when a surge in the Δψ is protective against

tobramycin by increasing RND-type efflux pump activity.

**Methyl-triclosan does not disrupt the membrane potential or synergize with tobramycin.**

To confirm that protonophore activity by triclosan is responsible for tobramycin

enhancement, we treated cells with the chemical derivative methyl-triclosan, which lacks the

hydroxyl moiety necessary for gaining and losing a proton and therefore cannot act as a

protonophore (35). In addition, without this hydroxyl group, methyl-triclosan cannot inhibit fatty

acid synthesis by forming a complex with the co-factors NADH/NAD+ (35).

We first performed biofilm susceptibility testing using methyl-triclosan alone and in

combination with tobramycin. Neither methyl-triclosan nor the combination were effective

against biofilms after 6-hrs of treatment over a range of concentrations (Fig. S3). Furthermore,

methyl-triclosan had no effect on the population of cells maintaining a Δψ compared to untreated

biofilms and did not reduce the tobramycin induced surge in $\Delta\psi$ after 2-hrs of treatment (Fig.

S4A). In addition, methyl-triclosan did not cause permeabilization of cells alone or in

combination with tobramycin (Fig. S4B). Likewise, methyl-triclosan was unable to significantly

reduce efflux pump activity (Fig. S5). These results support the conclusion that triclosan

protonophore activity is required for the depletion of the tobramycin induced $\Delta\psi$ surge and

inhibition of RND-type efflux pump activity.

**ATP Depletion does not render cells more sensitive to tobramycin.**

Another outcome of triclosan depleting the $\Delta\psi$ would be decreased energy generation,

which could contribute to the sensitivity of biofilms to tobramycin. We performed antimicrobial

susceptibility assays on 24-hr old biofilms grown in rich medium that were then DPBS-starved

for 5 days. We observed the same synergy of the combination in these conditions as treatment

with triclosan or tobramycin alone failed to cause bacterial killing whereas the combination of

triclosan and tobramycin caused significant killing (Fig. S6). These data suggest that the

disruption of the $\Delta\psi$ by triclosan does not induce synergy by depletion of ATP.

**Triclosan and tobramycin show enhanced efficacy in an *in vivo* wound model.**

To determine if triclosan and tobramycin are more effective against biofilms *in vivo,* we

tested their activity using a murine wound model (55, 56). In this model, a 1-day old wound on

the back of SKH-1 hairless mice were infected with ~$1\times10^9$ XEN 41 bioluminescent *P.*

*aeruginosa* cells growing as biofilms to allow imaging of the infection using the In Vivo

Imaging System.

2-day old biofilms were treated for 4-hrs using an agarose hydrogel imbedded with either

triclosan or tobramycin alone or in combination. Triclosan hydrogels had no effect compared to

control hydrogels while tobramycin hydrogels resulted in statistically non-significant 2.5-fold-

reduction in bioluminescence (Fig. 6). Hydrogels embedded with triclosan and tobramycin

resulted in statistically significant 4.5-fold-reduction in bioluminescence. The combination

hydrogels resulted in greater than 4-fold reduction in five mice whereas tobramycin treatment

only achieved this level of killing once.

**Discussion**

Here we report that triclosan acts as a protonophore uncoupler against *P. aeruginosa*

growing as biofilms (34-38). To our knowledge, this is the first report of protonophore activity

by triclosan against Gram-negative bacteria growing as biofilms both *in vitro* and *in vivo*.

Adaptive resistance, occurring within 2-hrs of exposure, reduces accumulation of

antimicrobials by inducing the activity of PMF driven RND-type efflux pumps and alterations to

the OM (8-12) . We previously demonstrated classic aminoglycoside adaptive resistance, finding

*P. aeruginosa* biofilms were refractory to tobramycin killing until ~6-hrs, with the greatest

resistance observed at 2-hrs. However, the addition of triclosan with tobramycin led to rapid

killing of *P. aeruginosa* biofilms (28). Here we show that triclosan reduces the activity of PMF

driven RND-type efflux pumps, resulting in greater tobramycin accumulation within cells (Figs.

3 and 4). Triclosan does this by inhibiting a tobramycin induced surge in $\Delta\psi$ occurring at 2-hrs

(Fig. 5). Together, our past and present data suggest that the disruption of the $\Delta\psi$ depletes the

energy required for effective efflux pump mediated adaptive resistance (Fig. 7) (28). In support

of this model, methyl-triclosan, which lacks a dissociable proton and thus cannot act as

protonophore, was completely ineffective alone and in combination with tobramycin (Figs. S3-

5).

In addition to protein-specific corruption of translation at ribosomes, the very process of

aminoglycoside uptake augments its bactericidal activity by non-specifically disrupting the OM

and IM  (18-20). In the first step of uptake, positively charged aminoglycosides bind to the

negative charged components of the OM, including lipopolysaccharide and phospholipids in

Gram-negative bacteria. This is followed by the displacement of magnesium ions responsible for

OM stabilization, promoting uptake by disrupting and increasing OM permeability (18-20). In

addition, aminoglycosides can diffuse through porins in the OM (18-20). In EDP-I,

aminoglycosides cross the IM via a mechanism that is influenced by the PMF, specifically the

$\Delta\psi$, in a concentration dependent manner (18-20). In EDP-II, aminoglycoside cause the

mistranslation of proteins by membrane associated ribosomes, resulting in the insertion of

misfolded proteins in the IM, which permeabilizes the cytoplasmic membrane accelerating its

own accumulation (18-20).

Several studies have shown that aminoglycoside-mediated killing of bacteria can be

enhanced by supplementing exogenous metabolites or host derived molecules, which increase

cellular respiration, PMF, and the uptake of aminoglycosides through EDP-I (57-62).  However,

these studies used aminoglycoside concentrations ranging from 2-25 µg/mL on cells growing

planktonically (57-62). Alternatively, the experiments described here used 250 µg/mL of

tobramycin on *P. aeruginosa* biofilms. This higher concentration of tobramycin is more

clinically relevant and similar to antipseudomonal therapeutic dosing in cystic fibrosis patients

(1-3, 21, 22).

Although our model might appear to contradict these prior studies, it has been

demonstrated that at concentrations of aminoglycosides greater than 30 µg/mL, as used here, the

PMF is less important for aminoglycoside uptake during EDP-I  (18-20).  In fact, we observed

that 250 µg/mL tobramycin can ultimately lead to biofilm killing at 24-hrs similar to the

combination, suggesting that slow uptake of tobramycin during the EDP-I phase in the absence

of triclosan eventually permeabilizes the IM stimulating EDP-II uptake in most of the biofilm

(Fig. S7) (4, 48). However, the addition of triclosan accelerates this process by uncoupling the

PMF such that maximal killing occurs at 4-hrs, as previously described (Fig. S7) (28). Together,

our results support a model where triclosan reduction of the $\Delta\psi$ reduces the PMF required for

active efflux leading to rapid accumulation of tobramycin in the cytoplasm and the stimulation of

EDP-II uptake, which triggers irreversible cytosolic tobramycin accumulation and cell killing

(4). In support of our model, other publications have demonstrated that bacteria can be sensitized

to antibiotics by molecules that dissipate the PMF (24-27, 63-66). Specifically, Verstraeten et al.,

found that depleting the $\Delta\psi$ by overexpressing Obg in *E. coli* increased intercellular

accumulation of tobramycin, likely through reduced efflux (65).

RND-type efflux pumps are intricately linked to antibiotic resistance, especially in

biofilms (4, 54, 67). RND-type efflux pumps consist of three proteins which span the inner

membrane and outer membrane (Fig. 7) (54). The inner membrane protein (IMP, e.g. MexX)

catalyzes the $H^+$ dependent efflux of compounds and provides antimicrobial specificity whereas

the periplasmic membrane fusion protein (MFP, e.g. MexY) connects the outer membrane

protein (OMP, e.g. OprM) to the IMP creating an exit channel (54). There are 12 RND-type

efflux pumps encoded in the genome of *P. aeruginosa* (4) and five are well characterized:

MexAB-OprM, MexCD-OprJ, MexEF-OprN, MexXY-OprM and MexJK-OpRM (4). In

particular, the MexXY-OprM pump is responsible for aminoglycoside efflux, playing an

essential role in adaptive and acquired resistance (9-12, 68). The most frequent class of

aminoglycoside resistant mutants in CF isolates of *P. aeruginosa* have lost function of the

repressor MexZ, leading to overproduction of the RND-type efflux pump MexXY (67, 69). We

previously demonstrated that the combination of tobramycin and triclosan exhibited significant

killing against a clinical CF *P. aeruginosa* isolate AMT0023_34 that is tobramycin resistant by

such a mechanism (28). Importantly, our results suggest that the triclosan addition can potentiate

tobramycin to treat *P. aeruginosa* infections that are tobramycin resistant due to the

overexpression of efflux pumps. Which of the multiple efflux pumps impacted by triclosan

addition is under investigation.

We show that triclosan reduces efflux pump activity through the reduction of the $\Delta\psi$, but

this activity does not potentiate the cells to triclosan itself, even though the RND-type efflux

pump TriABC is a triclosan resistance mechanism expressed by *P. aeruginosa* (31). We

hypothesize that although triclosan reduces efflux pump activity, this reduction is not sufficient

to render *P. aeruginosa* sensitive to triclosan because this species also encodes the triclosan

resistant FabI ortholog FabV (30). In addition, if triclosan reduces efflux pump activity, one may

predict it should broadly enhance multiple classes of antibiotics (70). However, our prior study

demonstrated that triclosan only enhanced aminoglycosides and to a lesser extent, tetracycline

(28). Because triclosan does not completely abolish the $\Delta\psi$, we hypothesize that triclosan only

enhances antibiotic killing by bacteria that are known to trigger adaptive resistance through

disrupting translation, such as aminoglycosides and tetracyclines (8-10, 12-15).

*P. aeruginosa* encodes two genes for elongation factor G (EF-G), *fusA1* and *fusA2* (71).

*In vitro* translation studies have suggested that *fusA2* is the primary EF-G responsible for

ribosome translocation while the function of *fusA1* remains unclear (71). *fusA1* and *fusA2* are

98% similar and 84% identical, but interestingly *fusA1* expression is induced in cells growing as

biofilms compared to cells growing planktonically (72). Recently, a new tobramycin resistance

mechanism caused by mutations in *fusA1* has been identified in CF *P. aeruginosa* isolates (73).

In fact, one of the mutations that we report here, L40Q, was identified in clinical isolates (73).

These results support our findings that mutations in *fusA1* reduce susceptibility to

aminoglycosides through an unknown mechanism(s).

In addition to mutations in the gene *fusA1*, all resistant mutants had SNPs or deletions in

the previously described genes *ptsP*, involved in the overproduction of the phenazine pyocyanin

responsible for the generation of reactive oxygen species (74), *mexT*, a repressor of the RND-

type efflux pump MexEF-OprN (75), and *wspF*, a regulatory of the diguanylate cyclase WspR

(76-78) (Table 1). Mutations in *mexT* were deletions or the SNP P170L (Table 1). These

mutations and deletions likely result in the overexpression of the RND-type efflux pump

MexEF-OprN due to loss of repression by MexT (75). This finding supports our model that the

inhibition of efflux pump activity by triclosan drives increased tobramycin accumulation and

activity. It is unclear what benefits would be provided by increased pyocyanin production, a

quorum sensing controlled metabolite that is toxic and plays numerous roles in *P. aeruginosa*

physiology and virulence (79). There is evidence to suggest pyocyanin may restrict social

cheating in biofilms and promote tolerance to oxidative stress, which may provide some

tolerance to tobramycin and triclosan treatment (80). Mutations in *wspF* are not surprising as

these promote the formation of highly tolerant biofilms by increasing intracellular cyclic di-GMP

stimulating increased EPS production that reduces the diffusion of tobramycin (76-78).

Owing to decades of overuse, the FDA has restricted the use of triclosan due to concerns

over bioaccumulation and the potential for induction of resistance to other antibiotics in bacteria.

However, triclosan maintains FDA approval for use in Colgate® total toothpaste at 100-times the

concentrations used in this study. In support of this decision, numerous toxicity studies and the

Scientific Committee on Consumer Safety published by the European Union has found that

triclosan is safe when used appropriately (40-42). Furthermore, envisioning triclosan as an

adjuvant for tobramycin therapy for CF, we previously performed acute and long-term triclosan

toxicity studies in rats and showed that direct delivery of triclosan to lungs did not elicit

significant toxicity (28).

More broadly, our results suggest that protonophores could be potent antibiotic adjuvants

that enhance antibiotic killing against bacterial biofilms. Protonophores are being developed as

novel antibiotics to target *Mtb*. Specifically, diphenyl ethers such as SQ109 are in phase II

clinical trials in humans (24-27, 64). Moreover, a HTS recently identified that the c-Jun N-

terminal kinase inhibitor SU3327, renamed halicin, dissipates the ΔpH and PMF, demonstrating

growth inhibitory properties both *in vitro* and *in vivo* against several pathogens including some

multi-drug resistant strains (66). Importantly, protonophore uncouplers are already routinely used

for parasitic infections, including oxyclozanide, which we previously found synergizes with

tobramycin much like triclosan (39). In addition to protonophores functioning as antimicrobials,

there is also renewed interest in using these compounds to target such diseases as diabetes and

cancer, as well as slow aging, suggesting compounds with this activity are druggable targets

when used appropriately (81-83). Together, our results and other published studies demonstrate

that the use of protonophore uncouplers to target bacterial energetics is a promising new strategy

to target antimicrobial resistant infections (24).

**Methods**

**Bacterial strains, culture conditions, and compounds.**

All strains used in this study are listed in Table 1. Bacterial strains were grown in glass

test tubes (18 X 150 mm) at 35ºC in cation adjusted Müeller-Hinton Broth II (MHB, Sigma-

Aldrich) with agitation at 210 revolutions per minute (RPM). Antibiotics, methyl-triclosan,

carbonyl cyanide 3-chlorophenylhydrazone (CCCP), and triclosan were obtained from Sigma-

Aldrich. AFN-1252 was obtained from MedChemExpress. Tobramycin sulfate, gentamicin

sulfate, and streptomycin sulfate were dissolved in autoclaved deionized water and filter

sterilized using 0.22 µM filter membranes (Thomas Scientific). Triclosan was dissolved in 100%

ethanol. Methyl-triclosan, CCCP, and AFN-1252 were dissolved in 100% dimethyl sulfoxide

(DMSO).

**Biofilm susceptibility testing using BacTiter-Glo™.**

To measure the antimicrobial susceptibility of biofilms, the MBEC Assay® (Innovotech)

was used as described (28). Briefly, an overnight culture was diluted and seeded into a MBEC

plate in 10% MHB v/v diluted in DPBS and incubated for 24-hrs at 35ºC with agitation at 150

RPM. The MBEC lid was then washed to remove non-adherent cells, transferred to a 96-well

treatment plate, and incubated for the indicated time at 35ºC without agitation. Following

treatment, the MBEC lid was washed and transferred to a black 96-well ViewPlate (PerkinElmer)

filled with 40% (v/v) BacTiter-Glo™ (Promega) diluted in DPBS to enumerate cell viability

using luminescence by an EnVison Multilabel Plate Reader (PerkenElmer, Waltham, MA). The

BacTiter-Glo™ Microbial Cell Viability Assay is a luminescent assay that determines the

number of viable cells based on quantification of ATP concentration. A calibration curve was

previously performed and it was found using a linear regression the coefficient of determination

was $r^2$=0.9884 for luminescence versus CFUs/mL (28).

**Selection for *P. aeruginosa* resistant mutants.**

To select mutants resistant to tobramycin and triclosan, we modified a protocol by

Lindsey and collages and serially passaged biofilms while treating them with ever increasing

concentrations of triclosan and tobramycin (Table S1) (44). 24-hr biofilms were formed as

described above and treated with triclosan and tobramycin for 24-hrs. After each treatment,

biofilms were sonicated for 15-mins (Branson Ultrasonics) to disperse surviving cells from the

pegs into 10% v/v MHB recovery media. Biofilms were then allowed to re-form on new pegs

overnight. The next day, passaged biofilms were treated at a slightly higher concentration of

triclosan and tobramycin. Following the treatment, the recovery process was repeated, and

biofilms were re-formed. We split the treatment groups into three regimens: gradual, moderate

and sudden (Table S1). The gradual treatment group was initially treated with 0.4 µM of

triclosan and 0.004 µM of tobramycin, which is 250x less than the MIC of tobramycin (28). The

moderate treatment group was initially treated with 8 µM of triclosan and 0.08 µM of

tobramycin, which is 12.5x less than the MIC of tobramycin (28). And then the two treatment

groups were treated in subsequent cycles with ever increasing concentrations of triclosan and

tobramycin at the same rate. The sudden treatment series was treated with 500 µM of triclosan

and tobramycin from cycle day 1 and was eradicated. By serially passaging biofilms and

gradually increasing the concentration of triclosan and tobramycin, two resistant mutant

populations were established, 30_B2 and 30_F1 (Cycle_Location on 96-well plate). Each mutant

pool was then streaked on LB agar plates for single colony isolation, and 96 single colony

mutants were isolated from each plate, totaling 192 single colony mutants.

**Whole genome sequencing.**

Genomic deoxyribonucleic acid (gDNA) was isolated from each mutant using the

Wizard® gDNA purification kit (Promega). Illumina NextSeq was then performed by the

Genomic Services Facility at Indiana University Center for Genomics and Bioinformatics. To

identify single nucleotide polymorphisms (SNPs), sequencing results were first verified for

quality using FASTQC and aligned to the PAO1 reference genome (84), which can be

downloaded from the *Pseudomonas* Genome Database (http://www.pseudomonas.com) using the

*Breseq* pipeline, which can be downloaded from (http://barricklab.org) (85). All identified SNPs

are listed in the supplemental spreadsheet.

**Ethidium bromide efflux pump assay.**

Intercellular accumulation of ethidium bromide, which is a substrate for RND-type efflux

pumps, was measured fluorometrically as previously described (49, 86). As described above, 24-

hr old biofilms were formed in 96-well black ViewPlate. 10 µg/mL of ethidium bromide was

added to each well along with various treatments. Fluorescence was recorded every 2-hrs for 6-

hrs using a SpectraMax M5 microplate spectrophotometer system ($\lambda_{excite}/\lambda_{emit}$ 530/600 nm).

**494** **BacLight™ membrane potential assay and cell permeabilization assay.**

**495**      To measure changes in membrane potential and cellular permeabilization the BacLight™

**496** Membrane Potential Assay was used in combination with the cell impermeable live/dead TO-

**497** Pro™-3 iodide stain, as previously described (39). Briefly, 24-hr old biofilms were formed in

**498** glass test tubes (18 x 150 mm) in 1 mL of 10% (v/v) MHB at 35°C and agitated at 150 RPM.

**499** Cells were washed in DPBS to remove non-adherent cells and treated with triclosan and

**500** tobramycin for 2-hrs. Following treatment, cells were washed in DPBS (without magnesium and

**501** calcium), and the biofilm was disrupted from the air-liquid interface. The cells were stained in 1

**502** mL of DPBS for 5-mins with DiOC2(3). This dye emits in the fluorescein isothiocyanate (FITC)

**503** channel within all cells. However, greater membrane potential drives accumulation and stacking

**504** of the dye in the cell cytoplasm, shifting emission to the phycoerythrin (PE) channel. TO-

**505** PRO™-3 iodide, which emits in the allophycocyanin (APC) channel, was used to identify

**506** permeabilized cells, which fluoresces in cells that have compromised membranes by

**507** intercalating DNA. Single cell flow cytometry was performed on an LSR II (BD Biosciences),

**508** with excitation from 488 mm and 640 mm lasers, and analyzed in FITC/PE and APC channels,

**509** respectively.

**510**

**511** **Tobramycin accumulation and extrusion assay.**

**512**      To measure the accumulation of tobramycin within cells in biofilms, tobramycin was

**513** conjugated to Texas red (Sigma-Aldrich) using an amine conjugation reaction, as previously

**514** described (50). Briefly, one milligram of Texas Red sulfonyl chloride (Thermo Fisher Scientific)

**515** was resuspended in 50 μL of anhydrous N,N-dimethylformamide (Sigma-Aldrich) on ice. The

solution was added slowly to 2.3 mL of 100 mM K2CO3 at pH 8.5, with or without 10 mg/mL

tobramycin (Sigma-Aldrich), on ice. Conjugated tobramycin was used at a concentration of 250

µg/mL (~500 µM) alone and in combination with 100 µM of triclosan against 24-hr old biofilms

formed in glass test tubes at the air-liquid interface, as described above. Following treatments,

biofilms were washed in DPBS and then disrupted with autoclaved wooden sticks into 1 mL of

0.2% Triton X-100 to lyse cells (Sigma-Aldrich). Lysed cells were then transferred to

spectrophotometer cuvettes (Thermo Fisher Scientific) and read using a SpectraMax M5

microplate spectrophotometer system ($\lambda_{excite}/\lambda_{emit}$ 595/615 nm). Extrusion assays were performed

in a similar fashion, except after 30-mins of treatment, biofilms were washed three times in

DPBS and then allowed to recover in treatment-free 1% MHB media for 30-mins. After

recovery, the media was read as described above.

**DPBS-starved *P. aeruginosa* biofilms.**

24-hr old biofilms were formed on MBEC™ plates as described above. After 24-hrs, the

lid was washed for 5-mins to remove non-adherent cells, transferred to a 96-well plate containing

DPBS, and incubated at 35ºC with agitation at 150 RPM for 5-days.

**Agarose hydrogels.**

Agarose hydrogels were made as previously described (56). Briefly, by dissolving 1 gm

of agarose (Sigma-Aldrich) into 200 mL of Tris-acetate-EDTA (TAE) buffer and heated to form

a homogenous solution using a microwave. The 0.5% agarose solution was then allowed to cool

and various treatments were added. The solution was then poured into 100 x 15 mm petri dishes

(Thermo Fisher Scientific) and stored at 4°C overnight. Prior to treatment, a 4 mm biopsy punch

(VWR) was used to create hydrogel wafers.

**Murine wound infection model.**

Wound surgery was performed on 8-9 week-old male and female SKH-1 mice (Charles

River) and infected with a bioluminescent derivative of PAO1, Xen41 (PerkinElmer), which

constitutively expresses *luxCDABE* gene*,* as previously described with the following

modifications (55, 56). Briefly, 24-hr old biofilms were formed on sterilized polycarbonate

membrane filters with a 0.2 µM pore size (Millipore Sigma) by diluting an overnight culture to

an $OD_{600}$ of 0.001 and pipetting 100 µl on 4 membranes on a tryptic soy agar (TSA) plate. 24-hr

old biofilms were scrapped using L-shaped spreaders (Sigma-Aldrich) from each membrane and

re-suspended in 500 µl of DPBS. 20 µl of the biofilm-suspension was inoculated into the 24-hr

old wounds previously formed on the dorsal side of the mouse midway between the head and the

base of the tail. 24-hrs later the biofilm within the wound was imaged using *in vivo* imaging

system (IVIS) (Perkin Elmer). Then the biofilm was treated by placing a 4 mm 0.5% agarose

hydrogel wafer either containing various treatments or only being made up of agarose alone on

top of the wound. To quantify antimicrobial activity, the wound was then imaged 4-hrs later

using IVIS and total flux (p/s) in radiance was used to quantify bacterial susceptibility. The loss

of radiance indicates fewer cells are present within the wound. Vertebrate research was approved

by the Michigan State University Institutional Animal Care and Use Committee application

03/18-036-00.

## Statistics

Significance was determined as described in accompanying figure legends by either a one-way ANOVA or two-way ANOVA followed by Sidak's or Bonferroni's multiple comparisons posttest, respectively. Unpaired t-test were also used as described in accompanying figure legends. The "n" indicates the number of biological replicates. For murine wound infection experiments, sample sizes were determined based on preliminary studies using groups that were sufficient to achieve a desired power of 0.85 and an alpha value of 0.05. All analyses were performed using GraphPad Prism version 8.4.1 (San Diego, CA) and a $p$-value of $< 0.05$ was considered statistically significant.

## Acknowledgments

This work was supported by grants from the Hunt for a Cure Foundation, the NSF BEACON center for evolution in action (DBI-0939454), NIH grants GM109259, GM110444, and AI143098 to C.M.W. and the Cystic Fibrosis Foundation Traineeship to M.M.M. In addition, M.M.M. was supported by a Wentworth Fellowship and Rudolf Hugh award from the MSU Department of Microbiology and Molecular Genetics and by a Dissertation Completion Fellowship from the College of National Science. We thank both Sandra O'Reilly and Louis King of the Michigan State Research Technology Support Facility for their assistance in the wound model and flow cytometry experiments, respectively. We also thank the Genomic Services Facility at Indiana University Center for Genomics and Bioinformatics. And we thank Lucas Demey, Mitchell Zachos and Emily Horning for their technical assistance.

582  **Table**

584  **Table 1. Bacterial strains used in this study.**

| Strain | Characteristics | Reference |
|--------|-----------------|-----------|
| PAO1 | *P. aeruginosa* Standard Reference Strain | (28) |
| Xen41 | Bioluminescent PAO1 derivative: constitutively expresses *luxCDABE* gene | PerkinElmer |
| 30_F1_5 | *fusA1*, L40Q (C**T**G→C**A**G)<br>*wspF,* Q132* (CAG→TAG)<br>*PtsP,* A718P (**G**CC→**C**CC)<br>*mexT,* P170L (C**C**G→C**T**G) | This Study |
| 30_F1_6 | *fusA1*, L40Q (C**T**G→C**A**G)<br>*wspF*, Q132* (CAG→TAG)<br>*PtsP,* coding (703/2280 nt)<br>*mexT,* coding (58-137/1044 nt) | This Study |
| 30_B2_8 | *fusA1*, T64A (**A**CC→**G**CC)<br>*wspF*, Q132* (CAG→TAG)<br>*PtsP,* coding (703/2280 nt)<br>*mexT,* coding (58-137/1044 nt) | This Study |
| 30_B2_30 | *fusA1*, T64A (**A**CC→**G**CC)<br>*wspF,* Q132* (CAG→TAG)<br>*PtsP,* A718P (**G**CC→**C**CC)<br>*mexT,* P170L (C**C**G→C**T**G) | This Study |
| 30_B2_57 | *fusA1*, T64A (**A**CC→**G**CC)<br>*wspF,* Q132* (CAG→TAG)<br>*PtsP,* coding (703/2280 nt)<br>*mexT,* coding (58-137/1044 nt) | This Study |
| 30_B2_82 | *fusA1*, T64A (**A**CC→**G**CC)<br>*wspF,* Q132* (CAG→TAG)<br>*PtsP,* A718P (**G**CC→**C**CC)<br>*mexT,* P170L (C**C**G→C**T**G) | This Study |

585  All additional SNPs in evolution mutants are listed in the supplemental spreadsheet.

	**Figure legends**

	**FIG 1. Triclosan in combination with tobramycin results in increased permeabilization of**

	**cells within biofilms.** 24-hr old biofilms were treated with triclosan (100 µM), or tobramycin

	(500 µM), alone and in combination for 2-hrs. Cells were stained with the live/dead cell indicator

	TO-PRO™-3 iodide to determine the number of cells that were permeabilized. The results are

	percent averages plus the standard error mean (SEM, n=4). Percent values indicate the average

	relative abundance of events within each gate normalized to the total number of events analyzed,

	excluding artifacts, aggregates and debris. A one-way analysis of variance (ANOVA) followed

	by Sidak's multiple comparison post-hoc test was used to determine statistical significance

	between each treatment and the untreated control or tobramycin treatment alone with the

	combination treatment. *, $p < 0.05$. NS, not significant.

	**FIG 2. Mutants from experimental evolution are resistant to tobramycin and the**

	**combination.** 24-hr old biofilms were treated with triclosan (100 µM) and tobramycin (500 µM)

	alone and in combination. The results represent the means plus SEM (n=6). A two-way ANOVA

	followed by Bonferroni's posttest was used to determine statistical significance between each

	treatment and the untreated control. *, $p < 0.05$. NS, not significant.

	**FIG 3. Triclosan reduces the activity of RND-type efflux pumps.** 24-hr biofilms were stained

	with ethidium bromide to measure accumulation. Biofilms were treated with CCCP (100 µM),

	triclosan (100 µM), tobramycin (500 µM) alone and in combination. Fluorescence was read at

	0,2,4, and 6-hrs. Results represent the average arbitrary fluorescence units ±SEM (n=6).

**FIG 4. Triclosan results in increased cellular accumulation and extrusion of Texas Red**

**conjugated tobramycin (TbTR).** 24-hr old biofilms grown in glass test tubes were treated for

30-mins with triclosan (100 µM), CCCP (100 µM) and TbTR (250 µg/mL). (A) Then cells were

disrupted from the biofilms and lysed with 0.2% Triton-X 100® to measure intercellular

accumulation of TbTR. (B) To measure extrusion, biofilms were first treated for 30-mins and

washed three-times in DPBS. Biofilms then recovered in treatment free media for 30-mins and

fluorescence of the media was measured. TbTR was measured by relative fluorescence units

using excitation $595_{nm}$ and emission $615_{nm}$. Results represent the average arbitrary fluorescence

units ±SEM (n=6). For panel A, a One-Way-ANOVA was performed comparing TbTR alone vs

CCCP and TbTR and vs triclosan and TbTR. For panel B, an unpaired t-test was performed

comparing TbTR versus triclosan and TbTR. *, $p < 0.05$. NS, not significant.

**FIG 5. Triclosan reduces the membrane potential surge induced by tobramycin at 2-hours.**

24-hr old biofilms were treated with triclosan (100 µM), or tobramycin (500 µM), alone and in

combination for 2-hrs. Cells were stained with DiOC2(3) to measure the membrane potential.

Dead or permeabilized cells were excluded from membrane potential analysis by parallel

staining with the dead cell indicator TO-PRO™-3 iodide. The results are percent averages plus

the SEM (n=4). Percent values indicate the average relative abundance of events within each

gate normalized to the total number of events analyzed, excluding dead cells, artifacts,

aggregates and debris. A one-way ANOVA followed by Sidak's multiple comparison post-hoc

test was used to determine statistical significance between each treatment and the untreated

control, and tobramycin treatment alone was compared with the combination. *, $p < 0.05$. NS, not

significant.

**FIG 6. Triclosan and tobramycin hydrogels are more effective than tobramycin hydrogels in a murine wound model.** 24-hr old bioluminescent biofilms formed within wounds were treated with triclosan (100 µM), or tobramycin (400 µM), alone and in combination for 4-hrs in a hydrogel. Reduction in the number of cells within biofilms was quantified using IVIS. The results are fold reduction of three separate experiments ±SEM, no treatment n=6, triclosan n=6, tobramycin n=10, triclosan and tobramycin n=9. A one-way ANOVA followed by Bonferroni's multiple comparison post-hoc test was used to determine statistical significance between each treatment and the untreated control and tobramycin treatment alone was compared with tobramycin and triclosan (*, $p<0.05$).

**FIG 7. Triclosan sensitizes *P. aeruginosa* to tobramycin by acting as a protonophore, inhibiting efflux pump activity, and abolishing adaptive resistance.** (1) Within 2-hrs of exposure to tobramycin, adaptive resistance occurs, (2) which is due to the induction of RND-type efflux pumps and surge in membrane potential ($\Delta\psi$), resulting in reduced accumulation of tobramycin within the cytosol. (3) Triclosan shuttles protons across the inner membrane, collapsing the proton motive force ($\Delta p$) and depolarizing the $\Delta\psi$. Consequently, efflux pump activity is reduced and there is enhanced accumulation of tobramycin within the cytosol. Finally, tobramycin binds to the A-site of the ribosome, corrupting protein synthesis and causing membrane permeabilization. Overall, triclosan accelerates and increase the effectiveness of tobramycin by reducing the $\Delta\psi$ and efflux pump activity. Proton gradient, $\Delta pH$. Outer membrane protein (OMP), membrane fusion protein (MFP), inner membrane protein (IMP).

**Supplemental Figure Legends**

**FIG S1. AFN-1252 alone or in combination with tobramycin is not effective against mature**

**biofilms.** 24-hr old biofilms grown on MBEC plates were treated for 6-hrs with AFN-1252 (400

µM), tobramycin (400 µM), alone and in combination in two-fold dilutions, and the number of

viable cells within the biofilms were quantified by BacTiter-Glo$^{TM}$. The assay was performed

twice in in duplicate. The results represent means ± the Standard Error Mean (SEM). A one-way

analysis of variance (ANOVA) followed by Bonferroni's multiple comparison post-hoc test was

used to determine statistical significance between tobramycin versus the combination (NS, not

significant).

**FIG S2. Initial tobramycin treatment is required for the combination to be effective.** 24-hr

old biofilms grown on MBEC plates were treated sequentially. First, biofilms were treated for 3-

hours with tobramycin (500 µM) and then washed three times in DPBS for 3-mins each, before

being treated with triclosan (100 µM) for 3-hours or vice versa. As a control, biofilms were also

treated for 6-hrs with triclosan and tobramycin. The number of viable cells within the biofilms

were quantified by BacTiter-Glo$^{TM}$. The assay was performed twice in in duplicate. The results

represent means plus the SEM. A one-way ANOVA followed by Bonferroni's multiple

comparison post-hoc test was used to determine statistical significance between each treatment

and the untreated control. *, $p < 0.05$. NS, not significant.

**FIG S3. Methyl-triclosan alone or in combination with tobramycin is not effective against**

**mature biofilms.** 24-hr old biofilms grown on MBEC plates were treated for 6-hrs with methyl-

triclosan (100 µM), tobramycin (400 µM), alone and in combination in two-fold dilutions, and

the number of viable cells within the biofilms were quantified by BacTiter-Glo[TM]. The assay was

performed twice in in duplicate. The results represent means ±SEM. A one-way ANOVA

followed by Bonferroni's multiple comparison post-hoc test was used to determine statistical

significance compared to tobramycin alone and the combination. NS, not significant.

**FIG S4. Methyl-triclosan alone and in combination with tobramycin does not reduced the**

**surge in membrane potential induced by tobramycin at 2-hours.** 24-hr old biofilms were

treated with methyl-triclosan (100 µM), or tobramycin (500 µM), alone and in combination for

2-hrs. Cells were stained with DiOC2(3) and TO-PRO™-3 iodide to determine the number of

cells that maintained a membrane potential (A) or were permeabilized (B), respectively. Dead or

permeabilized cells were excluded from membrane potential analysis and are shown in panel B.

The experiment was performed two separate times in duplicate. The results are percent averages

plus the SEM. Percent values indicate the average relative abundance of events within each gate

normalized to the total number of events analyzed, excluding artifacts, aggregates and debris. A

one-way ANOVA followed by Sidak's multiple comparison post-hoc test was used to determine

statistical significance between each treatment and an untreated control, and tobramycin

treatment alone was compared with the combination. *, $p<0.05$, NS, not significant.

**FIG S5. Methyl-triclosan does not reduced RND-type efflux pump activity.** Ethidium

bromide is a substrate of RND-type efflux pumps. 24-hr biofilms were stained with ethidium

bromide to measure accumulation. Biofilms were treated with methyl-triclosan (100 µM) and

tobramycin (500 µM) alone and in combination. Fluorescence was read at 0,2,4, and 6-hrs. The

assay was performed three times in triplicate. Results represent the average arbitrary

fluorescence units ±SEM.

**FIG S6. DPBS starved biofilms are not more sensitive to tobramycin alone.** 24-hour old

biofilms were starved of nutrients by replacing media with DPBS for 5-days. Starved biofilms

were treated with triclosan (100 µM), or tobramycin (500 µM), alone and in combination for 6-

hrs. The number of viable cells within the biofilms was quantified using the BacTiter-Glo™

assay. The assay was performed once using three biological replicates. The results represent the

means plus the SEM. A one-way ANOVA followed by Sidak's multiple comparison post hoc

test was used to determine statistical significance between the combination treatment and the

untreated control and between triclosan alone or tobramycin alone and the combination treatment

as indicated by the bars. *, $P < 0.05$.

**FIG S7. Tobramycin is as effective as the combination after 24-hrs of treatment.** 24-hr old

biofilms were treated for 24-hrs with triclosan (100 µM), tobramycin (500 µM), alone and in

combination, and the number of viable cells within the biofilms were quantified by BacTiter-

Glo™. The assay was performed at least three times in triplicate. The results represent means

plus the SEM. A one-way ANOVA followed by Bonferroni's multiple comparison post-hoc test

was used to determine statistical significance between untreated biofilms, tobramycin alone, and

the combination treatment. In addition, using the same analysis, a comparison was made between

tobramycin alone and the combination treatment. *, $P < 0.05$. NS, not significant. 4-hr

treatments were previously published and are shown for comparison (28).

**References:**

1.      Donlan RM, Costerton JW. Biofilms: survival mechanisms of clinically relevant microorganisms. Clin Microbiol Rev. 2002;15(2):167-93.

2.      O'Sullivan BP, Freedman SD. Cystic fibrosis. The Lancet. 2009;373(9678):1891-904.

3.      Omar A, Wright JB, Schultz G, Burrell R, Nadworny P. Microbial Biofilms and Chronic Wounds. Microorganisms. 2017;5(1):9.

4.      Poole K. *Pseudomonas aeruginosa*: resistance to the max. Front Microbiol. 2011;2:65.

5.      Livermore DM. Penicillin-binding proteins, porins and outer-membrane permeability of carbenicillin-resistant and -susceptible strains of *Pseudomonas aeruginosa*. J Med Microbiol. 1984;18(2):261-70.

6.      Morita Y, Tomida J, Kawamura Y. Responses of *Pseudomonas aeruginosa* to antimicrobials. Front Microbiol. 2014;4.

7.      Barclay ML, Begg EJ, Chambers ST, Thornley PE, Pattemore PK, Grimwood K. Adaptive resistance to tobramycin in *Pseudomonas aeruginosa* lung infection in cystic fibrosis. J Antimicrob Chemother. 1996;37(6):1155-64.

8.      Hay T, Fraud S, Lau CH, Gilmour C, Poole K. Antibiotic inducibility of the *mexXY*

multidrug efflux operon of *Pseudomonas aeruginosa*: involvement of the MexZ anti-repressor

ArmZ. PLoS One. 2013;8(2):e56858.

9.      Jeannot K, Sobel ML, El Garch F, Poole K, Plesiat P. Induction of the MexXY efflux pump

in *Pseudomonas aeruginosa* is dependent on drug-ribosome interaction. J Bacteriol.

2005;187(15):5341-6.

10.     Hocquet D, Vogne C, El Garch F, Vejux A, Gotoh N, Lee A, et al. MexXY-OprM efflux

pump is necessary for a adaptive resistance of *Pseudomonas aeruginosa* to aminoglycosides.

Antimicrobial agents and chemotherapy. 2003;47(4):1371-5.

11.     Karlowsky JA, Saunders MH, Harding GA, Hoban DJ, Zhanel GG. In vitro

characterization of aminoglycoside adaptive resistance in *Pseudomonas aeruginosa*. Antimicrob

Agents Chemother. 1996;40(6):1387-93.

12.     Yamamoto M, Ueda A, Kudo M, Matsuo Y, Fukushima J, Nakae T, et al. Role of MexZ

and PA5471 in transcriptional regulation of *mexXY* in *Pseudomonas aeruginosa*. Microbiology.

2009;155(Pt 10):3312-21.

13.     Barclay ML, Begg EJ. Aminoglycoside adaptive resistance: importance for effective

dosage regimens. Drugs. 2001;61(6):713-21.

14.    Barclay ML, Begg EJ, Chambers ST. Adaptive resistance following single doses of

gentamicin in a dynamic in vitro model. Antimicrob Agents Chemother. 1992;36(9):1951-7.

15.    Barclay ML, Begg EJ, Chambers ST, Peddie BA. The effect of aminoglycoside-induced

adaptive resistance on the antibacterial activity of other antibiotics against *Pseudomonas*

*aeruginosa* in vitro. J Antimicrob Chemother. 1996;38(5):853-8.

16.    Daikos GL, Lolans VT, Jackson GG. First-exposure adaptive resistance to aminoglycoside

antibiotics in vivo with meaning for optimal clinical use. Antimicrob Agents Chemother.

1991;35(1):117-23.

17.    Xiong YQ, Caillon J, Kergueris MF, Drugeon H, Baron D, Potel G, et al. Adaptive

resistance of *Pseudomonas aeruginosa* induced by aminoglycosides and killing kinetics in a rabbit

endocarditis model. Antimicrobial agents and chemotherapy. 1997;41(4):823-6.

18.    Taber HW, Mueller JP, Miller PF, Arrow AS. Bacterial uptake of aminoglycoside

antibiotics. Microbiol Rev. 1987;51(4):439-57.

19.    Davis BD. Mechanism of bactericidal action of aminoglycosides. Microbiol Rev.

1987;51(3):341-50.

20.    Hancock RE. Alterations in outer membrane permeability. Annu Rev Microbiol.

1984;38(1):237-64.

21.    Mogayzel PJ, Jr., Naureckas ET, Robinson KA, Brady C, Guill M, Lahiri T, et al. Cystic

Fibrosis Foundation pulmonary guideline. pharmacologic approaches to prevention and

eradication of initial *Pseudomonas aeruginosa* infection. Ann Am Thorac Soc. 2014;11(10):1640-

50.

22.    Chmiel JF, Aksamit TR, Chotirmall SH, Dasenbrook EC, Elborn JS, LiPuma JJ, et al.

Antibiotic management of lung infections in cystic fibrosis. I. The microbiome, methicillin-

resistant *Staphylococcus aureus*, gram-negative bacteria, and multiple infections. Ann Am Thorac

Soc. 2014;11(7):1120-9.

23.    Brown ED, Wright GD. Antibacterial drug discovery in the resistance era. Nature.

2016;529(7586):336-43.

24.    Hards K, Cook GM. Targeting bacterial energetics to produce new antimicrobials. Drug

Resist Updat. 2018;36:1-12.

25.    Andries K, Verhasselt P, Guillemont J, Gohlmann HW, Neefs JM, Winkler H, et al. A

diarylquinoline drug active on the ATP synthase of *Mycobacterium tuberculosis*. Science.

2005;307(5707):223-7.

26.    Hards K, Robson JR, Berney M, Shaw L, Bald D, Koul A, et al. Bactericidal mode of

action of bedaquiline. J Antimicrob Chemother. 2015;70(7):2028-37.

27.    Feng X, Zhu W, Schurig-Briccio LA, Lindert S, Shoen C, Hitchings R, et al. Antiinfectives

targeting enzymes and the proton motive force. Proc Natl Acad Sci U S A. 2015;112(51):E7073-

82.

28.    Maiden MM, Hunt AMA, Zachos MP, Gibson JA, Hurwitz ME, Mulks MH, et al.

Triclosan Is an Aminoglycoside Adjuvant for Eradication of *Pseudomonas aeruginosa* Biofilms.

Antimicrob Agents Chemother. 2018;62(6):e00146-18.

29.    Heath RJ, Rubin JR, Holland DR, Zhang E, Snow ME, Rock CO. Mechanism of triclosan

inhibition of bacterial fatty acid synthesis. J Biol Chem. 1999;274(16):11110-4.

30.    Zhu L, Lin J, Ma J, Cronan JE, Wang H. Triclosan resistance of *Pseudomonas aeruginosa*

PAO1 is due to FabV, a triclosan-resistant enoyl-acyl carrier protein reductase. Antimicrob Agents

Chemother. 2010;54(2):689-98.

31.    Mima T, Joshi S, Gomez-Escalada M, Schweizer HP. Identification and characterization

of TriABC-OpmH, a triclosan efflux pump of *Pseudomonas aeruginosa* requiring two membrane

fusion proteins. J Bacteriol. 2007;189(21):7600-9.

32.    Mitchell P. Coupling of phosphorylation to electron and hydrogen transfer by a chemi-

osmotic type of mechanism. Nature. 1961;191(4784):144-8.

33.    Mitchell P. Chemiosmotic coupling in oxidative and photosynthetic phosphorylation.

1966. Biochim Biophys Acta. 2011;1807(12):1507-38.

34.    Yoon DS, Choi Y, Cha DS, Zhang P, Choi SM, Alfhili MA, et al. Triclosan Disrupts SKN-

1/Nrf2-Mediated Oxidative Stress Response in C. elegans and Human Mesenchymal Stem Cells.

Sci Rep. 2017;7(1):12592.

35.    Weatherly LM, Shim J, Hashmi HN, Kennedy RH, Hess ST, Gosse JA. Antimicrobial

agent triclosan is a proton ionophore uncoupler of mitochondria in living rat and human mast cells

and in primary human keratinocytes. J Appl Toxicol. 2016;36(6):777-89.

36.    Popova LB, Nosikova ES, Kotova EA, Tarasova EO, Nazarov PA, Khailova LS, et al.

Protonophoric action of triclosan causes calcium efflux from mitochondria, plasma membrane

depolarization and bursts of miniature end-plate potentials. Biochim Biophys Acta Biomembr.

2018;1860(5):1000-7.

37.    Teplova VV, Belosludtsev KN, Kruglov AG. Mechanism of triclosan toxicity:

Mitochondrial dysfunction including complex II inhibition, superoxide release and uncoupling of

oxidative phosphorylation. Toxicol Lett. 2017;275:108-17.

38.    Ajao C, Andersson MA, Teplova VV, Nagy S, Gahmberg CG, Andersson LC, et al.

Mitochondrial toxicity of triclosan on mammalian cells. Toxicol Rep. 2015;2:624-37.

39.    Maiden MM, Zachos MP, Waters CM. The ionophore oxyclozanide enhances tobramycin

killing of *Pseudomonas aeruginosa* biofilms by permeabilizing cells and depolarizing the

membrane potential. J Antimicrob Chemother. 2019;74(4):894-906.

40.    Davison John SR, Maillard M, Pagès J-Y, Scenihr J-MO, et al. Scientific Committee on

Consumer Safety (SCCS): Opinion on triclosan antimicrobial resistance. DG Sanco Scientific

Committee on Consumer Safety. . 2010 22:1–56. .

41.    Rodricks JV, Swenberg JA, Borzelleca JF, Maronpot RR, Shipp AM. Triclosan: a critical

review of the experimental data and development of margins of safety for consumer products. Crit

Rev Toxicol. 2010;40(5):422-84.

42.    Fang JL, Stingley RL, Beland FA, Harrouk W, Lumpkins DL, Howard P. Occurrence,

efficacy, metabolism, and toxicity of triclosan. J Environ Sci Health C Environ Carcinog

Ecotoxicol Rev. 2010;28(3):147-71.

43.    Rao KN, Lakshminarasimhan A, Joseph S, Lekshmi SU, Lau MS, Takhi M, et al. AFN-

1252 is a potent inhibitor of enoyl-ACP reductase from *Burkholderia pseudomallei*--Crystal

structure, mode of action, and biological activity. Protein Sci. 2015;24(5):832-40.

44.    Lindsey HA, Gallie J, Taylor S, Kerr B. Evolutionary rescue from extinction is contingent

on a lower rate of environmental change. Nature. 2013;494(7438):463-7.

45.    Pal C, Papp B, Lazar V. Collateral sensitivity of antibiotic-resistant microbes. Trends

Microbiol. 2015;23(7):401-7.

46.    Rodnina MV, Savelsbergh A, Katunin VI, Wintermeyer W. Hydrolysis of GTP by

elongation factor G drives tRNA movement on the ribosome. Nature. 1997;385(6611):37-41.

47.    Arnaud Bolard PPs, and Katy Jeannota. Mutations in Gene fusA1 as a Novel Mechanism

of Aminoglycoside Resistance in Clinical Strains of *Pseudomonas aeruginosa*. Antimicrobial

Agents and Chemotherapy. 2018;62(2):e01835-17.

48.    Tseng BS, Zhang W, Harrison JJ, Quach TP, Song JL, Penterman J, et al. The extracellular

matrix protects *Pseudomonas aeruginosa* biofilms by limiting the penetration of tobramycin.

Environ Microbiol. 2013;15(10):2865-78.

49.    Blair JM, Piddock LJ. How to Measure Export via Bacterial Multidrug Resistance Efflux

Pumps. mBio. 2016;7(4):e00840-16.

50.    Sandoval R, Leiser J, Molitoris BA. Aminoglycoside antibiotics traffic to the Golgi

complex in LLC-PK1 cells. J Am Soc Nephrol. 1998;9(2):167-74.

51.    Booth IR. The Regulation of Intracellular pH in Bacteria. Chichester, UK: John Wiley &

Sons, Ltd.; 2007. 19-37 p.

52.    Campbell BD, Kadner RJ. Relation of aerobiosis and ionic strength to the uptake of

dihydrostreptomycin in *Escherichia coli*. Biochim Biophys Acta. 1980;593(1):1-10.

53.    Paulsen IT, Brown MH, Skurray RA. Proton-dependent multidrug efflux systems.

Microbiol Rev. 1996;60(4):575-608.

54.    Venter H, Mowla R, Ohene-Agyei T, Ma S. RND-type drug efflux pumps from Gram-

negative bacteria: molecular mechanism and inhibition. Front Microbiol. 2015;6:377.

55.    Agostinho Hunt AM, Gibson JA, Larrivee CL, O'Reilly S, Navitskaya S, Needle DB, et al.

A bioluminescent *Pseudomonas aeruginosa* wound model reveals increased mortality of type 1

diabetic mice to biofilm infection. J Wound Care. 2017;26(Sup7):S24-S33.

56.    Maiden MM, Zachos MP, Waters CM. Hydrogels Embedded With Melittin and

Tobramycin Are Effective Against *Pseudomonas aeruginosa* Biofilms in an Animal Wound

Model. Front Microbiol. 2019;10(1348):1348.

57.    Allison KR, Brynildsen MP, Collins JJ. Metabolite-enabled eradication of bacterial

persisters by aminoglycosides. Nature. 2011;473(7346):216-20.

58.    Barraud N, Buson A, Jarolimek W, Rice SA. Mannitol enhances antibiotic sensitivity of

persister bacteria in *Pseudomonas aeruginosa* biofilms. PLoS One. 2013;8(12):e84220.

59.    Crabbe A, Ostyn L, Staelens S, Rigauts C, Risseeuw M, Dhaenens M, et al. Host

metabolites stimulate the bacterial proton motive force to enhance the activity of aminoglycoside

antibiotics. PLoS Pathog. 2019;15(4):e1007697.

60.    Radlinski LC, Rowe SE, Brzozowski R, Wilkinson AD, Huang R, Eswara P, et al.

Chemical Induction of Aminoglycoside Uptake Overcomes Antibiotic Tolerance and Resistance

in *Staphylococcus aureus*. Cell Chem Biol. 2019;26(10):1355-64 e4.

61.    Su YB, Peng B, Han Y, Li H, Peng XX. Fructose restores susceptibility of multidrug-

resistant *Edwardsiella tarda* to kanamycin. J Proteome Res. 2015;14(3):1612-20.

62.    Su YB, Peng B, Li H, Cheng ZX, Zhang TT, Zhu JX, et al. Pyruvate cycle increases

aminoglycoside efficacy and provides respiratory energy in bacteria. Proc Natl Acad Sci U S A.

2018;115(7):E1578-E87.

63.    Farha MA, Verschoor CP, Bowdish D, Brown ED. Collapsing the proton motive force to

identify synergistic combinations against *Staphylococcus aureus*. Chem Biol. 2013;20(9):1168-

78.

64.    Foss MH, Pou S, Davidson PM, Dunaj JL, Winter RW, Pou S, et al. Diphenylether-

Modified 1,2-Diamines with Improved Drug Properties for Development against *Mycobacterium*

*tuberculosis*. ACS Infect Dis. 2016;2(7):500-8.

65.    Verstraeten N, Knapen WJ, Kint CI, Liebens V, Van den Bergh B, Dewachter L, et al. Obg

and Membrane Depolarization Are Part of a Microbial Bet-Hedging Strategy that Leads to

Antibiotic Tolerance. Mol Cell. 2015;59(1):9-21.

66.    Stokes JM, Yang K, Swanson K, Jin W, Cubillos-Ruiz A, Donghia NM, et al. A Deep

Learning Approach to Antibiotic Discovery. Cell. 2020;180(4):688-702 e13.

67.    Piddock LJ. Clinically relevant chromosomally encoded multidrug resistance efflux pumps

in bacteria. Clin Microbiol Rev. 2006;19(2):382-402.

68.    Morita Y, Tomida J, Kawamura Y. MexXY multidrug efflux system of *Pseudomonas*
*aeruginosa*. Front Microbiol. 2012;3(408):408.

69.    Mena A, Smith EE, Burns JL, Speert DP, Moskowitz SM, Perez JL, et al. Genetic
adaptation of *Pseudomonas aeruginosa* to the airways of cystic fibrosis patients is catalyzed by
hypermutation. J Bacteriol. 2008;190(24):7910-7.

70.    Askoura M, Mottawea W, Abujamel T, Taher I. Efflux pump inhibitors (EPIs) as new
antimicrobial agents against *Pseudomonas aeruginosa*. Libyan J Med. 2011;6.

71.    Palmer SO, Rangel EY, Hu Y, Tran AT, Bullard JM. Two homologous EF-G proteins from
*Pseudomonas aeruginosa* exhibit distinct functions. PLoS One. 2013;8(11):e80252.

72.    Anderson GG, Moreau-Marquis S, Stanton BA, O'Toole GA. In vitro analysis of
tobramycin-treated *Pseudomonas aeruginosa* biofilms on cystic fibrosis-derived airway epithelial
cells. Infect Immun. 2008;76(4):1423-33.

73.    Bolard A, Plesiat P, Jeannot K. Mutations in Gene fusA1 as a Novel Mechanism of
Aminoglycoside Resistance in Clinical Strains of *Pseudomonas aeruginosa*. Antimicrob Agents
Chemother. 2018;62(2).

74.    Xu H, Lin W, Xia H, Xu S, Li Y, Yao H, et al. Influence of ptsP gene on pyocyanin
production in *Pseudomonas aeruginosa*. FEMS Microbiol Lett. 2005;253(1):103-9.

75.      Juarez P, Broutin I, Bordi C, Plesiat P, Llanes C. Constitutive Activation of MexT by

Amino Acid Substitutions Results in MexEF-OprN Overproduction in Clinical Isolates of

*Pseudomonas aeruginosa*. Antimicrob Agents Chemother. 2018;62(5).

76.      D'Argenio DA, Calfee MW, Rainey PB, Pesci EC. Autolysis and autoaggregation in

*Pseudomonas aeruginosa* colony morphology mutants. J Bacteriol. 2002;184(23):6481-9.

77.      Hickman JW, Tifrea DF, Harwood CS. A chemosensory system that regulates biofilm

formation through modulation of cyclic diguanylate levels. Proc Natl Acad Sci U S A.

2005;102(40):14422-7.

78.      Starkey M, Hickman JH, Ma L, Zhang N, De Long S, Hinz A, et al. *Pseudomonas

aeruginosa* rugose small-colony variants have adaptations that likely promote persistence in the

cystic fibrosis lung. J Bacteriol. 2009;191(11):3492-503.

79.      Lau GW, Hassett DJ, Ran H, Kong F. The role of pyocyanin in *Pseudomonas aeruginosa*

infection. Trends Mol Med. 2004;10(12):599-606.

80.      Castaneda-Tamez P, Ramirez-Peris J, Perez-Velazquez J, Kuttler C, Jalalimanesh A,

Saucedo-Mora MA, et al. Pyocyanin Restricts Social Cheating in *Pseudomonas aeruginosa*. Front

Microbiol. 2018;9(1348):1348.

81.  Barbosa EJ, Lobenberg R, de Araujo GLB, Bou-Chacra NA. Niclosamide repositioning

for treating cancer: Challenges and nano-based drug delivery opportunities. Eur J Pharm

Biopharm. 2019;141:58-69.

82.  Chen W, Mook RA, Jr., Premont RT, Wang J. Niclosamide: Beyond an antihelminthic

drug. Cell Signal. 2018;41:89-96.

83.  Childress ES, Alexopoulos SJ, Hoehn KL, Santos WL. Small Molecule Mitochondrial

Uncouplers and Their Therapeutic Potential. J Med Chem. 2018;61(11):4641-55.

84.  Winsor GL, Griffiths EJ, Lo R, Dhillon BK, Shay JA, Brinkman FS. Enhanced annotations

and features for comparing thousands of *Pseudomonas* genomes in the *Pseudomonas* genome

database. Nucleic Acids Res. 2016;44(D1):D646-53.

85.  Deatherage DE, Barrick JE. Identification of mutations in laboratory-evolved microbes

from next-generation sequencing data using breseq. Methods Mol Biol. 2014;1151:165-88.

86.  Olmsted J, 3rd, Kearns DR. Mechanism of ethidium bromide fluorescence enhancement

on binding to nucleic acids. Biochemistry. 1977;16(16):3647-54.

**FIG 1. Triclosan in combination with tobramycin results in increased permeabilization of cells within biofilms.** 24-hr old biofilms were treated with triclosan (100 μM), or tobramycin (500 μM), alone and in combination for 2-hrs. Cells were stained with the live/dead cell indicator TO-PRO™-3 iodide to determine the number of cells that were permeabilized. The results are percent averages plus the standard error mean (SEM, n=4). Percent values indicate the average relative abundance of events within each gate normalized to the total number of events analyzed, excluding artifacts, aggregates and debris. A one-way analysis of variance (ANOVA) followed by Sidak's multiple comparison post-hoc test was used to determine statistical significance between each treatment and the untreated control or tobramycin treatment alone with the combination treatment. *, p<0.05. NS, not significant.

**FIG 2. Mutants from experimental evolution are resistant to tobramycin and the combination.** 24-hr old biofilms were treated with triclosan (100 μM) and tobramycin (500 μM) alone and in combination. The results represent the means plus SEM (n=6). A two-way ANOVA followed by Bonferroni's posttest was used to determine statistical significance between each treatment and the untreated control. *, p<0.05. NS, not significant.

**FIG 3. Triclosan reduces the activity of RND-type efflux pumps.** 24-hr biofilms were stained with ethidium bromide to measure accumulation. Biofilms were treated with CCCP (100 μM), triclosan (100 μM), tobramycin (500 μM) alone and in combination. Fluorescence was read at 0,2,4, and 6-hrs. Results represent the average arbitrary fluorescence units ±SEM (n=6).

**FIG 4. Triclosan results in increased cellular accumulation and extrusion of Texas Red conjugated tobramycin (TbTR).** 24-hr old biofilms grown in glass test tubes were treated for 30-mins with triclosan (100 µM), CCCP (100 µM) and TbTR (250 µg/mL). (A) Then cells were disrupted from the biofilms and lysed with 0.2% Triton-X 100® to measure intercellular accumulation of TbTR. (B) To measure extrusion, biofilms were first treated for 30-mins and washed three-times in DPBS. Biofilms then recovered in treatment free media for 30-mins and fluorescence of the media was measured. TbTR was measured by relative fluorescence units using excitation $595_{nm}$ and emission $615_{nm}$. Results represent the average arbitrary fluorescence units ±SEM (n=6). For panel A, a One-Way-ANOVA was performed comparing TbTR alone vs CCCP and TbTR and vs triclosan and TbTR. For panel B, an unpaired t-test was performed comparing TbTR versus triclosan and TbTR. *, $p < 0.05$. NS, not significant.

**FIG 5. Triclosan reduces the membrane potential surge induced by tobramycin at 2-hours.** 24-hr old biofilms were treated with triclosan (100 µM), or tobramycin (500 µM), alone and in combination for 2-hrs. Cells were stained with DiOC2(3) to measure the membrane potential. Dead or permeabilized cells were excluded from membrane potential analysis by parallel staining with the dead cell indicator TO-PRO™-3 iodide. The results are percent averages plus the SEM (n=4). Percent values indicate the average relative abundance of events within each gate normalized to the total number of events analyzed, excluding dead cells, artifacts, aggregates and debris. A one-way ANOVA followed by Sidak's multiple comparison post-hoc test was used to determine statistical significance between each treatment and the untreated control, and tobramycin treatment alone was compared with the combination. *, $p < 0.05$. NS, not significant.

**FIG 6. Triclosan and tobramycin hydrogels are more effective than tobramycin hydrogels in a murine wound model.** 24-hr old bioluminescent biofilms formed within wounds were treated with triclosan (100 µM), or tobramycin (400 µM), alone and in combination for 4-hrs in a hydrogel. Reduction in the number of cells within biofilms was quantified using IVIS. The results are fold reduction of three separate experiments ±SEM, no treatment n=6, triclosan n=6, tobramycin n=10, triclosan and tobramycin n=9. A one-way ANOVA followed by Bonferroni's multiple comparison post-hoc test was used to determine statistical significance between each treatment and the untreated control and tobramycin treatment alone was compared with tobramycin and triclosan (*, p<0.05).

**FIG 7. Triclosan sensitizes *P. aeruginosa* to tobramycin by acting as a protonophore, inhibiting efflux pump activity, and abolishing adaptive resistance.** (1) Within 2-hrs of exposure to tobramycin, adaptive resistance occurs, (2) which is due to the induction of RND-type efflux pumps and surge in membrane potential ($\Delta\psi$), resulting in reduced accumulation of tobramycin within the cytosol. (3) Triclosan shuttles protons across the inner membrane, collapsing the proton motive force ($\Delta p$) and depolarizing the $\Delta\psi$. Consequently, efflux pump activity is reduced and there is enhanced accumulation of tobramycin within the cytosol. Finally, tobramycin binds to the A-site of the ribosome, corrupting protein synthesis and causing membrane permeabilization. Overall, triclosan accelerates and increase the effectiveness of tobramycin by reducing the $\Delta\psi$ and efflux pump activity. Proton gradient, $\Delta pH$. Outer membrane protein (OMP), membrane fusion protein (MFP), inner membrane protein (IMP).

**FIG S1. AFN-1252 alone or in combination with tobramycin is not effective against mature biofilms.** 24-hr old biofilms grown on MBEC plates were treated for 6-hrs with AFN-1252 (400 µM), tobramycin (400 µM), alone and in combination in two-fold dilutions, and the number of viable cells within the biofilms were quantified by BacTiter-Glo™. The assay was performed twice in in duplicate. The results represent means ± the Standard Error Mean (SEM). A one-way analysis of variance (ANOVA) followed by Bonferroni's multiple comparison post-hoc test was used to determine statistical significance between tobramycin versus the combination (NS, not significant).

**FIG S2. Initial tobramycin treatment is required for the combination to be effective.** 24-hr old biofilms grown on MBEC plates were treated sequentially. First, biofilms were treated for 3-hours with tobramycin (500 µM) and then washed three times in DPBS for 3-mins each, before being treated with triclosan (100 µM) for 3-hours or vice versa. As a control, biofilms were also treated for 6-hrs with triclosan and tobramycin. The number of viable cells within the biofilms were quantified by BacTiter-Glo™. The assay was performed twice in in duplicate. The results represent means plus the SEM. A one-way ANOVA followed by Bonferroni's multiple comparison post-hoc test was used to determine statistical significance between each treatment and the untreated control. *, $p < 0.05$. NS, not significant.

**FIG S3. Methyl-triclosan alone or in combination with tobramycin is not effective against mature biofilms.** 24-hr old biofilms grown on MBEC plates were treated for 6-hrs with methyl-triclosan (100 µM), tobramycin (400 µM), alone and in combination in two-fold dilutions, and the number of viable cells within the biofilms were quantified by BacTiter-Glo™. The assay was performed twice in in duplicate. The results represent means ±SEM. A one-way ANOVA followed by Bonferroni's multiple comparison post-hoc test was used to determine statistical significance compared to tobramycin alone and the combination. NS, not significant.

**FIG S4. Methyl-triclosan alone and in combination with tobramycin does not reduced the surge in membrane potential induced by tobramycin at 2-hours.** 24-hr old biofilms were treated with methyl-triclosan (100 μM), or tobramycin (500 μM), alone and in combination for 2-hrs. Cells were stained with DiOC2(3) and TO-PRO™-3 iodide to determine the number of cells that maintained a membrane potential (A) or were permeabilized (B), respectively. Dead or permeabilized cells were excluded from membrane potential analysis and are shown in panel B. The experiment was performed two separate times in duplicate. The results are percent averages plus the SEM. Percent values indicate the average relative abundance of events within each gate normalized to the total number of events analyzed, excluding artifacts, aggregates and debris. A one-way ANOVA followed by Sidak's multiple comparison post-hoc test was used to determine statistical significance between each treatment and an untreated control, and tobramycin treatment alone was compared with the combination. *, $p < 0.05$, NS, not significant.

**FIG S5. Methyl-triclosan does not reduced RND-type efflux pump activity.** Ethidium bromide is a substrate of RND-type efflux pumps. 24-hr biofilms were stained with ethidium bromide to measure accumulation. Biofilms were treated with methyl-triclosan (100 µM) and tobramycin (500 µM) alone and in combination. Fluorescence was read at 0,2,4, and 6-hrs. The assay was performed three times in triplicate. Results represent the average arbitrary fluorescence units ±SEM.

**FIG S6. DPBS starved biofilms are not more sensitive to tobramycin alone.** 24-hour old biofilms were starved of nutrients by replacing media with DPBS for 5-days. Starved biofilms were treated with triclosan (100 µM), or tobramycin (500 µM), alone and in combination for 6-hrs. The number of viable cells within the biofilms was quantified using the BacTiter-Glo™ assay. The assay was performed once using three biological replicates. The results represent the means plus the SEM. A one-way ANOVA followed by Sidak's multiple comparison post hoc test was used to determine statistical significance between the combination treatment and the untreated control and between triclosan alone or tobramycin alone and the combination treatment as indicated by the bars. *, $P < 0.05$.

**FIG S7. Tobramycin is as effective as the combination after 24-hrs of treatment.** 24-hr old biofilms were treated for 24-hrs with triclosan (100 µM), tobramycin (500 µM), alone and in combination, and the number of viable cells within the biofilms were quantified by BacTiter-Glo™. The assay was performed at least three times in triplicate. The results represent means plus the SEM. A one-way ANOVA followed by Bonferroni's multiple comparison post-hoc test was used to determine statistical significance between untreated biofilms, tobramycin alone, and the combination treatment. In addition, using the same analysis, a comparison was made between tobramycin alone and the combination treatment. *, $P < 0.05$. NS, not significant. 4-hr treatments were previously published and are shown for comparison (28).

**Table S1.** Treatment regimens used for the selection of resistant mutants.

| Cycle | Tobramycin | Triclosan | Table Part A | | | Cycle | Tobramycin | Triclosan | Table Part B | | |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 0 | 0 | 0 | | | | 17 | 10 | 100 | Gradual & Moderate | | |
| 1 | 0.00474756 | 0.47475615 | Gradual Start | | | 18 | 12.77 | 110.59 | | | |
| 2 | 0.00678223 | 0.67822307 | | | | 19 | 16.30729 | 122.301481 | | | |
| 3 | 0.0096889 | 0.9688901 | | | | 20 | 20.8244093 | 135.253208 | | | |
| 4 | 0.01384129 | 1.38412872 | | | | 21 | 26.5927707 | 149.576523 | | | |
| 5 | 0.01977327 | 1.97732674 | | | | 22 | 33.9589682 | 165.416676 | | | |
| 6 | 0.02824752 | 2.82475249 | | | | 23 | 43.3656024 | 182.934302 | | | |
| 7 | 0.04035361 | 4.0353607 | | | | 24 | 55.3778743 | 202.307045 | | | |
| 8 | 0.05764801 | 5.764801 | | | | 25 | 70.7175454 | 223.731361 | | | |
| 9 | 0.0823543 | 8.23543 | Moderate Start | | | 26 | 90.3063055 | 247.424512 | | | |
| 10 | 0.117649 | 11.7649 | | | | 27 | 115.321152 | 273.626768 | | | |
| 11 | 0.16807 | 16.807 | | | | 28 | 147.265111 | 302.603843 | | | |
| 12 | 0.2401 | 24.01 | | | | 29 | 188.057547 | 334.64959 | | ↓ | ↓ |
| 13 | 0.343 | 34.3 | | | | 30 | 240.149488 | 370.088981 | End | | |
| 14 | 0.49 | 49 | | | | | | | | | |
| 15 | 0.7 | 70 | | | | | | | | | |
| 16 | 1* Tobramycin MIC | 100 | | ↓ | ↓ | | | | | | |

27 *P. aeruginosa* biofilms serially passaged in parallel were exposed to sudden, moderate, and gradual treatment regimens consisting of ever-increasing concentrations of tobramycin and triclosan spread out over longer and longer periods of time, as shown in the table above. Gradual and moderate treatment groups were started at different concentrations of tobramycin and triclosan. However, they were then treated in subsequent cycles with ever increasing concentrations of triclosan and tobramycin at the same rate. For example, the moderate treatment group reached 1 µM of tobramycin and 100 µM of triclosan in 8-cycles, whereas the gradual treatment group reached the same concentration in 16-cycles from their initial treatments. The sudden treatment series was treated with 500 µM of triclosan and tobramycin from cycle day 1 and was eradicated, thus, is not shown in the table. MIC, minimum inhibitory concentration.