On porcine reproductive and respiratory syndrome virus dissemination across pig production

systems in the United States

Running title: Evolutionary epidemiology of PRRSv

Manuel Jara[1], David A. Rasmussen[2,3], Cesar A. Corzo[4], Gustavo Machado[1*]

[1]Department of Population Health and Pathobiology, College of Veterinary Medicine, North Carolina

State University, Raleigh, NC, USA.

[2]Department of Entomology and Plant Pathology, North Carolina State University, Raleigh NC, USA

[3]Bioinformatics Research Center, North Carolina State University, Raleigh NC, USA

[4]Veterinary Population Medicine Department, College of Veterinary Medicine, University of Minnesota,

1365 Gortner Ave, St Paul, MN 55108, USA.

[*]**Corresponding author**: gmachad@ncsu.edu

Summary

Porcine reproductive and respiratory syndrome virus (PRRSv) remains widespread in the North American

pig population. Despite improvements in virus characterization, it is unclear whether PRRSv infections

are a product of viral circulation within a farm, within production systems (local) or across production

systems (external). Here we examined the dissemination dynamics of PRRSv and the processes

facilitating its spread within and among pig farms in three production systems. Overall, PRRSv genetic

diversity declined since 2018, while phylodynamic results support frequent transmission across-

production systems. We found that PRRSv dissemination occurred mostly through transmission between

farms of different production companies, which were predominant for several months, especially from

November until May when PRRSv tends to peak in the studied region. Within production systems,

dissemination occurred mainly through regular pig flow, (from sow to nursery and then to finisher farms);

nevertheless, an important flux of PRRSv dissemination from finisher to sow and nursery farms

highlighted the importance of downstream farms as sources of the virus. High density of farms and farms

near major roads were more likely to infect by PRRSv, whereas higher elevation and denser vegetation

acted as dissemination barriers. Although it is difficult to disentangle the exact route of PRRSv

transmission, results evidenced that dissemination among commercially unrelated pig production systems

was intense, reinforcing the importance of farm proximity on PRRSv spread. Thus, consideration of farm

location and their geographic characteristics may help to forecast dissemination. The understanding of

PRRSv transmission routes has the potential to inform targeted strategies for its prevention and control.

Further studies are needed to quantify the relative contribution of PRRSv transmission routes.

Keywords: molecular epidemiology, phylodynamics, virus dispersal, between-farm dissemination.

Introduction

Porcine reproductive and respiratory syndrome virus (PRRSv) remains among the most costly diseases in

North America (Neumann et al., 2005; Holtkamp et al., 2013; Pileri and Mateu, 2016). Despite the great

advances in reducing the incidence of PRRSv in some regions (Rathkjen and Dall, 2017) by enhanced

biosecurity (Silva et al., 2019) and immunization strategies (Corzo et al., 2010), every year approximately

20-30% of US breeding herds still become infected with PRRSv (MSHMP, 2020).

PRRS is caused by an enveloped positive single-stranded RNA virus in the family Arteriviridae,

order Nidovirales (Cavanagh et al., 1990). The genome of PRRSv is ~15 kb and consists of 9 open

reading frames (ORFs) that encode seven structural proteins and 14 non-structural (Dokland, 2010). The

mutation rate of PRRSv hinders the effectiveness of immune response against newly introduced variant

strains, especially in farms with recurrent outbreaks (Linhares et al., 2014; Sanhueza et al., 2019).

Likewise, recombination between North American wild-type virus strains and modified-live vaccines

represents an important mechanism in the historical evolution and spread of PRRSv (Yuan et al., 1999).

The phylodynamic patterns of PRRSv spread and dissemination among pig systems and farm

types have been previously described (Shi et al., 2010; Franzo et al., 2015; Alkhamis et al., 2016;

Alkhamis et al., 2017; Sun et al., 2019). While there was support for between-production system

dissemination (Alkhamis et al., 2016), the relevance of transmission in the maintenance of local

incidence, in comparison with within-production system circulation, remains unknown. Thus, the

question of whether PRRSv circulation is the product of local dissemination between farms that are part

of a common commercial network (production system) versus the result of external viral introductions

from nearby infected farms of distinct production systems still remains (Arruda et al., 2018).

Bayesian phylodynamic approaches have provided a breakthrough in tracking pathogen

dissemination dynamics (Baele et al., 2018; Faria et al., 2019; Hicks et al., 2020). Structured coalescent

models and other phylodynamic approaches allow the estimation of relative contribution of local

dissemination versus external introductions to the overall incidence of disease (Rasmussen et al., 2018;

Yang et al., 2019; Hicks et al., 2020). Novel methods have been developed to integrate spatial and

environmental covariates in phylodynamic analysis, to determine their influence on the spread of

epidemics (Dudas et al., 2017; Fountain-Jones et al., 2018; Rasmussen et al., 2018; Dellicour et al., 2019;

Yang et al., 2019). Thus far, such approaches have not been widely implemented to study the

dissemination of diseases in domestic animals, except for Foot-and-Mouth virus (Duchatel et al., 2019).s

In this study, we used phylodynamic methods to determine the viral dissemination rates among

pig production systems to assess whether different PRRSv variants are the product of local transmission

or external introductions and identified the factors that facilitate or impede PRRSv spread.

Material and methods

Data collection

In total, our dataset comprises 4,970 ORF-5 sequences collected from 2014 to 2019, obtained from three

commercially unrelated pig production systems (here coded as A, B, and C), in the US. Sequences were

generated through outbreak investigation and from regular surveillance activities performed by each

production system and shared with the Morrison Swine Health Monitoring Program (MSHMP). Each

sequence was attributed to a farm type (e.g., sow, nursery, finisher, and boar stud), geographic location,

pig production system (A, B, and C), and the date when the sequence was reported.

Preliminary phylogenetic analysis

Sequences were aligned using Mega X, available at www.megasoftware.net (Kumar et al., 2008). The

recombination detection program (RDP) v5.3 was used to search for recombination within our dataset

(Martin et al., 2015). The alignment was screened using five methods (BootScan, Chimaera, MaxChi,

RDP, and SiScan), evidencing 63 recombinant sequences, which were removed from the downstream

analysis (Table S1). To determine whether there was a sufficient temporal molecular evolutionary signal,

we used TempEst v1.5 (Rambaut et al., 2016). To calculate the *P*-values associated with the phylogenetic

signal analysis, we followed Murray et al. (2016) using 1,000 random permutations of the sequence

sampling dates (Navascués et al., 2010). The relationship found between phylogenetic tree root-to-tip

divergence and sampling dates (years) supported the molecular clock analysis for this study ($P < 0.05$).

Discrete and continuous phylogeographical analysis

Phylogenies were estimated by Bayesian inference through Markov chain Monte Carlo (MCMC),

implemented in BEAST v2.5.0 (Bouckaert et al., 2014). By using ModelFinder (Kalyaanamoorthy et al.,

2017) built into IQ-Tree version 1.6.1 (Nguyen et al., 2015), we compared different substitution models

to find the best fit for our genetic dataset. The marginal likelihood value supported the use of the general

time-reversible model (GTR) with gamma-distributed rate heterogeneity plus a proportion of invariable

sites (GTR+G+I) (Tavaré, 1986). To reconstruct PRRSv dissemination across systems and between farm

types (i.e., sow, nursery, finisher, boar stud), we used the discrete-trait phylogeographic models

implemented in BEASTv2.5.0 to reconstruct ancestral states (Bouckaert et al., 2014). To explore the most

important historical dissemination routes for PRRSv spread across farms, as well as among pig systems,

we used Bayesian stochastic search variable selection (BSSVS) (Lemey et al., 2009). Through BSSVS,

we identified the non-zero rates of change between each pair of discrete traits (production system and

farm types) based on its Bayes factor value (lower than 3). To perform this analysis, an asymmetric

migration rate matrix was assumed. To infer the intensity of directional transitions (forward and

backward) within the migration matrix, we used a Markov jump approach. To evaluate oversampling and

possible selection bias towards location and/or farm types with disproportionate amounts of samples (i.e.,

sow farms), we downsampled on each system to match the number of samples between the three

systems, maintaining the same proportion of farm type (system A -89%, B =-39.5%, and C=31.4 %).

To analyze the spatiotemporal spread of PRRSv, we performed a Bayesian continuous

phylogeographic analysis, where the nucleotide substitution model was similar to the discrete model

mentioned above. In addition, we used an uncorrelated relaxed molecular clock model with lognormal

distribution (Drummond et al., 2006). The coalescent tree prior used was Bayesian SkyGrid with

covariates (Gill et al., 2016). Phylogeographic inference of ancestral locations was performed under a

relaxed random clock model, considering a lognormal distribution. All analyses were run for 200 million

generations, sampling every 10,000th generation and removing 10% of the chain as burn-in. To visualize

the spatiotemporal spread of PRRSv, we used Spatial Phylogenetic Reconstruction of Evolutionary

Dynamics using Data-Driven Documents (D3) SPREAD3 software (Bielejec et al., 2016).

Spatiotemporal epidemiological statistics

To summarize PRRSv diffusion over time and space we analyzed data from each production system

separately and all the sequences as a whole, using the R package "seraphim" version 1.0 (Dellicour et al.,

2016). Spatiotemporal information about lineage locations was extracted from 100 input trees sampled at

regular intervals from the post-burn-in posterior distribution to account for phylogenetic uncertainty

(Dellicour, Rose and Pybus, 2016). Each phylogeny branch was summarized and represented as a distinct

vector defined by its starting geographic location, ending geographic location and dates (Dellicour, Rose

and Pybus, 2016; Dellicour et al., 2016). Each branch vector represents an independent viral lineage

dissemination event (Pybus et al., 2012; Laenen et al., 2016). Each vector was assigned to a diffusion

coefficient value (D) for subsequent visualization: as described by (Pybus et al., 2012), which accounts

for the time (in years) during which each specific vector (lineage) moved from its initial location until the

end location of each lineage (measured from the location of the tree root). These vectors were also used to

assess the invasion velocity of the sampled lineages. To determine and compare the PRRSv diffusivity

associated with the overall data and for each production system, we estimated the spatial diffusion

coefficient using D-weighted statistics (Trovão et al., 2015) since this index is less sensitive to extreme

values of short branches. These results were displayed via: i) dissemination epidemiological statistics that

estimated the mean branch dissemination velocity, and ii) kernel plots showing the lineage diffusion

coefficients of the mean and variation on the highest posterior density (HPD).

Between-production systems transmission: the contribution of within-system PRRSv circulation

(local) versus external introduction of virus from a commercially unrelated pig farm.

We estimated the contribution of local transmission (i.e., between farms within a production system)

versus external introductions (i.e., between farms across production systems) using a structured

coalescent model. Due to computational demand, we used PRRSv alignment from MEGA software

(Kumar et al., 2008) to produce a phylogenetic tree using PhyML version 3.0 (Guindon et al., 2010),

using a substitution rate = 0.00672 (estimated), and considering an HKY Hasegawa-Kishino-Yano + $\Gamma 4$

(Hasegawa et al., 1985; Yang, 1994), which was dated using Least Squares Dating (LSD) version 0.3 (To

et al., 2016). To infer the contribution of local versus external introductions the migration rates between

different pig production systems were estimated using the Marginal Approximation to the Structured

Coalescent (MASCOT) package (Müller et al., 2017, 2018) implemented in Beast2 (Bouckaert et al.,

2014). To infer the evolutionary rate from the time-calibrated tree, we used a strict clock model. Using

MASCOT, we estimated the effective population size in the local and external population and the

migration rates between them, which was allowed to be asymmetric. This analysis was replicated three

times, each time using one of the three pig systems as local and the other two as external. We also

calculated the posterior probability of each internal node in the tree being in each state (i.e., local or

external) using the forwards/backwards approach implemented in MASCOT (Müller et al., 2018).

All analyses were performed under 100 million MCMC runs, states were sampled every 10,000

and the first 10% of samples were discarded as burn-in using TreeAnnotator v. 2.3.0 (Rambaut and

Drummond, 2016). We reconstructed the number of migration events through time using the ancestral

locations reconstructed by MASCOT. A migration event was assumed to occur when the most probable

location of a lineage changed between its parent and child node, with the time of the migration event

assumed to be the date of the daughter node. Migration events (local and external) per month were plotted

using "baltic" (https://github.com/blab/baltic) in Python Software Foundation version 3.7, available at

http://www.python.org. Finally, we compared the differences in local vs external transmission between

155   PRRSv season and off-season through two-sample Wilcoxon test (Hogg et al., 2010), PRRSv season was

156   determined by the exponentially weighted moving average proposed by MSHMP (MSHMP, 2020).

Factors promoting or restricting PRRSv spread

To determine the effect of different predictors on the spread of PRRSv, we analyzed the results of the

continuous phylogeographic analysis using the R package "seraphim" version 1.0 (Dellicour et al., 2016).,

considering a continuous location approach (based on the geographic coordinate location of each pig

farm) and a random walk diffusion model. Spatiotemporal information was extracted from 100 trees

sampled at regular intervals from the posterior distribution, after burn-in, to account for phylogenetic

uncertainty. Each phylogenetic branch was considered a vector defined by its start and end location (e.g.

latitude and longitude), and its start and end dates. Statistical significance for the correlation between

phylogenetic and predictive factors was tested using 100 trees generated and expressed in the form of

Bayes factors (BF) (Dellicour et al., 2016). Based on the extracted spatio-temporal information contained

in each phylogenetic tree described above, we calculated the distances based on predictor variables

associated with every branch of each tree, which were used to determine the correlation between the

duration of each branch in the phylogeny and its related predictor variable distance, including the annual

temperature, annual precipitation, elevation, runoff (an index that quantity of water discharged in surface

streams), soil humidity (commonly used to quantify topographic control on hydrological processes)

(Sørensen et al., 2005), enhanced vegetation index (EVI), which can be used to quantify vegetation

greenness, as well as the pig density, and distance to the main roads (Fig. 1 and Table S3 for more

details). Briefly, the distance between each farm (geographic coordinates) to the nearest main road was

calculated based on the Euclidean distance. We calculated the statistic Q, which represents how much

variation in lineage movement is explained with spatial heterogeneity when each predictor variable is

considered (Dellicour, Rose and Pybus, 2016). Here, $Q = R^2_{env} - R^2_{null}$ where R^2_{env} is the coefficient of

determination obtained from the regression between branch duration against predictor (factors) distances,

while R^2_{null} represents the coefficient of determination for the regression between branch duration against

predictor distances, which were replaced by a null raster. Thus, the factors that were considered for

further Bayes factor (BF) analysis, were the ones that obtained positive Q values in at least 90% of the

tested trees (Jacquot et al., 2017). We approximated a BF value for all predictors tested that passed the

above-mentioned Q statistic. Predictive factors were treated as a conductance (variables that promote the

spread of the disease) or a resistance factor (variables that impede its spread). For the interpretation of BF,

values between 3-20 were considered "positively" supported, values between 20-150 were considered

"strong", and values >150 were considered " overwhelming support" (Kass and Raftery, 1995).

Results

A total of 63 recombinant PRRSv sequences were identified and removed from the database. Most

recombinants (52%) belonged to system B with the highest number of recombinants (82%) belonging to

sow farms (see supplementary table S1). Thus, 4,753 sequences remained for further discrete and

continuous phylogeographic analyses. Most sequences originated from system A = 3,748 (79%), while B

had 626 (13%), and C had 379 (8%). Overall, 2,958 sequences (62.2%) belonged to sow farms, 909

(19.1%) to finishers, 873 (18.4%) to nurseries, and 13 (0.3%) to boar studs (see supplementary Table S1).

Discrete phylogeography analysis

The results of PRRSv ancestral reconstruction showed that the most likely center of origin was production

system A with a root state posterior probability (RSPP) = 0.84, coming from a sow farm (RSPP = 0.75).

From there, it spread heterogeneously across the other systems, unrestricted to farm types (Fig. 2A). The

reconstructed phylogenetic relationship did not show evidence of any cluster related to production

systems. However, we observed a clear dominance of PRRSv originating from system A over the other

systems (Fig. 2A). Our sensitivity analysis on the downsample supported these results (Supplementary

Table S4). The relative genetic diversity expressed as the effective population size (e.g. number of

individuals contributing to a new PRRSv infection) over time showed a plateau until 2012. After that

time, we captured a steep increment that lasted until approximately 2016, when relative genetic diversity

decreased through June 2019 (Fig. 2B), the last month of data available for our study. The temporal

pattern shown by the sampled PRRSv sequences evidenced the typical annual trend of seasonal outbreaks

(Fig. 2B) (MSHMP, 2020). Phylogeographic results of system A showed that its most likely center of

origin was a sow farm (RSPP = 0.47), followed by an almost equal probability for origins in finisher farm

(RSPP = 0.27) and nursery farm (RSPP = 0.26), the boar stud farms were negligible (RSPP < 0.01) (Fig.

3A). In recent years, there was an evident increase in the proportion of PRRSv positive finisher farms,

reaching a similar level of representation than sow farms for some clusters (Fig. 3A). SkyGrid plot

revealed a relatively constant trend of relative genetic diversity until ~2011, showing a linear increase in

population size, which reached its highest levels ~2016 (Fig. 3B). We observed a similar trend when we

analyzed all systems together (Fig. 3B), 2015 and 2018 were the years with more samples sent to be

sequenced (Fig. 3B).

For production system B, a sow farm (RSPP = 0.4) was also the most likely origin; however, a

high probability was also linked to a nursery farm (RSPP = 0.33) (Fig. 4A). The temporal pattern shown

by the effective population size was static until 2012, after which it increased rapidly until 2015-2016,

when it started a linear decrease until June 2019, similar to the seasonal pattern shown by the number of

sequences (Fig. 4B). As observed in systems A and B, the most likely center of origin for system C

occurred inside a sow farm (RSPP = 0.52), followed by a high probability for originating in a finisher

farm (RSPP = 0.4). In contrast to the other systems, we observed a low probability of origin related to a

nursery farm (RSPP > 0.1) (Fig. 5A). The temporal trend in the number of sampled sequences showed the

mentioned annual seasonal pattern, with peak occurrence ~2015. SkyGrid plot evidenced a more stable

pattern, with higher levels of variation in effective population size during the period 2013-2014, then

decreasing until 2018, when it showed a small peak and then decreased through June 2019 (Fig. 5B).

Phylogeographic diffusion analysis and spread statistics

The reconstructed spatiotemporal diffusion of PRRSv was characterized by continuous expansion in the

geographic space, accompanied by diversification events. The comparison between each system and all

samples together showed two noticeable patterns. The first one was characterized by a radial expansion

over time, found when all samples were analyzed together and within samples of system A. In contrast,

systems B and C evidenced a directional trend, since for both systems the spread started in the southwest

region. In the case of system B, PRRSv spread to the northeast, while for system C, the spread was

mainly to the north, though spread did occur in all directions (see Supplementary Figs. S1-3).

The spatiotemporal analysis showed that when all sequences were analyzed together, the

estimated median value for the mean branch velocity was 39 km/year (95% HPD = 31.2-61.1). This

median value was lower than that shown by system A (49.2 km/year, 95% HPD= 42.9-60.4), and higher

than the values observed in systems B and C (32.7 km/year, 95% HPD= 29.1-43.3), and (26.4 km/year,

95% HPD= 20.1-41.0), respectively. Estimated diffusion coefficients were asymmetric for each system

and when all samples were analyzed together. System A showed the highest diffusivity (243.8 km^2/year);

however, its diffusion variation among branches was the lowest registered. The overall results (all

sequences) evidenced the highest among-branch variation, while its diffusion coefficient (150 km^2/year)

was lower than system A and substantially higher than the other systems. Systems B and C showed a

similar pattern in terms of diffusion coefficient variation among branches; however, B was higher than C

(97 km^2/year) and (51 km^2/year), respectively (Supplementary Fig. S4).

PRRSv dissemination among pig systems and farm types

Bayesian stochastic search variable selection (BSSVS) identified that the most significant disseminations

were highly related to system A, evidencing strong support by a Bayes Factor score > 20. The statistical

support or evidence for system A sourcing most PRRSv strains to system B, BF = 102.1 (average relative

migration rate "ARMR"= 1.79), followed by dissemination from A to C, BF = 47.8 (ARMR = 0.46); of

less importance was from B to A, BF= 27.4 (ARMR = 0.35). All the other disseminations between

production systems were also positive, but to a lesser degree than the ones mentioned above (Table 1). In

addition, a sensitivity analysis aimed to evaluate the bias related to oversampling showed that while

numerically it was a difference the interpretation remains the same, but we noticed that in the subsample

there was even more support for the dissemination from company B to C and company C to A (see

Supplementary Table S5 for more details). Our analysis reveals that when comparing PRRSv

dissemination between types of farms, the most frequent routes were linked with sow farms, especially

when the direction of spread was from sow to nursery (BF > 42). The highest support for this route

occurred when we analyzed all systems combined (BF = 109). The second most well-supported

dissemination occurred from sow to finisher farms (BF > 17.9), and as in the previous case, was best

supported when we analyzed all systems together. (BF = 80.5). Likewise, the dissemination between

nursery and finisher farms (in both directions) also showed positive support (BF > 6). However, all routes

of dissemination that involved boar stud farms were rare events (BF < 3) (Table 2). Furthermore, the

sensitivity analysis showed similar trends, where the results of the equal sample sizes these were less

sharp than the one using the original dataset. However, the transmissions that involved BoarStud were all

negligible (see Supplementary Table S6 for more details).

Within-production systems circulation (local) versus introduction of PRRSv from commercially

unrelated farms (external)

We tracked the origin of the migration events leading to the current PRRSv lineages circulating among

the three pig systems, to determine whether detected PRRSv strains originated from within the same

system (local) or from farms of non-commercially related pig systems (external introductions). For this

analysis, we considered each coalescent event in the tree as a proxy for the timing of a transmission event

and then classified each event as either a local transmission event if the parent and child lineages were

both reconstructed to be in the local population or an external introduction if the parent lineage resided

outside the local population. Overall, results showed a clear dominance of external transmission events

that were observed in all pig companies; this tendency was constant during the whole period of study

(Fig. 6). When looking at the difference between local transmission against external introductions, system

C showed the largest number of external introductions with 73.8% of all transmission events inferred to

be external introductions, while systems A and B exhibited 56.6% and 58.7% respectively (Table 3). In

addition, our results show that overall, the external introduction events tend to increase over time (Table

3). We also observed that coinciding with the month of higher incidence of PRRSv (MSHMP, 2020) (Fig.

6 and Supplementary Table S7). Two-sample Wilcoxon test indicated that the difference between the

within farm transmission and external introductions were statistically significantly higher during PRRSv

season (mean= 69.3) than the rest of the year (mean= 60.1) (Z= -9.9, $p < 0.001$).

284   **Factors promoting or restricting PRRSv dissemination direction**

285   To quantify the impact of each factor on PRRSv distribution, we calculated the association between

286   lineage duration and each covariate scaled distance through the Q statistic method. We found strong

287   support (Q value was positive) for pig density and distance to main roads treated as conductance factors,

288   while elevation and vegetation presented strong support as resistance factors (see Supplementary Table

289   S8 and Supplementary Figs. S5 and S6 for more details). These results also received statistical support

290   provided by Bayes factors (BF) since we observed that farms located in areas with a high density of pig

291   farms were more exposed to the introduction of PRRSv (BF > 12). Likewise, the distance between the

292   farm and the nearest major road also showed a positive association with PRRSv diffusion (BF > 7). In

293   contrast, the factors identified as negative drivers, variables that act as barriers for the spread of PRRSv,

294   were higher elevation and denser vegetation (BF > 3) (Supplementary Table S8).

295   **Discussion**

296   This analysis detailed the evolutionary epidemiology of PRRSv among commercially related and

297   unrelated farms through continuous and discrete phylogeographic analyses. Overall, a decline in PRRSv

298   genetic diversity was observed over time. Such findings could have different explanations, one being a

299   rapid and efficient disseminating strain that was introduced into sow farms which consequently was

300   transferred to growing pig sites through regular pig movement. Another potential explanation could be

301   related to the efficacy of current interventions that local production systems have implemented, which

302   reduced the volume and frequency of newly emerging strains, suggesting a more optimistic scenario than

303   proposed in a similar study done in another US region (Alkhamis et al., 2017). Given the economic

304   impact and herd health relevance of new PRRSv strain introductions, it is imperative to evaluate how

305   often PRRSv spreads among pig production systems, determine the role of each farm type, and identify

306   factors that may facilitate or keep PRRSv from spreading. We demonstrated that PRRSv circulation has

307   been mostly originated by transmission from outside of each pig production system, and this tendency is

308   more predominant during the PRRSv season (from November until May) (MSHMP, 2020).

Our findings suggest that virus transmission between pig systems is common. It is not well

understood which transmission mechanism plays a role in the dissemination across systems. Airborne

spread has been suggested as a potential route (Dee et al., 2009; Otake et al., 2010). However, Arruda et

al., (2018, 2019) did not reach that same conclusion. Pig farm density as expected arises as a risk factor in

several studies and in our study, our phylodynamic findings agree with a regional spread as an important

factor for PRRSv dissemination among pig farms. PRRSv spread events appear to be relatively common

in farms located in highly dense pig areas and near main roads, while farms at higher elevations and

surrounded by denser vegetation were significantly less likely to report new cases of PRRSv (Arruda et

al., 2017). It is important to mention that the source of our sequence data represented a limitation for our

phylodynamic analysis since it was a convenience sample that can introduce bias into our analysis as the

number of sequences does not directly represent all PRRSv cases circulating in this study region.

PRRSv spatiotemporal dynamics

The spatiotemporal dynamic pattern exhibited by PRRSv is the result of its transmission dynamics and

how circulating strains have been affected by viral selection (Pybus and Rambaut, 2009; Volz et al.,

2013). This can be observed in the shape of PRRSv phylodynamic patterns over the course of this study,

which shows a clear directional selection, which can result in the loss of genetic variation (Volz et al.,

2013). Based on MSHMP data (MSHMP, 2020), vaccine usage increased between 2012 and 2014, which

could have impacted PRRSv genetic diversity for system A after 2012. But the increment in the number

of farms using vaccines during 2014 did not show a noticeable effect on PRRSv genetic variation.

However, the increment in the application of MLV's in mid-2015 showed an undeniable positive effect

on decreasing PRRSv genetic diversity (as seen in Figs. 2-5), similarly to the observations made by (Sun

et al., 2012; Jeong et al., 2018). However, to increase the robustness of future studies it is necessary to

increase the farm coverage as well as the timeframe, not only related to the PRRSv genetic information

but also to the vaccination data.

The estimated speed in which PRRSv was spreading was 39 km/year which was slightly higher

than previous observations in a study that used data from one production system in another U.S. region

(34 km/year) (Alkhamis et al., 2017). This difference may be related to pig farm densities, being our

study region more populated with an average of 3 farms per 5 km^2, and a more loyal representation of

PRRSv dynamics since we compiled samples of more than 92% of all farms in our study area. When

compared to the diffusion of other viral diseases, the spread of PRRSv was similar to rabies spread in

North America (~37 km/year) (Dellicour, Rose and Pybus, 2016). Consequently, there are important

aspects that have to be considered while analyzing the speed at which PRRSv spreads, for example, those

related with the overall pig health (i.e., absence of coinfections) and factors related with on-farm

biosecurity (i.e., presence of a line of separation, cleaning, and disinfection station) (Rappole et al., 2006;

Altizer et al., 2011; Lycett et al., 2019). Thus, further studies are needed to evaluate how biosecurity

practices and overall pig health may contribute in slowing or accelerating PRRSv spread, results of these

assessments can be used to predict the most effective biosecurity measures and optimize control strategies

that go beyond PRRSv (Silva et al., 2019), not to mention current diseases that pose large-scale

biosecurity threats to the U.S swine health such as African Swine Fever.

PRRSv spread among and within farms of commercially unrelated production systems

Our analysis found relevant similarities in the spatiotemporal patterns of PRRSv spread among all farm

types of all production systems (Table 1, 2), which was similar to previously reported findings that used

data from one production system in the U.S. (Alkhamis et al., 2016). At farm type levels, sow farms were

by far the biggest spreader (Plain and Laurence, 2003) which was expected as they wean piglets on a

weekly basis and these can be transported several miles away to an off-site growing pig facility.

Phylogenetic results also showed that PRRSv introductions into sow and nursery farms likely originated

from finisher farms, which could be partially related to gilt development units sending replacement gilts

to sow farms (Dee, 1995; Perez et al., 2015). Despite the substantial efforts to improve farms' biosecurity

and immunization through the implementation of strategies such as, sow herd air filtration, gilt

acclimation, virus elimination strategies (i.e., load-close-expose and rollover) which successfully reduced

the number of PRRSv outbreaks (Corzo et al., 2010; Alonso et al., 2012; Velasova et al., 2012; Alonso et

al., 2013; Blanchette, 2015; Silva et al., 2019), we continue to observe significant PRRSv dissemination

by indirect routes (Corzo et al., 2010; Otake et al., 2010). Therefore, it is expected that changes in

immunization protocols, such as varying vaccine brands at farms of one specific system may have a

serious impact on PRRSv spread.

Within-production systems circulation (local) versus introduction of PRRSv from commercially

unrelated farms (external)

Between-farm animal movement has historically been associated with disease spread (Augusta et al.,

2019; Moon et al., 2019; Machado et al., 2020). Here, we used structured coalescent models to infer

migration rates among commercially unrelated pig farms, calculating PRRSv dissemination not related to

between-farm pig movement. Overall, PRRSv dissemination in this region has been mostly characterized

by transmission between non-commercially related farms (external introductions) rather than within a

production system (local). In addition, we observed that during PRRSv season (November-May), the

external transmissions tended to be higher than during the rest of the year. While our analyses do not

necessarily reveal where spread events occurred or how viral lineages first entered a population

(Rasmussen et al., 2018), they reinforce the importance of PRRSv collateral spread and open the question

of how its dissemination is happening. The external introduction of viruses can occur through different

routes, as contaminated vehicles that service multiple production systems (i.e., general maintenance,

vaccination, cleaning or loading) (Otake et al., 2003; Dee et al., 2004), shared equipment (Pitkin et al.,

2009), non-pig animal vectors (Otake et al., 2002; Schurrer et al., 2005), transport of semen (Christopher-

Hennings et al., 1995) and airborne transmission (Lager and Mengeling, 2000; Trincado et al., 2004; Cho

et al., 2007; Dee et al., 2009). Knowledge gaps regarding local dissemination of PRRSv still exist, hence,

studies that consider phylodynamics coupled with epidemiological models such as susceptible–infected–

recovered (SIR) to assess the relative contribution of between-farm PRRSv transmission routes are

needed. In addition, modeling allowed not only to answer "how it can spread" questions but also "what

for" issues, i.e. determining what effort is required for reducing transmission (Beaunée et al., 2017).

Factors promoting or restricting PRRSv dissemination direction

In our PRRSv dissemination models, directions of lineages revealed the potential importance of two

factors that limited the spread of PRRSv. Vegetation density and high elevation, which are similar to

previous observations made by Mahesh et al., (2015) and Arruda et al., (2017). These authors also

highlighted the important role that topography may play in mitigating the airborne spread of PRRSv. In

the same way, a study on foot-and-mouth disease, which also used continuous phylogeographic analysis,

identified that the presence of pure cropland or mixed land use both had a negative impact on virus

diffusion (Duchatel et al., 2019). Numerous studies have shown multiple positive impacts of having

vegetation barriers surrounding farms (i.e., trees, shrubs), from reducing the emission of odor and

pollution, as well as the airborne spread of different pathogens (Malone, 2004; Patterson et al., 2008; Van

Ryswyk et al., 2019), thus it should be recognized as a possible biosecurity strategy. Similarly, high

elevated areas have proven to act as a natural topographic barrier for the local spread of pathogens

(Jacquot et al., 2017), where the slope of the terrain was found to have an association with a lower PRRSv

incidence (Arruda et al., 2017). The identification of the main drivers associated with PRRSv

dissemination could help in deciding about new farm locations, especially for systems located in highly

dense pig areas (Yang et al., 2019). Farms located in areas with higher farm density were more likely to

sustain new PRRSv introductions, as has been previously described by Firkins and Weigel, (2004),

Arruda et al., (2017) and Alkhamis et al., (2017, 2018). In addition, our results identified distance to the

main road as associated with increased chances for PRRSv circulation. This has led concern by the swine

industry (Reicks, 2019), since vehicle movement has demonstrated to have a significant impact on

increasing the transmission of several other diseases, for example, African Swine Fever (ASF) (Mur et al.,

2012), Classical Swine Fever (CSF) (Bronsvoort et al., 2008), and for Foot and Mouth Disease (Muroga

et al., 2013).

Here, we showed that continuous phylogeographic reconstructions represent a useful tool to

describe and analyze the dissemination of PRRSv spread. Although the outcomes of phylodynamic

analyses can be exploited to investigate the impact of relevant factors on lineage dissemination diffusion,

it is important to note that such analyses depend on inferred viral lineage movement and thus, on the

spatial distribution of the sampled sequences (Dellicour et al., 2019). Finally, our samples represent the

circulating PRRS viral population, as data comes from 92% of pig farms in one highly dense pig region.

Conclusion

Our study revealed significant asymmetries in the phylodynamic patterns among farms of three pig

production systems. As a result of interventions that could be related to immunization and on-farm

biosecurity, PRRSv genetic diversity has consistently declined in this region, especially in the last two

years, which could explain why 2019 has had one of the lowest counts of PRRSv cases during its typical

high season in the US (Sanhueza et al., 2019). Our results evidenced that PRRSv transmission across each

system is most likely to originate from outside production systems. These results reiterate the importance

of PRRSv farm proximity as a relevant route for virus dissemination. We highlight characteristics of

locations to which PRRSv is more easily dispersed: highly dense pig population and proximity to main

roads. On the protective side, higher elevation and more vegetation would be preferred. Thus,

reinforcement of biosecurity may be the best tool. However, it raises the question of how much and which

biosecurity practices have relevant impact on PRRSv spread. Answering this question requires further

studies capable of quantifying how biosecurity practices impede or reduce PRRSv dissemination.

Acknowledgements

We acknowledge the Department of Population Health and Pathobiology: North Carolina State University

provided startup funds for Dr. Machado and CVM for the intramural grant which supported Dr. Jara. This

work was also supported by Critical Agricultural Research and Extension 2019-68008-29910 from the

USDA National Institute of Food and Agriculture. The Morrison Swine Health Monitoring Project is a

Swine Health Information Center funded project. Authors would like to acknowledge participating

systems and veterinarians.

Authors' contributions

MJ and GM conceived the paper ideas. MJ, DR, and GM participated in the design of the study. CC

coordinated the PRRSv data collection. MJ conducted data processing and cleaning. MJ performed the

phylodynamic analysis. MJ, DR, CAC, and GM wrote and edited the manuscript. All authors discussed

the results and critically reviewed the manuscript.

Conflict of interest

All authors confirm that there are no conflicts of interest to declare

Ethical statement

The authors confirm the ethical policies of the journal, as noted on the journal's author guidelines page.

Since this work did not involve animal sampling nor questionnaire data collection by the researchers there

was no need for ethics permits.

Data Availability Statement

The data that support the findings of this study are not publicly available and are protected by confidential

agreements, therefore, are not available.

References

Alkhamis, M., A. Perez, M. Murtaugh, X. Wang, and R. Morrison, 2016: Applications of Bayesian

phylodynamic methods in a recent U.S. porcine reproductive and respiratory syndrome virus outbreak.

Front. Microbiol. **7**.

Alkhamis, M.A., A.G. Arruda, R.B. Morrison, and A.M. Perez, 2017: Novel approaches for Spatial and

Molecular Surveillance of Porcine Reproductive and Respiratory Syndrome Virus (PRRSv) in the United

States. *Sci. Rep.* **7**.

Alkhamis, M.A., A.G. Arruda, C. Vilalta, R.B. Morrison, and A.M. Perez, 2018: Surveillance of porcine

reproductive and respiratory syndrome virus in the United States using risk mapping and species

distribution modeling. *Prev. Vet. Med.* **150**, 135–142.

Alonso, C., M.P. Murtaugh, S.A. Dee, and P.R. Davies, 2013: Epidemiological study of air filtration

systems for preventing PRRSV infection in large sow herds. *Prev. Vet. Med.* **112**, 109–117.

Alonso, C., S. Otake, P. Davies, and S. Dee, 2012: An evaluation of interventions for reducing the risk of

PRRSV introduction to filtered farms via retrograde air movement through idle fans. *Vet. Microbiol.* **157**,

304–310.

Altizer, S., R. Bartel, and B.A. Han, 2011: Animal migration and infectious disease risk. *Science (80).* **331**,

296–302.

Arruda, A.G., J. Sanhueza, C. Corzo, and C. Vilalta, 2018: Assessment of area spread of porcine

reproductive and respiratory syndrome (PRRS) virus in three clusters of swine farms. *Transbound. Emerg.*

Dis. **65**, 1282–1289.

Arruda, A.G., S. Tousignant, J. Sanhueza, C. Vilalta, Z. Poljak, M. Torremorell, C. Alonso, and C.A. Corzo,

2019: Aerosol Detection and Transmission of Porcine Reproductive and Respiratory Syndrome Virus

(PRRSV): What Is the Evidence, and What Are the Knowledge Gaps? *Viruses* **11**, 712.

Arruda, A.G., C. Vilalta, A. Perez, and R. Morrison, 2017: Land altitude, slope, and coverage as risk factors

for Porcine Reproductive and Respiratory Syndrome (PRRS) outbreaks in the United States. *PLoS One* **12**.

Augusta, C., G.W. Taylor, and R. Deardon, 2019: Dynamic contact networks of swine movement in

Manitoba, Canada: Characterization and implications for infectious disease spread. *Transbound. Emerg.*

Dis. **66**, 1910–1919.

Baele, G., S. Dellicour, M.A. Suchard, P. Lemey, and B. Vrancken, 2018: Recent advances in

computational phylodynamics. *Curr. Opin. Virol.* **31**, 24–32.

Beaunée, G., E. Vergu, A. Joly, and P. Ezanno, 2017: Controlling bovine paratuberculosis at a regional

scale: Towards a decision modelling tool. *J. Theor. Biol.* **435**, 157–183.

Bielejec, F., G. Baele, B. Vrancken, M. Suchard, A. Rambaut, and P. Lemey, 2016: SpreaD3: interactive

visualization of spatiotemporal history and trait evolutionary processes. *Mol. Biol. Evol.* **33**, 2167–2169.

Blanchette, A., 2015: Herding species: Biosecurity, posthuman labor, and the american industrial pig. *Cult.*

Anthropol. **30**, 640–669.

Bouckaert, R., J. Heled, D. Kühnert, T. Vaughan, C.H. Wu, D. Xie, M.A. Suchard, A. Rambaut, and A.J.

Drummond, 2014: BEAST 2: A Software Platform for Bayesian Evolutionary Analysis. (Andreas Prlic,

Ed.) *PLoS Comput. Biol.* **10**, e1003537.

Bronsvoort, B.M.D.C., S. Parida, I. Handel, S. McFarland, L. Fleming, P. Hamblin, and R. Kock, 2008:

Serological survey for foot-and-mouth disease virus in wildlife in eastern Africa and estimation of test

parameters of a nonstructural protein enzyme-linked immunosorbent assay for buffalo. *Clin. Vaccine*

Immunol. **15**, 1003–11.

Cavanagh, D., D.A. Brian, L. Enjuanes, K. V. Holmes, M.M.C. Lai, H. Laude, S.G. Siddell, W. Spaan, F.

Taguchi, and P.J. Talbot, 1990: Recommendations of the coronavirus study group for the nomenclature of

the structural proteins, mRNAs, and genes of coronaviruses. *Virology* **176**, 306–307.

Cho, J.G., J. Deen, and S.A. Dee, 2007: Influence of isolate pathogenicity on the aerosol transmission of

Porcine reproductive and respiratory syndrome virus. *Can. J. Vet. Res.* **71**, 23–27.

Christopher-Hennings, J., 2001: Monitoring for porcine reproductive and respiratory syndrome virus

(PRRSV) in the boar stud. *J. Swine Heal. Prod.* **9**, 186–188.

Christopher-Hennings, J., E.A. Nelson, R.J. Hines, J.K. Nelson, S.L. Swenson, J.J. Zimmerman, C.L.

Chase, M.J. Yaeger, and D.A. Benfield, 1995: Persistence of porcine reproductive and respiratory syndrome

virus in serum and semen of adult boars. *J. Vet. Diagn. Invest.* **7**, 456–64.

Corzo, C.A., E. Mondaca, S. Wayne, M. Torremorell, S. Dee, P. Davies, and R.B. Morrison, 2010: Control

and elimination of porcine reproductive and respiratory syndrome virus. *Virus Res.* **154**, 185–192.

Dee, S.A., 1995: Controlling the spread of PRRS virus in the breeding herd through management of the gilt

pool. *Swine Heal. Prod.* **3**, 64–69.

Dee, S.A., J. Deen, and C. Pijoan, 2004: Evaluation of 4 intervention strategies to prevent the mechanical

transmission of porcine reproductive and respiratory syndrome virus. *Can. J. Vet. Res.* **68**, 19–26.

Dee, S.A., S. Otake, S. Oliveira, and J. Deen, 2009: Evidence of long distance airborne transport of porcine

reproductive and respiratory syndrome virus and Mycoplasma hyopneumoniae. *Vet. Res.* **40**, 39.

Dellicour, S., P. Lemey, J. Artois, T.T. Lam, A. Fusaro, I. Monne, G. Cattoli, D. Kuznetsov, I. Xenarios,

G. Dauphin, W. Kalpravidh, S. Von Dobschuetz, F. Claes, S.H. Newman, M.A. Suchard, G. Baele, and M.

Gilbert, 2019: Incorporating heterogeneous sampling probabilities in continuous phylogeographic inference

- Application to H5N1 spread in the Mekong region. *Bioinformatics* 1–7.

Dellicour, S., R. Rose, N.R. Faria, P. Lemey, and O.G. Pybus, 2016: SERAPHIM: Studying environmental

rasters and phylogenetically informed movements. *Bioinformatics* **32**, 3204–3206.

Dellicour, S., R. Rose, and O.G. Pybus, 2016: Explaining the geographic spread of emerging epidemics: a

framework for comparing viral phylogenies and environmental landscape data. *BMC Bioinformatics* **17**,

82.

Dellicour, S., C. Troupin, F. Jahanbakhsh, A. Salama, S. Massoudi, M.K. Moghaddam, G. Baele, P. Lemey,

A. Gholami, and H. Bourhy, 2019: Using phylogeographic approaches to analyse the dispersal history,

velocity and direction of viral lineages - Application to rabies virus spread in Iran. *Mol. Ecol.* **28**, 4335–

4350.

Dokland, T., 2010: The structural biology of PRRSV. *Virus Res.* **154**, 86–97.

Drummond, A.J., S.Y.W. Ho, M.J. Phillips, and A. Rambaut, 2006: Relaxed phylogenetics and dating with

confidence. (David Penny, Ed.)*PLoS Biol.* **4**, e88.

Duchatel, F., M. Bronsvoort, and S. Lycett, 2019: Phylogeographic analysis and identification of factors

impacting the diffusion of Foot-and-Mouth disease virus in Africa. *Front. Ecol. Evol.* **7**, 371.

Dudas, G., L.M. Carvalho, T. Bedford, A.J. Tatem, G. Baele, N.R. Faria, and F. Bielejec, 2017: Virus

genomes reveal factors that spread and sustained the Ebola epidemic. *Nature* **544**, 309–315.

Faria, N.R., N. Vidal, J. Lourenco, J. Raghwani, K.C.E. Sigaloff, A.J. Tatem, D.A.M. Van De Vijver, A.C.

Pineda-Peña, R. Rose, C.L. Wallis, S. Ahuka-Mundeke, J.J. Muyembe-Tamfum, J. Muwonga, M.A.

Suchard, T.F.R. De Wit, R.L. Hamers, N. Ndembi, G. Baele, M. Peeters, O.G. Pybus, P. Lemey, and S.

Dellicour, 2019: Distinct rates and patterns of spread of the major HIV-1 subtypes in Central and East

Africa. *PLoS Pathog.* **15**, e1007976–e1007976.

Firkins, L.D., and R.M. Weigel, 2004: A retrospective study of risk factors for porcine reproductive and

respiratory syndrome virus infection and clinical disease in swine herds in Illinois during the early years of

the pandemic. *J. Swine Heal. Prod.* **12**, 23–28.

Fountain-Jones, N.M., W.D. Pearse, L.E. Escobar, A. Alba-Casals, S. Carver, T.J. Davies, S. Kraberger,

M. Papeş, K. Vandegrift, K. Worsley-Tonks, and M.E. Craft, 2018: Towards an eco-phylogenetic

framework for infectious disease ecology. *Biol. Rev.* **93**, 950–970.

Franzo, G., G. Dotto, M. Cecchinato, D. Pasotto, M. Martini, and M. Drigo, 2015: Phylodynamic analysis

of porcine reproductive and respiratory syndrome virus (PRRSV) in Italy: Action of selective pressures and

interactions between different clades. *Infect. Genet. Evol.* **31**, 149–157.

Gill, M.S., P. Lemey, S.N. Bennett, R. Biek, and M.A. Suchard, 2016: Understanding Past Population

Dynamics: Bayesian Coalescent-Based Modeling with Covariates. *Syst. Biol.* **65**, 1041–1056.

Guindon, S., J.F. Dufayard, V. Lefort, M. Anisimova, W. Hordijk, and O. Gascuel, 2010: New algorithms

and methods to estimate maximum-likelihood phylogenies: Assessing the performance of PhyML 3.0. *Syst.*

Biol. **59**, 307–321.

Hasegawa, M., H. Kishino, and T. Yano, 1985: Dating of the human-ape splitting by a molecular clock of

mitochondrial DNA. *J. Mol. Evol.* **22**, 160–174.

Hicks, J.T., D.H. Lee, V.R. Duvuuri, M.K. Torchetti, D.E. Swayne, and J. Bahl, 2020: Agricultural and

geographic factors shaped the North American 2015 highly pathogenic avian influenza H5N2 outbreak.

PLoS Pathog. **16**, e1007857.

Hogg, R. V, E.A. Tanis, D.L. Zimmerman, B. Columbus, I. New, Y. San, F. Upper, S. River, A. Cape, T.

Dubai, L. Madrid, M. Munich, P. Montreal, T. Delhi, M.C. São, P. Sydney, H. Kong, S. Singapore, and T.

Tokyo, 2010: Probability and Statistical Inference. Pearson/Prentice Hall.

Holtkamp, D.J., J.B. Kliebenstein, E. Neumann, J.J. Zimmerman, H. Rotto, T.K. Yoder, C. Wang, P. Yeske,

C.L. Mowrer, C.A. Haley, E.J. Neumann, H.F. Rotto, and P.E. Yeske, 2013: Assessment of the economic

impact of porcine reproductive and respiratory syndrome virus on United States pork producers Assessment

of the economic impact of porcine reproductive and respiratory syndrome virus on United States pork

producers Assessment of. *J. Swine Heal. Prod.* **21**.

Jacquot, M., K. Nomikou, M. Palmarini, P. Mertens, and R. Biek, 2017: Bluetongue virus spread in Europe

is a consequence of climatic, landscape and vertebrate host factors as revealed by phylogeographic

inference. *Proc. R. Soc. B Biol. Sci.* **284**.

Jeong, J., S. Kim, C. Park, I. Kang, K.H. Park, H.J. Ham, and C. Chae, 2018: Effect of vaccination with a

porcine reproductive and respiratory syndrome subunit vaccine on sow reproductive performance in

endemic farms. *Vet. Rec.* **182**, 602.

Kalyaanamoorthy, S., B.Q. Minh, T.K.F. Wong, A. Von Haeseler, and L.S. Jermiin, 2017: ModelFinder:

Fast model selection for accurate phylogenetic estimates. *Nat. Methods* **14**, 587–589.

Kass, R.E., and A.E. Raftery, 1995: Bayes Factors. *J. Am. Stat. Assoc.* **90**, 773–795.

Kumar, S., M. Nei, J. Dudley, and K. Tamura, 2008: MEGA: A biologist-centric software for evolutionary

analysis of DNA and protein sequences. *Brief. Bioinform.* **9**, 299–306.

Laenen, L., S. Dellicour, V. Vergote, I. Nauwelaers, S. De Coster, I. Verbeeck, B. Vanmechelen, P. Lemey,

and P. Maes, 2016: Spatio-temporal analysis of Nova virus, a divergent hantavirus circulating in the

European mole in Belgium. *Mol. Ecol.* **25**, 5994–6008.

Lager, K.M., and W.L. Mengeling, 2000: Experimental aerosol transmission of pseudorabies virus and

porcine reproductive and respiratory syndrome virus. *Proc. Am. Assoc. Swine Pract.*409–410.

Lemey, P., A. Rambaut, A.J. Drummond, and M.A. Suchard, 2009: Bayesian Phylogeography Finds Its

Roots. (Christophe Fraser, Ed.)*PLoS Comput. Biol.* **5**, e1000520.

Linhares, D.C.L., J.P. Cano, M. Torremorell, and R.B. Morrison, 2014: Comparison of time to PRRSv-

stability and production losses between two exposure programs to control PRRSv in sow herds. *Prev. Vet.*

Med. **116**, 111–119.

Lycett, S.J., V.N. Tanya, M. Hall, D. King, S. Mazeri, V. Mioulet, N. Knowles, J. Wadsworth, K. Bachanek-

Bankowska, N.N. Victor, K.L. Morgan, and M. Bronsvoort, 2019: The evolution and phylodynamics of

serotype A and SAT2 foot-and-mouth disease viruses in endemic regions of Africa. *Sci. Rep.* **9**, 5614.

Machado, G., J.A. Galvis, F.P.N. Lopes, J. Voges, A.A.R. Medeiros, and N.C. Cárdenas, 2020: Quantifying

the dynamics of pig movements improves targeted disease surveillance and control plans.

*bioRxiv*2019.12.29.886408.

Mahesh, K.C., J. Bhoj Raj, S. Swoyam, Prakash, M. Prajapati, D. Kathayat, and S. Dhakal, 2015: Sero-

Prevalence of Porcine Reproductive and Respiratory Syndrome (PRRS) in Pigs of Different Developmental

Regions of Nepal. *Int. J. Appl. Sci. Biotechnol.* **3**, 218–222.

Malone, B., 2004: Using trees to reduce dust and odor emissions from poultry farms*Poult. Inf. Exch.*

Martin, D.P., B. Murrell, M. Golden, A. Khoosal, and B. Muhire, 2015: RDP4: Detection and analysis of

recombination patterns in virus genomes. *Virus Evol.* **1**.

Moon, S.A., T. Ferdousi, A. Self, and C.M. Scoglio, 2019: Estimation of swine movement network at farm

level in the US from the Census of Agriculture data. *Sci. Rep.* **9**.

MSHMP, 2020: PRRS charts [Online] Available at https://www.vetmed.umn.edu/centers-programs/swine-

program/outreach-leman-mshmp/mshmp/mshmp-prrs-figures (accessed 01/10/2020).

Müller, N.F., D. Rasmussen, and T. Stadler, 2018: MASCOT: Parameter and state inference under the

marginal structured coalescent approximation. *Bioinformatics* **34**, 3843–3848.

Müller, N.F., D.A. Rasmussen, and T. Stadler, 2017: The structured coalescent and its approximations.

Mol. Biol. Evol. **34**, 2970–2981.

Mur, L., B. Martínez-López, and J.M. Sánchez-Vizcaíno, 2012: Risk of African swine fever introduction

into the European Union through transport-associated routes: returning trucks and waste from international

ships and planes. *BMC Vet. Res.* **8**, 149.

Muroga, N., S. Kobayashi, T. Nishida, Y. Hayama, T. Kawano, T. Yamamoto, and T. Tsutsui, 2013: Risk

factors for the transmission of foot-and-mouth disease during the 2010 outbreak in Japan: a case--control

study. *BMC Vet. Res.* **9**, 150.

Murray, G.G.R., F. Wang, E.M. Harrison, G.K. Paterson, A.E. Mather, S.R. Harris, M.A. Holmes, A.

Rambaut, and J.J. Welch, 2016: The effect of genetic structure on molecular dating and tests for temporal

signal. *Methods Ecol. Evol.* **7**, 80–89.

Navascués, M., F. Depaulis, and B. Emerson, 2010: Combining contemporary and ancient DNA in

population genetic and phylogeographical studies. *Mol. Ecol. Resour.* **10**, 760–772.

Neumann, E.J., J.B. Kliebenstein, C.D. Johnson, J.W. Mabry, E.J. Bush, A.H. Seitzinger, A.L. Green, and

J.J. Zimmerman, 2005: Assessment of the economic impact of porcine reproductive and respiratory

syndrome on swine production in the United States. *J. Am. Vet. Med. Assoc.* **227**, 385–392.

Nguyen, L.T., H.A. Schmidt, A. Von Haeseler, and B.Q. Minh, 2015: IQ-TREE: A fast and effective

stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol. Biol. Evol.* **32**, 268–274.

Otake, S., S. Dee, C. Corzo, S. Oliveira, and J. Deen, 2010: Long-distance airborne transport of infectious

PRRSV and Mycoplasma hyopneumoniae from a swine population infected with multiple viral variants.

Vet. Microbiol. **145**, 198–208.

Otake, S., S. Dee, K. Rossow, R. Moon, and C. Pijoan, 2002: Mechanical transmission of porcine

reproductive and respiratory syndrome virus by mosquitoes, Aedes vexans (Meigen). *Can. J. Vet. Res.* **66**,

191–195.

Otake, S., S.. Otake, S., R.D. Dee, K.D. Moon, C. Roscow, C. Trincado, and C. Pijoan, 2003: Evaluation of mosquitoes, Aedes vexans, as biological vectors of porcine reproductive and respiratory syndrome virus. *Can. J. Vet. Res.* **67**, 265–270.

Patterson, P.H., A. Adrizal, R.M. Hulet, R.M. Bates, C.A.B. Myers, G.P. Martin, R.L. Shockey, M. Van Der Grinten, D.A. Anderson, and J.R. Thompson, 2008: Vegetative buffers for fan emissions from poultry farms: 2. Ammonia, dust and foliar nitrogen. *J. Environ. Sci. Heal. - Part B Pestic. Food Contam. Agric. Wastes* **43**, 96–103.

Perez, A.M., P.R. Davies, C.K. Goodell, D.J. Holtkamp, E. Mondaca-Fernández, Z. Poljak, S.J. Tousignant, P. Valdes-Donoso, J.J. Zimmerman, and R.B. Morrison, 2015: Lessons learned and knowledge gaps about the epidemiology and control of porcine reproductive and respiratory syndrome virus in North America. *J. Am. Vet. Med. Assoc.* **246**, 1304–1317.

Pileri, E., and E. Mateu, 2016: Review on the transmission porcine reproductive and respiratory syndrome virus between pigs and farms and impact on vaccination. *Vet. Res.* **47**, 1–13.

Pitkin, A., J. Deen, and S. Dee, 2009: Use of a production region model to assess the airborne spread of porcine reproductive and respiratory syndrome virus. *Vet. Microbiol.* **136**, 1–7.

Plain, R.L., and J.D. Laurence, 2003: Swine production. *Vet. Clin. North Am. Food Anim. Pract.* **19**, 319–337.

Pybus, O., M. Suchard, P. Lemey, F. Bernardin, A. Rambaut, F. Crawford, R. Gray, N. Arinaminpathy, S. Stramer, M. Busch, and E. Delwart, 2012: Unifying the spatial epidemiology and molecular evolution of emerging epidemics. *Natl. Acad Sci.* 1–6.

Pybus, O.G., and A. Rambaut, 2009: Evolutionary analysis of the dynamics of viral infectious disease. *Nat. Rev. Genet.* **10**, 540–550.

Rambaut, A., and A. Drummond, 2016: TreeAnnotator v.2.4. 3. . Part of the BEAST package.

Rambaut, A., T. Lam, L. Max Carvalho, and O. Pybus, 2016: Exploring the temporal structure of heterochronous sequences using TempEst (formerly Path-O-Gen). *Virus Evol.* **2**, vew007.

Rappole, J.H., B.W. Compton, P. Leimgruber, J. Robertson, D.I. King, and S.C. Renner, 2006: Modeling

movement of West Nile virus in the Western Hemisphere. *Vector-Borne Zoonotic Dis.* **6**, 128–139.

Rasmussen, D.A., E. Wilkinson, A. Vandormael, F. Tanser, D. Pillay, T. Stadler, and T. de Oliveira, 2018:

Tracking external introductions of HIV using phylodynamics reveals a major source of infections in rural

KwaZulu-Natal, South Africa. *Virus Evol.* **4**, vey037.

Rathkjen, P.H., and J. Dall, 2017: Control and eradication of porcine reproductive and respiratory syndrome

virus type 2 using a modified-live type 2 vaccine in combination with a load, close, homogenise model: An

area elimination study. *Acta Vet. Scand.* **59**, 4.

Reicks, D.L., 2019: Effective biosecurity to protect North American studs and clients from emerging

infectious disease. *Theriogenology* **137**, 82–87.

Sanhueza, J.M., C. Vilalta, C. Corzo, and A.G. Arruda, 2019: Factors affecting Porcine Reproductive and

Respiratory Syndrome virus time-to-stability in breeding herds in the Midwestern United States.

Transbound. Emerg. Dis. **66**, 823–830.

Sanhueza, J.M., C. Vilalta, M. Kikuti, E. Geary, and C. Corzo, 2019: Comparison of PRRS incidence rate

in status IV breeding herds during 2009-2019.

Schurrer, J., S. Dee, R. Moon, M. Murthaugh, C. Finnegan, J. Deen, S. Kleiboeker, and C. Pijoan, 2005:

Retention of ingested porcine reproductive and respiratory syndrome virus in houseflies. *Am Vet Med Assoc*

66, 1517–1525.

Shi, M., T. Tsan-Yuk Lam, C.-C. Hon, R. Kin-Hei Hui, K.S. Faaberg, T. Wennblom, M.P. Murtaugh, T.

Stadejek, and F. Chi-Ching Leung, 2010: Molecular epidemiology of PRRSV: A phylogenetic perspective.

Virus Res. **154**, 7–17.

Silva, G.S., G. Machado, K.L. Baker, D.J. Holtkamp, and D.C.L. Linhares, 2019: Machine-learning

algorithms to identify key biosecurity practices and factors associated with breeding herds reporting PRRS

outbreak. *Prev. Vet. Med.* **171**.

Sørensen, R., U. Zinko, and J. Seibert, 2005: On the calculation of the topographic wetness index:

evaluation of different methods based on field observations On the calculation of the topographic wetness

index: evaluation of different methods based on field observations Evaluation of TWI calculation Vol.

2*Hydrol. Earth Sys. Sci. Discuss*.

Sun, R.Q., R.J. Cai, Y.Q. Chen, P.S. Liang, D.K. Chen, and C.X. Song, 2012: Outbreak of porcine epidemic

diarrhea in suckling piglets, China. *Emerg. Infect. Dis.* **18**, 161–163.

Sun, Y.K., X.L. Han, Y.F. Wei, Z.Q. Yu, C.H. Ji, Q. Li, G. Lu, L. Shen, C.Q. Ma, H. Wang, and G.H.

Zhang, 2019: Phylogeography, phylodynamics and the recent outbreak of lineage 3 porcine reproductive

and respiratory syndrome viruses in China. *Transbound. Emerg. Dis.* **66**, 2152–2162.

Tavaré, S., 1986: Some probabilistic and statistical problems in the analysis of dna sequences. *Lect Math*

Life Sci. **17**, 57–86.

To, T.H., M. Jung, S. Lycett, and O. Gascuel, 2016: Fast Dating Using Least-Squares Criteria and

Algorithms. *Syst. Biol.* **65**, 82–97.

Trincado, C., S. Dee, L. Jacobson, S. Otake, K. Rossow, and C. Pijoan, 2004: Attempts to transmit porcine

reproductive and respiratory syndrome virus by aerosols under controlled field conditions. *Vet. Rec.* **154**,

294–297.

Trovão, N.S., M.A. Suchard, G. Baele, M. Gilbert, and P. Lemey, 2015: Bayesian inference reveals host-

specific contributions to the epidemic expansion of influenza A H5N1. *Mol. Biol. Evol.* **32**, 3264–3275.

Van Ryswyk, K., N. Prince, M. Ahmed, E. Brisson, J.D. Miller, and P.J. Villeneuve, 2019: Does urban

vegetation reduce temperature and air pollution concentrations? Findings from an environmental

monitoring study of the Central Experimental Farm in Ottawa, Canada. *Atmos. Environ.* **218**.

Velasova, M., P. Alarcon, S. Williamson, and B. Wieland, 2012: Risk factors for porcine reproductive and

respiratory syndrome virus infection and resulting challenges for effective disease surveillance. *BMC Vet.*

Res. **8**, 184.

Volz, E.M., K. Koelle, and T. Bedford, 2013: Viral Phylodynamics. *PLoS Comput. Biol.* **9**.

Yang, J., N.F. Müller, R. Bouckaert, B. Xu, and A.J. Drummond, 2019: Bayesian phylodynamics of avian

influenza A virus H9N2 in Asia with time-dependent predictors of migration. *PLoS Comput. Biol.* **15**.

Yang, Z., 1994: Maximum likelihood phylogenetic estimation from DNA sequences with variable rates

over sites: Approximate methods. *J. Mol. Evol.* **39**, 306–314.

Yuan, S., C.J. Nelsen, M.P. Murtaugh, B.J. Schmitt, and K.S. Faaberg, 1999: Recombination between North

American strains of porcine reproductive and respiratory syndrome virus. *Virus Res.* **61**, 87-98.

Table list

Table 1. PRRSv dissemination between systems. Bayes factor results represent the level of support in the
rates of the spread between systems, where BF > 20 is strong support, BF = 3-19 is positive support and
BF< 3 is negligible.

Origin	Destination	Average relative migration rate	Bayes factor	Interpretation
A	B	1.79 ± 0.92	102.1	Strong support
A	C	0.46 ± 0.39	47.8	Strong support
B	A	0.35 ± 0.37	27.4	Strong support
B	C	0.33 ± 0.13	7.9	Positive support
C	A	0.41 ± 0.22	14.7	Positive support
C	B	0.39 ± 0.16	10.3	Positive support

712   **Table 2.** PRRSv dissemination between types of farms. Bayes factor results represent the level of support
713   in the rates of the spread between types of farms, where BF > 20 is strong support, BF = 3-19 is positive
714   support and BF< 3 is negligible.

Farm type of origin	Farm type of destination	All systems (Bayes factor)	System A (Bayes factor)	System B (Bayes factor)	System C (Bayes factor)
Sow	Nursery	109	56.2	42.2	51.7
Sow	Finisher	80.5	36.6	28.1	17.9
Sow	Boar stud	0.1	2.2	0.6	1.1
Nursery	Sow	7.3	5.8	4.9	4.6
Nursery	Finisher	27.5	33.7	10.9	16.6
Nursery	Boar stud	1.9	0.1	0.7	1.3
Finisher	Sow	18.9	10.9	14.1	5.3
Finisher	Nursery	12	16.7	13.2	6.4
Finisher	Boar stud	0.2	1.4	0.1	0.6
Boar stud	Sow	0.5	1.1	0.3	0.1
Boar stud	Nursery	3.9	2.5	2.1	2.4
Boar stud	Finisher	0.4	2	0.8	1.5

716   **Table 3.** Number of transmission events that have been originated in farms that belong to the same pig
717   system (local) compared to the events coming from farms integrated into another pig system (external).

System	Year	Local		External		Total
		Events	%	Events	%	

	Year					
A	2014	807	54.9	663	45.1	1470
	2015	629	60.5	410	39.5	1039
	2016	598	59.2	412	40.8	1010
	2017	527	60.3	347	39.7	874
	2018	215	58.4	153	41.6	368
B	2014	1030	90.7	106	9.3	1136
	2015	851	89.5	100	10.5	951
	2016	1432	90.4	152	9.6	1584
	2017	771	91.5	72	8.5	843
	2018	322	94.4	19	5.6	341
C	2014	1100	89.8	125	10.2	1225
	2015	826	89.0	102	11.0	928
	2016	789	92.2	67	7.8	856
	2017	684	92.1	59	7.9	743
	2018	333	95.1	17	4.9	350

719 **Figure legend**

721 **Figure 1.** Diagram of factors promoting or restricting PRRSv spread. Arrows represent the relationship

722 between the predictor and PRRSv occurrence based on Bayes factor analysis, where green and orange

723 lines represent positive and negative drivers respectively. Black represent negligible support, and dashed

724 lines show relatedness between environmental predictors affecting PRRSv through other variables.

Figure 2. Dissemination history of PRRSv, inferred by discrete phylogeographic analysis. A) Maximum

clade credibility phylogeny, colored according to pig production systems (A, B, C). The probabilities of

ancestral states (inferred from the Bayesian discrete trait analysis) are shown in pie charts on each node,

representing the most likely type of farms. B) Spatiotemporal patterns in the relative genetic diversity

represented through the Bayesian SkyGrid plot and the number of sequences per month. The left y-axis

summarizes the effective population size over time, the mean estimate is represented by the dark blue

line, while the shaded light blue regions correspond to the 95% highest posterior density (HPD), and in

the right y-axis shows the number of sampled sequences over time.

Figure 3. PRRSv dissemination within system A. Panel A) Maximum clade credibility phylogeny colored

according to the different farm types (sow, nursery, finisher, and boar stud). Probabilities of ancestral

states shown in pie charts in nodes, representing the most likely type of farm origin. Panel B)

Spatiotemporal patterns in the relative genetic variation through Bayesian SkyGrid plot and number of

sequences per month. The left y-axis summarizes the effective population size over time. Mean estimate

is represented by the dark blue line. Shaded light blue regions correspond to the 95% highest posterior

density (HPD), and in the right y-axis shows the number of sampled sequences over time.

Figure 4. PRRSv dissemination within system B. Panel A) Maximum clade credibility phylogeny colored

according to the different farm types (sow, nursery, finisher, and boar stud). Probabilities of ancestral

states shown in pie charts in nodes, representing the most likely type of farm origin. Panel B)

Spatiotemporal patterns in the relative genetic variation through Bayesian SkyGrid plot and number of

sequences per month. The left y-axis summarizes the effective population size over time. Mean estimate

is represented by the dark blue line. Shaded light blue regions correspond to the 95% highest posterior

density (HPD), and in the right y-axis shows the number of sampled sequences over time.

Figure 5. PRRSv dissemination within system C. Panel A) Maximum clade credibility phylogeny colored

according to the different farm types (sow, nursery, finisher, and boar stud). Probabilities of ancestral

states shown in pie charts in nodes, representing the most likely type of farm origin. Panel B)

Spatiotemporal patterns in the relative genetic variation through Bayesian SkyGrid plot and number of

sequences per month. The left y-axis summarizes the effective population size over time. Mean estimate

is represented by the dark blue line. Shaded light blue regions correspond to the 95% highest posterior

density (HPD), and in the right y-axis shows the number of sampled sequences over time.

Figure 6. PRRSv migration events among pig systems are inferred through a structured coalescent model. Number of dissemination events originated in farms from the same pig system compared to events from farms integrated into another pig system. Grey background represents the months where the exponentially weighted moving average epidemic monitor surpassed the established PRRSv epidemic threshold based on the February 2020 MSHMP report (MSHMP, 2020), while dashed lines represent the beginning of each year.