Porcine reproductive and respiratory syndrome virus dissemination across pig production systems

in the United States

Running title: Evolutionary epidemiology of PRRSV

Manuel Jara[1], David A. Rasmussen[2,3] ,Cesar A. Corzo[4], Gustavo Machado[1*]

[1]Department of Population Health and Pathobiology, College of Veterinary Medicine, North Carolina

State University, Raleigh, NC, USA.

[2]Department of Entomology and Plant Pathology, North Carolina State University, Raleigh NC, USA

[3]Bioinformatics Research Center, North Carolina State University, Raleigh NC, USA

[4]Veterinary Population Medicine Department, College of Veterinary Medicine, University of Minnesota,

1365 Gortner Ave, St Paul, MN 55108, USA.

[*]**Corresponding author**: gmachad@ncsu.edu

Summary

Porcine reproductive and respiratory syndrome virus (PRRSV) remains widespread in the North

American pig population. Despite improvements in virus characterization, it is unclear whether PRRSV

infections are a product of viral circulation within a farm, within production systems (local) or across

production systems (external). Here we examined the dissemination dynamics of PRRSV and the

processes facilitating its spread within and among pig farms in three production systems. Overall, PRRSV

genetic diversity declined since 2018, while phylodynamic results support frequent transmission across-

production systems. We found that PRRSV dissemination occurred mostly through transmission between

farms of different production companies, which were predominant for several months, especially from

November until May when PRRSV tends to peak in the studied region. Within production systems,

dissemination occurred mainly through regular pig flow (from sow to nursery and then to finisher farms);

nevertheless, an important flux of PRRSV dissemination from finisher to sow and nursery farms

highlighted the importance of downstream farms as sources of the virus. Farms at areas with pig density

from 500 to 1000 pig/km^2 and farms located at a range within 0.5 km and 0.7 km from major roads were

more likely to infect by PRRSV, whereas farms at elevation between 41 and 61 meters and denser

vegetation acted as dissemination barriers. Although remains a challenge, there is a need to disentangle

the route of PRRSV transmission, results evidenced that dissemination among commercially unrelated pig

production systems was intense, reinforcing the importance of farm proximity on PRRSV spread. Thus,

consideration of farm location and their geographic characteristics may help to forecast dissemination.

The understanding of PRRSV transmission routes has the potential to inform targeted strategies for its

prevention and control. Further studies are needed to quantify the relative contribution of PRRSV

transmission routes.

Keywords: molecular epidemiology, phylodynamics, virus dispersal, between-farm dissemination.

Introduction

Porcine reproductive and respiratory syndrome virus (PRRSV) remains among the most costly diseases in

North America (Neumann et al., 2005; Holtkamp et al., 2013; Pileri and Mateu, 2016). Despite the great

advances in reducing the incidence of PRRSv in some regions (Rathkjen and Dall, 2017) by enhanced

biosecurity (Silva et al., 2019) and immunization strategies (Corzo et al., 2010), every year approximately

20-30% of US breeding herds still become infected with PRRSV (MSHMP, 2020).

PRRS is caused by an enveloped positive single-stranded RNA virus in the family Arteriviridae,

order Nidovirales (Cavanagh et al., 1990). The genome of PRRSV is ~15 kb and consists of 9 open

reading frames (ORFs) that encode seven structural proteins and 14 non-structural (Dokland, 2010). The

mutation rate of PRRSV hinders the effectiveness of immune response against newly introduced variant

strains, especially in farms with recurrent outbreaks (Linhares et al., 2014; Sanhueza et al., 2019).

Likewise, recombination between North American wild-type virus strains and modified-live vaccines

represents an important mechanism in the historical evolution and spread of PRRSV (Yuan et al., 1999).

The phylodynamic patterns of PRRSV spread and dissemination among pig systems and farm

types have been previously described (Shi et al., 2010; Franzo et al., 2015; Alkhamis et al., 2016;

Alkhamis et al., 2017; Sun et al., 2019). While there was support for between-production system

dissemination (Alkhamis et al., 2016), the relevance of transmission in the maintenance of local

incidence, in comparison with within-production system circulation, remains unknown. Thus, the

question of whether PRRSV circulation is the product of local dissemination between farms that are part

of a common commercial network (production system) versus the result of external viral introductions

from nearby infected farms of distinct production systems still remains (Arruda et al., 2018).

Bayesian phylodynamic approaches have provided a breakthrough in tracking pathogen

dissemination dynamics (Baele et al., 2018; Faria et al., 2019; Hicks et al., 2020). Structured coalescent

models and other phylodynamic approaches allow the estimation of relative contribution of local

dissemination versus external introductions to the overall incidence of disease (Rasmussen et al., 2018;

Yang et al., 2019; Hicks et al., 2020). Novel methods have been developed to integrate spatial and

environmental covariates in phylodynamic analysis, to determine their influence on the spread of

epidemics (Dudas et al., 2017; Fountain-Jones et al., 2018; Rasmussen et al., 2018; Dellicour et al., 2019;

Yang et al., 2019). Thus far, such approaches have not been widely implemented to study the

dissemination of diseases in domestic animals, except for Foot-and-Mouth virus (Duchatel et al., 2019).

In this study, we used phylodynamic methods to determine the viral dissemination rates among

pig production systems to assess whether different PRRSV variants are the product of local transmission

or external introductions and identified the factors that facilitate or impede PRRSV spread.

Material and methods

Data collection

In total, our dataset comprises 4,970 ORF-5 sequences collected from 2014 to 2019, obtained from three

commercially unrelated pig production systems (here coded as A, B, and C), in the US. Production

system A, B and C do not shared feed truck or traded pig among themselves. Sequences were generated

through outbreak investigation and from regular surveillance activities performed by each production

system and shared with the Morrison Swine Health Monitoring Program (MSHMP). Each sequence was

attributed to a farm type (e.g., sow, nursery, finisher, and boar stud), geographic location, pig production

system (A, B, and C), and the date when the sequence was reported.

Preliminary phylogenetic analysis

Sequences were aligned using Mega X, available at www.megasoftware.net (Kumar et al., 2018). The recombination detection program (RDP) v5.3 was used to search for recombination within our dataset (Martin et al., 2015). The alignment was screened using five methods (BootScan, Chimaera, MaxChi, RDP, and SiScan), evidencing 63 recombinant sequences, which were removed from the downstream analysis (Table S1). To determine whether there was a sufficient temporal molecular evolutionary signal, we used TempEst v1.5 (Rambaut et al., 2016). To calculate the P-values associated with the phylogenetic signal analysis, we followed Murray et al. (2016) using 1,000 random permutations of the sequence sampling dates (Navascués et al., 2010). The relationship found between phylogenetic tree root-to-tip divergence and sampling dates (years) supported the molecular clock analysis for this study ($P < 0.05$).

Discrete and continuous phylogeographical analysis

Phylogenies were estimated by Bayesian inference through Markov chain Monte Carlo (MCMC), implemented in BEAST v2.5.0 (Bouckaert et al., 2014). By using ModelFinder (Kalyaanamoorthy et al., 2017) built into IQ-Tree version 1.6.1 (Nguyen et al., 2015) Supplementary Table S2, we compared different substitution models to find the best fit for our genetic dataset. The marginal likelihood value supported the use of the general time-reversible model (GTR) with gamma-distributed rate heterogeneity plus a proportion of invariable sites (GTR+G+I) (Tavaré, 1986). To reconstruct PRRSV dissemination across systems and between farm types (i.e., sow, nursery, finisher, boar stud), we used the discrete-trait phylogeographic models implemented in BEASTv2.5.0 to reconstruct ancestral states (Bouckaert et al., 2014). To explore the most important historical dissemination routes for PRRSV spread across farms, as well as among pig systems, we used Bayesian stochastic search variable selection (BSSVS) (Lemey et al., 2009). Through BSSVS, we identified the non-zero rates of change between each pair of discrete traits (production system and farm types) based on its Bayes factor value (lower than 3). To perform this analysis, an asymmetric migration rate matrix was assumed. To infer the intensity of directional transitions (forward and backward) within the migration matrix, we used a Markov jump approach. To evaluate oversampling and possible selection bias towards location and/or farm types with

disproportionate amounts of samples (i.e., sow farms), we downsampled on each system to match the

number of samples between the three systems, maintaining the same proportion of farm type (system A -

89%, B =-39.5%, and C=31.4 %).

To analyze the spatiotemporal spread of PRRSV, we performed a Bayesian continuous

phylogeographic analysis, where the nucleotide substitution model was similar to the discrete model

mentioned above. In addition, we used an uncorrelated relaxed molecular clock model with lognormal

distribution (Drummond et al., 2006). The coalescent tree prior used was Bayesian SkyGrid with

covariates (Gill et al., 2016). Phylogeographic inference of ancestral locations was performed under a

relaxed random clock model, considering a lognormal distribution. All analyses were run for 200 million

generations, sampling every 10,000th generation and removing 10% of the chain as burn-in. To estimate

the relative genetic diversity, we used skygrid which is based on a nonparametric coalescent model, and is

utilized to calculate the effective population size over time (Hill and Baele, 2019). To visualize the

spatiotemporal spread of PRRSV, we used Spatial Phylogenetic Reconstruction of Evolutionary

Dynamics using Data-Driven Documents (D3) SPREAD3 software (Bielejec et al., 2016). To evaluate the

impact generated by the imbalance distribution of sequences among systems and production types, a

sensitivity analysis was performed. A subsample taken from the systems and farm types with the highest

number of sequences was used to match the sample size in the systems with the least number of

sequenced, thus this smaller sample was reanalyzed (Supplementary material S6).

Spatiotemporal epidemiological statistics

To summarize PRRSV diffusion over time and space we analyzed data from each production system

separately and all the sequences as a whole, using the R package "seraphim" version 1.0 (Dellicour et al.,

2016). Spatiotemporal information about lineage locations was extracted from 100 input trees sampled at

regular intervals from the post-burn-in posterior distribution to account for phylogenetic uncertainty

(Dellicour, Rose and Pybus, 2016). Each phylogeny branch was summarized and represented as a distinct

vector defined by its starting geographic location, ending geographic location and dates (Dellicour, Rose

and Pybus, 2016; Dellicour et al., 2016). Each branch vector represents an independent viral lineage

dissemination event (Pybus et al., 2012; Laenen et al., 2016). Each vector was assigned to a diffusion

coefficient value (*D*) for subsequent visualization: as described by (Pybus et al., 2012), which accounts

for the time (in years) during which each specific vector (lineage) moved from its initial location until the

end location of each lineage (measured from the location of the tree root). These vectors were also used to

assess the invasion velocity of the sampled lineages. To determine and compare the PRRSV diffusivity

associated with the overall data and for each production system, we estimated the spatial diffusion

coefficient using *D*-weighted statistics (Trovão et al., 2015) since this index is less sensitive to extreme

values of short branches. These results were displayed via: i) dissemination epidemiological statistics that

estimated the mean branch dissemination velocity, and ii) kernel plots showing the lineage diffusion

coefficients of the mean and variation on the highest posterior density (HPD).

Between-production systems transmission: the contribution of within-system PRRSV circulation

(local) versus external introduction of virus from a commercially unrelated pig farm.

We estimated the contribution of local transmission (i.e., between farms within a production system)

versus external introductions (i.e., between farms across production systems) using a structured

coalescent model. Due to computational demand, we used PRRSV alignment from MEGA software

(Kumar et al., 2008) to produce a phylogenetic tree using PhyML version 3.0 (Guindon et al., 2010),

using a substitution rate = 0.00672 (estimated), and considering an HKY Hasegawa-Kishino-Yano + $\Gamma 4$

(Hasegawa et al., 1985; Yang, 1994), which was dated using Least Squares Dating (LSD) version 0.3 (To

et al., 2016). To infer the contribution of local versus external introductions the migration rates between

different pig production systems were estimated using the Marginal Approximation to the Structured

Coalescent (MASCOT) package (Müller et al., 2017, 2018) implemented in Beast2 (Bouckaert et al.,

2014). To infer the evolutionary rate from the time-calibrated tree, we used a strict clock model. Using

MASCOT, we estimated the effective population size in the local and external population and the

migration rates between them, which was allowed to be asymmetric. This analysis was replicated three

times, each time using one of the three pig systems as local and the other two as external. We also

calculated the posterior probability of each internal node in the tree being in each state (i.e., local or

external) using the forwards/backwards approach implemented in MASCOT (Müller et al., 2018).

All analyses were performed under 100 million MCMC runs, states were sampled every 10,000 and the

first 10% of samples were discarded as burn-in using TreeAnnotator v. 2.3.0 (Rambaut and Drummond,

2016). We reconstructed the number of migration events through time using the ancestral locations

reconstructed by MASCOT. A migration event was assumed to occur when the most probable location of

a lineage changed between its parent and child node, with the time of the migration event assumed to be

the date of the daughter node. Migration events (local and external) per month were plotted using "baltic"

(https://github.com/blab/baltic) in Python Software Foundation version 3.7, available at

http://www.python.org. Finally, we compared the differences in local vs external transmission between

PRRSV season and off-season through two-sample Wilcoxon test (Hogg et al., 2010), PRRSV season was

determined by the exponentially weighted moving average proposed by MSHMP (MSHMP, 2020). In the

same way as we described in the section above, a sensitivity analysis with a subsample was evaluated for

consistency of the results (Supplementary material S8).

Factors promoting or restricting PRRSv spread

To determine the effect of different predictors on the spread of PRRSV, we analyzed the results of the

continuous phylogeographic analysis using the R package "seraphim" version 1.0 (Dellicour et al., 2016),

considering a continuous location approach (based on the geographic coordinate location of each pig

farm) and a random walk diffusion model. Spatiotemporal information was extracted from 100 trees

sampled at regular intervals from the posterior distribution, after burn-in, to account for phylogenetic

uncertainty. Each phylogenetic branch was considered a vector defined by its start and end location (e.g.

latitude and longitude), and its start and end dates. Statistical significance for the correlation between

phylogenetic and predictive factors was tested using 100 trees generated and expressed in the form of

Bayes factors (BF) (Dellicour et al., 2016). Based on the extracted spatio-temporal information contained

in each phylogenetic tree described above, we calculated the distances based on predictor variables

associated with every branch of each tree, which were used to determine the correlation between the

182 duration of each branch in the phylogeny and its related predictor variable distance, including the annual

183 temperature, annual precipitation, elevation, runoff (an index that quantity of water discharged in surface

184 streams), soil humidity (commonly used to quantify topographic control on hydrological processes)

185 (Sørensen et al., 2005), enhanced vegetation index (EVI), which can be used to quantify vegetation

186 greenness, as well as the pig density, and distance to the main roads (Fig. 1 and Supplementary Table S3

187 for more details). Briefly, the distance between each farm (geographic coordinates) to the nearest main

188 road was calculated based on the Euclidean distance. We calculated the statistic Q, which represents how

189 much variation in lineage movement is explained with spatial heterogeneity when each predictor variable

190 is considered (Dellicour, Rose and Pybus, 2016). Here, $Q= R^2_{env} - R^2_{null}$ where R^2_{env} is the coefficient of

191 determination obtained from the regression between branch duration against predictor (factors) distances,

192 while R^2_{null} represents the coefficient of determination for the regression between branch duration against

193 predictor distances, which were replaced by a null raster. Thus, the factors that were considered for

194 further Bayes factor (BF) analysis, were the ones that obtained positive Q values in at least 90% of the

195 tested trees (Jacquot et al., 2017). We approximated a BF value for all predictors tested that passed the

196 above-mentioned Q statistic. Predictive factors were treated as a conductance (variables that promote the

197 spread of the disease) or a resistance factor (variables that impede its spread). For the interpretation of BF,

198 values between 3-20 were considered "positively" supported, values between 20-150 were considered

199 "strong", and values >150 were considered " overwhelming support" (Kass and Raftery, 1995). For all

200 variables that showed a significant support either as conductance of resistance factors (considering Q

201 statistic and Bayes factor) (Supplementary Table S10), each raster was subdivided into four alternative

202 ranges based on Jenks natural breaks, for example, for elevation raster values below 20 meters constituted

203 the low elevation areas, from 21-40 meters medium-low elevation, 41-60 medium-high elevation and >61

204 high elevation (Supplementary Table S11). The same continuous phylogeographic analysis using the R

205 package "seraphim" was applied individually for each range raster as detailed above.

206 **Results**

207 **Study population**

A total of 63 recombinant PRRSV sequences were identified and removed from the database. Most

recombinants (52%) belonged to system B with the highest number of recombinants (82%) belonging to

sow farms (Supplementary Table S1). Thus, 4,753 sequences remained for further discrete and continuous

phylogeographic analyses. Most sequences originated from system A = 3,748 (79%), while B had 626

(13%), and C had 379 (8%). Overall, 2,958 sequences (62.2%) belonged to sow farms, 909 (19.1%) to

finishers, 873 (18.4%) to nurseries, and 13 (0.3%) to boar studs (Supplementary Table S1). The nearest-

neighbor distances among farms of distinct production systems had a median value of 7.6 km (1st

quartile= 3.6 km and 3rd quartile= 19.3 km), the closest related production systems were A and C with

median distance among all pairwise farms of 1.8 km (1st quartile= 1.2 km and 3rd quartile= 2.5 km), and

finisher farms of system A and C were the closest with median of 1.9 km (1st quartile= 1.3 km, 3rd

quartile= 3.2 km) (Supplementary Table S4 and S5).

Discrete phylogeography analysis

The results of PRRSV ancestral reconstruction showed that the most likely center of origin was

production system A with a root state posterior probability (RSPP) = 0.84, coming from a sow farm

(RSPP = 0.75). From there, it spread heterogeneously across the other systems, unrestricted to farm types

(Fig. 2A). The reconstructed phylogenetic relationship did not show evidence of any cluster related to

production systems. However, we observed a clear dominance of PRRSV originating from system A over

the other systems (Fig. 2A). Our sensitivity analysis on the downsample supported these results

(Supplementary Table S6). The relative genetic diversity expressed as the effective population size (e.g.

number of individuals contributing to a new PRRSV infection) over time showed a plateau until 2012.

After that time, we captured a steep increment that lasted until approximately 2016, when relative genetic

diversity decreased through June 2019 (Fig. 2B), the last month of data available for our study. The

temporal pattern shown by the sampled PRRSV sequences evidenced the typical annual trend of seasonal

outbreaks (Fig. 2B) (MSHMP, 2020). Phylogeographic results of system A showed that its most likely

center of origin was a sow farm (RSPP = 0.47), followed by an almost equal probability for origins in

finisher farm (RSPP = 0.27) and nursery farm (RSPP = 0.26), the boar stud farms were negligible (RSPP

< 0.01) (Fig. 3A). In recent years, there was an evident increase in the proportion of PRRSV positive

finisher farms, reaching a similar level of representation than sow farms for some clusters (Fig. 3A).

SkyGrid plot revealed a relatively constant trend of relative genetic diversity until ~2011, showing a

linear increase in population size, which reached its highest levels ~2016 (Fig. 3B). We observed a

similar trend when we analyzed all systems together (Fig. 3B), 2015 and 2018 were the years with more

samples sent to be sequenced (Fig. 3B).

For production system B, a sow farm (RSPP = 0.4) was also the most likely origin; however, a

high probability was also linked to a nursery farm (RSPP = 0.33) (Fig. 4A). The temporal pattern shown

by the effective population size was static until 2012, after which it increased rapidly until 2015-2016,

when it started a linear decrease until June 2019, similar to the seasonal pattern shown by the number of

sequences (Fig. 4B). As observed in systems A and B, the most likely center of origin for system C

occurred inside a sow farm (RSPP = 0.52), followed by a high probability for originating in a finisher

farm (RSPP = 0.4). In contrast to the other systems, we observed a low probability of origin related to a

nursery farm (RSPP > 0.1) (Fig. 5A). The temporal trend in the number of sampled sequences showed the

mentioned annual seasonal pattern, with peak occurrence ~2015. SkyGrid plot evidenced a more stable

pattern, with higher levels of variation in effective population size during the period 2013-2014, then

decreasing until 2018, when it showed a small peak and then decreased through June 2019 (Fig. 5B).

Phylogeographic diffusion analysis and spread statistics

The reconstructed spatiotemporal diffusion of PRRSV was characterized by continuous expansion in the

geographic space, accompanied by diversification events. The comparison between each system and all

samples together showed two noticeable patterns. The first one was characterized by a radial expansion

over time, found when all samples were analyzed together and within samples of system A. In contrast,

systems B and C evidenced a directional trend, since for both systems the spread started in the southwest

region. In the case of system B, PRRSV spread to the northeast, while for system C, the spread was

mainly to the north, though spread did occur in all directions (Supplementary Figs. S1-3).

The spatiotemporal analysis showed that when all sequences were analyzed together, the

estimated median value for the mean branch velocity was 39 km/year (95% HPD = 31.2-61.1). This

median value was lower than that shown by system A (49.2 km/year, 95% HPD= 42.9-60.4), and higher

than the values observed in systems B and C (32.7 km/year, 95% HPD= 29.1-43.3), and (26.4 km/year,

95% HPD= 20.1-41.0), respectively. Estimated diffusion coefficients were asymmetric for each system

and when all samples were analyzed together. System A showed the highest diffusivity (243.8 km^2/year);

however, its diffusion variation among branches was the lowest registered. The overall results (all

sequences) evidenced the highest among-branch variation, while its diffusion coefficient (150 km^2/year)

was lower than system A and substantially higher than the other systems. Systems B and C showed a

similar pattern in terms of diffusion coefficient variation among branches; however, B was higher than C

(97 km^2/year) and (51 km^2/year), respectively (Supplementary Fig. S4).

PRRSv dissemination among pig systems and farm types

Bayesian stochastic search variable selection (BSSVS) identified that the most significant disseminations

were highly related to system A, evidencing strong support by a Bayes Factor score > 20. The statistical

support or evidence for system A sourcing most PRRSV strains to system B, BF = 102.1 (average relative

migration rate "ARMR"= 1.79), followed by dissemination from A to C, BF = 47.8 (ARMR = 0.46); of

less importance was from B to A, BF= 27.4 (ARMR = 0.35). All the other disseminations between

production systems were also positive, but to a lesser degree than the ones mentioned above (Table 1). In

addition, a sensitivity analysis aimed to evaluate the bias related to oversampling showed that while

numerically it was a difference the interpretation remains the same, but we noticed that in the subsample

there was even more support for the dissemination from company B to C and company C to A

(Supplementary Table S7 for more details). Our analysis reveals that when comparing PRRSV

dissemination between types of farms, the most frequent routes were linked with sow farms, especially

when the direction of spread was from sow to nursery (BF > 42). The highest support for this route

occurred when we analyzed all systems combined (BF = 109). The second most well-supported

dissemination occurred from sow to finisher farms (BF > 17.9), and as in the previous case, was best

supported when we analyzed all systems together. (BF = 80.5). Likewise, the dissemination between

nursery and finisher farms (in both directions) also showed positive support (BF > 6). However, all routes

of dissemination that involved boar stud farms were rare events (BF < 3) (Table 2). Furthermore, the

sensitivity analysis showed similar trends, where the results of the equal sample sizes these were less

sharp than the one using the original dataset. However, the transmissions that involved BoarStud were all

negligible (Supplementary Table S8 for more details).

Within-production systems circulation (local) versus introduction of PRRSV from commercially

unrelated farms (external)

We tracked the origin of the migration events leading to the current PRRSV lineages circulating among

the three pig systems, to determine whether detected PRRSV strains originated from within the same

system (local) or from farms of non-commercially related pig systems (external introductions). For this

analysis, we considered each coalescent event in the tree as a proxy for the timing of a transmission event

and then classified each event as either a local transmission event if the parent and child lineages were

both reconstructed to be in the local population or an external introduction if the parent lineage resided

outside the local population. Overall, results showed a clear dominance of external transmission events

that were observed in all pig companies; this tendency was constant during the whole period of study

(Fig. 6). System C predominantly had external introductions which peaked in 2015 with 94.7% of the

transmission events, while systems A and B exhibited 82.0% in 2018 and 82.4% in 2017, respectively

(Table 3). In addition, our results show that overall, the external introduction events tend to increase over

time (Table 3). We also observed that coinciding with the month of higher incidence of PRRSV

(MSHMP, 2020) (Fig. 6 and Supplementary Table S9). Two-sample Wilcoxon test indicated that the

difference between the within farm transmission and external introductions were statistically significantly

higher during PRRSV season (mean= 69.3) than the rest of the year (mean= 60.1) (Z= -9.9, p< 0.001).

Factors promoting or restricting PRRSV dissemination direction

To quantify the impact of each factor on PRRSV distribution, we calculated the association between

lineage duration and each covariate scaled distance through the Q statistic method. We found strong

support (Q value was positive) for pig density and distance to main roads treated as conductance factors,

while elevation and vegetation presented strong support as resistance factors (Supplementary Table S10

and Supplementary Figs. S5 and S6 for more details). These results also received statistical support

provided by Bayes factors (BF) since we observed that farms located in areas with a high density of pig

farms were more exposed to the introduction of PRRSV (BF > 12). Likewise, the distance between the

farm and the nearest major road also showed a positive association with PRRSV diffusion (BF > 7). In

contrast, the factors identified as negative drivers, variables that act as barriers for the spread of PRRSV,

were higher elevation and denser vegetation (BF > 3) (Supplementary Table S10). With respect to the

analysis of the variables with significantly supported as conductance, analyzed as ranges (Supplementary

Table S11), the density of pigs ranging from 501-1.000 number of pigs/km^2 was sufficient to facilitate the

spread of PRRSV when all sequences were analyzed together and for system A, while farms at B and C

were more likely infected when pig density was above 1.000 pigs/km^2. Likewise, farms located at a

distance range of 1 km to 2 km of main roads showed to favor PRRSV introductions, while when systems

were analyzed in separately the dangerous distance was between 0.5 km and 1 km (Supplementary Table

S11). On the other hand, for variables acting as resistance factors, our results were consistent in which at

elevation above 61 meters was sufficient to slower PRRSV movement, but for system C range of 41-61

meter was sufficient. The results of vegetation index were in disagreement, while for all systems together

there was no analytical support, but farms at system A, B and C an EVI index between 41 and 45 acted as

a resistance range (Supplementary Table S11).

Discussion

This analysis detailed the evolutionary epidemiology of PRRSV among commercially related and

unrelated farms through continuous and discrete phylogeographic analyses. Overall, a decline in PRRSV

genetic diversity was observed over time. Such findings could have different explanations, one being a

rapid and efficient disseminating strain that was introduced into sow farms which consequently was

transferred to growing pig sites through regular pig movement. Another potential explanation could be

related to the efficacy of current interventions that local production systems have implemented, which

reduced the volume and frequency of newly emerging strains, suggesting a more optimistic scenario than

proposed in a similar study done in another US region (Alkhamis et al., 2017). Given the economic

impact and herd health relevance of new PRRSV strain introductions, it is imperative to evaluate how

often PRRSV spreads among pig production systems, determine the role of each farm type, and identify

factors that may facilitate or keep PRRSV from spreading. We demonstrated that PRRSV circulation has

been mostly originated by transmission from outside of each pig production system, and this tendency is

more predominant during the PRRSV season (from November until May) (MSHMP, 2020).

Our findings suggest that virus transmission between pig systems is common. It is not well

understood which transmission mechanism plays a role in the dissemination across systems. Airborne

spread has been suggested as a potential route (Dee et al., 2009; Otake et al., 2010). However, Arruda et

al., (2018, 2019) did not reach that same conclusion. Pig farm density as expected arises as a risk factor in

several studies and in our study, our phylodynamic findings agree with a regional spread as an important

factor for PRRSV dissemination among pig farms. PRRSV spread events appear to be relatively common

in farms located in highly dense pig areas and near main roads, while farms at higher elevations and

surrounded by denser vegetation were significantly less likely to report new cases of PRRSv (Arruda et

al., 2017). It is important to mention that the source of our sequence data represented a limitation for our

phylodynamic analysis since it was a convenience sample that can introduce bias into our analysis as the

number of sequences does not directly represent all PRRSV cases circulating in this study region. There

was important asymmetry in the number of sequences from boar studs, thus results of this farm type

should be interpreted with caution.

PRRSV spatiotemporal dynamics

The spatiotemporal dynamic pattern exhibited by PRRSV is the result of its transmission dynamics and

how circulating strains have been affected by viral selection (Pybus and Rambaut, 2009; Volz et al.,

2013). This can be observed in the shape of PRRSV phylodynamic patterns over the course of this study,

which shows a clear directional selection, which can result in the loss of genetic variation (Volz et al.,

2013). Based on MSHMP data (MSHMP, 2020), vaccine usage increased between 2012 and 2014, which

could have impacted PRRSV genetic diversity for system A after 2012. But the increment in the number

of farms using vaccines during 2014 did not show a noticeable effect on PRRSV genetic variation.

However, the increment in the application of MLV's in mid-2015 showed an undeniable positive effect

on decreasing PRRSV genetic diversity (as seen in Figs. 2-5), similarly to the observations made by other

authors (Sun et al., 2012; Jeong et al., 2018). However, to increase the robustness of future studies it is

necessary to increase the farm coverage as well as the timeframe, not only related to the PRRSV genetic

information but also to the vaccination data.

The estimated speed in which PRRSV was spreading was 39 km/year which was slightly higher

than previous observations in a study that used data from one production system in another U.S. region

(34 km/year) (Alkhamis et al., 2017). This difference may be related to pig farm densities, being our

study region more populated with an average of 3 farms per 5 km^2, and a more loyal representation of

PRRSV dynamics since we compiled samples of more than 92% of all farms in our study area..

Consequently, there are important aspects that have to be considered while analyzing the speed at which

PRRSV spreads, for example, those related with the overall pig health (i.e., absence of coinfections) and

factors related with on-farm biosecurity (i.e., presence of a line of separation, cleaning, and disinfection

station) (Rappole et al., 2006; Altizer et al., 2011; Lycett et al., 2019). Thus, further studies are needed to

evaluate how biosecurity practices and overall pig health may contribute in slowing or accelerating

PRRSV spread, results of these assessments can be used to predict the most effective biosecurity

measures and optimize control strategies that go beyond PRRSV (Silva et al., 2019), not to mention

current diseases that pose large-scale biosecurity threats to the U.S swine health such as African Swine

Fever.

PRRSV spread among and within farms of commercially unrelated production systems

Our analysis found relevant similarities in the spatiotemporal patterns of PRRSV spread among all farm

types of all production systems (Table 1, 2), which was similar to previously reported findings that used

data from one production system in the U.S. (Alkhamis et al., 2016). At farm type levels, sow farms were

by far the biggest spreader (Plain and Laurence, 2003) which was expected as they wean piglets on a

weekly basis and these can be transported several miles away to an off-site growing pig facility.

Phylogenetic results also showed that PRRSV introductions into sow and nursery farms likely originated

from finisher farms, which could be partially related to gilt development units sending replacement gilts

to sow farms (Dee, 1995; Perez et al., 2015). Despite the substantial efforts to improve farms' biosecurity

and immunization through the implementation of strategies such as, sow herd air filtration, gilt

acclimation, virus elimination strategies (i.e., load-close-expose and rollover) which successfully reduced

the number of PRRSV outbreaks (Corzo et al., 2010; Alonso et al., 2012; Velasova et al., 2012; Alonso et

al., 2013; Blanchette, 2015; Silva et al., 2019), we continue to observe significant PRRSv dissemination

by indirect routes (Corzo et al., 2010; Otake et al., 2010). Therefore, it is expected that changes in

immunization protocols, such as varying vaccine brands at farms of one specific system may have a

serious impact on PRRSV spread.

Within-production systems circulation (local) versus introduction of PRRSV from commercially

unrelated farms (external)

Between-farm animal movement has historically been associated with disease spread (Augusta et al.,

2019; Moon et al., 2019; Machado et al., 2020). Here, we used structured coalescent models to infer

migration rates among commercially unrelated pig farms, calculating PRRSV dissemination not related to

between-farm pig movement. Overall, PRRSV dissemination in this region has been mostly characterized

by transmission between non-commercially related farms (external introductions) rather than within a

production system (local). In addition, we observed that during PRRSV "season" (November-May), the

external transmissions tended to be greater than during the rest of the year. While our analyses was not

aimed to explore the route of transmission events or how viral lineages first entered a population

(Rasmussen et al., 2018), they reinforce the importance of PRRSV collateral spread and open the question

of how this dissemination is happening. Indeed, a recent work (Galvis et al., 2020) showed that for sow

farms more than 50% of the between-farm transmission occurred by proximity among infected and

susceptible farms, while 20% and 53% of the nurseries and finisher farms were infect by the same route,

respectively. While little is known about this collateral PRRSV propagations, at some level the external

introduction have been associated with contaminated vehicles that service multiple production systems

(i.e., general maintenance, vaccination, cleaning or loading) (Otake et al., 2003; Dee et al., 2004), shared

equipment (Pitkin et al., 2009), non-pig animal vectors (Otake et al., 2002; Schurrer et al., 2005),

transport of semen (Christopher-Hennings et al., 1995) and airborne transmission (Lager and Mengeling,

2000; Trincado et al., 2004; Cho et al., 2007; Dee et al., 2009). At some degree, a contracted farm will

start producing pigs under a new production system, thus it is possible that a resident virus would be

introduced and become widespread within the new production system network. Knowledge gaps

regarding local dissemination of PRRSV still exist, hence, studies that consider phylodynamics coupled

with epidemiological models such as susceptible–infected–recovered to assess the relative contribution of

between-farm PRRSV transmission routes are needed. In addition, the proposed modeling should allowed

not only to answer "how it can spread" questions but also "what for", i.e. determining what effort is

required for reducing transmission (Beaunée et al., 2017).

Factors promoting or restricting PRRSV dissemination direction

In our PRRSV dissemination models, directions of lineages revealed the potential importance of two

factors that limited the spread of PRRSV. Vegetation density and high elevation, which are similar to

previous observations made by Mahesh et al., (2015), Arruda et al., (2017), and (Balka et al., 2018).

These authors also highlighted the important role that topography may play in mitigating the airborne

spread of PRRSV. In the same way, a study on foot-and-mouth disease, which also used continuous

phylogeographic analysis, identified that the presence of pure cropland or mixed land use both had a

negative impact on virus diffusion (Duchatel et al., 2019). Numerous studies have shown multiple

positive impacts of having vegetation barriers surrounding farms (i.e., trees, shrubs) over the airborne

spread of different pathogens (Malone, 2004; Patterson et al., 2008; Van Ryswyk et al., 2019), thus it

should be recognized as a possible biosecurity strategy. This study provided the first look at which

amount of healthy vegetation needed to protect farms from new introductions, consistently EVI index

between 41 and 45 showed to be sufficient. These values correspond to a dense tree coverage that is

highly similar to the values exhibited by evergreen broadleaf forest, trees within this category are live oak

(*Quercus virginiana*), laurel oak (*Quercus laurifolia*) and loblolly pine (*Pinus taeda*) (Zhang et al., 2017).

Similarly, high elevated areas have proven to act as a natural topographic barrier for the local spread of

pathogens (Jacquot et al., 2017), where the slope of the terrain was found to have an association with a

lower PRRSV incidence (Arruda et al., 2017). We were able to identify a safe range of elevation that was

consistent above 61 meter from the sea level, while this can help the local production systems, other

regions of the US would need to be analyzed separately due to the differences possible elevation ranges.

The identification of the main drivers associated with PRRSV dissemination could help in deciding about

new farm locations, especially for systems located in highly dense pig areas (Yang et al., 2019). Farms

located in areas with higher farm density were more likely to sustain new PRRSV introductions, we

highlight areas in which the volume of pigs by km^2 above 500 animals, as has been previously described

by Firkins and Weigel, (2004), Arruda et al., (2017) and Alkhamis et al., (2017, 2018). In addition, our

results identified distance to the main road as associated with increased chances for PRRSV circulation.

This has led concern by the swine industry (Reicks, 2019), since vehicle movement has demonstrated to

have a significant impact on increasing the transmission of several other diseases, for example, African

Swine Fever (ASF) (Mur et al., 2012), Classical Swine Fever (CSF) (Bronsvoort et al., 2008), and for

Foot and Mouth Disease (Muroga et al., 2013). Our findings indicate that farms located at 0.5 km to 0.7

km from major roads were more easily exposed to PRRSV, therefore this distance could be used to make

decisions about routes used for pig movement, the enhancement of on-farm biosecurity of sites within this

range of distances to major roads, and in future expansions of operations new buildings could be built

beyond 0.7 km from the main road.

Here, we showed that continuous phylogeographic reconstructions represent a useful tool to

describe and analyze the dissemination of PRRSV spread. Although the outcomes of phylodynamic

analyses can be exploited to investigate the impact of relevant factors on lineage dissemination diffusion,

it is important to note that such analyses depend on inferred viral lineage movement and thus, on the

spatial distribution of the sampled sequences (Dellicour et al., 2019). Our samples represent the

circulating PRRS viral population, as data comes from ~ 92% of pig farms in one highly dense pig region,

therefore can be directly applied in the region. Given that most commercial pig operation in North

America are similar but not the same our finding could also help the swine industry decision making on

other parts of the county, conditional to data availability this analysis can be replicated at each relevant

region or production system.

Our study was subject to a variety of limitations. The results of distance to main roads as a proxy

for the exposure to PRRSV was inconsistent thus need to be interpreted with caution, when we analyzed

all production systems together the distance of 1 km to 2 km was the only dangerous range, even though

when analysis of individual systems showed that 0.7 km was enough this result need to the interpreted

with caution. In the same why swine densities above > 1000 pigs/km^2 were not supported as conductance

while range of 500- 1000 pigs/km^2, in both cases distance to main road and swine density, one reason for

both inconsistence results could be related to the number of farm in such range intervals. For that reason,

there is an urgent need to expand the current phylodynamic framework to allow for the calculation of

point estimates of the likelihood of PRRSV spread driven by one and more predictor variables more

directly, which would also considered categorical variables allowing us to determine relative safe or

dangerous distances to main roads, for example (Dellicour et al., 2020).

Conclusion

Our study revealed significant asymmetries in the phylodynamic patterns among farms of three pig

production systems. As a result of interventions that could be related to immunization and on-farm

biosecurity, PRRSV genetic diversity has consistently declined in this region, especially in the last two

years, which could explain why 2019 has had one of the lowest counts of PRRSV cases during its typical

high season in the US (Sanhueza et al., 2019). Our results evidenced that PRRSV transmission across

each system is most likely to originate from outside production systems. These results reiterate the

importance of PRRSV farm proximity as a relevant route for virus dissemination. We highlight

characteristics of locations to which PRRSV is more easily dispersed: highly dense pig population and

proximity to main roads. On the protective side, higher elevation and more vegetation would be preferred.

Thus, reinforcement of biosecurity may be the best tool. However, it raises the question of how much and

which biosecurity practices have relevant impact on PRRSV spread. Answering this question requires

further studies capable of quantifying how biosecurity practices impede or reduce PRRSV dissemination.

Acknowledgements

We acknowledge the Department of Population Health and Pathobiology: North Carolina State University

provided startup funds for Dr. Machado and CVM for the intramural grant which supported Dr. Jara. This

work was also supported by Critical Agricultural Research and Extension 2019-68008-29910 from the

USDA National Institute of Food and Agriculture. The Morrison Swine Health Monitoring Project is a

Swine Health Information Center funded project. Authors would like to acknowledge participating

systems and veterinarians.

Authors' contributions

MJ and GM conceived the paper ideas. MJ, DR, and GM participated in the design of the study. CC

coordinated the PRRSV data collection. MJ conducted data processing and cleaning. MJ performed the

phylodynamic analysis. MJ, DR, CAC, and GM wrote and edited the manuscript. All authors discussed

the results and critically reviewed the manuscript.

Conflict of interest

All authors confirm that there are no conflicts of interest to declare

Ethical statement

The authors confirm the ethical policies of the journal, as noted on the journal's author guidelines page.

Since this work did not involve animal sampling nor questionnaire data collection by the researchers there

was no need for ethics permits.

Data Availability Statement

The data that support the findings of this study are not publicly available and are protected by confidential

agreements, therefore, are not available.

References

Alkhamis, M., A. Perez, M. Murtaugh, X. Wang, and R. Morrison, 2016: Applications of Bayesian

phylodynamic methods in a recent U.S. porcine reproductive and respiratory syndrome virus

outbreak. *Front. Microbiol.* **7**, DOI: 10.3389/fmicb.2016.00067.

Alkhamis, M.A., A.G. Arruda, R.B. Morrison, and A.M. Perez, 2017: Novel approaches for Spatial and

Molecular Surveillance of Porcine Reproductive and Respiratory Syndrome Virus (PRRSv) in the

United States. *Sci. Rep.* **7**, DOI: 10.1038/s41598-017-04628-2.

Alkhamis, M.A., A.G. Arruda, C. Vilalta, R.B. Morrison, and A.M. Perez, 2018: Surveillance of porcine

reproductive and respiratory syndrome virus in the United States using risk mapping and species

distribution modeling. *Prev. Vet. Med.* **150**, 135–142, DOI: 10.1016/j.prevetmed.2017.11.011.

Alonso, C., M.P. Murtaugh, S.A. Dee, and P.R. Davies, 2013: Epidemiological study of air filtration

systems for preventing PRRSV infection in large sow herds. *Prev. Vet. Med.* **112**, 109–117, DOI:

10.1016/j.prevetmed.2013.06.001.

Alonso, C., S. Otake, P. Davies, and S. Dee, 2012: An evaluation of interventions for reducing the risk of

PRRSV introduction to filtered farms via retrograde air movement through idle fans. *Vet. Microbiol.*

157, 304–310, DOI: 10.1016/j.vetmic.2012.01.010.

Altizer, S., R. Bartel, and B.A. Han, 2011: Animal migration and infectious disease risk. *Science (80-.).*

331, 296–302, DOI: 10.1126/science.1194694.

Arruda, A.G., J. Sanhueza, C. Corzo, and C. Vilalta, 2018: Assessment of area spread of porcine

reproductive and respiratory syndrome (PRRS) virus in three clusters of swine farms. *Transbound.*

Emerg. Dis. **65**, 1282–1289, DOI: 10.1111/tbed.12875.

Arruda, A.G., S. Tousignant, J. Sanhueza, C. Vilalta, Z. Poljak, M. Torremorell, C. Alonso, and C.A.

Corzo, 2019: Aerosol Detection and Transmission of Porcine Reproductive and Respiratory

Syndrome Virus (PRRSV): What Is the Evidence, and What Are the Knowledge Gaps? *Viruses* **11**,

712, DOI: 10.3390/v11080712.

Arruda, A.G., C. Vilalta, A. Perez, and R. Morrison, 2017: Land altitude, slope, and coverage as risk

factors for Porcine Reproductive and Respiratory Syndrome (PRRS) outbreaks in the United States.

PLoS One **12**, DOI: 10.1371/journal.pone.0172638.

Augusta, C., G.W. Taylor, and R. Deardon, 2019: Dynamic contact networks of swine movement in

Manitoba, Canada: Characterization and implications for infectious disease spread. *Transbound.*

Emerg. Dis. **66**, 1910–1919, DOI: 10.1111/tbed.13220.

Baele, G., S. Dellicour, M.A. Suchard, P. Lemey, and B. Vrancken, 2018: Recent advances in

computational phylodynamics. *Curr. Opin. Virol.* **31**, 24–32, DOI:

10.1016/J.COVIRO.2018.08.009.

Balka, G., K. Podgórska, M.S. Brar, Á. Bálint, D. Cadar, V. Celer, L. Dénes, Z. Dirbakova, A. Jedryczko,

L. Márton, D. Novosel, T. Petrović, I. Sirakov, D. Szalay, I. Toplak, F.C.C. Leung, and T. Stadejek,

2018: Genetic diversity of PRRSV 1 in Central Eastern Europe in 1994-2014: Origin and evolution

of the virus in the region. *Sci. Rep.* **8**, 1–12, DOI: 10.1038/s41598-018-26036-w.

Beaunée, G., E. Vergu, A. Joly, and P. Ezanno, 2017: Controlling bovine paratuberculosis at a regional

scale: Towards a decision modelling tool. *J. Theor. Biol.* **435**, 157–183, DOI:

10.1016/j.jtbi.2017.09.012.

Bielejec, F., G. Baele, B. Vrancken, M. Suchard, A. Rambaut, and P. Lemey, 2016: SpreaD3: interactive

visualization of spatiotemporal history and trait evolutionary processes. *Mol. Biol. Evol.* **33**, 2167–

2169.

Blanchette, A., 2015: Herding species: Biosecurity, posthuman labor, and the american industrial pig.

Cult. Anthropol. **30**, 640–669, DOI: 10.14506/ca30.4.09.

Bouckaert, R., J. Heled, D. Kühnert, T. Vaughan, C.H. Wu, D. Xie, M.A. Suchard, A. Rambaut, and A.J.

Drummond, 2014: BEAST 2: A Software Platform for Bayesian Evolutionary Analysis. (Andreas

Prlic, Ed.)*PLoS Comput. Biol.* **10**, e1003537, DOI: 10.1371/journal.pcbi.1003537.

Bronsvoort, B.M.D.C., S. Parida, I. Handel, S. McFarland, L. Fleming, P. Hamblin, and R. Kock, 2008:

Serological survey for foot-and-mouth disease virus in wildlife in eastern Africa and estimation of

test parameters of a nonstructural protein enzyme-linked immunosorbent assay for buffalo. *Clin.*

Vaccine Immunol. **15**, 1003–11, DOI: 10.1128/CVI.00409-07.

Cavanagh, D., D.A. Brian, L. Enjuanes, K. V. Holmes, M.M.C. Lai, H. Laude, S.G. Siddell, W. Spaan, F.

Taguchi, and P.J. Talbot, 1990: Recommendations of the coronavirus study group for the

nomenclature of the structural proteins, mRNAs, and genes of coronaviruses. *Virology* **176**, 306–

307, DOI: 10.1016/0042-6822(90)90259-T.

Cho, J.G., J. Deen, and S.A. Dee, 2007: Influence of isolate pathogenicity on the aerosol transmission of

Porcine reproductive and respiratory syndrome virus. *Can. J. Vet. Res.* **71**, 23–27.

Christopher-Hennings, J, E.A. Nelson, R.J. Hines, J.K. Nelson, S.L. Swenson, J.J. Zimmerman, C.L.

Chase, M.J. Yaeger, and D.A. Benfield, 1995: Persistence of porcine reproductive and respiratory

syndrome virus in serum and semen of adult boars. *J. Vet. Diagn. Invest.* **7**, 456–64, DOI:

10.1177/104063879500700406.

Christopher-Hennings, Jane, 2001: Monitoring for porcine reproductive and respiratory syndrome virus

(PRRSV) in the boar stud. *J. Swine Heal. Prod.* **9**, 186–188.

Corzo, C.A., E. Mondaca, S. Wayne, M. Torremorell, S. Dee, P. Davies, and R.B. Morrison, 2010:

Control and elimination of porcine reproductive and respiratory syndrome virus. *Virus Res.* **154**,

185–192, DOI: 10.1016/j.virusres.2010.08.016.

De Smit, A.J., A. Bouma, C. Terpstra, and J.T. Van Oirschot, 1999: Transmission of classical swine fever

virus by artificial insemination. *Vet. Microbiol.* **67**, 239–249, DOI: 10.1016/S0378-1135(99)00045-

0.

Dee, S. A., 1995: Controlling the spread of PRRS virus in the breeding herd through management of the

gilt pool. *Swine Heal. Prod.* **3**, 64–69.

Dee, S. A., S. Otake, S. Oliveira, and J. Deen, 2009: Evidence of long distance airborne transport of

porcine reproductive and respiratory syndrome virus and Mycoplasma hyopneumoniae. *Vet. Res.* **40**,

39, DOI: 10.1051/vetres/2009022.

Dee, Scott A., J. Deen, and C. Pijoan, 2004: Evaluation of 4 intervention strategies to prevent the

mechanical transmission of porcine reproductive and respiratory syndrome virus. *Can. J. Vet. Res.*

68, 19–26.

Dellicour, S., D. Desmecht, J. Paternostre, C. Malengreaux, A. Licoppe, M. Gilbert, and A. Linden, 2020:

Unravelling the dispersal dynamics and ecological drivers of the African swine fever outbreak in

Belgium. (Andrew Park, Ed.)*J. Appl. Ecol.*1365-2664.13649, DOI: 10.1111/1365-2664.13649.

Dellicour, S., P. Lemey, J. Artois, T.T. Lam, A. Fusaro, I. Monne, G. Cattoli, D. Kuznetsov, I. Xenarios,

G. Dauphin, W. Kalpravidh, S. Von Dobschuetz, F. Claes, S.H. Newman, M.A. Suchard, G. Baele,

and M. Gilbert, 2019: Incorporating heterogeneous sampling probabilities in continuous

phylogeographic inference - Application to H5N1 spread in the Mekong region. *Bioinformatics*1–7,

DOI: 10.1093/bioinformatics/btz882.

Dellicour, S., R. Rose, N.R. Faria, P. Lemey, and O.G. Pybus, 2016: SERAPHIM: Studying

environmental rasters and phylogenetically informed movements. *Bioinformatics* **32**, 3204–3206,

DOI: 10.1093/bioinformatics/btw384.

Dellicour, S., R. Rose, and O.G. Pybus, 2016: Explaining the geographic spread of emerging epidemics: a

framework for comparing viral phylogenies and environmental landscape data. *BMC Bioinformatics*

17, 82, DOI: 10.1186/s12859-016-0924-x.

Dellicour, S., C. Troupin, F. Jahanbakhsh, A. Salama, S. Massoudi, M.K. Moghaddam, G. Baele, P.

Lemey, A. Gholami, and H. Bourhy, 2019: Using phylogeographic approaches to analyse the

dispersal history, velocity and direction of viral lineages - Application to rabies virus spread in Iran.

Mol. Ecol. **28**, 4335–4350, DOI: 10.1111/mec.15222.

Dokland, T., 2010: The structural biology of PRRSV. *Virus Res.* **154**, 86–97, DOI:

10.1016/j.virusres.2010.07.029.

Drummond, A.J., S.Y.W. Ho, M.J. Phillips, and A. Rambaut, 2006: Relaxed phylogenetics and dating

with confidence. (David Penny, Ed.)*PLoS Biol.* **4**, e88, DOI: 10.1371/journal.pbio.0040088.

Duchatel, F., M. Bronsvoort, and S. Lycett, 2019: Phylogeographic analysis and identification of factors

impacting the diffusion of Foot-and-Mouth disease virus in Africa. *Front. Ecol. Evol.* **7**, 371.

Dudas, G., L.M. Carvalho, T. Bedford, A.J. Tatem, G. Baele, N.R. Faria, and F. Bielejec, 2017: Virus

genomes reveal factors that spread and sustained the Ebola epidemic. *Nature* **544**, 309–315, DOI:

10.1097/01.qai.0000532609.26906.45.

Faria, N.R., N. Vidal, J. Lourenco, J. Raghwani, K.C.E. Sigaloff, A.J. Tatem, D.A.M. Van De Vijver,

A.C. Pineda-Peña, R. Rose, C.L. Wallis, S. Ahuka-Mundeke, J.J. Muyembe-Tamfum, J. Muwonga,

M.A. Suchard, T.F.R. De Wit, R.L. Hamers, N. Ndembi, G. Baele, M. Peeters, O.G. Pybus, P.

Lemey, and S. Dellicour, 2019: Distinct rates and patterns of spread of the major HIV-1 subtypes in

Central and East Africa. *PLoS Pathog.* **15**, e1007976–e1007976, DOI:

10.1371/journal.ppat.1007976.

Firkins, L.D., and R.M. Weigel, 2004: A retrospective study of risk factors for porcine reproductive and

respiratory syndrome virus infection and clinical disease in swine herds in Illinois during the early

years of the pandemic. *J. Swine Heal. Prod.* **12**, 23–28.

Fountain-Jones, N.M., W.D. Pearse, L.E. Escobar, A. Alba-Casals, S. Carver, T.J. Davies, S. Kraberger,

M. Papeş, K. Vandegrift, K. Worsley-Tonks, and M.E. Craft, 2018: Towards an eco-phylogenetic

framework for infectious disease ecology. *Biol. Rev.* **93**, 950–970, DOI: 10.1111/brv.12380.

Franzo, G., G. Dotto, M. Cecchinato, D. Pasotto, M. Martini, and M. Drigo, 2015: Phylodynamic analysis

of porcine reproductive and respiratory syndrome virus (PRRSV) in Italy: Action of selective

pressures and interactions between different clades. *Infect. Genet. Evol.* **31**, 149–157, DOI:

10.1016/j.meegid.2015.01.026.

Galvis, J.A., J. Prada, C. Jones, and G. Machado, 2020: Transmission and control strategies for porcine

reproductive and respiratory syndrome virus. *bioRxiv*2020.05.23.112946, DOI:

10.1101/2020.05.23.112946.

Gill, M.S., P. Lemey, S.N. Bennett, R. Biek, and M.A. Suchard, 2016: Understanding Past Population

Dynamics: Bayesian Coalescent-Based Modeling with Covariates. *Syst. Biol.* **65**, 1041–1056, DOI:

10.1093/sysbio/syw050.

Guindon, S., J.F. Dufayard, V. Lefort, M. Anisimova, W. Hordijk, and O. Gascuel, 2010: New algorithms

and methods to estimate maximum-likelihood phylogenies: Assessing the performance of PhyML

3.0. *Syst. Biol.* **59**, 307–321, DOI: 10.1093/sysbio/syq010.

Hasegawa, M., H. Kishino, and T. Yano, 1985: Dating of the human-ape splitting by a molecular clock of

mitochondrial DNA. *J. Mol. Evol.* **22**, 160–174, DOI: 10.1007/BF02101694.

Hicks, J.T., D.H. Lee, V.R. Duvuuri, M.K. Torchetti, D.E. Swayne, and J. Bahl, 2020: Agricultural and

geographic factors shaped the North American 2015 highly pathogenic avian influenza H5N2

outbreak. *PLoS Pathog.* **16**, e1007857, DOI: 10.1371/journal.ppat.1007857.

Hill, V., and G. Baele, 2019: Bayesian estimation of past population dynamics in BEAST 1.10 using the

Skygrid coalescent model. *Mol. Biol. Evol.* **36**, 2620–2628.

Hogg, R. V, E.A. Tanis, D.L. Zimmerman, B. Columbus, I. New, Y. San, F. Upper, S. River, A. Cape, T.

Dubai, L. Madrid, M. Munich, P. Montreal, T. Delhi, M.C. São, P. Sydney, H. Kong, S. Singapore,

and T. Tokyo, 2010: Probability and Statistical Inference. Pearson/Prentice Hall.

Holtkamp, D.J., J.B. Kliebenstein, E. Neumann, J.J. Zimmerman, H. Rotto, T.K. Yoder, C. Wang, P.

Yeske, C.L. Mowrer, C.A. Haley, E.J. Neumann, H.F. Rotto, and P.E. Yeske, 2013: Assessment of

the economic impact of porcine reproductive and respiratory syndrome virus on United States pork

producers Assessment of the economic impact of porcine reproductive and respiratory syndrome

virus on United States pork producers Assessment of. *J. Swine Heal. Prod.* **21**.

Jacquot, M., K. Nomikou, M. Palmarini, P. Mertens, and R. Biek, 2017: Bluetongue virus spread in

Europe is a consequence of climatic, landscape and vertebrate host factors as revealed by

phylogeographic inference. *Proc. R. Soc. B Biol. Sci.* **284**, DOI: 10.1098/rspb.2017.0919.

Jeong, J., S. Kim, C. Park, I. Kang, K.H. Park, H.J. Ham, and C. Chae, 2018: Effect of vaccination with a

porcine reproductive and respiratory syndrome subunit vaccine on sow reproductive performance in

endemic farms. *Vet. Rec.* **182**, 602, DOI: 10.1136/vr.104547.

Kalyaanamoorthy, S., B.Q. Minh, T.K.F. Wong, A. Von Haeseler, and L.S. Jermiin, 2017: ModelFinder:

Fast model selection for accurate phylogenetic estimates. *Nat. Methods* **14**, 587–589, DOI:

10.1038/nmeth.4285.

Kass, R.E., and A.E. Raftery, 1995: Bayes Factors. *J. Am. Stat. Assoc.* **90**, 773–795, DOI:

10.1080/01621459.1995.10476572.

Kumar, S., M. Nei, J. Dudley, and K. Tamura, 2008: MEGA: A biologist-centric software for

evolutionary analysis of DNA and protein sequences. *Brief. Bioinform.* **9**, 299–306, DOI:

10.1093/bib/bbn017.

Kumar, Sudhir, G. Stecher, M. Li, C. Knyaz, and K. Tamura, 2018: MEGA X: Molecular Evolutionary

Genetics Analysis across Computing Platforms | Molecular Biology and Evolution | Oxford

Academic. *Mol. Biol. Evol.* **35**, 1547–1549.

Laenen, L., S. Dellicour, V. Vergote, I. Nauwelaers, S. De Coster, I. Verbeeck, B. Vanmechelen, P.

Lemey, and P. Maes, 2016: Spatio-temporal analysis of Nova virus, a divergent hantavirus

circulating in the European mole in Belgium. *Mol. Ecol.* **25**, 5994–6008, DOI: 10.1111/mec.13887.

Lager, K.M., and W.L. Mengeling, 2000: Experimental aerosol transmission of pseudorabies virus and

porcine reproductive and respiratory syndrome virus. *Proc. Am. Assoc. Swine Pract.*409–410.

Lemey, P., A. Rambaut, A.J. Drummond, and M.A. Suchard, 2009: Bayesian Phylogeography Finds Its

Roots. (Christophe Fraser, Ed.)*PLoS Comput. Biol.* **5**, e1000520, DOI:

10.1371/journal.pcbi.1000520.

Linhares, D.C.L., J.P. Cano, M. Torremorell, and R.B. Morrison, 2014: Comparison of time to PRRSv-

stability and production losses between two exposure programs to control PRRSv in sow herds.

Prev. Vet. Med. **116**, 111–119, DOI: 10.1016/j.prevetmed.2014.05.010.

Lycett, S.J., V.N. Tanya, M. Hall, D. King, S. Mazeri, V. Mioulet, N. Knowles, J. Wadsworth, K.

Bachanek-Bankowska, N.N. Victor, K.L. Morgan, and M. Bronsvoort, 2019: The evolution and

phylodynamics of serotype A and SAT2 foot-and-mouth disease viruses in endemic regions of

Africa. *Sci. Rep.* **9**, 5614, DOI: 10.1101/572198.

Machado, G., J.A. Galvis, F.P.N. Lopes, J. Voges, A.A.R. Medeiros, and N.C. Cárdenas, 2020:

Quantifying the dynamics of pig movements improves targeted disease surveillance and control

plans. *bioRxiv*2019.12.29.886408, DOI: 10.1101/2019.12.29.886408.

Mahesh, K.C., J. Bhoj Raj, S. Swoyam, Prakash, M. Prajapati, D. Kathayat, and S. Dhakal, 2015: Sero-

Prevalence of Porcine Reproductive and Respiratory Syndrome (PRRS) in Pigs of Different

Developmental Regions of Nepal. *Int. J. Appl. Sci. Biotechnol.* **3**, 218–222, DOI:
10.3126/ijasbt.v3i2.12539.

Malone, B., 2004: Using trees to reduce dust and odor emissions from poultry farms*Poult. Inf. Exch.*

Martin, D.P., B. Murrell, M. Golden, A. Khoosal, and B. Muhire, 2015: RDP4: Detection and analysis of
recombination patterns in virus genomes. *Virus Evol.* **1**, DOI: 10.1093/ve/vev003.

Moon, S.A., T. Ferdousi, A. Self, and C.M. Scoglio, 2019: Estimation of swine movement network at
farm level in the US from the Census of Agriculture data. *Sci. Rep.* **9**, DOI: 10.1038/s41598-019-
42616-w.

MSHMP, 2020: PRRS charts [Online] Available at https://www.vetmed.umn.edu/centers-
programs/swine-program/outreach-leman-mshmp/mshmp/mshmp-prrs-figures (accessed January 10,
2020).

Müller, N.F., D. Rasmussen, and T. Stadler, 2018: MASCOT: Parameter and state inference under the
marginal structured coalescent approximation. *Bioinformatics* **34**, 3843–3848, DOI:
10.1093/bioinformatics/bty406.

Müller, N.F., D.A. Rasmussen, and T. Stadler, 2017: The structured coalescent and its approximations.
Mol. Biol. Evol. **34**, 2970–2981, DOI: 10.1093/molbev/msx186.

Mur, L., B. Martínez-López, and J.M. Sánchez-Vizcaíno, 2012: Risk of African swine fever introduction
into the European Union through transport-associated routes: returning trucks and waste from
international ships and planes. *BMC Vet. Res.* **8**, 149, DOI: 10.1186/1746-6148-8-149.

Muroga, N., S. Kobayashi, T. Nishida, Y. Hayama, T. Kawano, T. Yamamoto, and T. Tsutsui, 2013: Risk
factors for the transmission of foot-and-mouth disease during the 2010 outbreak in Japan: a case--
control study. *BMC Vet. Res.* **9**, 150, DOI: 10.1186/1746-6148-9-150.

Murray, G.G.R., F. Wang, E.M. Harrison, G.K. Paterson, A.E. Mather, S.R. Harris, M.A. Holmes, A.
Rambaut, and J.J. Welch, 2016: The effect of genetic structure on molecular dating and tests for
temporal signal. *Methods Ecol. Evol.* **7**, 80–89, DOI: 10.1111/2041-210X.12466.

Navascués, M., F. Depaulis, and B. Emerson, 2010: Combining contemporary and ancient DNA in

population genetic and phylogeographical studies. *Mol. Ecol. Resour.* **10**, 760–772, DOI:

10.1111/j.1755-0998.2010.02895.x.

Neumann, E.J., J.B. Kliebenstein, C.D. Johnson, J.W. Mabry, E.J. Bush, A.H. Seitzinger, A.L. Green, and

J.J. Zimmerman, 2005: Assessment of the economic impact of porcine reproductive and respiratory

syndrome on swine production in the United States. *J. Am. Vet. Med. Assoc.* **227**, 385–392, DOI:

10.2460/javma.2005.227.385.

Nguyen, L.T., H.A. Schmidt, A. Von Haeseler, and B.Q. Minh, 2015: IQ-TREE: A fast and effective

stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol. Biol. Evol.* **32**, 268–274,

DOI: 10.1093/molbev/msu300.

Otake, S., S. Dee, K. Rossow, R. Moon, and C. Pijoan, 2002: Mechanical transmission of porcine

reproductive and respiratory syndrome virus by mosquitoes, Aedes vexans (Meigen). *Can. J. Vet.*

Res. **66**, 191–195.

Otake, Satoshi., S. Dee, C. Corzo, S. Oliveira, and J. Deen, 2010: Long-distance airborne transport of

infectious PRRSV and Mycoplasma hyopneumoniae from a swine population infected with multiple

viral variants. *Vet. Microbiol.* **145**, 198–208, DOI: 10.1016/j.vetmic.2010.03.028.

Otake, Satoshi, S.. Otake, S., R.D. Dee, K.D. Moon, C. Roscow, 265–270. Trincado Pijoan, 2003:

Evaluation of mosquitoes, Aedes vexans, as biological vectors of porcine reproductive and

respiratory syndrome virus. Can. J. Vet. Res. 67, S.A. Dee, R.D. Moon, K.D. Roscow, C. Trincado,

and C. Pijoan, 2003: Evaluation of mosquitoes, Aedes vexans, as biological vectors of porcine

reproductive and respiratory syndrome virus. *Can. J. Vet. Res.* **67**, 265–270.

Passafaro, T.L., A.F.A. Fernandes, B.D. Valente, N.H. Williams, and G.J.M. Rosa, 2020: Network

analysis of swine movements in a multi-site pig production system in Iowa, USA. *Prev. Vet. Med.*

174, DOI: 10.1016/j.prevetmed.2019.104856.

Patterson, P.H., A. Adrizal, R.M. Hulet, R.M. Bates, C.A.B. Myers, G.P. Martin, R.L. Shockey, M. Van

Der Grinten, D.A. Anderson, and J.R. Thompson, 2008: Vegetative buffers for fan emissions from

poultry farms: 2. Ammonia, dust and foliar nitrogen. *J. Environ. Sci. Heal. - Part B Pestic. Food*

Contam. Agric. Wastes **43**, 96–103, DOI: 10.1080/03601230701735078.

Perez, A.M., P.R. Davies, C.K. Goodell, D.J. Holtkamp, E. Mondaca-Fernández, Z. Poljak, S.J.

Tousignant, P. Valdes-Donoso, J.J. Zimmerman, and R.B. Morrison, 2015: Lessons learned and

knowledge gaps about the epidemiology and control of porcine reproductive and respiratory

syndrome virus in North America. *J. Am. Vet. Med. Assoc.* **246**, 1304–1317, DOI:

10.2460/javma.246.12.1304.

Pileri, E., and E. Mateu, 2016: Review on the transmission porcine reproductive and respiratory syndrome

virus between pigs and farms and impact on vaccination. *Vet. Res.* **47**, 1–13, DOI: 10.1186/s13567-

016-0391-4.

Pitkin, A., J. Deen, and S. Dee, 2009: Use of a production region model to assess the airborne spread of

porcine reproductive and respiratory syndrome virus. *Vet. Microbiol.* **136**, 1–7, DOI:

10.1016/j.vetmic.2008.10.013.

Plain, R.L., and J.D. Laurence, 2003: Swine production. *Vet. Clin. North Am. Food Anim. Pract.* **19**, 319–

337.

Pybus, O., M. Suchard, P. Lemey, F. Bernardin, A. Rambaut, F. Crawford, R. Gray, N. Arinaminpathy, S.

Stramer, M. Busch, and E. Delwart, 2012: Unifying the spatial epidemiology and molecular

evolution of emerging epidemics. *Natl. Acad Sci.* 1–6, DOI: 10.1073/pnas.1206598109/-

/DCSupplemental.

Pybus, O.G., and A. Rambaut, 2009: Evolutionary analysis of the dynamics of viral infectious disease.

Nat. Rev. Genet. **10**, 540–550, DOI: 10.1038/nrg2583.

Rambaut, A., and A. Drummond, 2016: TreeAnnotator v.2.4. 3. . Part of the BEAST package.

Rambaut, A., T. Lam, L. Max Carvalho, and O. Pybus, 2016: Exploring the temporal structure of

heterochronous sequences using TempEst (formerly Path-O-Gen). *Virus Evol.* **2**, vew007, DOI:

10.1093/ve/vew007.

Rappole, J.H., B.W. Compton, P. Leimgruber, J. Robertson, D.I. King, and S.C. Renner, 2006: Modeling

movement of West Nile virus in the Western Hemisphere. *Vector-Borne Zoonotic Dis.* **6**, 128–139,

779 DOI: 10.1089/vbz.2006.6.128.

780 Rasmussen, D.A., E. Wilkinson, A. Vandormael, F. Tanser, D. Pillay, T. Stadler, and T. de Oliveira,

781  2018: Tracking external introductions of HIV using phylodynamics reveals a major source of

782  infections in rural KwaZulu-Natal, South Africa. *Virus Evol.* **4**, vey037, DOI: 10.1093/ve/vey037.

783 Rathkjen, P.H., and J. Dall, 2017: Control and eradication of porcine reproductive and respiratory

784  syndrome virus type 2 using a modified-live type 2 vaccine in combination with a load, close,

785  homogenise model: An area elimination study. *Acta Vet. Scand.* **59**, 4, DOI: 10.1186/s13028-016-

786  0270-z.

787 Reicks, D.L., 2019: Effective biosecurity to protect North American studs and clients from emerging

788  infectious disease. *Theriogenology* **137**, 82–87, DOI: 10.1016/j.theriogenology.2019.05.041.

789 Sanhueza, J.M., C. Vilalta, M. Kikuti, E. Geary, and C. Corzo, 2019: Comparison of PRRS incidence rate

790  in status IV breeding herds during 2009-2019.

791 Sanhueza, Juan M., C. Vilalta, C. Corzo, and A.G. Arruda, 2019: Factors affecting Porcine Reproductive

792  and Respiratory Syndrome virus time-to-stability in breeding herds in the Midwestern United States.

793  *Transbound. Emerg. Dis.* **66**, 823–830, DOI: 10.1111/tbed.13091.

794 Schurrer, J., S. Dee, R. Moon, M. Murthaugh, C. Finnegan, J. Deen, S. Kleiboeker, and C. Pijoan, 2005:

795  Retention of ingested porcine reproductive and respiratory syndrome virus in houseflies. *Am Vet*

796  *Med Assoc* **66**, 1517–1525, DOI: 10.2460/ajvr.2005.66.1517.

797 Shi, M., T. Tsan-Yuk Lam, C.-C. Hon, R. Kin-Hei Hui, K.S. Faaberg, T. Wennblom, M.P. Murtaugh, T.

798  Stadejek, and F. Chi-Ching Leung, 2010: Molecular epidemiology of PRRSV: A phylogenetic

799  perspective. *Virus Res.* **154**, 7–17, DOI: 10.1016/j.virusres.2010.08.014.

800 Silva, G.S., G. Machado, K.L. Baker, D.J. Holtkamp, and D.C.L. Linhares, 2019: Machine-learning

801  algorithms to identify key biosecurity practices and factors associated with breeding herds reporting

802  PRRS outbreak. *Prev. Vet. Med.* **171**, DOI: 10.1016/j.prevetmed.2019.104749.

803 Sørensen, R., U. Zinko, and J. Seibert, 2005: On the calculation of the topographic wetness index:

804  evaluation of different methods based on field observations On the calculation of the topographic

805 wetness index: evaluation of different methods based on field observations Evaluation of TWI

806 calculation Vol. 2*Hydrol. Earth Sys. Sci. Discuss*.

807 Sun, R.Q., R.J. Cai, Y.Q. Chen, P.S. Liang, D.K. Chen, and C.X. Song, 2012: Outbreak of porcine

808 epidemic diarrhea in suckling piglets, China. *Emerg. Infect. Dis.* **18**, 161–163, DOI:

809 10.3201/eid1801.111259.

810 Sun, Y.K., X.L. Han, Y.F. Wei, Z.Q. Yu, C.H. Ji, Q. Li, G. Lu, L. Shen, C.Q. Ma, H. Wang, and G.H.

811 Zhang, 2019: Phylogeography, phylodynamics and the recent outbreak of lineage 3 porcine

812 reproductive and respiratory syndrome viruses in China. *Transbound. Emerg. Dis.* **66**, 2152–2162,

813 DOI: 10.1111/tbed.13269.

814 Tavaré, S., 1986: Some probabilistic and statistical problems in the analysis of dna sequences. *Lect Math*

815 *Life Sci.* **17**, 57–86.

816 To, T.H., M. Jung, S. Lycett, and O. Gascuel, 2016: Fast Dating Using Least-Squares Criteria and

817 Algorithms. *Syst. Biol.* **65**, 82–97, DOI: 10.1093/sysbio/syv068.

818 Trincado, C., S. Dee, L. Jacobson, S. Otake, K. Rossow, and C. Pijoan, 2004: Attempts to transmit

819 porcine reproductive and respiratory syndrome virus by aerosols under controlled field conditions.

820 *Vet. Rec.* **154**, 294–297, DOI: 10.1136/vr.154.10.294.

821 Trovão, N.S., M.A. Suchard, G. Baele, M. Gilbert, and P. Lemey, 2015: Bayesian inference reveals host-

822 specific contributions to the epidemic expansion of influenza A H5N1. *Mol. Biol. Evol.* **32**, 3264–

823 3275, DOI: 10.1093/molbev/msv185.

824 Van Ryswyk, K., N. Prince, M. Ahmed, E. Brisson, J.D. Miller, and P.J. Villeneuve, 2019: Does urban

825 vegetation reduce temperature and air pollution concentrations? Findings from an environmental

826 monitoring study of the Central Experimental Farm in Ottawa, Canada. *Atmos. Environ.* **218**, DOI:

827 10.1016/j.atmosenv.2019.116886.

828 Velasova, M., P. Alarcon, S. Williamson, and B. Wieland, 2012: Risk factors for porcine reproductive

829 and respiratory syndrome virus infection and resulting challenges for effective disease surveillance.

830 *BMC Vet. Res.* **8**, 184, DOI: 10.1186/1746-6148-8-184.

Volz, E.M., K. Koelle, and T. Bedford, 2013: Viral Phylodynamics. *PLoS Comput. Biol.* **9**, DOI:

10.1371/journal.pcbi.1002947.

Yang, J., N.F. Müller, R. Bouckaert, B. Xu, and A.J. Drummond, 2019: Bayesian phylodynamics of avian

influenza A virus H9N2 in Asia with time-dependent predictors of migration. *PLoS Comput. Biol.*

15, DOI: 10.1371/journal.pcbi.1007189.

Yang, Z., 1994: Maximum likelihood phylogenetic estimation from DNA sequences with variable rates

over sites: Approximate methods. *J. Mol. Evol.* **39**, 306–314, DOI: 10.1007/BF00160154.

Yuan, S., C.J. Nelsen, M.P. Murtaugh, B.J. Schmitt, and K.S. Faaberg, 1999: Recombination between

North American strains of porcine reproductive and respiratory syndrome virus. *Virus Res.* **61**, 87–

98, DOI: 10.1016/S0168-1702(99)00029-5.

Zhang, X., S. Wu, X. Yan, and Z. Chen, 2017: A global classification of vegetation based on NDVI,

rainfall and temperature. *Int. J. Climatol.* **37**, 2318–2324, DOI: 10.1002/joc.4847.

Table list

Table 1. PRRSV dissemination between systems. Bayes factor results represent the level of support in the

rates of the spread between systems, where BF > 20 is strong support, BF = 3-19 is positive support and

BF< 3 is negligible.

Origin	Destination	Average relative migration rate	Bayes factor	Interpretation
A	B	1.79 ± 0.92	102.1	Strong support
A	C	0.46 ± 0.39	47.8	Strong support
B	A	0.35 ± 0.37	27.4	Strong support
B	C	0.33 ± 0.13	7.9	Positive support
C	A	0.41 ± 0.22	14.7	Positive support
C	B	0.39 ± 0.16	10.3	Positive support

849   **Table 2.** PRRSV dissemination between types of farms. Bayes factor results represent the level of support

850   in the rates of the spread between types of farms, where BF > 20 is strong support, BF = 3-19 is positive

851   support and BF< 3 is negligible.

Farm type of origin	Farm type of destination	All systems (Bayes factor)	System A (Bayes factor)	System B (Bayes factor)	System C (Bayes factor)
Sow	Nursery	109	56.2	42.2	51.7
Sow	Finisher	80.5	36.6	28.1	17.9
Sow	Boar stud	0.1	2.2	0.6	1.1
Nursery	Sow	7.3	5.8	4.9	4.6
Nursery	Finisher	27.5	33.7	10.9	16.6
Nursery	Boar stud	1.9	0.1	0.7	1.3
Finisher	Sow	18.9	10.9	14.1	5.3
Finisher	Nursery	12	16.7	13.2	6.4
Finisher	Boar stud	0.2	1.4	0.1	0.6
Boar stud	Sow	0.5	1.1	0.3	0.1
Boar stud	Nursery	3.9	2.5	2.1	2.4
Boar stud	Finisher	0.4	2	0.8	1.5

853 **Table 3.** Number of transmission events that have been originated in farms that belong to the same pig

854 system (local) compared to the events coming from farms integrated into another pig system (external).

System	Year	Local		External		Total
		Events	%	Events	%	
A	2014	349	25.9%	996	74.1%	1345
	2015	294	25.9%	840	74.1%	1134
	2016	292	25.8%	839	74.2%	1131
	2017	330	28.3%	836	71.7%	1166
	2018	109	18.0%	498	82.0%	607
B	2014	299	22.1%	1055	77.9%	1354
	2015	310	27.5%	819	72.5%	1129
	2016	262	23.1%	871	76.9%	1133
	2017	205	17.6%	963	82.4%	1168
	2018	117	18.5%	517	81.5%	634
C	2014	228	16.9%	1125	83.1%	1353
	2015	60	5.3%	1075	94.7%	1135
	2016	126	11.2%	1002	88.8%	1128
	2017	80	6.8%	1088	93.2%	1168
	2018	54	8.9%	551	91.1%	605

859 **Figure legend**

861 **Figure 1.** Diagram of factors promoting or restricting PRRSV spread. Arrows represent the relationship

862 between the predictor and PRRSV occurrence based on Bayes factor analysis, where green and orange

863 lines represent positive and negative drivers respectively. Black represent negligible support, and dashed

864 lines show relatedness between environmental predictors affecting PRRSV through other variables.

Figure 2. Dissemination history of PRRSV, inferred by discrete phylogeographic analysis. A) Maximum

clade credibility phylogeny, colored according to pig production systems (A, B, C). The probabilities of

ancestral states (inferred from the Bayesian discrete trait analysis) are shown in pie charts on each node,

representing the most likely type of farms. B) Spatiotemporal patterns in the relative genetic diversity

represented through the Bayesian SkyGrid plot and the number of sequences per month. The left y-axis

summarizes the effective population size over time, the mean estimate is represented by the dark blue

line, while the shaded light blue regions correspond to the 95% highest posterior density (HPD), and in

the right y-axis shows the number of sampled sequences over time.

Figure 3. PRRSV dissemination within system A. Panel A) Maximum clade credibility phylogeny

colored according to the different farm types (sow, nursery, finisher, and boar stud). Probabilities of

ancestral states shown in pie charts in nodes, representing the most likely type of farm origin. Panel B)

Spatiotemporal patterns in the relative genetic variation through Bayesian SkyGrid plot and number of

sequences per month. The left y-axis summarizes the effective population size over time. Mean estimate

is represented by the dark blue line. Shaded light blue regions correspond to the 95% highest posterior

density (HPD), and in the right y-axis shows the number of sampled sequences over time.

Figure 4. PRRSV dissemination within system B. Panel A) Maximum clade credibility phylogeny

colored according to the different farm types (sow, nursery, finisher, and boar stud). Probabilities of

ancestral states shown in pie charts in nodes, representing the most likely type of farm origin. Panel B)

Spatiotemporal patterns in the relative genetic variation through Bayesian SkyGrid plot and number of

sequences per month. The left y-axis summarizes the effective population size over time. Mean estimate

is represented by the dark blue line. Shaded light blue regions correspond to the 95% highest posterior

density (HPD), and in the right y-axis shows the number of sampled sequences over time.

Figure 5. PRRSV dissemination within system C. Panel A) Maximum clade credibility phylogeny

colored according to the different farm types (sow, nursery, finisher, and boar stud). Probabilities of

ancestral states shown in pie charts in nodes, representing the most likely type of farm origin. Panel B)

Spatiotemporal patterns in the relative genetic variation through Bayesian SkyGrid plot and number of

sequences per month. The left y-axis summarizes the effective population size over time. Mean estimate

is represented by the dark blue line. Shaded light blue regions correspond to the 95% highest posterior

density (HPD), and in the right y-axis shows the number of sampled sequences over time.

Figure 6. PRRSV migration events among pig systems are inferred through a structured coalescent

model. Number of dissemination events originated in farms from the same pig system compared to events

from farms integrated into another pig system. Grey background represents the months where the

exponentially weighted moving average epidemic monitor surpassed the established PRRSv epidemic

threshold based on the February 2020 MSHMP report (MSHMP, 2020), while dashed lines represent the

beginning of each year.