Cytoplasmic mRNA decay represses RNA polymerase II transcription during early

apoptosis

Christopher Duncan-Lewis[1], Ella Hartenian[1], Valeria King[1], and Britt A. Glaunsinger*[1,2,3]

[1] Department of Molecular and Cell Biology; University of California, Berkeley; Berkeley,

CA 94720, USA

[2]Department of Plant and Microbial Biology; University of California, Berkeley; Berkeley,

CA 94720, USA

[3]Howard Hughes Medical Institute, Berkeley, CA 94720, USA

*Correspondence: glaunsinger@berkeley.edu

ABSTRACT

RNA abundance is generally sensitive to perturbations in decay and synthesis rates, but crosstalk between RNA polymerase II transcription and cytoplasmic mRNA degradation often leads to compensatory changes in gene expression. Here, we reveal that widespread mRNA decay during early apoptosis represses RNAPII transcription, indicative of positive (rather than compensatory) feedback. This repression requires active cytoplasmic mRNA degradation, which leads to impaired recruitment of components of the transcription preinitiation complex to promoter DNA. Importin α/β-mediated nuclear import is critical for this feedback signaling, suggesting that proteins translocating between the cytoplasm and nucleus connect mRNA decay to transcription. We also show that an analogous pathway activated by viral nucleases similarly depends on nuclear protein import. Collectively, these data demonstrate that accelerated mRNA decay leads to the repression of mRNA transcription, thereby amplifying the shutdown of gene expression. This highlights a conserved gene regulatory mechanism by which cells respond to threats.

IMPACT STATEMENT: Human cells respond to cytoplasmic mRNA depletion during early apoptosis by inhibiting RNA polymerase II transcription, thereby magnifying the gene expression shutdown during stress.

INTRODUCTION

Gene expression is often depicted as a unidirectional flow of discrete stages: DNA is first transcribed by RNA polymerase II (RNAPII) into messenger RNA (mRNA), which is processed and exported to the cytoplasm where it is translated and then degraded. However, there is a growing body of work that reveals complex cross talk

between the seemingly distal steps of transcription and decay. For example, the yeast

Ccr4-Not deadenylase complex, which instigates basal mRNA decay by removing the

poly(A) tails of mRNAs (Tucker et al., 2001), was originally characterized as a

transcriptional regulator (Collart & Stmhp, 1994; Denis, 1984). Other components of

transcription such as RNAPII subunits and gene promoter elements have been linked to

cytoplasmic mRNA decay (Bregman et al., 2011; Lotan et al., 2005; Lotan, Goler-Baron,

Duek, Haimovich, & Choder, 2007), while the activity of cytoplasmic mRNA degradation

machinery such as the cytoplasmic 5'-3' RNA exonuclease Xrn1 can influence the

transcriptional response (Haimovich et al., 2013; Sun et al., 2012).

The above findings collectively support a model in which cells engage a buffering

response to reduce transcription when mRNA decay is slowed, or reduce mRNA decay

when transcription is slowed to preserve the steady state mRNA pool (Haimovich et al.,

2013; Hartenian & Glaunsinger, 2019). While much of this research has been performed

in yeast, the buffering model is also supported by studies in mouse and human cells

(Helenius et al., 2011; Singh et al., 2019). In addition to bulk changes to the mRNA

pool, compensatory responses can also occur at the individual gene level to buffer

against aberrant transcript degradation. Termed "nonsense-induced transcription

compensation" (NITC; Wilkinson, 2019), this occurs when nonsense-mediated mRNA

decay leads to transcriptional upregulation of genes with some sequence homology to

the aberrant transcript (El-Brolosy et al., 2019; Ma et al., 2019).

A theme that unites much of the research linking mRNA decay to transcription is

homeostasis; perturbations in mRNA stability are met with an opposite transcriptional

response in order to maintain stable mRNA transcript levels. However, there are cellular

contexts in which homeostasis is not beneficial, for example during viral infection. Many

viruses induce widespread host mRNA decay (Narayanan & Makino, 2013) and co-opt

the host transcriptional machinery (Harwig, Landick, & Berkhout, 2017) in order to

express viral genes. Indeed, infection with mRNA decay-inducing herpesviruses or

expression of broad-acting viral ribonucleases in mammalian cells causes RNAPII

transcriptional repression in a manner linked to accelerated mRNA decay (Abernathy,

Gilbertson, Alla, & Glaunsinger, 2015; Hartenian, Gilbertson, Federspiel, Cristea, &

Glaunsinger, 2020). It is possible that this type of positive feedback represents a

protective cellular shutdown response, perhaps akin to the translational shutdown

mechanisms that occur upon pathogen sensing (D. Walsh, Mathews, & Mohr, 2013). A

central question, however, is whether transcriptional inhibition upon mRNA decay is

restricted to infection contexts, or whether it is also engaged upon other types of stimuli.

The best-defined stimulus known to broadly accelerate cytoplasmic mRNA

decay outside of viral infection is induction of apoptosis. Overall levels of poly(A) RNA

are reduced rapidly after the induction of extrinsic apoptosis via accelerated degradation

from the 3' ends of transcripts (Thomas et al., 2015). The onset of accelerated mRNA

decay occurs coincidentally with mitochondrial outer membrane depolarization (MOMP)

and requires release of the mitochondrial exonuclease polyribonucleotide

nucleotidyltransferase 1 (PNPT1) into the cytoplasm. PNPT1 then coordinates with

other 3' end decay machinery such as DIS3L2 and terminal uridylyltransferases

(TUTases; Liu et al., 2018; Thomas et al., 2015). Notably, mRNA decay occurs before

other hallmarks of apoptosis including phosphatidylserine (PS) externalization and DNA

fragmentation, but likely potentiates apoptosis by reducing the expression of unstable

anti-apoptotic proteins such as MCL1 (Thomas et al., 2015).

Here, we used early apoptosis as a model to study the impact of accelerated

cytoplasmic mRNA decay on transcription. We reveal that under conditions of increased

mRNA decay, there is a coincident decrease in RNAPII transcription, indicative of

positive feedback between mRNA synthesis and degradation. Using decay factor

depletion experiments, we demonstrate that mRNA decay is required for the

transcriptional decrease and further show that transcriptional repression is associated

with reduced RNAPII polymerase occupancy on promoters. This phenotype requires

ongoing nuclear-cytoplasmic protein transport, suggesting that protein trafficking may

provide the signal linking cytoplasmic decay to transcription. Collectively, our findings

elucidate a distinct gene regulatory mechanism by which cells sense and respond to

threats.

RESULTS

mRNA decay during early apoptosis is accompanied by reduced synthesis of

RNAPII transcripts

To induce widespread cytoplasmic mRNA decay, we initiated rapid extrinsic

apoptosis in HCT116 colon carcinoma cells by treating them with TNF-related apoptosis

inducing ligand (TRAIL). TRAIL treatment causes a well-characterized progression of

apoptotic events including caspase cleavage and mitochondrial outer membrane

permeabilization or "MOMP" (Albeck et al., 2008; Thomas et al., 2015). It is MOMP that

stimulates mRNA decay in response to an apoptosis-inducing ligand (Figure 1A), partly

by releasing the mitochondrial 3'-5' RNA exonuclease PNPT1 into the cytoplasm (Liu et

al., 2018; Thomas et al., 2015). A time course experiment in which cells were treated

with 100 ng/mL TRAIL for increasing 30 min increments showed activation of caspase 8

(CASP8) and caspase 3 (CASP3) by 1.5 hr (Figure 1B), as measured by disappearance

of the full-length zymogen upon cleavage (Kim et al., 2000; Thomas et al., 2015). In

agreement with Liu et al. (2018), RT-qPCR performed on total RNA revealed a

coincident decrease in the mRNA levels of several housekeeping genes (*ACTB*,

GAPDH, *EEF1A*, *PPIA*, *CHMP2A*, *DDX6*, *RPB2*, *and RPLP0*) beginning 1.5 hr after

TRAIL was applied (Figure 1C, Supplementary Figure 1A). Fold changes were

calculated in reference to 18S rRNA, which has been shown to be stable during early

apoptosis (Houge et al., 1995; Thomas et al., 2015). As expected, this decrease was

specific to RNAPII transcripts, as the RNA polymerase III (RNAPIII)-transcribed ncRNAs

U6, 7SK, 7SL, and 5S did not show a similarly progressive decrease. The *U6* transcript

was instead upregulated, possibly suggesting post-transcriptional regulation of *U6* as

alluded to in a previous study (Noonberg, Scott, & Benz, 1996). These data confirm that

mRNA depletion occurs by 1.5-2 hr during TRAIL-induced apoptosis.

To monitor whether apoptosis also influenced transcription, we pulse labeled the

cells with 4-thiouridine (4sU) for 20 min at the end of each TRAIL treatment. 4sU is

incorporated into actively transcribing RNA and can be subsequently coupled to HPDP-

biotin and purified over streptavidin beads, then quantified by RT-qPCR to measure

nascent transcript levels (Dölken, 2013). 4sU-labeled RNA levels were also normalized

to 18S rRNA, which was produced at a constant level in the presence and absence of

TRAIL (Supplementary Figure 1B). In addition to a reduction in steady state mRNA

abundance, TRAIL treatment caused a decrease in RNAPII-driven mRNA production,

while RNAPIII transcription was largely either unaffected or enhanced (Figure 1D,

Supplementary Figure 1C). Thus, TRAIL triggers mRNA decay and decreases nascent

mRNA production in HCT116 cells but does not negatively impact RNAPIII transcript

abundance or production.

RNAPII transcription is globally repressed during early apoptosis

z-VAD-fmk (zVAD), a pan-caspase inhibitor, was used to confirm that TRAIL-

induced mRNA decay and transcriptional arrest were associated with apoptosis and not

due to an off-target effect of TRAIL. HCT116 cells were pre-treated with 40 μM zVAD or

equal volume of vehicle (DMSO) for 1 hr prior to TRAIL treatment. The effectiveness of

zVAD treatment was confirmed by showing it blocked the cleavage of the CASP8 target

BID and blocked the degradation of the CASP3 substrate PARP1 (Supplementary

Figure 2A). The decreases in total and nascent 4sU-labeled mRNA abundance upon

TRAIL treatment were rescued in the presence of zVAD (Figure 2A-B), confirming the

role of canonical apoptotic signaling in both phenotypes.

Liu et al. (2018) reported that the mRNA degradation that occurs in HCT116 cells

shortly after TRAIL treatment is widespread. In order to quantify the extent of the

associated transcriptional arrest, we sequenced the 4sU-labeled nascent transcriptome

of non-apoptotic and apoptotic cells. HCT116 cells were pre-treated with DMSO (i.e.

allowing for apoptosis signaling to proceed) or zVAD (to prevent apoptosis) prior to

addition of TRAIL (or vehicle) for 2 hr. Nascent RNA was depleted of rRNA before

sequencing libraries were prepared, and >95% of the resultant reads mapped to

mRNAs and other predominantly RNAPII transcripts such as lncRNAs (Marchese,

Raimondi, & Huarte, 2017) and snoRNAs (Dieci, Preti, & Montanini, 2009)

(Supplementary Tables 1 and 2). Significantly more nascent transcripts decreased than

increased upon TRAIL treatment in cells pre-treated with DMSO ($p \cong 0$, chi-squared

test), while significantly more increased than decreased with TRAIL treatment in the

presence of zVAD ($p = 5.234\text{e-}143$, chi-squared test). Markedly, 71.2% of the ~28,000

unique transcripts detected were downregulated more than 2-fold in TRAIL-treated cells

with DMSO (Figure 2C), while only 14.7% of ~32,000 transcripts were repressed upon

TRAIL treatment when caspases were inhibited with zVAD (Figure 2D). By contrast,

fewer than 20% of transcripts were transcriptionally upregulated by the same amount

during apoptosis in both conditions (Figure 2C-D). To validate that RNA production was

accurately captured in the sequencing libraries, fold changes for a representative

transcript (*ACTB*) in the original 4sU-labeled RNA samples were assessed by RT-qPCR

using both exonic and intronic primers (Supplementary Figure 2B), showing good

agreement between the three quantification methods.

Of the transcripts changed by 2-fold or greater in TRAIL-treated cells in the

absence of zVAD, 745 decreased and 69 increased in a statistically significant manner

across 2 biological replicates (Supplementary Figure 2C). By contrast, only 56

transcripts were significantly downregulated upon TRAIL treatment in the presence of

zVAD, while 364 were upregulated (Supplementary Figure 2D). Gene ontology

enrichment analysis revealed that the genes upregulated upon TRAIL treatment were

disproportionately involved in cellular responses to chemokines and cell death (Figure

2E, Supplementary Table 3), while no statistically significant enrichments were

observed in the downregulated transcripts (Supplementary Table 4). Interestingly, the

induced genes were more likely to be regulated by transcription factors implicated in

apoptosis such as CSRNP1 (Ye et al., 2017), FOSB (Baumann et al., 2003), NR4A3

(Fedorova et al., 2019), NFKB2 (Keller et al., 2010), JUNB (Gurzov, Ortis, Bakiri,

Wagner, & Eizirik, 2008), JUN (Wisdom, Johnson, & Moore, 1999), EGR1 (Pignatelli,

Luna-Medina, Pérez-Rendón, Santos, & Perez-Castillo, 2003), FOS (Preston et al.,

1996), KFL6 (Huang, Li, & Guo, 2008), RELB (Guerin et al., 2002), and BATF3 (Qiu,

Khairallah, Romanov, & Sheridan, 2020), indicating that the dataset reflects established

apoptotic transcriptional dynamics (Figure 2F, Supplementary Table 5).

Transcriptional repression during early apoptosis requires MOMP, but not

necessarily caspase activity

We next sought to determine whether any hallmark features of apoptosis

underlay the observed transcriptional repression. These include the limited proteolysis

by the "initiator" CASP8, mass proteolysis by the "executioner" CASP3, the

endonucleolytic cleavage of the genome by a caspase-activated DNase (CAD) that

translocates into the nucleus during late stages of apoptosis (Enari et al., 1998) and

MOMP (which instigates mRNA decay).

Small interfering RNA (siRNA) knockdowns (Figure 3A) were performed to first

determine if the rescue of mRNA production caused by zVAD was due specifically to

the inhibition of the initiator CASP8 or the executioner CASP3. Both accelerated mRNA

decay (Supplementary Figure 3A) and RNAPII transcriptional repression (Figure 3B)

upon 2 hr TRAIL treatment required the MOMP-inducing CASP8 but not CASP3,

suggesting that mass proteolysis by CASP3 does not significantly contribute to either

phenotype. Knockdown of CAD did not affect the reduction in 4sU incorporation

observed during early apoptosis (Figure 3B), and DNA fragmentation (as measured by

TUNEL assay) was not detected in TRAIL-treated HCT116 cells until 4-8 hr post-

treatment (Supplementary Figure 3C). These findings are in agreement with the prior

study showing that MOMP and mRNA decay occur before DNA fragmentation begins

during extrinsic apoptosis (Thomas et al., 2015).

Although CASP3 is dispensable for apoptotic transcriptional repression, a

previous report suggests that CASP8 may cleave RPB1, the largest subunit of RNAPII

(Lu, Luo, & Bregman, 2002). We therefore measured the protein expression of RPB1

and the 3 next largest RNAPII subunits (RBP2-4) during early apoptosis to determine if

degradation of these subunits might explain the observed TRAIL-induced reduction in

RNAPII transcription. Expression of RPB1-4 was relatively unaffected by 2 hr TRAIL

treatment in the presence or absence of zVAD, with the exception of a small decrease

in the amount of RPB2 that was rescued upon zVAD treatment (Supplementary Figure

3C). Thus, RNAPII depletion is unlikely to underlie the transcriptional repression

phenotype.

Our above observations suggest that MOMP activation, for example through

CASP8, is necessary to drive apoptotic mRNA decay and the ensuing transcriptional

repression. To test this hypothesis, we attenuated MOMP in TRAIL-treated cells by

depleting the mitochondrial pore-forming proteins BAX and BAK (Figure 3C). Indeed,

siRNA-mediated depletion of BAX and BAK rescued cytoplasmic mRNA abundance

(Supplementary Figure 3D) and RNAPII transcription (Figure 3D) of the *ACTB* and

GAPDH transcripts in the presence of TRAIL, even though CASP8 and CASP3 were

still cleaved (Figure 3D). Thus, CASP8 likely participates in this pathway only to the

extent that it activates MOMP, as MOMP appears to be the main driver of mRNA decay

and transcriptional repression.

Finally, to confirm that MOMP is sufficient to drive this phenotype, we used a

small molecule, raptinal, that bypasses CASP8 to intrinsically induce MOMP (Heimer,

Knoll, Schulze-Osthoff, & Ehrenschwender, 2019; Palchaudhuri et al., 2015). HeLa cells

treated with 10 μM raptinal for 4 hr underwent MOMP, as measured by cytochrome c

release into the cytoplasm, in the presence or absence of zVAD (Supplementary Figure

3E). Steady state levels (Figure 3E) and transcription (Figure 3F) of the aforementioned

mRNAs was reduced upon raptinal treatment. The fact that this reduction in mRNA

abundance and synthesis was maintained upon caspase inhibition by zVAD treatment

indicates that the caspases are not required to drive these phenotypes outside of their

role in MOMP activation.. Taken together, these data confirm that the mRNA

degradation and ensuing transcriptional repression observed during early apoptosis are

driven by MOMP.

Cytoplasmic 3'- but not 5'-RNA exonucleases are required for apoptotic RNAPII

transcriptional repression

Based on connections between virus-activated mRNA decay and RNAPII

transcription (Abernathy et al., 2015; Gilbertson, Federspiel, Hartenian, Cristea, &

Glaunsinger, 2018), we hypothesized that TRAIL-induced mRNA turnover was

functionally linked to the concurrent transcriptional repression. Apoptotic mRNA decay

occurs from the 3' end by the actions of the cytoplasmic 3'-RNA exonuclease DIS3L2

and the mitochondrial 3'-RNA exonuclease PNPT1, which is released into the

cytoplasm by MOMP (Liu et al., 2018; Thomas et al., 2015). This stands in contrast to

basal mRNA decay, which occurs predominantly from the 5' end by XRN1 (Jones,

Zabolotskaya, & Newbury, 2012). We therefore set out determine if 3' or 5' decay

factors were required for apoptosis-linked mRNA decay and the ensuing repression of

mRNA transcription. Depletion of DIS3L2, PNPT1, or the cytoplasmic 3' RNA exosome

subunit EXOSC4 individually did not reproducibly rescue the total levels of either the

ACTB or *GAPDH* mRNA during early apoptosis (Supplementary Figure 4A), nor did

they affect the relative production of these transcripts (Supplementary Figure 4B). Given

the likely redundant nature of the multiple 3' end decay factors (Houseley & Tollervey,

2009), we instead performed concurrent knockdowns of DIS3L2, EXOSC4, and PNPT1

to more completely inhibit cytoplasmic 3' RNA decay. We also knocked down the

predominant 5'-3' RNA exonuclease XRN1 to check the involvement of 5' decay (Figure

4A). Depletion of the 3'-5' but not the 5'-3' decay machinery attenuated the apoptotic

decrease in total RNA levels and largely restored RNAPII transcription (Figure 4B-C).

Importantly, there was only a minor reduction in CASP3 activation in cells depleted of

3'-5' decay factors and this was not significantly different from that observed upon

XRN1 knockdown (Figure 4A). These observations suggest that decreased RNAPII

transcription occurs as a consequence of accelerated 3' mRNA degradation in the

cytoplasm during early apoptosis.

Apoptosis causes reduced RNAPII promoter occupancy in an mRNA decay-

dependent manner

272   RNAPII is recruited to promoters in an unphosphorylated state, but its

subsequent promoter escape and elongation are governed by a series of

phosphorylation events in the heptad $(Y_1S_2P_3T_4S_5P_6S_7)_n$ repeats of the RPB1 C-terminal

domain (CTD). To determine the stage of transcription impacted by mRNA decay, we

performed RNAPII ChIP-qPCR and western blots using antibodies recognizing the

different RNAPII phosphorylation states. Occupancy of hypophosphorylated RNAPII at

the *ACTB* and *GAPDH* promoters was significantly reduced after 2 hr TRAIL treatment

(Figure 3D). In accordance with the 4sU labeling results, siRNA-mediated knockdowns

showed that loss of RNAPII occupancy in response to TRAIL requires CASP8 but not

CASP3 (Supplementary Figure 4C), as well as cytoplasmic 3'-5' RNA decay factors but

not the 5'-3' exonuclease XRN1 (Figure 4D). Impaired binding of the TATA-binding

protein (TBP), which nucleates the formation of the RNAPII pre-initiation complex (PIC)

at promoters (Buratowski, Hahn, Guarente, & Sharp, 1989; Louder et al., 2016),

mirrored that of RNAPII (Figure 4E). These changes are not driven by alterations in the

stability of RPB1 or TBP, since the expression of these proteins remained relatively

constant during early apoptosis regardless of the presence of mRNA decay factors

(Supplementary Figure 4D-E). The relative proportion of initiating RPB1 CTD

phosphorylated at the serine 5 position and the ratio of serine 5 to serine 2

phosphorylation, which decreases during transcriptional elongation (Shandilya &

Roberts, 2012), also remain unchanged during early apoptosis regardless of the

presence of mRNA decay factors (Figure 4F). These data suggest that the decay-

dependent reduction in mRNA synthesis occurs at or before PIC formation, rather than

during transcriptional initiation and elongation.

Importin α/β transport links mRNA decay and transcription

Finally, we sought to determine how TRAIL-induced cytoplasmic mRNA decay

signals to the nucleus to induce transcriptional repression. Data from viral systems

suggest that this signaling involves differential trafficking of RNA binding proteins

(RBPs), many of which transit to the nucleus in response to virus-induced cytoplasmic

mRNA decay (Gilbertson et al., 2018). We therefore sought to test the hypothesis that

nuclear import of RBPs underlies how cytoplasmic mRNA decay is sensed by the

RNAPII transcriptional machinery.

To determine whether RBP redistribution occurs during early apoptosis, we

analyzed the subcellular distribution of cytoplasmic poly(A) binding protein PABPC1, an

RBP known to shuttle to the nucleus in response to virus-induced RNA decay

(Gilbertson et al., 2018; Kumar & Glaunsinger, 2010; Lee & Glaunsinger, 2009). We

performed cell fractionations of HCT116 cells to measure PABPC1 levels in the nucleus

versus the cytoplasm upon induction of apoptosis. Indeed, 2 hr after TRAIL treatment,

PABPC1 protein levels increased in the nuclear fraction of cells but not in the

cytoplasmic fraction, indicative of relocalization (Figure 5A). This increase in nuclear

PABPC was dependent on the presence of CASP8 but not CASP3, mirroring the

incidence of mRNA decay and reduced RNAPII transcription. Thus, similar to viral

infection, PABPC1 relocalization occurs coincidentally with increased mRNA decay and

transcriptional repression in the context of early apoptosis.

We next sought to directly evaluate the role of protein shuttling in connecting

cytoplasmic mRNA decay to transcriptional repression. The majority of proteins ~60

kilodaltons (kDa) and larger cannot passively diffuse through nuclear pores; they require

assistance by importins to enter the nucleus from the cytoplasm (Gö, 1998). Classical

nuclear transport occurs by importin α binding to a cytoplasmic substrate, which is then

bound by an importin β to form a tertiary complex that is able to move through nuclear

pore complexes (Stewart, 2007). To test if this mode of nuclear transport is required for

transcriptional feedback, HCT116 cells were pretreated with ivermectin, a specific

inhibitor of importin α/β transport (Wagstaff, Sivakumaran, Heaton, Harrich, & Jans,

2012), before the 2 hr TRAIL treatment. The efficacy of ivermectin was validated by an

observed decrease in the nuclear levels of the RBP nucleolin (NCL), a known importin

α/β substrate (Figure 5B). Ivermectin pretreatment rescued RNAPII transcription upon

TRAIL treatment (Figure 5C) without significantly affecting the extent of mRNA decay

(Supplementary Figure 5A), suggesting that protein trafficking to the nucleus provides

signals connecting cytoplasmic mRNA decay to transcription.

Importantly, ivermectin did not diminish the extent of CASP3 cleavage during

early apoptosis (Figure 5B), nor did it decrease baseline levels of transcription in non-

apoptotic cells (Supplementary Figure 5B). Interestingly, it also did not prevent PABPC1

import (Supplementary Figure 5C), suggesting that PABPC1 translocation likely occurs

via an importin α/β-independent pathway and is not sufficient to repress RNAPII

transcription.

Finally, we evaluated whether importin α/β transport is also required for feedback

between viral nuclease-driven mRNA decay and RNAPII transcription, as this would

suggest that the underlying mechanisms involved in activating this pathway may be

conserved. We used the mRNA specific endonuclease muSOX from the

gammaherpesvirus MHV68, as muSOX expression has been shown to cause

widespread mRNA decay and subsequent transcriptional repression (Abernathy et al.,

2014, 2015). HEK-293T cell lines were engineered to stably express dox-inducible wild-

type muSOX or the catalytically inactive D219A mutant (Abernathy et al., 2015). These

cells were treated with ivermectin for 3 hr and the resultant changes in RNAPII promoter

occupancy were measured by ChIP-qPCR. As expected, expression of WT (Figure 5D-

E) but not D219A muSOX (Supplementary Figure 5E-F) caused mRNA decay and

transcriptional repression. Notably, inhibiting nuclear import with ivermectin

(Supplementary Figure 5D) rescued RNAPII promoter occupancy (Figure 5E) without

altering the extent of mRNA decay in muSOX expressing cells (Figure 5D). Thus,

importin α/β nuclear transport plays a key role in linking cytoplasmic mRNA decay to

nuclear transcription, both during early apoptosis and upon viral nuclease expression.

DISCUSSION

mRNA decay and synthesis rates are tightly regulated in order to maintain

appropriate levels of cellular mRNA transcripts (Braun & Young, 2014). It is well

established that when cytoplasmic mRNA is stabilized, for example by the depletion of

RNA exonucleases, transcription often slows in order to compensate for increased

transcript abundance (Haimovich et al., 2013; Helenius et al., 2011; Singh et al., 2019;

Sun et al., 2012). Eukaryotic cells thus have the capacity to "buffer" against reductions

in mRNA turnover or synthesis. Here, we revealed that a buffering response does not

occur under conditions of elevated cytoplasmic mRNA degradation stimulated during

early apoptosis. Instead, cells respond to cytoplasmic mRNA depletion by decreasing

RNAPII promoter occupancy and transcript synthesis, thereby amplifying the magnitude

of the gene expression shut down. Nuclear import of cytoplasmic proteins is required for

this "transcriptional feedback", suggesting a pathway of gene regulation in which

enhanced mRNA decay prompts cytoplasmic proteins to enter the nucleus and halt

mRNA production. Notably, similar transcriptional feedback is elicited during virus-

induced mRNA decay (Abernathy et al., 2015; Gilbertson et al., 2018; Hartenian et al.,

2020), indicating that distinct cellular stresses can converge on this pathway to

potentiate a multi-tiered shutdown of gene expression.

Multiple experiments support the conclusion that the TRAIL-induced

transcriptional repression phenotype is a consequence cytoplasmic decay triggered by

MOMP, rather than caspase activity. XRN1-driven 5'-3' end decay is the major pathway

involved in basal mRNA decay (Łabno, Tomecki, & Dziembowski, 2016), but MOMP-

induced mRNA decay is primarily driven by 3' exonucleases such as PNPT1 and

DIS3L2 (Liu et al., 2018; Thomas et al., 2015). Accordingly, co-depletion of 3' decay

factors but not XRN1 restored RNAPII promoter occupancy and transcription during

early apoptosis. In contrast to that of 3' mRNA decay factors, depletion of CASP3 (or

the caspase activated DNase CAD) did not block mRNA degradation or transcriptional

repression, even though CASP3 is responsible for the vast majority of proteolysis that is

characteristic of apoptotic cell death (J. G. Walsh et al., 2008). Additionally, the initiator

CASP8 was required only under conditions where its activity was needed to induce

MOMP. These findings reinforce the idea that mRNA decay and transcriptional

repression are very early events that are independent of the subsequent cascade of

caspase-driven phenotypes underlying many of the hallmark features of apoptosis.

A key open question is what signal conveys cytoplasmic mRNA degradation

information to the nucleus to cause transcriptional repression. Our data are consistent

with a model in which the signal is provided by one or more proteins imported into the

nucleus in response to accelerated mRNA decay (Figure 6). Indeed, many cytoplasmic

RNA binding proteins undergo nuclear-cytoplasmic redistribution under conditions of

viral nuclease-induced mRNA decay, including PABPC (Gilbertson et al., 2018; Kumar

& Glaunsinger, 2010; Kumar, Shum, & Glaunsinger, 2011). We propose that a certain

threshold of mRNA degradation is necessary to elicit protein trafficking and

transcriptional repression. Presumably, normal levels of basal mRNA decay and regular

cytoplasmic repopulation result in a balanced level of mRNA-bound versus unbound

proteins. However, if this balance is tipped during accelerated mRNA decay, an excess

of unbound RNA binding proteins could accumulate and be transported into the

nucleus.

PABPC1 is likely among the first proteins released from mRNA transcripts

undergoing 3' end degradation, and its nuclear translocation is driven by a poly(A)-

masked nuclear localization signal that is exposed upon RNA decay (Kumar et al.,

2011). Indeed, we found that PABPC1 undergoes nuclear translocation during early

apoptosis. However, the fact that ivermectin blocks transcriptional repression but not

PABPC1 import indicates that while PABPC1 redistribution is a marker of mRNA decay,

it is not sufficient to induce transcriptional repression in this system. Instead, we

hypothesize that the observed RNAPII transcriptional repression occurs as a result of

the cumulative nuclear import of multiple factors via importin α/β. Our group previously

reported that at least 66 proteins in addition to PABPC1 are selectively enriched in the

nucleus upon transfection with the viral endonuclease muSOX (but not with the

catalytically-dead D219A mutant), 22 of which are known to be RNA-binding proteins

and 45 of which shuttle in a manner dependent on the cytoplasmic mRNA exonuclease

primarily responsible for clearing endonuclease cleavage fragments, XRN1 (Gilbertson

et al., 2018). Future studies in which changes in the nuclear and cytoplasmic proteome

upon apoptosis induction and viral endonuclease expression in the presence and

absence of ivermectin will likely provide insight into which additional protein or proteins

play a role in connecting cytoplasmic mRNA turnover to RNAPII transcription.

Nonetheless, the requirement for importin α/β–mediated nuclear transport in both

apoptosis-induced and viral nuclease-induced transcriptional repression suggests that

these two stimuli may activate a conserved pathway of gene regulation. Whether other

types of cell stress elicit a similar response remains an important question for future

investigation.

The mRNA 3' end decay-dependent decrease in RNA synthesis is accompanied

by a reduction in TBP and RNAPII occupancy at promoters, consistent with

transcriptional inhibition occurring upstream of RNAPII initiation and elongation

(Gilbertson et al., 2018; Hartenian et al., 2020). The mechanism driving such a defect is

yet to be defined, but possibilities include changes to transcript processing pathways,

transcriptional regulators, or the chromatin state that directly or indirectly impact

formation of the preinitiation complex. In this regard, mRNA processing and

transcription are functionally linked (Bentley, 2014) and nuclear accumulation of

PABPC1 has been shown to affect mRNA processing by inducing hyperadenylation of

nascent transcripts (Kumar & Glaunsinger, 2010; Lee & Glaunsinger, 2009).

Furthermore, TBP interacts with the cleavage–polyadenylation specificity factor

(Dantonel, Murthy, Manjey, & Tora, 1997), providing a possible link between RNAPII

preinitiation complex assembly and polyadenylation. Such a mechanism could also

explain the specificity of the observed transcriptional repression to RNAPII transcripts.

Chromatin architecture could also be influenced by shuttling of RNA binding proteins, as

for example the yeast nucleocytoplasmic protein Mrn1 is implicated in the function of

chromatin remodeling complexes (Düring et al., 2012). Interestingly, the polycomb

repressive chromatin complex 2 (PRC2) and DNA methyltransferase 1 (DNMT1) have

been shown to bind both nuclear mRNA and chromatin in a mutually exclusive manner,

(Beltran et al., 2016; Di Ruscio et al., 2013; Garland et al., 2019), evoking the possibility

that a nuclear influx of RBPs could secondarily increase the chromatin association of

such transcriptional repressors. Future work will test these possibilities in order to

mechanistically define the connection between nuclear import and RNAPII transcription

under conditions of enhanced mRNA decay.

There are several potential benefits to dampening mRNA transcription in

response to accelerated mRNA turnover. Debris from apoptotic cells is usually cleared

by macrophages, but inefficient clearance of dead cells can lead to the development of

autoantibodies to intracellular components such as histones, DNA, and

ribonucleoprotein complexes. This contributes to autoimmune conditions such as

systemic lupus erythematosus (Caruso & Poon, 2018; Nagata, Hanayama, & Kawane,

2010). In the context of infection, many viruses require the host RNAPII transcriptional

machinery to express viral genes (Harwig et al., 2017; Rivas, Schmaling, & Gaglia,

2016; Walker & Fodor, 2019). It may therefore be advantageous for the cell to halt

transcription in attempt to pre-empt a viral takeover of mRNA synthesis. That said, as

with antiviral translational shutdown mechanisms (D. Walsh et al., 2013), viruses have

evolved strategies to evade transcriptional repression (Hartenian et al., 2020; Harwig et

al., 2017), as well as inhibit cell death via apoptosis and/or co-opt apoptotic signaling

cascades (Suffert et al., 2011; Tabtieng, Degterev, & Gaglia, 2018; Zhang, Hildreth, &

Colberg-Poley, 2013; Zhou, Jiang, Liu, Liu, & Liang, 2017). In either case, the shutdown

of transcription in a cell under stress may protect surrounding cells from danger,

particularly in an in vivo context. If true, this type of response is likely to be more

relevant in multicellular compared with unicellular organisms.

ACKNOWLEDGEMENTS

This work was supported by NIH grant R01CA136367 to B.G. and the HHMI

Gilliam Fellowship for Advanced Study to C.D. B.G. is an investigator of the Howard

Hughes Medical Institute. We thank the UC Berkeley Cell Culture Facility for providing

the cell lines used in this study, in addition to the UC Berkeley DNA Sequencing Facility

and all members of the Glaunsinger and Coscoy labs for providing valuable feedback.

AUTHOR CONTRIBUTIONS

C.D. conceived and performed the experiments, analyzed and visualized the

data, acquired funding, and drafted the manuscript. E.H. created the stable cell lines

and drafted the corresponding section of the manuscript, as well as directed and

assisted with analysis of the 4sU-seq data. V.K. analyzed the 4sU-seq data. B.G.

conceived and supervised the study, acquired funding, and edited the manuscript.

DECLARATION OF INTERESTS

The authors declare no competing interests.

FIGURE LEGENDS

Figure 1. mRNA decay during early apoptosis is accompanied by reduced synthesis of

RNAPII transcripts. (A) Schematic representation of how the extrinsic apoptotic pathway

accelerates mRNA decay. (B) Western blot of HCT116 lysates showing the depletion of

full-length caspase 8 (CASP8) and caspase 3 (CASP3) over a time course of 100 ng/μL

TRAIL treatment. Vinculin (VCL) serves as a loading control. Blot representative of

those from 4 biological replicates. (C, D) RT-qPCR quantification of total (C) and

nascent 4sU pulse-labeled (D) RNA at the indicated times post TRAIL treatment of

HCT116 cells (n = 4). Also see Supplementary Figure 1. No biotin control quantifies

RNA not conjugated to biotin that is pulled down with strepdavidin selection beads. Fold

changes were calculated from Ct values normalized to 18S rRNA in reference to mock

treated cells. Graphs display mean +/- SEM with individual biological replicates

represented as dots. Statistically significant deviation from a null hypothesis of 1 was

determined using one sample t test; *p<0.05, **p<0.01, ***p<0.001.

Figure 2. RNAPII transcription is globally repressed during early apoptosis. (A, B) RT-

qPCR measurements of total (A) and nascent 4sU-labeled (B) RNA fold changes after 2

hr TRAIL treatment of HCT116 cells, including a 1 hr pre-treatment with either 40 μM

zVAD or an equal volume of DMSO ("mock"). Also see Supplementary Figure 2A. RNA

fold change values were calculated in reference to *18S* rRNA. Bar graphs display mean

+/- SEM with individual biological replicates (*n* = 4) represented as dots. Statistically

significant deviation from a null hypothesis of 1 was determined using one sample t test

and indicated with asterisks directly above bars, while student's t tests were performed

to compare mean fold change values for mock inhibitor or scramble treated cells to

those treated with inhibitor or a targeting siRNA and indicated with brackets. *p<0.05,

**p<0.0.1, ***p<0.001. (C, D) rRNA-depleted cDNA sequencing libraries were reverse

transcribed from 4sU-labeled RNA isolated from cells under the conditions described in

(A, B). Transcripts that aligned to genes in the human genome are graphed with

differential log2 fold change expression values (log2FC) on the y axis and fragments per

kilobase per million reads (FKPM) expression values on the x axis. All values were

averaged from 2 biological replicates. Data points for transcripts upregulated or

downregulated by 2-fold or greater are colored green and purple, respectively.

Percentages of transcripts in each expression class are indicated with an arrow and in

their corresponding colors. Also see Supplementary Figure 2C-D and Supplementary

Tables 1-2. (E) Top statistically significant hits from gene ontology analysis performed

for the list of (69) transcripts that were upregulated upon TRAIL treatment with DMSO in

a statistically significant manner across biological duplicates. Also see Supplementary

Table 3. (F) Top hits from transcription factor (TF) enrichment analysis for the same list

of genes as above. The lower the MeanRank value, the more statistically significant

enrichment for genes regulated by the indicated TF. Also see Supplementary Table 4.

Figure 3 Transcriptional repression during early apoptosis requires MOMP, but not

necessarily caspase activity. (A) Western blots showing the efficacy of CASP3, CASP8,

and caspase-activated DNase (CAD) protein depletion following nucleofection with the

indicated siRNAs, with or without 2 hr TRAIL treatment. α-Tubulin (TUBA1A) serves as

a loading control. Blot representative of those from 4 biological replicates. (B) RT-qPCR

measurements of 4sU-labeled nascent transcripts with or without 2 hr TRAIL treatment

in cells nucleofected with the indicated the siRNAs ($n = 3$). Also see Supplementary

Figures 3A-C. (C) Western blot detecting the indicated proteins in cells nucleofected

with the indicated siRNA in the presence or absence of TRAIL. TUBA1A serves as a

loading control. Blot representative of 4 biological replicates. The cleavage of CASP3

and CASP8 (as measured by the disappearance of the full-length form of each

zymogen upon 2 hr TRAIL treatment, normalized to TUBA1A) is graphed below ($n = 4$).

(D) 4sU-labeled RNA levels measured by RT-qPCR in HCT116 cells nucleofected with

the indicated siRNAs, with or without 2 hr TRAIL treatment ($n = 4$). Also see

Supplementary Figure 3D. (E, F) Total (E) and 4sU-labeled (F) RNA levels measured

by RT-qPCR in HeLa cells after 10 μM raptinal treatment for 4 hr, with or without a 1 hr

pre-treatment of 20 μM zVAD ($n = 4$). Also see Supplementary Figure 2E. Fold changes

were calculated in reference to *U6* snRNA since its production was more stable after 4

hr raptinal than that of *18S* rRNA (see Supplementary Figure 2F). All RNA fold changes

were calculated from Ct values normalized to *18S* or *U6* RNA, then normalized to non-

apoptotic cells ("no TRAIL") under otherwise identical conditions. Graphs display mean

+/- SEM with individual biological replicates represented as dots. Statistically significant

deviation from a null hypothesis of 1 was determined using one sample t test and

indicated with asterisks directly above bars, while student's t tests were performed to

compare mean fold change values for mock inhibitor or scramble treated cells to those

treated with zVAD or a targeting siRNA and indicated with brackets. The Holm-Sidak

correction for multiple comparisons was applied in the student's t tests represented in

(B) *$p<0.05$, **$p<0.0.1$, ***$p<0.001$.

Figure 4. Apoptosis causes reduced RNAPII transcriptional output and promoter

occupancy in an mRNA decay-dependent manner. (A) Western blots performed with

HCT116 cell lysates depleted of the indicated decay factors with and without 2 hr TRAIL

treatment. Blot representative of 3 biological replicates. Apoptosis induction was

confirmed by disappearance of the full-length CASP3 band, quantified in the graph

below by measuring band intensity normalized to an α-tubulin (TUBA1A) loading control

($n = 3$). (B, C) Changes in total (B) and nascent 4sU-labeled (C) RNA upon 2 hr TRAIL

treatment in cells nucleofected with the indicated siRNAs were quantified by RT-qPCR

($n = 4$). Fold changes were calculated from Ct values normalized to *18S* rRNA. Also see

Supplementary Figure 4A-B. (D, E) Chromatin immunoprecipitation (ChIP)-qPCR was

used to measure occupancy of the indicated promoters by hypophosphorylated RNAPII

(D) or TBP (E) under cellular conditions described in (A). Rabbit and mouse IgG

antibodies were included in parallel immunoprecipitation reactions with chromatin from

scramble siRNA-treated non-apoptotic cells in lieu of TBP and RNAPII antibodies,

respectively, as a control. Also see Supplementary Figure 4D-E. (F) Relative band

intensity ratios from 4 replicates of the representative western blots depicted in

Supplementary Figure 4D, using primary antibodies specific to the indicated RPB1 CTD

phosphorylation state under cellular conditions described in (A). Band intensity values

were first normalized to a vinculin (VCL) loading control in each blot. All bar graphs

display mean +/- SEM with individual biological replicates represented as dots.

Statistically significant deviation from a null hypothesis of 1 was determined using one

sample t test and indicated with asterisks directly above bars, while student's t tests with

the Holm-Sidak correction for multiple comparisons were performed to compare mean

values between groups indicated with brackets. *$p<0.05$, **$p<0.0.1$, ***$p<0.001$.

Figure 5. Importin α/β transport links mRNA decay and transcription. (A) Western blots

showing the indicated proteins in the nuclear and cytoplasmic fractions of apoptotic and

non-apoptotic HCT116 cells nucleofected with the indicated siRNAs. The nuclear and

cytoplasmic fractions of PABPC1 were imaged on the same membrane section but

cropped and edited separately to visualize the low nuclear expression of this canonically

cytoplasmic protein. Protein expression (right) was calculated by band intensity in

reference to lamin B1 (LMNB1) or α-tubulin (TUBA1A) loading controls, for the nuclear

and cytoplasmic fractions, respectively. (B) Nuclear and cytoplasmic expression of the

indicated proteins in apoptotic and non-apoptotic cells with a 1 hr pre-treatment with 25

$\square$M ivermectin or an equal volume of EtOH ("mock"). Nuclear levels of the importin

α/β substrate nucleolin (NCL) were quantified by band intensity in reference to a LMNB1

loading control, while disappearance of full-length CASP3 was quantified in the

cytoplasm in reference to TUBA1A. Also see Supplementary Figure 5A (C) Levels of

nascent 4sU-labeled RNA were quantified by RT-qPCR under the cellular conditions

described in (B). Also see Supplementary Figure 5B-C. (D) RT-qPCR quantification of

total RNA levels in HEK293T cells stably expressing a doxycycline (dox)-inducible form

of muSOX endonuclease, cultured with or without 1 μg/mL dox for 24 hr. Cells were

treated with 25 μM ivermectin or an equal volume of EtOH 2 hr before harvesting. (E)

RNAPII promoter occupancy at the ACTB and GAPDH promoters was determined by

ChIP-qPCR under cellular conditions described in (D). Also see Supplementary Figure

5D-F. RNA fold changes were calculated from Ct values normalized to 18S rRNA. All

bar graphs display mean +/- SEM with individual biological replicates represented as

dots. Statistically significant deviation from a null hypothesis of 1 was determined using

one sample t test and indicated with asterisks directly above bars, while student's t tests

were performed to compare mean fold change values for mock inhibitor or scramble

treated cells to those treated with ivermectin or a targeting siRNA and indicated with

brackets. The Holm-Sidak correction for multiple comparisons was applied in the

student's t tests represented in (A) . *p<0.05, **p<0.0.1, ***p<0.001.

Figure 6. Schematic representation of the cellular events connecting apoptosis

induction with mRNA decay and RNAPII transcription.

Supplementary Figure 1. mRNA decay during early apoptosis is accompanied by

reduced synthesis of RNAPII transcripts. (A) RT-qPCR quantification of total RNA at the

indicated times post 100 ng/μL TRAIL treatment of HCT116 cells (n = 4). (B) Stained

agarose gel depicting a 200 nt RT-PCR product of 4sU-labeled 18S rRNA, extracted

and isolated from an equal number of cells treated with 3 hr vehicle ("mock") or 100

ng/μL TRAIL. Gel representative of that from 3 biological replicates. (C) RT-qPCR

quantification of 4sU-labeled RNA at the indicated times post 100 ng/μL TRAIL

treatment of HCT116 cells (n = 4). Fold changes were calculated from Ct values

normalized to *18S* rRNA in reference to mock treated cells. Graphs display mean +/-

SEM with individual biological replicates represented as dots. Statistically significant

deviation from a null hypothesis of 1 was determined using one sample t test; *p<0.05,

**p<0.01, ***p<0.001.

Supplementary Figure 2. RNAPII transcription is globally repressed during early

apoptosis. (A) Western blot of HCT116 lysates showing cleavage (or lack thereof) of the

caspase 8 (CASP8) targets BID and caspase 3 (CASP3), as well as the CASP3 target

PARP1 upon 2 hr 100 ng/µL TRAIL treatment in the presence or absence of 40 µM

zVAD. Blot representative of that from 3 biological replicates. (B) Log2 fold change

($\log_2$FC) in abundance of the ACTB transcript upon 2 hr TRAIL treatment in the

presence of either zVAD or DMSO vehicle ("mock"). Fold changes from the same 4sU-

labeled RNA samples were quantified by next-generation sequencing and differential

expression analysis, as well as RT-qPCR normalized to 18S rRNA using both intronic

and exonic gene-specific primers. Graph displays mean +/- SEM with individual

biological replicates ($n = 2$) represented as dots. Student's t tests were performed to

compare the $\log_2$FC values upon TRAIL treatment in the presence of DMSO vehicle or

zVAD. **$p<0.0.1$ (C, D) Volcano plot depicting the $-\log_{10}p$ values across biological

duplicates for the fragment per kilobase per million read fold changes in the sequencing

libraries depicted in Figures 2C, D. Each point represents a transcript mapped to the

human genome. Upregulated and downregulated transcripts with $p < 0.05$ are colored in

green and purple, respectively. Number of transcripts in each expression class are

indicated with an arrow and in their corresponding colors.

Supplementary Figure 3. Transcriptional repression during early apoptosis requires

MOMP, but not necessarily caspase activity. (A) RT-qPCR quantification of the change

in total RNA levels in HCT116 cells treated with the indicated siRNAs upon 2 hr TRAIL

treatment. Fold change were calculated from Ct values normalized to *18S* rRNA. (B)

Quantification of BrdUTP-labeled HCT116 cells treated with TRAIL for the indicated

times in a TUNEL assay ($n = 4$)*.* Percent positive cells calculated by flow cytometry

using FloJo imaging software to count the number of cells with elevated FL2

fluorescence. (C) Western blot of HCT116 lysates from cells under conditions described

in (A) depicting levels of the indicated RNAPII subunits and CASP3 activation. α-Tubulin

(TUBA1A) serves as a loading control. Graph to the right quantifies the change in

RNAPII subunit levels upon apoptosis induction by TUBA1A-normalized band density.

(D) Total RNA levels measured by RT-qPCR in HCT116 cells nucleofected with the

indicated siRNAs, with or without 2 hr TRAIL treatment ($n = 4$). (E) Western blot of the

cytoplasmic and mitochondrial fractions of HeLa cells treated with 10 μM raptinal or

equal volume of DMSO, in the presence of 20 μM zVAD or additional volume of DMSO.

Efficacy of raptinal treatment was demonstrated by cytochrome c (CYTC) release into

the cytoplasm in the presence and absence of zVAD, which effectively blocked the

cleavage and activation of CASP3. TUBA1A and VDAC1 served as cytoplasmic and

mitochondrial loading controls, respectively. Blot representative of that from 3 biological

replicates. (F) Stained agarose gel depicting a 200 nt and 101 nt RT-PCR product of

4sU-labeled *18S* and *U6* rRNA, respectively, extracted and isolated from an equal

number of cells treated with 4 hr 10 μM raptinal or equal volume of DMSO. Gel

representative of that from 3 biological replicates. All bar graphs display mean +/- SEM

with individual biological replicates represented as dots. Statistically significant deviation

from a null hypothesis of 1 was determined using one sample t test and indicated with

asterisks directly above bars, while student's t tests were performed to compare mean

fold change values for mock inhibitor or scramble treated cells to those treated with

zVAD or a targeting siRNA and indicated with brackets. The Holm-Sidak correction for

multiple comparisons was applied in the student's t tests represented in (B). *p<0.05,

**p<0.0.1, ***p<0.001.

Supplementary Figure 4. Apoptosis causes reduced RNAPII transcriptional output and

promoter occupancy in an mRNA decay-dependent manner. (A, B) RT-qPCR

quantification of the change in total (A) and 4sU-labeled (B) RNA levels in HCT116 cells

treated with the indicated siRNAs upon 2 hr TRAIL treatment ($n = 4$). Fold changes

were calculated from Ct values normalized to *18S* rRNA. (C) Chromatin

immunoprecipitation (ChIP)-qPCR was used to measure the change in

hypophosorylated RNAPII occupancy at the indicated promoters in cells nucleofected

with the indicated siRNAs ($n = 4$). Mouse IgG antibody was included in parallel

immunoprecipitation reactions with chromatin from scramble siRNA-treated non-

apoptotic cells in lieu of RNAPII antibody as a control. (D) Representative western blots

on lysates of HCT116 cells depleted of either 3' DFs (DIS3L2, EXOSC4, and PNPT1) or

XRN1 before and after 2 hr TRAIL treatment, probing for the indicated protein or RPB1

C-terminal domain (CTD) phosphorylation state. Apoptosis induction was confirmed by

observing CASP3 cleavage. A vinculin (VCL) loading control was imaged for the blots of

each phosphorylation state, but a single representative panel is shown. Quantification of

RPB1 phosphorylation states from 4 replicates displayed in Figure 4F. (E) Band

intensity fold changes of hypophosphorylated RPB1 and TBP derived from 4 biological

replicates of the experiment described in (D). Bar graph displays mean +/- SEM with

individual biological replicates represented as dots. Statistically significant deviation

from a null hypothesis of 1 was determined using one sample t test and indicated with

asterisks directly above bars, while student's t tests with the Holm-Sidak correction for

multiple comparisons were performed to compare mean values between groups

indicated with brackets. *p<0.05, **p<0.0.1, ***p<0.001.

Supplementary Figure 5. Importin α/β transport links mRNA decay and transcription.

(A) Western blot showing the indicated proteins in the nuclear and cytoplasmic fractions

of HCT116 cells with or without 2 hr TRAIL treatment and/or 3 hr ivermectin. Lamin B1

(LMNB1) and TUBA1A served as nuclear and cytoplasmic loading controls, repectively.

Blot representative of that from 3 biological replicates. (B) Changes in total RNA levels

after 2 hr TRAIL treatment in HCT116 cells pretreated with either 25 μM ivermectin or

an equal volume of ethanol ("mock") for 1 hr (n = 3). (C) Alternative analysis of data

presented in Figure 5C, instead quantifying the fold change in nascent transcription in

non-apoptotic cells in the presence or absence of 25 μM ivermectin (n = 3). (D) Western

blot showing nuclear and cytoplasmic expression of the indicated proteins in HEK293T

expressing the viral endonuclease muSOX with a 3 hr treatment of 25 μM ivermectin or

equal volume of ethanol. Nuclear and cytoplasmic levels (right) of nucleolin (NCL) were

quantified by band intensity in reference to a lamin B1 (LMNB1) or glyceraldehyde 3-

phosphate dehydrogenase (GAPDH) loading control, respectively. (E) RT-qPCR

quantification of the change in total RNA levels upon dox-inducible expression of the

catalytically dead D219A mutant of muSOX with a 3 hr treatment of 25 μM ivermectin or

equal volume of ethanol. (F) RNAPII occupancy at the *ACTB* and *GAPDH* promoters

under cellular conditions described in (E). RNA fold change values were calculated in

reference to *18S* rRNA. All bar graphs display mean +/- SEM with individual biological

replicates represented as dots. Statistically significant deviation from a null hypothesis

of 1 was determined using one sample t test and indicated with asterisks directly above

bars, while student's t tests were performed to compare mean fold change values for

mock inhibitor or scramble treated cells to those treated with inhibitor or a targeting

siRNA and indicated with brackets. $*p<0.05$, $**p<0.0.1$, $***p<0.001$.

MATERIALS AND METHODS

Lead Contact

717 Further information and requests for resources and reagents should be directed

718 to and will be fulfilled by the Lead Contact, Britt Glaunsinger

719 (glaunsinger@berkeley.edu).

Materials availability

721 The dox-inducible muSOX-expressing cell line and rabbit polyclonal anti-muSOX

722 antibody generated in this study are both available upon request.

Data and Code Availability

724 Sequencing data generated in this study are publicly available on GEO repository

725 (accession number GSE163923).

Cells and culture conditions

Wild-type HCT116, HEK293T, and HeLa cells (all from ATCC) were obtained

from the UC Berkeley Tissue Culture Facility. Cell lines were authenticated by STR

analysis and determined to be free of mycoplasma by PCR screening. HEK293Ts were

made to stably express doxycycline(dox)-inducible wild-type muSOX and its

catalytically-dead D219A mutant by PCR amplifying the aforementioned coding

sequences from Addgene plasmids 131702 and 131704 using the muSOX F/R primers

(see Key Resources Table) and InFusion cloning these fragments into the Lenti-X^{TM}

Tet-One^{TM} Inducible Expression System digested with AgeI. Lentivirus was made for

both constructs by transfecting 293T cells with second generation packaging plasmids

and spinfected onto 293T cells at a low multiplicity of infection (MOI) as previously

described (Hartenian et al., 2020). 24 hr later, 350 μg/ml zeocin was added to select for

transduced cells. HEK293T and HeLa cells were grown in Dulbecco's Modified Eagle

Medium (ThermoFisher Scientific) supplemented with 10% fetal bovine serum (FBS)

and 1 U/mL penicillin-streptomycin (pen-strep). HCT116 cells were maintained in

McCoy's (modified) 5A medium (ThermoFisher Scientific) with 10% FBS and 1 U/mL

pen-strep. All cells were incubated at 37 ^{o}C with 5% CO_2. Cells were maintained in

culture in 10 cm^2 plates and 1x10^6 cells were seeded into 6 well plates for all

experiments except for chromatin immunoprecipitations, in which 5x10^6 cells were

seeded into 10 cm^2 plates, and 4sU-sequencing, in which 5x10^6 cells were seeded into

15 cm^2 plates.

siRNA nucleofections

Protein knockdowns were performed using the Neon Transfection System with

siRNA pools for the following targets: non-targeting control pool (scramble or scr

siRNA), CASP3, CASP8, DFFB, DIS3L2, EXOSC4, PNPT1, and XRN1. For all siRNA

pools, cells were nucleofected according to manufacturer protocols for HCT116 cells

and immediately seeded into plates containing media supplemented with 10% FBS but

lacking pen-strep to improve cell viability post-nucleofection. A 50 nM final siRNA

concentration was used for pools targeting CASP3, CASP8, and DFFB (CAD), while

200 nM was used for individual knockdown of XRN1, DIS3L2, EXOSC4, and PNPT1, as

well as the concurrent knockdowns of DIS3L2/EXOSC4/PNPT1 (66.7 nM each) and

BAX/BAK (100 nM each). For all experiments involving siRNA knockdowns, cells were

treated and/or harvested once 80-90% confluent in each plate or well, approximately 72

hr post-nucleofection, and protein knockdown was confirmed by western blot. Cell

populations of siRNA-transfected cells were split into two wells or plates 24 hr pre-

harvesting to allow for the direct comparison between apoptotic and non-apoptotic cells

in the same genetic background.

Apoptosis induction

100 ng/mL TNF-related apoptosis-inducing ligand (TRAIL) was used to induce

rapid extrinsic apoptosis in HCT116 cells. Where indicated, HCT116 cells were pre-

treated for 1 hr with 40 μM caspase Inhibitor Z-VAD-FMK (zVAD) or 25 μM ivermectin

before TRAIL treatment. Extrinsic apoptosis induction was confirmed by observing

CASP8 and CASP3 cleavage on western blots. CASP3/8 cleavage was quantified by

disappearance in intensity of the full-sized band normalized to TUBA1A or VCL using

Bio-Rad ImageLab software. Intrinsic apoptosis was induced in HeLa cells with a 4 hr

treatment of 10 μM raptinal, with or without 1 hr pre-treatment with 20 μM zVAD.

Induction was confirmed by evidence of cytochrome c release into the cytoplasm with

cell fractionation and western blot. "Mock" treatments consisted of an equal volume of

vehicle used to dissolve each reagent: TRAIL storage and dilution buffer, DMSO, and

ethanol for TRAIL, zVAD, and ivermectin, respectively.

RNA, DNA, and protein extractions

Total RNA, and protein extractions were performed according to manufacturer's

instructions after cells were harvested with Trizol^{TM} reagent. RNA pellets were dissolved

in DEPC water and protein pellets were dissolved in 1% SDS overnight at 50 ^{o}C before

spinning down insoluble material at 10,000 x g for 10 min.

TUNEL Assay

Terminal deoxynucleotidyl transferase Br-dUTP nick end labeling was performed

on TRAIL-treated cells using a TUNEL assay kit according to manufacturer protocol. Br-

dUTP incorporation was quantified by flow cytometry, analyzing the elevated peak in

FL2 fluorescence on a BD Accuri Flow Cytometry System using FlowJo analysis

software.

Quantitative reverse transcription PCR (RT-qPCR)

Extracted RNA was DNAse-treated treated, primed with random nonamers, and

reverse-transcribed to cDNA with Avian Myeloblastosis Virus Reverse Transcriptase

according to manufacturer protocols. Genes were quantified by RT-qPCR using iTaq

Universal SYBR Master Mix and primers specific to each gene of interest. RNA fold

change values were calculated in reference to *18S* or *U6* ncRNAs, as indicated on

figure axes.

Western blotting

Protein samples were quantified by Bradford assay according to manufacturer

instructions. 12.5-50 µg of protein was mixed with one third volume of 4X Laemelli

Sample Buffer (Bio-Rad Laboratories 1610747) and boiled at 100 ^{o}C before loading into

a polyacrylamide gel alongside either the PageRuler™ or PageRuler™ Plus Prestained

Protein Ladder. Proteins were separated by electrophoresis and transferred to a

nitrocellulose membrane, which was then cut into sections surrounding the size of the

protein of interest, allowing for multiple proteins to be quantified from one gel.

Membranes were then blocked with 5% non-fat dry milk in TBST (1X Tris-buffered

saline with 0.2% [v/v] Tween 20) at RT for 1 hour then incubated with relevant primary

antibodies diluted with 1% milk in TBST overnight at 4 $^{\circ}$C. All primary antibodies were

applied at a 1:1000 dilution with the exception of the following (target, dilution): RPB1,

1:500; TUBA1A, 1:500; RPB2, 1:500; RPB3, 1:10000; LMNB1, 1:1000; GAPDH,

1:5000; and CYTC, 1:500. After three five min TBST washes, species-specific

secondary antibodies were diluted 1:5000 with 1% milk in TBST and incubated for one

hour at RT. Blots were then developed, after three additional five min TBST washes,

with Clarity Western ECL Substrate for five min and imaged using a ChemiDoc^{TM} MP

Imaging System (Bio-Rad Laboratories). Each membrane section was imaged and

processed separately. Band intensity was quantified using Bio-Rad Image Lab software,

and relative expression changes were calculated after normalizing to an α-tubulin

(TUBA1A), vinculin (VCL), or lamin-B1 (LMNB1) loading control. For all blots that

appear in figures, auto-contrast was applied in Image Lab for each membrane section

before the resultant image was exported for publication. When appropriate, membrane

sections were stripped with 25 mM glycine in 1% SDS, pH 2 and washed 2 times for 10

min with TBST before being blocked and re-probed as previously described with a

primary antibody targeting a protein of similar size.

Antibody generation

Polyclonal rabbit anti-muSOX antibody was made and purified by YenZym

antibodies, LLC from recombinant muSOX protein.

Cell Fractionations

For experiments performed in apoptotic HCT116 and HeLa cells, nuclear,

cytoplasmic, and mitochondrial fractions were isolated using the Abcam Cell

Fractionation Kit according to manufacturer's instructions. Cytoplasmic and nuclear

fractions of samples in experiments that did not involve apoptosis were separated using

the REAP method (Suzuki, Bose, Leong-Quong, Fujita, & Riabowol, 2010). 1/5[th] of the

total cell lysate was reserved and diluted to the same volume of the cell and nuclear

fractions for whole cell lysate samples. Protein was extracted from 200 μL of each

fraction from both methods using Trizol[TM] LS reagent and analyzed by western blot as

described above.

4-thiouridine- (4sU)- pulse labeling

4sU-pulse labeling was used to measure nascent transcription concurrently with

mRNA decay. 50 μM 4sU was added to cells 20 min before harvesting lysates for RNA

and/or protein extraction. Labeled transcripts contained in 25 μg of total extracted RNA

were biotinylated as described by Dölken (2013), using 50 μg HPDP biotin. Biotinylated

RNA was conjugated to Dynabeads™ MyOne™ Streptavidin C1 magnetic beads for 1

hour in the dark, then the beads were washed four times (twice at 65 $^{\circ}$C and twice at

RT) with wash buffer (100 mM Tris-HCl, 10 mM EDTA, 1 M sodium chloride, and 0.1%

Tween 20) before eluting RNA off of the beads twice with 5% BME in DEPC water. RNA

was precipitated by adding 1/10[th] volume of 3M sodium acetate and 2.5 volumes of

ethanol and spun down at full speed in a 4°C benchtop centrifuge. After a 75% ethanol

wash, the 4sU-labeled RNA was resuspended in 20 μL DEPC water for use in in RT-

qPCR.

Reverse transcription PCR (RT-PCR)

4sU-labeled RNA from HCT116 cells treated with 100 ng/uL TRAIL, 10 μM

raptinal, or their respective mock treatments was isolated as previously described. 2 μL

4sU RNA from each sample was reverse transcribed into cDNA and amplified with

primers targeting regions of the 18S rRNA and/or U6 snRNA using the QIAGEN

OneStep RT-PCR Kit according to manufacturer instructions. 25 μL of each resultant

PCR product was combined with 5 μL 6X DNA loading dye and loaded onto a 1%

agarose (in 1X Tris-borate EDTA) electrophoresis gel stained with SYBR[TM] Safe DNA

Gel Stain alongside a DNA ladder. Gels were imaged on ChemiDoc[TM] MP Imaging

System.

4sU-sequencing (4sU-seq)

$5x10^6$ HCT116 cells were seeded onto 15 cm^2 plates and 24 hr later, were pre-

treated with either DMSO or 20 μM zVAD, and treated with storage and dilution buffer or

100 ng/μL TRAIL for 2 hr. This process was repeated to generate 2 biological replicates.

50 μM 4sU was added to cells 20 min before harvesting. Cells were suspended in 2 mL

Trizol and RNA extracted as previously described. Biotinylation and strepdavidin

selection was performed on 200 μg of total RNA, scaling up the previously detailed

protocol by 8X. 125 ng of 4sU-labeled RNA was used to synthesize rRNA-depleted

sequencing libraries using KAPA-stranded RNA-Seq Kit with Ribo-Erase, HMR

according to manufacturer's instructions. ERCC RNA Spike-in Mix 1 was added at a

1:100 dilution to each RNA sample immediately prior to library preparation normalize

read counts to RNA input across samples. Libraries were submitted for analysis on a

Bioanalyzer to ensure ~400 bp fragment lengths, then submitted for sequencing on a

Nova-Seq 6000 with 100 bp paired-end reads at the QB3-Berkeley Genomics

Sequencing Core.

Bioinformatics analysis was conducted using the UC Berkeley High Performance

Computing Cluster. Paired end sequence FASTQ files were downloaded and checked

for quality using FastQC. Reads were then trimmed of adaptors using Sickle/1.33.

Reads were mapped to human reference genome hg19 and ERCC spike-in list was

obtained using STAR genome aligner (Dobin et al., 2013). Differential expression upon

TRAIL treatment for each gene were calculated using Cuffdiff 2 (Trapnell et al., 2013)

on samples in the DMSO condition with their replicates, and on zVAD samples and their

replicates. Differential expression values for each gene were normalized to ERCC

spike-in controls. Normalized fold change values were calculated and analyzed using

Microsoft Excel. Statistically significant >2-fold upregulated genes upon TRAIL

treatment in the DMSO condition were input into PANTHER-GO Slim gene ontology

analysis (Mi, Muruganujan, Ebert, Huang, & Thomas, 2019) and ChEA3 transcription

factor enrichment analysis (Keenan et al., 2019).

Chromatin immunoprecipitation (ChIP)

HCT116 cells nucleofected with the relevant siRNAs were seeded onto 10 cm^2

plates. 72 hr post-nucleofection, cells were washed with PBS, trypsinized, washed with

PBS again, and fixed in 1% formaldehyde for 2.5 min before quenching with 125 mM

glycine. After an additional PBS wash, cells were lysed for 10 min at 4^{o}C in lysis buffer

(50 mM HEPES pH 7.9, 140 mM NaCl, 1 mM EDTA, 10% [v/v] glycerol, 0.5% NP40,

0.25% Triton X-100) then washed with wash buffer (10 mM Tris Cl pH 8.1, 100 mM

NaCl, 1 mM EDTA pH 8.0) at 4^{o}C for 10 min. Cells were then suspended in 1 mL

shearing buffer (50 mM Tris Cl pH 7.5, 10 mM EDTA, 0.1% [v/v] SDS) and sonicated in

a Covaris S220 sonicator (Covaris, Inc.) with the following parameters: peak power,

140.0; duty factor, 5.0; cycle/burst, 200; and duration, 300 sec. Insoluble material in the

shearing buffer was then spun down at full speed in a 4 $^{\circ}$C benchtop centrifuge to yield

the chromatin supernatant. 10 µg chromatin was rotated overnight at 4 $^{\circ}$C with 2.5 µg or

4 µg ChIP-grade primary antibodies targeting RPB1 and TBP, respectively, in 500 µL

dilution buffer (1.1% [v/v] Triton-X-100, 1.2mM EDTA, 6.7mM Tris-HCl pH 8.0, 167mM

NaCl). 5 µL of each IP was reserved as an input sample before antibody was added. 20

µL of either Dynabeads™ Protein G or a 1:1 mixture Protein A and Protein G beads, for

anti-mouse and anti-rabbit antibodies, respectively, were added to each reaction and

rotated at 4 $^{\circ}$C for at least 2 hr. The beads were then sequentially washed with low salt

immune complex wash buffer (0.1% [v/v] SDS, 1% [v/v] Triton-X-100, 2 mM EDTA, 20

mM Tris-HCl pH 8.0, 150 mM NaCl), high low salt immune complex wash buffer (0.1%

[v/v] SDS, 1% [v/v] Triton-X-100, 2 mM EDTA, 20 mM Tris-HCl pH 8, 500 mM NaCl),

LiCl immune complex buffer (0.25 M LiCl, 1% [v/v] NP40, 1% [v/v] deoxycholic acid,

1mM EDTA, 10mM Tris-HCl pH 8.0), and TE buffer (10 mM Tris-HCl pH 8.0, 1 mM

EDTA). All washes were 5 min in duration and performed at 4 $^{\circ}$C. Beads and input

samples were then suspended in 100 µL elution buffer (150 mM NaCl, 50ug/ml

proteinase K) and incubated at 55 $^{\circ}$C for 2 hr then at 65 $^{\circ}$C for 12 hr in a thermal cycler.

DNA fragments were purified with an oligonucleotide clean and concentrator kit and %

input values were quantified by RT-qPCR as previously described using primers

complementary to the locus of interest.

Data visualization

Bar graphs were created using GraphPad Prism 8 software and the graphical abstract

was created using the Bio-Render online platform.

Quantification and statistical analysis

Biological replicates were defined as experiments performed separately on

biologically distinct (i.e. from cells cultured at different times in different flasks or wells)

samples representing identical conditions and/or time points. See figures and figure

legends for the number of biological replicates performed for each experiment. Criteria

for the inclusion of data was based on the performance of positive and negative controls

within each experiment. No outliers were eliminated in this study. One-sample t-tests

were performed on control and experimental groups for which mean fold change values

were calculated, comparing these values to the null hypothesis of 1. Student's T tests

(corrected for multiple comparisons with the Holm-Sidak method when appropriate)

were also performed comparing means between control and experimental groups,

signified by brackets spanning the two groups being compared. All statistical analyses

were performed using GraphPad Prism 8 unless otherwise noted.

Key Resources Table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Anti-RNA polymerase II CTD repeat YSPTSPS antibody (clone 8WG16)	Abcam	Cat#ab817
Anti-RNA polymerase II CTD repeat YSPTSPS (phospho S2) antibody	Abcam	Cat#ab5095
Anti-RNA polymerase II CTD repeat YSPTSPS (phospho S5) antibody	Abcam	Cat#ab5131
Anti-alpha Tubulin antibody	Abcam	Cat#ab7291
POLR2B (RPB2) Antibody (clone E-12)	Santa Cruz Biotechnology	Cat#sc-166803
Anti-RPB3 antibody (clone EPR13294B)	Abcam	Cat#ab182150
POLR2D (RPB4) Polyclonal Antibody	ThermoFisher Scientific	Cat#PA5-35954
Anti-Vinculin antibody	Abcam	Cat#ab91459
Anti-PNPT1 antibody	Abcam	Cat#ab96176

Anti-RRP41 (EXOSC4) antibody	Abcam	Cat#ab137250
DIS3L2 Antibody	Novus Biologicals	Cat#NBP184740
Anti-Lamin B1 antibody	Abcam	Cat# ab133741
Anti-GAPDH antibody (clone 6C5)	Abcam	Cat# ab8245
Caspase-8 (clone D35G2) Rabbit mAb	Cell Signaling Technology	Cat#4790
Caspase-3 Antibody	Cell Signaling Technology	Cat#9662
Bak Antibody	Cell Signaling Technology	Cat# 3814S
Bax Antibody	Cell Signaling Technology	Cat#2772S
XRN1 Antibody	Bethyl Laboratories	Cat#A400-433A
Anti-DFFB antibody	Abcam	Cat#ab69438
PABP1 Antibody	Cell Signaling Technology	Cat#4992
PABP4 Antibody	Bethyl Laboratories	Cat#A301-466A
C23 (NCL) Antibody (clone MS-3)	Santa Cruz Biotechnology	Cat#sc-8031
PARP Antibody	Cell Signaling Technology	Cat#9542
BID Antibody (clone 5C9)	Santa Cruz Biotechnology	Cat#sc-56025
Anti-TATA binding protein TBP antibody (clone mAbcam 51841)	Abcam	Cat#ab51841
Cytochrome c (clone D18C7) Rabbit mAb	Cell Signaling Technology	Cat#11940
Anti-VDAC1 / Porin antibody	Abcam	Cat# ab15895
Polyclonal rabbit ant-muSOX antibody	This paper	N/A
Chemicals, Peptides, and Recombinant Proteins		
Trizol™ Reagent	ThermoFisher Scientific	Cat#15596026
Trizol™ LS Reagent	ThermoFisher Scientific	Cat#10296028
TURBO™ DNase	ThermoFisher Scientific	Cat#AM2238
Avian Myeloblastosis Virus Reverse Transcriptase	Promega Corporation	Cat#M5108
iTaq Universal SYBR Master Mix	Bio-Rad Laboratories	Cat#1725122
Dynabeads™ Protein G	ThermoFisher Scientific	Cat#10003D
Dynabeads™ Protein A	ThermoFisher Scientific	Cat#10002D
Dynabeads® MyOne™ Streptavidin C1	ThermoFisher Scientific	Cat#
EZ-link™ HPDP-biotin	ThermoFisher Scientific	Cat#21341
SuperKillerTRAIL® Protein (soluble) (human), (recombinant)	Enzo Life Sciences	Cat# ALX-201-115-3010
KillerTRAIL™ Storage and Dilution Buffer	Enzo Life Sciences	Cat# ALX-505-005-R500
Caspase Inhibitor Z-VAD-FMK	Promega Corporation	Cat#G7231
Ivermectin	Millipore Sigma	Cat#I8898
Raptinal	Millipore Sigma	Cat#SML1745
Dulbecco's Modified Eagle Medium	ThermoFisher Scientific	Cat#12800082
McCoy's (modified) 5A medium	ThermoFisher Scientific	Cat#16600082
Fetal Bovine Serum	VWR	Cat#89510-186
Trypsin-EDTA (0.05%), phenol red	ThermoFisher Scientific	Cat# 25300120
PageRuler™ Prestained Protein Ladder	ThermoFisher Scientific	Cat#26616
PageRuler™ Plus Prestained Protein Ladder	ThermoFisher Scientific	Cat#26620

Quick-Load® Purple 1 kb Plus DNA Ladder	New England BioLabs	Cat#N0550S
Clarity Western ECL Substrate	Bio-Rad Laboratories	Cat#1705061
4x Laemmli Sample Buffer	Bio-Rad Laboratories	Cat#1610747
Gel Loading Dye, Purple (6X), no SDS	New England BioLabs	B7025S
Critical Commercial Assays		
TUNEL Assay Kit - BrdU-Red	Abcam	Cat#ab66110
OneStep RT-PCR Kit	QIAGEN	Cat#210210
Cell Fractionation Kit	Abcam	Cat#ab109719
Bio-Rad Protein Assay Kit II	Bio-Rad Laboratories	Cat#5000002
Oligo Clean & Concentrator Kit	Zymo Research	Cat#D4060
In-Fusion® HD Cloning Kit	Takara Bio USA	Cat#639650
Lenti-X™ Tet-On® 3G Inducible Expression System	Takara Bio USA	Cat#631187
Neon Transfection System	ThermoFisher Scientific	Cat#MPK5000
KAPA Stranded RNA-Seq Kit with RiboErase (HMR)	Roche	Cat#KK8484
ERCC RNA Spike-In Mix	ThermoFisher Scientific	Cat# 4456740
Experimental Models: Cell Lines		
Human: HCT116 cells	ATCC	CCL-247
Human: 293T/17 cells	ATCC	CRL-11268
Human: HeLa Cells	ATCC	CCL-2
Oligonucleotides		
muSOX F	This paper	TCCCGTATACACCGGTATGTGGAGCCACCCC
muSOX R	This paper	ATCCGCCGGCACCGGTTTAGGGGGTTATGGG
ON-TARGETplus Non-targeting Control Pool	Horizon Discovery Group	Cat#D-001810-10
SMARTpool: ON-TARGETplus DIS3L2 siRNA	Horizon Discovery Group	Cat#L-018715-01
SMARTpool: ON-TARGETplus Human EXOSC4 siRNA	Horizon Discovery Group	Cat#L-013760-00
SMARTpool: ON-TARGETplus Human PNPT1 siRNA	Horizon Discovery Group	Cat#L-019454-01
SMARTpool: ON-TARGETplus XRN1 siRNA	Horizon Discovery Group	Cat#L-013754-01
SMARTpool: ON-TARGETplus CASP3 siRNA	Horizon Discovery Group	Cat#L-004307-00
SMARTpool: ON-TARGETplus CASP8 siRNA	Horizon Discovery Group	Cat#L-003466-00
ON-TARGETplus DFFB siRNA SMARTpool	Horizon Discovery Group	Cat#L-004425-00
SMARTpool: ON-TARGETplus Human BAX siRNA	Horizon Discovery Group	Cat# L-003308-01

SMARTpool: ON-TARGETplus Human BAK1 siRNA	Horizon Discovery Group	Cat# L-003305-00
See Supplementary Table 7 for RT-(q)PCR primers		
Software and Algorithms		
Prism 8	GraphPad	https://www.graphpad.com/scientific-software/prism/
FlowJo	BD	https://www.flowjo.com/solutions/flowjo
Image Lab Software	Bio-Rad Laboratories	Cat#1709690
FastQC	Babraham Bioinformatics	http://www.bioinformatics.babraham.ac.uk/projects/fastqc
Sickle version 1.33	N/A	https://github.com/najoshi/sickle
STAR	Dobin et al. 2012	https://doi.org/10.1093/bioinformatics/bts635
Cuffdiff 2	Trapnell et al. 2013	https://doi.org/10.1038/nbt.2450
PANTHER GO-slim	Muruganujan et al. 2019	http://pantherdb.org/
ChEA3	Keenan et al. 2019	https://maayanlab.cloud/chea3/

REFERENCES

Abernathy, E., Clyde, K., Yeasmin, R., Krug, L. T., Burlingame, A., Coscoy, L., &

Glaunsinger, B. (2014). Gammaherpesviral Gene Expression and Virion

Composition Are Broadly Controlled by Accelerated mRNA Degradation. *PLoS*

Pathogens, *10*(1), e1003882. https://doi.org/10.1371/journal.ppat.1003882

Abernathy, E., Gilbertson, S., Alla, R., & Glaunsinger, B. (2015). Viral nucleases induce

an mRNA degradation-transcription feedback loop in mammalian cells. *Cell Host*

and Microbe, *18*(2), 243–253. https://doi.org/10.1016/j.chom.2015.06.019

Albeck, J. G., Burke, J. M., Aldridge, B. B., Zhang, M., Lauffenburger, D. A., & Sorger,

P. K. (2008). Quantitative Analysis of Pathways Controlling Extrinsic Apoptosis in

Single Cells. *Molecular Cell*, *30*(1), 11–25.

https://doi.org/10.1016/j.molcel.2008.02.012

Baumann, S., Hess, J., Eichhorst, S. T., Krueger, A., Angel, P., Krammer, P. H., &

Kirchhoff, S. (2003). An unexpected role for FosB in activation-induced cell death of

T cells. *Oncogene*, *22*(9), 1333–1339. https://doi.org/10.1038/sj.onc.1206126

Beltran, M., Yates, C. M., Skalska, L., Dawson, M., Reis, F. P., Viiri, K., … Jenner, R. G.

(2016). The interaction of PRC2 with RNA or chromatin is mutually antagonistic.

Genome Research, *26*(7), 896–907. https://doi.org/10.1101/gr.197632.115

Bentley, D. L. (2014). Coupling mRNA processing with transcription in time and space.

Nature Reviews Genetics, *15*(3), 163–175. https://doi.org/10.1038/nrg3662

Braun, K. A., & Young, E. T. (2014). Coupling mRNA Synthesis and Decay. *Molecular*

and Cellular Biology, *34*(22), 4078–4087. https://doi.org/10.1128/MCB.00535-14

Bregman, A., Avraham-kelbert, M., Barkai, O., Duek, L., Guterman, A., & Choder, M.

(2011). Promoter Elements Regulate Cytoplasmic mRNA Decay. *Cell*, *147*(7),

1473–1483. https://doi.org/10.1016/j.cell.2011.12.005

Buratowski, S., Hahn, S., Guarente, L., & Sharp, P. A. (1989). Five intermediate

complexes in transcription initiation by RNA polymerase II. *Cell*, *56*(4), 549–561.

https://doi.org/10.1016/0092-8674(89)90578-3

Caruso, S., & Poon, I. K. H. (2018). Apoptotic cell-derived extracellular vesicles: More

than just debris. *Frontiers in Immunology*, *9*, 1486.

https://doi.org/10.3389/fimmu.2018.01486

Collart, M. A., & Stmhp, K. (1994). *NOTI (CDC39), NOT2 (CDC36), NOT3 , and N0T4*

encode a global-negative regulator of transcription that differentially affects TATA-

element utilization. 2, 525–537.

Dantonel, J. C., Murthy, K. G. K., Manjey, J. L., & Tora, L. (1997). Transcription factor

TFIID recruits factor CPSF for formation of 3 end of mRNA. *Nature, 389*(6649),

399–402. https://doi.org/10.1038/38763

Denis, C. L. (1984). Identification of new genes involved in the regulation of yeast

alcohol dehydrogenase II. *Genetics, 108*(4), 833–844.

Di Ruscio, A., Ebralidze, A. K., Benoukraf, T., Amabile, G., Goff, L. A., Terragni, J., …

Tenen, D. G. (2013). DNMT1-interacting RNAs block gene-specific DNA

methylation. *Nature, 503*(7476), 371–376. https://doi.org/10.1038/nature12598

Dieci, G., Preti, M., & Montanini, B. (2009). Eukaryotic snoRNAs: A paradigm for gene

expression flexibility. *Genomics, 94*(2), 83–88.

https://doi.org/10.1016/j.ygeno.2009.05.002

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., … Gingeras,

T. R. (2013). STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics, 29*(1), 15–

21. https://doi.org/10.1093/bioinformatics/bts635

Dölken, L. (2013). High Resolution Gene Expression Profiling of RNA Synthesis,

Processing, and Decay by Metabolic Labeling of Newly Transcribed RNA Using 4-

Thiouridine. In S. M. Bailer & D. Lieber (Eds.), *Virus-Host Interactions: Methods*

and Protocols (pp. 91–100). https://doi.org/10.1007/978-1-62703-601-6_6

Düring, L., Thorsen, M., Petersen, D. S. N., Køster, B., Jensen, T. H., & Holmberg, S.

(2012). MRN1 Implicates Chromatin Remodeling Complexes and Architectural

Factors in mRNA Maturation. *PLoS ONE, 7*(9), 1–13.

https://doi.org/10.1371/journal.pone.0044373

El-Brolosy, M. A., Kontarakis, Z., Rossi, A., Kuenne, C., Günther, S., Fukuda, N., …

Stainier, D. Y. R. (2019). Genetic compensation triggered by mutant mRNA

degradation. *Nature*, *568*(7751), 193–197. https://doi.org/10.1038/s41586-019-

1064-z

Enari, M., Sakahira, H., Yokoyama, H., Okawa, K., Iwamatsu, A., & Nagata, S. (1998).

Cleavage of CAD inhibitor in CAD activation and DNA degradation during

apoptosis. *Nature*, *391*(6662), 43–50.

Fedorova, O., Petukhov, A., Daks, A., Shuvalov, O., Leonova, T., Vasileva, E., …

Barlev, N. A. (2019). Orphan receptor NR4A3 is a novel target of p53 that

contributes to apoptosis. *Oncogene*, *38*(12), 2108–2122.

https://doi.org/10.1038/s41388-018-0566-8

Garland, W., Comet, I., Wu, M., Radzisheuskaya, A., Rib, L., Vitting-Seerup, K., …

Jensen, T. H. (2019). A Functional Link between Nuclear RNA Decay and

Transcriptional Control Mediated by the Polycomb Repressive Complex 2. *Cell*

Reports, *29*(7), 1800-1811.e6. https://doi.org/10.1016/j.celrep.2019.10.011

Gilbertson, S., Federspiel, J. D., Hartenian, E., Cristea, I. M., & Glaunsinger, B. (2018).

Changes in mRNA abundance drive shuttling of RNA binding proteins, linking

cytoplasmic RNA degradation to transcription. *ELife*, *7*, 1–26.

https://doi.org/10.7554/eLife.37663

Gö, D. (1998). MEDAL REVIEW Transport into and out of the cell nucleus. *The EMBO*

Journal, *17*(10), 2721–2727.

Guerin, S., Baron, M. L., Valero, R., Herrant, M., Auberger, P., & Naquet, P. (2002).

RelB reduces thymocyte apoptosis and regulates terminal thymocyte maturation.

European Journal of Immunology, *32*(1), 1–9. https://doi.org/10.1002/1521-

4141(200201)32:1<1::AID-IMMU1>3.0.CO;2-S

Gurzov, E. N., Ortis, F., Bakiri, L., Wagner, E. F., & Eizirik, D. L. (2008). JunB inhibits

ER stress and apoptosis in pancreatic beta cells. *PLoS ONE*, *3*(8).

https://doi.org/10.1371/journal.pone.0003030

Haimovich, G., Medina, D. A., Causse, S. Z., Garber, M., Millán-Zambrano, G., Barkai,

O., … Choder, M. (2013). Gene expression is circular: Factors for mRNA

degradation also foster mRNA synthesis. *Cell*, *153*(5).

https://doi.org/10.1016/j.cell.2013.05.012

Hartenian, E., Gilbertson, S., Federspiel, J. D., Cristea, I. M., & Glaunsinger, B. A.

(2020). RNA decay during gammaherpesvirus infection reduces RNA polymerase II

occupancy of host promoters but spares viral promoters. *PLoS Pathogens*, *16*(2),

e1008269–e1008269. https://doi.org/10.1371/journal.ppat.1008269

Hartenian, E., & Glaunsinger, B. A. (2019). Feedback to the central dogma: cytoplasmic

mRNA decay and transcription are interdependent processes. *Critical Reviews in*

Biochemistry and Molecular Biology, *54*(4), 385–398.

https://doi.org/10.1080/10409238.2019.1679083

Harwig, A., Landick, R., & Berkhout, B. (2017). The battle of RNA synthesis: Virus

versus host. *Viruses*, *9*(10), 1–18. https://doi.org/10.3390/v9100309

Heimer, S., Knoll, G., Schulze-Osthoff, K., & Ehrenschwender, M. (2019). Raptinal

bypasses BAX, BAK, and BOK for mitochondrial outer membrane permeabilization

and intrinsic apoptosis. *Cell Death and Disease*, *10*(8).

https://doi.org/10.1038/s41419-019-1790-z

Helenius, K., Yang, Y., Tselykh, T. V., Pessa, H. K. J., Frilander, M. J., & Mäkelä, T. P. (2011). Requirement of TFIIH kinase subunit Mat1 for RNA Pol II C-terminal domain Ser5 phosphorylation, transcription and mRNA turnover. *Nucleic Acids Research*, *39*(12), 5025–5035. https://doi.org/10.1093/nar/gkr107

Houge, G., Robaye, B., Eikhom, T. S., Golstein, J., Mellgren, G., Gjertsen, B. T., … Døskeland, S. O. (1995). Fine mapping of 28S rRNA sites specifically cleaved in cells undergoing apoptosis. *Molecular and Cellular Biology*, *15*(4), 2051–2062. https://doi.org/10.1128/MCB.15.4.2051

Houseley, J., & Tollervey, D. (2009). The Many Pathways of RNA Degradation. *Cell*, *136*(4), 763–776. https://doi.org/10.1016/j.cell.2009.01.019

Huang, X., Li, X., & Guo, B. (2008). KLF6 induces apoptosis in prostate cancer cells through up-regulation of ATF3. *Journal of Biological Chemistry*, *283*(44), 29795–29801. https://doi.org/10.1074/jbc.M802515200

Jones, C. I., Zabolotskaya, M. V., & Newbury, S. F. (2012). The 5′ → 3′ exoribonuclease XRN1/Pacman and its functions in cellular processes and development. *Wiley Interdisciplinary Reviews: RNA*, *3*(4), 455–468. https://doi.org/10.1002/wrna.1109

Keenan, A. B., Torre, D., Lachmann, A., Leong, A. K., Wojciechowicz, M. L., Utti, V., … Ma'ayan, A. (2019). ChEA3: transcription factor enrichment analysis by orthogonal omics integration. *Nucleic Acids Research*, *47*(W1), W212–W224. https://doi.org/10.1093/nar/gkz446

Keller, U., Huber, J., Nilsson, J. A., Fallahi, M., Hall, M. A., Peschel, C., & Cleveland, J. L. (2010). Myc suppression of Nfkb2 accelerates lymphomagenesis. *BMC Cancer*, *10*. https://doi.org/10.1186/1471-2407-10-348

Kim, Y. M., Kim, T. H., Chung, H. T., Talanian, R. V., Yin, X. M., & Billiar, T. R. (2000). Nitric oxide prevents tumor necrosis factor α-induced rat hepatocyte apoptosis by the interruption of mitochondrial apoptotic signaling through S-nitrosylation of caspase-8. *Hepatology*, *32*(4 I), 770–778. https://doi.org/10.1053/jhep.2000.18291

Kimura, M., Kose, S., Okumura, N., Imai, K., Furuta, M., Sakiyama, N., … Imamoto, N. (2013). Identification of cargo proteins specific for the nucleocytoplasmic transport carrier transportin by combination of an in vitro transport system and stable isotope labeling by amino acids in cell culture (SILAC)-based quantitative proteomics. *Molecular and Cellular Proteomics*, *12*(1), 145–157. https://doi.org/10.1074/mcp.M112.019414

Kumar, G. R., & Glaunsinger, B. A. (2010). Nuclear Import of Cytoplasmic Poly(A) Binding Protein Restricts Gene Expression via Hyperadenylation and Nuclear Retention of mRNA. *Molecular and Cellular Biology*, *30*(21), 4996–5008. https://doi.org/10.1128/MCB.00600-10

Kumar, G. R., Shum, L., & Glaunsinger, B. A. (2011). Importin -Mediated Nuclear Import of Cytoplasmic Poly(A) Binding Protein Occurs as a Direct Consequence of Cytoplasmic mRNA Depletion. *Molecular and Cellular Biology*, *31*(15), 3113–3125. https://doi.org/10.1128/MCB.05402-11

Łabno, A., Tomecki, R., & Dziembowski, A. (2016). Cytoplasmic RNA decay pathways - Enzymes and mechanisms. *Biochimica et Biophysica Acta - Molecular Cell Research*, *1863*(12), 3125–3147. https://doi.org/10.1016/j.bbamcr.2016.09.023

Lee, Y. J., & Glaunsinger, B. A. (2009). Aberrant herpesvirus-induced polyadenylation correlates with cellular messenger RNA destruction. *PLoS Biology*, *7*(5), e1000107.

https://doi.org/10.1371/journal.pbio.1000107

Liu, X., Fu, R., Pan, Y., Meza-sosa, K. F., Zhang, Z., Liu, X., … Lieberman, J. (2018).

PNPT1 Release from Mitochondria during Apoptosis Triggers Decay of Poly(A)

RNAs. *Cell*, 1–15. https://doi.org/10.1016/j.cell.2018.04.017

Lotan, R., Bar-On, V. G., Harel-Sharvit, L., Duek, L., Melamed, D., & Choder, M. (2005).

The RNA polymerase II subunit Rpb4p mediates decay of a specific class of

mRNAs. *Genes and Development*, *19*(24), 3004–3016.

https://doi.org/10.1101/gad.353205

Lotan, R., Goler-Baron, V., Duek, L., Haimovich, G., & Choder, M. (2007). The Rpb7p

subunit of yeast RNA polymerase II plays roles in the two major cytoplasmic mRNA

decay mechanisms. *Journal of Cell Biology*, *178*(7), 1133–1143.

https://doi.org/10.1083/jcb.200701165

Louder, R. K., He, Y., López-Blanco, J. R., Fang, J., Chacón, P., & Nogales, E. (2016).

Structure of promoter-bound TFIID and model of human pre-initiation complex

assembly. *Nature*, *531*(7596), 604–609. https://doi.org/10.1038/nature17394

Lu, Y., Luo, Z., & Bregman, D. B. (2002). Rna polymerase II large subunit is cleaved by

caspases during DNA damage-induced apoptosis. *Biochemical and Biophysical*

Research Communications, *296*(4), 954–961. https://doi.org/10.1016/S0006-

291X(02)02028-4

Ma, Z., Zhu, P., Shi, H., Guo, L., Zhang, Q., Chen, Y., … Chen, J. (2019). PTC-bearing

mRNA elicits a genetic compensation response via Upf3a and COMPASS

components. *Nature*, *568*(7751), 259–263. https://doi.org/10.1038/s41586-019-

1057-y

Marchese, F. P., Raimondi, I., & Huarte, M. (2017). The multidimensional mechanisms

of long noncoding RNA function. *Genome Biology*, *18*(1), 1–13.

https://doi.org/10.1186/s13059-017-1348-2

Mi, H., Muruganujan, A., Ebert, D., Huang, X., & Thomas, P. D. (2019). PANTHER

version 14: More genomes, a new PANTHER GO-slim and improvements in

enrichment analysis tools. *Nucleic Acids Research*, *47*(D1), D419–D426.

https://doi.org/10.1093/nar/gky1038

Nagata, S., Hanayama, R., & Kawane, K. (2010). Autoimmunity and the Clearance of

Dead Cells. *Cell*, *140*(5), 619–630. https://doi.org/10.1016/j.cell.2010.02.014

Narayanan, K., & Makino, S. (2013). Interplay between viruses and host mRNA

degradation. *Biochimica et Biophysica Acta - Gene Regulatory Mechanisms*,

1829(6–7), 732–741. https://doi.org/10.1016/j.bbagrm.2012.12.003

Noonberg, S. B., Scott, G. K., & Benz, C. C. (1996). Evidence of post-transcriptional

regulation of U6 small nuclear RNA. *Journal of Biological Chemistry*, *271*(18),

10477–10481. https://doi.org/10.1074/jbc.271.18.10477

Palchaudhuri, R., Lambrecht, M. J., Botham, R. C., Partlow, K. C., van Ham, T. J., Putt,

K. S., … Hergenrother, P. J. (2015). A Small Molecule that Induces Intrinsic

Pathway Apoptosis with Unparalleled Speed. *Cell Reports*, *13*(9), 2027–2036.

https://doi.org/10.1016/j.celrep.2015.10.042

Pignatelli, M., Luna-Medina, R., Pérez-Rendón, A., Santos, A., & Perez-Castillo, A.

(2003). The transcription factor early growth response factor-1 (EGR-1) promotes

apoptosis of neuroblastoma cells. *Biochemical Journal*, *373*(3), 739–746.

https://doi.org/10.1042/BJ20021918

Preston, G. A., Lyon, T. T., Yin, Y., Lang, J. E., Solomon, G., Annab, L., … Barrett, J. C.

(1996). Induction of apoptosis by c-Fos protein . These include☐: Induction of

Apoptosis by c-Fos Protein. *American Society for Microbiology*, *16*(1), 211–218.

Qiu, Z., Khairallah, C., Romanov, G., & Sheridan, B. S. (2020). Cutting Edge: Batf3

Expression by CD8 T Cells Critically Regulates the Development of Memory

Populations . *The Journal of Immunology*, *205*(4), 901–906.

https://doi.org/10.4049/jimmunol.2000228

Rivas, H. G., Schmaling, S. K., & Gaglia, M. M. (2016). Shutoff of host gene expression

in influenza A virus and herpesviruses: Similar mechanisms and common themes.

Viruses, *8*(4), 1–26. https://doi.org/10.3390/v8040102

Shandilya, J., & Roberts, S. G. E. (2012). The transcription cycle in eukaryotes: From

productive initiation to RNA polymerase II recycling. *Biochimica et Biophysica Acta*

- Gene Regulatory Mechanisms, *1819*(5), 391–400.

https://doi.org/10.1016/j.bbagrm.2012.01.010

Singh, P., James, R. S., Mee, C. J., Morozov, I. Y., Singh, P., James, R. S., …

Morozov, I. Y. (2019). mRNA levels are buffered upon knockdown of RNA decay

and translation factors via adjustment of transcription rates in human HepG2 cells

adjustment of transcription rates in human HepG2 cells. *RNA Biology*, *16*(9), 1147–

1155. https://doi.org/10.1080/15476286.2019.1621121

Stewart, M. (2007). Molecular mechanism of the nuclear protein import cycle. *Nature*

Reviews Molecular Cell Biology, *8*(3), 195–208. https://doi.org/10.1038/nrm2114

Suffert, G., Malterer, G., Hausser, J., Viiliäinen, J., Fender, A., Contrant, M., … Pfeffer,

S. (2011). Kaposi's sarcoma herpesvirus microRNAs target caspase 3 and regulate

apoptosis. *PLoS Pathogens*, *7*(12), e1002405.

https://doi.org/10.1371/journal.ppat.1002405

Sun, M., Schulz, D., Pirkl, N., Etzold, S., Larivie, L., Maier, K. C., … Cramer, P. (2012).

Comparative dynamic transcriptome analysis (cDTA) reveals mutual feedback

between mRNA synthesis and degradation. *Genome Research*, *22*(7), 1350–1359.

https://doi.org/10.1101/gr.130161.111.1350

Suzuki, K., Bose, P., Leong-Quong, R. Y., Fujita, D. J., & Riabowol, K. (2010). REAP: A

two minute cell fractionation method. *BMC Research Notes*, *3*, 294.

https://doi.org/10.1186/1756-0500-3-294

Tabtieng, T., Degterev, A., & Gaglia, M. M. (2018). Caspase-Dependent Suppression of

Type I Interferon Signaling. *Virology*, *92*(10), 1–17.

Thomas, M. P., Liu, X., Whangbo, J., McCrossan, G., Sanborn, K. B., Basar, E., …

Lieberman, J. (2015). Apoptosis Triggers Specific, Rapid, and Global mRNA Decay

with 3' Uridylated Intermediates Degraded by DIS3L2. *Cell Reports*, *11*(7), 1079–

1089. https://doi.org/10.1016/j.celrep.2015.04.026

Trapnell, C., Hendrickson, D. G., Sauvageau, M., Goff, L., Rinn, J. L., & Pachter, L.

(2013). Differential analysis of gene regulation at transcript resolution with RNA-

seq. *Nature Biotechnology*, *31*(1), 46–53. https://doi.org/10.1038/nbt.2450

Tucker, M., Valencia-Sanchez, M. A., Staples, R. R., Chen, J., Denis, C. L., & Parker,

R. (2001). The transcription factor associated Ccr4 and Caf1 proteins are

components of the major cytoplasmic mRNA deadenylase in Saccharomyces

cerevisiae. *Cell*, *104*(3), 377–386. https://doi.org/10.1016/S0092-8674(01)00225-2

Wagstaff, K. M., Sivakumaran, H., Heaton, S. M., Harrich, D., & Jans, D. A. (2012).

Ivermectin is a specific inhibitor of importin α/β-mediated nuclear import able to inhibit replication of HIV-1 and dengue virus. *Biochemical Journal*, *443*(3), 851–856. https://doi.org/10.1042/BJ20120150

Walker, A. P., & Fodor, E. (2019). Interplay between Influenza Virus and the Host RNA Polymerase II Transcriptional Machinery. *Trends in Microbiology*, *27*(5), 398–407. https://doi.org/10.1016/j.tim.2018.12.013

Walsh, D., Mathews, M. B., & Mohr, I. (2013). Tinkering with Translation: Protein Synthesis. *Cold Spring Harbor Perspectives in Biology*, *5*(1), 1–30. https://doi.org/10.1101/cshperspect.a012351

Walsh, J. G., Cullen, S. P., Sheridan, C., Lüthi, A. U., Gerner, C., & Martin, S. J. (2008). Executioner caspase-3 and caspase-7 are functionally distinct proteases. *Proceedings of the National Academy of Sciences of the United States of America*, *105*(35), 12815–12819. https://doi.org/10.1073/pnas.0707715105

Wilkinson, M. F. (2019). Genetic paradox explained by nonsense (NITC: nonsense-induced transcriptional compentsation). *Nature*, *568*(7751), 179–180. https://doi.org/10.1038/d41586-019-00823-5

Wisdom, R., Johnson, R. S., & Moore, C. (1999). c-Jun regulates cell cycle progression and apoptosis by distinct mechanisms. *EMBO Journal*, *18*(1), 188–197. https://doi.org/10.1093/emboj/18.1.188

Ye, X., Lin, J., Lin, Z., Xue, A., Li, L., Zhao, Z., … Cong, B. (2017). Axin1 up-regulated 1 accelerates stress-induced cardiomyocytes apoptosis through activating Wnt/β-catenin signaling. *Experimental Cell Research*, *359*(2), 441–448. https://doi.org/10.1016/j.yexcr.2017.08.027

Zhang, A., Hildreth, R. L., & Colberg-Poley, A. M. (2013). Human Cytomegalovirus

Inhibits Apoptosis by Proteasome-Mediated Degradation of Bax at Endoplasmic

Reticulum-Mitochondrion Contacts. *Journal of Virology*, *87*(10), 5657–5668.

https://doi.org/10.1128/jvi.00145-13

Zhou, X., Jiang, W., Liu, Z., Liu, S., & Liang, X. (2017). Virus infection and death

receptor-mediated apoptosis. *Viruses*, *9*(11), 316. https://doi.org/10.3390/v9110316

1201

A

Ligand binding death receptor

BID → tBID

proCASP8 ("initiator") → CASP8*

CASP8* → tBID

tBID → MOMP

MOMP → mRNA decay

proCASP3 ("executioner") → CASP3*

CASP3* → **APOPTOSIS**

B

TRAIL (hrs): 0 1 1.5 2 2.5 3

αCASP8

size (kDa)

αCASP3
← inactive
← active

130 αVCL

C Total RNA

mock
TRAIL

mRNA fold chg/18S

RNAPII genes: *ACTB*, *GAPDH*, *EEF1A*, *PP1A*
RNAPIII genes: *U6*, *7SK*

treatment (hrs): 0 1 1.5 2 2.5 3

D Nascent (4sU) RNA

no biotin
mock
TRAIL

RNA fold chg/18S

RNAPII genes: *ACTB*, *GAPDH*, *EEF1A*, *PP1A*
RNAPIII genes: *U6*, *7SK*

treatment (hrs): - 0 1 1.5 2 2.5 3

A

| zVAD: | − | − | + | + |
| 2 hr TRAIL: | − | + | − | + |

size (kDa)

130 / 100 — αPARP1

35 / 25 — αCASP3

25 / 15 — αBID

55 — αTUBA1A

B Nascent (4sU) ACTB mRNA

☐ 2 hr TRAIL mock ▨ 2 hr TRAIL zVAD

$\log_2$FC

4sU-seq FPKM   qPCR (exon)   qPCR (intron)

** ** **

C 2 hr TRAIL +DMSO

$-\log_{10}p$

↓745 transcripts   ↑69 transcripts

$\log_2$FC

D 2 hr TRAIL +zVAD

$-\log_{10}p$

↓56 transcripts   ↑364 transcripts

$\log_2$FC