Comparative ACE2 variation and primate COVID-19 risk

Amanda D. Melin[1,2,3*+], Mareike C. Janiak[1,3*], Frank Marrone III[4], Paramjit S. Arora[4], & James P.
Higham[5,6+]

1 Department of Anthropology and Archaeology, University of Calgary, CA
2 Department of Medical Genetics, University of Calgary, CA
3 Alberta Children's Hospital Research Institute, University of Calgary, CA
4 Department of Chemistry, New York University, US
5 Department of Anthropology, New York University, US
6 New York Consortium in Evolutionary Primatology, New York, US

* These authors contributed equally to this work

+Authors for correspondence:

Dr Amanda Melin, Department of Anthropology and Archaeology, University of Calgary, 2500
University Drive, Calgary, Alberta T2N 1N4, CA

Dr James Higham, Department of Anthropology, New York University, 25 Waverly Place, New
York, NY 10003, US

Abstract

The emergence of the novel coronavirus SARS-CoV-2, which in humans is highly infectious and
leads to the potentially fatal disease COVID-19, has caused hundreds of thousands of deaths and
huge global disruption. The viral infection may also represent an existential threat to our closest
living relatives, the nonhuman primates, many of which are endangered and often reduced to
small populations. The virus engages the host cell receptor, angiotensin-converting enzyme-2
(ACE2), through the receptor binding domain (RBD) on the spike protein. The contact surface of
ACE2 displays amino acid residues that are critical for virus recognition, and variations at these
critical residues are likely to modulate infection susceptibility across species. While infection
studies are emerging and have shown that some primates, such as rhesus macaques and vervet
monkeys, develop COVID-19-like symptoms when exposed to the virus, the susceptibility of many
other nonhuman primates is unknown. Here, we show that all apes, including chimpanzees,

bonobos, gorillas, and orangutans, and all African and Asian monkeys (catarrhines), exhibit the

same set of twelve key amino acid residues as human ACE2. Monkeys in the Americas, and

some tarsiers, lemurs and lorisoids, differ at significant contact residues, and protein modeling

predicts that these differences should greatly reduce the binding affinity of the ACE2 for the virus,

hence moderating their susceptibility for infection. Other lemurs are predicted to be closer to

catarrhines in their susceptibility. Our study suggests that apes and African and Asian monkeys,

as well as some lemurs are all likely to be highly susceptible to SARS-CoV-2, representing a

critical threat to their survival. Urgent actions have been undertaken to limit the exposure of Great

Apes to humans, and similar efforts may be necessary for many other primate species.

Introduction

In late 2019 a novel coronavirus SARS-CoV-2 emerged in China. In humans, this virus can lead

to the respiratory disease COVID-19, which can be fatal[1,2]. Since then, SARS-CoV-2 has spread

around the world, causing widespread mortality, and with major impacts on societies and

economies. While the virus and its resulting disease represent a major humanitarian disaster,

they also represent a potential existential risk to our closest living relatives, the nonhuman

primates. Transmission incidences of bacteria and viruses - including another coronavirus (H-

CoV-OC43) - from humans to wild populations of nonhuman primates have previously caused

outbreaks of Ebola, yellow fever, and fatal respiratory diseases, leading in some cases to mass

mortality[3–9]. Such past events raise considerable concerns among the global conservation

community with respect to the impact of the current pandemic[10].

Infection studies of rhesus monkeys, longtailed macaques, and vervets as biomedical

models have made it clear that at least some nonhuman primate species are permissive to SARS-

CoV-2 infection and develop symptoms in response to infection that resemble those of humans

following the development of COVID-19, including similar age-related effects[11–16]. Recognizing

the potential danger of COVID-19 to nonhuman primates, the International Union for the

Conservation of Nature (IUCN), together with the Great Apes section of the Primate Specialist

Group, released a joint statement on precautions that should be taken for researchers and

caretakers when interacting with great apes[17]. However, the risk for many primate taxa remains

unknown. Here we begin to assess the potential likelihood that our closest living relatives are

susceptible to SARS-CoV-2 infection.

While the biology underlying susceptibility to SARS-CoV-2 infection remains to be fully

elucidated, the viral target is well established. The SARS-CoV-2 virus binds to the cellular receptor

protein angiotensin-converting enzyme-2 (ACE2), which is expressed on the extracellular surface

of endothelial cells of diverse bodily tissues, including the lungs, kidneys, small intestine and renal

tubes[18]. ACE2 is a carboxypeptidase whose activities include regulation of blood pressure and

inflammatory response through its role in cleaving the vasoconstrictor angiotensin II to produce

angiotensin 1-7 and triggering varied downstream responses[19–22]. ACE2 is made up of a signal

sequence at the N-terminus (residues 1-17), a transmembrane sequence at the C-terminus

(residues 741-762), and an extracellular region, which contains a zinc metallopeptidase domain

(residues 19–611) and a collectrin homolog (residues 612-740)[23,24].

Characterizations of the infection dynamics of SARS-CoV-2 have demonstrated that the

binding affinity for the human ACE2 receptor is high, which is a key factor in determining the

susceptibility and transmission dynamics. When compared to SARS-CoV, which caused a serious

global outbreak of disease in 2002-2003[25,26], the binding affinity between SARS-CoV2 and ACE2

is estimated to be between 4-fold[27–30] and 10- to 20-fold greater[31]. Recent reports describing

structural characterization of ACE2 in complex with the SARS-CoV2 spike protein receptor

binding domain (RBD)[27–30] allow identification of the key binding residues that enable the host-

pathogen protein-protein recognition. Following initial binding of the virus to the ACE2 receptor,

humans experience a great deal of variation in response to infection, with some individuals

experiencing relatively mild symptoms, while others experience major breathing problems and

organ failures, which can lead to death. Some of this response is known to be linked to variation

in how the immune system responds to infection, with some individuals experiencing a

hyperinflammatory 'cytokine storm', which in turn aggravates respiratory failures and increases

mortality risk[32,33]. There may also be some variation among humans in initial susceptibility to

infection, such that approaches examining variation in ACE2 tissue expression and gene

sequences can offer insight into variation in human susceptibility to COVID-19[34–37]. Similarly, we

can use such an approach to compare sequence variation across species, and hence try to predict

the likely interspecific variation in susceptibility to initial infection. Previous analysis of comparative

variation at these sites enabled estimates of the affinity of the ACE2 receptor for SARS-CoV in

nonhuman species (bats)[38].

Here, we undertake such an analysis for SARS-CoV-2 across the primate radiation. Our

aim is to investigate the likelihood of initial susceptibility to infection for different major radiations

and species, while recognizing that down-stream processes such as immune responses are likely

to determine the extent to which species and individuals develop symptoms and pathologies in

response to infection. We compiled *ACE2* gene sequence data from 29 primate species for which

genomes are publicly available, covering primate taxonomic breadth. For comparison, we

assessed 4 species of other mammals that have been tested directly for SARS-CoV2
susceptibility in laboratory infection studies[39]. We also included in our analysis the amino acid
sequence variation at these sites for horseshoe bats, thought to be the original vector of the virus,
and pangolins, a potential intermediate host, where viral recombination may have led to the novel
viral form SARS-CoV-2[40]. We assessed the variation at amino acid residues identified as critical
for ACE2 recognition by CoV RBD, and undertook analysis of positive selection and protein
modeling to gauge the potential for adaptive differences and the likely effects of protein variation.
Our aim was to develop predictions about the susceptibility of our closest living relatives to SARS-
CoV-2 as a resource for stakeholders, including researchers, caretakers, practitioners,
conservationists, and governmental and nongovernmental agencies.
113
Methods
Variation in ACE2 sequences
We compiled *ACE2* gene sequences for 16 catarrhine primates: 4 species from all 3 genera of
great ape (*Gorilla*, *Pan*, *Pongo*), 2 genera of gibbons (*Hylobates*, *Nomascus*), and 10 species of
African and Asian monkeys in 7 genera (*Cercocebus, Chlorocebus, Macaca, Mandrillus, Papio,*
Rhinopithecus, Piliocolobus, Theropithecus); 6 genera of platyrrhines (monkeys from the
Americas: *Alouatta, Aotus, Callithrix, Cebus, Saimiri, Sapajus*); 1 species of tarsier (*Carlito*
syrichta); and 5 genera of strepsirrhines (lemurs and lorisoids: *Eulemur*, *Daubentonia*,
Microcebus, *Propithecus*, *Otolemur*) (Suppl. Table S1). We also included 4 species of mammals
that have been tested clinically for susceptibility to SARS-CoV-2 infection[39], including the
domestic cat (*Felis catus*), dog (*Canis lupus familiaris*), pig (*Sus scrofa*), and ferret (*Mustela*
putorius furo). Finally, we included the pangolin (*Manis javanica*) and several bat species,
including horseshoe bats (*Rhinolophus* spp., *Hipposideros pratti*, *Myotis daubentonii*). Sequences
were retrieved from NCBI, either from annotations of published genomes or from GenBank
entries[38]. We manually checked annotations by performing tblastn searches of the human ACE2
protein sequence against each genome. We identified one misannotation for exon 15 in
Microcebus murinus, which we manually corrected. The *ACE2* nucleotide sequence for *Alouatta*
palliata was obtained from an unpublished draft genome, via tblastn searches using the *Cebus*
capucinus ACE2 protein sequence as a query and default search settings. Accession numbers
for sequences retrieved from NCBI and GenBank are provided in Supplemental Table S1 and the
Alouatta palliata sequence is available in the supplemental materials.
Coding sequences were translated using Geneious Version 9.1.8 and we aligned both
nucleotide and amino acid sequences with MAFFT[41]. Amino acids were aligned with the

BLOSUM62 scoring matrix, while the 200 PAM scoring matrix was used for nucleotides. A 1.53

gap open penalty and an offset value of 0.123 were used for both. We manually inspected and

corrected any misalignments, and verified the absence of indels and premature stop codons.

To visualize patterns of gene conservation across taxa and identify the congruence of the

ACE2 gene tree with currently accepted phylogenetic relationships among species, we

reconstructed trees using both Bayesian (MrBayes 3.2.6[42]) and Maximum Likelihood (RAxML

8.2.11[43]) methods with 200,000 MCMC cycles and 1,000 bootstrap replicates, respectively (code

available on GitHub[44]). Gene trees were compared to a current species phylogeny assembled

using TimeTree[45], which is also used to illustrate the evolutionary relationships between study

species in Figure 1. Parsimony-informative sites along the *ACE2* sequence were identified with

the pis function in the R package ips v. 0.0.11[46,47].

Identification of critical binding residues and species-specific ACE2–RBD interactions

Critical ACE2 protein contact sites for the viral spike protein receptor binding domain (RBD) have

been identified using cryo EM and X-ray crystallography structural analysis methods[27–30]. The

ACE2-RBD complex is characteristic of protein-protein interactions (PPIs) that feature extended

interfaces spanning a multitude of binding residues. Experimental and computational analyses of

PPIs have shown that a handful of contact residues can dominate the binding energy landscape[48].

Alanine scanning mutagenesis provides an assessment of the contribution of each residue to

complex formation[49–51]. Critical binding residues can be computationally identified by assessing

the change in binding free energy of complex formation upon mutation of the particular residue to

alanine, which is the smallest residue that may be incorporated without significantly impacting the

protein backbone conformation[52]. Our computational modeling utilizes the human SARS

RBD/ACE2 high resolution structures, and we make the implicit assumption that the overall

conformation of ACE2 is conserved among different species. This assumption, which is rooted in

the high sequence similarity between ACE2 sequences, allows us to use the structure of the

complex to predict the impact of mutations at the protein-protein interface.

We defined critical residues as those that upon mutation to alanine decrease the binding

energy by a threshold value $\Delta\Delta G_{bind}$ ≥1.0 kcal/mol. Nine of the 21 residues identified by alanine

scanning as involved in the ACE2-RBD complex met this criterion (Suppl. Table S2). There was

large congruence in the sites identified with those highlighted by other methods. Each of the eight

sites implicated by cryo EM[27], were also detected by alanine modelling; five residues were ≥1.0

kcal/mol threshold and 3 were below this threshold. To be cautious, in addition to the the 9 critical

ACE2 sites we identified through alanine scanning, we also examined residue variation at the 3

sites that fell below the ≥1.0 kcal/mol threshold but that were identified as important by structural

analyses[27–30] for a total of 12 critical sites. All computational alanine scanning mutagenesis

analyses were performed using Rosetta software[52]. The alanine mutagenesis approach has been

extensively evaluated and used to analyze PPIs and design their inhibitors, including by members

of the present authorship[53,54].

We utilized the SSIPe program[55] to predict how ACE2 amino acid differences in each

species would affect the relative binding energy of the ACE2/SARS-Cov-2 interaction. Using

human ACE2 bound to the SARS-Cov-2 RBD as a benchmark (PDB 6M0J), the program mutates

selected residues and compares the binding energy to that of the original. Using this algorithm,

we studied interactions of all primates across the full suite of amino acid changes occurring at

critical binding sites for each species. To more thoroughly assess the impact of each amino acid

substitution, we also examined the predicted effect of individual amino acid changes (in isolation)

on protein-binding affinity.

Adaptive evolution of ACE2 sequences

We further investigated *ACE2* and how selective pressures in different clades might be shaping

variation at the binding sites, using codeml clade C and branch-site models in PAML[56]. We first

tested if selection acting on *ACE2* is divergent between the major clades in our sample

(platyrrhine, catarrhine, and strepsirrhine primates, non-primate mammals) with the codeml clade

model C, which was compared to the null model (M2a_rel) with a likelihood ratio test[57]. This test

shows whether there is divergent selection (dN/dS ratio = ω) across all clades, but not which

clades are experiencing positive selection. We, therefore, followed the clade model with a series

of branch-site models, which allow one clade at a time to be designated as a set of "foreground"

branches and test whether this clade has experienced episodes of positive selection compared

to the remaining sets of "background" branches ($\omega_{foreground} > \omega_{background}$). Branch-site models are

compared to a null model that fixes ω at 1 with a likelihood ratio test. In the case of the alternative

model having a significantly better fit than the null model, indicating positive selection, potential

sites under positive selection are identified with a Bayes Empirical Bayes (BEB) approach[58]. We

completed branch-site models for each primate clade (platyrrhine, strepsirrhine, and catarrhine),

as well as bats, because previous research has identified *ACE2* to be under positive selection in

this clade, potentially in response to coronaviruses[59]. We had to exclude *Hipposideros pratti* and

Myotis daubentonii from PAML analyses, because only a partial *ACE2* sequence was available

for these two species. Input files and control files for PAML codeml analyses are available in the
GitHub repository[44].

Results

Variation in ACE2 sequences

The *ACE2* gene (2418 bp) and translated protein (805 amino acids) sequences are strongly

conserved across primates. The average pairwise identity across 29 primate species is 93.6% for

the *ACE2* nucleotide sequence and 90.8% for the protein sequence, with a pairwise similarity

(BLOSUM62 ≥ 1) of 95.3% (Suppl. Tables S3-5). Out of 2418 bp, 1631 bp (67.5%) are identical,

while 401 bp (16.58%) are phylogenetically informative sites for primates, and gene trees we

generated (Suppl. Fig. S1a,b) closely recapitulate the currently accepted phylogeny of primates

(Figure 1). In particular, the twelve sites in the ACE2 protein that are critical for binding of the

SARS-CoV-2 virus are invariant across the Catarrhini, which includes great apes, gibbons, and

monkeys of Africa and Asia (Figure 1). Furthermore, catarrhines do not vary at any of the 21 sites

identified by alanine scanning (Suppl. Table S2, Suppl. Fig. S2). The other major radiation of

monkeys, those found in the Americas (Platyrrhini), have ACE2 sequences that are less similar

to humans across the length of the protein (91.68-92.55% identical to *H. sapiens*, Suppl. Table

S4) but conserved within their clade (average pairwise identity 97.2%, Suppl. Table S4). They

share nine of twelve critical amino acid residues with catarrhine primates; the three sites that vary

from catarrhines, H41, E42 and T82, are conserved within the platyrrhines. Strepsirrhine primates

and tarsiers, were more variable in the binding sites and less similar to the human protein across

the length of the sequence (81.86-86.93% pairwise identity, Suppl. Table S4). Like platyrrhines,

the tarsier (*Carlito syrichta*), mouse lemur *(Microcebus murinus)*, and galago (*Otolemur garnettii*)

have an H41 residue, while the sifaka (*Propithecus coquereli*), aye-aye (*Daubentonia*

madagascariensis), and the blue-eyed black lemur (*Eulemur flavifrons*) have the same allele as

humans and other catarrhines, Y41.

In non-primate mammals, a higher number of amino acid substitutions are evident (77.37-

85.22% pairwise identity to *H. sapiens*, Suppl. Table S4), including at critical binding sites. All

species possess a different residue to primates at site 24. Bats are exceptionally variable within

the binding sites, with the genus *Rhinolophus* alone encompassing all of the variation seen in the

rest of the non-primate mammals. Where primates have glutamine (Q24), bats have glutamate

(E24), lysine (K24), leucine (L24), or arginine (R24) (Figure 1). All fasta alignments of ACE2 gene

and protein sequences are available in the supplemental materials, a full-length protein alignment

is shown in Suppl. Figure S2, and distance matrices are provided in Suppl. Table S3-5.

Analysis of species-specific residues on ACE2–RBD interactions

The ACE2 receptors of all catarrhines have identical residues to humans at the RBD/ACE2

binding interface across all 12 critical sites and are predicted to have similar binding affinity for

SARS-CoV-2. Platyrrhines diverge from catarrhines at three of the twelve critical amino acid

residues. Compared to catarrhine ACE2, the platyrrhines' ACE2 is predicted to bind SARS-CoV2

RBD with roughly 400-fold reduced affinity ($\Delta\Delta G_{bind}$=3.5 kcal/mol) (Table 1a). In particular, the

change at site 41 from Y to H found in monkeys in the Americas has the largest impact of any

residue change examined (Table 1b), which alone is predicted to lead to a 25-fold decrease in

the binding affinity to SARS-CoV-2 (Figure 2). This single mutation combined with additional

substitutions, especially Q42E, found in platyrrhines is predicted to significantly reduce the

likelihood of successful viral binding (Table 1b). Of the other primates modeled, two of the three

strepsirhines, and tarsiers, also have the H41 residue and furthermore have additional protein

sequence differences leading to further decreases in predicted binding affinity. The predicted

binding affinity of tarsier ACE2 is the most dissimilar to humans and this primate might be the

least susceptible of the species we examine. In contrast, Coquerel's sifaka (*Propithecus*

coquereli), the aye-aye (*Daubentonia madagascariensis*), and blue-eyed black lemur (*Eulemur*

flavifrons) share the same residue as humans and other catarrhines at site 41 and have projected

affinities that are near to humans (Table 1b). Other mammals included in our study - ferrets, cats,

dogs, pigs, pangolin and two of the seven bat species (*R. pusillus* and *R. macrotis)* - show the

same residue as humans (Y) at site 41, with accompanying strong affinities for SARS-CoV-2. The

remaining five sister species of bats possess H41 and lower binding affinities (Table 1b).

Adaptive evolution of ACE2 sequences

We find evidence that the selective pressures acting on *ACE2* are not equivalent across the major

clades in our analysis. The codeml clade model C provided a better fit than the null model (LRT

= 26.726, p < 0.001; Table 2, Suppl. Table S6). Branch-site models indicate that the catarrhine

primate clade (LRT = 14.546, p < 0.001) and bat clade (LRT = 42.649, p < 0.001) are both under

positive selection, while platyrrhines (LRT = 0.633, p = 0.427) and strepsirrhines (LRT = 0.833, p

= 0.361) are not. The six positively selected sites in the bat clade include the binding site 24 and

two others adjacent to known binding sites (Table 2). In catarrhines, the three positively selected

sites identified by BEB calculations are not near the binding sites for SARS-Cov-2 (residues 249,
653, and 658; Table 2).

Discussion

Our results strongly suggest that catarrhines - all apes, and all monkeys of Africa and Asia, are
likely to be susceptible to infection by SARS-CoV-2. There is high conservancy in the protein
sequence of the target receptor, ACE2, including uniformity at all identified and tested major
binding sites. Indeed, even among the 21 residues identified in our full list of potential binding
points, catarrhines are invariant (Suppl. Table 2, Suppl. Fig. S2). Consistent with our results,
infection studies show that rhesus monkeys (*Macaca mulatta*), longtailed macaques (*M.
fascicularis*) and vervets (*Chlorocebus sabaeus*) are permissive to infection by SARS-CoV-2, and
go on to develop COVID-19 like symptoms[11–14,16]. Our results based on protein modeling offer
potentially better news for monkeys in the Americas (platyrrhines). There are three differences in
amino acid residues between platyrrhines and catarrhines, and two of these, H41Y and E42Q
show strong evidence of being impactful changes. These amino acid changes are modeled to
reduce the binding affinity between SARS-CoV-2 and ACE2 by ca. 400-fold. Recent clinical
analysis of viral shedding, viremia, and histopathology in catarrhine (macaque) versus platyrrhine
(marmoset, *Callithrix jacchus*) responses to inoculation with SARS-CoV-2, show much more
severe presentation of disease symptoms in the former, strongly supporting our results[13]. Similar
reduced susceptibility is predicted for tarsiers, and two of the five lemurs and lorisoids
(strepsirrhines). What is concerning is three of the analyzed lemurs spanning divergent lineages
- the Coquerel's sifaka, the aye-aye, and the blue-eyed black lemur - are more similar to
catarrhines at important binding sites, including possessing the high risk residue variant at site
41, and as such are also predicted to be susceptible. Nonetheless, these are only predicted
results based on amino acid residues, and protein-protein interaction models. We urge extreme
caution in using our analyses as the basis for relaxing policies regarding the protection of
platyrrhines, tarsiers or any strepsirrhines. Experimental assessment of synthetic protein
interactions can now occur in the laboratory e.g.[60], and confirmation of our model predictions
should be sought before any firm conclusions are reached.

Emerging evidence in experimental mammalian models appears to support our results;
dogs, ferrets, pigs, and cats have all shown some susceptibility to SARS-CoV-2 but have
demonstrated variation in disease severity and presentation, including across studies[39,61].
Substitutions at binding sites might be at least partially protective against COVID-19 in these
mammals. For example, the limited experimental evidence to date suggests that while cats -

which have the same residue as humans at site 34 - are not strongly symptomatic, they present
lung lesions, while dogs - which have a substitution at this site - do not[39]. The amino acid residue
at site 24 differs from primates in all other mammalian species examined. However, our models
suggest that the variant residues may confer relatively minor reductions in binding affinity. Other
sources of variation may affect ACE2 protein stability[35]. Our results are also consistent with
previous reports that ACE2 genetic diversity is greater among bats than that observed among
mammals susceptible to SARS-CoV-type viruses. This variation has been suggested to indicate
that bat species may act as a reservoir of SARS-CoV viruses or their progenitors[38]. Intriguingly,
all but 2 bat species we examined have the putatively protective variant, H41. Additionally, results
of our codeml branch-site analysis support previous findings of *ACE2* in bats being under positive
selection, including sites within the binding domain of SARS-CoV and SARS-CoV-2[59], which may
be evidence of host-virus coevolution. Sites showing evidence of positive selection within
catarrhine *ACE2* sequences were not in or near known CoV binding sites (Table 2, Figure 1). Two
(residues 653, 658) fall within the cleavage site (residues 652-659) utilized by the sheddase
ADAM17, known to interact with ACE2[62]. However, neither of the residues under selection are
the amino acids targeted by ADAM17[63] leaving the functional significance of evolution at these
sites uncertain. Further clinical and laboratory study is needed to fully understand infection
dynamics.

There are a number of important caveats to our study. Firstly, all of our predictions are
based on interpretations of gene and resultant amino acid sequences, rather than based on direct
assessment of individual responses to induced infection. Nonetheless, the overall pattern of our
results is being borne out by infection studies on a few species that are used as biomedical
models. So far, all catarrhine species tested by infection studies, including rhesus macaques,
longtailed macaques, and vervet monkeys[13,14,64] have exhibited COVID-19-like symptoms in
response to infection, including large lung and other organ lesions[13] and cytokine storms[14]. In
contrast, marmosets did not exhibit major symptoms in response to infection[13]. While these results
support and validate our findings based on ACE2 sequence interpretation, the number of primate
species that can and will be tested directly by infection studies will be restricted to just a handful.
Our study enhances this picture, by allowing inferences to be made across the primate radiation,
backed up by the published infection studies on a few target model species.

Some of our results, such as the uniform conservation of *ACE2* among catarrhines,
backed up by the demonstrated high susceptibility of humans and other catarrhines to SARS-
CoV-2, should give a good degree of confidence of high levels of risk.

Given the identical residues of humans to other apes and monkeys in Asia and Africa at
the target site, it seems unlikely that the ACE2 receptor and the SARS-CoV-2 proteins would not
readily bind. Our results for other taxa are dependent on modeling, hence should be treated more
cautiously. This includes all interpretations of the susceptibility of platyrrhines and strepsirrhines,
where the effects of residue differences on binding affinities have been estimated based on
protein-protein interaction modeling. Another caveat is that we have modeled only interactions at
binding sites, and not predictions based on full residue sequence variation. Residues that are not
in direct contact may still affect binding allosterically. Other factors, including proteases necessary
for viral entry, and other viral targets, may also impact disease susceptibility and responses[35].
More generally, if adhering to the precautionary principle, then our results highlighting higher risks
to some species should be taken with greater gravity than our results that predict potential lower
risks to others. Another limitation of our study is that we have looked at only 29 primate species,
albeit with broad taxonomic scope. Analysis of additional species is important, especially among
strepsirrhine species, where our coverage is relatively scant. In particular, the residue overlap at
important binding sites in the sequences of Coquerel's sifaka, the aye-aye, and blue-eyed black
lemur with those of catarrhines suggests many lemurs may be highly vulnerable and we
underscore the need to assess a wider diversity of lemur species. Furthermore, we examine only
one individual per species, and intraspecific variation across populations should be considered;
however, studies on intraspecific ACE2 variation with humans and vervet monkeys suggest ACE2
variants are low in frequency[65–67]. Finally, it is also important to remember that our study assesses
only the potential for initial binding of the virus to the target site. Downstream consequences of
infection may differ drastically based on species-specific proteases, genomic variants,
metabolism, and immune system responses[68,69]. In humans, the development of COVID-19 can
lead to a pro-inflammatory cytokine storm of hyperinflammation, which may lead to some of the
more severe impacts of infection[32,70]. Nonetheless, it is evidence from the hundreds of thousands
of deaths and global lockdown that humans are highly susceptible to SARS-CoV-2 infection, and
our results suggest that all apes and monkeys in Africa and Asia are similarly susceptible.

Many endangered primate species are now only found in very small population sizes[71].
For example, there are believed to be only around 1000 mountain gorillas left in their entire
range[72]. With such small populations, the introduction of a new highly infectious disease is a
potential extinction-level event. Re-opening access to habituated great ape groups for tourism
purposes, which may be critical to local economies[73], may be fraught with issues. IUCN best
practices recommend that tourists stay at least 7 metres away from great apes[74], but in practice,
almost all tourists get far closer than this - for example, the average distance that tourists get from

mountain gorillas at the Bwindi Impenetrable National Park in Uganda is just 2.76 metres[75].
Concerted effort may be required by all stakeholders to try to avoid the introduction of SARS-
CoV-2 into wild primate populations[10]. Recent measures suggested by the IUCN for researchers
and caretakers of great ape populations include: ensuring that all individuals wear clean clothing
and disinfected footwear; providing hand-washing facilities; requiring that a surgical face mask be
worn by anyone coming within 10 meters of great apes; ensuring that individuals needing to cough
or sneeze ideally leave the area, or at least cough/sneeze into the crux of their elbows; imposing
a 14-day quarantine for all people arriving into great ape areas who will come into frequent close
proximity with them[17]. The IUCN's 'Best Practice Guidelines for Health Monitoring and Disease
Control in Great Ape Populations' should also be followed[76].

Our results suggest that dozens of nonhuman primate species, including all of our closest
relatives, are likely to be highly susceptible to SARS-CoV-2 infection, and vulnerable to its effects.
Major actions may be needed to limit the exposure of many wild primate populations to humans.
This is likely to require coordinated input from all stakeholders, including local communities,
international and national governmental agencies, nongovernmental conservation and
development organizations, and academics and researchers. While the focus of many at this time
is rightly on mitigating the humanitarian devastation of COVID-19, we also have a duty to ensure
that our closest living relatives do not suffer extinctions, or massive population declines, in
response to yet another human-induced catastrophe.

Data Availability Statement

Nucleotide and protein sequences used in this study are available from NCBI and are also
available as fasta files and alignments in the supplemental material and on Github
(https://github.com/MareikeJaniak/ACE2). All code used in this project is available in the same
repository.

Acknowledgements

MCJ was funded by a National Sciences and Engineering Council of Canada Discovery
Accelerator Supplement to ADM and by a postdoctoral fellowship from the Alberta Children's
Hospital Research Institute. PSA thanks the National Institutes of Health (R35GM130333) for
financial support. We thank four reviewers for constructive comments, which improved the
manuscript considerably.

Author Contributions

ADM, JPH, and MCJ designed the study. ADM and JPH wrote the paper with input and edits from

MCJ and PSA. MCJ conducted genetic analyses with input from ADM. FM ran the protein

substitution models with input from PSA. All authors have approved the final submission for

publication.

Literature Cited

1. Zhu, N. *et al.* A novel coronavirus from patients with pneumonia in China, 2019. *N. Engl. J. Med.* **382**, 727–733 (2020).

2. Poon, L. L. M. & Peiris, M. Emergence of a novel human coronavirus threatening human health. *Nat. Med.* **26**, 317–319 (2020).

3. Holzmann, I. *et al.* Impact of yellow fever outbreaks on two howler monkey species (*Alouatta guariba clamitans* and *A. caraya*) in Misiones, Argentina. *Am. J. Primatol.* **72**, 475–480 (2010).

4. Bermejo, M. *et al.* Ebola outbreak killed 5000 gorillas. *Science* **314**, 1564 (2006).

5. Köndgen, S. *et al.* Pandemic human viruses cause decline of endangered great apes. *Curr. Biol.* **18**, 260–264 (2008).

6. Kaur, T. *et al.* Descriptive epidemiology of fatal respiratory outbreaks and detection of a human-related metapneumovirus in wild chimpanzees (*Pan troglodytes*) at Mahale Mountains National Park, Western Tanzania. *Am. J. Primatol.* **70**, 755–765 (2008).

7. Goldberg, T. L., Gillespie, T. R., Rwego, I. B., Estoff, E. L. & Chapman, C. A. Forest fragmentation as cause of bacterial transmission among nonhuman primates, humans, and livestock, Uganda. *Emerg. Infect. Dis.* **14**, 1375–1382 (2008).

8. Palacios, G. *et al.* Human metapneumovirus infection in wild mountain gorillas, Rwanda. *Emerg. Infect. Dis.* **17**, 711–713 (2011).

9. Patrono, L. V. *et al.* Human coronavirus OC43 outbreak in wild chimpanzees, Côte d´Ivoire, 2016. *Emerg. Microbes Infect.* **7**, 118 (2018).

10. Gillespie, T. R. & Leendertz, F. H. COVID-19: protect great apes during human pandemics. *Nature* **579**, 497 (2020).

11. Shan, C. *et al.* Infection with novel coronavirus (SARS-CoV-2) causes pneumonia in the rhesus macaques. *ResearchSquare* doi:10.21203/rs.2.25200/v1.

12. Rockx, B. *et al.* Comparative pathogenesis of COVID-19, MERS, and SARS in a nonhuman

primate model. *Science* **368**, 1012–1015 (2020).

13. Lu, S. *et al.* Comparison of SARS-CoV-2 infections among 3 species of non-human

primates. *bioRxiv* 2020.04.08.031807 (2020) doi:10.1101/2020.04.08.031807.

14. Blair, R. V. *et al.* ARDS and cytokine storm in SARS-CoV-2 infected Caribbean vervets.

bioRxiv 2020.06.18.157933 (2020) doi:10.1101/2020.06.18.157933.

15. Yu, P. *et al.* Age-related rhesus macaque models of COVID-19. *Animal Model Exp Med* **3**,

93–97 (2020).

16. Deng, W. *et al.* Primary exposure to SARS-CoV-2 protects against reinfection in rhesus

macaques. *Science* (2020) doi:10.1126/science.abc5343.

17. IUCN SSC Wildlife Health Specialist Group & Primate Specialist Group, Section on Great

Apes. Great apes, COVID-19 and the SARS CoV-2 joint statement of the IUCN SSC

Wildlife Health Specialist Group and the Primate Specialist Group, Section on Great Apes.

Great apes and SARS CoV 2 Communique (2020).

18. Hamming, I. *et al.* Tissue distribution of ACE2 protein, the functional receptor for SARS

coronavirus. A first step in understanding SARS pathogenesis. *J. Pathol.* **203**, 631–637

(2004).

19. Vickers, C. *et al.* Hydrolysis of biological peptides by human angiotensin-converting

enzyme-related carboxypeptidase. *J. Biol. Chem.* **277**, 14838–14843 (2002).

20. Donoghue, M. *et al.* Heart block, ventricular tachycardia, and sudden death in *ACE2*

transgenic mice with downregulated connexins. *J. Mol. Cell. Cardiol.* **35**, 1043–1053

(2003).

21. Rodrigues Prestes, T. R., Rocha, N. P., Miranda, A. S., Teixeira, A. L. & Simoes-E-Silva, A.

C. The anti-inflammatory potential of ACE2/Angiotensin-(1-7)/Mas receptor axis: Evidence

from basic and clinical research. *Curr. Drug Targets* **18**, 1301–1313 (2017).

22. Verdecchia, P., Cavallini, C., Spanevello, A. & Angeli, F. The pivotal link between ACE2

deficiency and SARS-CoV-2 infection. *Eur. J. Intern. Med.* **76**, 14–20 (2020).

23. Tipnis, S. R. *et al.* A human homolog of angiotensin-converting enzyme. Cloning and

functional expression as a captopril-insensitive carboxypeptidase. *J. Biol. Chem.* **275**,

33238–33243 (2000).

24. Towler, P. *et al.* ACE2 X-ray structures reveal a large hinge-bending motion important for

inhibitor binding and catalysis. *J. Biol. Chem.* **279**, 17996–18007 (2004).

25. May, R. M., McLean, A. R., Pattison, J., Weiss, R. A. & Heymann, D. L. The international

response to the outbreak of SARS in 2003. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **359**,

1127–1129 (2004).

26. Demmler, G. J. & Ligon, B. L. Severe acute respiratory syndrome (SARS): a review of the

history, epidemiology, prevention, and concerns for the future. *Semin. Pediatr. Infect. Dis.*

14, 240–244 (2003).

27. Yan, R. *et al.* Structural basis for the recognition of SARS-CoV-2 by full-length human

ACE2. *Science* **367**, 1444–1448 (2020).

28. Shang, J. *et al.* Structural basis of receptor recognition by SARS-CoV-2. *Nature* **581**, 221–

224 (2020).

29. Lan, J. *et al.* Structure of the SARS-CoV-2 spike receptor-binding domain bound to the

ACE2 receptor. *Nature* **581**, 215–220 (2020).

30. Wang, Q. *et al.* Structural and functional basis of SARS-CoV-2 entry by using human

ACE2. *Cell* **181**, 894–904.e9 (2020).

31. Wrapp, D. *et al.* Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation.

Science **367**, 1260–1263 (2020).

32. Chen, G. *et al.* Clinical and immunologic features in severe and moderate Coronavirus

Disease 2019. *J. Clin. Invest.* **130**, 2620–2629 (2020).

33. Ragab, D., Salah Eldin, H., Taeimah, M., Khattab, R. & Salem, R. The COVID-19 Cytokine

Storm; What We Know So Far. *Front. Immunol.* **11**, 1446 (2020).

34. Benetti, E. *et al.* ACE2 variants underlie interindividual variability and susceptibility to

COVID-19 in Italian population. *medRxiv* (2020) doi:10.1101/2020.04.03.20047977.

35. Lukassen, S. *et al.* SARS-CoV-2 receptor ACE2 and TMPRSS2 are primarily expressed in

bronchial transient secretory cells. *EMBO J.* **39**, e105114 (2020).

36. Hussain, M. *et al.* Structural variations in human ACE2 may influence its binding with

SARS-CoV-2 spike protein. *J. Med. Virol.* (2020) doi:10.1002/jmv.25832.

37. Gibson, W. T., Evans, D. M., An, J. & Jones, S. J. M. ACE 2 coding variants: A potential X-

linked risk factor for COVID-19 disease. *bioRxiv* 2020.04.05.026633 (2020)

doi:10.1101/2020.04.05.026633.

38. Hou, Y. *et al.* Angiotensin-converting enzyme 2 (ACE2) proteins of different bat species

confer variable susceptibility to SARS-CoV entry. *Arch. Virol.* **155**, 1563–1569 (2010).

39. Shi, J. *et al.* Susceptibility of ferrets, cats, dogs, and other domesticated animals to SARS-

coronavirus 2. *Science* **368**, 1016–1020 (2020).

40. Wong, M. C., Javornik Cregeen, S. J., Ajami, N. J. & Petrosino, J. F. Evidence of

recombination in coronaviruses implicating pangolin origins of nCoV-2019. *bioRxiv*

doi:10.1101/2020.02.07.939207.

41. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

42. Huelsenbeck, J. P. & Ronquist, F. MRBAYES: Bayesian inference of phylogenetic trees.

Bioinformatics **17**, 754–755 (2001).

43. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large

phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

44. Janiak, M. C. Supporting material and code for primate comparative ACE2 project. *GitHub*

https://github.com/MareikeJaniak/ACE2 (2020).

45. Kumar, S., Stecher, G., Suleski, M. & Hedges, S. B. TimeTree: A resource for timelines,

timetrees, and divergence times. *Mol. Biol. Evol.* **34**, 1812–1819 (2017).

46. R Core Team. *R: A language and environment for statistical computing*. (2019).

47. Heibl, C., Cusimano, N. & Krah, F.-S. *Interfaces to Phylogenetic Software in R.*

48. Clackson, T. & Wells, J. A. A hot spot of binding energy in a hormone-receptor interface.

Science **267**, 383–386 (1995).

49. Bogan, A. A. & Thorn, K. S. Anatomy of hot spots in protein interfaces. *J. Mol. Biol.* **280**, 1–

9 (1998).

50. Massova, I. & Kollman, P. A. Computational alanine scanning to probe protein- protein

interactions: a novel approach to evaluate binding free energies. *J. Am. Chem. Soc.* **121**,

8133–8143 (1999).

51. Kortemme, T. & Baker, D. A simple physical model for binding energy hot spots in protein–

protein complexes. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 14116–14121 (2002).

52. Kortemme, T., Kim, D. E. & Baker, D. Computational alanine scanning of protein-protein

interfaces. *Sci. STKE* **2004**, l2 (2004).

53. Jochim, A. L. & Arora, P. S. Systematic analysis of helical protein interfaces reveals targets

for synthetic inhibitors. *ACS Chem. Biol.* **5**, 919–923 (2010).

54. Raj, M., Bullock, B. N. & Arora, P. S. Plucking the high hanging fruit: A systematic approach

for targeting protein–protein interactions. *Bioorg. Med. Chem.* **21**, 4051–4057 (2013).

55. Huang, X., Zheng, W., Pearce, R. & Zhang, Y. SSIPe: accurately estimating protein–protein

binding affinity change upon mutations using evolutionary profiles in combination with an

optimized physical energy function. *Bioinformatics* **36**, 2429–2437 (2020).

56. Yang, Z. PAML 4: Phylogenetic Analysis by Maximum Likelihood. *Mol. Biol. Evol.* **24**, 1586–

1591 (2007).

57. Weadick, C. J. & Chang, B. S. W. An improved likelihood ratio test for detecting site-

specific functional divergence among clades of protein-coding genes. *Mol. Biol. Evol.* **29**,

1297–1300 (2012).

58. Yang, Z., Wong, W. S. W. & Nielsen, R. Bayes empirical bayes inference of amino acid

sites under positive selection. *Mol. Biol. Evol.* **22**, 1107–1118 (2005).

59. Demogines, A., Farzan, M. & Sawyer, S. L. Evidence for ACE2-utilizing coronaviruses

(CoVs) related to severe acute respiratory syndrome CoV in bats. *J. Virol.* **86**, 6350–6353

(2012).

60. Guy, J. L., Jackson, R. M., Jensen, H. A., Hooper, N. M. & Turner, A. J. Identification of

critical active-site residues in angiotensin-converting enzyme-2 (ACE2) by site-directed

mutagenesis. *FEBS J.* **272**, 3512–3520 (2005).

61. Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat

origin. *Nature* **579**, 270–273 (2020).

62. Mukerjee, S. *et al.* ACE2 and ADAM17 interaction regulates the activity of presympathetic

neurons. *Hypertension* **74**, 1181–1191 (2019).

63. Heurich, A. *et al.* TMPRSS2 and ADAM17 cleave ACE2 differentially and only proteolysis

by TMPRSS2 augments entry driven by the Severe Acute Respiratory Syndrome

Coronavirus spike protein. *J. Virol.* **88**, 1293–1307 (2014).

64. Hartman, A. L. *et al.* SARS-CoV-2 infection of African green monkeys results in mild

respiratory disease discernible by PET/CT imaging and prolonged shedding of infectious

virus from both respiratory and gastrointestinal tracts. *bioRxiv* 2020.06.20.137687 (2020)

doi:10.1101/2020.06.20.137687.

65. Schmitt, C. A. *et al. ACE2* and *TMPRSS2* variation in savanna monkeys (*Chlorocebus*

spp.): Potential risk for zoonotic/anthroponotic transmission of SARS-CoV-2 and a potential

model for functional studies. *PLoS One* **15**, e0235106 (2020).

66. Stawiski, E. W. *et al.* Human ACE2 receptor polymorphisms predict SARS-CoV-2

susceptibility. *bioRxiv* 2020.04.07.024752 (2020) doi:10.1101/2020.04.07.024752.

67. Cao, Y. *et al.* Comparative genetic analysis of the novel coronavirus (2019-nCoV/SARS-

CoV-2) receptor ACE2 in different populations. *Cell Discov* **6**, 11 (2020).

68. Li, J. *et al.* Virus-host interactome and proteomic survey of PMBCs from COVID-19 patients

reveal potential virulence factors influencing SARS-CoV-2 pathogenesis. *bioRxiv*

2020.03.31.019216 (2020) doi:10.1101/2020.03.31.019216.

69. Hoffmann, M. *et al.* SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell* **181**, 271–280.e8 (2020).

70. Pedersen, S. F. & Ho, Y.-C. SARS-CoV-2: A storm is raging. *J. Clin. Invest.* **130**, 2202–2205 (2020).

71. Estrada, A. *et al.* Impending extinction crisis of the world's primates: Why primates matter. *Science Advances* **3**, e1600946 (2017).

72. Granjon, A. *et al.* Estimating abundance and growth rates in a wild mountain gorilla population. *Animal Conservation* (2020) doi:10.1111/acv.12559.

73. Sandbrook, C. G. Putting leakage in its place: The significance of retained tourism revenue in the local context in Rural Uganda. *Journal of International Development* **22**, 124–136 (2010).

74. Macfie, E. J. & Williamson, E. A. *Best Practice Guidelines for Great Ape Tourism*. (IUCN, 2010).

75. Sandbrook, C. & Semple, S. The rules and the reality of mountain gorilla *Gorilla beringei beringei* tracking: how close do tourists get? *Oryx* **40**, 428–433 (2006).

76. Gilardi, K. V. *et al.* Best practice guidelines for health monitoring and disease control in great ape populations. *Occasional Papers of the IUCN Species Survival Commission* **56**, (2015).

581 **Table 1.** Results of computational protein-protein interaction experiments predicting impact of
582 amino acid changes, relative to human ACE2 residues, at critical binding sites with SARS-CoV-2
583 receptor binding domain. Impacts of changes across the full complement of critical binding sites
584 are presented in (A), single residue replacements are presented in (B).

586 **A)**

Species	Mutations	ΔΔG (kcal/mol)[a]
Carlito syrichta	H34Q, Y41H, M82S, K353N	5.506
Microcebus murinus	D30E, H34N, Y41H, M82T	4.001
Propithecus coquereli	M82T	0.938
Otolemur garnettii	H34R, D38E, Y41H, M82T	3.815
Monkeys (Americas)	Y41H, Q42E, M42T	3.506

588 **B)**

Mutation	ΔΔG (kcal/mol)[a]
Y41H	1.929
Q42E	0.954
M82T	0.938
D38E	0.651
Q24L	-0.753
H34L	-0.566
H34Y	-0.139
D30E	0.692

589 [a]Mutations were analyzed with SSIPe server (https://zhanglab.ccmb.med.umich.edu/SSIPe/)
590 and PDB file 6M0J.

592   **Table 2.** Results of codeml analyses of adaptive evolution across ACE2 gene sequences.

Model	Foreground Branch	ω	proportion of sites	LRT	p	positively selected sites[a,b]
clade C	n/a	$\omega_0 = 0.059$, $\omega_1 = 1.000$, $\omega_2 = 0.081$, $\omega_3 = 1.123$, $\omega_4 = 0.236$, $\omega_5 = 1.346$	$p_0 = 0.581$, $p_1 = 0.331$, $p_{2\text{-}5} = 0.089$	26.726	<0.001	n/a
branch-site	platyrrhines	background: $\omega_0 = 0.076$, $\omega_1 = 1.000$, $\omega_{2a} = 0.076$, $\omega_{2b} = 1.000$; foreground: $\omega_0 = 0.076$, $\omega_1 = 1.000$, $\omega_{2a} = 6.218$, $\omega_{2b} = 6.218$	$p_0 = 0.638$, $p_1 = 0.359$, $p_{2a} = 0.002$, $p_{2b} = 0.001$	0.633	0.427	none
	catarrhines	background: $\omega_0 = 0.075$, $\omega_1 = 1.000$, $\omega_{2a} = 0.075$, $\omega_{2b} = 1.000$; foreground: $\omega_0 = 0.075$, $\omega_1 = 1.000$, $\omega_{2a} = 8.988$, $\omega_{2b} = 8.988$	$p_0 = 0.631$, $p_1 = 0.356$, $p_{2a} = 0.009$, $p_{2b} = 0.005$	14.546	0.0001	249M (0.962*), 653A (0.958*), 658V (0.957*)
	strepsirrhines	background: $\omega_0 = 0.072$, $\omega_1 = 1.000$, $\omega_{2a} = 0.072$, $\omega_{2b} = 1.000$; foreground: $\omega_0 = 0.072$, $\omega_1 = 1.000$, $\omega_{2a} = 1.384$, $\omega_{2b} = 1.384$	$p_0 = 0.607$, $p_1 = 0.316$, $p_{2a} = 0.051$, $p_{2b} = 0.027$	0.833	0.361	none
	bats	background: $\omega_0 = 0.075$, $\omega_1 = 1.000$, $\omega_{2a} = 0.075$, $\omega_{2b} = 1.000$; foreground: $\omega_0 = 0.075$, $\omega_1 = 1.000$, $\omega_{2a} = 10.535$, $\omega_{2b} = 10.535$	$p_0 = 0.626$, $p_1 = 0.338$, $p_{2a} = 0.024$, $p_{2b} = 0.013$	42.649	<0.001	24Q (0.998**), 31E (0.959*), 35E (0.974*), 298V (0.959*), 568L (0.998**), 575G (0.965*)

593   [a]Sites with posterior probability >0.95 are shown. [b]The amino acid shown reflects the residue

594   present at the site in the first sequence of the alignment (*Alouatta palliata*).

Figure 1. ACE2 protein sequence alignment and evolutionary relationships of study species. Branch lengths represent evolutionary distance (time, in millions of years) estimated from TimeTree[45]. We outline amino acid residues at critical binding sites for the SARS-CoV-2 spike receptor binding domain. Solid outlines highlight sites predicted to have the most substantial impact on viral binding affinity. Notably, protein sequences of catarrhine primates are highly conserved, including uniformity among amino acids at all binding sites. Primate species that are able to be successfully infected with COVID-19 are indicated in red. Predicted susceptibility to COVID-19 for other primates is additionally coded by terminal branch colors.

Figure 2. Model of human ACE2 in complex with SARS-CoV-2 RBD. Key ACE2 interfacial residues are highlighted. (A). Interactions at critical binding sites 41 and 42 are shown for the residues found in all catarrhines (apes and monkeys in Africa and Asia); (B), and for the residues found in all platyrrhines (monkeys in the Americas) (C). The dashed lines indicate predicted hydrogen bonding interactions. Y41 participates in extensive van der Waals and hydrogen bonding interactions with RBD; these interactions are abrogated with histidine. Q42 side chain amide serves as a hydrogen acceptor and donor to contact RBD; change to glutamic acid diminishes the hydrogen bonding interactions.