bHLH11 inhibits bHLH IVc proteins by recruiting the TOPLESS/TOPLESS-RELATED corepressors in Arabidopsis

Yang Li[a,b,1], Rihua Lei[a,b,1], Mengna Pu[a,b,c], Yuerong Cai[a,b,c], Chengkai Lu[a,b], Zhifang Li[d], Gang Liang[a, b, c, 2]

[a]CAS Key Laboratory of Tropical Plant Resources and Sustainable Use, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Kunming, Yunnan 650223, China

[b]Center of Economic Botany, Core Botanical Gardens, Chinese Academy of Sciences, Mengla, Yunnan 666303, China

[c]The College of Life Sciences, University of Chinese Academy of Sciences, Beijing 100049, China

[d]State Key Laboratory of Cotton Biology, State Key Laboratory of Crop Stress Adaptation and Improvement, School of Life Sciences, Henan University, No. 85 Minglun Street, Kaifeng, Henan 475001, China

[1]These authors contributed equally to this work

[2]Correspondence: lianggang@xtbg.ac.cn

Running title: bHLH11 acts an active repressor

One-sentence summary: bHLH IVc proteins promote the bHLH11 protein accumulation in the nucleus where bHLH11 inhibits the transcriptional activation ability of bHLH IVc via its EAR motifs recruiting the TOPLESS/TOPLESS-RELATED corepressors.

ABSTRACT

Iron (Fe) homeostasis is essential for plant growth and development. Many transcription factors play pivotal roles in the maintenance of Fe homeostasis. bHLH11 was identified as a negative transcription factor regulating Fe homeostasis, however, the underlying molecular mechanism remains elusive. We generated two loss-of-function *bhlh11* mutants which display the enhanced sensitivity to Fe excess, the increased Fe accumulation and the elevated expression of Fe deficiency responsive genes. bHLH11 protein, localized in both the cytoplasm and nucleus, decreases in response to Fe deficiency. Coexpression assays indicate that bHLH IVc transcription factors (TFs) (bHLH34, bHLH104, bHLH105, and bHLH115) facilitate the nuclear accumulation of bHLH11 protein. Further analysis indicates that bHLH11 represses the transactivity of bHLH IVc TFs towards bHLH Ib genes (*bHLH38*, *bHLH39*, *bHLH100*, and *bHLH101*). bHLH11 contains two EAR motifs which are responsible for the repression function by recruiting the TOPLESS/TOPLESS-RELATED (TPL/TPRs) corepressors. Correspondingly, the expression of Fe uptake genes increases in the *tpr1 tpr4 tpl* mutant. Moreover, genetic analysis reveals that bHLH11 has functions independent of FIT. This study provides insights into the complicate Fe homeostasis signaling network.

Introduction

Iron (Fe) is an indispensable microelement for plant growth and development. Plants acquire Fe from the soil, which has low concentration of Fe available, especially in alkaline environments (Jeong and Guerinot, 2009). As about one-third of the world's cultivated land is calcareous (alkaline), iron deficiency is common for plants (Grotz and Guerinot, 2006). Fe functions in many physiological processes, such as photosynthesis, respiration, hormone biosynthesis, and nitrogen fixation. Fe deficiency causes symptoms including delayed growth and leaf chlorosis and can affect the yield and nutritional quality of crops (Kobayashi and Nishizawa, 2014). Although Fe is required for plant growth and development, Fe excess can be toxic to plants because Fe can cause the production of reactive oxygen radicals that are harmful to plant cells (Quinet et al., 2012). Therefore, maintaining Fe homeostasis in plant cells is crucial for their normal growth and development.

Plants have evolved a set of molecular mechanisms for iron absorption, transport, distribution, and storage that ensure appropriate Fe concentrations in cells under low Fe availability. Dicotyledonous and non-gramineous monocotyledonous plants take up Fe using a reduction strategy (strategy I). In *Arabidopsis thaliana*, this strategy involves rhizosphere acidification, ferric iron reduction, and ferrous iron transport. H^+-ATPases such as the P-type ATPase AHA2/AHA7 release protons into the soil, which improves the solubility of Fe in the soil (Santi and Schmidt, 2009; Kobayashi and Nishizawa, 2012). Then, the root surface Fe chelate reductase FERRIC REDUCTION OXIDASE2 (FRO2) catalyzes the reduction of Fe^{3+} to Fe^{2+} (Robinson et al., 1999). IRON-REGULATED TRANSPORTER1 (IRT1) transports Fe^{2+} into roots (Henriques et al., 2002; Varotto et al., 2002; Vert et al., 2002). By contrast, gramineous plants employ a chelation strategy (strategy II) in which high-affinity Fe chelators of the mugineic acid family, also known as phytosiderophores, are secreted into the rhizosphere and facilitate the uptake of the Fe^{3+}-phytosiderophore complex. Recent studies suggest that secretion

of Fe-chelating compounds is also important for the survival of non-gramineous plants such as Arabidopsis in alkaline soil (Rodríguez-Celm et al., 2013; Schmidt et al., 2014; Fourcroy et al., 2014, 2016; Siwinska et al., 2018; Tsai et al., 2018).

To maintain Fe homeostasis, plants must sense the environmental Fe concentration and fine-tune the expression of Fe uptake-associated genes accordingly. BRUTUS (BTS) interacts with the basic helix-loop-helix transcription factors bHLH105 and bHLH115 and promotes their degradation (Selote et al., 2015). IRON MAN (IMA), a class of peptides, interact with and inhibit BTS, facilitating the accumulation of bHLH105 and bHLH115 (Grillet et al., 2018; Li et al., 2021). bHLH105 and bHLH115 belong to the bHLH IVc group, which contains four members. The other two members are bHLH34 and bHLH104. These four members regulate the expression of *FER-LIKE IRON-DEFICIENCY-INDUCED TRANSCRIPTION FACTOR* (*FIT*), bHLH Ib genes (*bHLH38*, *bHLH39*, *bHLH100*, and *bHLH101*) and *POPEYE* (*PYE*) (Zhang et al., 2015; Li et al., 2016; Liang et al., 2017). bHLH121 interacts with bHLH IVc proteins and is required for the maintenance of Fe homeostasis (Kim et al., 2019; Gao et al., 2020; Lei et al., 2020). Downstream of bHLH IVc and bHLH121, FIT interacts with bHLH Ib TFs to promote the expression of Fe-uptake associated genes *IRT1* and *FRO2* (Yuan et al., 2008; Wang et al., 2013). In contrast, PYE and bHLH11 are negative regulators of Fe homeostasis (Long et al., 2010; Tanabe et al., 2019). Additionally, bHLH105 also functions as a negative regulator when it interacts with PYE (Tissot et al., 2019). There is also a similar Fe deficiency response signaling network in rice (Ogo et al., 2007; Kobayashi 2013, 2019; Zhang et al., 2017, 2020; Wang et al., 2020; Li et al., 2020).

The overexpression of *bHLH11* causes the dramatic decline of Fe uptake genes including *IRT1* and *FRO2*, and the severe Fe deficiency symptoms (Tanabe et al., 2019). However, the molecular mechanism by which bHLH11 regulates Fe homeostasis remains elusive. In the present study, we

characterized the roles of bHLH11 in the maintenance of Fe homeostasis in Arabidopsis. bHLH11 is localized in both the cytoplasm and nucleus and is exclusively in the nucleus when bHLH IVc TFs are abundant. bHLH11 also interacts with and inhibits bHLH IVc TFs. bHLH11 exerts its transcriptional repression function by its two EAR motifs recruiting the transcriptional TOPLESS/TOPLESS-RELATED corepressors.

Results

Loss-of-function of *bHLH11* impairs Fe homeostasis

It was reported that the overexpression of *bHLH11* leads to the enhanced sensitivity to Fe deficiency (Tanabe et al., 2019). To further explore the molecular mechanism of bHLH11 regulating Fe homeostasis, we employed the CRISPR-Cas9 system to edit *bHLH11*. Two single guide RNAs were designed to specifically target exons 4 and 3 of *bHLH11* and respectively integrated into the binary vector with a Cas9 (Liang et al., 2016) which were then used for transformation of wild type plants. We identified two homozygous mutants (*bhlh11-1* and *bhlh11-2*), the former containing a 1-bp insertion in exon 4 and the latter containing a 2-bp deletion in exon 3 (Figure S1A), both of which caused a frameshift mutation (Figure S1B). Expression analysis indicated that bHLH11 mRNA levels did not change in these two mutants (Figure S1C). When grown on Fe0 (Fe free) or Fe100 (100 μM Fe^{2+}) medium, no visible difference was observed between the *bhlh11* mutants and wild type (Figure 1A). By contrast, when grown on Fe300 (300 μM Fe^{2+}) medium, the *bhlh11* mutants produced low shoot biomass (Figure 1A, B). The *bHLH11* driven by its native promoter rescued the sensitivity of *bhlh11-1* to Fe excess (Figure S1D). Fe content analysis suggested that the Fe concentration of *bhlh11* mutants was higher than that of the wild type (Figure 1C). These data suggest that the loss-of-function of *bHLH11* leads to the enhanced sensitivity to Fe excess.

To further investigate the effect of bHLH11 on the Fe signaling network, we examined the expression of several Fe homeostasis associated genes (Figure

1D). The expression of *IRT1*, *FRO2*, *bHLH38*, *bHLH39*, and *bHLH100* was higher in the *bhlh11* mutants than in the wild type regardless of Fe status. These data further support the negative regulation function of *bHLH11* in the Fe homeostasis.

bHLH11 expression and subcellular localization

To investigate the response of *bHLH11* to Fe status, RT-qPCR was used to determine the expression of *bHLH11* in response to Fe status, showing that *bHLH11* mRNA increased in the roots with an increase of Fe concentration in the growth medium (Figure 2A), which is in consistence with the previous study (Tanabe et al., 2019). To examine the response of bHLH11 protein to Fe status, the *bHLH11* overexpression construct was introduced into wild type plants. In agreement with the previous report (Tanabe et al., 2019), *bHLH11* overexpression plants were more sensitive to Fe deficiency compared with wild type (Figure S2). One-week-old *bHLH11* overexpression plants grown on Fe100 medium were transferred to Fe0 or Fe300 medium, and root samples were harvested after 1, 2, and 3 days. Immunoblot analysis showed that bHLH11 increased with an increase in Fe concentration and decreased with a decrease in Fe concentration (Figure 2B).

Several Fe-homeostasis associated bHLH TFs were found outside the nucleus (Gratz et al., 2019; Trofimov et al., 2019; Lei et al., 2020; Wang et al., 2020; Liang et al., 2020). To determine the subcellular localization of bHLH11, we generated the *35S:bHLH11-mCherry* construct, in which the mCherry tag was fused in frame with the C end of bHLH11. When this construct was transiently expressed in tobacco leaves, mCherry was mainly observed in the cytoplasm and nucleus, which is very similar to that of free mCherry (Figure 2C). The cytoplasmic localization of bHLH11 was unexpected because transcription factors are known to function in the nucleus. Thus, we examined whether bHLH11 can be retained in the cytoplasm due to a lack of a nuclear localization signal (NLS). NLS prediction was conducted by cNLS Mapper with

a   cutoff   score   =   4   (Kosugi   et   al.,   2009;

http://nls-mapper.iab.keio.ac.jp/cgi-bin/NLS_Mapper_y.cgi).   No   NLS   was

found in bHLH11. When bHLH11 was fused with NLS-mCherry, which contains

an NLS from the SV40 virus, bHLH11-NLS-mCherry was exclusively localized

in the nucleus (Figure 2C). These data suggest that the lack of an NLS causes

the cytoplasmic localization of bHLH11.

bHLH11 interacts with bHLH IVc TFs in the nucleus

Considering that TFs usually function in the nucleus and an NLS allows

bHLH11 to remain in the nucleus, we hypothesized that bHLH11 might be

recruited to the nucleus by its nuclear-localized interaction partners. Recent

studies revealed that bHLH121, the closest homolog of bHLH11, interacts with

bHLH IVc TFs (Kim et al., 2019; Gao et al., 2020; Lei et al., 2020). Therefore,

we employed the yeast two-hybrid system to test whether bHLH11 interacts

with bHLH IVc TFs. The bHLH11 protein was fused with the GAL4 DNA

binding domain in the pGBKT7 vector as the bait (BD-bHLH11). bHLH IVc TFs

were cloned to the pGADT7 vector as the preys. As expected, bHLH11

interacts with all four bHLH IVc TFs in yeast (Figure 3A).

To confirm that bHLH IVc TFs interact with bHLH11 in plant cells, we

employed the tripartite split-GFP system (Liu et al., 2018). The GFP10

fragment was fused with bHLH IVc proteins in their N-end (GFP10-bHLH IVc)

and the GFP11 was fused with bHLH11 in the C-end (bHLH11-GFP11). As a

control, bHLH121 was fused with the GFP10. When GFP10-bHLH IVc and

bHLH11-GFP11 were transiently co-expressed with GFP1-9 in tobacco leaves,

the GFP signal was visible in the nucleus of transformed cells (Figure 3B). By

contrast, the combination GFP10-bHLH121/bHLH11-GFP11/GFP1-9 did not

result in visible GFP signal. The negative controls did not result in a GFP signal

in the cells (Figure S3).

We next performed a coimmunoprecipitation (Co-IP) assay to confirm the

interactions between bHLH IVc TFs and bHLH11 (Figure 3C). MYC tag-fused

bHLH IVc TFs and HA tag-fused bHLH11 were transiently co-expressed in tobacco leaves. The total proteins were incubated with MYC antibody and A/G-agarose beads and then separated on SDS-PAGE for immunoblotting with HA antibody. Consistent with the results from the yeast two-hybrid and tripartite split-GFP assays, bHLH IVc and bHLH11 were present in the same protein complex. These data suggest that bHLH IVc TFs physically interact with bHLH11.

bHLH IVc TFs affect the subcellular localization of bHLH11

Having confirmed that bHLH11 interacts with bHLH IVc TFs in the nucleus, we wondered whether the bHLH IVc TFs have an impact on the subcellular localization of bHLH11. When any one of these four GFP tagged proteins was respectively co-expressed with bHLH11-mCherry, bHLH11-mCherry accumulated exclusively in the nucleus (Figure 4A). By contrast, co-expression of the free GFP did not affect the subcellular localization of bHLH11-mCherry (Figure 4A).

To further confirm the distribution of bHLH11 in the cytoplasm and nucleus, we used immunoblot to measure the expression of the bHLH11 protein. As shown in Figure 4B, bHLH11 protein was detected both in the nucleus and cytoplasm, and both its nuclear and cytoplasmic counterparts were responsive to Fe status.

bHLH11 antagonizes the transactivity of bHLH IVc TFs

The bHLH Ib genes are activated directly by the bHLH IVc TFs (Zhang et al., 2015; Li et al., 2016; Liang et al., 2017). Our expression analyses also suggested that the bHLH Ib genes were upregulated in the *bhlh11* mutants, implying that bHLH11 is a negative regulator of bHLH Ib genes. Because bHLH11 interacts with the bHLH IVc TFs, we proposed that bHLH11 could antagonize the functions of the bHLH IVc TFs. To confirm this hypothesis, transient expression assays were conducted in Arabidopsis protoplasts (Figure

5A). The reporter construct *ProbHLH38:LUC*, in which the LUC reporter was fused with the promoter of *bHLH38*, and different effectors in which the 35S promoter was used to drive GFP, bHLH11 or bHLH IVc, were used in the transient assays. Compared with GFP, bHLH IVc TFs activated the expression of *ProbHLH38:LUC*, whereas bHLH11 had no significant effect. When bHLH11 and bHLH IVc were co-expressed, the LUC/REN ratio declined significantly. These data suggest that bHLH11 inhibits the transactivity of bHLH IVc TFs towards *bHLH38*.

To further investigate whether bHLH11 inhibits the functions of bHLH IVc TFs by direct protein–protein interaction, we employed the *pGAL4* promoter. In the reporter construct, GFP fused with an NLS sequence was driven by *pGAL4* containing the minimal CaMV 35S promoter with five repeats of the GAL4 binding motif (Figure 5B). In the effectors, the DNA binding domain (BD) of GAL4 was fused in frame with either bHLH104 or bHLH105 and driven by the 35S promoter. Consistent with the fact that bHLH IVc TFs are transcriptional activators, the chimeric BD-bHLH104 or bHLH105 activated the expression of *GFP*. When bHLH11 was co-expressed with BD-bHLH104 or BD-bHLH105, the expression of *GFP* was significantly repressed. These data suggest that bHLH11 antagonizes the transcriptional activation ability of bHLH IVc TFs through direct protein interaction.

To further confirm the antagonistic role of bHLH11 to bHLH IVc TFs, we generated *bHLH11-OX/bHLH104-OX* plants by crossing *bHLH11-OX-20* with *bHLH104-OX*. Compared with *bHLH104-OX*, the tolerance of *bHLH11-OX/bHLH104-OX* to Fe deficiency was reduced (Figure S4). Taken together, our data suggest that bHLH11 can antagonizes the functions of bHLH IVc TFs.

It is reported that *bHLH11* overexpression causes the increased expression of bHLH Ib genes (Tanabe et al., 2019), which seems to be contrary to the results above that bHLH11 represses the expression of *bHLH38* (Figure 5A). We reasoned that the upregulation of bHLH Ib genes was not a direct result

from the *bHLH11* overexpression, but caused by a secondary effect of the

disrupted Fe homeostasis in *bHLH11-OX* plants. To avoid the secondary effect,

we generated transgenic plants containing a *pER8-bHLH11* construct, in which

the *HA-bHLH11* fusion gene was under the control of an inducible promoter,

activated by estradiol. Under Fe deficient conditions, the *pER8-bHLH11* plants

grew as well as the wild type, and the *bHLH11* transcript level was similar

between the wild type and *pER8-bHLH11*. After treatment with estradiol, the

bHLH11 gene was overexpressed in the *pER8-bHLH11* plants (Figure S5A).

As expected, the *pER8-bHLH11* plants displayed the enhanced sensitivity to

Fe deficiency when gown on Fe0 + estradiol medium (Figure S5B). To

examine the expression of Fe deficiency-responsive genes, seven-day-old

plants grown on Fe0 medium were transferred to Fe0 medium with or without

estradiol for 6 h and roots were isolated for analysis. We found that the

expression of bHLH Ib genes was downregulated in the *pER8-bHLH11* plants

after treatment with estradiol (Figure 5C). Taken together, our data suggest

that bHLH11 antagonizes the transcriptional activation ability of bHLH IVc TFs

to bHLH Ib genes.

The repression function of bHLH11 requires its EAR motifs

Considering that bHLH11 negatively regulates the expression of Fe

deficiency-responsive genes, we wanted to know how bHLH11 acts as a

transcriptional repressor. There are two typical ethylene-responsive element

binding factor-associated amphiphilic repression (EAR) motifs (LxLxL) in the

C-terminal region of bHLH11 (Figure 6A). It is known that EAR motifs account

for the repression function of many transcription factors (Kagale et al., 2010).

To investigate whether the EAR motifs are required for the repression function

of bHLH11, we conducted reporter–effector transient expression assays in

which bHLH105 was used as an effector to activate *ProbHLH38-LUC* (Figure

6B). We compared the effects of GFP, bHLH11, bHLH11dm (a version

containing two mutated EAR motifs), and bHLH11dm-VP16 (VP16, an

established activation domain) on bHLH105. Compared with the significant repression effect of bHLH11 on bHLH105, bHLH11dm had no significant effect, whereas bHLH11dm-VP16 enhanced the transactivation function of bHLH105 (Figure 6C). These data suggest that bHLH11 functions as a transcriptional repressor and that this function is dependent on its EAR motifs.

To assess the consequences of disrupting the repression functions of bHLH11 *in vivo*, we generated *bHLH11dm-VP16* transgenic plants. The *bHLH11dm-VP16* plants showed the enhanced tolerance to Fe deficiency compared to *bHLH11-OX* plants (Figure 6D). Correspondingly, the expression of *IRT1* and *FRO2* was activated in the *bHLH11dm-VP16* plants (Figure S6). Taken together, the EAR motifs are needed for the repression function of bHLH11.

bHLH11 interacts with the transcription corepressors TPL/TPRs

The EAR motif is a characteristic of proteins interacting with the TPL/TPRs which function as transcription corepressors (Szemenyei et al., 2008; Pauwels et al., 2010; Causier et al., 2012). Thus, we determined whether bHLH11 interacts with TPL/TPRs. Yeast two-hybrid assays indicated that bHLH11 interacts with TPL/TPRs (Figure 7A). To further investigate whether the EAR motifs are required for the interaction, the various EAR-mutated versions, bHLH11m1, bHLH11m2, and bHLH11dm, were tested. The results indicated that the interaction between bHLH11 and TPL/TPRs was dependent on the EAR motifs, as the mutation of both EAR motifs abolished the interaction between bHLH11 and TPL/TPRs.

Next, we wanted to know whether TPL/TPRs participate in the regulation of Fe homeostasis. To this aim, the *tpr1 tpr4 tpl* triple mutant plants were used for phenotypic analysis. When grown on Fe0 or Fe100 medium, no visible difference was observed. In contrast, when grown on Fe300 medium, the shoot biomass of the *tpr1 tpr4 tpl* triple mutant plants was higher than that of wild type (Figure 7B, C). The measurement of Fe concentration showed that

the *tpr1 tpr4 tpl* triple mutant plants had higher Fe concentration than wild type (Figure 7D). We also examined the expression of Fe deficiency responsive genes, finding that the expression of *IRT1* and *FRO2* was higher in the *tpr1 tpr4 tpl* than that in the wild type (Figure 7E). These data suggest that TPL/TPRs negatively regulate the expression of Fe uptake genes.

Genetic relationship between *bHLH11* and *FIT*

Tanabe et al. (2019) reported that bHLH11 negatively regulates Fe uptake by repressing *FIT* transcription. To explore whether the function of bHLH11 depends on FIT, we conducted genetic analysis. The *bHLH11-OX-20* line was crossed into the *fit-2* mutant, and homozygous *bHLH11-OX-20/fit-2* plants were identified. When grown on Fe100 or Fe300 medium, no visible difference was observed among the wild type, *fit-2*, *bHLH20-OX-20* and *bHLH11-OX-20/fit-2* plants (Figure 8A; Figure 7A). However, when grown on Fe0 medium, the *bHLH11-OX-20/fit-2* plants were more sensitive to Fe deficiency than *fit-2* and *bHLH11-OX-20* plants, as shown by the shorter roots and bleached leaves (Figure 8A; Figure S7B). These data suggest that bHLH11 has roles independent of FIT.

Subsequently, we examined the expression of *IRT1*, *FRO2* and bHLH Ib (Figure 8B). Under Fe deficient conditions, the expression of *IRT1* and *FRO2* was lower in the *fit-2* than that in the *bHLH11-OX-20,* but as low as that in the *bHLH11-OX-20/fit-2.* In contrast, the expression of bHLH Ib genes was similar among *fit-2, bHLH20-OX-20* and *bHLH11-OX-20/fit-2* under Fe deficient conditions. Under Fe sufficient conditions, although the expression of bHLH Ib genes was higher in the *fit-2*, the introduction of *bHLH11-OX-20* significantly repressed their expression. These data suggest that bHLH11 represses the expression of bHLH Ib genes independently of FIT under Fe sufficient conditions.

Discussion

Plants sense Fe-deficient environments and activate a signal transduction

cascade that ultimately results in the transcriptional regulation of downstream

effector genes of the Fe uptake system. The expression of Fe

homeostasis-associated genes is tightly regulated by Fe availability, including

environmental Fe availability and local Fe availability in developing tissues and

organs. However, this mechanism is not an on-off process but rather a

fine-tuned one, with multiple layers of transcription regulations. Considerable

progress has been made in deciphering the signal transduction pathways that

maintain Fe homeostasis, leading to the identification of many signaling

components. Here, we show that bHLH11 acts an active repressor by

recruiting TPL/TPRs. bHLH11 contributes to Fe homeostasis by repressing

bHLH IVc TFs.

The antagonistic regulation between positive and negative TFs is prevalent

in plants. For example, the positive TFs MYC2/MYC3/MYC4 and the negative

TFs bHLH3/bHLH13/bHLH14/bHLH17 antagonistically modulate jasmonic

acid signaling (Fernandez-Calvo et al., 2011; Song et al., 2013). In Fe

homeostasis signaling, the bHLH IVa TFs (bHLH18, bHLH19, bHLH20, and

bHLH25) antagonize the bHLH Ib TFs to regulate FIT protein stability under Fe

deficiency (Cui et al., 2018). We show the antagonistic function of bHLH11 to

bHLH IVc TFs, which may explain why *bHLH11-OX* plants display the severe

Fe deficiency phenotypes similar to those of bHLH IVc mutants (Liang et al.,

2017). In addition to transcriptional regulation, the protein degradation is

another type of regulation in Fe homeostasis signaling. As reported previously,

bHLH105 and bHLH115 are degraded by BTS (Selote et al., 2015), and FIT by

BTSL1 and BTSL2 (Rodríguez-Celma et al., 2019). We found that bHLH11

protein decreased under Fe deficient conditions (Figure 2B), which may

benefit plants by alleviating the repression of bHLH11 to Fe uptake associated

genes. Further investigation is required to understand the post-transcription

regulation of *bHLH11* in response to Fe deficiency. These coordinated

regulations of transcription and post-transcription may help plants adapt to

their various Fe-nutrition habitats.

Although bHLH Ib genes were upregulated in both the *bHLH11-OX* and

bhlh11 mutant plants, we provide evidence supporting that bHLH11 negatively

regulates bHLH Ib genes: (1) bHLH11 represses the promoter of *bHLH38* in

the transient expression assays (Figure 5A); (2) the transient induction of

bHLH11 reduced the abundance of bHLH Ib genes in the *pER8-bHLH11* plants

(Figure 5C); (3) the overexpression of *bHLH11* reduced the levels of bHLH Ib

genes in the *fit-2* under Fe sufficient conditions (Figure 8B). We proposed that

the upregulation of bHLH Ib genes may result from a feedback regulation that

the severe Fe deficient status of *bHLH11-OX* plants activates bHLH Ib genes.

In fact, feedback regulations are universal in Fe homeostasis. For example,

bHLH Ib genes are also activated in the *irt1*, *frd3*, *opt3* and *fit* which are

defective in Fe uptake or transportation (Wang et al., 2007). Therefore, our

data suggest that the abundance of bHLH Ib genes are balanced by bHLH IVc

TFs and bHLH11. We demonstrated that bHLH11 negatively regulates the

expression of *IRT1* and *FRO2*, which is in consistence with the conclusion that

bHLH11 represses the FIT-dependent Fe uptake (Tanabe et al., 2019).

Meanwhile, bHLH11 also plays roles independent of FIT since its

overexpression made *fit-2* more sensitive to Fe deficiency (Figure 8A), which is

reasonable because bHLH11 can inhibit bHLH IVc TFs.

bHLH11 has no canonical NLS sequence. bHLH11 protein exists in the

nucleus and cytoplasm, and it accumulates in the nucleus with the assistance

of bHLH IVc TFs (Figure 4A). This interaction dependent nuclear localization of

bHLH11 might contribute to the maintenance of Fe homeostasis. bHLH11

inhibits the activation ability of bHLH IVc TFs and restricts the expression of Fe

uptake-associated genes. The repression function of bHLH11 may help plants

avoid Fe toxicity and adapt to environments with excessive Fe by reducing the

rate of Fe uptake. This putative protein shuttling strategy enables plants to

respond quickly to Fe fluctuation.

Two types of transcriptional repressors exist: active and passive (Krogan

and Long, 2009). Generally, active repressors repress transcription by recruiting transcriptional repression components, whereas passive repressors indirectly influence transcription by competitively interfering with activators. TPL/TPRs are a class of corepressors and usually recruited by the EAR motif containing transcriptional repressors (Kagale et al., 2010; Causier et al., 2012). bHLH11 functions negatively and has two EAR motifs, both of which can interact with TPL/TPRs, suggesting that bHLH11 is an active repressor. The observation that the EAR motif is conserved in bHLH11 homologs across different plant species, such as *Zea mays*, *Oryza sativa*, and *Brassica rapa* (Figure S8), implies that different plant species employ a conserved repression strategy to fine-tune Fe homeostasis. In Fe homeostasis signaling pathway, PYE is a negative regulator of Fe homeostasis associated genes *ZIF1*, *FRO3* and *NAS4* (Long et al., 2010), and an EAR motif was found in its C-terminal region (Kagale et al., 2010). ZAT12 (ZINC FINGER OF ARABIDOPSIS THALIANA 12) functions as a negative regulator of Fe uptake, and contains an EAR motif responsible for the interaction with FIT (Le et al., 2016). By contrast, EIN3 (ETHYLENE-INSENSITIVE 3), also containing an EAR motif (Kagale et al., 2010), is a positive regulator of Fe homeostasis by interacting with FIT (Lingam et al., 2011). It is likely that these TFs also recruit TPL/TPRs. In support of this hypothesis, the *tpr1 tpr4 tpl* plants are also sensitive to Fe excess to a lesser extent compared with the *bhlh11* plants, as shown by the shoot biomass (Figure 1B; Figure 7C).

This study expands our knowledge of the Fe homeostasis transcription network mediated by bHLH Ib and IVc proteins. Based on our findings, we propose a putative working model for bHLH11 (Figure 9). When Fe is sufficient, bHLH11 mRNAs increase and its proteins are stable. To limit Fe uptake, bHLH IVc proteins facilitate the accumulation of bHLH11 in the nucleus, where bHLH11 recruits TPL/TPR corepressors to repress the activation of bHLH IVc proteins to bHLH Ib genes, and then the reduction of bHLH Ib genes results in the down-regulation of Fe uptake genes *IRT1* and *FRO2.* When Fe is limited,

bHLH11 proteins decrease rapidly and few bHLH11 proteins enter the nucleus

to inhibit bHLH IVc. Finally, the bHLH Ib proteins accumulate and promote the

expression of *IRT1* and *FRO2*. This enables plants to control Fe uptake and

maintain Fe homeostasis. Our study provides experimental support for the

existence of an elaborate system that allows plants to respond dynamically to

Fe status. This mechanism is based on an equilibrium between the activation

of Fe uptake-associated genes by bHLH IVc and their repression by bHLH11.

Materials and methods

Plant materials and growth conditions

Arabidopsis thaliana ecotype Col-0 was used as the wild type in this study. *fit-2*

was described previously (Lei et al., 2020). *tpr1 tpr4 tpl* (N72353) was obtained

from NASC. Plants were grown in long photoperiods (16-hour light/8-hour dark)

or short photoperiods (10-hour light/14-hour dark) at 22°C. Surface sterilized

seeds were stratified at 4°C for 2 d before being planted on medium. Half

Murashige and Skoog (MS) medium with 1% sucrose, 0.8% agar A and the

indicated Fe^{2+}EDTA concentration were used. Fe0 (0 µM Fe^{2+}), Fe50 (50 µM

Fe^{2+}), Fe100 (100 µM Fe^{2+}) and Fe300 (300 µM Fe^{2+}).

Generation of CRISPR/Cas9-edited *bHLH11*

For CRISPR/Cas9-mediated editing of *bHLH11*, two target sites were

designed by CRISPR-GE (Xie et al., 2017) to target the third and fourth exon

of bHLH11, which were driven by the AtU3b promoter and respectively cloned

into the pMH-SA binary vector carrying the Cas9 (Liang et al., 2016). The wild

type plants were transformed and positive transgenic plants were selected on

half-strength MS medium containing 20 µg/mL hygromycin. The positive

transformants were sequenced and the homozygous mutants without the Cas9

were selected for further analysis.

Generation of transgenic plants

HA-tag or VP16 domain were fused in frame with the full-length coding

sequence of *bHLH11* to generate *35S:HA-bHLH11* and *35S:bHLH11dm-VP16*

in the pOCA30 binary vector. *HA-bHLH11* was cloned into pER8 vector (Zuo et

al., 2001). These constructs were introduced into *Agrobacterium tumefaciens*

(EHA105) respectivley and then used for transformation in the wild type

Arabidopsis. For complementation of *bhlh11-1*, the 3kb DNA fragment

upstream of *bHLH11* translation start site was used to drive *HA-bHLH11-GFP*

and then introduced into the *bhlh11-1* by *A. tumefaciens* mediated

transformation.

Yeast-two-hybrid assays

Full-length bHLH11 was cloned into pGBKT7 as a bait. The full-length of bHLH

IVc in the pGADT7 was described previously (Li et al., 2016). Growth was

determined as described in the Yeast Two-Hybrid System User Manual

(Clontech).

Subcellular localization

For the construction of *35S:bHLH11-mCherry*, mCherry-tag was fused with

bHLH11. *35S:bHLH34-GFP*, *35S:bHLH104-GFP*, *35S:bHLH105-GFP*,

35S:bHLH115-GFP, and *35S:GFP* were described previously (Lei et al., 2020).

35S:bHLH11-mCherry was co-expressed with various GFP-containing vectors

in tobacco cells. Epidermal cells were recorded on an OLYMPUS confocal

microscope. Excitation laser wave lengths of 488 nm and 563 nm were used

for imaging GFP and mCherry signals, respectively.

Fluorescence complementation assays

The tripartite split-GFP fluorescence complementation assay was described as

previously (Lei et al., 2020). The C-end of bHLH11 was fused with the GFP11

tag. The N-end of bHLH IVc and bHLH121 was fused with the GFP10 tag. All

vectors were introduced into *A. tumefaciens* (strain EHA105) and the various

combinations of Agrobacterial cells were infiltrated into leaves of *Nicotiana*

benthamiana by an infiltration buffer (0.2 mM acetosyringone, 10 mM $MgCl_2$,

and 10 mM MES, PH 5.6). Gene expression was induced 1 day after

agroinfiltration by injecting 20 μM *β*-estradiol in the abaxial side of the leaves.

Fluorescence of epidermal cells was recorded on a Carl Zeiss Microscopy.

Co-immunoprecipitation assay

HA-bHLH11 and MYC-bHLH IVc or MYC-GFP were transiently expressed in

the *N. benthamiana* leaves and the leaves were infiltrated with MG132 12

hours before harvesting. 2 g leaf samples were used for protein extraction in 2

ml IP buffer (50 m M Tris-HCl, pH 7.4, 150 mM NaCl, 1 mM $MgCl_2$, 20%

glycerol, 0.2% NP-40, 1 X protease inhibitor cocktail and 1 X phosphatase

inhibitor cocktail from Roche). Lysates were clarified by centrifugation at 20,

000 g for 15 min at 4 °C and were immunoprecipitated using MYC antibody. IP

proteins were analyzed by immunoblot using anti-HA and anti-MYC antibody

respectively (Affinity Biosciences).

Gene expression analysis

Total root RNA was extracted by the use of the Trizol reagent (Invitrogen). For

the reverse transcription reaction, 1 μg total RNA was used for cDNA synthesis

by oligo(dT)18 primer according to the manufacturer's protocol (Takara). The

resulting cDNA was subjected to relative quantitative PCR using the ChamQ^{TM}

SYBR qPCR Master Mix (Vazyme Biotech Co.,Ltd) on a Roche LightCycler

480 real-time PCR machine, according to the manufacturer's instructions. For

gene expression analysis in Arabidopsis plants, the relative level of genes was

normalized to *ACT2* and *TUB2*. For the quantification of each gene, three

biological replicates were used. The primers used for quantitative reverse

transcription-PCR are listed in Table S1.

Fe concentration measurement

To determine Fe concentration, rosette leaves from three-week-old seedlings grown in soil were harvested and dried at 65 °C for 3 days. About 100 mg dry weight was wet-ashed with 5 ml of 11 M HNO_3 and 1 ml of 12 M $HClO_4$ for 20 min at 220°C. Each sample was diluted to 16 ml with 18 MΩ water and Fe concentration was analyzed on a Thermo SCIENTIFIC ICP-MS(iCAP6300).

Transient expression assays in Arabidopsis protoplasts

Arabidopsis mesophyll protoplasts preparation and subsequent transfection were performed as described previously (Wu et al., 2009). The promoter sequence of *bHLH38* was amplified from genomic DNA and cloned into pGreenII 0800-LUC vector which contains a renillia luciferase encoding gene *REN* driven by the 35S promoter. The coding sequences of GFP and various kinds of bHLHs (bHLH34, bHLH104, bHLH105, bHLH115, bHLH11, bHLH11dm and bHLH11dm-VP16) were respectively cloned into the pGreenII 62-SK vector under control of 35S promoter. For the reporter and effectors, 10 μg plasmid for each construct was used. After protoplast preparation and subsequent transfection, firefly luciferase (LUC) and REN activities were measured using the Dual-Luciferase Reporter Assay System (Promega) following the manufacturer's instructions. Relative (LUC) activity was calculated by normalizing against the REN activity.

Transient expression assays in tobacco

Agrobacterium tumefaciens strains EHA105 was used in the transient expression experiments in tobacco. *pGAL4* promoter and BD domain were described previously (Li et al., 2016). *pGAL4* promoter was fused with NLS-GFP and cloned into the pOCA28 binary vector. 35S:BD, 35S:BD-bHLH104, 35S:BD-bHLH105 and 35S:HA-bHLH11 were constructed in the pOCA30 binary vector. For co-infiltration, different constructs were mixed prior to infiltration. Leaf infiltration was conducted in 3-week-old *N. benthamiana*. *NPTII* gene in the pOCA28 vector was used as the internal

control. *GFP* transcript abundance was normalized to that of *NPTII.*

Immunoblotting

For total protein extraction, roots were ground to a fine powder in liquid

nitrogen and then resuspended and extracted in RIPA buffer (50 mM Tris, 150

mM NaCl, 1% NP-40, 0.5% sodium deoxycholate, 0.1% SDS, 1 mM PMSF, 1 x

protease inhibitor cocktail [pH 8.0]). Isolation of cytoplasmic and nuclear

proteins was performed as described previously (Li et al., 2018). Sample was

loaded onto 12% SDS-PAGE gels and transferred to nitrocellulose membranes.

The membrane was blocked with TBST (10 mM Tris-Cl, 150 mM NaCl, and

0.05% Tween 20, pH8.0) containing 5% nonfat milk (TBSTM) at room

temperature for 60 min and incubated with primary antibody in TBSTM

(overnight at 4°C). Membranes were washed with TBST (three times for 5 min

each) and then incubated with the appropriate horseradish

peroxidase-conjugated secondary antibodies in TBSTM at room temperature

for 1.5 h. After washing three times, bound antibodies were visualized with

ECL substrate.

Supplemental data

Supplemental Figure 1. Identification of *bhlh11* mutants.

Supplemental Figure 2. Identification of *bHLH11* overexpression plants.

Supplemental Figure 3. Negative controls of tripartite split-sfGFP
complementation assays.

Supplemental Figure 4. Antagonism between bHLH104 and bHLH11.

Supplemental Figure 5. Phenotypes of *pER8-bHLH11* transgenic plants.

Supplemental Figure 6. Expression of *IRT1* and *FRO2* in the

bHLH11dm-VP16 plants.

Supplemental Figure 7. Analysis of *bHLH11-OX-20/fit-2* plants.

Supplemental Figure 8. Conserved EAR motifs in the bHLH11 homologs from

various plants species.

Supplemental Table 1. Primers used in this paper.

Acknowledgements

We thank the Biogeochemical Laboratory and Central Laboratory (Xishuangbanna Tropical Botanical Garden) for assistance in the determination of metal contents. We also thank Germplasm Bank of Wild Species in Southwest China for confocal laser scanning microscopy.

Finding

This work was supported by the National Natural Science Foundation of China (31770270) and the Applied Basic Research Project of Yunnan Province (2019FB028 and 202001AT070131).

Author contributions

G.L. conceived the project. Y.L., R.L., M.P., C.L., Z. L., and G.L. constructed plasmids, M.P. and Y.C generated transgenic plants, and Y.L. and R.L. characterized plants, determined gene and protein expression, and conducted cellular assays. Y.L. and G.L. wrote the manuscript and all authors discussed and approved the manuscript.

Conflict of interest statement. None declared.

References

Causier B, Ashworth M, Guo W, Davies B (2012) The TOPLESS interactome: A framework for gene repression in Arabidopsis. Plant Physiol **158:** 423-438

Cui Y, Chen CL, Cui M, Zhou WJ, Wu HL, Ling HQ (2018) Four IVa bHLH Transcription Factors Are Novel Interactors of FIT and Mediate JA Inhibition of Iron Uptake in Arabidopsis. Mol Plant **11:** 1166-1183

Fernández-Calvo P, Chini A, Fernández-Barbero G, Chico JM, Gimenez-Ibanez S, Geerinck J, Eeckhout D, Schweizer F, Godoy M, Franco-Zorrilla JM, et al. (2011) The Arabidopsis bHLH transcription factors MYC3 and MYC4 are targets of JAZ repressors and act additively with MYC2 in the activation of jasmonate responses. Plant Cell **23:**701-715

Fourcroy P, Sisó-Terraza P, Sudre D, Sairón M, Reyt G, Gaymard F, Abadía A, Abadia J, Alvarez-Fernández A, Briat J (2014) Involvement of the ABCG37 transporter in secretion of scopoletin and derivatives by Arabidopsis roots in response to iron deficiency. New Phytol **201:** 155-167

Fourcroy P, Tissot N, Gaymard F, Briat JF, Dubos C (2016) Facilitated Fe Nutrition by Phenolic Compounds Excreted by the Arabidopsis ABCG37/PDR9 Transporter Requires the IRT1/FRO2 High-Affinity Root Fe(2+) Transport System. Mol Plant **9:** 485-488

Gao F, Robe K, Bettembourg M, Navarro N, Rofidal V, Santoni V, Gaymard F, Vignols F, Roschzttardtz H, Izquierdo E, Dubos C (2020) The transcription factor bHLH121 interacts with bHLH105 (ILR3) and its closest homologs to regulate iron homeostasis in Arabidopsis. Plant Cell **32:** 508-524

Gratz R, Manishankar P, Ivanov R, Köster P, Mohr I, Trofimov K, Steinhorst L, Meiser J, Mai HJ, Drerup M, et al. (2019) CIPK11-dependent phosphorylation modulates fit activity to promote arabidopsis iron acquisition in response to calcium signaling. Dev Cell **48:** 726-740

Grillet L, Lan P, Li W, Mokkapati G, Schmidt W (2018) IRON MAN is a ubiquitous family of peptides that control iron transport in plants. Nat Plants **4:** 953-963

Grotz N, Guerinot ML (2006) Molecular aspects of Cu, Fe and Zn homeostasis in plants. Biochim Biophys Acta. 1763:595-608.

Henriques R, Jásik J, Klein M, Martinoia E, Feller U, Schell J, Pais MS, Koncz C (2002) Knock-out of Arabidopsis metal transporter gene IRT1 results in iron deficiency accompanied by cell differentiation defects. Plant Mol Biol **50:** 587–597

Jeong J, Guerinot ML (2009) Homing in on iron homeostasis in plants. Trends Plant Sci **14:** 280-285

Kagale S, Links MG, Rozwadowski K (2010) Genome-wide analysis of Ethylene responsive element binding factor-associated Amphiphilic Repression motif-containing transcriptional regulators in Arabidopsis. Plant Physiol **152:** 1109-1134

Kim SA, LaCroix IS, Gerber SA, Guerinot ML (2019) The iron deficiency response in Arabidopsis thaliana requires the phosphorylated transcription factor URI. Proc Natl Acad Sci USA **116:** 24933-24942

Kosugi S, Hasebe M, Tomita M, Yanagawa H (2009) Systematic identification of yeast cell cycle-dependent nucleocytoplasmic shuttling proteins by prediction of composite motifs. Proc Natl Acad Sci USA **106:** 10171-10176

Krogan NT, Long JA (2009) Why so repressed? Turning off transcription during plant growth and development. Curr Opin Plant Biol **12:** 628-636

Kobayashi T, Nishizawa NK (2012) Iron uptake, translocation, and regulation in higher plants. **Annu Rev Plant Biol** 63: 131–152

Kobayashi T, Ozu A, Kobayashi S, An G, Jeon JS, Nishizawa NK (2019) OsbHLH058 and OsbHLH059 transcription factors positively regulate iron deficiency responses in rice. Plant Mol Biol **101:** 471-486

Lei R, Li Y, Cai Y, Li C, Pu M, Lu C, Yang Y, Liang G (2020) bHLH121 Functions as a Direct Link that Facilitates the Activation of FIT by bHLH IVc Transcription Factors for Maintaining Fe Homeostasis in Arabidopsis. Mol Plant **13:** 634-649

Li C, Liu X, Qiang X, Li X, Li X, Zhu S, Wang L, Wang Y, Liao H, Luan S, Yu F (2018) EBP1 nuclear accumulation negatively feeds back on FERONIA-mediated RALF1 signaling. PLoS Biol **16:** e2006340

Li X, Zhang H, Ai Q, Liang G, Yu D (2016) Two bHLH transcription factors, bHLH34 and bHLH104, regulate iron homeostasis in *Arabidopsis thaliana*. Plant Physiol **170:** 2478-2493

Li Y, Lu CK, Li CY, Lei RH, Pu MN, Zhao JH, Peng F, Ping HQ, Wang D, Liang G (2021) IRON MAN interacts with BRUTUS to maintain iron homeostasis in Arabidopsis. Proc Natl Acad Sci USA **118:**e2109063118

Liang G, Zhang HM, Lou DJ, Yu DQ (2016) Selection of highly efficient sgRNAs for CRISPR/Cas9-based plant genome editing. Sci Rep **6:** 21451

Liang G, Zhang HM, Li XL, Ai Q, Yu D (2017) bHLH transcription factor bHLH115 regulates iron homeostasis in Arabidopsis thaliana. J Exp Bot **68:** 1743-1755

Liang G, Zhang H, Li Y, Pu M, Yang Y, Li C, Lu C, Xu P, Yu D (2020) Oryza sativa Fer-like fe deficiency-induced transcription factor (OsFIT/OsbHLH156) interacts with OsIRO2 to regulate iron homeostasis. J Integr Plant Biol **62:** 668-689

Lingam S, Mohrbacher J, Brumbarova T, Potuschak T, Fink-Straube C, Blondet E, Genschik P, Bauer P (2011) Interaction between the bHLH transcription factor FIT and ETHYLENE INSENSITIVE3/ETHYLENE INSENSITIVE3-LIKE1 reveals molecular linkage between the regulation of iron acquisition and ethylene signaling in Arabidopsis. Plant Cell **23:**1815-1829

Liu L, Zhang Y, Tang S, Zhao Q, Zhang Z, Zhang H, Dong L, Guo H, Xie Q (2010) An efficient system to detect protein ubiquitination by agroinfiltration in

Nicotiana benthamiana. Plant J **61:** 893-903

Liu TY, Chou WC, Chen WY, Chu CY, Dai CY, Wu PY (2018) Detection of
membrane protein-protein interaction in planta based on dual-intein-coupled
tripartite split-GFP association. Plant J **94:** 426-438

Le CT, Brumbarova T, Ivanov R, Stoof C, Weber E, Mohrbacher J,
Fink-Straube C, Bauer P (2016) ZINC FINGER OF ARABIDOPSIS
THALIANA12 (ZAT12) Interacts with FER-LIKE IRON
DEFICIENCY-INDUCED TRANSCRIPTION FACTOR (FIT) Linking Iron
Deficiency and Oxidative Stress Responses. Plant Physiol. 170(1):540-557.

Long TA, Tsukagoshi H, Busch W, Lahner B, Salt DE, Benfey PN (2010)
The bHLH transcription factor POPEYE regulates response to iron
deficiency in Arabidopsis roots. Plant Cell **22:** 2219-2236

Ogo Y, Itai RN, Nakanishi H, Kobayashi T, Takahashi M, Mori S, Nishizawa
NK (2007) The rice bHLH protein OsIRO2 is an essential regulator of the
genes involved in Fe uptake under Fe-deficient conditions. Plant J **51:**
366-377

Pauwels L, Barbero GF, Geerinck J, Tilleman S, Grunewald W, Pérez AC,
Chico JM, Bossche RV, Sewell J, Gil E, et al. (2010) NINJA connects the
co-repressor TOPLESS to jasmonate signalling. Nature **464:**788-791

Quinet M, Vromman D, Clippe A, Bertin P, Lequeux H, Dufey I, Lutts S,
Lefèvre I (2012) Combined transcriptomic and physiological approaches
reveal strong differences between short- and long-term response of rice
(Oryza sativa) to iron toxicity. Plant Cell Environ **35:**1837–1859

Robinson NJ, Procter CM, Connolly EL, Guerinot ML (1999) A ferricchelate
reductase for iron uptake from soils. Nature **397:** 694–697

Rodríguez-Celma J, Lin W-D, Fu G-M, Abadía J, López-Millán, Schmidt W
(2013) Mutually exclusive alterations in secondary metabolism are critical for
the uptake of insoluble iron compounds by Arabidopsis and Medicago
truncatula. Plant Physiol **162:** 1473-1485

Rodríguez-Celma J, Connorton JM, Kruse I, Green RT, Franceschetti M,
Chen YT, Cui Y, Ling HQ, Yeh KC, Balk J (2019) Arabidopsis
BRUTUS-LIKE E3 ligases negatively regulate iron uptake by targeting
transcription factor FIT for recycling. Proc Natl Acad Sci USA
116:17584-17591

Samira R, Li B, Kliebenstein D, Li C, Davis E, Gillikin JW, Long TA (2018)
The bHLH transcription factor ILR3 modulates multiple stress responses in
Arabidopsis. Plant Mol Biol **97:** 297-309

Santi S, Schmidt W (2009) Dissecting iron deficiency-induced proton
extrusion in Arabidopsis roots. New Phytol **183:** 1072–1084

Selote D, Samira R, Matthiadis A, Gillikin JW, Long TA (2015) Iron-binding
E3 ligase mediates iron response in plants by targeting basic helix-loop-helix
transcription factors. Plant Physiol **167:** 273-286

Schmid NB, Giehl RF, Döll S, Mock HP, Strehmel N, Scheel D, Kong X,
Hider RC, von Wirén (2014) Feruloyl-CoA 69-Hydroxylase1-dependent

coumarins mediate iron acquisition from alkaline substrates in Arabidopsis.
Plant Physiol **164:** 160-172

Shahbazian MD, Grunstein M (2007) Functions of site-specific histone
acetylation and deacetylation. Annu Rev Biochem **76:**75-100

Siwinska J, Siatkowska K, Olry A, Grosjean J, Hehn A, Bourgaud F,
Meharg AA, Carey M, Lojkowska E, Ihnatowicz A (2018) Scopoletin
8-hydroxylase: a novel enzyme involved in coumarin biosynthesis and
iron-deficiency responses in Arabidopsis. J Exp Bot **69:**1735-1748

Song S, Qi T, Fan M, Zhang X, Gao H, Huang H, Wu D, Guo H, Xie D (2013)
The bHLH subgroup IIId factors negatively regulate jasmonate-mediated
plant defense and development. PLoS Genet **9:** e1003653

Szemenyei H, Hannon M, Long JA (2008) TOPLESS mediates auxin
dependent transcriptional repression during Arabidopsis embryogensis.
Science **319:** 1384-138

Kobayashi T, Nishizawa NK (2014) Iron sensors and signals in response to
iron deficiency. Plant Sci **224:**36-43

Tanabe N, Noshi M, Mori D, Nozawa K, Tamoi M, Shigeoka S (2019) The
basic helix-loop-helix transcription factor, bHLH11 functions in the
iron-uptake system in Arabidopsis thaliana. J Plant Res **132:** 93-105

Tissot N, Robe K, Gao F, Grant-Grant S, Boucherez J, Bellegarde F,
Maghiaoui A, Marcelin R (2019) Transcriptional integration of the
responses to iron availability in Arabidopsis by the bHLH factor ILR3. New
Phytol **223:**1433-1446

Trofimov K, Ivanov R, Eutebach M, Acaroglu B, Mohr I, Bauer P,
Brumbarova T (2019) Mobility and localization of the iron
deficiency-induced transcription factor bHLH039 change in the presence of
FIT. Plant Direct **3:**e00190

Tsai HH, Rodríguez-Celma J, Lan P, Wu YC, Vélez-Bermúdez IC, Schmidt
W (2018) Scopoletin 8-Hydroxylase-Mediated Fraxetin Production Is Crucial
for Iron Mobilization. Plant Physiol **177:**194-207

Wang HY, Klatte M, Jakoby M, Bäumlein H, Weisshaar B, Bauer P (2007)
Iron deficiency-mediated stress regulation of four subgroup Ib BHLH genes
in Arabidopsis thaliana. Planta 226: 897-908

Wang N, Cui Y, Liu Y, Fan H, Du J, Huang Z, Yuan Y, Wu H, Ling HQ (2013)
Requirement and functional redundancy of Ib subgroup bHLH proteins for
iron deficiency responses and uptake in Arabidopsis thaliana. Mol Plant **6:**
503-513

Wang S, Li L, Ying Y, Wang J, Shao JF, Yamaji N, Whelan J, Ma JF, Shou H
(2020) A transcription factor OsbHLH156 regulates Strategy II iron acquisition
through localising IRO2 to the nucleus in rice. New Phytol **225:**1247-1260

Wu FH, Shen SC, Lee LY, Lee SH, Chan MT, Lin CS (2009) Tape-Arabidopsis
Sandwich-a simpler Arabidopsis protoplast isolation method. Plant Methods
5:16

Yuan Y, Wu H, Wang N, Li J, Zhao W, Du J, Wang D, Ling HQ (2008) FIT

interacts with AtbHLH38 and AtbHLH39 in regulating iron uptake gene expression for iron homeostasis in Arabidopsis. Cell Res **18:** 385–397

Zhang H, Li Y, Yao X, Liang G, Yu D (2017) POSITIVE REGULATOR OF IRON HOMEOSTASIS1, OsPRI1, facilitates iron homeostasis. Plant Physiol **175:** 543-554

Zhang J, Liu B, Li M, Feng D, Jin H, Wang P, Liu J, Xiong F, Wang J, Wang HB (2015) The bHLH Transcription Factor bHLH104 Interacts with IAA-LEUCINE RESISTANT3 and Modulates Iron Homeostasis in Arabidopsis. Plant Cell **27:** 787-805

Zheng L, Ying Y, Wang L, Wang F, Whelan J, Shou H (2010) Identification of a novel iron regulated basic helix-loop-helix protein involved in Fe homeostasis in Oryza sativa. BMC Plant Biol **10:**166

Zuo J, Niu QW, Chua NH (2001) Technical advance: An estrogen receptor-based transactivator XVE mediates highly inducible gene expression in transgenic plants. Plant J **24:** 265-273

Figure Legends

Figure 1. *bhlh11* mutants are sensitive to Fe excess.

(A) Phenotypes of *bhlh11* mutants. Two-week-old seedlings grown on Fe0, Fe100 or Fe300 medium.

(B) Shoot biomass of *bhlh11* mutants. Fresh weight of two-week-old shoots grown on Fe0, Fe100 or Fe300 medium. Three biological duplicates, each of which contains 15 plants, were analyzed.

(C) Fe concentration of rosette leaves of three-week-old wild type and *bhlh11* plants grown in soil.

(D) Expression of *IRT1*, *FRO2* and bHLH Ib genes. Four-day-old plants grown on Fe100 medium were transferred to Fe0, Fe100 or Fe300 medium for three days, and root samples were harvested and used for RNA extraction and RT-qPCR.

(B-D)The different letters above each bar indicate statistically significant differences as determined by one-way ANOVA followed by Tukey's multiple comparison test (P < 0.05).

Figure 2. Response of bHLH11 to Fe status

(A) RT-qPCR analysis of *bHLH11* expression. Four-day-old plants grown on Fe100 medium were shifted to Fe0, Fe50, Fe100 or F300 medium for 3 days. Roots were used for RNA extraction and RT-qPCR. The different letters above each bar indicate statistically significant differences as determined by one-way ANOVA followed by Tukey's multiple comparison test (P < 0.05).

(B) Degradation of bHLH11 in response to Fe deficiency. Seven-day-old wild type and *bHLH11-OX-20* seedlings grown on Fe100 medium were transferred to Fe0 or Fe300 medium, and root samples were harvested after 1, 2, and 3 days. anti-HA was used to detect HA-bHLH11. β-tubulin was used as a loading control.

(C) Subcellular localization of bHLH11. The free mCherry, bHLH11-mCherry or bHLH11-NLS-mCherry were respectively expressed in *N. benthamiana* leaves.

Figure 3. bHLH11 physically interacts with bHLH IVc TFs.

(A) Yeast two-hybrid analysis of the interaction between bHLH11 and bHLH IVc TFs. Yeast cotransformed with different BD and AD plasmid combinations was spotted on synthetic dropout medium lacking Leu/Trp (SD–W/L) or Trp/Leu/His/Ade (SD –W/L/H/A).

(B) Interaction of bHLH11 and bHLH IVc TFs in plant cells. Tripartite split-sfGFP complementation assays were performed. bHLH34, bHLH104, bHLH105, bHLH115, and bHLH121 were fused with GFP10, and bHLH11 was fused with GFP11. The constructs were introduced into agrobacterium respectively, and the indicated combinations were co-expressed in *N. benthamiana* leaves.

(C) Co-IP analysis of the interaction between bHLH11 and bHLH IVc TFs. Total proteins from different combinations of HA-bHLH11 and MYC-GFP,

MYC-bHLH34, MYC-bHLH104, MYC-bHLH105, or MYC-bHLH115 were immunoprecipitated with anti-MYC followed by immunoblotting with the indicated antibodies. MYC-GFP was used as a negative control. Protein molecular weight (in kD) is indicated to the left of the immunoblot.

Figure 4 Change of bHLH11 subcellular localization.
(A) Location of bHLH11 in the absence or presence of bHLH IVc. bHLH11-mCherry was co-expressed with bHLH IVc TFs. The combination of bHLH11-GFP and free mCherry was used as a negative control. Transient expression assays were performed in tobacco leaves.
(B) Immunoblot analysis of bHLH11 protein distribution in the cytoplasm and nuclear fractions. Seven-day-old *bHLH11-OX-20* seedlings grown on Fe100 medium were transferred to Fe0, Fe100 or Fe300 medium. Root samples were harvested after 3 days, and cytoplasmic and nuclear proteins were extracted and subjected to immunoblot analysis with the indicated antibodies. Ratio indicates the relative protein abundance of HA-bHLH11.

Figure 5. bHLH11 antagonizes the transcriptional activation ability of bHLH IVc TFs.
(A) bHLH11 represses the functions of bHLH IVc TFs. Schematic diagram of the constructs transiently expressed in Arabidopsis protoplasts. The LUC/REN ratio represents the LUC activity relative to the internal control REN. The asterisk indicates a significant difference as determined by Student's t Test.
(B) bHLH11 inhibits the functions of bHLH IVc TFs by direct protein-protein interaction. The schematic diagram shows the constructs used in the transient expression assays in tobacco leaves. The abundance of *GFP* was normalized to that of *NPTII*. The asterisk indicates a significant difference as determined by Student's t Test.
(C) Expression of bHLH Ib genes in *pER8-bHLH11* plants. Seven-day-old plants grown on Fe0 medium were transferred to Fe0 medium with or without 4 µM estradiol for 6 h, and root samples were harvested and used for RNA extraction and RT-qPCR. The different letters above each bar indicate statistically significant differences as determined by one-way ANOVA followed by Tukey's multiple comparison test (P < 0.05).

Figure 6. bHLH11 acts as a repressor dependently on its two EAR motifs.
(A) Schematic diagram of the various mutated versions of bHLH11. The mutated amino acid is indicated in red. bHLH11m1, the first EAR mutated. bHLH11m2, the second EAR mutated. bHLH11dm, both double EARs mutated. bHLH11dm-VP16, bHLH11dm fused with the VP16 domain.
(B) The schematic diagram shows the constructs used in the transient expression assays in (C).
(C) The EAR motifs are required for the repression of bHLH11. Arabidopsis protoplasts were used for transient expression assays. The abundance of *GFP*

was normalized to that of *NPTII*. The different letters above each bar indicate statistically significant differences as determined by one-way ANOVA followed by Tukey's multiple comparison test (P < 0.05).

(D) Phenotypes of *bHLH11dm-VP16* and *bHLH11-OX* plants. Seven-day-old seedlings grown on Fe0 or Fe100 medium are shown.

Figure 7. Enhanced Fe deficiency response in *tpr1 tpr4 tpl*.

(A) The EAR motifs are required for the interaction between bHLH11 and TPL/TPRs. Yeast cotransformed with different BD and AD plasmid combinations was spotted on synthetic dropout medium lacking Leu/Trp (SD –T/L) or Trp/Leu/His/Ade (SD –T/L/H/A).

(B) Phenotypes of *tpr1 tpr4 tpl*. Two-week-old seedlings grown on Fe0, Fe100 or Fe300 medium are shown.

(C) Shoot biomass of *tpr1 tpr4 tpl*. Fresh weight of two-week-old shoots grown on Fe0, Fe100 or Fe300 medium. Three biological duplicates, each of which contains 15 plants, were analyzed.

(D) Fe concentration of rosette leaves of 3-week-old wild type and *tpr1 tpr4 tpl* plants grown in soil.

(E) Expression of *IRT1*, *FRO2* and bHLH Ib genes. Four-day-old plants grown on Fe100 medium were transferred to Fe0, Fe100 or Fe300 medium for three days, and root samples were harvested and used for RNA extraction and RT-qPCR.

(C-D) The asterisks indicate that the values are significantly different from the corresponding wild type value by Student's t Test (P < 0.05).

Figure 8. Genetic interaction between *bHLH11* and *FIT*.

(A) Phenotypes of *bHLH11-OX-20/fit-2* plants. Seven-day-old seedlings grown on Fe0 or Fe100 medium are shown.

(B) Expression of *IRT1*, *FRO2* and bHLH Ib genes. Four-day-old plants grown on Fe100 medium were transferred to Fe0 or Fe100 medium for three days, and root samples were harvested and used for RT-qPCR. The different letters above each bar indicate statistically significant differences as determined by one-way ANOVA followed by Tukey's multiple comparison test (P < 0.05).

Figure 9. A working model of bHLH11 in Fe homeostasis.

bHLH11 functions as an active repressor by recruiting TPL/TPR corepressors, and bHLH IVc TFs promote the nuclear accumulation of bHLH11. Under Fe-sufficient conditions, *bHLH11* message is activated and its protein accumulates. bHLH11 inhibits the transactivity of bHLH IVc TFs to bHLH Ib genes. The repression function of bHLH11 allows plants to avoid Fe toxicity. Under Fe-deficient conditions, unknown proteins repress the transcription of *bHLH11* and its protein decreases, which alleviates the bHLH11-mediated repression to bHLH IVc TFs. bHLH IVc TFs promote the transcription of bHLH Ib genes, and then bHLH Ib proteins interact with FIT to activate the expression of the Fe uptake genes *IRT1* and *FRO2*. The question signs

indicate unknown proteins.

A

	mCherry	GFP	Bright	Merged
bHLH11-mCherry GFP				
bHLH11-mCherry bHLH34-GFP				
bHLH11-mCherry bHLH104-GFP				
bHLH11-mCherry bHLH105-GFP				
bHLH11-mCherry bHLH115-GFP				

B

	Cytoplasm			Nucleus		
	Fe0	Fe100	Fe300	Fe0	Fe100	Fe300
Ratio	1.00	1.57	1.72	0.13	0.20	0.24
anti-HA						
anti-HSP						
anti-H3						