Generation of a novel SARS-CoV-2 sub-genomic RNA due to the R203K/G204R variant

in nucleocapsid: homologous recombination has potential to change SARS-CoV-2 at

both protein and RNA level

Shay Leary[1,¶], Silvana Gaudieri[1,2,3,¶], Matthew D. Parker[4,¶], Abha Chopra[1], Ian James[1], Suman

Pakala[3], Eric Alves[2], Mina John[1,5], Benjamin B. Lindsey[6,7], Alexander J Keeley[6,7], Sarah L.

Rowland-Jones[6,7], Maurice S. Swanson[8], David A. Ostrov[9], Jodi L. Bubenik[8], Suman Das[3],

John Sidney[10], Alessandro Sette[10,11], COVID-19 Genomics UK (COG-UK) consortium,

Thushan I. de Silva[6,7,*], Elizabeth Phillips[1,3,*], Simon Mallal[1,3,#*]

[1]Institute for Immunology and Infectious Diseases, Murdoch University, Murdoch, Western

Australia, Australia.

[2]School of Human Sciences, University of Western Australia, Crawley, Western Australia,

Australia.

[3]Division of Infectious Diseases, Department of Medicine, Vanderbilt University Medical

Center, Nashville, Tennessee, USA.

[4]Sheffield Biomedical Research Centre, Sheffield Bioinformatics Core, The University of

Sheffield, Sheffield, UK.

[5]Department of Clinical Immunology, Royal Perth Hospital, Perth, Western Australia,

Australia.

[6]Sheffield Teaching Hospitals NHS Foundation Trust, Sheffield, UK.

[7]Department of Infection, Immunity and Cardiovascular Disease and The Florey Institute for

Host-Pathogen Interactions, Medical School, University of Sheffield, Sheffield, UK.

[8]Department of Molecular Genetics and Microbiology, Center for NeuroGenetics and the

Genetics Institute, University of Florida, Gainesville, Florida, USA.

9Department of Pathology, Immunology and Laboratory Medicine, University of Florida, Gainesville, Florida, USA.

10Center for Infectious Disease and Vaccine Research, La Jolla Institute for Immunology, La Jolla, California, USA.

11Department of Medicine, Division of Infectious Diseases and Global Public Health, University of California, San Diego, La Jolla, California, USA.

¶These authors contributed equally to this work.

*These authors also contributed equally to this work.

#Corresponding author

Prof. Simon Mallal

Email: s.mallal@vumc.org

Keywords: COVID-19; SARS-CoV-2; homologous recombination; sub-genomic RNA transcript; transcription-regulating sequence; viral polymorphism

Abstract

Background: Genetic variations across the SARS-CoV-2 genome may influence transmissibility of the virus and the host's anti-viral immune response, in turn affecting the frequency of variants over-time. In this study, we examined the adjacent amino acid polymorphisms in the nucleocapsid (R203K/G204R) of SARS-CoV-2 that arose on the background of the spike D614G change and describe how strains harboring these changes became dominant circulating strains globally. **Methods:** Deep sequencing data of SARS-CoV-2 from public databases and from clinical samples were analyzed to identify and map genetic variants and sub-genomic RNA transcripts across the genome. **Results:** Sequence analysis suggests that the three adjacent nucleotide changes that result in the K203/R204 variant have arisen by homologous recombination from the core sequence (CS) of the leader transcription-regulating sequence (TRS) rather than by stepwise mutation. The resulting sequence changes generate a novel sub-genomic RNA transcript for the C-terminal dimerization domain of nucleocapsid. Deep sequencing data from 981 clinical samples confirmed the presence of the novel TRS-CS-dimerization domain RNA in individuals with the K203/R204 variant. Quantification of sub-genomic RNA indicates that viruses with the K203/R204 variant may also have increased expression of sub-genomic RNA from other open reading frames. **Conclusions:** The finding that homologous recombination from the TRS may have occurred since the introduction of SARS-CoV-2 in humans resulting in both coding changes and novel sub-genomic RNA transcripts suggests this as a mechanism for diversification and adaptation within its new host.

Introduction

It is believed SARS-CoV-2 originated from a bat coronavirus transmitted to humans, likely via an intermediate host such as a pangolin, acquiring a furin-cleavage site in the process. This new motif allows cleavage at the boundary of the S1 and S2 domains of the spike protein in virus-producing cells (1). A SARS-CoV-2 variant in the spike protein, D614G (B.1 lineage), emerged early in the epidemic and has rapidly became dominant in virtually all areas of the world where it has circulated (2). Several studies have shown this variant to be associated with higher viral RNA levels in the upper respiratory tract, higher titers in pseudoviruses in-vitro (2, 3) and increased infectivity (4, 5). More recently, emerging lineages from this genetic background (B.1.1.7 – 'Alpha or UK variant', B.1.351 – 'Beta or South African variant', or B.1.617.2 - 'Delta variant') have been identified with reported rapid local expansions of these viruses.

The diversification of coronaviruses can occur via point mutations and recombination events (6, 7) that can result in increased prevalence due to selective advantage related to increased infectiousness and transmission of the virus or by chance. Evidence of viral adaptation to selective pressures as a virus spreads among diverse human populations has important implications for the ongoing potential for changes in viral fitness over time, which in turn may impact transmissibility, disease pathogenesis and immunogenicity. Furthermore, the functional impact of new genetic changes need to be considered in the performance of diagnostic tests, ongoing public health measures to contain infection around the world and the development of universal vaccines and antiviral therapies including monoclonal antibodies.

Here we examined a variant of SARS-CoV-2 that emerged within the subset of sequences

harboring the D614G variant and contains three adjacent nucleotide changes spanning two

residues of the nucleocapsid protein (R203K/G204R; B.1.1 lineage) that has resulted in a

novel sub-genomic RNA transcript. Sequence analysis suggests these changes are the result

of homologous recombination from the core sequence (CS) of the leader transcription-

regulating sequence (TRS). This event introduced a new TRS between the RNA binding and

dimerization domains of nucleocapsid providing the template for the generation of a novel

sub-genomic RNA transcript. Further novel sub-genomic RNA transcripts arising in

association with incorporation of leader sequence and TRS were also observed, suggesting

homologous recombination from this region as a potential mechanism for SARS-CoV-2

diversification and adaptation within its new host.

Methods

Study Design

This study utilized deposited SARS-CoV-2 genomic sequences in public databases, with a further 981 Oxford Nanopore Technology genomes and clinical metadata from Sheffield, UK, as a validation set, to identify and map genetic variants and sub-genomic RNA transcripts across the genome. Accession numbers and links to datasets are in Supplementary Material.

SARS-CoV-2 sequence generation from patients with COVID-19

SARS-CoV-2 sequences, with matched clinical metadata, were generated using samples taken for routine clinical diagnostic use from 981 individuals presenting with COVID-19 disease to Sheffield Teaching Hospitals NHS Foundation Trust, Sheffield, UK. This work was performed under approval by the Public Health England Research Ethics and Governance Group for the COVID-19 Genomics UK consortium (R&D NR0195). Following extraction, samples were processed using the ARTIC Network SARS-CoV-2 protocol. After RT-PCR, SARS-CoV-2 specific PCR and library preparation with Oxford Nanopore LSK-109 and barcoding expansion packs NBD-104 and NBD-114 samples were sequenced on an Oxford Nanopore GridION X5 using R9.4.1D flow cells. Bases were called with either fast or high accuracy guppy with demultiplexing enabled and set to --require-both-ends. Samples were then analyzed using ARTIC Network pipeline v1.1.0rc1.

SARS-CoV-2 sequence acquisition from public repositories

Complete SARS-CoV-2 genome sequences were downloaded from the GISAID EpiCoV repository on 24th January 2021 (https://www.gisaid.org/). The complete dataset of 455,774 sequences with coverage across the genome were aligned in CLCbio Genomics Workbench

12 (QIAGEN Bioinformatics) to the GenBank reference sequence NC_045512.2. Aligned

sequences were exported in FASTA format and imported into Visual Genomics Analysis

Studio (VGAS), an in-house program for visualizing and analyzing sequencing data

(http://www.iiid.com.au/software/vgas). The chronological appearance of the sequences was

generated using the collection dates for each of the sequences. Of note, our current

knowledge of the global circulating variants is dependent on the ability of laboratories in

different countries to deposit full genome length SARS-CoV-2 sequences and may be subject

to ascertainment bias. As such, the frequencies of specific variants shown may not reflect the

size of the outbreak. However, the data does provide the opportunity to predict the presence

of specific variants in areas given the known epidemiology within different countries and

regions. A subset of subjects also had individual deep sequence reads deposited in the

Sequence Read Archive (SRA) at www.ncbi.nlm.nih/sra. These sequence reads were

downloaded and aligned as indicated above.

Identification of amino acid substitutions

Codon usage output allowed for identification of amino acid substitutions across the SARS-

Cov-2 genome. A cut-off of 5% frequency within the consensus SARS-CoV-2 protein

sequences was set to obtain the codon usage across all sequences and as shown in S1 Table.

The viral polymorphisms detected are present in viral variants sequenced using different

NGS platforms (e.g. nanopore, Illumina) and the Sanger-based sequencing method making it

unlikely that the new changes are sequence or alignment errors. In addition, different

laboratories around the world have deposited sequences with these polymorphisms in the

database and examination of individual sequences in the region failed to uncover obvious

insertions/deletions likely representing alignment issues or homopolymer slippage.

HLA peptide binding prediction

The region containing the adjacent amino acid polymorphisms in the nucleocapsid was divided into sliding windows of 8-14 amino acids. NetMHC 4.0 (http://www.cbs.dtu.dk/services/NetMHC/) and NetMHCpan 4.0 (http://www.cbs.dtu.dk/services/NetMHCpan/) with default settings were utilized to predict HLA-class I binding scores and binding differences across all HLA class-I alleles for the original 2003 SARS and current SARS-CoV-2 sequences harboring the R203/G204 and K203/R204 polymorphisms in the nucleocapsid (output listed in S2 Table).

HLA peptide binding assays

MHC was purified from the Steinlin EBV transformed homozygous cell line (IHWG ID: 9087; A*01:01, B*08:01 and C*07:01) using the B123.2 (anti-HLA-B, C) and W6/32 (anti-class I) monoclonal antibodies, and classical MHC-peptide inhibition of binding assays performed, as previously described (8). To develop an HLA C*07:01-specific binding assay, the IEDB was utilized to identify candidate peptides reported as HLA-C*07:01 epitopes or eluted ligands. One peptide (3424.0028; sequence IRSSYIRVL, Macaca mulatta and Homo sapiens DNA replication licensing factor MCM5 289-297) was radiolabeled and found in direct binding assays to yield a strong signal with as little as 0.5 nM MHC. Subsequent inhibition of binding assays established that 3424.0028 bound with an affinity of 0.21 nM. To establish that the putative assay was specific for C*07:01, and not co-purified B*08:01, two additional peptides previously reported as HLA-C*07:01 ligands were also tested, with one found to bind with high affinity (IC50 67 nM) and the other with intermediate (IC50 1600 nM). At the same time, a panel of known B*08:01 ligands were not found to have the capacity to inhibit binding of radiolabeled 3424.0028 (S3 Table). By contrast, when the same panel of peptides was tested in the previously validated B*08:01 assay (9), 3424.0028 was

found to bind with about 1500-fold lower affinity, all of the known B*08:01 ligands bound

with IC50s <10 nM, and the C*07:01 ligands with affinities >1000 nM.

Sub-genomic RNA classification & quantification in the Validation Dataset

We developed a tool, "periscope" (v0.0.0), to classify and quantify sub-genomic RNA in the

Sheffield ARTIC network Nanopore dataset (10). The tool can be downloaded from git-hub

at https://github.com/sheffield-bioinformatics-core/periscope. Briefly, this tool uses local

alignment to identify putative sub-genomic RNA supporting reads and uses genomic reads

from the same amplicon to normalize.

RNA structure modeling

The RNAfold program from the ViennaRNA Web Server (http://rna.tbi.univie.ac.at/) was

used for structural predictions using the default settings and the minimum free energy

structures were acquired using the base-pairing probability color scheme. The Dot-bracket

folding notations were obtained for each of the R203K/G204R sequences and used for

Junction   Explorer   (nature.njit.edu/biosoft/Junction-Explorer/)   and   CHS-align

(nature.njit.edu/biosoft/CHSalign/).

Statistical Analysis

Fisher exact test was used to compare the proportion of subjects with specific sub-genomic RNA transcripts. P values less than 0.05 was used as the statistical threshold. Comparisons between sub-genomic and genomic RNA expression in R203/G204 compared to K203/R204 containing sequences was made using the Mann-Whitney U test, corrected for multiple comparisons using the Holm method. Logistic and linear regression modeling used to explore the impact of K203/R204 and other co-variates on hospitalization, CT values and sub-genomic RNA expression.

Results and Discussion

Adjacent nucleocapsid polymorphisms emerged from the existing spike protein D614G variant

We utilized publicly available SARS-CoV-2 sequences from the GISAID database (available on the 24[th] of January 2021; www.gisaid.org) to identify amino acid polymorphisms arising in global circulating forms of the virus in relation to region and time of collection. Of the 455,774 circulating variants there were 29 amino acid polymorphisms present in >5% of the deposited sequences (of a total of 9413 sites; S1 Table) including the spike D614G variant (B.1 lineage) that emerged early in the pandemic and the adjacent R203K/G204R variants (B.1.1 lineage) in the nucleocapsid protein (11) that formed one of the main variants emerging from Europe in early 2020. As of the end of January 2021, the K203/R204 variant comprises 37.4% of globally reported SARS-CoV-2 sequences (Fig 1) and almost exclusively occurs on the D614G genetic background (S4 Table).

Although the D614G change rapidly increased in prevalence in almost all regions, the prevalence rates of the K203/R204 subset of the D614G variant are variable in different geographic areas and over-time (Fig 2). For example, an almost complete replacement of D614 by G614 was noted in South America between March and April 2020 and a similar replacement pattern was seen with the K203/R204 variant most marked in Chile, Argentina and Brazil (12). A closer examination of the deposited sequences in the UK shows the K203/R204 variant increasing in prevalence early in 2020 but the second wave later in the year shows a shift in the proportion of deposited sequences with the R203/G204 subset of the D614G variant (B.1.177 lineage) until the recent appearance of the B.1.1.7 'Alpha or UK variant' that harbors the K203/R204 polymorphisms (S1 Fig and S4 Table); supporting a likely increased infectivity of this variant.

Amino acid polymorphisms due to three adjacent nucleotide changes in the

nucleocapsid likely due to homologous recombination

Of the publicly available sequences examined with the two amino acid polymorphisms

K203/R204, all showed the three adjacent nucleotide changes from A<u>GG</u> <u>GG</u>A to A<u>AA</u>

<u>C</u>GA. There was no differential codon usage for the K203/R204 variant in the database.

However, there was evidence of low frequency alternative codon usage for arginine at 203

(AGA) for the R203/G204 variant and for lysine (AAG) at 203 for the K203/G204 variant

(S5 Table). Overall, circulating variants that contain the intermediate codon as the consensus

that could facilitate a single step from the AGG arginine codon to the AAA lysine codon at

position 203 appear rare among captured variants to date (S5 Table). Furthermore, a K203

polymorphism alone was seen in 0.3% and an R204 polymorphism alone seen in only 0.02%

of sequences (S5 Table). The low frequency K203/L204 and K203/P204 variants are both

one nucleotide step from the K203/R204 variant, have been deposited into the public

databases (November 2020) well after the emergence of the K203/R204 variant (February

2020) and accordingly likely arose from this genetic background.

The rapid emergence of closely linked polymorphisms in viruses can also reflect strong

selection pressure on this region of the genome in which the original mutation incurred a

replicative capacity, or other fitness cost, which could be restored by a linked compensatory

mutation. Evidence for such adaptations with closely linked compensatory mutations are

known to occur under host immune pressure as is well established for other RNA viruses

such as HIV (13-15) and Hepatitis C virus (16). In the absence of anti-viral treatment, these

viruses have such a high rate of viral replication, error-prone polymerases and lack associated

proofreading, mismatch repair, and other nucleic acid repair pathways generating a swarm of

viral variants with ongoing recombination between variants (in the case of HIV) being

generated continuously. As a result, selection pressure exerted by immune responses or other

selective pressures effectively operate on each separate residue independently (15). In

contrast, coronaviruses encode proofreading machinery and have a propensity to adapt by

homologous recombination between viruses (6) rather than necessarily by classic stepwise

individual mutations driven by selective pressures effectively operating on individual viral

residues. Furthermore, a simulation based on the nucleocapsid genomic region and allowing

up to 10 random mutations indicates the likelihood of observing three consecutive nucleotide

changes is less than 0.0005. These findings argue against stepwise change of the nucleotides

for the R203K/G204R variant.

The introduction of the AAACGA motif by homologous or heterologous recombination is a

more parsimonious mechanistic explanation and would have immediately resulted in both an

R to K change and adjacent G to R change at the positions 203 and 204, respectively. It is

critical to determine if the introduction of the AAACGA motif has induced any replicative or

other fitness change for the virus as a result of either structural or functional changes in the

RNA or the concomitant change of amino acids from R203/G204 to K203/R204 and any

related structural or functional impact on the nucleocapsid protein.

SARS-CoV-2 itself as likely source for homologous recombination

To identify possible viral sources for homologous recombination with SARS-CoV-2,

we initially performed a search of the motif in the nucleocapsid in related beta coronaviruses

from human and other species in the public databases and only found the presence of the

R203/G204 combination. We performed a similar search in our metatranscriptome data

generated from a cohort study consisting of 65 subjects of whom 43 had acute respiratory

infections and 22 were asymptomatic. From the data we assembled near complete and coding

complete viral genomes of the Coronavirus (NL63 - alpha, OC43 - beta, 229E - alpha), RSV

(A, B), Rhinovirus (A, B, C), Influenza (A - H3N2), and Bocavirus family. None of the alpha

coronaviruses had the R203/G204 or K203/R204 combination or indeed any variation at

these sites (n=14; sequence depth >3000). We then performed a search for stretches of

similarity using varying window sizes (>14 base-pair (bp) including the motif) in all

sequences. A 14bp window was selected as 14bp has been shown to be the minimum amount

of homology required for homologous recombination in mammalian cells (17). No significant

hits were identified. However, the AAACGA sequence encoding the K203/R204 amino acids

overlaps with the CTAA<u>ACGAAC</u> motif of the leader transcription-regulating sequences

(TRS; core underlined) (18) of SARS-CoV-2 itself and this core sequence motif is also found

near the start codon of the protein for surface glycoprotein (S), ORF3a, E, M, ORF6, ORF7a,

ORF8, ORF10 and nucleocapsid, in keeping with its known roles in mediating template

switching and discontinuous transcription (18).

Deep sequencing confirms quasi-species with the leader sequence linked to known or

introduced TRS region

Discontinuous transcription of SARS-CoV-2 results in sub-genomic RNA (sgRNA)

transcripts containing 5'-leader sequence-TRS-start codon-ORF-3'. These RNA transcripts

should also be captured from reads generated from NGS platforms. We therefore reasoned we

should be able to find such sequences within deep sequencing reads at the sites of known

sub-genomic regions (corresponding to the ORFs) and adjacent to position 203/204 of the

nucleocapsid in subjects infected with the K203/R204 variant but not in those with the

R203/G204 variant (Fig 3).

We searched for sgRNAs in sequence data generated from n=981 patients with COVID-19

based on the ARTIC network protocol (www.artic.network/ncov-2019; Fig 3) and subsequent

Nanopore sequencing in Sheffield, UK. As expected, the most frequent sgRNA transcripts in

each subject, irrespective of variant, corresponded to the known regions containing the start

codon of the SARS-CoV-2 proteins (Fig 4A). However, out of a total of 550 K203/R204

sequences, 231 had evidence (>=1 read containing leader sequence at the novel TRS site) of

the non-canonical nucleocapsid sgRNA (42%) but only 1 out of a total of 431 R203/G204

subjects had evidence of the novel sgRNA (likely a false positive as described in S2 Fig).

We confirmed the presence of the novel non-canonical nucleocapsid sgRNA in 27/45

individuals with the K203/R204 variant but in none of 45 individuals with the R203/G204

variant (Fisher test, p=5.0e-11; S6 Table) from the sequence read archive (SRA) database

(www.ncbi.nlm.nih/sra). Interestingly, we also found the presence of 23 other non-canonical

sgRNA transcripts with the 5'-leader-TRS-start codon-3' at low frequency in the 90 subjects

(irrespective of variant) due to multiple adjacent changes to the consensus sequence across

the genome generating new core TRS motifs (including with minor mismatches) (S6 Table).

It should be noted that none of these changes are present in the consensus sequence of the

SARS-CoV-2 genomes downloaded and represent low frequency quasispecies within

individuals. It does, however, suggest other instances of the introduction of the core

sequences from the leader TRS elsewhere in the SARS-CoV-2 genome.

SARS-CoV-2 viruses with K203/R204 are not associated with greater hospitalization

with COVID-19 or higher virus levels in the upper respiratory tract

The same dataset from COVID-19 patients in Sheffield, UK, was used to explore whether the

K203/R204 variant had any association with clinical outcome. The median age of this cohort

was 54 years (IQR 38 to 74) and 59.8% were female. Of these, 440 (44.9%) were

hospitalized COVID-19 patients and 42 (4.3%) subsequently required critical care support. A

multivariable logistic regression model including 203/204 status, age and sex showed no

association of K203/R204 with hospitalization (OR 0.82, 95% confidence intervals (CI 0.58 –

1.16), p=0.259). As expected, higher age and male sex were significantly association with

hospitalization with COVID-19 (OR 1.09, 95% CI 1.08 – 1.11, p <2e-16 for age and OR

4.47, 95% CI 3.13 – 6.43, p=2.91e-16 for male sex). Male sex, but not age or 203/204 status,

was associated with risk of critical care admission (S7 Table).

We explored whether K203/R204 was associated with greater virus levels in the upper

respiratory tract as estimated by cycle threshold (CT) values from the diagnostic RT-PCR. As

day of illness will impact CT value, we focused on a subset of the cohort (n=478) where this

information was available (all non-hospitalized patients, median symptom day 3, range 1 – 13

days). Data were analyzed with sequences stratified by spike 614 and nucleocapsid 203/204

status (D614/R203/G204, G614/R203/G204 and G614/K203/R204). Multivariable linear

regression models showed no impact of G614/K203/R204 compared to G614/R203/G204

status on CT values (p= 0.83, S6B Table), but as expected, later day of symptom onset was

significantly associated with higher CT values, therefore lower viral load (S8 Table,

p=2.05E-05). Consistent with recent findings (2), presence of a spike D614G variant was

significantly associated with lower CT values (higher viral loads) in the same subset of

individuals, even when day of illness at sampling is included in the model (S8A Table,

D614/R203/G204 vs G614/R203/G204, p=0.00011, Fig 5A & B).

SARS-CoV-2 viruses with K203/R204 have evidence of higher sub-genomic RNA

expression

We hypothesized that the amount of sgRNA at each of the ORF TRS positions in the SARS-

CoV-2 genome in ARTIC nanopore sequencing data could serve as a proxy for expression

levels of each of the ORFs due to their positions in the amplicons (Fig 3). To test this

hypothesis we developed a tool, periscope (19), which quantifies the number of sgRNA and

genomic RNA reads at each ORF TRS position in ARTIC network nanopore sequencing

data. We applied periscope to the 981 sequences in the Sheffield validation dataset. To

control for the sequencing depth differences evident between amplicons, we determined the

amplicon that shares the 3' primer with the sgRNA reads and used the total count of genomic

RNA at this amplicon to calculate the proportion of sgRNA for each ORF. The N ORF

sgRNA is expressed at high levels in all samples. ORF10 sgRNA was absent as others have

shown (20). A significant increase in sgRNA levels for several ORFs in samples with

K203/R204 compared to R203/G204 samples is apparent (Fig 4B). N is the most striking

example (Fig 4C, Mann-Whitney U test p value, adjusted for multiple testing p = 2.06e-37),

but sgRNA from ORFs E, M and ORF6 are also significantly increased. There is no

significant difference in genomic RNA levels (Fig 4D, normalized to total mapped reads)

between these two groups.

As discussed above, the K203/R204 variants appear to have emerged within the subset of

SARS-CoV-2 sequences with a D614G variant in the spike protein, which has been

associated with infections with a higher viral load in the upper respiratory tract. To explore

whether the differences between K203/R204 and R203/G204 sequences in sgRNA quantities

were due to D614 compared to G614 variant differences, we repeated the comparisons

following further stratification of sequences. Interestingly, G614/R203/G204 variants showed

lower total sgRNA expression than D614/R203/G204 samples (S3 Fig). Of note, sgRNA for

spike (S), membrane (M) and envelope (E) ORFs were significantly higher in samples with

D614/R203/G204 compared to those with G614/R203/G204 (adjusted p values 1.02e-4 for S,

0.0495 for M and 0.00696 for E). Total sgRNA in G614/K203/R204-containing samples was

still significantly higher than in G614/R203/G204 samples (S3A Fig, Mann-Whitney U test p

value, adjusted for multiple testing p = 3.5e-6). Similar increases in some individual ORF

sgRNA quantities in G614/K203/R204 compared to G614/R203/G204 sequences were also

seen, most notably for nucleocapsid (S3B Fig, adjusted p value 1.34e-12).

To ensure that the increase in sgRNA in K203/R204-containing sequences was not due to

confounding by differences in sampling date compared to date of symptom onset, we

evaluated the impact of K203/R204 and day of illness on sgRNA expression in a

multivariable linear regression model using the subset of 478 sequences described above

(stratified by D614/R203/G204, G614/R203/G204 and G614/K203/R204 status). Higher

sgRNA levels were significantly associated with later day from symptom onset (S9 Table,

p=9.9E-08). G614/R203/G204 compared to D614/R203/G204 was again associated with a

reduction in sgRNA levels (p=0.011, S9A Table), whereas a K203/R204 change on the

background of spike G614-containing sequences was associated with a significant increase in

sub-genomic RNA (p=4.51E-05, S9B Table). Spike canonical sub-genomic RNA was higher

in D614/R203/G204 samples, whereas nucleocapsid canonical sub-genomic RNA was higher

in G614/K203/R204 samples (Fig 5C and D, S3 Fig).

RT-PCR assays have been developed to directly assess sub-genomic mRNA (sgRNA) as a

measure of replicative intermediates of SARS-CoV-2 representing putative replication in

cells rather than RNA packaged in virions or residual viral RNA (21, 22). A decline in

sgRNA in sputum typically occurs from day 10 to 11 after onset of symptoms (22). Our

finding that a variant can emerge that is associated with a novel sub-genomic RNA or may

differentially impact the level of different sgRNAs suggest that the viral sequences should be

analyzed to ensure the primers or probes used are appropriate and analysis of short read deep

sequences with the periscope tool considered to help interpret results obtained from different

variants.

Potential impact of introduced TRS sequences on RNA structure

Modeling of the region around the mRNA encoding position 203 and 204 of the nucleocapsid

using RNAfold (23) predicts the presence of a three-way junction in the RNA (S4 Fig),

which was also predicted using Junction-Explorer (24). Three-way junction motifs are

common throughout biology and are found both in pure RNAs, such as riboswitches or

ribozymes, and in RNA-protein complexes, including the ribosome (25). RNA three-way

junctions are often stabilized via terminal loop interactions with distant tertiary contacts

while the junctions act like flexible hinges. These attributes allow these structures to sample

unusual conformational spaces and they often form platforms for interactions with other

molecules such as proteins, RNAs or small molecule ligands (25), and these folds often have

an essential role in either the function or assembly of the molecules in which they are

contained.

RNAfold predicts the mutation from A<u>GGGG</u>A to A<u>AAC</u>GA strongly disrupts this structure

as the lengths of the predicted helices and each of the junctions are altered and the stability of

Helix 2 is undermined (S4 Fig). A comparison of the two-modeled sequences using

CHSalign (26) also indicates that none of the junctions are maintained. Given these

widespread alterations, this modeling predicts that the A<u>GGG</u>GA to A<u>AAC</u>GA mutation

would have a strong impact on the local RNA structure of this region, and likely impacts the

normal function of this three-way junction motif. Interestingly, the RNA modeling shown in

S4 Fig also suggests that pairing of specific nucleotides to maintain these RNA structures

may require the preferential codon usage by RG (AGGGGA) and KR (AAACGA) and be a

contributory factor to preferential codon usage in RNA viruses more generally even in

protein coding regions.

While it is not possible to determine the impact of this proposed structural alteration on

SARS-CoV-2 without a defined function for this structure, there are precedents where minor

changes in a three-way junction have large functional consequences for their host viruses. For

example, Flaviviruses such as Dengue and West Nile virus utilize the host cell machinery to

degrade viral genomes until they encounter structures near the 3' end that are resistant to

XRN1 5'-3' exonuclease (27). The resulting small flaviviral RNAs (sfRNAs) are non-coding

RNAs that induce cytopathicity and pathogenicity. The resistance of sfRNA to XRN1 is

dependent on the structure of a three-way junction and a single nucleotide change at the

junction alters the fold sufficiently to prevent the accumulation of disease-related sfRNAs.

Thus, small changes at the nucleotide level can have profound functional consequences for

viral RNA three-way junctions.

Lack of evidence that the RG to KR change at positions 203 and 204 of nucleocapsid

was driven by HLA-restricted immune selective pressure

Selection of viral adaptations to polymorphic host responses mediated by T cells, NK-cells

and antibodies are well described for other RNA viruses such as HIV and HCV (15, 28).

HIV-1 adaptations to human leucocyte antigen (HLA)-restricted T-cell responses have also

been shown to be transmitted and accumulate over time (29, 30). As previously shown for

SARS-CoV, T-cell responses against SARS-CoV-2 are likely to target the nucleocapsid (31).

Notably, SARS-CoV-2 R203K/G204R polymorphisms modify the predicted binding of

putative HLA-restricted T-cell epitopes containing these residues (S2 Table). One of the

predicted T-cell epitopes is restricted by the HLA-C*07 allele; and we therefore considered

whether escape from HLA-C-restricted T-cell responses may conceivably confer a fitness

advantage for SARS-CoV-2, particularly in European populations where HLA-C*07 is

prevalent and carried by >40% of the population (www.allelefrequencies.net). However,

using HLA-C*07:01 purified from the Steinlin cell line (IHWG ID: 9087; A*01:01, B*08:01

and C*07:01) and the anti-HLA Class I B123.2 mAb in inhibition assays we were not able to

detect binding of either of the SARS-CoV-2 peptides S<u>RG</u>TSPARM or S<u>KR</u>TSPARM (S3

Table). We therefore have, as yet no evidence of any impact or selective advantage to the

virus at the protein level of a change at position 203/204 from the RG to KR residues.

SARS-CoV-2 and Host Adaptation: Implications for global viral dynamics,

pathogenesis and immunogenicity

Currently the possible functional effect(s) of the introduction of the AAACGA motif from the

leader TRS into the RNA encoding position 203 and 204 of the nucleocapsid at the RNA and

protein level are not known. TRS sites are usually intergenic and it has been assumed that

recombination events at such sites are more likely to be viable. It has also been shown

recently that recombination breakpoint hotspots in coronaviruses are more frequently co-

located with TRS-B sites than expected (32). Our findings suggest that a novel TRS-B site

can be introduced in a recombination breakpoint from the leader TRS, and that this can occur

within an ORF and remain viable. The exact mechanism by which the AAA CGA codons

could have been incorporated from the TRS-L into the nucleocapsid is not known but may

have first required the AAACGA to be captured from the TRS-L and then for replication to

be reinitiated at the nucleocapsid to generate a full-length genomic RNA.

The nucleocapsid protein is a key structural protein critical to viral transcription and

assembly (33), suggesting that changes in this protein could either increase or decrease

replicative fitness. The K203/R204 polymorphism is located between the RNA

binding/serine-rich domains and the dimerization structural domain (S5 Fig) in a part of the

protein that has not been characterized in terms of 3-dimensional structure. The sequence of

this region is not similar enough to solved structures to allow prediction of the influence of

the K203/R204 polymorphisms on the structure or function of the protein. However, it is

known that SARS-CoV-2 is exquisitely sensitive to interferons and that it depends on the

nucleocapsid and M proteins to maintain interferon antagonism (34, 35). Specifically the C

terminus (aa 362 to 422) of the nucleocapsid, which is predicted to be expressed at higher

levels in those with the KR variant and novel sgRNA, has been shown to interact with the

SPRY domain of TRIM25 disturbing its interaction with CARDs of RIG-I inhibiting RIG-I

ubiquitination and Type 1 interferon signaling (36). Importantly the cells expressing the C-

terminal nucleocapsid protein in that study produced lower viral titer, suggesting the

incorporation of this protein into the nucleocapsid may reduce the formation of functional

virus. This raises the possibility that any enhancement of inhibition of interferon signaling

associated with the novel K203/R204 sgRNA may be offset by less efficient replication,

potentially accounting for the lack of association with higher viral load in the upper

respiratory tract and absence of epidemiologic evidence of increased transmission. It is also

possible that the increase in sgRNA directly inhibits RIG-I signaling and downstream Type I

interferon responses as has been described for Dengue serotype 2 (37). Finally, the central

region of coronavirus nucleocapsid (aa 117 to 268) has been shown to have RNA chaperone

activity that enhances template switching and efficient transcription possibly accounting for

the increase in sgRNA for the E and M proteins and ORF6 in KR-sequences compared to

RG-sequences (38). Note we cannot exclude that the novel sgRNA may also use the

downstream ATG in the ORF9c reading frame.

The adaptive potential of differential expression of sgRNAs is supported by a recent study by

Thorne and colleagues that demonstrates that the B.1.1.7 ('Alpha' or UK variant) isolate

containing the R203K/G204R substitutions is associated with enhanced antagonism of the

innate immune response (39). Specifically, this study showed that in-vitro infection of human

lung epithelial (Calu-3) cells by B.1.1.7 isolates showed diminished RNA and protein

expression of IFNβ and reduced induction of interferon sensitive genes relative to other

isolates without these defining mutations in the nucleocapsid (normalized for intracellular

viral RNA). This effect was independent of the reduced sensitivity to type I and III IFNs

described for isolates carrying the D614G spike mutation (40). Further evaluation of this

system showed that infection with the B.1.1.7 isolate resulted in significant changes in

protein expression of known innate immune regulators such as ORF9b (41), ORF6 (42) and

nucleocapsid (36, 43), as well as increased levels of the N* sgRNA described in this study

and was again confirmed to be unique to those isolates with the R203K/G204R mutations.

These increased levels of sgRNAs and protein support the findings in this study showing

increased sgRNA levels for N, ORF6 and N* in clinical samples from B.1.1.7-infected

subjects relative to subjects infected with other SARS-CoV-2 isolates. Interestingly, the

increased levels of ORF9b may be due to the D3L mutation in the nucleocapsid that we have

proposed to have arisen similarly to the R203G/G204R mutations and is associated with

increased levels of B.1.1.7 sgRNA encoding ORF9b in clinical samples (44).

The B.1.617.2 ('Delta') variant appears to be more transmissible even in the context of

previous vaccination and is now replacing other variants. This variant has acquired an

R203M substitution as a result of a single nucleotide change while retaining an arginine (G)

at position 204. This raises the possibility that the residue 203 is critical to the interaction of

nucleocapsid with TRIM25 decreasing the Type 1 interferon response or increases

transmissibility in some other way (36).

Other contemporary concerns include the fall in antibody levels following infection or

vaccination, the potential limited durability of protection afforded by currently available

vaccines and the risk of reinfection by variants after vaccination (45, 46). At low levels of

antibodies, the lungs appear to remain relatively protected against severe disease presumably

by some combination of antibodies and amnestic responses restimulated by the time the lung

is involved. In contrast, the early establishment of infection in the upper respiratory tract

appears possible if antibody levels are low (47). We therefore postulate that variants that are

more effective in interfering with Type 1 interferon responses would be more transmissible,

but not necessarily cause severe disease in the context of waning immunity at an individual or

population level.

Conclusion

Marked viral diversity and adaptation of other RNA viruses such as HIV, HCV and influenza

to host selective pressures have been a barrier to successful treatment and vaccination to date.

Although SARS-CoV-2 is less diverse and adaptable, the D614G variant and the K203/R204

and Delta variants have emerged by either nucleotide mutation or homologous recombination

during its rapid, widespread global spread and do appear to have functional impact. It will

therefore be critical to continue molecular surveillance of the virus and elucidate the

functional consequences of any newly emerging viral genetic changes to guide development

of diagnostics, antivirals and universal vaccines and to target conserved and potentially less

mutable SARS-CoV-2 elements. The ability of SARS-CoV-2 to introduce new TRS motifs

throughout its genome with the potential to introduce both novel sub-genomic RNA

transcripts and coding changes in its proteins may add to these challenges.

Acknowledgments: We thank colleagues at the Institute for Immunology and Infectious

Diseases, Murdoch University, Australia and the Department of Medicine, Division of

Infectious Diseases, Vanderbilt University Medical Center, USA. We would like to

acknowledge additional members of the Sheffield COVID-19 Genomic Group who

contributed to the generation of the sequence data: Adrienne Angyal, Rebecca L. Brown,

Laura Carrilero, Cariad M Evans, Luke R. Green, Danielle C. Groves, Katie J Johnson, Paul J

Parsons, David Partridge, Mohammad Raza, Rachel M. Tucker, Dennis Wang, Matthew D.

Wyles. Ethics approval and consent to participate (COG-UK CONSORTIUM; R&D

NR0195).

We thank the reviewers and editor for their helpful comments.

Funding Sources: SG, SL and EA were supported by a grant awarded by the National

Health and Medical Research Council (NHMRC; APP1148284). SM was supported by a

National Institutes of Health (NIH)-funded Tennessee Center for AIDS Research (P30

AI110527). MDP was funded by the NIHR Sheffield Biomedical Research Centre (BRC - IS-

BRC-1215-20017). Sequencing of SARS-CoV-2 samples was undertaken by the Sheffield

COVID-19 Genomics Group as part of the COG-UK CONSORTIUM. COG-UK and

supported by funding from the Medical Research Council (MRC) part of UK Research &

Innovation (UKRI), the National Institute of Health Research (NIHR) and Genome Research

Limited, operating as the Wellcome Sanger Institute. TIdS is supported by a Wellcome Trust

Intermediate Clinical Fellowship (110058/Z/15/Z).

Conflicts of interests: The authors declare that they have no conflicts of interests.

References:

1.   Zhang T, Wu Q, Zhang Z. Probable Pangolin Origin of SARS-CoV-2 Associated with the COVID-19 Outbreak. Curr Biol. 2020;30(7):1346-51 e2. Epub 2020/03/21. doi: 10.1016/j.cub.2020.03.022. PubMed PMID: 32197085; PMCID: PMC7156161.

2.   Korber B, Fischer WM, Gnanakaran S, Yoon H, Theiler J, Abfalterer W, Hengartner N, Giorgi EE, Bhattacharya T, Foley B, Hastie KM, Parker MD, Partridge DG, Evans CM, Freeman TM, de Silva TI, Sheffield C-GG, McDanal C, Perez LG, Tang H, Moon-Walker A, Whelan SP, LaBranche CC, Saphire EO, Montefiori DC. Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G Increases Infectivity of the COVID-19 Virus. Cell. 2020. Epub 2020/07/23. doi: 10.1016/j.cell.2020.06.043. PubMed PMID: 32697968; PMCID: PMC7332439.

3.   Grubaugh ND, Hanage WP, Rasmussen AL. Making Sense of Mutation: What D614G Means for the COVID-19 Pandemic Remains Unclear. Cell. 2020. Epub 2020/07/23. doi: 10.1016/j.cell.2020.06.040. PubMed PMID: 32697970; PMCID: PMC7332445.

4.   Yurkovetskiy L, Pascal KE, Tompkins-Tinch C, Nyalile T, Wang Y, Baum A, Diehl WE, Dauphin A, Carbone C, Veinotte K, Egri SB, Schaffner SF, Lemieux JE, Munro J, Sabeti PC, Kyratsous C, Shen K, Luban J. SARS-CoV-2 Spike protein variant D614G increases infectivity and retains sensitivity to antibodies that target the receptor binding domain. bioRxiv. 2020. Epub 2020/07/09. doi: 10.1101/2020.07.04.187757. PubMed PMID: 32637944; PMCID: PMC7337374.

5.   Zhang L, Jackson CB, Mou H, Ojha A, Rangarajan ES, Izard T, Farzan M, Choe H. The D614G mutation in the SARS-CoV-2 spike protein reduces S1 shedding and increases infectivity. bioRxiv. 2020. Epub 2020/06/27. doi: 10.1101/2020.06.12.148726. PubMed PMID: 32587973; PMCID: PMC7310631.

6.   Graham RL, Baric RS. Recombination, reservoirs, and the modular spike:

mechanisms of coronavirus cross-species transmission. Journal of virology. 2010;84(7):3134-

46. Epub 2009/11/13. doi: 10.1128/JVI.01394-09. PubMed PMID: 19906932; PMCID:

2838128.

7.   Ji W, Wang W, Zhao X, Zai J, Li X. Cross-species transmission of the newly

identified coronavirus 2019-nCoV. Journal of medical virology. 2020;92(4):433-40. Epub

2020/01/23. doi: 10.1002/jmv.25682. PubMed PMID: 31967321.

8.   Sidney J, Southwood S, Moore C, Oseroff C, Pinilla C, Grey HM, Sette A.

Measurement of MHC/peptide interactions by gel filtration or monoclonal antibody capture.

Curr Protoc Immunol. 2013;Chapter 18:Unit 18 3. Epub 2013/02/09. doi:

10.1002/0471142735.im1803s100. PubMed PMID: 23392640; PMCID: PMC3626435.

9.   Sidney J, del Guercio MF, Southwood S, Engelhard VH, Appella E, Rammensee HG,

Falk K, Rotzschke O, Takiguchi M, Kubo RT, et al. Several HLA alleles share overlapping

peptide specificities. Journal of immunology. 1995;154(1):247-59. Epub 1995/01/01.

PubMed PMID: 7527812.

10.   Parker MD, Lindsey BB, Leary S, Gaudieri S, Chopra A, Wyles M, Angyal A, Green

LR, Parsons P, Tucker RM, Brown R, Groves D, Johnson K, Carrilero L, Heffer J, Partridge

DG, Evans C, Raza M, Keeley AJ, Smith N, Filipe ADS, Shepherd JG, Davis C, Bennett S,

Sreenu VB, Kohl A, Aranday-Cortes E, Tong L, Nichols J, Thomson EC, Wang D, Mallal S,

De Silva TI. Subgenomic RNA identification in SARS-CoV-2 genomic sequencing data.

Genome Research. 2021;31(4):645-58. doi: 10.1101/gr.268110.120.

11.   Jenjaroenpun P, Wanchai V, Ono-Moore KD, Laudadio J, James LP, Adams SH,

Prior F, Nookaew I, Ussery DW, Wongsurawat T. Two SARS-CoV-2 Genome Sequences of

Isolates from Rural U.S. Patients Harboring the D614G Mutation, Obtained Using Nanopore

Sequencing. Microbiol Resour Announc. 2020;10(1). Epub 2020/12/19. doi: 10.1128/MRA.01109-20. PubMed PMID: 33334896.

12. Franco-Muñoz C, Álvarez-Díaz DA, Laiton-Donato K, Wiesner M, Escandón P, Usme-Ciro JA, Franco-Sierra ND, Flórez-Sánchez AC, Gómez-Rangel S, Rodríguez-Calderon LD, Barbosa-Ramirez J, Ospitia-Baez E, Walteros DM, Ospina-Martinez ML, Mercado-Reyes M. Substitutions in Spike and Nucleocapsid proteins of SARS-CoV-2 circulating in South America. Infect Genet Evol. 2020;85:104557-. Epub 2020/09/17. doi: 10.1016/j.meegid.2020.104557. PubMed PMID: 32950697.

13. Leslie A, Kavanagh D, Honeyborne I, Pfafferott K, Edwards C, Pillay T, Hilton L, Thobakgale C, Ramduth D, Draenert R, Le Gall S, Luzzi G, Edwards A, Brander C, Sewell AK, Moore S, Mullins J, Moore C, Mallal S, Bhardwaj N, Yusim K, Phillips R, Klenerman P, Korber B, Kiepiela P, Walker B, Goulder P. Transmission and accumulation of CTL escape variants drive negative associations between HIV polymorphisms and HLA. J Exp Med. 2005;201(6):891-902. Epub 2005/03/23. doi: 10.1084/jem.20041455. PubMed PMID: 15781581; PMCID: 2213090.

14. Leslie AJ, Pfafferott KJ, Chetty P, Draenert R, Addo MM, Feeney M, Tang Y, Holmes EC, Allen T, Prado JG, Altfeld M, Brander C, Dixon C, Ramduth D, Jeena P, Thomas SA, St John A, Roach TA, Kupfer B, Luzzi G, Edwards A, Taylor G, Lyall H, Tudor-Williams G, Novelli V, Martinez-Picado J, Kiepiela P, Walker BD, Goulder PJ. HIV evolution: CTL escape mutation and reversion after transmission. Nature medicine. 2004;10(3):282-9. Epub 2004/02/11. doi: 10.1038/nm992. PubMed PMID: 14770175.

15. Moore CB, John M, James IR, Christiansen FT, Witt CS, Mallal SA. Evidence of HIV-1 adaptation to HLA-restricted immune responses at a population level. Science. 2002;296(5572):1439-43. Epub 2002/05/25. doi: 10.1126/science.1069660. PubMed PMID: 12029127.

16.   Fitzmaurice K, Petrovic D, Ramamurthy N, Simmons R, Merani S, Gaudieri S, Sims

S, Dempsey E, Freitas E, Lea S, McKiernan S, Norris S, Long A, Kelleher D, Klenerman P.

Molecular footprints reveal the impact of the protective HLA-A*03 allele in hepatitis C virus

infection. Gut. 2011;60(11):1563-71. Epub 2011/05/10. doi: 10.1136/gut.2010.228403.

PubMed PMID: 21551190; PMCID: 3184218.

17.   Rubnitz J, Subramani S. The minimum amount of homology required for homologous

recombination in mammalian cells. Molecular and cellular biology. 1984;4(11):2253-8. Epub

1984/11/01. doi: 10.1128/mcb.4.11.2253. PubMed PMID: 6096689; PMCID: 369052.

18.   Sola I, Moreno JL, Zuniga S, Alonso S, Enjuanes L. Role of nucleotides immediately

flanking the transcription-regulating sequence core in coronavirus subgenomic mRNA

synthesis. Journal of virology. 2005;79(4):2506-16. Epub 2005/02/01. doi:

10.1128/JVI.79.4.2506-2516.2005. PubMed PMID: 15681451; PMCID: 546574.

19.   Parker MD, Lindsey BB, Leary S, Gaudieri S, Chopra A, Wyles M, Angyal A, Green

LR, Parsons P, Tucker RM, Brown R, Groves D, Johnson K, Carrilero L, Heffer J, Partridge

DG, Evans C, Raza M, Keeley AJ, Smith N, Filipe ADS, Shepherd JG, Davis C, Bennett S,

Sreenu VB, Kohl A, Aranday-Cortes E, Tong L, Nichols J, Thomson EC, Consortium C-GU,

Wang D, Mallal S, de Silva TI. Subgenomic RNA identification in SARS-CoV-2 genomic

sequencing data. Genome Res. 2021;31(4):645-58. Epub 2021/03/17. doi:

10.1101/gr.268110.120. PubMed PMID: 33722935; PMCID: PMC8015849.

20.   Kim D, Lee JY, Yang JS, Kim JW, Kim VN, Chang H. The Architecture of SARS-

CoV-2 Transcriptome. Cell. 2020;181(4):914-21 e10. Epub 2020/04/25. doi:

10.1016/j.cell.2020.04.011. PubMed PMID: 32330414; PMCID: PMC7179501.

21.   Chandrashekar A, Liu J, Martinot AJ, McMahan K, Mercado NB, Peter L, Tostanoski

LH, Yu J, Maliga Z, Nekorchuk M, Busman-Sahay K, Terry M, Wrijil LM, Ducat S,

Martinez DR, Atyeo C, Fischinger S, Burke JS, Slein MD, Pessaint L, Van Ry A,

Greenhouse J, Taylor T, Blade K, Cook A, Finneyfrock B, Brown R, Teow E, Velasco J,

Zahn R, Wegmann F, Abbink P, Bondzie EA, Dagotto G, Gebre MS, He X, Jacob-Dolan C,

Kordana N, Li Z, Lifton MA, Mahrokhian SH, Maxfield LF, Nityanandam R, Nkolola JP,

Schmidt AG, Miller AD, Baric RS, Alter G, Sorger PK, Estes JD, Andersen H, Lewis MG,

Barouch DH. SARS-CoV-2 infection protects against rechallenge in rhesus macaques.

Science. 2020;369(6505):812-7. doi: 10.1126/science.abc4776.

22.   Wölfel R, Corman VM, Guggemos W, Seilmaier M, Zange S, Müller MA, Niemeyer

D, Jones TC, Vollmar P, Rothe C, Hoelscher M, Bleicker T, Brünink S, Schneider J, Ehmann

R, Zwirglmaier K, Drosten C, Wendtner C. Virological assessment of hospitalized patients

with COVID-2019. Nature. 2020;581(7809):465-9. doi: 10.1038/s41586-020-2196-x.

23.   Lorenz R, Bernhart SH, Honer Zu Siederdissen C, Tafer H, Flamm C, Stadler PF,

Hofacker IL. ViennaRNA Package 2.0. Algorithms for molecular biology : AMB. 2011;6:26.

Epub 2011/11/26. doi: 10.1186/1748-7188-6-26. PubMed PMID: 22115189; PMCID:

3319429.

24.   Laing C, Wen D, Wang JT, Schlick T. Predicting coaxial helical stacking in RNA

junctions. Nucleic acids research. 2012;40(2):487-98. Epub 2011/09/16. doi:

10.1093/nar/gkr629. PubMed PMID: 21917853; PMCID: 3258123.

25.   de la Pena M, Dufour D, Gallego J. Three-way RNA junctions with remote tertiary

contacts: a recurrent and highly versatile fold. Rna. 2009;15(11):1949-64. Epub 2009/09/11.

doi: 10.1261/rna.1889509. PubMed PMID: 19741022; PMCID: 2764472.

26.   Hua L, Song Y, Kim N, Laing C, Wang JT, Schlick T. CHSalign: A Web Server That

Builds upon Junction-Explorer and RNAJAG for Pairwise Alignment of RNA Secondary

Structures with Coaxial Helical Stacking. PloS one. 2016;11(1):e0147097. Epub 2016/01/21.

doi: 10.1371/journal.pone.0147097. PubMed PMID: 26789998; PMCID: 4720362.

27.   Chapman EG, Moon SL, Wilusz J, Kieft JS. RNA structures that resist degradation by

Xrn1 produce a pathogenic Dengue virus RNA. eLife. 2014;3:e01892. Epub 2014/04/03. doi:

10.7554/eLife.01892. PubMed PMID: 24692447; PMCID: 3968743.

28.   Gaudieri S, Rauch A, Park LP, Freitas E, Herrmann S, Jeffrey G, Cheng W, Pfafferott

K, Naidoo K, Chapman R, Battegay M, Weber R, Telenti A, Furrer H, James I, Lucas M,

Mallal SA. Evidence of viral adaptation to HLA class I-restricted immune pressure in chronic

hepatitis C virus infection. J Virol. 2006;80(22):11094-104. Epub 2006/10/31. doi:

10.1128/JVI.00912-06. PubMed PMID: 17071929; PMCID: 1642167.

29.   Brumme ZL, Kinloch NN, Sanche S, Wong A, Martin E, Cobarrubias KD, Sandstrom

P, Levett PN, Harrigan PR, Joy JB. Extensive host immune adaptation in a concentrated

North American HIV epidemic. Aids. 2018;32(14):1927-38. Epub 2018/07/27. doi:

10.1097/QAD.0000000000001912. PubMed PMID: 30048246; PMCID: 6125742.

30.   Katoh J, Kawana-Tachikawa A, Shimizu A, Zhu D, Han C, Nakamura H, Koga M,

Kikuchi T, Adachi E, Koibuchi T, Gao GF, Brumme ZL, Iwamoto A. Rapid HIV-1 Disease

Progression in Individuals Infected with a Virus Adapted to Its Host Population. PloS one.

2016;11(3):e0150397. Epub 2016/03/10. doi: 10.1371/journal.pone.0150397. PubMed

PMID: 26953793; PMCID: 4783116.

31.   Peng H, Yang LT, Wang LY, Li J, Huang J, Lu ZQ, Koup RA, Bailer RT, Wu CY.

Long-lived memory T lymphocyte responses against SARS coronavirus nucleocapsid protein

in SARS-recovered patients. Virology. 2006;351(2):466-75. Epub 2006/05/13. doi:

10.1016/j.virol.2006.03.036. PubMed PMID: 16690096.

32.   Yang Y, Yan W, Hall B, Jiang X. Characterizing transcriptional regulatory sequences

in coronaviruses and their role in recombination. bioRxiv. 2020. Epub 2020/06/27. doi:

10.1101/2020.06.21.163410. PubMed PMID: 32587968; PMCID: PMC7310624.

33.   Sola I, Almazan F, Zuniga S, Enjuanes L. Continuous and Discontinuous RNA

Synthesis in Coronaviruses. Annual review of virology. 2015;2(1):265-88. Epub 2016/03/10.

doi: 10.1146/annurev-virology-100114-055218.   PubMed   PMID: 26958916;   PMCID:

6025776.

34.   Kopecky-Bromberg SA, Martinez-Sobrido L, Frieman M, Baric RA, Palese P. Severe

acute respiratory syndrome coronavirus open reading frame (ORF) 3b, ORF 6, and

nucleocapsid proteins function as interferon antagonists. Journal of virology. 2007;81(2):548-

57.   Epub 2006/11/17.   doi: 10.1128/JVI.01782-06.   PubMed   PMID: 17108024;   PMCID:

PMC1797484.

35.   Lokugamage KG, Hage A, Schindewolf C, Rajsbaum R, Menachery VD. SARS-

CoV-2 is sensitive to type I interferon pretreatment. bioRxiv. 2020. Epub 2020/06/09. doi:

10.1101/2020.03.07.982264. PubMed PMID: 32511335; PMCID: PMC7239075.

36.   Hu Y, Li W, Gao T, Cui Y, Jin Y, Li P, Ma Q, Liu X, Cao C. The Severe Acute

Respiratory Syndrome Coronavirus Nucleocapsid Inhibits Type I Interferon Production by

Interfering with TRIM25-Mediated RIG-I Ubiquitination. Journal of virology. 2017;91(8).

Epub   2017/02/06.   doi: 10.1128/JVI.02143-16.   PubMed   PMID: 28148787;   PMCID:

PMC5375661.

37.   Manokaran G, Finol E, Wang C, Gunaratne J, Bahl J, Ong EZ, Tan HC, Sessions OM,

Ward AM, Gubler DJ, Harris E, Garcia-Blanco MA, Ooi EE. Dengue subgenomic RNA

binds TRIM25 to inhibit interferon expression for epidemiological fitness. Science.

2015;350(6257):217-21. Epub 2015/07/04. doi: 10.1126/science.aab3369. PubMed PMID:

26138103; PMCID: PMC4824004.

38.   Zuniga S, Cruz JL, Sola I, Mateos-Gomez PA, Palacio L, Enjuanes L. Coronavirus

nucleocapsid protein facilitates template switching and is required for efficient transcription.

Journal of virology. 2010;84(4):2169-75. Epub 2009/12/04. doi: 10.1128/JVI.02011-09.

PubMed PMID: 19955314; PMCID: PMC2812394.

39. Thorne LG, Bouhaddou M, Reuschl A-K, Zuliani-Alvarez L, Polacco B, Pelin A,

Batra J, Whelan MVX, Ummadi M, Rojc A, Turner J, Obernier K, Braberg H, Soucheray M,

Richards A, Chen K-H, Harjai B, Memon D, Hosmillo M, Hiatt J, Jahun A, Goodfellow IG,

Fabius JM, Shokat K, Jura N, Verba K, Noursadeghi M, Beltrao P, Swaney DL, Garcia-

Sastre A, Jolly C, Towers GJ, Krogan NJ. Evolution of enhanced innate immune evasion by

the SARS-CoV-2 B.1.1.7 UK variant. bioRxiv. 2021.

40. Guo K, Barrett BS, Mickens KL, Hasenkrug KJ, Santiago ML. Interferon Resistance

of Emerging SARS-CoV-2 Variants. bioRxiv. 2021.

41. Jiang HW, Zhang HN, Meng QF, Xie J, Li Y, Chen H, Zheng YX, Wang XN, Qi H,

Zhang J, Wang PH, Han ZG, Tao SC. SARS-CoV-2 Orf9b suppresses type I interferon

responses by targeting TOM70. Cell Mol Immunol. 2020;17(9):998-1000. Epub 2020/07/31.

doi: 10.1038/s41423-020-0514-8. PubMed PMID: 32728199; PMCID: PMC7387808.

42. Miorin L, Kehrer T, Sanchez-Aparicio MT, Zhang K, Cohen P, Patel RS, Cupic A,

Makio T, Mei M, Moreno E, Danziger O, White KM, Rathnasinghe R, Uccellini M, Gao S,

Aydillo T, Mena I, Yin X, Martin-Sancho L, Krogan NJ, Chanda SK, Schotsaert M, Wozniak

RW, Ren Y, Rosenberg BR, Fontoura BMA, Garcia-Sastre A. SARS-CoV-2 Orf6 hijacks

Nup98 to block STAT nuclear import and antagonize interferon signaling. Proc Natl Acad

Sci U S A. 2020;117(45):28344-54. Epub 2020/10/25. doi: 10.1073/pnas.2016650117.

PubMed PMID: 33097660; PMCID: PMC7668094.

43. Oh SJ, Shin OS. SARS-CoV-2 Nucleocapsid Protein Targets RIG-I-Like Receptor

Pathways to Inhibit the Induction of Interferon Response. Cells. 2021;10(3). Epub

2021/04/04. doi: 10.3390/cells10030530. PubMed PMID: 33801464; PMCID: PMC7999926.

44.   Parker MD, Lindsey BB, Shah DR, Hsu S, Keeley AJ, Partridge DG, Leary S, Cope

A, State A, Johnson K, Ali N, Raghei R, Heffer J, Smith N, Zhang P, Gallis M, Louka SF,

Whiteley M, Foulkes BH, Christou S, Wolverson P, Pohare M, Hansford SE, Green LR,

Evans C, Raza M, Wang D, Gaudieri S, Mallal S, , de Silva TI. Altered Subgenomic RNA

Expression in SARS-CoV-2 B.1.1.7 Infections. bioRxiv. 2021.

45.   Earle KA, Ambrosino DM, Fiore-Gartland A, Goldblatt D, Gilbert PB, Siber GR,

Dull P, Plotkin SA. Evidence for antibody as a protective correlate for COVID-19 vaccines.

Vaccine. 2021;39(32):4423-8. doi: 10.1016/j.vaccine.2021.05.063.

46.   Khoury DS, Cromer D, Reynaldi A, Schlub TE, Wheatley AK, Juno JA, Subbarao K,

Kent SJ, Triccas JA, Davenport MP. Neutralizing antibody levels are highly predictive of

immune protection from symptomatic SARS-CoV-2 infection. Nature medicine.

2021;27(7):1205-11. doi: 10.1038/s41591-021-01377-8.

47.   Bergwerk M, Gonen T, Lustig Y, Amit S, Lipsitch M, Cohen C, Mandelboim M, Gal

Levin E, Rubin C, Indenbaum V, Tal I, Zavitan M, Zuckerman N, Bar-Chaim A, Kreiss Y,

Regev-Yochay G. Covid-19 Breakthrough Infections in Vaccinated Health Care Workers.

New England Journal of Medicine. 2021. doi: 10.1056/nejmoa2109072.

FIGURES

798
799
Fig 1. Proportion of weekly deposited SARS-CoV-2 sequences globally (n=455774). The
801 D614G (B.1) variant has become one of the dominant forms globally. Note a small
802 proportion of deposited sequences did not include information regarding specific collection
803 date and as such were excluded.
804
805

Fig 2. Proportion of weekly deposited SARS-CoV-2 sequences by region. The proportion
of R203/G204 to K203/R204 sub-variants of the D614G variant differs in different regions
with recent increases in the frequency of new variants.

Fig 3. The configuration of canonical sgRNAs and the novel non-canonical nucleocapsid sgRNA (N*) in SARS-CoV-2. The bottom bar illustrates the presence of the leader sequence (blue text) followed by the transcription-regulating sequence (TRS; red text) within the genomic sequence that continues into the first ORF 1a. The presence of other canonical sgRNA transcripts in which the leader sequence and TRS precede the start codon (methionine; pink) of the other proteins are shown. The presence of the novel non-canonical sgRNA transcript containing the K203/R204 polymorphisms (N*) is shown. The ARTIC primer locations and resultant amplicons are shown.

833
834
835

Fig 4. Exploration of sgRNAs in 981 samples from Sheffield, UK. A. A heatmap showing presence or absence of sgRNAs from different ORFs. K203/R204 (KR)-containing sequences have evidence of the novel truncated N ORF sgRNA (N*, red, 233/553, 42%). An ORF sgRNA was deemed present if we could find >=1 read in support. Heatmap is ordered by the presence or absence of the novel sgRNA. There were a total of 448 R203/G204 (RG)-containing sequences and 1 had evidence of a novel sgRNA (likely false positive, Fig S2). **B**. Significantly higher (Mann-Whitney U p < 2.2e-16) total sgRNA in KR-containing compared to RG-containing sequences. **C**. Sub-genomic RNA is significant increased in KR-containing

compared to RG-containing sequences for a number of ORFs, most notably nucleocapsid (N;
Mann-Whitney U p = 2.06e-37 corrected for multiple testing using the Holm method). Y-axis
denotes square root transformed sub-genomic reads normalized to 100,000 genomic reads
from the same ARTIC amplicon. **D.** There is no difference in genomic RNA levels
(normalized to total mapped reads) between KR- and RG-containing sequences. *novel
sgRNA, ORF10 and ORF1a are excluded from this analysis due to ORF10 not being
expressed, difficulty in discriminating ORF1a sgRNA from genomic RNA and the novel
truncated N sgRNA is only being present in KR-containing sequences. ***** < 0.001, ** <**
0.01, * < 0.05. All p values shown are following correction for multiple testing with the
Holm method.
854
855
856

858
859
860
Fig 5. Spike 614 and Nucleocapsid 203/204 Status, Diagnostic Metrics and level of sub-
genomic and genomic RNA. A. E gene cycle threshold (CT) normalized to RNAseP CT
stratified by variant status in N = 478 individuals from Sheffield dataset with day of symptom
onset data available. This normalization was done to combine and display E gene CT data
from two different extraction protocols. Y-axis reversed to aid interpretation, as lower
normalized CT values equal higher virus levels. **B.** Normalized E gene CT vs the day of
sampling from day of symptom onset. P values provided are from a generalized multivariable
linear regression model (GLM) for the difference in normalized E gene CT value between
samples containing each variant, with extraction method and day of illness included in the
model (Table S6) **C.** Normalized (per 1000 genomic reads) sgRNA levels for ORFs S and N.
D. Normalized (per 100,000 mapped reads) genomic RNA levels for ORFs S and N.

872
873
874