Title: Pattern-recognition receptors are required for NLR-mediated plant immunity

Authors: Minhang Yuan[1,2], Zeyu Jiang[1,2], Guozhi Bi[3,4], Kinya Nomura[5], Menghui Liu[6], Sheng Yang He[5,7], Jian-Min Zhou[3,4] and Xiu-Fang Xin[1,8]*

Affiliations:

[1]National key Laboratory of Plant Molecular Genetics, CAS Center for Excellence in Molecular Plant Sciences, Institute of Plant Physiology and Ecology, Chinese Academy of Sciences, Shanghai, China.

[2]University of the Chinese Academy of Sciences, Beijing, China.

[3]State Key Laboratory of Plant Genomics, Institute of Genetics and Developmental Biology, Innovation Academy for Seed Design, Chinese Academy of Sciences, Beijing, China.

[4]CAS Center for Excellence in Biotic Interactions, University of Chinese Academy of Sciences, Beijing, China.

[5]Department of Energy Plant Research Laboratory, Plant Resilient Institute, Michigan State University, East Lansing, MI, USA.

[6]Key Laboratory of Plant Stress Biology, State Key Laboratory of Cotton Biology, School of Life Sciences, Henan University, Kaifeng, China.

[7]Howard Hughes Medical Institute, Michigan State University, East Lansing, MI, USA.

[8]CAS-JIC Center of Excellence for Plant and Microbial Sciences (CEPAMS), Institute of Plant Physiology and Ecology, Chinese Academy of Sciences, Shanghai, China.

*Correspondence to Xiu-Fang Xin (xinxf@sippe.ac.cn)

Abstract

The plant immune system is fundamental to the survival and productivity of plants in crop fields and natural ecosystems. Substantial evidence supports the prevailing notion that plants have evolved two branches of innate immune signaling, called pattern-triggered immunity (PTI) and effector-triggered immunity (ETI). PTI is triggered by conserved microbial patterns via surface-localized pattern-recognition receptors (PRRs), whereas ETI is activated by pathogen effector proteins via mostly intracellularly-localized receptors called nucleotide-binding, leucine-rich

repeat proteins (NLRs)[1-4]. PTI and ETI have traditionally been considered to be separate immune

signaling pathways and have evolved sequentially[5,6]. Here we show that, contrary to the

perception of PTI and ETI being separate immune signaling pathways, Arabidopsis PRR/co-

receptor mutants, *fls2/efr/cerk1* and *bak1/bkk1/cerk1* triple mutants, are greatly compromised in

ETI responses triggered by *Pseudomonas syrinage* effectors. We further identified the NADPH

oxidase (RBOHD)-mediated production of reactive oxygen species as a critical early signaling

event connecting PRR and NLR signaling cascades and show that PRR-mediated activation of

RBOHD is necessary for full activation of RBOHD during ETI. Furthermore, NLR signaling

rapidly augments the transcript and protein levels of key PTI components at an early stage and in

a salicylic acid-independent manner. Our study supports an alternative model in which PTI is in

fact an indispensable component of ETI during bacterial infection, implying that ETI halts

pathogen infection, in part, by directly co-opting the anti-pathogen mechanisms proposed for

PTI. This alternative model conceptually unites two major immune signaling pathways in the

plant kingdom and mechanistically explains the long-observed similarities in downstream

defense outputs between PTI and ETI.

Main

It is generally believed that PRRs and NLRs, with largely distinct activation mechanisms and

subcellular localizations, mediate different immune signaling pathways. PRRs are surface-

localized receptor-like kinases/proteins (RLKs/RLPs) with extracellular ligand-binding domain

to sense conserved molecular patterns, ranging from bacterial flagellin and fungal chitin

molecules from both pathogenic and nonpathogenic microbes. NLRs, on the other hand, are

intracellular proteins that sense pathogen-derived effector proteins inside the plant cell and can

be further classified into the coiled coil (CC)-type or Toll/interleukin-1 receptor (TIR)-type,

depending on their N-terminal domain[7]. However, PRR- and NLR-mediated signaling pathways

result in many similar downstream immune outputs, including defense gene expression,

production of reactive oxygen species (ROS) and callose deposition at the plant cell wall[8,9]. The

underlying reason is not clear and mechanistic relationship between the two immune pathways

remains enigmatic. Notably, while many PRR signaling components have been identified and

anti-pathogen mechanisms described, the downstream signaling events in ETI and how ETI halts

pathogen growth remain largely elusive, despite recent breakthroughs in the understanding of

NLR protein structures and activities[10-13].

Requirement of PRR/co-receptors for ETI

Using *Arabidopsis thaliana-Pseudomonas syringae* pathosystem, we accidentally discovered a

striking and unexpected role of PRR/co-receptors in ETI. Specifically, an "avirulent", ETI-

eliciting bacterial strain, *P. syringae* pv. *tomato* (*Pst*) DC3000(*avrRpt2*), which activates RPS2

(Resistance to *P. syringae* 2)-dependent ETI in wild-type Col-0 plants[14,15], failed to elicit

effective ETI in two separate PRR/co-receptor Arabidopsis mutants, *fls2/efr/cerk1* (*fec*) and

bak1/bkk1/cerk1 (*bbc*) mutants, which are mutated in major PRR/co-receptors recognizing

bacteria-associated molecular patterns[16]. As shown in Fig. 1a, the *fec* and *bbc* mutants did not

mount an effective ETI against *Pst* DC3000(*avrRpt2*). To determine whether a requirement of

PRR/co-receptors for ETI is specific to *Pst* DC3000(*avrRpt2*) or is a more general phenomenon,

we tested two other ETI-triggering "avirulent" effectors, AvrPphB and AvrRps4, which are

recognized by RPS5[17] and RPS4[18], respectively, in Arabidopsis Col-0 accession. We found that

the compromised ETI phenotype in *fec* and *bbc* mutants held true for both AvrPphB and

AvrRps4 (Extended Data Fig. 1), suggesting a potentially broad role of PRR/co-receptors in ETI

pathways. We subsequently focused on AvrRpt2-triggered ETI for in-depth characterization.

Hypersensitive response (HR), manifested by fast cell death under high bacterial inoculum, is a

hallmark of ETI. We tested HR phenotype in *fec* and *bbc* mutants in response to *Pst*

DC3000(*avrRpt2*) and found that, even though HR cell death eventually occurs in these mutants,

the rate was delayed, as shown by the compromised tissue collapse 7h after bacteria infiltration

(Fig. 1b).

For the past several decades, classical studies of ETI triggered by *Pst* DC3000 carrying

"avirulent" effector genes have been performed in the presence of all 36 endogenous effector

genes in *Pst* DC3000. Because many of the endogenous effectors in *Pst* DC3000 are linked to

interference of PTI and/or ETI via unclear mechanisms[19,20], it is not always easy to clearly

interpret the relationship between PTI and ETI during infection by avirulent *Pst* DC3000 strains.

We took advantage of the recent availability of *Pst* DC3000 strain D36E[21], in which all 36

effector genes are deleted and therefore is expected to activate only PTI, and D36E(*avrRpt2*)

strain, which delivers only AvrRpt2 and activates both PTI and RPS2-mediated ETI with no

interference from any endogenous *Pst* DC3000 effectors. Although D36E is greatly reduced in

virulence compared to *Pst* DC3000 (Fig. 1a), we could still observe a robust AvrRpt2-induced

ETI in Col-0 plants, with strain D36E(*avrRpt2*) growing significantly less than strain D36E (Fig.

1c). We found that AvrRpt2-triggered ETI was no longer detectable in both *fec* and *bbc* mutants

(Fig. 1c), demonstrating that the requirement of PRR/co-receptors for AvrRpt2-triggered ETI

was not caused by some hidden interactions between endogenous *Pst* DC3000 effectors and

Arabidopsis *fec* and *bbc* mutants.

Dual role of RBOHD in ROS during PTI and ETI

Previous studies have shown that AvrRpt2 cleaves the plant protein RIN4 (RPM1-interacting

protein 4), leading to activation of the NLR protein RPS2[14,15]. To understand why AvrRpt2-

mediated ETI is lost in *fec* and *bbc* mutants, we examined the cleavage of the RIN4 protein by

AvrRpt2. Results showed that D36E(*avrRpt2*)-induced RIN4 protein depletion was normal in the

fec and *bbc* mutants compared to that in Col-0 plants (Fig. 1d). Gene expression analysis showed

that *RPS2* transcript level is also similar in all genotypes after bacteria inoculation (Extended

Data Fig. 2). We then sought to assay downstream signaling events of AvrRpt2-triggered ETI

and examined MAPK phosphorylation level, which has been reported to be stronger and more

sustained after AvrRpt2 recognition by RPS2, compared to that induced during PTI[22]. As shown

in Fig. 1e, we observed strong ETI-associated MPK3/6 phosphorylation (i.e., at 4 or 8 h post

inoculation) in Col-0 plants, which however remained intact in *fec* and *bbc* mutants.

Another important immune response associated with both PTI and ETI is production of ROS,

including superoxide and hydrogen peroxide (H_2O_2), which have been proposed to act as defense

molecules that kill pathogens and signaling molecules that further activate immune responses[23].

Using luminol-horseradish peroxidase (HRP)-based method, we examined PTI- and ETI-

associated ROS production in a transgenic *avrRpt2* plant, in which *avrRpt2* expression is driven

by a dexamethasone (DEX)-inducible promoter[24]. In this system, PTI and ETI activation can be

initiated separately or in combination using PAMP (e.g., flg22, a 22-aa peptide derived from

bacterial flagellin) and DEX treatments in the absence of bacterial infection. As shown in Fig. 1f,

flg22 alone triggered a fast and transient ROS burst as reported before[25], while DEX-induced

expression of AvrRpt2 alone triggered only a weak and kinetically slower ROS burst.

Interestingly, co-treatment of flg22 and DEX triggered a strong and sustained second-phase ROS

burst, peaking at 2h to 3h after treatment, and lasted for several hours (Fig. 1f, g), a profile that

bears a remarkable similarity to previously observed during bacteria-triggered ETI[26,27]. This

result raises an intriguing possibility that activation of PTI may be required for the production of

a strong and sustained ROS characteristic of ETI. To test this hypothesis, we generated

bbc/DEX::avrRpt2 plants by transforming the *DEX::avrRpt2* construct into the *bbc* mutant plant

(as well as Col-0 plant as control). Independent lines in which the *avrRpt2* expression level was

similar or even higher than that in Col-0/*DEX::avrRpt2* plants were chosen (Extended Data Fig.

3) for further analysis. As shown in Fig. 1h, i, in the *bbc/DEX::avrRpt2* plants, not only flg22-

induced first-phase ROS is absent, but also the second-phase AvrRpt2-triggered ROS burst is

almost completely abolished, clearly demonstrating a requirement of PRR/co-receptors for ETI-

associated ROS production.

To examine whether PTI- and ETI-associated ROS bursts are produced at the same or different

subcellular compartments, ROS production was monitored with the fluorescent dye H_2DCFDA,

which can cross the plasma membrane of the plant cell and detect both apoplastic and cytosolic

ROS[28]. As shown in Fig. 2a, strong fluorescent signal was detected in the apoplastic space of

Col-0 leaves 5h post infiltration of D36E(*avrRpt2*). The apoplastic signal was much weaker in

the *bbc* mutant plant, which was indistinguishable compared to the *rps2* control plant infiltrated

with D36E(*avrRpt2*) or Col-0 plant infiltrated with D36E (Fig. 2a). Two classes of enzymes, the

NADPH oxidases such as respiratory burst oxidase homolog D (RBOHD) and peroxidases, have

been shown to be involved in generating apoplastic ROS in pathogen-infected plant leaves[29,30].

We therefore investigated which class is involved in generating AvrRpt2-triggered ROS by using

chemical inhibitors diphenylene iodonium (DPI), which inhibits NADPH oxidases, and

salicylhydroxamic acid (SHAM) and sodium azide, which inhibit peroxidase activities[26,31]. As

shown in Extended Data Fig. 4a-c, co-treatment of DPI, but not SHAM or sodium azide, with

flg22 and DEX almost completely blocked ETI-associated ROS and greatly compromised PTI-

associated ROS. Interestingly, when we added these inhibitors at 40 min after flg22+DEX

treatment (i.e., after PTI-associated ROS and before the start of ETI-associated ROS), still only

DPI, but not SHAM or sodium azide, almost completely blocked ETI-ROS (Fig. 2b), indicating

that NADPH oxidases mediate ETI-associated ROS. We further tested whether NADPH oxidase

RBOHD, which has been shown to play a prominent role in generating pathogen-induced

ROS[29,32,33], mediates the ETI-associated ROS. As shown in Fig. 2c, D36E(*avrRpt2*)-induced

apoplastic ROS, as detected *in planta* by the H_2DCFDA dye, was completely lost in the *rbohd*

plant. Consistently, we detected a much compromised ETI resistance against *Pst*

DC3000(*avrRpt2*) in the *rbohd* mutant plant (Fig. 2d, Extended Data Fig. 5b). Notably, *rbohd*

mutant plants grew to sizes that were similar to wild-type Col-0 plants under optimized growth

conditions and, under these conditions, *rbohd* mutant plants showed similar or even slightly

enhanced susceptibility to *Pst* DC3000 (Fig. 2d, Extended Data Fig. 5a, b). Altogether, our

results suggest a critical dual role of RBOHD in PTI and ETI.

Requirement of PTI activation of RBOHD for ROS during ETI

We next assayed the transcript and protein level of RBOHD and found that the transcript (Fig.

2e) and protein level (Fig. 2f) of RBOHD are induced both by D36E and, interestingly, to a much

higher level, by D36E(*avrRpt2*) inoculation in Col-0 plant. Surprisingly, the strong induction of

RBOHD transcript and protein by D36E(*avrRpt2*) occurred at a comparable level in *bbc* mutant

plants (Fig. 2e, f). These results indicate that neither *RBOHD* transcript nor RBOHD protein

accumulation accounts for the compromised ROS production in the *bbc* mutant and suggests an

involvement of PRRs/PTI in post-translational regulation of RBOHD during ETI. Previous

studies have reported several classes of kinases, including calcium dependent protein kinases

(CPKs) and *Botrytis*-induced kinase 1 (BIK1), involved in phosphorylating RBOHD for ROS

production[32-36]. We examined the ETI-ROS level in the *cpk4/5/6/11* quadruple mutant and *bik1*

mutant plants, and found that ETI-associated ROS was reduced in *bik1* mutant but did not seem

to be affected in *cpk4/5/6/11* mutant plant (Extended Data Fig. 6), suggesting that BIK1

contributes to the production of ETI-ROS. BIK1 was reported to phosphorylate RBOHD at

multiple sites including S39, S343 and S347 during PTI activation[32,33]. We therefore examined

RBOHD phosphorylation levels in protoplasts prepared from Col-0/*DEX::avrRpt2* and

bbc/*DEX::avrRpt2* plants and transformed with an DNA construct expressing FLAG-RBOHD.

PTI and ETI in these protoplasts were activated by flg22 and DEX, respectively. A 35S promoter

was used to express FLAG-RBOHD to ensure similar protein levels during various treatments.

While no phosphorylation at S343/S347 in Col-0 protoplasts was detected 2.5h after treatment

with flg22, which is known to induce a PTI-associated transient phosphorylation on RBOHD,

DEX alone reproducibly induced a weak phosphorylation of S343/S347 in Col-0 protoplasts 2.5h

after treatment (Fig. 2g). Strikingly, a flg22+DEX treatment induced a much stronger

phosphorylation on S343/S347 in Col-0 background 2.5h after treatment (Fig. 2g), suggesting a

synergistic effect of PTI and ETI in phosphorylating RBOHD. In contrast, no phosphorylation

was detected in the *bbc* background with flg22, DEX or flg22+DEX treatment, confirming the

requirement of PRR/co-receptors for RBOHD phosphorylation during ETI. Collectively, our

results demonstrate that two classes of immune receptors coordinate to ensure the abundance (i.e.

by RPS2) and activity (i.e. by PRRs) of RBOHD for generating robust ETI-ROS. Interestingly,

S343/S347 phosphorylation of RBOHD has previously been shown to be important for ETI

resistance and restriction of bacterial growth[37].

PRRs control a subset of ETI-associated transcriptome

The requirement of PRR/co-receptors for activation of RBOHD and a strong up-regulation of

RBOHD during ETI (Fig. 2e, f) are intriguing and suggest that ETI may have evolved to co-opt

RBOHD and other components of the PTI pathway as part of its mechanism. We therefore

examined the expression patterns of other components of the PTI pathway and the rest of

Arabidopsis transcriptome by RNAseq (Fig. 3a). Bacteria were infiltrated at a high dose (i.e.,

~2×10^7 cfu/mL) and expression was examined at early time points (i.e. 3h and 6h post

infiltration) to ensure similar bacterial populations in Col-0 and *bbc* plants at the sampling times

(Extended Data Fig. 7a). We found that, at 3h post infiltration, D36E(*avrRpt2*) already induced

significantly different expression of many genes (i.e., more than 4,000 genes) compared to D36E

in Col-0 plant (Extended Data Fig. 7b), suggesting that 3h is sufficient for delivery of AvrRpt2

into the plant cell and triggering strong ETI-associated gene expression. We therefore focused

analysis on 3h time point in order to reveal early, and likely more direct, changes of ETI-

associated gene expression. Many genes are differentially regulated at this early time point

between Col-0 and *bbc* plants in response to PTI-inducing D36E (Fig. 3b), as expected.

Interestingly, these genes show similar expression pattern in Col-0 and *bbc* plants after

D36E(*avrRpt2*) inoculation to elicit ETI (Fig. 3b), suggesting that ETI can largely restore PTI-

associated global gene expression in the *bbc* plant. Similar trends were observed for genes

associated with salicylic acid, jasmonate and ethylene pathways (Extended Data Fig. 7c-e).

Despite the overall similarity in global gene expression patterns between Col-0 and *bbc* mutant

infected by D36E(*avrRpt2*), we did notice that a subset of 272 genes were differentially

expressed (Supplementary table 1). GO analysis of the 209 down-regulated genes (i.e., higher

expression in Col-0 compared to that in the *bbc* mutant) revealed several immune-related terms,

such as "response to chitin" and "response to hydrogen peroxide" (Extended Data Fig. 8a).

Interestingly, we found that a cluster of *WRKY* genes including *WRKY22/29* and *FRK1* (*Flg22-*

induced Receptor-like Kinase 1), which are canonical marker genes of flg22-induced PTI

pathway[38], are down-regulated in the *bbc* plant during AvrRpt2-triggered ETI (Extended Data

Fig. 8b, c). This suggests that, despite the general rescue of global PTI-associated gene

expression by ETI in the *bbc* mutant, the WRKY-FRK1 branch represents a unique immune

branch, the activation of which during ETI requires PRR/co-receptors.

Augmentation of key PTI components by ETI in an SA-independent manner

We noticed an interesting expression pattern for many PTI signaling genes. As shown in Fig. 3c,

while PTI-inducing D36E can induce, to a moderate level, many key PTI components, namely

BAK1, *BIK1*, *XLG2/AGB1/AGG2*[39], *MKK4/5* and *MPK3*, ETI-inducing D36E(*avrRpt2*) induced

these genes to a much higher level. Importantly, the strong activation of these PTI components

by ETI is independent of PRR/co-receptors, since it occurs in the *bbc* mutant. Noticeably, BIK1

and some other PBLs, but not PBL1, are strongly induced after D36E(*avrRpt2*) inoculation

(Extended Data Fig. 9), suggesting differential contribution of different members of the

BIK1/PBL family to ETI. Quantitative RT-PCR was performed to confirm the RNAseq results

(Fig. 3d), and western blot further confirmed, at the protein level, the ETI up-regulation of

several key components, including BAK1, BIK1 and MPK3 (Fig. 3e). Our results suggest that

part of the AvrRpt2-ETI response is to ensure rapid high-level expression of key components of

the PTI pathway, consistent with PTI being an essential component of ETI. We propose that this

ETI-mediated up-regulation of PTI components is also likely an important part of a mechanism

to overcome the negative regulation of PTI by endogenous "braking" systems of plants and

exogenous pathogen effectors during infections[2,40].

A previous study showed that SA treatment could up-regulate PRR protein level[41]. We therefore

tested gene expression in the SA-deficient *sid2* plant and found that PTI components were still

up-regulated by D36E(*avrRpt2*) inoculation (Supplementary Fig. 10a). In addition, we examined

in our RNAseq dataset the expression pattern of responsive genes to SA and N-hydroxy-

pipecolic acid (NHP), which have been shown to function synergistically in plant immunity[42,43].

Our results showed that both SA- (Extended Data Fig. 7c) and NHP- (Extended Data Fig. 10b)

responsive genes had similar transcript level in the Col-0 and *bbc* plants after D36E(*avrRpt2*)

inoculation, suggesting intact SA/NHP signaling in the *bbc* mutant during AvrRpt2-ETI.

Therefore, ETI seems to "re-enforce" the PTI pathway in a SA/NHP-independent manner.

Discussion

Our study reveals a surprising requirement of PRR/co-receptors for effective ETI responses and

suggests a model in which ETI co-opts the PTI machinery, including the BIK1-RBOHD module,

as an indispensable component (Fig. 4). In particular, we found that PRRs and NLRs, the two

primary classes of plant immune receptors, function synergistically to ensure a robust level as

well as the fully active status of a key immune component RBOHD, which mediates ETI-ROS

generation and full disease resistance. We identified PRR-mediated RBOHD phosphorylation at

S343/S347 sites as one of the mechanistic links between PTI and ETI.

Our study sheds light on a long-standing puzzle in the field of plant immunity with respect to the

enigmatic similarities in many PTI- and ETI-associated cellular defense features. Our model is

supported by a parallel study by Ngou et al. (see back-to-back submission., Manuscript number

"BIORXIV/2020/034173"), who focus on the *P. syringae* effector AvrRps4 recognized by TIR-

type NLRs (RPS4/RRS1), while we focus on RPS2, a CC-type NLR. Our complementary data

suggest conservation of the discovered mechanism for two different types of NLRs, which

account for the vast majority of pathogen-sensing NLRs in plants. Intriguingly, synergistic

interaction between surface and intracellular immune receptors in animals and humans has also

been reported[44-47], suggesting a possible conceptual parallel in immune receptor functions across

kingdoms.

The demonstration of PTI as an integral component of ETI has significant implications in

understanding how ETI resistance mechanisms prevent pathogen growth. Specifically, several

PTI-associated anti-pathogen mechanisms have been described recently, including suppression of

bacterial type III secretion[48-50], inhibition of bacterial motility by lignification[51] and restriction of

nutrient acquisition[52]. Our study suggests that ETI co-opts these PTI mechanisms to halt

pathogen growth. Our work likely has significant practical implications as well, as it suggests the

possibility for carefully controlled augmentation of PTI components as a new strategy to broadly

increase the effectiveness of ETI against numerous plant diseases in crop fields.

References

1   Yu, X., Feng, B., He, P. & Shan, L. From Chaos to Harmony: Responses and Signaling upon
Microbial Pattern Recognition. *Annu Rev Phytopathol* **55**, 109-137, doi:10.1146/annurev-phyto-
080516-035649 (2017).
2   Couto, D. & Zipfel, C. Regulation of pattern recognition receptor signalling in plants. *Nat Rev*
Immunol **16**, 537-552, doi:10.1038/nri.2016.77 (2016).
3   Spoel, S. H. & Dong, X. How do plants achieve immunity? Defence without specialized immune
cells. *Nat Rev Immunol* **12**, 89-100, doi:10.1038/nri3141 (2012).
4   Cui, H., Tsuda, K. & Parker, J. E. Effector-triggered immunity: from pathogen perception to
robust defense. *Annual review of plant biology* **66**, 487-511, doi:10.1146/annurev-arplant-
050213-040012 (2015).
5   Jones, J. D. & Dangl, J. L. The plant immune system. *Nature* **444**, 323-329,
doi:10.1038/nature05286 (2006).
6   Chisholm, S. T., Coaker, G., Day, B. & Staskawicz, B. J. Host-microbe interactions: shaping the
evolution of the plant immune response. *Cell* **124**, 803-814, doi:10.1016/j.cell.2006.02.008
(2006).
7   Jones, J. D., Vance, R. E. & Dangl, J. L. Intracellular innate immune surveillance devices in plants
and animals. *Science* **354**, doi:10.1126/science.aaf6395 (2016).
8   Tsuda, K. & Katagiri, F. Comparing signaling mechanisms engaged in pattern-triggered and
effector-triggered immunity. *Current opinion in plant biology* **13**, 459-465,
doi:10.1016/j.pbi.2010.04.006 (2010).
9   Peng, Y., van Wersch, R. & Zhang, Y. Convergent and Divergent Signaling in PAMP-Triggered
Immunity and Effector-Triggered Immunity. *Mol Plant Microbe Interact* **31**, 403-409,
doi:10.1094/MPMI-06-17-0145-CR (2018).

10 Wang, J. *et al.* Reconstitution and structure of a plant NLR resistosome conferring immunity.
Science **364**, doi:10.1126/science.aav5870 (2019).
11 Wang, J. *et al.* Ligand-triggered allosteric ADP release primes a plant NLR complex. *Science* **364**,
doi:10.1126/science.aav5868 (2019).
12 Wan, L. *et al.* TIR domains of plant immune receptors are NAD(+)-cleaving enzymes that
promote cell death. *Science* **365**, 799-803, doi:10.1126/science.aax1771 (2019).
13 Horsefield, S. *et al.* NAD(+) cleavage activity by animal and plant TIR domains in cell death
pathways. *Science* **365**, 793-799, doi:10.1126/science.aax1911 (2019).
14 Axtell, M. J. & Staskawicz, B. J. Initiation of RPS2-specified disease resistance in Arabidopsis is
coupled to the AvrRpt2-directed elimination of RIN4. *Cell* **112**, 369-377, doi:10.1016/s0092-
8674(03)00036-9 (2003).
15 Mackey, D., Belkhadir, Y., Alonso, J. M., Ecker, J. R. & Dangl, J. L. Arabidopsis RIN4 is a target of
the type III virulence effector AvrRpt2 and modulates RPS2-mediated resistance. *Cell* **112**, 379-
389, doi:10.1016/s0092-8674(03)00040-0 (2003).
16 Xin, X. F. *et al.* Bacteria establish an aqueous living space in plants crucial for virulence. *Nature*
539, 524-529, doi:10.1038/nature20166 (2016).
17 Shao, F. *et al.* Cleavage of Arabidopsis PBS1 by a bacterial type III effector. *Science* **301**, 1230-
1233, doi:10.1126/science.1085671 (2003).
18 Gassmann, W., Hinsch, M. E. & Staskawicz, B. J. The Arabidopsis RPS4 bacterial-resistance gene
is a member of the TIR-NBS-LRR family of disease-resistance genes. *Plant J* **20**, 265-277,
doi:10.1046/j.1365-313x.1999.t01-1-00600.x (1999).
19 Xin, X. F., Kvitko, B. & He, S. Y. Pseudomonas syringae: what it takes to be a pathogen. *Nat Rev*
Microbiol **16**, 316-328, doi:10.1038/nrmicro.2018.17 (2018).
20 Toruno, T. Y., Stergiopoulos, I. & Coaker, G. Plant-Pathogen Effectors: Cellular Probes Interfering
with Plant Defenses in Spatial and Temporal Manners. *Annu Rev Phytopathol* **54**, 419-441,
doi:10.1146/annurev-phyto-080615-100204 (2016).
21 Wei, H. L. *et al.* Pseudomonas syringae pv. tomato DC3000 Type III Secretion Effector
Polymutants Reveal an Interplay between HopAD1 and AvrPtoB. *Cell Host Microbe* **17**, 752-762,
doi:10.1016/j.chom.2015.05.007 (2015).
22 Tsuda, K. *et al.* Dual regulation of gene expression mediated by extended MAPK activation and
salicylic acid contributes to robust innate immunity in Arabidopsis thaliana. *PLoS Genet* **9**,
e1004015, doi:10.1371/journal.pgen.1004015 (2013).
23 Qi, J., Wang, J., Gong, Z. & Zhou, J. M. Apoplastic ROS signaling in plant immunity. *Current*
opinion in plant biology **38**, 92-100, doi:10.1016/j.pbi.2017.04.022 (2017).
24 McNellis, T. W. *et al.* Glucocorticoid-inducible expression of a bacterial avirulence gene in
transgenic Arabidopsis induces hypersensitive cell death. *Plant J* **14**, 247-257,
doi:10.1046/j.1365-313x.1998.00106.x (1998).
25 Felix, G., Duran, J. D., Volko, S. & Boller, T. Plants have a sensitive perception system for the
most conserved domain of bacterial flagellin. *Plant J* **18**, 265-276, doi:10.1046/j.1365-
313x.1999.00265.x (1999).
26 Levine, A., Tenhaken, R., Dixon, R. & Lamb, C. H2O2 from the oxidative burst orchestrates the
plant hypersensitive disease resistance response. *Cell* **79**, 583-593, doi:10.1016/0092-
8674(94)90544-4 (1994).
27 Chandra, S., Martin, G. B. & Low, P. S. The Pto kinase mediates a signaling pathway leading to
the oxidative burst in tomato. *Proc Natl Acad Sci U S A* **93**, 13393-13397,
doi:10.1073/pnas.93.23.13393 (1996).
28 Tian, S. *et al.* Plant Aquaporin AtPIP1;4 Links Apoplastic H2O2 Induction to Disease Immunity
Pathways. *Plant physiology* **171**, 1635-1650, doi:10.1104/pp.15.01237 (2016).

29 Torres, M. A., Dangl, J. L. & Jones, J. D. Arabidopsis gp91phox homologues AtrbohD and AtrbohF
are required for accumulation of reactive oxygen intermediates in the plant defense response.
Proc Natl Acad Sci U S A **99**, 517-522, doi:10.1073/pnas.012452499 (2002).
30 Daudi, A. *et al.* The apoplastic oxidative burst peroxidase in Arabidopsis is a major component of
pattern-triggered immunity. *The Plant cell* **24**, 275-287, doi:10.1105/tpc.111.093039 (2012).
31 Li, Y. *et al.* Glucose triggers stomatal closure mediated by basal signaling through HXK1 and
PYR/RCAR receptors in Arabidopsis. *J Exp Bot* **69**, 1471-1484, doi:10.1093/jxb/ery024 (2018).
32 Kadota, Y. *et al.* Direct regulation of the NADPH oxidase RBOHD by the PRR-associated kinase
BIK1 during plant immunity. *Mol Cell* **54**, 43-55, doi:10.1016/j.molcel.2014.02.021 (2014).
33 Li, L. *et al.* The FLS2-associated kinase BIK1 directly phosphorylates the NADPH oxidase RbohD to
control plant immunity. *Cell Host Microbe* **15**, 329-338, doi:10.1016/j.chom.2014.02.009 (2014).
34 Dubiella, U. *et al.* Calcium-dependent protein kinase/NADPH oxidase activation circuit is
required for rapid defense signal propagation. *Proc Natl Acad Sci U S A* **110**, 8744-8749,
doi:10.1073/pnas.1221294110 (2013).
35 Gao, X. *et al.* Bifurcation of Arabidopsis NLR immune signaling via Ca(2)(+)-dependent protein
kinases. *PLoS pathogens* **9**, e1003127, doi:10.1371/journal.ppat.1003127 (2013).
36 Zhang, M. *et al.* The MAP4 Kinase SIK1 Ensures Robust Extracellular ROS Burst and Antibacterial
Immunity in Plants. *Cell Host Microbe* **24**, 379-391 e375, doi:10.1016/j.chom.2018.08.007
(2018).
37 Kadota, Y. *et al.* Quantitative phosphoproteomic analysis reveals common regulatory
mechanisms between effector- and PAMP-triggered immunity in plants. *New Phytol* **221**, 2160-
2175, doi:10.1111/nph.15523 (2019).
38 Asai, T. *et al.* MAP kinase signalling cascade in Arabidopsis innate immunity. *Nature* **415**, 977-
983, doi:10.1038/415977a (2002).
39 Liang, X. *et al.* Arabidopsis heterotrimeric G proteins regulate immunity by directly coupling to
the FLS2 receptor. *Elife* **5**, e13568, doi:10.7554/eLife.13568 (2016).
40 Macho, A. P. & Zipfel, C. Targeting of plant pattern recognition receptor-triggered immunity by
bacterial type-III secretion system effectors. *Curr Opin Microbiol* **23**, 14-22,
doi:10.1016/j.mib.2014.10.009 (2015).
41 Tateda, C. *et al.* Salicylic acid regulates Arabidopsis microbial pattern receptor kinase levels and
signaling. *The Plant cell* **26**, 4171-4187, doi:10.1105/tpc.114.131938 (2014).
42 Sun, T. *et al.* Redundant CAMTA Transcription Factors Negatively Regulate the Biosynthesis of
Salicylic Acid and N-Hydroxypipecolic Acid by Modulating the Expression of SARD1 and CBP60g.
Mol Plant **13**, 144-156, doi:10.1016/j.molp.2019.10.016 (2020).
43 Kim, Y., Gilmour, S. J., Chao, L., Park, S. & Thomashow, M. F. Arabidopsis CAMTA Transcription
Factors Regulate Pipecolic Acid Biosynthesis and Priming of Immunity Genes. *Mol Plant* **13**, 157-
168, doi:10.1016/j.molp.2019.11.001 (2020).
44 Rathinam, V. A. *et al.* TRIF licenses caspase-11-dependent NLRP3 inflammasome activation by
gram-negative bacteria. *Cell* **150**, 606-619, doi:10.1016/j.cell.2012.07.007 (2012).
45 Baroja-Mazo, A. *et al.* The NLRP3 inflammasome is released as a particulate danger signal that
amplifies the inflammatory response. *Nature immunology* **15**, 738-748, doi:10.1038/ni.2919
(2014).
46 Franklin, B. S. *et al.* The adaptor ASC has extracellular and 'prionoid' activities that propagate
inflammation. *Nature immunology* **15**, 727-737, doi:10.1038/ni.2913 (2014).
47 Cao, X. Self-regulation and cross-regulation of pattern-recognition receptor signalling in health
and disease. *Nat Rev Immunol* **16**, 35-50, doi:10.1038/nri.2015.8 (2016).

48   Crabill, E., Joe, A., Block, A., van Rooyen, J. M. & Alfano, J. R. Plant immunity directly or indirectly
403   restricts the injection of type III effectors by the Pseudomonas syringae type III secretion
404   system. *Plant physiology* **154**, 233-244, doi:10.1104/pp.110.159723 (2010).
405   49   Anderson, J. C. *et al.* Decreased abundance of type III secretion system-inducing signals in
406   Arabidopsis mkp1 enhances resistance against Pseudomonas syringae. *Proceedings of the*
407   *National Academy of Sciences* **111**, 6846-6851, doi:10.1073/pnas.1403248111 (2014).
408   50   Nobori, T. *et al.* Transcriptome landscape of a bacterial pathogen under plant immunity.
409   *Proceedings of the National Academy of Sciences* **115**, E3055-E3064,
410   doi:10.1073/pnas.1800529115 (2018).
411   51   Lee, M. H. *et al.* Lignin-based barrier restricts pathogens to the infection site and confers
412   resistance in plants. *Embo j* **38**, e101948, doi:10.15252/embj.2019101948 (2019).
413   52   Yamada, K., Saijo, Y., Nakagami, H. & Takano, Y. Regulation of sugar transporter activity for
414   antibacterial defense in Arabidopsis. *Science* **354**, 1427-1430, doi:10.1126/science.aah5692
415   (2016).

Methods

Plant materials and growth conditions

Arabidopsis thaliana plants used in this study are in Col-0 ecotype background. The *fls2/efr/cerk1*[53], *bak1/bkk1/cerk1*[16], *rps2*[54], *rbohd*[29], *bik1*[55], *cpk4/5/6/11*[35] mutants were reported previously. Plants were grown in potting soil in environmentally-controlled growth chambers, with relative humidity set at 60% and temperature at 22 ℃ with a 12h light/12h dark photoperiod unless stated otherwise. Four- to five-week-old plants were used for all experiments in this study. To generate the *bbc*/*DEX::avrRpt2* and Col-0/*DEX::avrRpt2* transgenic plants, the *avrRpt2* gene was cloned into pBUD-DEX (pBD) vector in the *Xho*I/*Spe*I restriction enzyme sites, and the expression cassette was introduced into Col-0 or *bbc* plants by *Agrobacterium*-mediated transformation.

Bacterial disease and HR assays

The *Pst* DC3000 strains carrying *avrRpt2*, *avrRps4* and *avrPphB* were published previously[56-58]. The D36E*(avrRpt2)* strain was generated by transforming the *avrRpt2* expression plasmid into D36E strain by electroporation. For bacterial inoculation, *Pst* strains were cultured in Luria-Marine (LM) medium overnight at 30 ℃ to a cell density of OD_{600}=0.8-1.0. Bacteria were collected by centrifugation and washed once with sterile water, and adjusted to a cell density of OD_{600}=0.2. For disease assay, bacterial suspension was further diluted to a cell density of OD_{600}=0.001-0.002. Bacteria were infiltrated into leaves with a needleless syringe, and inoculated plants were kept under ambient humidity for about 1h to allow evaporation of excess water from the leaf and then covered with a transparent plastic dome to keep high humidity for 3-4 days. For quantification of bacteria, four leaf discs from two different leaves (after surface sterilization) were taken using a cork borer (7.5mm in diameter) as one biological repeat, and 3-4 repeats were taken for each treatment. Leaf discs were ground and diluted in sterile water, and the extraction solutions were then plated on LM agar plates supplemented with rifampicin (at 50mg/L). Colonies were counted with a stereoscope 24h after incubation at 30 ℃. For HR assay, *Pst* DC3000(*avrRpt2*) suspension was prepared as described above and bacterial suspension at the cell density of OD_{600}=0.2 was syringe-infiltrated into leaves. Plants were then kept under ambient humidity for about 7h before tissue collapse was recorded.

RIN4 cleavage assays

Arabidopsis plant leaves were infiltrated with *Pst* D36E(*avrRpt2*) (at OD$_{600}$=0.1), and samples

were collected at 0, 2, 4, 8h after infiltration by snap-freezing in liquid nitrogen. Three leaves

were collected as one biological repeat. Total proteins were extracted in protein extraction buffer

(50mM Tris-HCl pH 7.5, 150mM NaCl, 5mM EDTA pH 7.5, 1mM DTT, 1% Triton X-100,

1mM Phenylmethylsulfonyl fluoride) supplemented with 1 x plant protease inhibitor cocktail

(Complete EDTA-free, Roche). Cell lysates were centrifuged at 12,000 x *g* for 15min at 4°C, and

the pellet was discarded. Protein concentration of the supernatant ("total protein extract") was

determined by Bradford protein assay kit (Bio-Rad). An equal amount of total protein was

loaded on 12% SDS acrylamide gels (Bio-Rad) for SDS-PAGE. RIN4 protein was detected by

anti-RIN4 antibody at a dilution of 1:1000[59]. Total proteins were stained by Coomassie Brilliant

Blue (CBB) to show equal loading.

MAPK kinase activity assay

Four-week-old plant leaves were infiltrated with *Pst* D36E(*avrRpt2*) (at OD$_{600}$=0.1), and leaves

were collected at different time points by snap-freezing in liquid nitrogen. Proteins were

extracted in protein extraction buffer (50mM Tris-HCl pH 7.5, 150mM NaCl, 5mM EDTA pH

7.5, 1mM DTT, 1% Triton X-100, 1mM Phenylmethylsulfonyl fluoride) supplemented with 1 x

plant protease inhibitor cocktail (Complete EDTA-free, Roche) and 1 x phosphatase inhibitor

cocktail (PhosSTOP, Roche). Total protein concentration was determined with Bradford protein

assay kit (Bio-Rad). An equal amount of protein was loaded onto 12% SDS-PAGE gel for

western blot. Phosphorylated MPK3 and MPK6 proteins were detected by anti-Phospho-p44/42

antibody (Cell Signaling Technology).

Protein extraction and immunoblotting for PTI signaling components

Four-week-old plant leaves were infiltrated with sterile water (mock) or different *Pst* strains at

OD$_{600}$=0.02, and samples were collected at 0.5, 3, 6, 8h after infiltration. Three to four leaves

from different plants were collected as one sample. Protein was extracted using Plasma

Membrane Protein Isolation Kit (Invent) according to the manufacturer's protocol. Concentration

of the cytosolic protein was determined with Bradford protein assay kit (Bio-Rad). An equal

amount of protein was loaded onto SDS-PAGE gel for western blot. Different PTI components

were detected by following antibodies with indicated dilution: anti-RBOHD (Agrisera), 1:1000;

anti-BAK1 (Agrisera), 1:5000; anti-BIK1 (Agrisera), 1:3000; anti-MPK3 (Sigma-Aldrich),

1:2500; anti-MPK6 (Sigma-Aldrich), 1:5000.

Protoplast transformation and detection of RBOHD phosphorylation

Protoplasts were prepared from Col-0/*DEX::avrRpt2* and *bbc/DEX::avrRpt2* plants (4-5 weeks

old; grown under 10h light/14h dark photoperiod) and transfected with FLAG-RBOHD plasmid.

After overnight incubation to allow protein accumulation, protoplasts were treated with 100nM

flg22, 5μM DEX or 100nM flg22+5μM DEX and incubated for 2.5h. Total protein was extracted

with protein extraction buffer (50 mM HEPES [pH 7.5], 150 mM KCl, 1 mM EDTA, 0.5%

Trition-X100, 1 mM DTT, protease inhibitor cocktail), and then incubated with 50μL anti-FLAG

M2 agarose beads (Sigma-Aldrich) for 2 h at 4 ℃. The bound protein was eluted with 50μL of

0.5mg/mL 3xFLAG peptide for 30 min. RBOHD phosphorylation was detected by

immunoblotting with RBOHD-pS343/347 antibody published previously[33].

ROS detection Assays

ROS measurement with luminol-based approach was performed as previously described with

minor modification[32]. Briefly, leaf discs of four-week-old *Arabidopsis* plants were harvested

using a cork borer (5.5mm in diameter) and floated on 200μL sterile water in a 96-well plate, and

then incubated overnight at room temperature under continuous light. On the next day, water was

replaced with a solution containing 30mg/L (w/v) luminol (Sigma-Aldrich) and 20mg/L (w/v)

peroxidase from horseradish (Sigma-Aldrich) with 100nM flg22 only, 5μM DEX only or 100nM

flg22+5μM DEX. The luminescence was detected for 5-6h with a signal integration time of 2min

using Varioskan Flash plate reader (Thermo Fisher Scientific). For determining the effects of

chemical inhibitors, 10μM diphenyleneiodonium (DPI; Sigma-Aldrich), 15μM

salicylhydroxamic acid (SHAM; Sigma-Aldrich) or 1μM sodium azide was added to the

elicitation solution at indicated time points and luminescence was recorded as described above.

For detection of ROS production by 2',7'-Dichlorofluorescein diacetate (H_2DCFDA) under

confocal microscopy, plants were infiltrated with *Pst* D36E (OD_{600}=0.02) or D36E(*avrRpt2*)

(OD_{600}=0.02), air-dried and put back into the plant growth room. ROS was detected at 4-5h post

infiltration. Ten μM of H_2DCFDA solution was infiltrated into the leaf and fluorescence signal

was detected 10 min later. Images were captured using a Leica SP8 microscope with a 488nm

excitation and 501-550nm emission, and chlorophyll auto-fluorescence was detected at 640-

735nm.

RNA extraction and qRT-PCR analysis of gene expression

To analyze gene expression levels, four-week-old *Arabidopsis* plant leaves were infiltrated with

sterile water (mock) or different *Pst* strains at OD_{600}=0.04, and then harvested at indicated time

points. Three leaves from different plants were collected as one biological replicate and 4

replicates were collected for each treatment. Samples were frozen and ground in liquid nitrogen.

Total mRNA was extracted using Trizol reagent (Invitrogen) according to the manufacturer's

protocol. One μg of RNA was used for reverse transcription using the ReverTra Ace® qPCR RT

Master Mix with gDNA remover (TOYOBO). Real-time qPCR analysis was carried out with the

SYBR®Green Realtime PCR Master Mix (TOYOBO) on a CFX real-time machine (Bio-Rad).

Two technical repeats were performed for each sample. The plant *U-box* gene was used as

reference gene for normalization. Primer sequences for qPCR are listed in Supplementary Table

2.

cDNA library generation and RNAseq

For RNAseq experiments, bacterial inoculation and sample collection were performed as

described above. Two leaves from different plants were harvested as one replicate, and four

biological replicates were collected for each treatment/time point. Total mRNA was extracted

using Trizol reagent (Invitrogen). Total RNA was then treated with DNase I (Invitrogen) to

remove DNA and purified RNA was recovered with RNeasy® MinElute™ Cleanup kit

(QIAGEN) according to the manufacturer's instructions. Library construction and RNA

sequencing were performed by Novogene company. Briefly, RNA purity and integrity was

examined using the NanoPhotometer® spectrophotometer (IMPLEN) and the RNA Nano 6000

Assay Kit of the Bioanalyzer 2100 system (Agilent Technologies). RNA concentration was

measured with Qubit® RNA Assay Kit in Qubit® 2.0 Flurometer (Life Technologies). One μg

RNA per sample was used as input material for library preparation and sequencing. Sequencing

libraries were generated using NEBNext® Ultra™ RNA Library Prep Kit for Illumina® (NEB),

following the manufacturer's recommendations and sequenced on Illumina Hiseq platform and

150 bp paired-end reads were generated.

Data analysis of RNA-seq

Clean raw data were obtained by removing reads containing adapter sequences or ploy-N and low-quality reads and were then mapped to the Arabidopsis genome (TAIR10). Gene expression levels were calculated using the TPM method (Transcripts per Kb of exon model per Million mapped reads). Differential expression analysis was performed using the DEGSeq R package (1.18.0). The resulting P-values were adjusted using the Benjamini and Hochberg's approach for controlling the false discovery rate. Genes with an q-value < 0.05 and $\log_2$(Fold change) > 1 found by DESeq were assigned as differentially expressed.

Statistical analysis

All statistical analyses were performed by one-way or two-way analysis of variance (ANOVA) with GraphPad software or two sided student's t-test with Office Excel software. Each experiment was repeated at least three times and data were represented as the mean $\pm$ standard error of mean (s.e.m.) or standard deviation (s.d.) as indicated.

Data availability

All data is available in the main text or the supplementary materials.

Gimenez-Ibanez, S., Ntoukakis, V. & Rathjen, J. P. The LysM receptor kinase CERK1 mediates bacterial perception in Arabidopsis. *Plant signaling & behavior* **4**, 539-541, doi:10.4161/psb.4.6.8697 (2009).

Mindrinos, M., Katagiri, F., Yu, G. L. & Ausubel, F. M. The A. thaliana disease resistance gene RPS2 encodes a protein containing a nucleotide-binding site and leucine-rich repeats. *Cell* **78**, 1089-1099, doi:10.1016/0092-8674(94)90282-8 (1994).

Veronese, P. *et al.* The membrane-anchored BOTRYTIS-INDUCED KINASE1 plays distinct roles in Arabidopsis resistance to necrotrophic and biotrophic pathogens. *The Plant cell* **18**, 257-273, doi:10.1105/tpc.105.035576 (2006).

Mudgett, M. B. & Staskawicz, B. J. Characterization of the Pseudomonas syringae pv. tomato AvrRpt2 protein: demonstration of secretion and processing during bacterial pathogenesis. *Mol Microbiol* **32**, 927-941, doi:10.1046/j.1365-2958.1999.01403.x (1999).

Hinsch, M. & Staskawicz, B. Identification of a new Arabidopsis disease resistance locus, RPs4, and cloning of the corresponding avirulence gene, avrRps4, from Pseudomonas syringae pv. pisi. *Mol Plant Microbe Interact* **9**, 55-61, doi:10.1094/mpmi-9-0055 (1996).

Aarts, N. *et al.* Different requirements for EDS1 and NDR1 by disease resistance genes define at least two R gene-mediated signaling pathways in Arabidopsis. *Proc Natl Acad Sci U S A* **95**, 10306-10311, doi:10.1073/pnas.95.17.10306 (1998).

Lee, D., Bourdais, G., Yu, G., Robatzek, S. & Coaker, G. Phosphorylation of the Plant Immune Regulator RPM1-INTERACTING PROTEIN4 Enhances Plant Plasma Membrane H(+)-ATPase

572		Activity and Inhibits Flagellin-Triggered Immune Responses in Arabidopsis. *The Plant cell* **27**,
573		2042-2056, doi:10.1105/tpc.114.132308 (2015).
574	60	Pajerowska-Mukhtar, K. M. *et al.* The HSF-like transcription factor TBF1 is a major molecular
575		switch for plant growth-to-defense transition. *Curr Biol* **22**, 103-112,
576		doi:10.1016/j.cub.2011.12.015 (2012).
577	61	Hickman, R. *et al.* Architecture and Dynamics of the Jasmonic Acid Gene Regulatory Network.
578		*The Plant cell* **29**, 2086-2105, doi:10.1105/tpc.16.00958 (2017).
579	62	Nemhauser, J. L., Hong, F. & Chory, J. Different plant hormones regulate similar processes
580		through largely nonoverlapping transcriptional responses. *Cell* **126**, 467-475,
581		doi:10.1016/j.cell.2006.05.050 (2006).
582	63	Hartmann, M. *et al.* Flavin Monooxygenase-Generated N-Hydroxypipecolic Acid Is a Critical
583		Element of Plant Systemic Immunity. *Cell* **173**, 456-469 e416, doi:10.1016/j.cell.2018.02.049
584		(2018).

Acknowledgments:

We would like to thank Xin lab members for helpful discussions. We thank the Greenhouse and

Confocal Microscopy Imaging facilities at Institute of Plant Physiology and Ecology, Chinese

Academy of Sciences for plant growth and technical support. Dr. Gitta Coaker from University

of California, Davis, USA kindly provided RIN4 antibody. We thank Bruno Pok Man Ngou and

Pingtao Ding from Jonathan Jones' lab at the Sainsbury Laboratory, UK, for insightful

discussions during manuscript preparation. This research was supported by Chinese Academy of

Sciences, Center for Excellence in Molecular Plant Sciences/Institute of Plant Physiology and

Ecology, National key Laboratory of Molecular Plant Genetics and Chinese Academy of

Sciences Strategic Priority Research Program (Type-B; Project number: XDB27040211). Guozhi

Bi was supported by the Youth Program of National Natural Science Foundation of China

(NSFC) (Project number: 31900222).

Author contributions: With initial observation of PRR dependency for ETI resistance made by

X-F. X. while at Michigan State University, supported by the US National Institute of General

Medical Sciences (GM109928), M. Y. and X-F. X. designed all subsequent experiments at

Institute of Plant Physiology and Ecology, Chinese Academy of Sciences; M. Y., Z. J., G. B., K.

N. and M. L. performed all the experiments described; M. Y. and X-F. X. wrote the paper and S.

Y. H. and J-M. Z. edited the paper.

Competing interests: The authors declare no competing interests.

Materials & Correspondence: Correspondence and material requests should be addressed to

xinxf@sippe.ac.cn.

Supplementary Table 2. Primers used in this study

Primer name	Sequences (5'-3')	Purpose
FRK1 qRT-FP	ATCTTCGCTTGGAGCTTCTC	**For RT-qPCR**
FRK1 qRT-RP	TGCAGCGCAAGGACTAGAG	
WRKY29 qRT-FP	CTCCATACCCAAGGAGTTATTACAG	
WRKY29 qRT-RP	CGGGTTGGTAGTTCATGATTG	
U-box qRT-FP	TGCGCTGCCAGATAATACACTATT	
U-box qRT-RP	TGCTGCCCAACATCAGGTT	
RPS2 qp-FP	GTGAGTAAATGTGGAGGATTGC	
RPS2 qp-RP	GTAGCTGAATTTCAAAAGGGCA	
RbohD qRT-FP	GATCAAGGTGGCTGTTTACCC	
RbohD qRT-RP	TCGGCAGTTCACCAACATGA	
BAK1 qRT-FP	AGGTGTTCTCTTGGAGACTAGGA	
BAK1 qRT-RP	AGAGATCCAGAACTTGTAGCGT	
BIK1 qRT-FP	TTCTTCACAGCGATCCCGTC	
BIK1 qRT-RP	TTGCGTTGTAGTCCGCATCA	
XLG2 qRT-FP	AGTCTTCAGGCAATGTGGGAG	
XLG2 qRT-RP	CTCACGGCTTGATGGTGGAA	
MKK4 qRT-FP	CTTATCTCCATAGCCGTCACAT	
MKK4 qRT-RP	GAGCCAAGATCCTACTAACTCC	
MKK5 qRT-FP	CGCCGCTAAAAGCTTATCC	
MKK5 qRT-RP	CGTCTCACGGTATCTTCGTG	
MAPK3 qRT-FP	CCAAGAAGCCATAGCACTCA	
MAPK3 qRT-RP	AGCCATTCGGATGGTTATTG	
AvrRpt2 qRT-FP	CTTTTCACGATCCCCGACAGGG	
AvrRpt2 qRT-RP	GCGGTAGAGCATTGCGTGTGG	
pBD-AvrRpt2-F	CCGCTCGAGATGAAAATTGCTCCAGTTGC	**For Cloning**
pBD-AvrRpt2-R	CGGACTAGTTTAGCGGTAGAGCATTGCGT	

Fig. 1 | See next page for caption.

Fig.1| PTI-associated PRR/co-receptors are required for ETI responses and resistance. a, Two independent triple PRR/co-receptor mutant plants were hyper-susceptible to *Pst* DC3000 (*avrRpt2*) infection. Bacteria were infiltrated into Arabidopsis leaves at OD_{600}=0.002 and bacterial populations inside leaves were determined 3 days post infection. **b,** HR cell death (indicated by leaf collapse in Col-0) was compromised in PRR/co-receptor mutants. *Pst* DC3000 (*avrRpt2*) bacteria were infiltrated into Arabidopsis leaves at OD_{600}=0.2 and pictures were taken ~7 h after infiltration. Numbers of dead and total infiltrated leaves were counted. **c,** The triple PRR/co-receptor mutant plants were hyper-susceptible to D36E(*avrRpt2*) infection. Bacteria were infiltrated into Arabidopsis leaves at OD_{600}=0.002-0.004 and bacterial populations inside leaves were determined 3-4 days post infection. Different letters in **a,** and **c,** indicate statistically significant differences in bacterial population, as determined by two-way ANOVA (mean ± s.d.; $n \geq 3$; $p < 0.05$). **d, e,** RIN4 cleavage **d** and MPK3/6 phosphorylation **e** in Col-0 and the PRR/co-receptor mutants after D36E(*avrRpt2*) inoculation. CBB, Coomassie Brilliant Blue staining. An equal amount of total protein was loaded in each lane. **f, g,** ROS burst detected by luminol-HRP approach in the DEX-inducible *avrRpt2* transgenic plant after treatment of 100nM flg22, 5μM DEX, or both. RLUs, relative luminescence units. Total photon counts are calculated from **f** ($n \geq 27$). Different letters indicate statistically significant differences as analyzed by one-way ANOVA ($p < 0.05$). **h, i,** ROS burst in Col-0/*DEX::avrRpt2* and *bbc/DEX::avrRpt2* plants after treatment of 5μM DEX, or 100nM flg22+5μM DEX. Total photon counts are calculated from **h** ($n \geq 5$). Different letters indicate statistically significant differences as analyzed by two-way ANOVA ($p < 0.05$). Box plots: centre line, median; box limits, lower and upper quartiles; whiskers, highest and lowest data points. Dots indicate individual data points. All experiments were repeated at least three times with similar trends.

Fig. 2 | See next page for caption.

Fig.2| AvrRpt2-triggered ROS is mediated by RBOHD and requires PRR/co-receptors for activation. a, ROS burst detected with fluorescent dye H_2DCFDA in Col-0, *bbc*, *rps2* and *rbohd* leaves 5h after infiltration of D36E(*avrRpt2*) or in Col-0 leaves 5h after infiltration of D36E strain. White arrows indicate the apoplast space in the leaf. Scale bars = 25μm. **b,** ETI-associated ROS burst is inhibited by DPI, an NADPH oxidase inhibitor. ROS was detected in Col-0/*DEX::avrRpt2* plants after treatment of 100nM flg22 and 5μM DEX. Chemical inhibitors (DPI, SHAM or NaN_3) were added after the first ROS burst (about 40min after addition of flg22 and DEX). Data are displayed as mean ± s.e.m. ($n \geq 6$). **c,** ROS burst detected with fluorescent dye H_2DCFDA in Col-0 and *rbohd* leaves 5h after infiltration of D36E(*avrRpt2*). Scale bars = 25μm. **d,** The *rbohd* mutant plant is compromised in ETI resistance against *Pst* DC3000 (*avrRpt2*). Bacteria were infiltrated into Arabidopsis leaves at OD_{600}=0.001 and bacterial populations inside leaves were determined 2 days post infection. **, student's *t*-test, two-tailed, $p < 0.001$. Data are displayed as mean ± s.d. ($n = 3$). **e,** RBOHD transcript levels in Col-0 and *bbc* plants 3h after inoculation of bacterial strains indicated. Data are displayed by mean ± s.e.m. ($n \geq 3$). Different letters indicate statistically significant differences, as analyzed by two-way ANOVA ($p < 0.05$). **f,** RBOHD protein levels in Col-0 and *bbc* plants at different time points after inoculation of bacterial strains indicated. The numbers indicate band intensity relative to that of Ponceau S, quantified by ImageJ, and red indicates strong induction. **g,** Phosphorylation of RBOHD protein at S343/S347 sites requires PRR/co-receptors. FLAG-RBOHD was transformed into protoplasts prepared from Col-0/*DEX::avrRpt2* and *bbc*/*DEX::avrRpt2* plants. Protoplasts were treated with elicitors (F, 100nM flg22; D, 5μM DEX; FD, 100nM flg22+5μM DEX) and harvested 2.5h later for FLAG-RBOHD immunoprecipitation. Phosphorylated and total RBOHD proteins were detected by RBOHD pS343/347-specific antibody and FLAG antibody, respectively. All experiments were repeated at least three times with similar trends.

Fig.3| ETI upregulates key components of the PTI pathway. a, A diagram showing the RNAseq design in this study. **b,** A heat-map of the expression pattern of D36E/PTI-responsive genes in the RNAseq experiment. **c,** A heat map of the expression pattern of PTI pathway genes, showing restoration and up-regulation of expression of major PTI components during AvrRpt2-triggered ETI. **d,** qRT-PCR results of representative PTI pathway genes. Col-0 and *bbc* plants were infiltrated with different strains indicated, and leaves were harvested 3h post infiltration for transcript analysis (mean ± s.e.m; $n \geq 3$). Different letters indicate statistically significant differences as analyzed by two-way ANOVA ($p < 0.05$). **e,** Protein levels of BAK1, BIK1, MPK3 and MPK6 in Col-0 plants at different time points after inoculation of bacterial strains indicated. BAK1 is detected in the immunoblot of total membrane fraction and other proteins are detected in the immunoblot of total protein extracts. Equal amounts of total proteins were loaded into gel lanes. MPK6 protein is not induced by ETI and serves as an internal control. The numbers indicate band intensity relative to that of Ponceau S, quantified by ImageJ, and red indicates strong induction. All experiments were repeated at least three times with similar trends.

Fig.4| A model depicting findings from this study showing PTI as a key component of ETI.
Grey color indicates mutated (i.e. FLS2 and BAK1) or inactive (i.e. RBOHD and BIK1) proteins. In wild-type plant, PTI is integrated into ETI in that RPS2 activation leads to protein accumulation of PTI components such as RBOHD and PRR/co-receptors are required for "activating" it by phosphorylation. In the absence of PRR/co-receptors (left panel), although NLR activation strongly induces protein accumulation of PTI components, many of these components, such as BIK1 and RBOHD, are inactive (shown by the grey arrows), leading to compromised ETI resistance.

Extended Data Fig.1| PRR/co-receptors are required for ETI elicited by different *P. syringae* avirulent effectors. AvrPphB- and AvrRps4-mediated ETI are also compromised in *fec* and *bbc* mutants. Plants were infiltrated with different strains at OD_{600}=0.002. Bacterial populations were determined 3 days post inoculation. Different letters indicate statistically significant differences in bacterial populations as determined by two-way ANOVA. (mean ± s.d.; $n \geq 3$; $p < 0.05$). Experiments were repeated at least three times with similar trends.

Extended Data Fig.2| The transcript levels of *RPS2* are not altered in the *fec* and *bbc* mutant plants. *RPS2* transcript levels in the *fec* and *bbc* mutant plants were similar to those in Col-0 plants after inoculation of bacterial strains indicated. Different letters indicate statistically significant differences as analyzed by two-way ANOVA (mean ± s.e.m; $n = 3$; $p < 0.05$). Experiments were repeated at least three times with similar trends.

Extended Data Fig.3| Characterization of different lines of *bbc/DEX::avrRpt2* plants.
Expression levels of the *avrRpt2* transgene in different transgenic lines 2h after infiltration with
5μM DEX. Different letters indicate statistically significant differences as analyzed by one-way
ANOVA (mean ± s.e.m.; $n \geq 3$; $p < 0.05$). Experiments were repeated at least three times with
similar trends.

Extended Data Fig.4| AvrRpt2-triggered ETI-ROS depends on NADPH oxidase. a-c, ROS production in Col-0/*DEX::avrRpt2* L1 plants was inhibited by NADPH oxidase inhibitor DPI. Leaf discs were treated with 100nM flg22 and 5μM DEX. DPI, SHAM and NaN$_3$ were added at the beginning of measurement (mean ± s.e.m.; $n \geq 6$). **b-c,** Total photon counts are calculated from **a** at the PTI phase (0-30min) or ETI phase (60-200min). Different letters indicate statistically significant difference as analyzed by one-way ANOVA ($p < 0.05$). Box plots: centre line, median; box limits, lower and upper quartiles; dots, individual data points; whiskers, highest and lowest data points. Experiments were repeated at least three times with similar trends.

Extended Data Fig.5| The *rbohd* mutant plant is compromised in ETI resistance against *Pst* DC3000(*avrRpt2*). a, Appearance of the 5 week-old *rbohd* mutant plants before bacteria inoculation. **b,** Disease symptom of Col-0 and *rbohd* mutant plant 2 days after *Pst* DC3000 and *Pst* DC3000 (*avrRpt2*) infiltration. Experiments were repeated at least three times with similar trends.

D36E (*avrRpt2*)

Extended Data Fig.6| The AvrRpt2 ETI-associated ROS burst is partially mediated by BIK1.
ROS was detected in the *bik1* and *cpk4/5/6/11* mutant plants by H_2DCFDA dye 4.5 h after D36E
(*avrRpt2*) inoculation. Scale bars = 25 μm. Experiments were repeated at least three times with
similar trends.

Extended Data Fig.7| Transcriptomic analysis of RNAseq experiments. a, Bacterial population in Arabidopsis leaves at 3h or 6h post infiltration. Data are displayed by mean ± s.d. (n = 3). **b,** A Venn diagram showing numbers of differentially expressed genes (DEGs) 3h after D36E or D36E(*avrRpt2*) infection in Col-0 plants. **c-e,** Heat-maps of SA- **(c**; genes extracted from Karolina et al., 2012[60]**),** jasmonate- **(d;** genes extracted from Hickman et al., 2017[61]**)** and ethylene-**(e;** genes extracted from Nemhauser et al.,2006[62]**)** responsive genes.

Extended Data Fig.8| Genome-wide expression profiling reveals interesting features of gene regulation attributed to PTI and ETI. a, GO enrichment analysis of the 209 down-regulated genes in the *bbc* plant compared to Col-0 plant after D36E(*avrRpt2*) infection (among the total 272 DEGs). **b**, Heat map of the 272 DEGs in the *bbc* plant compared to Col-0 plant after D36E (*avrRpt2*) infection, with the canonical PTI pathway genes highlighted in red. **c,** qRT-PCR of *FRK1* and *WRKY29* expression level in Col-0 and *bbc* plants 3h after infiltration with different strains or Mock. (mean ± s.e.m.; $n = 3$; statistical analysis by two-way ANOVA; $p < 0.05$; different letters indicate statistically significant difference). Experiments were repeated at least three times with similar trends.

	Mock		D36E		D36E (*avrRpt2*)		
	Col-0	*bbc*	Col-0	*bbc*	Col-0	*bbc*	
	0.00	0.00	0.00	0.03	0.04	0.00	*PBL25*
	0.00	0.00	0.00	0.00	0.02	0.00	*PBL43*
	16.65	10.85	18.61	15.82	14.75	18.17	*PBL37*
	62.48	45.94	122.15	66.82	146.22	124.60	*PBL17*
	21.32	19.60	37.49	25.41	32.57	33.34	*PBL7*
	45.83	27.01	234.57	118.99	240.40	298.86	*PBL39*
	24.92	17.81	85.39	49.95	57.73	50.70	*PBL1*
	63.34	37.39	93.03	62.95	77.15	73.88	*PBL3*
	1.73	0.84	3.36	2.30	24.62	27.50	*PBL33*
	1.86	1.63	2.19	1.86	3.10	3.38	*PBL6*
	21.34	14.95	31.45	24.79	94.39	131.63	*PBL19*
	0.01	0.00	0.00	0.00	0.20	0.24	*PBL22*
	0.77	0.62	0.69	0.88	2.67	2.64	*PBL18*
	0.00	0.00	0.00	0.00	0.05	0.04	*PBL42*
	0.08	0.09	0.16	0.08	0.41	0.21	*PBL12*
	0.04	0.05	0.06	0.01	0.15	0.13	*PBL29*
	0.01	0.00	0.01	0.00	0.02	0.07	*PBL4*
	0.35	0.12	1.32	1.50	1.80	2.96	*PBL20*
	0.92	0.97	1.55	1.11	1.87	2.23	*PBL38*
	0.87	0.41	3.43	1.65	7.81	6.93	*PBL13*
	33.42	15.53	172.10	56.65	476.20	388.81	*PBL14*
	20.84	12.05	48.68	28.09	128.08	136.06	*PBL30*
	36.61	23.03	90.19	55.27	180.79	195.64	*PBL32*
	55.05	47.22	100.52	76.31	125.87	146.74	*PBL31*
	80.14	57.92	163.05	94.83	214.99	250.21	*PBL11*
	24.42	19.27	68.43	33.81	101.72	96.92	*PBL40*
	15.97	12.44	71.13	24.89	152.50	173.87	*BIK1*
	41.99	28.98	122.92	56.52	230.00	262.82	*PBL2*
	7.11	5.59	6.06	7.12	5.21	6.09	*PBL16*
	1.06	0.67	0.46	0.95	0.33	0.20	*PBL28*
	63.05	68.76	31.68	56.66	12.03	16.83	*PBL8*
	39.85	47.90	10.26	28.18	1.78	2.17	*PBL21*
	23.94	31.12	44.00	29.43	12.68	7.73	*PBL10*
	8.01	6.59	7.66	8.45	5.11	5.56	*PBL5*
	9.43	10.05	9.46	10.62	4.30	4.37	*PBL35*
	14.67	13.73	14.76	15.18	9.51	10.44	*PBL34*
	44.90	44.22	44.65	41.83	20.49	24.77	*PBL36*
	0.00	0.04	0.12	0.10	0.08	0.03	*PBL24*
	31.45	29.33	38.61	34.14	31.12	29.48	*PBS1*
	0.15	0.08	0.32	0.23	0.05	0.05	*PBL23*
	17.59	20.39	31.28	31.20	17.74	19.84	*PBL9*
	0.28	0.38	0.51	0.40	0.20	0.34	*PBL41*
	0.00	0.04	0.03	0.03	0.00	0.05	*PBL26*
	18.37	20.30	20.84	18.16	19.57	20.02	*PBL27*

Extended Data Fig.9| Heat map of *BIK1/PBL* family gene expression in the RNAseq experiment. Numerical values indicate expression level calculated by TPM (Transcripts per Kb of exon model per Million). Genes labeled in red show significant up-regulation after D36E(*avrRpt2*) inoculation, compared to mock and D36E inoculation, in Col-0 and *bbc* plants. Arrows indicate *BIK1* and *PBL1* genes.

Extended Data Fig.10| Up-regulation of key PTI component genes by AvrRpt2-triggered ETI seems to be independent of SA/NHP. a, qRT-PCR analysis of *BIK1, XLG2, MKK4, MKK5* and *MPK3* expression levels in Col-0 and *sid2* plants 3h after infiltration with D36E or D36E(*avrRpt2*). Different letters indicate statistically significant differences, as analyzed by two-way ANOVA (mean ± s.e.m.; n ≥ 3; p < 0.05). Experiments were repeated at least three times with similar trends. **b,** Heat-maps of NHP-responsive genes (extracted from Hartmann et al., 2018[63], defined by genes that are responsive to pipecolic acid and depend on FMO1 for expression) in the Col-0 and *bbc* plants in our RNAseq experiment.