1

Temporal patterns in electrical nerve stimulation: burst gap code shapes tactile frequency perception

Kevin K. W. Ng[1,2]*, Christoffer Olausson[2,3], Richard M. Vickery[1,2], Ingvars Birznieks[1,2]

[1] School of Medical Sciences, UNSW Sydney, Sydney, Australia

[2] Neuroscience Research Australia, Sydney, Australia

[3] Faculty of Medicine and Health Sciences, Linköping University, Linköping, Sweden

* Corresponding author

E-mail: kevin.ng@unsw.edu.au

Abstract

We have described a novel temporal encoding mechanism in the somatosensory system, where

tactile pulses grouped into periodic bursts create a perceived frequency based on the duration of the

silent gap between bursts, rather than the mean rate or the periodicity. This coding strategy may

offer new opportunities for transmitting information to the brain using various sensory neural

prostheses and brain-machine interfaces. However, it was not known whether the same coding

mechanisms apply when electrical, rather than mechanical, stimulation is used. Here, we

demonstrate that the predictions of the burst gap coding model for frequency perception apply to

burst stimuli delivered with electrical pulses, re-emphasising the importance of the temporal

structure of spike patterns in neural processing and perception of tactile stimuli. Reciprocally, the

electrical stimulation data confirm that the results observed with mechanical stimulation do indeed

depend on neural processing mechanisms in the central nervous system and are not due to

mechanical effects in the skin or afferent population activation patterns. Our results suggest that

spike intervals contribute to perceived frequency with perceptual weights that depend on their

duration, and that a generalised frequency processing system would explain the constancy of

vibrotactile frequency perception across different receptor types.

Introduction

Temporal features of neural spike activity play a major role in encoding tactile information [1-4]. In a

previous study, we used pulsatile mechanical stimulation to investigate the neural basis of frequency

perception in the flutter range. When we grouped pulsatile stimulation patterns into periodic bursts,

we found that perceived frequency was determined by the duration of the silent gap between bursts

of spikes in peripheral afferents, irrespective of the number of spikes within a burst or the burst rate

[5]. This stands in contrast to the mean spike rate or periodicity code that are often proposed to be

neural codes for frequency perception in higher neural centres [6-9]. Effective deployment of this

burst coding may be useful in restoring the sense of touch in amputees using a prosthesis through

electrical stimulation [10-12] or in the treatment of movement disorders through deep brain

stimulation as several studies have established [13-16].

The discovery of the burst gap code for perceived frequency depended on brief mechanical pulses

that only evoke a single spike in responding afferents, as had been verified using microneurography,

allowing us to precisely control the spiking pattern of responding tactile afferents [5, 17, 18]. On the

other hand, electrical stimulation activates peripheral axons in a manner different than that of

mechanical stimulation [19], by bypassing the specialised mechanoreceptors in the skin and directly

stimulating the tactile afferent axons [20, 21]. Electrocutaneous stimulation non-selectively activates

the different types of tactile afferents (associated with different skin receptors) [22, 23], which

contrasts with mechanical stimulation where the type of afferent recruited varies depending on the

sensitivity of the fibre type to stimulus characteristics such as the vibration frequency of a

mechanical sinusoid [24, 25].

A single tap on a finger generates a complex propagating wave on the fingertip skin [26, 27] that will

evoke complex and potentially variable population spike patterns due to sporadically responding

afferents [28]. In particular, such a stimulus would reliably activate afferents whose receptive fields

are directly under the probe, but this activation would become stochastic at some distance from the

locus of stimulation [25, 29]. This raises the possibility that the burst gap code that we observed with

mechanical stimulation is in some way dependent on these spatial activation patterns and/or

predominant activation of certain afferent types. As an alternative, electrical stimulation is a robust

method to reliably activate axons, and a simple technique to implement in brain-computer

interfaces or haptic displays [30]. Thus, in this study we used electrical stimuli to directly evoke spike

trains in tactile afferent fibres while avoiding complex mechanical skin phenomena. We aimed to

establish that the burst gap encoding for frequency perception is genuinely caused by neural

processing mechanisms in the central nervous system and not the result of a complex population

effect, and to demonstrate the efficacy of this technique for use in interfaces with the nervous

system.

Materials and methods

Subjects

Two separate experiments were conducted. Twelve subjects (aged 21-26, 5 females) participated in

Experiment 1. Eight subjects (aged 21-25, 5 females) participated in Experiment 2. Three of the

subjects participated in both experiments. All subjects were healthy and without any known history

of altered tactile function. The experimental protocols were approved by the Human Research Ethics

Committee (HC16245) of UNSW Sydney and written consent was obtained from subjects before

experimentation began.

Apparatus

Stimulus patterns were generated via a CED Power1401 mk II data acquisition system using Spike2

(Cambridge Electronic Design, Cambridge, UK) and MATLAB (Mathworks, Natick, USA) software.

These patterns triggered constant current electrical pulses from a DS7AH stimulator (Digitimer,

Welwyn Garden City, UK). The electrical stimuli were delivered to the right index finger of subjects

using Kendall 200 Series foam electrodes (Covidien, Mansfield, MA, USA), with one electrode placed

on the distal interphalangeal joint and another on the proximal phalanx, to stimulate the digital

nerve. A button box was used by subjects to indicate their responses to the psychophysical task.

Compound action potentials were recorded using a PowerLab data acquisition system and LabChart

software (ADInstruments, Sydney, Australia).

Stimulation patterns

Experiment 1

The stimulus timing patterns used were based on that of our previous study [5]. In our first experiment, the aim was to determine whether burst gap coding would best describe the perception of frequency in subjects. The test stimuli consisted of one of four trains with periodic bursts of 2-4 pulses (Fig 1A), such as would be generated in the afferent response to high-amplitude vibration [25, 31]. The individual pulses were spaced either 4.35 ms (stimuli a-c) or 8.7 ms (stimulus d) apart within the burst. The period between repeating bursts was fixed at 43.5 ms, resulting in a burst rate of 23 Hz. However, the mean pulse rate varied considerably between the four stimuli, ranging from 46 to 92 Hz.

Experiment 2

In our second experiment, we measured the critical value of the inter-pulse interval within a burst that determined how the pulses contribute to perceived frequency. The test stimuli were one of 13 trains of a 2-pulse burst (doublet) with a short interval (t_s) and a long interval (t_L) (Fig 1B). The short inter-pulse interval (t_s) ranged from 4.35 ms to 56.55 ms, increasing in steps of 4.35 ms. The long interval was fixed at 87 ms. The burst gap code predicts that when two pulses are close enough in time, they will be treated as a burst and accordingly the perceived frequency will be determined by the burst gap (approximately 11.5 Hz in this experiment).

Fig 1. Schematic of stimulus patterns and experimental protocol, each red vertical line represents an electrical pulse. (A) In Experiment 1, test stimuli have the same periodicity (burst rate), but different overall pulse rates. (B) In Experiment 2, test stimuli consisted of a short interval doublet (t_s), followed by an 87 ms long inter-pulse interval (t_L). The short inter-pulse interval was varied in

114 steps of 4.35 ms, from 4.35 ms up to 56.55 ms. (C) Two interval forced choice procedure used to

115 determine perceived frequency of test stimuli.

Electrical stimulation parameters

117 Electrical stimuli were 0.1 ms wide square pulses. Prior to the start of each experiment, the electrical

118 current level was optimised for the subject so that the electrical pulses were perceived clearly as a

119 series of taps without discomfort. Current levels used ranged from 6.5 to 11.5 mA across subjects

120 and were kept at a fixed level within a session.

Verifying the stable activation of a population of afferents

Sensory nerve action potentials (SNAPs) were obtained in two subjects by stimulating the sensory

nerves distally and recording the compound action potential more proximally [32]. The stimulating

electrodes were positioned on the finger as described above to stimulate the digital nerves which

innervate the index finger. The recording electrodes were located on the wrist to record activity in

the median nerve, while the ground electrode was placed on the thenar eminence. These recordings

were used to verify that the afferent axons fired reliably to each electrical pulse within a burst in our

closely spaced stimulation patterns.

Measuring point of subjective equality for frequency

To determine the perceived frequency of each test stimulus, a two-interval forced choice paradigm

was used as in our previous study [5]. Test stimuli were compared against 6-8 different comparison

stimuli, which were regular trains of pulses with repetition rates ranging from 14-54 Hz for

Experiment 1 and 7-21 Hz for Experiment 2. For each trial, a test stimulus and a comparison stimulus

were presented for 1 s each, in a random order and separated by 500 ms (Fig 1C). The subject

indicated which of the two stimuli had the higher frequency. To obtain psychometric curves, each

test condition was compared 20 times against each of the regular comparison frequencies, giving

rise to 120-160 trials per test stimulus. Following optimisation of the electrical current, a practice

round was conducted which comprised a selection of test and comparison stimuli to familiarise

subjects with the task, before the actual test procedure began.

At each comparison frequency, we determined the proportion of times the subject indicated that

the comparison stimulus had a higher frequency than the test stimulus (P_H). To obtain a linear

psychometric function, the logit transformation $\ln(P_H/(1-P_H))$ was applied to the data [33]. The value

of the comparison frequency that is equally likely to be judged as higher or lower than the test

stimulus was taken as the point of subjective equality (PSE). This value corresponds to the frequency

where the zero crossing of the logit axis occurs for the regression line fitted to the logit transformed

data (example in S1 Fig).

Statistical analysis

For Experiment 1, linear regression was performed to determine if the slope of the line fitted to the

data would be better explained by the burst gap code or the periodicity, which was constant at 23

Hz. A two-way ANOVA (repeated measures within the same subject for the different stimulation

methods) was used to determine if there were differences between the data collected here using

electrical stimulation and the mechanical data collected in our previous study. For Experiment 2, a

joinpoint regression model created using Joinpoint Trend Analysis Software (National Cancer

Institute, Bethesda, MD, USA) was used to determine the trend that best fitted the psychophysical

data. An alpha level of 0.05 was used for all statistical tests.

Results

Stable activation of the afferent population

Electrical stimulation with patterns used in Experiment 1 (Fig 1A) resulted in SNAPs that were of

consistent amplitude during a burst irrespective of inter-pulse intervals used or number of pulses

within a burst (Fig 2). Fig 3 illustrates the consistency of the slopes for each averaged SNAP

waveform within a burst in both subjects, indicating that the afferent population recruited remained

stable between pulses in a burst.

Fig 2. Sensory nerve action potential (SNAP) recordings. Average of 200 compound action

potentials recorded from the median nerve in response to Experiment 1 stimulation patterns

illustrated in Fig 1A. The red circles indicate SNAPs, while the clipped waveform occurring before

each SNAP represents the electrical stimulation artefact.

Fig 3. Consistency of evoked SNAPs within a burst. Each panel represents the slopes measured from

the downward phase of the averaged waveform (n = 200) for evoked SNAPs within a burst in a single

subject (n = 2). Error bars denote standard deviation.

How is perceived frequency encoded?

In Experiment 1, the aim was to determine which coding mechanism could best describe the

perception of frequency in subjects when the electrical pulses were grouped into bursts. The mean

data from the 12 subjects are shown in Fig 4. The observed perceived frequencies were 2- to 4-fold

lower than that predicted by the mean pulse rate (Fig 4A), indicating that a simple rate code could

not explain perceived frequency in subjects. Linear regression was performed between the

reciprocal of the burst gap (t_L) and the perceived frequency for each stimulus (Fig 4B). The regression

line had a slope that was non-zero (1.2, 95% CI 1.0-1.4; R^2 = 0.76; p < 0.0001), which does not

correspond to that of the periodicity (expected slope 0), but is a good match to the reciprocal of the

burst gap (t_L). The slope here closely resembles that of the mechanical data from our previous study

(slope = 1.3, 95% CI 1.0-1.5; R^2 = 0.68; p < 0.0001), which also best matched $1/t_L$ [5].

Fig 4. Possible coding schemes for frequency perception of electrical burst stimuli. (A) Boxplots

with whiskers represent the subjective perceived frequency of the electrical stimuli used in

Experiment 1 (n = 12). Green and pink arrowheads indicate predicted PSE values if the frequency

judgement had been based on the mean pulse rate and the burst rate (periodicity), respectively. (B)

Fitted regression line (black line) indicating that the mean electrical PSE data (black circles; n = 12)

from this study is a close match to the reciprocal of the burst gap t_L. The regression line fitted to

mechanical data from our previous study [5] is shown for comparison (orange line). The dotted line

is the line of identity, where the PSE would equal the reciprocal of the burst gap t_L. Error bars denote

95% confidence intervals.

A two-way ANOVA indicated that there was a statistically significant main effect by the stimulus

pattern (p < 0.0001, $F_{2,44}$ = 143, 71% of total variation) and the method of stimulation (p = 0.0084,

$F_{1,22}$ = 8.4, 4.7% of total variation). However, there was no interaction effect (p = 0.36, $F_{3,66}$ = 1.1),

suggesting that varying the stimulation pattern had the same effect on frequency perception

regardless of whether mechanical or electrical stimulation was used.

200 What defines an inter-pulse interval as part of a burst?

For Experiment 2, we varied the duration of the short interval to determine which inter-pulse

intervals (t_S) would be treated as a burst, so that the perceived frequency depended only on the

longer interval (t_L). The results from the 8 subjects are shown in Fig 5 which plots the apparent

frequency of each stimulus. A joinpoint regression model was fitted, which indicated that the data

was best explained by three linear segments. The first segment extended from 4.35 to 13.05 ms,

with a slope that was not significantly different from zero (p = 0.64). The next two segments had

slopes that were significantly non-zero. The second segment spanned 13.05 to 34.8 ms and had a

slope of 0.2 (p = 0.0002). The last segment had a range between 34.8 and 56.55 ms, with a slope of -

0.08 (p = 0.0002).

The slope for the segments here were similar to those reported in our previous study involving

mechanical stimulation (Fig 5, Mechanical PSE), though the joinpoint between the second and third

segments occurred at 30.45 ms in that study [5], thus at a slightly shorter inter-pulse interval than in

the current study (34.8 ms). Nevertheless, the same basic findings were observed in both studies, as

shown by the confidence intervals overlapping with the joinpoint model based on the mechanical

data and an identical first jointpoint. This suggests that the same definition of a burst can be applied

to both mechanical and electrical stimulation.

Fig 5. Relationship between the short inter-pulse interval and perceived frequency. PSE values

obtained using Experiment 2 stimuli, indicated by black circles (Electrical PSE; n = 8). The black line

signifies the three fitted linear segments as determined by the joinpoint regression model. The green

and blue lines represent predicted PSE values if frequency judgement had been based on the mean

pulse rate and the burst gap code, respectively. The orange line (Mechanical PSE) is the joinpoint

model based on the mechanical data from our previous study [5]. Error bars denote 95% confidence

intervals.

226 Discussion

The main objective of the present study was to verify whether the burst gap coding model for

frequency perception could be observed with electrical stimulation. This is an important step which

had to be demonstrated due to the potential value of this coding strategy for implementation in

bionic systems for providing somatosensory information. Our results show that the burst gap code

could reliably predict perceived frequency of complex bursting spike patterns when the spikes were

elicited by electrical pulse stimulation of axons. This means that the same overall spike rate in an

axon can produce a different frequency percept, depending on how those spikes are arranged in

time. This reinforces the importance of the temporal structure of spiking patterns in neural

processing and sensory perception.

Electrical stimulation bypasses the complex viscoelastic properties of the skin that may have affected

previous work using mechanical stimulation, and so provides strong independent confirmation of

the burst gap code. We verified that the efficiency of the electrical stimulus was not affected by the

repetition rate with SNAP recordings, which showed that even the shortest inter-pulse interval (4.35

ms) resulted in reliable and consistent activation of tactile afferents. As we used a reasonably strong

stimulation current at a supra-threshold level, it was expected that a majority, if not all, of the large

myelinated afferents of the digital nerve would be reliably activated regardless of the inter-pulse

interval that was chosen.

Although the results in our first experiment, where we used electrical pulses to create bursting trains

of the same periodicity but different number of spikes, were best explained by the burst gap code

rather than a periodicity or rate code, the mean PSEs were lower than that would be predicted

simply from the burst gap duration. One explanation for this bias towards lower frequencies may be

that electrical stimulation recruits all afferent types non-selectively, so that the slowly adapting (SA)

afferents would be recruited in addition to the fast adapting (FA) afferents, whereas the mechanical

stimulus predominantly activated FA afferents [5]. Although previous research has found that SA

afferents and their corresponding cortical sensory neurons do not contribute to vibrotactile

frequency perception [34-36], their activation and the corresponding sense of pressure they convey

[24] may bias subject judgements in psychophysics tests. However, although the difference in PSEs

between electrical and mechanical stimulation was statistically significant, it is unknown whether

this difference indicates any physiological importance, as the difference is well within the Weber

fraction of ~0.2-0.3 that has been previously reported in the literature [11, 17, 37-40].

The data from the second experiment, where doublets of different inter-pulse durations were

tested, clearly indicate that the frequency of periodic doublets is perceived in a similar manner for

both mechanical and electrical stimuli, as the criterion defining a burst observed with mechanical

stimuli [5] could also be applied to electrical stimulation. When the inter-pulse interval is sufficiently

brief (<15 ms), the pulses are interpreted as belonging to a single burst event and the burst gap

determines the perceived frequency. As the separation of the two pulses is increased beyond 15 ms,

the propensity for two consecutive pulses to be regarded as a burst decreases and the short interval

appears to have a partial weighted contribution to frequency perception [41]. Beyond 30-35 ms,

each individual pulse becomes a discrete event and the perceived frequency resembles the mean

pulse rate.

These findings lend weight to another recent study which revealed that spiking pattern determined

the perceived frequency of mechanical stimuli, even when driven by different receptor types [17]. In

that study, we found that low-frequency spike trains in Pacinian (FA2) afferents could induce a

vibratory precept, even though Meissner (FA1) afferents had been traditionally thought to be solely

responsible for perception of low frequency flutter vibrations (<60 Hz), whereas higher frequency

activation of Pacinian (FA2) afferents would evoke a different percept of vibratory hum [25]. A

generalised frequency processing system that operates across a variety of receptor type inputs

would explain these findings, and the constancy of vibrotactile frequency perception across different

skin regions innervated by different afferent types [37, 42], e.g. hairy skin lack Meissner corpuscles

[43, 44]. The congruence of the present results using electrical stimulation whereby all types of

afferents are recruited non-selectively, with the previous results using mechanical stimuli that

activated predominantly FA afferents, supports the suggestion of a generalised frequency processing

mechanism.

Natural stimuli encountered during object manipulation and surface exploration generally evoke

complex discharge patterns in tactile afferents [2, 45-47]. Additionally, numerous studies have

implicated the importance of frequency content in skin vibrations for encoding tactile information

such as texture [45, 48, 49]. Accordingly, the frequency perception mechanism substantiated here

may represent a more universal method of encoding complex naturalistic patterns of vibration,

which may not necessarily have a simple fixed periodicity. For instance, we previously showed that

this sensory coding strategy also applied to stimuli that are irregular or aperiodic in nature, as it

relies on the processing of individual inter-pulse intervals by weighting their perceptual contribution

to perceived frequency depending on their duration, rather than detection of periodicity or spike

counting [41]. Hence, this neural coding strategy may provide a good way to manipulate frequency

perception while providing sensory feedback in settings such as neural prostheses or haptic displays,

through electrical stimulation of the tactile system.

References

1.   Saal HP, Wang X, Bensmaia SJ. Importance of spike timing in touch: an analogy with hearing?

Curr Opin Neurobiol. 2016;40:142-9. Epub 2016/08/10. doi: 10.1016/j.conb.2016.07.013. PubMed

PMID: 27504741; PubMed Central PMCID: PMCPMC5315566.

2.   Mackevicius EL, Best MD, Saal HP, Bensmaia SJ. Millisecond precision spike timing shapes

tactile perception. J Neurosci. 2012;32(44):15309-17. doi: 10.1523/JNEUROSCI.2161-12.2012.

PubMed PMID: 23115169; PubMed Central PMCID: PMCPMC3752122.

3.   Johansson RS, Birznieks I. First spikes in ensembles of human tactile afferents code complex

spatial fingertip events. Nat Neurosci. 2004;7(2):170-7. Epub 2004/01/20. doi: 10.1038/nn1177.

PubMed PMID: 14730306.

4.   Saal HP, Vijayakumar S, Johansson RS. Information about complex fingertip parameters in

individual human tactile afferent neurons. J Neurosci. 2009;29(25):8022-31. Epub 2009/06/26. doi:

10.1523/JNEUROSCI.0665-09.2009. PubMed PMID: 19553442; PubMed Central PMCID:

PMCPMC6666033.

5.   Birznieks I, Vickery RM. Spike timing matters in novel neuronal code involved in vibrotactile

frequency perception. Curr Biol. 2017;27(10):1485-90.e2. Epub 2017/05/10. doi:

10.1016/j.cub.2017.04.011. PubMed PMID: 28479322.

6.   Salinas E, Hernandez A, Zainos A, Romo R. Periodicity and firing rate as candidate neural

codes for the frequency of vibrotactile stimuli. J Neurosci. 2000;20(14):5503-15. Epub 2000/07/07.

doi: 10.1523/JNEUROSCI.20-14-05503.2000. PubMed PMID: 10884334; PubMed Central PMCID:

PMCPMC6772326.

7.   Mountcastle VB, Talbot WH, Sakata H, Hyvärinen J. Cortical neuronal mechanisms in flutter-

vibration studied in unanesthetized monkeys. Neuronal periodicity and frequency discrimination. J

Neurophysiol. 1969;32(3):452-84. Epub 1969/05/01. doi: 10.1152/jn.1969.32.3.452. PubMed PMID:

4977839.

8.   Luna R, Hernandez A, Brody CD, Romo R. Neural codes for perceptual discrimination in

primary somatosensory cortex. Nat Neurosci. 2005;8(9):1210-9. Epub 2005/08/02. doi:

10.1038/nn1513. PubMed PMID: 16056223.

9.   Romo R, Salinas E. Flutter discrimination: neural codes, perception, memory and decision

making. Nat Rev Neurosci. 2003;4(3):203-18. Epub 2003/03/04. doi: 10.1038/nrn1058. PubMed

PMID: 12612633.

10.   Tan DW, Schiefer MA, Keith MW, Anderson JR, Tyler J, Tyler DJ. A neural interface provides

long-term stable natural touch perception. Sci Transl Med. 2014;6(257):257ra138. Epub 2014/10/10.

doi: 10.1126/scitranslmed.3008669. PubMed PMID: 25298320; PubMed Central PMCID:

PMCPMC5517305.

11.   Graczyk EL, Schiefer MA, Saal HP, Delhaye BP, Bensmaia SJ, Tyler DJ. The neural basis of

perceived intensity in natural and artificial touch. Sci Transl Med. 2016;8(362):362ra142. Epub

2016/11/01. doi: 10.1126/scitranslmed.aaf5187. PubMed PMID: 27797958; PubMed Central PMCID:

PMCPMC5713478.

12.   Schiefer MA, Graczyk EL, Sidik SM, Tan DW, Tyler DJ. Artificial tactile and proprioceptive

feedback improves performance and confidence on object identification tasks. PLoS One.

2018;13(12):e0207659. Epub 2018/12/06. doi: 10.1371/journal.pone.0207659. PubMed PMID:

30517154; PubMed Central PMCID: PMCPMC6281191.

13.   Grill WM. Temporal pattern of electrical stimulation is a new dimension of therapeutic

innovation. Curr Opin Biomed Eng. 2018;8:1-6. Epub 2019/03/25. doi:

10.1016/j.cobme.2018.08.007. PubMed PMID: 30906909; PubMed Central PMCID:

PMCPMC6426311.

14.   Brocker DT, Swan BD, Turner DA, Gross RE, Tatter SB, Koop MM, et al. Improved efficacy of

temporally non-regular deep brain stimulation in Parkinson's disease. Exp Neurol. 2013;239:60-7.

Epub 2012/10/02. doi: 10.1016/j.expneurol.2012.09.008. PubMed PMID: 23022917; PubMed Central

PMCID: PMCPMC3547657.

15.   Baker KB, Zhang J, Vitek JL. Pallidal stimulation: effect of pattern and rate on bradykinesia in

the non-human primate model of Parkinson's disease. Exp Neurol. 2011;231(2):309-13. Epub

2011/07/20. doi: 10.1016/j.expneurol.2011.06.012. PubMed PMID: 21767534; PubMed Central

PMCID: PMCPMC3536492.

16.   Hess CW, Vaillancourt DE, Okun MS. The temporal pattern of stimulation may be important

to the mechanism of deep brain stimulation. Exp Neurol. 2013;247:296-302. Epub 2013/02/13. doi:

10.1016/j.expneurol.2013.02.001. PubMed PMID: 23399890; PubMed Central PMCID:

PMCPMC3742615.

17.   Birznieks I, McIntyre S, Nilsson HM, Nagi SS, Macefield VG, Mahns DA, et al. Tactile sensory

channels over-ruled by frequency decoding system that utilizes spike pattern regardless of receptor

type. eLife. 2019;8. Epub 2019/08/07. doi: 10.7554/eLife.46510. PubMed PMID: 31383258; PubMed

Central PMCID: PMCPMC6684274.

18.   Gardner EP, Palmer CI. Simulation of motion on the skin. I. Receptive fields and temporal

frequency coding by cutaneous mechanoreceptors of OPTACON pulses delivered to the hand. J

Neurophysiol. 1989;62(6):1410-36. Epub 1989/12/01. doi: 10.1152/jn.1989.62.6.1410. PubMed

PMID: 2600632.

19.   Kaczmarek KA, Tyler ME, Brisben AJ, Johnson KO. The afferent neural response to

electrotactile stimuli: preliminary results. IEEE Trans Rehabil Eng. 2000;8(2):268-70. Epub

2000/07/15. doi: 10.1109/86.847828. PubMed PMID: 10896199.

20.   Rollman GB. Electrocutaneous stimulation: Psychometric functions and temporal

integration. Percept Psychophys. 1969;5(5):289-93. doi: 10.3758/bf03209565.

21.   Rollman GB. Electrocutaneous stimulation. In: Geldard FA, editor. Conference on cutaneous

communication systems and devices. Austin, Texas: Psychonomic Society; 1974. p. 38-51.

22.   Steenbergen P, Buitenweg JR, Trojan J, van der Heide EM, van den Heuvel T, Flor H, et al. A

system for inducing concurrent tactile and nociceptive sensations at the same site using

electrocutaneous stimulation. Behav Res Methods. 2012;44(4):924-33. Epub 2012/07/19. doi:

10.3758/s13428-012-0216-y. PubMed PMID: 22806702; PubMed Central PMCID: PMCPMC3509324.

23.   van der Heide EM, Buitenweg JR, Marani E, Rutten WL. Single pulse and pulse train

modulation of cutaneous electrical stimulation: a comparison of methods. J Clin Neurophysiol.

2009;26(1):54-60. Epub 2009/01/20. doi: 10.1097/WNP.0b013e3181942cd2. PubMed PMID:

19151619.

24.   Johansson RS, Landström U, Lundström R. Responses of mechanoreceptive afferent units in

the glabrous skin of the human hand to sinusoidal skin displacements. Brain Res. 1982;244(1):17-25.

Epub 1982/07/22. doi: 10.1016/0006-8993(82)90899-x. PubMed PMID: 6288178.

25.   Talbot WH, Darian-Smith I, Kornhuber HH, Mountcastle VB. The sense of flutter-vibration:

comparison of the human capacity with response patterns of mechanoreceptive afferents from the

monkey hand. J Neurophysiol. 1968;31(2):301-34. Epub 1968/03/01. doi: 10.1152/jn.1968.31.2.301.

PubMed PMID: 4972033.

26.   Manfredi LR, Baker AT, Elias DO, Dammann JF, 3rd, Zielinski MC, Polashock VS, et al. The

effect of surface wave propagation on neural responses to vibration in primate glabrous skin. PLoS

One. 2012;7(2):e31203. Epub 2012/02/22. doi: 10.1371/journal.pone.0031203. PubMed PMID:

22348055; PubMed Central PMCID: PMCPMC3278420.

27.   Dandu B, Shao Y, Stanley A, Visell Y. Spatiotemporal Haptic Effects from a Single Actuator via

Spectral Control of Cutaneous Wave Propagation. 2019 IEEE World Haptics Conference (WHC); 2019

9-12 July 2019; Tokyo, Japan. doi: 10.1109/WHC.2019.8816149.

28.   Johnson KO. Reconstruction of population response to a vibratory stimulus in quickly

adapting mechanoreceptive afferent fiber population innervating glabrous skin of the monkey. J

Neurophysiol. 1974;37(1):48-72. Epub 1974/01/01. doi: 10.1152/jn.1974.37.1.48. PubMed PMID:

4204567.

29.   Bensmaia SJ. Tactile intensity and population codes. Behav Brain Res. 2008;190(2):165-73.

Epub 2008/04/19. doi: 10.1016/j.bbr.2008.02.044. PubMed PMID: 18420286; PubMed Central

PMCID: PMCPMC2659538.

30.   Kajimoto H. Electro-tactile Display: Principle and Hardware. In: Kajimoto H, Saga S, Konyo M,

editors. Pervasive Haptics: Science, Design, and Application. Tokyo: Springer Japan; 2016. p. 79-96.

doi: 10.1007/978-4-431-55772-2_5.

31.   von Békésy G. Synchronism of neural discharges and their demultiplication in pitch

perception on the skin and in hearing. J Acoust Soc Am. 1959;31(3):338-49. doi: 10.1121/1.1907722.

32.   Mallik A, Weir AI. Nerve conduction studies: essentials and pitfalls in practice. J Neurol

Neurosurg Psychiatry. 2005;76(Suppl 2):ii23-ii31. Epub 2005/06/18. doi: 10.1136/jnnp.2005.069138.

PubMed PMID: 15961865; PubMed Central PMCID: PMCPMC1765692.

33.   Vickery RM, Morley JW, Rowe MJ. The role of single touch domes in tactile perception. Exp

Brain Res. 1993;93(2):332-4. Epub 1993/01/01. doi: 10.1007/bf00228402. PubMed PMID: 8491272.

34.   Vallbo ÅB, Olsson KÅ, Westberg KG, Clark FJ. Microstimulation of single tactile afferents from

the human hand. Sensory attributes related to unit type and properties of receptive fields. Brain.

1984;107(3):727-49. Epub 1984/09/01. doi: 10.1093/brain/107.3.727. PubMed PMID: 6478176.

35.   Romo R, Hernandez A, Zainos A, Brody CD, Lemus L. Sensing without touching:

psychophysical performance based on cortical microstimulation. Neuron. 2000;26(1):273-8. Epub

2000/05/08. doi: 10.1016/s0896-6273(00)81156-3. PubMed PMID: 10798410.

36.   Merzenich MM, Harrington TH. The sense of flutter-vibration evoked by stimulation of the

hairy skin of primates: comparison of human sensory capacity with the responses of

mechanoreceptive afferents innervating the hairy skin of monkeys. Exp Brain Res. 1969;9(3):236-60.

Epub 1969/01/01. doi: 10.1007/bf00234457. PubMed PMID: 4984464.

37.   Mahns DA, Perkins NM, Sahai V, Robinson L, Rowe MJ. Vibrotactile frequency discrimination

in human hairy skin. J Neurophysiol. 2006;95(3):1442-50. Epub 2005/12/02. doi:

10.1152/jn.00483.2005. PubMed PMID: 16319219.

38.   Rothenberg M, Verrillo RT, Zahorian SA, Brachman ML, Bolanowski SJ, Jr. Vibrotactile

frequency for encoding a speech parameter. J Acoust Soc Am. 1977;62(4):1003-12. Epub

1977/10/01. doi: 10.1121/1.381610. PubMed PMID: 908786.

39.   Goff GD. Differential discrimination of frequency of cutaneous mechanical vibration. J Exp

Psychol. 1967;74(2):294-9. Epub 1967/06/01. doi: 10.1037/h0024561. PubMed PMID: 6048472.

40.   Li M, Zhang D, Chen Y, Chai X, He L, Chen Y, et al. Discrimination and Recognition of Phantom

Finger Sensation Through Transcutaneous Electrical Nerve Stimulation. Front Neurosci. 2018;12:283.

Epub 2018/05/16. doi: 10.3389/fnins.2018.00283. PubMed PMID: 29760647; PubMed Central

PMCID: PMCPMC5937010.

41.   Ng KKW, Birznieks I, Tse ITH, Andersen J, Nilsson S, Vickery RM. Perceived Frequency of

Aperiodic Vibrotactile Stimuli Depends on Temporal Encoding. In: Prattichizzo D, Shinoda H, Tan HZ,

Ruffaldi E, Frisoli A, editors. Haptics: Science, Technology, and Applications. EuroHaptics 2018.

Lecture Notes in Computer Science. 10893. Cham: Springer International Publishing; 2018. p. 199-

208. doi: 10.1007/978-3-319-93445-7_18.

42.   McIntyre S, Birznieks I, Andersson R, Dicander G, Breen PP, Vickery RM. Temporal

Integration of Tactile Inputs from Multiple Sites. In: Bello F, Kajimoto H, Visell Y, editors. Haptics:

Perception, Devices, Control, and Applications. Eurohaptics 2016. Lecture Notes in Computer

Science. 9774. Cham: Springer International Publishing; 2016. p. 204-13. doi: 10.1007/978-3-319-

42321-0_19.

43.   Vallbo ÅB, Olausson H, Wessberg J, Kakuda N. Receptive field characteristics of tactile units

with myelinated afferents in hairy skin of human subjects. J Physiol. 1995;483(3):783-95. Epub

1995/03/15. doi: 10.1113/jphysiol.1995.sp020622. PubMed PMID: 7776258; PubMed Central

PMCID: PMCPMC1157818.

44.   Bolton CF, Winkelmann RK, Dyck PJ. A quantitative study of Meissner's corpuscles in man.

Neurology. 1966;16(1):1-9. Epub 1966/01/01. doi: 10.1212/wnl.16.1.1. PubMed PMID: 5948004.

45.   Manfredi LR, Saal HP, Brown KJ, Zielinski MC, Dammann JF, 3rd, Polashock VS, et al. Natural

scenes in tactile texture. J Neurophysiol. 2014;111(9):1792-802. Epub 2014/02/14. doi:

10.1152/jn.00680.2013. PubMed PMID: 24523522.

46.   Hollins M, Bensmaia SJ, Roy EA. Vibrotaction and texture perception. Behav Brain Res.

2002;135(1-2):51-6. Epub 2002/10/03. doi: 10.1016/s0166-4328(02)00154-7. PubMed PMID:

12356433.

47.   Johansson RS, Flanagan JR. Coding and use of tactile signals from the fingertips in object

manipulation tasks. Nat Rev Neurosci. 2009;10(5):345-59. Epub 2009/04/09. doi: 10.1038/nrn2621.

PubMed PMID: 19352402.

48.   Delhaye B, Hayward V, Lefevre P, Thonnard JL. Texture-induced vibrations in the forearm

during tactile exploration. Front Behav Neurosci. 2012;6:37. Epub 2012/07/12. doi:

10.3389/fnbeh.2012.00037. PubMed PMID: 22783177; PubMed Central PMCID: PMCPMC3390558.

49.   Weber AI, Saal HP, Lieber JD, Cheng JW, Manfredi LR, Dammann JF, 3rd, et al. Spatial and

temporal codes mediate the tactile perception of natural textures. Proc Natl Acad Sci U S A.

2013;110(42):17107-12. Epub 2013/10/02. doi: 10.1073/pnas.1305509110. PubMed PMID:

24082087; PubMed Central PMCID: PMCPMC3800989.

Supporting information

S1 Fig. Example psychometric function. An example of a logistic regression fit applied to psychometric data from a single subject in one test condition (Experiment 1, stimulus d). PSE (point of subjective equality) is determined as the frequency that gives logit = 0 from the fitted regression line. In this case, PSE was estimated to be 38.1 Hz.

S2 Dataset. Individual subject PSE values for each experiment.

a

b

c

d