A novel mapping strategy utilizing mouse chromosome substitution strains identifies multiple epistatic interactions that regulate complex traits

Anna K. Miller[*], Anlu Chen[†1], Jacquelaine Bartlett[‡], Li Wang[*], Scott M. Williams[*‡§] and David A. Buchner[*†§]

[*]Department of Genetics and Genome Sciences, Case Western Reserve University School of Medicine, Cleveland, OH 44106

[†]Department of Biochemistry, Case Western Reserve University School of Medicine, Cleveland, OH 44106

[‡]Department of Population and Quantitative Health Sciences, Case Western Reserve University School of Medicine, Cleveland, OH 44106

§ These authors contributed equally

[1]Division of Endocrinology, Diabetes, and Metabolism, Beth Israel Deaconess Medical Center, and Harvard Medical School, Boston, MA 02115

Data availability:
RNA-Seq data is available in GEO
Pipeline scripts are available in protocols.io

30   Running title: Epistasis in complex traits
31
32   Key words: QTL, complex trait, missing heritability, genetic architecture, epistasis
33
34   Corresponding author's contact information:
35

36   **David A. Buchner, Case Western Reserve University School of Medicine, 10900 Euclid**
37   **Ave., Cleveland, OH, 44106-4935, 216-368-1816, dab22@case.edu**

Abstract

The genetic contribution of additive versus non-additive (epistatic) effects in the regulation of complex traits is unclear. While genome-wide association studies typically ignore gene-gene interactions, in part because of the lack of statistical power for detecting them, mouse chromosome substitution strains (CSSs) represent an alternate and powerful model for detecting epistasis given their limited allelic variation. Therefore, we utilized CSSs to identify and map both additive and epistatic loci that regulate a range of hematologic- and metabolism-related traits, as well as hepatic gene expression. Quantitative trait loci (QTLs) were identified using a CSS-based backcross strategy involving the segregation of variants on the A/J-derived substituted chromosomes 4 and 6 on an otherwise C57BL/6J genetic background. In the liver transcriptomes of offspring from this cross, we identified and mapped additive QTLs regulating the hepatic expression of 768 genes, and epistatic QTL pairs for 519 genes. Similarly, we identified additive QTLs for fat pad weight, platelets, and the percentage of granulocytes in blood, as well as epistatic QTL pairs controlling the percentage of lymphocytes in blood and red cell distribution width. The variance attributed to the epistatic QTL pairs was approximately equal to that of the additive QTLs; however, the SNPs in the epistatic QTL pairs that accounted for the largest variances were undetected in our single locus association analyses. These findings highlight the need to account for epistasis in association studies, and more broadly demonstrate the importance of identifying genetic interactions to understand the complete genetic architecture of complex traits.

Introduction

The severity, presentation, and prevalence of complex traits and diseases are influenced by many

genetic variants (FU *et al.* 2013; BOYLE *et al.* 2017). However, it remains unclear whether the

variants work together in an additive manner or have non-linear effects on the phenotype, referred

to as epistasis (CORDELL 2002; CARLBORG AND HALEY 2004; JASNOS AND KORONA 2007;

MANOLIO *et al.* 2009; BLOOM *et al.* 2013). Epistasis has been extensively observed in model

organisms including *S. cerevisiae* (JASNOS AND KORONA 2007; BLOOM *et al.* 2013; FORSBERG *et*

al. 2017), *C. elegans* (LEHNER *et al.* 2006), *D. melanogaster* (HUANG *et al.* 2012), and *M. musculus*

(SHAO *et al.* 2008; TYLER *et al.* 2017). In humans, epistasis has been more difficult to detect with

standard genetic analyses. This is potentially due to low allele frequencies, limited sample sizes,

complexity of interactions, insufficient effect sizes, diversity of genetic backgrounds, and

methodological limitations due to the fact that in humans evidence for genetic effects is almost

always statistical (MACKAY 2014; HUANG AND MACKAY 2016). However, despite these inherent

difficulties, epistasis in humans has been detected in genome-wide interaction-based association

studies and other methods for Crohn's disease, Glaucoma, Behçet's disease, multiple sclerosis,

Hirschbrung disease, among others (LIU *et al.* 2011; HU *et al.* 2013; KIRINO *et al.* 2013; HEMANI

et al. 2014; HUANG *et al.* 2015; VERMA *et al.* 2016; GALARZA-MUNOZ *et al.* 2017; CHATTERJEE

AND CHAKRAVARTI 2019; TYLER *et al.* 2020). Assuming these examples generalize, discovering

and understanding epistatic interactions will be critical for improving the predicting of phenotypes

from genetic data as well as understanding of pathophysiology, thereby better guiding precision

medicine-based decisions (MOORE AND WILLIAMS 2009; COLE *et al.* 2017).

One particularly powerful method of detecting epistasis in a mammalian system is based on studies

of mouse chromosome substitution strains (CSSs) (NADEAU *et al.* 2000). The genomes of CSSs

are comprised of one chromosome derived from a donor strain, with the remainder derived from a

host strain such that when two CSSs are crossed together, any genetic interactions between the two

non-homologous donor chromosomes can be more readily detected. This system provides an

efficient model to map interacting quantitative trait loci (QTLs) on a fixed genetic background.

This is in contrast to populations with genome-wide allelic variation that segregate in mapping

populations such as the Diversity Outbred mouse population or typical analyses in humans

(CHURCHILL *et al.* 2012). Specifically, as it relates to epistasis, what might otherwise be a rare

allelic combination in a segregating cross can be fixed in the genome of a CSS, allowing for

detailed and reproducible studies of specific allelic combinations. Initial studies of CSSs derived

from strains C57BL/6J and A/J, which differ with respect to metabolic disease and cancer risk,

indicated that epistatic interactions were a dominant feature of many complex traits (SHAO *et al.*

2008).

More recent work on CSS combinations with two substituted chromosomes highlighted the

importance of epistasis in the regulation of complex traits and gene expression, and built upon

previous studies by directly identifying pairs of QTLs (chromosomes) with non-additive

phenotypic effects. These studies found that in mice carrying two A/J-derived chromosomes on an

otherwise C57BL/6J genetic background, the A/J chromosomes frequently showed evidence of

non-additive interactions (CHEN *et al.* 2017). The patterns of interactions were largely in the

direction of negative (or suppressing) epistasis, such that traits that were altered due to the effects

of one of the substituted chromosomes were returned to baseline levels due to the combined effects

of the two interacting substituted chromosomes (CHEN *et al.* 2017). These findings indicated that

epistatic interactions can be a common feature of complex traits as well as important for

maintaining homeostasis.

A major limitation to previous analyses of epistasis using CSSs is that the resolution of QTLs

mapped was at the size of the entire substituted chromosomes. However, to identify the molecular

nature of such QTLs requires identification of the underlying genes, which necessitates higher

resolution mapping studies. Therefore, to identify novel epistatic QTLs that regulate complex traits

and gene expression levels with greater mapping resolution, we generated and analyzed the N2

generation mice from a modified backcross based on using some of the same CSSs in Chen *et al*.

(2017). The resulting analyses demonstrate the ability of a CSS-based backcross to detect and fine-

map epistatic QTLs for complex traits. The analyses discovered significant epistatic interactions

that control lymphocyte quantity and the distribution of red cell width, and the expression of

hundreds of genes in the liver. Studies of this type are essential to make progress towards the

molecular characterization of epistatic interactions that will provide insight into the biological

pathways of disease pathophysiology as well as improve our understanding of trait heritability and

genetic architecture.

Materials and Methods

Mice

Strains C57BL/6J-Chr6$^{A/J}$/NaJ (B6.A6) (stock #004384), C57BL/6J-Chr4$^{A/J}$/NaJ (B6.A4) (stock

#004382), and C57BL/6J (B6) (stock #000664) were purchased from The Jackson Laboratory.

Mice were maintained by brother-sister matings with offspring weaned at 3 weeks of age. Mice

were housed in ventilated racks with access to food and water *ad libitum* and maintained at 21ºC

on a 12-hour light/12-hour dark cycle. All mice were cared for as described under the Guide for

the Care and Use of Animals, eighth edition (2011) and all experiments were approved by IACUC

(protocol #2016-0064) and carried out in an AAALAC approved facility. Mice were anesthetized

with isoflurane prior to retro-orbital bleeding and subsequently euthanized under anesthesia by

cervical dislocation for tissue collection.

Genotyping

Genomic DNA was isolated from mouse tail tissue using DNeasy Blood and Tissue Kits (Qiagen)

following digestion with proteinase K. N2 offspring of the CSS backcross were genotyped with

57 SNP markers on Chromosome 4 and 50 SNP markers on Chromosome 6 (~1 marker per 1.5

cM) using the PlexSeq technology (Agriplex Genomics Inc.) (KAYIMA *et al.* 2017; SOONG *et al.*

2018). Genotyping results are listed in Supplemental Table 1. Of the 150 samples, one sample did

not pass the PlexSeq quality control (<66% of genotypes called) and was not used in these

analyses.

Mouse Phenotyping

Five-week old mice were fasted overnight prior to phenotypic analysis. Mice were anesthetized

with isoflurane and measurements were collected for total body weight and nose to anus length.

Body mass index (BMI) was calculated as g/cm^2. Blood was collected retro-orbitally and glucose

levels were measured using an OneTouch Ultra2 meter (LifeScan, Milpitas, CA, USA). Mice were

subsequently euthanized by cervical dislocation, gonadal fat pads were removed and weighed, and

the caudate lobe of the liver was collected and immediately placed in RNAlater Stabilization

Solution (Thermo Fisher Scientific). Complete blood counts were performed on either a Heska

HemaTrue or Drew Scientific Hemavet 950 blood analyzer and included measurements of white

blood count (wbc), neutrophils (ne), lymphocytes (ly), monocytes (mo), red blood count (rbc),

hemoglobin (hb), hematocrit (hct), mean corpuscular volume (mcv), mean corpuscular

hemoglobin (mch), mean corpuscular hemoglobin concentration (mchc), red cell distribution

width (rdw), platelet count (plt), mean platelet volume (mpv), granulocyte % (gran%), granulocyte

KuL (gran KuL), and red cell distribution width (absolute) (rdwa).

Transcriptome analysis

Total RNA was isolated from liver using PureLink RNA mini kit (Thermo Fisher Scientific). RNA

quality was analyzed using an Agilent TapeStation with all resulting RNA Integrity Numbers

(RIN) greater than 8.5. Sequencing libraries were generated from total RNA (1 ug) using the

TruSeq stranded mRNA kit (Illumina Corp.) and were sequenced using a NovaSeq6000 S4.

Paired-end sequencing was performed at an average read number of $51,404,587 \pm 12,403,503$

reads per sample with an average of 92.7% of bases reaching quality scores of at least Q30. RNA

quality control, library construction, and sequencing were performed by Macrogen Inc (now

Psomagen) of Rockville, Maryland.

Prior to gene expression analysis, Fastqc (version 0.11.8) was used to determine the sequence

quality of the fastq files before and after adapters were trimmed with Trimmomatic (version 0.36)

(ANDREWS 2010; BOLGER *et al.* 2014). Sequencing reads were aligned to the C57BL/6J mouse

genome by Hisat2 (version 2.1.0) using the Ensembl GRCm38.p5 primary DNA assembly

reference sequence (KIM *et al.* 2015; CUNNINGHAM *et al.* 2019). On average, the overall alignment

rate was 98.1% ± 0.3% [range: 96.9-98.6%]. Sequence alignment files were written into bam files,

sorted, and indexed by Samtools (version 1.9.0) and genomic reads were counted by HTSeq

(version 0.11.1) (ANDERS *et al.* 2015; MORGAN *et al.* 2019). Genes with genomic read counts of 0

or a low yield were identified and removed from the analysis by EdgeR (version 3.10) as low yield

genes provide little evidence for differential expression, followed by normalization for the

remaining expressed genes (MCCARTHY *et al.* 2012).

QTL mapping

Trait data was analyzed from offspring of the CSS backcross (n=149) (Figure 1). To generate N2

offspring for analysis, B6.A4 mice and B6.A6 mice were first separately backcrossed to B6 mice.

The resulting (B6.A4 x B6)F1 and (B6.A6 x B6)F1 offspring were then intercrossed to each other

(Figure 1A). Their offspring, which we refer to as the N2 generation, carried one chromosome 4

and one chromosome 6 that was entirely of B6 origin and one chromosome for each of these

homologues that was recombinant, partially of B6 and partially of A/J origin (Figure 1B). The

remainder of the genome, excluding chromosomes 4 and 6, is entirely from B6. Trait data was

measured and analyzed for 149 N2 offspring at 5 weeks of age. All trait data for all samples can

be found in Supplemental Table 2. Additionally, summary statistics, Student t-tests, and Wilcoxon

tests were performed in R (version 3.5.3). Gene expression was analyzed from a subset of the

backcross offspring (n=49).

QTL linkage analysis was performed using R/qtl software (v.1.44-9), freely-available statistical

software that performs single and multiple QTL mapping analyses and allows for the inclusion of

covariates (BROMAN *et al.* 2003). One-dimensional (one locus) scans for main effects and two-

dimensional (two loci) scans for epistatic effects were conducted for both the trait data and the

transcriptome data to detect and map main and epistatic effects using the R/qtl functions "scanone"

and "scantwo", respectively. As some metabolism- and blood-related traits vary by sex, the scans

were performed with and without sex as a covariate using logistic regression with the HK

algorithm (one-dimensional) or the EM algorithm (two-dimensional). Two-dimensional scans

report the interaction with the largest LOD score for each possible chromosomal interaction:

Chromosome 4 to Chromosome 4, Chromosome 6 to Chromosome 6, and Chromosome 4 to

Chromosome 6. Trait data was also performed with BMI as an additive covariate to account for

variation due to adiposity (SAMOCHA-BONET *et al.* 2008; MARGINEAN *et al.* 2019). LOD

significance thresholds were obtained from permutation tests for both additive and epistatic

models. The R/qtl permutation test permutes the phenotypes relative to the genotype data, applies

the QTL mapping method to the shuffled version of the data to obtain a set of LOD curves, and

derives the genome-wide maximum LOD score. Permutation tests for each trait were performed

with 10,000 permutations; thus, concerns of multiple testing related to the number of statistical

tests performed for each trait are accounted for by using permutation p-values (BROMAN 2009).

QTL locations were updated in the context of a multiple QTL model based on maximum likelihood

using an iterative scan with the "refineqtl" function in R/qtl (BROMAN *et al.* 2019). Intra-

chromosomal interacting QTLs regulating gene expression within 30 MB of each other were

removed as visual inspection of the data suggested a high frequency of suspected false positives,

which is also consistent with the limited mapping resolution given the study's sample size.

Significant interactions were visualized in three-dimensional space using the R packages

"circularize" (GU 2014) and "rgl" (ADLER 2019).

Data availability

221 All RNA sequencing files can be found at the NCBI gene expression omnibus. All genotype

222 data can be found in Supplemental Table 1 and all phenotype data can be found in Supplemental

223 Table 2. The complete QTL analysis pipeline can be found on protocols.io.

Results

Identification of additive QTLs for complex traits

B6.A6 and B6.A4 mice have previously been found to carry allelic variation on their respective

substituted chromosomes that non-linearly interacted with each other to regulate plasma glucose

levels. This particular strain combination demonstrated the third most statistically significant

interaction regulating glucose levels among 15 strain combinations tested, with evidence of this

interaction found in both male and female mice. Additionally, mRNA expression of 83 genes were

identified that were regulated by epistatic interactions between these two CSSs (CHEN *et al.* 2017).

Beyond the epistatic interactions regulating glucose and gene expression, previous studies also

found evidence of QTLs mapped to these chromosomes for other complex traits, including a

number of hematology-related traits that can quickly, quantitatively, and cost-effectively be

measured in hundreds of mice (LAKE J. *et al.*). Studies of hematology-related traits in the B6.A6

strain in particular identified as many or more QTLs mapped to that substituted chromosome (Chr.

6) than were found on any other chromosome. Thus, given the previous findings of epistasis

between allelic variation on Chromosomes 4 and 6, as well as the evidence for additional QTLs

regulating clinically relevant traits that could be efficiently studied in a large mapping population,

this particular strain combination was chosen for additional studies. To further map the genetic

loci that control plasma glucose levels and other traits, a total of 17 metabolism- and blood-related

traits were analyzed in N2 offspring from a modified backcross between B6.A6 and B6.A4 (Figure

1).

A single-locus QTL genome scan was performed to test for main (additive) effects in the 149 N2

mice (BROMAN *et al.* 2003). Unadjusted main effect QTL analysis identified significant QTLs for

two traits: platelet count (plt) (LOD score > 2.04, p-value < 0.05) and granulocyte % (gran%)

(LOD score > 2.02, p-value < 0.05). To test whether including sex as a covariate might affect QTL

detection, we first tested for phenotypic differences in metabolism- and blood-related traits

between male and female mice from the same crosses. There was a significant difference between

sexes for 8 of the 17 factors tested: fasted body weight, fasting glucose, length, fat pad weight,

mcv, mchc, plt, and rdwa (p < 0.05, Table S3). Thus, sex was included as a covariate to better

account for sex-specific trait differences. Sex-adjusted main effect QTLs were identified for three

traits: plt and gran% as before, plus an additional QTL for fat pad weight (Table 1). The 1-LOD

support QTL intervals for these traits were located between 38.9 - 104.8 Mb on Chr. 6 (fat pad

weight), 83.2 - 128.8 Mb on Chr. 4 (plt), and 125.0 - 146.5 Mb on Chr. 6 (gran%) (Table 1). These

three loci accounted for 28.3% (fat pad weight), 29.4% (plt), and 29.0% (gran %) of their

respective trait variances.

Main effect QTL analyses were also performed with BMI in the model to account for variation

due to adiposity. QTL analyses that included sex and BMI resulted in the same significant main

effect QTLs as sex alone. However, there was an increase in significance for the plt QTL when

BMI was added to the model (sex p value = 0.018 vs. sex and BMI p value=0.010) (Table 1),

which aligns with the previously seen association between platelet count and BMI (SAMOCHA-

BONET *et al.* 2008; MARGINEAN *et al.* 2019).

Identification of epistatic QTLs for complex traits

To test for epistatic QTL interactions, a two-dimensional (two-locus) genome scan with sex as a

covariate was performed for the metabolism- and blood-related trait data from 149 N2 mice.

Epistatic QTL pairs were identified for two of the traits: lymphocytes (ly%) and red cell

distribution width (rdw). An inter-chromosomal interaction was discovered between loci on

chromosomes 4 and 6 that regulated ly% and an intra-chromosomal interaction between two

distinct loci on chromosome 6 regulated rdw (Table 2). The 1-LOD support QTL intervals for ly%

were between 5.09 – 104.51 Mb on Chr. 4 and between 38.85 – 93.25 Mb on Chr. 6. The 1-LOD

support interval for ly% on Chr. 4 contains a second peak (Figure 2D), however the direction of

the effect on lymphocyte levels is the same for both regions, and given the overlap between the 1-

LOD support intervals the possibility remains that these seemingly distinct peaks represent one

signal. The 1-LOD support QTL intervals for rdw were between 3.49 – 16.06 Mb and 14.36 Mb -

17.72 Mb on Chr. 6 (Table 2).

For ly%, the most significant interacting SNPs within the QTL intervals were rs13477644 (Chr.

4) and rs13478739 (Chr. 6). Testing for effects of these SNP genotypes individually on ly%

revealed no marginal effect on lymphocyte count (Figure 2A, B). Accordingly, a main effect QTL

analysis of these regions individually found no evidence for a QTL at either locus (Figures 2C,

S4). However, when accounting for both SNP genotypes taken together, association with

lymphocyte count became readily apparent (Figure 2D), with the resulting QTL analysis for

interacting loci revealing a strong epistatic QTL (Figure 2E). The epistatic regulation of ly%

represents a case of negative epistasis, as each individual A/J-derived allele at rs13477644 and

rs13478739 is associated with a modest increase in ly% (Figure 2E, middle two sections relative

to the left section); however, the combination of A/J-alleles at both rs13477644 and rs13478739

(Figure 2E, far right section) together reduce ly% back towards the control levels. The context

dependent effects of this interaction are graphically illustrated in Figure 2F, as shown by the

crossing lines connecting the mean values for each genotype combination indicating suppression

epistasis. Remarkably, this epistatic interaction alone accounted for 12.1% of the variance in this

trait.

For rdw, the most significant interacting SNPs within the QTL intervals were rs13478633 (Chr. 6)

and rs13478641 (Chr. 6). Testing for effects of these SNP genotypes individually on rdw revealed

no marginal effect on red cell distribution width (Figure S1A, B) and a main effect QTL analysis

found no evidence for a QTL at either locus (Figures S1C, S4). However, when both SNP

genotypes were jointly analyzed (Figure S1D), the association with red cell distribution width was

identified (Figure S1E). The epistatic regulation of rdw again represents a case of negative

epistasis, as each A/J-derived allele at rs13478633 and rs13478641 is associated with a modest

decrease in rdw (Figure S1D, middle two sections relative to the left section); however, the

combination of A/J-alleles at both rs13478633 and rs13478641 (Fig. S1D far right section) revert

rdw back towards the control levels. The context dependent effects of this interaction are

graphically illustrated in Figure S1F. This epistatic interaction accounted for 10.7% of variance in

this trait.

Identification of QTLs that regulate hepatic gene expression

In addition to the metabolism and blood traits, we measured the liver transcriptomes in a subset

(n=49) of the offspring from the B6.A4 and B6.A6 backcross to identify additive and epistatic

QTLs regulating hepatic gene expression. Linkage analyses of the RNA-Seq data revealed 768

main effect QTLs (single locus) (Table S5). Among the main effect QTLs, there were 159 genes

within the QTL interval on either chromosome 4 or 6, and are thus cis-QTLs, whereas 609 were

located elsewhere in the genome and are therefore trans-QTLs. There were 519 epistatic QTL

pairs regulating hepatic gene expression (LOD Score > 2.44, p-value < 0.05) (Table S6). As the

R/qtl mapping software used only identifies the most significant inter-chromosomal epistatic

interaction between chromosome 4 and 6 and the most significant intra-chromosomal epistatic

interactions for each chromosome, if anything these numbers represent an underestimate of the

true number of epistatic QTL pairs, thus demonstrating the pervasive effects of epistasis in the

regulation of gene expression. In addition to the detection of widespread epistasis, the N2 cross

strategy used in this study (Figure 1) enabled higher resolution QTL mapping relative to prior CSS

studies in which the mapping resolution was limited to entire chromosomes (CHEN *et al.* 2017).

For example, previous studies mapped main effect QTLs regulating the expression of *Extl1* to

Chromosome 4, *Cadps2* to Chromosome 6, and an epistatic interaction controlling the expression

of *Tmem245* to Chromosomes 4 and 6, all of which were mapped with higher resolution with the

N2 cross (Figures S2 and S3).

The relationship between main effect QTL LOD scores and interaction QTL LOD scores shows

that a large interaction QTL LOD score does not predict a large main effect QTL LOD score, and

vice versa (Figure 3). Of the 748 unique genes with significant main effects and 510 unique genes

with significant epistatic effects, 727 genes had only main effects, 489 had only epistatic effects,

respectively, and 21 genes had both significant main and epistatic effects. Thus, there appears to

be little overlap between marginal and epistatic effector loci (Figure 3). The percent variance

explained by interaction QTLs (mean 24.5%; range 20.5% - 85.6%) was similar to that explained

by single locus QTLs (mean 28.5%; range 20.5% - 44.4%). The detection of so many epistatic

QTLs accounting for such a high proportion of the variance was not unexpected, as our previous

data suggested the widespread importance of epistasis in regulating gene expression (CHEN *et al.*

2017). Of 519 epistatic QTLs (the 510 unique genes and genes regulated by multiple eQTLs), 243

(46.8%) exhibited positive epistasis whereas 276 (53.2%) exhibited negative epistasis. That a

slight majority of QTLs exhibited negative epistasis indicates that epistasis can have a profound

effect on maintaining homeostasis, although the proportion of negative epistatic QTL pairs in this

study was not as high as previously detected using CSSs (>95%) (CHEN *et al.* 2017), However, the

current proportion of negative epistasis is more in line with other studies comparing negative and

positive interactions (SEGRE *et al.* 2005; COSTANZO *et al.* 2016; GUERRERO *et al.* 2017).

The most statistically significant epistatic QTL regulating gene expression was for *Arhgap25*, with

LOD peaks at the SNPs rs13478003 (Chr. 4) and rs13478976 (Chr. 6). As seen in the analysis of

ly%, analysis of *Arhgap25* expression associated with each individual SNP failed to reveal a main

effect of genotype (Figure 4A, B), and neither resulted in a significant QTL association (Figure

4C). However, when *Arhgap25* expression was analyzed with SNP-SNP interactions, a clear

genetic effect of the combination of rs13478003 and rs13478976 was detected (Figure 4D-F). This

interaction accounted for a remarkable 41.7% of the variance for this trait, but was completely

absent in a standard single locus association analysis (Table S6). Beyond *Arhgap25*, 518 additional

genes were identified with significant epistatic QTLs. Interaction plots of the next 5 most

significant regulated genes (after *Arhgap25*) are shown in Figure 5A-E and include *Irf2bpl*, *Gnmt*,

Rasa3, *Flnb*, and *Capn10*, all demonstrating interactions that act to maintain homeostatic levels of

gene expression (negative epistasis), rather than exacerbate strain differences in expression levels

(Figure 5). None of the interacting loci that associated with the regulation of these transcripts was

marginally associated with controlling gene expression levels.

Interacting loci on chromosomes 4 and 6 were widely distributed along each chromosome (Figure

6A). Of the 186 intra-chromosomal interactions, 37 were located on Chromosome 4 and 149 on

Chromosome 6 (Figure 6B). The remaining 333 interactions were inter-chromosomal (Figure 6C).

Of the genes whose expression was regulated by an epistatic QTL pair, 19 were located within the

QTL interval on either chromosome 4 or 6, and are thus likely controlled by cis-QTLs. This

represents a 2.0-fold enrichment of cis-QTLs based on the size of the QTL intervals and relative

to the number of cis-QTLs expected by chance (p = 0.006). The remaining 500 interacting eQTLs

were located elsewhere in the genome and are therefore trans-QTLs. Although there appeared to

be few if any QTL hotspots, the genes that were regulated by epistatic QTL pairs were enriched

for a number of biological pathways (Supplemental Table 7). Gene ontology analysis discovered

that the most significantly enriched pathways tended to be non-specific pathways related to various

metabolic processes. Other pathways identified were related to basic cellular processes including

the regulation of gene expression and signal transduction, RNA processing, and stress responses.

All this is to say that the genes regulated by epistasis collectively have broad cellular functions

and represent the most fundamental cellular biological processes.

Discussion

N2 CSS mice with allelic variation from two substituted chromosomes were used to study epistatic

interactions and map loci controlling complex traits and gene expression, as these CSSs have a

greatly simplified genetic background relative to other commonly used mapping resources in

mammalian systems. The resulting analyses identified widespread epistatic interactions, including

specific interacting loci that contributed to the heritable variation in lymphocyte percentage (ly%),

red cell distribution width (rdw), and hepatic gene expression of over 500 genes. Intra-

chromosomal interactions were detected in rdw and hepatic gene expression data that could not be

identified in studies of the B6.A4 and B6.A6 CSS strains carrying the entire nonrecombinant A/J-

derived substituted chromosomes (CHEN *et al.* 2017). That this number of traits and genes were

identified with detectable regulation by epistasis is remarkable considering that only a single time

point was examined, gene expression was only measured in the liver, and only one pairwise strain

combination of CSSs was examined. Given that ly% and rdw are associated with diagnosis and

prognosis in cancer patients as well as cardiometabolic disease, identifying a more complete

genetic model that incorporates interaction may improve our ability to predict those at greatest risk

or modulate these traits based on improved knowledge of the biological pathways that underlie

these interactions (ISEKI *et al.* 2017; ZHAO *et al.* 2017; PILLING *et al.* 2018).

Among the most significant genes whose expression is regulated by epistatic interactions, *Capn10*

is particularly interesting given its controversial association with T2D risk. It was the first T2D

susceptibility gene identified in linkage studies (HORIKAWA *et al.* 2000), with multiple subsequent

GWAS or candidate gene studies having also identified an association between variants near

CAPN10 and T2D risk and other diabetes-related phenotypes (BAIER *et al.* 2000; EVANS *et al.*

2001; OROZCO *et al.* 2014; IBRAHIM *et al.* 2015; CUI *et al.* 2016; ZHAO *et al.* 2016; BAYRAMCI *et*

al. 2017; HOU *et al.* 2017; CASTRO-MARTINEZ *et al.* 2018). However, many other studies failed to

replicate these findings (HEGELE *et al.* 2001; TSAI *et al.* 2001; KHAN *et al.* 2014; AL-SINANI *et al.*

2015; PLENGVIDHYA *et al.* 2015; ZHANG *et al.* 2019). It is tempting to speculate that one potential

reason for the inconsistent results could be an incomplete understanding of epistatic interactions

between *CAPN10* and other loci. To illustrate how this could happen, previous simulation studies

of interacting loci revealed that small differences in allelic frequencies between populations can

lead to a failure to replicate GWAS results (GREENE *et al.* 2009). Therefore, it is of note that one

study of ~1,400 T2D and control individuals found no evidence of main effects in *CAPN10*, but

did identify evidence for gene-gene interactions involving *CAPN10* influencing T2D risk (UMA

JYOTHI AND REDDY 2015). This is remarkably similar to what we have shown with ly% (Figure 3)

and *Arhgap25* expression (Figure 4), where a failure to account for interactions can mask an

otherwise strong association.

The ~500 genes whose hepatic expression was regulated by epistasis is significantly more than

were first identified in studies of the B6.A4 and B6.A6 CSS strains carrying the entire

nonrecombinant A/J-derived substituted chromosomes (CHEN *et al.* 2017). This may be, in part,

because the recombination events within chromosomes 4 and 6 decoupled intra-chromosomal

interactions in order to reveal previously undetected intra- and inter-chromosomal epistatic QTLs.

This phenomenon has been frequently observed in previous studies of CSS for a wide range of

traits, with numerous previously hidden QTLs revealed upon deconstruction of the substituted

chromosome (BUCHNER AND NADEAU 2015). The high prevalence of epistatic interactions

provides evidence for complex molecular models underlying the genetics of many quantitative

traits (GERKE *et al.* 2009; GERKE *et al.* 2010; CHOW 2016; SACKTON AND HARTL 2016). In

addition, it is of note that there is little overlap between the genes that affect gene expression via

marginal versus epistatic actions. This means that conditioning tests for epistasis based on prior

detected marginal effects, as is often done, may underestimate the importance of epistasis in

complex traits (GREENE *et al.* 2009).

Beyond the identification of specific epistatic QTLs, this study has broad implications for

understanding the genetic architecture of complex traits. For instance, among the traits where an

epistatic QTL was detected, those QTLs accounted for a considerable proportion of the variance

for that trait (11.4% for blood-related traits, 24.5% for gene expression). The gene expression

QTLs revealed a similar number of loci with comparable variance explained (interaction: mean

24.5%; range 20.5% - 85.6%, marginal: mean 28.5%; range 20.5% - 44.4%). This demonstrates

that epistatic interactions can have large effect sizes. However, it is important to note that studies

in CSSs have consistently found QTLs that explain larger proportions of the variance relative to

other mapping populations such as the hybrid mouse diversity panel and the diversity outbred

mapping populations (BUCHNER AND NADEAU 2015). The high variance accounted for by QTLs

in studies based on CSSs has been attributed in part to the limited allelic variation within each

CSS, the limited power for detecting smaller effects given study sample sizes, and the study of

genetically identical replicates within each CSS strain to improve genotype-phenotype correlations

and reduce "noise" in the analysis, although the later does not apply to these data given the unique

genotypes of each N2 offspring (Figure 1). Therefore, caution must be applied to directly

extrapolating these effect sizes to human genetic studies or even studies in other mammalian

systems. Nonetheless, within this CSS-based study, these same factors apply equally to main

effects and interaction effects, and the fact that the variance explained was similar between these

two classes of QTLs contributes to the growing body of evidence that epistatic interactions play

an important role in the genetic architecture of complex traits. This finding has broad implications

for the genetic architecture of complex traits, where most studies to date have exclusively

identified main effect QTLs. Despite this limitation, clinical applications based on these studies,

such as the use of polygenic risk scores, are beginning to be clinically evaluated with some

promising early results (TORKAMANI *et al.* 2018). Our results highlight that these models are likely

to remain severely handicapped until they account for interactions, as the individual and

cumulative effects of the interacting loci can be equivalent to the main effects, but remain

unaccounted for in the models.

While it is relatively easy to lament the absence of epistatic interactions in modeling genetic

architecture, the problem is not simply that the models do not account for interactions, but rather

that experimentally or statistically identifying such interactions has proven exceedingly difficult.

Herein lies the power of this new CSS interaction-based mapping paradigm, which enabled the

detection and higher-resolution mapping of otherwise hidden QTLs, revealing the genomic

locations of numerous novel loci that interact to control a number of complex traits and the

expression of hundreds of genes that were not found using only a marginal analysis strategy. This

was all accomplished with a relatively small number of mice for a mapping population, including

just ~150 mice for the trait data and ~50 mice for the gene expression analyses. Thus, the use of

additional CSS crosses and larger mapping populations promises to facilitate the widespread

identification of the specific genes and alleles that underlie a broad spectrum of complex traits.

These results point to the importance of searching for epistasis in as simplified a genetic context

as possible, knowing that even within the limited allelic variation of a single substituted

chromosome there is a remarkable genetic complexity contributing to phenotypic variation. As we

attempt to move towards more personalized medicine, it will no doubt require a more

comprehensive understanding of the many interacting loci throughout the genome, which should

be greatly facilitated by this CSS-based mapping paradigm.

Figure Legends

Figure 1. **Diagram of the CSS backcross strategy to map epistatic QTL pairs.** (A) (B6.A4 x

B6)F1 mice were crossed with (B6 x B6.A6)F1 mice to generate 149 "N2" offspring. (B) Seven

hypothetical "N2" offspring are shown. The recombinant chromosomes 4 and 6 are derived from

A/J or C57BL/6J as indicated. All other chromosomes are B6- derived in all mice and therefore

lack any allelic variation in this cross.

Figure 2. Epistatic interaction between loci on chromosomes 4 and 6 regulate the percentage

of lymphocyte cells in blood. Lymphocyte % based on genotype at SNP markers (A) rs13477644

on chromosome 4 and (B) rs13478739 on chromosome 6. Each dot represents one mouse. (C) QTL

mapping results for main effects on chromosomes 4 and 6. The LOD threshold for significance (p

< 0.05) was calculated by permutation testing (n=10,000) and is indicated by a horizontal line. Mb

position is indicated for both chromosomes 4 and 6 along the x-axis. No significant main effect

QTLs were detected. (D) Lymphocyte % based on the combined genotypes at rs13477644 on

chromosome 4 and rs13478739 on chromosome 6. (E) QTL mapping results for interaction effects

on chromosomes 4 and 6. The LOD threshold for significance (p < 0.05) was calculated by

permutation testing (n=10,000) and is indicated by a horizontal line. The most significant

interaction QTL pairs were detected with peaks centered at rs13477644 (17.3 Mb) and rs13478739

(22.6 Mb), with a potentially second peak on chromosome 4 centered at rs13477796 (37.6 Mb).

Mb position is indicated for both chromosomes 4 and 6 along the x-axis. (F) Context-dependent

effects of the BB and AB genotypes at markers rs13477644 and rs13478739. Mean and standard

error are shown for each genotype combination. An "A" genotype indicates A/J-derived. A "B"

genotype indicates C57BL/6J-derived.

Figure 3. Little overlap between main effect QTLs and epistatic QTL pairs. (A,B) LOD scores

for peak SNPs defining significant epistatic QTL pairs were plotted against the corresponding

main effect LOD scores for each SNP within the epistatic pair. (C) LOD scores for peak SNPs

defining significant main effect QTLs were plotted against the most significant interaction LOD

score for that SNP. The solid horizontal red line in panels A and B indicates the threshold level for

significance for epistatic QTL pairs and the vertical dashed red line in panel C indicates the

threshold level for significance for main effect QTLs.

Figure 4. Epistatic interaction between loci on chromosomes 4 and 6 regulates *Arhgap25*

mRNA expression. *Arhgap25* mRNA expression based on genotype at SNP markers (A)

rs13478003 on chromosome 4 and (B) rs13478976 on chromosome 6. Each dot represents one

mouse. (C) QTL mapping results for main effects on chromosomes 4 and 6. The LOD threshold

for significance ($p < 0.05$) was calculated using R/qtl and is indicated by a horizontal line. Mb

position is indicated for both chromosomes 4 and 6 along the x-axis. No significant QTLs were

detected. (D) *Arhgap25* mRNA expression based on the combined genotypes at rs13478003 on

chromosome 4 and rs13478976 on chromosome 6. (E) QTL mapping results for interaction effects

on chromosomes 4 and 6. The LOD threshold for significance ($p < 0.05$) was calculated by R/qtl

using 10,000 permutations and is indicated by a horizontal line. The most significant interaction

QTLs were detected with peaks at rs13478003 (69.1 Mb) and rs13478976 (52.2 Mb). Mb position

is indicated for both chromosomes 4 and 6 along the x-axis. (F) Context- dependent effects of the

BB and AB genotypes at rs13478003 and rs13478976. Mean and standard error are shown for each

genotype combination. An "A" genotype indicates A/J-derived. A "B" genotype indicates

C57BL/6J-derived.

Figure 5. **Epistatic interactions control gene expression.** Context dependent effects on gene

expression for (A) *Irf2bpl*, (B) *Gnmt*, (C) *Rasa3*, (D) *Flnb*, and (E) *Capn10*. The y-axis represents

the number of unique sequencing reads per gene after normalizing read depth across samples.

Mean and standard error are shown for each genotype combination. An "A" genotype indicates

A/J-derived. A "B" genotype indicates C57BL/6J-derived.

Figure 6. Inter-chromosomal and intra-chromosomal epistatic interactions that control gene

expression are widely distributed along chromosomes 4 and 6. (A) The circos plots illustrates

the location along chromosomes 4 and 6 for the peak SNPs for all epistatic QTL pairs regulating

gene expression. Chromosome 6 and the intra-chromosomal epistatic QTL pairs on that

chromosome (n=37) are shown in red. Chromosome 4 and the intra-chromosomal epistatic QTL

pairs on that chromosome (n=149) are shown in blue. Numbering around the circle plots indicates

the Mb position on each chromosome. (B) Inter-chromosomal epistatic QTL pairs between

chromosomes 4 and 6 (n=333) are shown in grey.

Supplemental Figure 1. Epistatic interaction between loci on chromosome 6 regulate red cell

distribution width in blood. Red cell distribution width based on genotype at SNP markers (A)

rs13478633 and (B) rs13478641 on chromosome 6. Each dot represents one mouse. (C) QTL

mapping results for main effects on chromosome 6. The LOD threshold for significance (p < 0.05)

was calculated by permutation testing (n=10,000) and is indicated by a horizontal line. Mb position

is indicated for both chromosomes 4 and 6 along the x-axis. No significant main effect QTLs were

detected. (D) Red cell distribution width based on the combined genotypes at rs13478633 and

rs13478641 on chromosome 6. (E) QTL mapping results for interaction effects on chromosome 6.

The LOD threshold for significance (p < 0.05)) was calculated by permutation testing (n=10,000)

and is indicated by a horizontal line. The most significant interaction QTLs were detected with

peaks at rs13478633 (5.2 Mb) and rs13478641 (7.2 Mb). Mb position is indicated for Chromosome

6 along the x-axis. (F) Context-dependent effects of the BB and AB genotypes at rs13478633 and

rs13478641. Mean and standard error are shown for each genotype combination. An "A" genotype

indicates A/J-derived. A "B" genotype indicates C57BL/6J-derived.

Supplemental Figure 2. Higher resolution mapping of main effect QTLs regulating the

mRNA expression of *Extl1* and *Cadps2*. The main effects for QTLs regulating the expression of

Extl1 to Chromosome 4 and *Cadps2* to Chromosome 6 are denoted by LOD peaks. In our previous

study expression of these two genes mapped the chromosome 4 and chromosome 6 in their entirety

(CHEN *et al.* 2017), but in the current study we were able to map them to a single locus on each of

the chromosomes as shown using N2 progeny. (A) The LOD score plot for the *Extl1* QTL is shown,

with the LOD score peak located at rs13477965 (126.7 Mb). (B) The LOD score plot for the

Cadps2 QTL is shown, with the LOD score peak located at rs13478695 (32.3 Mb).

Supplemental Figure 3. Higher resolution mapping of an interaction effect QTL pair

regulating the mRNA Expression of *Tmem245*. Using the N2 crosses, we mapped expression of

Tmem245 to interacting loci, one on chromosome 4 and one on chromosome 6. This provided

increased mapping resolution in comparison to our previous study where we mapped interacting

loci only to the entire chromosomes (CHEN *et al.* 2017). The LOD score plot for the *Tmem245*

interaction QTL is shown, with the LOD score peaks located at rs13477555 (10.2 Mb) on

Chromosome 4 and rs3023840 (147.7 Mb) on Chromosome 6.

References
580
Adler, D. M., D., 2019 rgl: 3D Visualization Using OpenGL. R package version 0.100.30.
https://CRAN.R-project.org/package=rgl.
Al-Sinani, S., N. Woodhouse, A. Al-Mamari, O. Al-Shafie, M. Al-Shafaee *et al.*, 2015
Association of gene variants with susceptibility to type 2 diabetes among Omanis. World
J Diabetes 6**:** 358-366.
Anders, S., P. T. Pyl and W. Huber, 2015 HTSeq--a Python framework to work with high-
throughput sequencing data. Bioinformatics 31**:** 166-169.
Andrews, S., 2010 FastQC: a quality control tool for high throughput sequence data. Available
online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc.
Baier, L. J., P. A. Permana, X. Yang, R. E. Pratley, R. L. Hanson *et al.*, 2000 A calpain-10 gene
polymorphism is associated with reduced muscle mRNA levels and insulin resistance. J
Clin Invest 106**:** R69-73.
Bayramci, N. S., L. Acik, C. Kalkan and I. Yetkin, 2017 Investigation of glucocorticoid receptor
and calpain-10 gene polymorphisms in Turkish patients with type 2 diabetes mellitus.
Turk J Med Sci 47**:** 1568-1575.
Bloom, J. S., I. M. Ehrenreich, W. T. Loo, T. L. Lite and L. Kruglyak, 2013 Finding the sources
of missing heritability in a yeast cross. Nature 494**:** 234-237.
Bolger, A. M., M. Lohse and B. Usadel, 2014 Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics 30**:** 2114-2120.
Boyle, E. A., Y. I. Li and J. K. Pritchard, 2017 An Expanded View of Complex Traits: From
Polygenic to Omnigenic. Cell 169**:** 1177-1186.
Broman, K. W., 2009 *A Guide to QTL MApping with R/qtl*. Springer Science & Business Media.
Broman, K. W., D. M. Gatti, P. Simecek, N. A. Furlotte, P. Prins *et al.*, 2019 R/qtl2: Software
for Mapping Quantitative Trait Loci with High-Dimensional Data and Multiparent
Populations. Genetics 211**:** 495-502.
Broman, K. W., H. Wu, S. Sen and G. A. Churchill, 2003 R/qtl: QTL mapping in experimental
crosses. Bioinformatics 19**:** 889-890.
Buchner, D. A., and J. H. Nadeau, 2015 Contrasting genetic architectures in different mouse
reference populations used for studying complex traits. Genome Res 25**:** 775-791.
Carlborg, O., and C. S. Haley, 2004 Epistasis: too often neglected in complex trait studies? Nat
Rev Genet 5**:** 618-625.
Castro-Martinez, A. G., J. Sanchez-Corona, A. P. Vazquez-Vargas, A. G. Garcia-Zapien, A.
Lopez-Quintero *et al.*, 2018 Association analysis of calpain 10 gene variants/haplotypes
with gestational diabetes mellitus among Mexican women. Cell Mol Biol (Noisy-le-
grand) 64**:** 81-86.
Chatterjee, S., and A. Chakravarti, 2019 A gene regulatory network explains RET-EDNRB
epistasis in Hirschsprung disease. Hum Mol Genet 28**:** 3137-3147.
Chen, A., Y. Liu, S. M. Williams, N. Morris and D. A. Buchner, 2017 Widespread epistasis
regulates glucose homeostasis and gene expression. PLoS Genet 13**:** e1007025.
Chow, C. Y., 2016 Bringing genetic background into focus. Nat Rev Genet 17**:** 63-64.
Churchill, G. A., D. M. Gatti, S. C. Munger and K. L. Svenson, 2012 The Diversity Outbred
mouse population. Mamm Genome 23**:** 713-718.
Cole, B. S., M. A. Hall, R. J. Urbanowicz, D. Gilbert-Diamond and J. H. Moore, 2017 Analysis
of Gene-Gene Interactions. Curr Protoc Hum Genet 95**:** 1 14 11-11 14 10.

Cordell, H. J., 2002 Epistasis: what it means, what it doesn't mean, and statistical methods to detect it in humans. Hum Mol Genet 11**:** 2463-2468.

Costanzo, M., B. VanderSluis, E. N. Koch, A. Baryshnikova, C. Pons *et al.*, 2016 A global genetic interaction network maps a wiring diagram of cellular function. Science 353.

Cui, J., X. Xu, S. Yin, F. Chen, P. Li *et al.*, 2016 Meta-analysis of the association between four CAPN10 gene variants and gestational diabetes mellitus. Arch Gynecol Obstet 294**:** 447-453.

Cunningham, F., P. Achuthan, W. Akanni, J. Allen, M. R. Amode *et al.*, 2019 Ensembl 2019. Nucleic Acids Res 47**:** D745-D751.

Evans, J. C., T. M. Frayling, P. G. Cassell, P. J. Saker, G. A. Hitman *et al.*, 2001 Studies of association between the gene for calpain-10 and type 2 diabetes mellitus in the United Kingdom. Am J Hum Genet 69**:** 544-552.

Forsberg, S. K., J. S. Bloom, M. J. Sadhu, L. Kruglyak and O. Carlborg, 2017 Accounting for genetic interactions improves modeling of individual quantitative trait phenotypes in yeast. Nat Genet 49**:** 497-503.

Fu, W., T. D. O'Connor and J. M. Akey, 2013 Genetic architecture of quantitative traits and complex diseases. Curr Opin Genet Dev 23**:** 678-683.

Galarza-Munoz, G., F. B. S. Briggs, I. Evsyukova, G. Schott-Lerner, E. M. Kennedy *et al.*, 2017 Human Epistatic Interaction Controls IL7R Splicing and Increases Multiple Sclerosis Risk. Cell 169**:** 72-84 e13.

Gerke, J., K. Lorenz and B. Cohen, 2009 Genetic interactions between transcription factors cause natural variation in yeast. Science 323**:** 498-501.

Gerke, J., K. Lorenz, S. Ramnarine and B. Cohen, 2010 Gene-environment interactions at nucleotide resolution. PLoS Genet 6**:** e1001144.

Greene, C. S., N. M. Penrod, S. M. Williams and J. H. Moore, 2009 Failure to replicate a genetic association may provide important clues about genetic architecture. PLoS One 4**:** e5639.

Gu, Z., 2014 circlize implements and enhances circular visualization in R. Bioinformatics. DOI: 10.1093/bioinformatics/btu393.

Guerrero, R. F., C. D. Muir, S. Josway and L. C. Moyle, 2017 Pervasive antagonistic interactions among hybrid incompatibility loci. PLoS Genet 13**:** e1006817.

Hegele, R. A., S. B. Harris, B. Zinman, A. J. Hanley and H. Cao, 2001 Absence of association of type 2 diabetes with CAPN10 and PC-1 polymorphisms in Oji-Cree. Diabetes Care 24**:** 1498-1499.

Hemani, G., K. Shakhbazov, H. J. Westra, T. Esko, A. K. Henders *et al.*, 2014 Detection and replication of epistasis influencing transcription in humans. Nature 508**:** 249-253.

Horikawa, Y., N. Oda, N. J. Cox, X. Li, M. Orho-Melander *et al.*, 2000 Genetic variation in the gene encoding calpain-10 is associated with type 2 diabetes mellitus. Nat Genet 26**:** 163-175.

Hou, Z., M. Li and Y. Cao, 2017 TCF7L2, CAPN10 polymorphisms are associated with gestational diabetes mellitus (GDM) risks: a meta-analysis. Gynecol Endocrinol 33**:** 399-404.

Hu, T., Y. Chen, J. W. Kiralis, R. L. Collins, C. Wejse *et al.*, 2013 An information-gain approach to detecting three-way epistatic interactions in genetic association studies. J Am Med Inform Assoc 20**:** 630-636.

Huang, W., and T. F. Mackay, 2016 The Genetic Architecture of Quantitative Traits Cannot Be Inferred from Variance Component Analysis. PLoS Genet 12**:** e1006421.

Huang, W., S. Richards, M. A. Carbone, D. Zhu, R. R. Anholt *et al.*, 2012 Epistasis dominates the genetic architecture of Drosophila quantitative traits. Proc Natl Acad Sci U S A 109**:** 15553-15559.

Huang, Y., C. Wang, Y. Yao, X. Zuo, S. Chen *et al.*, 2015 Molecular Basis of Gene-Gene Interaction: Cyclic Cross-Regulation of Gene Expression and Post-GWAS Gene-Gene Interaction Involved in Atrial Fibrillation. PLoS Genet 11**:** e1005393.

Ibrahim, A. T., A. Hussain, M. A. Salih, O. A. Ibrahim, S. E. Jamieson *et al.*, 2015 Candidate gene analysis supports a role for polymorphisms at TCF7L2 as risk factors for type 2 diabetes in Sudan. J Diabetes Metab Disord 15**:** 4.

Iseki, Y., M. Shibutani, K. Maeda, H. Nagahara, T. Tamura *et al.*, 2017 The impact of the preoperative peripheral lymphocyte count and lymphocyte percentage in patients with colorectal cancer. Surg Today 47**:** 743-754.

Jasnos, L., and R. Korona, 2007 Epistatic buffering of fitness loss in yeast double deletion strains. Nat Genet 39**:** 550-554.

Kayima, J., J. Liang, Y. Natanzon, J. Nankabirwa, I. Ssinabulya *et al.*, 2017 Association of genetic variation with blood pressure traits among East Africans. Clin Genet 92**:** 487-494.

Khan, I. A., S. Movva, N. A. Shaik, S. Chava, P. Jahan *et al.*, 2014 Investigation of Calpain 10 (rs2975760) gene polymorphism in Asian Indians with Gestational Diabetes Mellitus. Meta Gene 2**:** 299-306.

Kim, D., B. Langmead and S. L. Salzberg, 2015 HISAT: a fast spliced aligner with low memory requirements. Nat Methods 12**:** 357-360.

Kirino, Y., G. Bertsias, Y. Ishigatsubo, N. Mizuki, I. Tugal-Tutkun *et al.*, 2013 Genome-wide association analysis identifies new susceptibility loci for Behcet's disease and epistasis between HLA-B*51 and ERAP1. Nat Genet 45**:** 202-207.

Lake J., L. Donahue and M. Davisson, Multi-system analysis of mouse physiology, C57BL/6J-Chr#A/J/NaJ chromosome substitution panel, pp., edited by M. P. D. w. r. (RRID:SCR_003212), The Jackson Laboratory, Bar Harbon, Maine USA, https://phenome.jax.org.

Lehner, B., C. Crombie, J. Tischler, A. Fortunato and A. G. Fraser, 2006 Systematic mapping of genetic interactions in Caenorhabditis elegans identifies common modifiers of diverse signaling pathways. Nat Genet 38**:** 896-903.

Liu, Y., H. Xu, S. Chen, X. Chen, Z. Zhang *et al.*, 2011 Genome-wide interaction-based association analysis identified multiple new susceptibility Loci for common diseases. PLoS Genet 7**:** e1001338.

Mackay, T. F., 2014 Epistasis and quantitative traits: using model organisms to study gene-gene interactions. Nat Rev Genet 15**:** 22-33.

Manolio, T. A., F. S. Collins, N. J. Cox, D. B. Goldstein, L. A. Hindorff *et al.*, 2009 Finding the missing heritability of complex diseases. Nature 461**:** 747-753.

Marginean, C. O., L. E. Melit, D. V. Ghiga and M. O. Marginean, 2019 Early Inflammatory Status Related to Pediatric Obesity. Front Pediatr 7**:** 241.

McCarthy, D. J., Y. Chen and G. K. Smyth, 2012 Differential expression analysis of multifactor RNA-Seq experiments with respect to biological variation. Nucleic Acids Res 40**:** 4288-4297.

Moore, J. H., and S. M. Williams, 2009 Epistasis and its implications for personal genetics. Am J Hum Genet 85**:** 309-320.

Morgan, M., H. Pagès, V. Obenchain and N. Hayden, 2019 Rsamtools: Binary alignment
(BAM), FASTA, variant call (BCF), and tabix file import. R package version 2.0.1,
http://bioconductor.org/packages/Rsamtools.
Nadeau, J. H., J. B. Singer, A. Matin and E. S. Lander, 2000 Analysing complex genetic traits
with chromosome substitution strains. Nat Genet 24: 221-225.
Orozco, A. C., A. M. Munoz, C. M. Velasquez, R. M. Uscategui, M. V. Parra *et al.*, 2014
Variant in CAPN10 gene and environmental factors show evidence of association with
excess weight among young people in a Colombian population. Biomedica 34: 546-555.
Pilling, L. C., J. L. Atkins, G. A. Kuchel, L. Ferrucci and D. Melzer, 2018 Red cell distribution
width and common disease onsets in 240,477 healthy volunteers followed for up to 9
years. PLoS One 13: e0203504.
Plengvidhya, N., K. Chanprasert, W. Tangjittipokin, W. Thongnoppakhun and P. T.
Yenchitsomanus, 2015 Detection of CAPN10 copy number variation in Thai patients
with type 2 diabetes by denaturing high performance liquid chromatography and real-
time quantitative polymerase chain reaction. J Diabetes Investig 6: 632-639.
Sackton, T. B., and D. L. Hartl, 2016 Genotypic Context and Epistasis in Individuals and
Populations. Cell 166: 279-287.
Samocha-Bonet, D., D. Justo, O. Rogowski, N. Saar, S. Abu-Abeid *et al.*, 2008 Platelet counts
and platelet activation markers in obese subjects. Mediators Inflamm 2008: 834153.
Segre, D., A. Deluna, G. M. Church and R. Kishony, 2005 Modular epistasis in yeast
metabolism. Nat Genet 37: 77-83.
Shao, H., L. C. Burrage, D. S. Sinasac, A. E. Hill, S. R. Ernest *et al.*, 2008 Genetic architecture
of complex traits: large phenotypic effects and pervasive epistasis. Proc Natl Acad Sci U
S A 105: 19910-19914.
Soong, T. R., B. E. Howitt, A. Miron, N. S. Horowitz, F. Campbell *et al.*, 2018 Evidence for
lineage continuity between early serous proliferations (ESPs) in the Fallopian tube and
disseminated high-grade serous carcinomas. J Pathol 246: 344-351.
Torkamani, A., N. E. Wineinger and E. J. Topol, 2018 The personal and clinical utility of
polygenic risk scores. Nat Rev Genet 19: 581-590.
Tsai, H. J., G. Sun, D. E. Weeks, R. Kaushal, M. Wolujewicz *et al.*, 2001 Type 2 diabetes and
three calpain-10 gene polymorphisms in Samoans: no evidence of association. Am J Hum
Genet 69: 1236-1244.
Tyler, A., J. M. Mahoney and G. W. Carter, 2020 Genetic Interactions Affect Lung Function in
Patients with Systemic Sclerosis. G3 (Bethesda) 10: 151-163.
Tyler, A. L., B. Ji, D. M. Gatti, S. C. Munger, G. A. Churchill *et al.*, 2017 Epistatic Networks
Jointly Influence Phenotypes Related to Metabolic Disease and Gene Expression in
Diversity Outbred Mice. Genetics 206: 621-639.
Uma Jyothi, K., and B. M. Reddy, 2015 Gene-gene and gene-environment interactions in the
etiology of type 2 diabetes mellitus in the population of Hyderabad, India. Meta Gene 5:
9-20.
Verma, S. S., J. N. Cooke Bailey, A. Lucas, Y. Bradford, J. G. Linneman *et al.*, 2016 Epistatic
Gene-Based Interaction Analyses for Glaucoma in eMERGE and NEIGHBOR
Consortium. PLoS Genet 12: e1006186.
Zhang, X., C. Shi, L. Wei, F. Sun and L. Ji, 2019 The Association Between the rs2975760 and
rs3792267 Single Nucleotide Polymorphisms of Calpain 10 (CAPN10) and Gestational
Diabetes Mellitus. Med Sci Monit 25: 5137-5142.

Zhao, F., D. Mamatyusupu, Y. Wang, H. Fang, H. Wang *et al.*, 2016 The Uyghur population and
genetic susceptibility to type 2 diabetes: potential role for variants in CAPN10, APM1
and FUT6 genes. J Cell Mol Med 20**:** 2138-2147.
Zhao, W., P. Wang, H. Jia, M. Chen, X. Gu *et al.*, 2017 Lymphocyte count or percentage: which
can better predict the prognosis of advanced cancer patients following palliative care?
BMC Cancer 17**:** 514.
768

A

Chr. 4

X

Chr. 6

B

Chr. 4

Chr. 6

1   2   3   4   5   6   7

A/J-derived

C57BL/6J-derived

TABLE 1. 1-LOD SUPPORT INTERVALS FOR MAIN EFFECT TRAIT QTLS

	Chromosome	Marker	Position (Mb)	LOD	p-value	BMI LOD	BMI p-value
Fat Pad Weight	6	rs13478720	38.85	1.17		1.19	
	6	**rs13478811**	**70.31**	**2.18**	**0.030**	**2.24**	**0.030**
	6	rs13478947	104.84	1.06		1.08	
Platelets	4	rs13477808	83.18	1.56		1.55	
	4	**rs13477911**	**111.29**	**2.61**	**0.018**	**2.64**	**0.010**
	4	rs13477973	128.75	1.41		1.36	
Granulocyte (%)	6	rs3023094	125.04	1.61		1.49	
	6	**rs13479068**	**137.48**	**2.64**	**0.006**	**2.57**	**0.02**
	6	rs3090690	146.46	1.55		1.55	

*Peak SNPs are in bold with the SNPs defining the 1-LOD support interval shown above and below the peak SNP. The LOD and P-value columns include sex in the model, The BMI LOD and BMI p-value columns include sex and BMI in the model

TABLE 2. 1-LOD SUPPORT INTERVALS FOR EPISTATIC TRAIT QTLS

	Chromosome	Marker (nearest SNP)	Position (Mb)	LOD	p-value
Lymphocytes (%)					**0.032**
Interval 1					
	4	rs13477536	5.09	2.86	
	4	**rs13477644**	**35.28**	**4.23**	
	4	rs3695339	104.51	2.72	
Interval 2					
	6	rs13478720	38.85	2.29	
	6	**rs13478739**	**47.03**	**3.85**	
	6	rs13478897	93.25	1.95	
Red Cell Distribution Width					**0.036**
Interval 1					
	6	rs13459096	3.49	3.10	
	6	**rs13478633**	**12.25**	**5.69**	
	6	rs13478641	16.06	0.90	
Interval 2					
	6	rs13478638	14.36	3.70	
	6	**rs13478641**	**16.06**	**5.95**	
	6	rs3023064	17.72	3.91	

*Peak SNPs are in bold with the SNPs defining the 1-LOD support interval shown above and below both peak SNPs

TABLE S3. TRAIT DATA SUMMARY STATISTICS

Trait	Female (mean(sd))	Male (mean(sd))	T-Test	Wilcoxon Test
Fasted Weight (g)	14.09 (0.99)	16.94 (1.73)	<2.2E-16	<2.2E-16
Fasting Glucose (mg/dL)	83.58 (20.39)	98.79 (24.30)	6.26E-05	6.19E-06
Length (cm)	88.80 (1.95)	92.28 (2.73)	1.28E-15	2.04E-14
Fat Pad Weight (g)	0.043 (0.02)	0.078 (0.04)	1.54E-11	1.88E-10
White Blood Count (K/uL)	6.31 (2.20)	5.95 (2.33)	3.60E-01	3.03E-01
Lymphocytes (K/uL)	5.15 (1.62)	4.90 (1.93)	4.14E-01	2.30E-01
Monocytes (K/uL)	0.31 (0.14)	0.29 (0.14)	4.86E-01	4.89E-01
Lymphocytes (%)	83.03 (4.47)	82.28 (4.14)	3.20E-01	3.36E-01
Basophils (%)	0.12 (0.13)	0.17 (0.18)	5.42E-01	5.00E-01
Red Blood Count (M/uL)	9.32 (1.00)	9.28 (0.97)	7.88E-01	7.54E-01
Hemoglobin (g/dL)	14.79 (1.58)	14.56 (1.73)	4.30E-01	4.83E-01
Hematocrit (%)	43.63 (4.08)	44.34 (4.29)	3.31E-01	9.99E-02
Mean Corpus Volume (fL)	46.88 (1.68)	47.84 (1.69)	1.28E-03	1.81E-04
Mean Corpuscular Hemoglobin (pg)	15.85 (0.36)	15.68 (0.88)	1.59E-01	8.23E-01
Mean Corpuscular Hemoglobin Concentration (g/dL)	33.87 (1.30)	32.86 (2.46)	3.44E-03	1.73E-05
Red Cell Distribution Width (%)	19.11 (1.00)	19.24 (1.09)	4.88E-01	3.14E-01
Platelets (K/uL)	562.90 (125.60)	656.14 (152.13)	1.73E-04	6.25E-06
Mean Platelet Volume (fL)	5.19 (0.39)	5.16 (0.44)	7.06E-01	5.19E-01

Granulocyte (K/uL)	0.75 (0.38)	0.75 (0.37)	9.95E-01	7.48E-01
Granulocyte (%)	12.6 (4.06)	13.17 (3.88)	4.11E-01	1.47E-01
RDWa	33.62 (1.00)	34.77 (1.41)	2.26E-06	6.63E-06

TABLE S4. MAIN EFFECT LOD SCORES FOR THE INTERVALS CONTAINING EPISTATIC TRAIT QTLS

Trait	Chromosome	SNP	Position (Mb)	LOD	p value
Lymphocytes (%)	4	rs13477644	35.28	0.393	0.182
	6	rs13478739	47.03	9.535e-5	0.982
Red Cell Distribution Width	6	rs13478633	12.25	0.316	0.966
	6	rs13478641	16.06	0.345	0.926