MULTIOBJECTIVE OPTIMIZATION IDENTIFIES CANCER-SELECTIVE COMBINATION THERAPIES

A PREPRINT

Otto I. Pulkkinen[*]
Institute for Molecular Medicine Finland (FIMM)
University of Helsinki, FI-00290 Helsinki Finland;
Helsinki Institute for Information Technology (HIIT), Department of Computer Science
FI-00014 University of Helsinki and FI-00076 Aalto, Finland;
Computational Physics Laboratory
Tampere University, Tampere FI-33720, Finland
otto.i.pulkkinen@gmail.com

Prson Gautam
Institute for Molecular Medicine Finland (FIMM)
University of Helsinki, FI-00290 Helsinki Finland

Ville Mustonen[*][†]
Helsinki Institute for Information Technology (HIIT), Department of Computer Science
FI-00014 University of Helsinki and FI-00076 Aalto, Finland;
Organismal and Evolutionary Biology Research Programme,
Institute of Biotechnology, University of Helsinki, 00014 Helsinki, Finland
v.mustonen@helsinki.fi

Tero Aittokallio[*][†]
Institute for Molecular Medicine Finland (FIMM)
University of Helsinki, FI-00290 Helsinki Finland;
Institute for Cancer Research, Department of Cancer Genetics
Oslo University Hospital, N-0310 Oslo, Norway;
Oslo Centre for Biostatistics and Epidemiology (OCBE), Faculty of Medicine
University of Oslo, N-0317 Oslo, Norway;
Department of Mathematics and Statistics, University of Turku
Quantum, FI-20014 Turku, Finland
tero.aittokallio@helsinki.fi

April 10, 2020

ABSTRACT

Combinatorial therapies are required to treat patients with advanced cancers that have become resistant to monotherapies through rewiring of redundant pathways. Due to a massive number of potential drug combinations, there is a need for systematic approaches to identify safe and effective combinations for each patient, using cost-effective methods. Here, we developed an exact multiobjective optimization method for identifying pairwise or higher-order combinations that show maximal cancer-selectivity. The prioritization of patient-specific combinations is based on Pareto-optimization in the search

---

[*]To whom correspondence should be addressed
[†]TA and VM contributed equally to this work

space spanned by the therapeutic and nonselective effects of combinations. We demonstrate the performance of the method in the context of BRAF-V600E melanoma treatment, where the optimal solutions predicted a number of co-inhibition partners for vemurafenib, a selective BRAF- V600E inhibitor, approved for advanced melanoma. We experimentally validated many of the predictions in BRAF-V600E melanoma cell line, and the results suggest that one can improve selective inhibition of BRAF-V600E melanoma cells by combinatorial targeting of MAPK/ERK and other compensatory pathways using pairwise and third-order drug combinations. Our mechanism-agnostic optimization method is widely applicable to various cancer types, and it takes as input only measurements of a subset of pairwise drug combinations, without requiring target information or genomic profiles. Such data-driven approaches may become useful for functional precision oncology applications that go beyond the cancer genetic dependency paradigm to optimize cancer-selective combinatorial treatments.

significance statement

Cancer is diagnosed in nearly 40% of people in the U.S at some point during their lifetimes. Despite decades of research to lower cancer incidence and mortality, cancer remains a leading cause of deaths worldwide. Therefore, new targeted therapies are required to further reduce the death rates and toxic effects of treatments. Here we developed a mathematical optimization framework for finding cancer-selective treatments that optimally use drugs and their combinations. The method uses multiobjective optimization to identify drug combinations that simultaneously show maximal therapeutic potential and minimal non-selectivity, to avoid severe side effects. Our systematic search approach is applicable to various cancer types and it enables optimization of combinations involving both targeted therapies as well as standard chemotherapies.

keywords: Combination therapy | Mathematical optimization | Precision oncology

Combination therapies have become the standard of care to treat many complex diseases, including advanced cancers, which have developed resistance to monotherapies. Due to the heterogeneity of the disease, there is an increasing need to identify targeted combinatorial therapies for each cancer patient individually based on patient-derived cell models, such as ex-vivo cell cultures or patient-derived organoids and xenograft models. Systematic approaches to finding new combinatorial treatments are often based on high-throughput screening of pre-clinical models that emphasize combination efficacy and/or synergy as the key determinants of the drug combination performance [1, 2, 3, 4]. A combination is called synergistic if it has a higher than expected effect on cancer cell inhibition, where the expected combination effect is calculated based on monotherapy responses and a reference model, such as the highest single agent, Loewe additivity or Bliss independence [5].

The mechanistic basis of therapeutically beneficial combinations remain still poorly understood, but recent findings in cancer cell line screens suggest that targets of highly synergistic drug combinations are likely to interact at the level of signaling pathways, either targeting the same or functionally opposite pathways, in such a way that the combined effect is greater than the sum of the individual effects [6]. This can occur for example by combinatorial inhibition of two orthogonal pathways or by targeting compensatory mechanisms via pathway cross-talks [7]. However, the same combination mechanism may affect not only the cancerous cells, but manifest itself also as unintended adverse effects in healthy cells [8]. In this case, the combination is nonselective against the particular cancer cells, and may result in a degree of toxic effects as well. Therefore, combination synergy and expected efficacy alone are not sufficient determinants when designing safe and effective treatment options.

Identifying cancer-selective drug combinations amounts to maximizing combination efficacy in the cancer cells, while minimizing potential toxic effects to healthy cells. Such drug combination design can be considered as a multiobjective optimization problem (MOP), in which the two goals are pursued simultaneously [9, 10]. Even if multiobjective optimization has been used as a tool in planning radiotherapies [11, 12], to our knowledge, the study of Matlock et al. [13] is the only attempt to apply the mathematical optimization framework to designing drug combinations. They proposed numerical algorithms that converge to most effective and least toxic combinations of drug targets based on so-called target inhibition maps [14]. Even though such target-centric approach provides clues about the targets to inhibit or avoid in a given cell context, it cannot optimize the drugs to be used in the combination, and is not applicable to non-targeted therapies, such as standard chemotherapies or emerging immunotherapies.

In this work, we develop a novel drug-centric mathematical optimization framework for maximizing drug combination efficacy and minimizing its nonselectivity against a given cancer type. Nonselectivity is defined through the average effect of a drug compound across a large number of cells from various cancer types, and it is used as a surrogate of toxic effects. This definition also makes it possible to apply the framework using drug sensitivity measurements in the cancer cells alone, without requiring measurements on healthy cells, nor information on targets of the drugs or genomic profiles of the cancer cells. To exemplify the framework, we use pairwise combination measurements of 104

drug compounds in 60 cancer cell lines from the NCI-ALMANAC resource [15], and subsequently experimentally validate the most interesting predictions in BRAF-V600E melanoma cell line. We propose a new quantity of selective synergy, Bliss multiexcess, a multiobjective generalization of the traditional Bliss excess measure.

Results

Joint modeling of therapeutic and nonselective effects

The reduction in the rate of growth of a cell line by a drug compound is typically measured as a growth fraction Q, the relative number of live cells with respect to untreated control after a certain time (e.g., inhibition % after 72h of treatment). *The therapeutic effect E of a drug i at concentration c_i, or a combination of a pair of drugs i, j at concentrations c_i, c_j, on a target cell line l is here defined as the negative logarithm of growth fraction Q,*

$$
\begin{align}
E_i(c_i; l) &= -\log Q_i(c_i; l) \tag{1} \\
E_{ij}(c_i, c_j; l) &= -\log Q_{ij}(c_i, c_j; l). \tag{2}
\end{align}
$$

Hence, the higher the therapeutic effect, the less viable the cells are after the treatment. One advantage of the logarithmic formulation is that the effect is additive, in that the predicted effect of two drugs acting independently, i.e. without interaction in the sense of Bliss independence [16], is given by the sum $E_i + E_j$ [17, 18]. We assume that higher order combination measurements are not available, so all modeling needs to rely on the measured effects of single drugs and pairs of drugs [19, 20]. To this end, we define the therapeutic effect of a combination $\boldsymbol{c} = (c_i)_{i \in \mathcal{D}}$ from a set $\mathcal{D}$ of N possible drugs according to what is known as a regression model [21, 22]:

$$
E(\boldsymbol{c}; l) = \underbrace{\sum_{i=1}^{N} E_i(c_i; l)}_{=E_B(\boldsymbol{c};l)} + \underbrace{\sum_{j=1}^{N-1} \sum_{k=j+1}^{N} E_{j,k}^{XS}(c_j, c_k; l)}_{=E_B^{XS}(\boldsymbol{c};l)}, \tag{3}
$$

where $E_{ij}^{XS}(c_i, c_j; l) = E_{ij}(c_i, c_j; l) - E_i(c_i; l) - E_j(c_j; l)$ is the effect excess of the pair over Bliss independence model, or *Bliss excess* for short. The first sum in [3] yields the Bliss model effect $E_B(\boldsymbol{c}; l)$ of the combination $\boldsymbol{c}$, and the sum over pairs of drugs is its Bliss excess $E_B^{XS}(\boldsymbol{c}; l)$.

We model the nonselectivity of a drug compound by the mean effect over the set of all cell lines $\mathcal{L}$ in the data,

$$
\overline{E}(\boldsymbol{c}) = \frac{1}{|\mathcal{L}|} \sum_{l \in \mathcal{L}} E(\boldsymbol{c}; l). \tag{4}
$$

For a large number of patient-derived samples from various tissue types, the average in Eq. [4] comprises many somatically different cell lines. Hence, we can consider it as a proxy for the degree of nonselective toxic effects on an average (cancerous or non-cancerous) tissue. As a drug combination should not benefit from additional drugs with hardly no effect, we add to each $\overline{E}_i$ a small regularizing parameter δ (we use the value 0.1) that counterbalances any arbitrarily small increase in therapeutic effect or decrease in the mean effect. We hence define the *nonselective effect* of a drug combination with $m(\boldsymbol{c})$ components as

$$
\overline{E}_\delta(\boldsymbol{c}) = \delta \cdot m(\boldsymbol{c}) + \overline{E}(\boldsymbol{c}). \tag{5}
$$

In the spirit of Eq. [3], the nonselective effect too can be split into nonselective effects $\overline{E}_i + \delta$ of isolated compounds, and mean excesses $\overline{E}_{ij}^{XS}$, which stem from the drug interactions.

To find optimal combination treatments of at most M drugs for a cancer cell line l that simultaneously maximize the therapeutic effect and minimize the nonselective effect is a multi-objective optimization problem (MOP) [9, 10]. Formulated using the so-called ϵ-constraint method, the task is to find the set of concentration vectors

$$
C(M; l) = \bigcup_{\Theta \in \mathbb{R}_+} \left\{ \operatorname{argmax} E(\boldsymbol{c}; l) : \overline{E}_\delta(\boldsymbol{c}) \leq \Theta, m(\boldsymbol{c}) \leq M \right\}. \tag{6}
$$

Here, Θ denotes the tolerance to nonselective effect, and a vector of drug concentrations that maximizes the therapeutic effect corresponds to each Θ. The set C of solutions of this continuous MOP traces for each M a *Pareto front*, a curve in the plane of nonselective and therapeutic effects, the points of which are called *Pareto optimal*. For each point on the Pareto front, no combination of drug concentrations with lower nonselective effect and higher therapeutic effect exists when M is fixed.

The fact that the concentrations $\boldsymbol{c}$ can be adjusted on a continuous scale means that the number of solutions at the Pareto front is infinite. The Pareto front is therefore a continuous curve in the plane of therapeutic and nonselective effects. Solving this continuum MOP is obviously a numerically demanding task, and a successful solution very much depends on availability of reliable combination measurements at a multitude of concentrations of all drug compounds, so that a faithful continuous approximation to the effect in the plane of concentrations can be found for each pair of drugs. To lower the risk of false predictions due to measurement error and/or insufficient data resolution, we inspect a discrete set of concentrations instead. The two-drug combination effects are typically quantified by n-by-n square matrices (*e.g.* $n = 4, 8$, see Fig. 1) with $n - 1$ nonzero concentrations for both drugs in a combination. However, these concentrations are not necessarily the same for a single drug in each of those combination measurements, which complicates the analysis. Here we choose to reduce the problem further: We use binary variables $N_i = 1$ or 0, to determine whether a drug compound $i \in \mathcal{D}$ is present in the combination or not, respectively. The effect of the whole drug combination is then predicted by the regression model to be

$$E(\boldsymbol{c}; l) = \sum_{j=1}^{N} \varepsilon_i(l) N_i + \sum_{j=1}^{N-1} \sum_{k=j+1}^{N} \varepsilon_{jk}^{XS}(l) N_j N_k \ , \tag{7}$$

where

$$\varepsilon_i(l) = \max_{c_i > 0} E_i(c_i; l) \tag{8}$$

is the strongest effect of drug i computed from a fitted Hill curve to growth fractions Q_i (see Fig. 1 and SI), and, with $(c_i^*, c_j^*) = \mathrm{argmax}_{c_i, c_j > 0} E_{ij}(c_i, c_j; l)$,

$$\varepsilon_{ij}^{XS}(l) = E_{ij}(c_i^*, c_j^*; l) - E_{ij}(c_i^*, 0; l) - E_{ij}(0, c_j^*; l) \tag{9}$$

is the Bliss excess at the maximal measured therapeutic effect of the combination i, j. This way of defining the parameters of the binary problem was found to constitute a robust model. In particular, as high-throughput drug screens often show variability within and between plates on which the cells are drugged, the strongest effect from a fitted Hill curve for each cell line yields a reliable single number for the therapeutic effect of a drug compound in monotherapy. Furthermore, the way the pairwise interaction constants ε_{ij}^{XS} are computed per matrix from measurements on the same plate, and not against the fitted Hill curves, respects plate-to-plate variance and significantly reduces the the risk of misestimating the drug interaction strengths.

In the binary model, the nonselective effect is again given by Eq. [5] with the mean effect over all cell lines computed using the binary effect function. The notions of Bliss independence and Bliss excess of the whole combination carry over to the new definition in a natural way.

A consequence of dimensional reduction through Eqs. [7 – 9] is that the concentration of a drug in combination with another drug gets fixed, and a different concentration may get chosen for the same drug in another pairwise combination. The combination triplet spanned by these two pairs would then paradoxically contain a single drug in two different concentrations, which may lead to overestimation of the combinatorial therapeutic effect. However, should this occur, the nonselective effect is likewise affected, and the MOP is stable against these changes.

We consider our solution to MOP *a posteriori* method [9], in which a human decision maker decides, which of the Pareto optimal points provided by the algorithm are clinically relevant and worth testing further in the laboratory. For example, the combinations taken into consideration should not be much more nonselective than broadly toxic standard chemotherapies. Even if this condition is controlled by the nonselective effect, the set of Pareto optimal or nearly optimal combinations may become large. Therefore ways of further delineating the benefits achieved by a particular combination in comparison to other Pareto optimal solutions are needed.

Therapeutic dominance

In multiobjective optimization, a vector is said to dominate another vector if it performs at least equally well as the other one in all features, *and* it outperforms the other vector at least in one feature [23]. Inspired by this concept, we define the *therapeutic dominance* of a Pareto optimal combination with M components and nonselective effect $\overline{E}_\delta$ as the difference of its therapeutic effect and that of the Pareto optimal combination with $M - 1$ components and the highest nonselective effect not exceeding $\overline{E}_\delta$. This directly quantifies the therapeutic advance gained by adding a drug to a Pareto optimal combination, or achieved by a monotherapy in comparison to no therapy at all.

Selective synergy

A drug combination can be Pareto optimal for several reasons. First, the individual therapeutic effects of the drug components may be strong enough to bring the combination to the Pareto front without any synergistic interaction. In

this case, the Bliss excess $E_B^{XS}(\boldsymbol{c}; l)$ of the combination is small. Second, the combination may exhibit a strong effect due to synergistic interaction of the drugs. However, synergism occurs in a large number of cell lines, leading to a comparable but not excessive increase in nonselective effect. This scenario is characterized by a higher-than-average Bliss excess, and a mean Bliss excess $\overline{E}_B^{XS}(\boldsymbol{c})$ of the same order. Third, the combination may show high *selective synergy*, in that the drug interaction is manifested as an increase in therapeutic effect, and only little, or even as a negative change, in the mean effect over cell lines. Selective synergy is evidenced by high Bliss excess and low mean Bliss excess.

The degree of selective synergy of a combination can be boiled down to a single number, *Bliss multiexcess*

$$\mathcal{E}_B^{XS}(\boldsymbol{c}, \phi; l) = E_B^{XS}(\boldsymbol{c}; l) - \phi \overline{E}_B^{XS}(\boldsymbol{c}),\tag{10}$$

where ϕ weighs the importance of adverse effects. A motivation for definition [10], and a systematic way of choosing the value of ϕ comes from the scalarized (*i.e.* projected to one dimension), convex optimization problem corresponding to our MOP: Bliss multiexcess [10] equals the interaction part of its objective function [23]

$$\mathcal{E}(\boldsymbol{c}, \phi; l) = E(\boldsymbol{c}; l) - \phi \overline{E}_\delta(\boldsymbol{c}),\tag{11}$$

which is to be maximized. Thus the combinations that are scalar-optimal for synergistic reasons have high Bliss multiexcess. This argument can be extended to combinations outside the convex hull by appropriately choosing the weighing parameter ϕ. As the Pareto optimal combinations near the convex hull can be seen as optional solutions to the scalarized problem, we use a single value of ϕ for all Pareto optimal combinations between two convex solutions—the smallest ϕ that yields the higher effect combination as an optimal solution of the scalarized problem.

Application to melanoma

We applied the optimization framework to designing new combination therapies for malignant melanoma. The optimization is based on the sensitivity data of 104 drug compounds and their pairwise combinations in the NCI-60 cell lines available in the NCI ALMANAC resource [24]. In this data, the drug-induced growth inhibition is tested in four-by-four matrices, as depicted in Fig. 1 A. Hierarchical clustering of drug compounds and the cell lines based on monotherapy sensitivities reveals high variation across the cell lines (Fig. 1 B). In particular, vemurafenib shows a strong effect in most of the melanoma cell lines, whereas in other cancer types the effect is less potent. This is expected, as vemurafenib is a selective inhibitor of V600E-mutant BRAF, and this mutation is present in approximately 60% of melanomas. Vemurafenib has become one of the main treatments for late-stage melanoma since its discovery in 2008 and consequent FDA approval in 2011 [25, 26, 27]. Despite the initial efficacy of vemurafenib in many patients, the prognoses still remain poor as many patients develop resistance to vemurafenib, often within 6 to 8 months from therapy initiation, due to reactivation of the MAPK/ERK and other pathways. This has led to interest in combinatorial inhibition of the MAPK/ERK pathway components.

Fig. 1 C shows the solution of the MOP for all 104 drug compounds in the MALME-3M melanoma cell line with the BRAF-V600E mutation. The Pareto front traces out the optimal combinations in the plane of nonselective versus therapeutic effects. Vemurafenib stands out as a highly selective and effective treatment, as expected. Vitamin A derivative tretinoin has a lower nonselective effect than vemurafenib, but its therapeutic effect is much smaller. On the other hand, gefitinib, an EGFR inhibitor upstream of BRAF in the MAPK/ERK signaling pathway, has slightly smaller therapeutic effect than vemurafenib but it is more nonselective. Notably, the combination of vemurafenib and gefitinib (Fig. 1 C, top) shows an equally strong therapeutic effect as the most potent monotherapy, mithramycin (plicamycin), and the combination is synergistic, in that the effect is higher than that predicted by the Bliss independence model.

The solutions of the MOPs for combinations of up to two drugs (Fig. 2 A) and higher-order combinations (Fig. 2 B) confirm that vemurafenib-gefitinib combination is Pareto optimal regardless of the number of components allowed in combinations. Other Pareto optimal combinations involve vemurafenib with anastrozole, tretinoin, thalidomide and sirolimus as partners. Of these, the anti-inflammatory drug thalidomide and the aromatase inhibitor anastrozole have little effect when used alone, indicating a true synergistic interaction with vemurafenib. The same compounds appear in the Pareto optimal combinations of three and higher order in Fig. 2 B. With a few exceptions, Pareto optimal combinations consist mostly of monotherapies and duplets at smaller therapeutic and nonselective effects than vemurafenib-gefitinib. A cloud of triplets that all include vemurafenib and gefitinib as a pairwise subcombination dominates at higher effects in Fig. 2 B.

Therapeutic dominance and the measure of selective synergy were used to further investigate the selected combinations (Fig 3). Vemurafenib monotherapy, along with vemurafenib-gefitinib and vemurafenib-gefitinib-sirolimus combinations are clearly the most dominant therapies, since they yield the most therapeutic advance in comparison to a lower order Pareto optimal combinations. However, such advance can be associated with a significant increase in nonselective

Table 1: NCI ALMANAC and FIMM drug assays on MALME-3M.

Treatment	E_{NCI}	$E_{B,NCI}^{XS}$	E_{FIMM}	$E_{B,FIMM}^{XS}$
vemurafenib	1.60	-	2.10(10)	-
gefitinib	1.13	-	1.39(6)	-
anastrozole	0.02	-	0.02(1)	-
vismodegib	0.04	-	0.03(1)	-
vem+gef	2.99	0.26	3.57	0.08
vem+ana	1.91	0.29	2.33	0.21
vem+vis	1.53	-0.11	2.41	0.28
gef+ana	1.57	0.42	1.73	0.32
gef+vis	2.13	0.96	2.15	0.73
vem+gef+ana(1 μM)	3.73	0.98	3.74	0.23
vem+gef+vis(1 μM)	3.89	1.12	3.73	0.21

E_{NCI}: Therapeutic effect predicted by the binary regression model based on NCI ALMANAC data.
E_{FIMM}: Therapeutic effect measured at FIMM. The figure in parenthesis is the error of the last digit from fitting the Hill curve.
E_B^{XS}: Bliss excesses computed either from the first or the third column.

effect in the case of combinations. For instance, the duplet of vemurafenib and gefitinib is synergistic, but not selectively synergistic. The vemurafenib duplets with anastrozole, thalidomide, tretinoin and sirolimus, and the triplets of vemurafenib, gefitinib and anastrozole, or sirolimus are truly selectively synergistic combinations with significant therapeutic dominance over their lower order combinations.

In-house validations

To further test the most interesting predictions from the MOP solution, we experimentally validated in-house two Pareto optimal drug triplets and their subcombinations in the BRAF-V600E melanoma cell line MALME-3M. This validation experiment was performed to test the accuracy of the predicted therapeutic effect in other experimental assay (FIMM) than that used for model predictions (NCI-60). The in-house drug combination assay was performed in 8x8 matrix format, similar to previous studies (25) (see Materials and Methods for details).

Table 1 shows the maximal effect over the measured concentrations in both assays. In accord to predictions, the triplet combinations tested in-house (E_{FIMM}) show stronger effect than any of the duplets, and the observed therapeutic effects of triplets are very close to the predicted values (E_{NCI}). The vemurafenib monotherapy and its combinations with gefitinib had stronger effects in FIMM assay than those reported in NCI ALMANAC. This makes the Bliss excesses smaller in the in-house assays. However, as the concentration of the third component was fixed in the in-house assays, we lack the full three-dimensional optimization over concentrations, and hence the most synergistic vectors may have been missed.

As the pairwise matrices in our in-house experiments have higher resolution (8x8), than in the NCI ALMANAC assays (4x4), we were able to observe interesting nonlinearities in the growth inhibition patterns. For example, the combination of vemurafenib and gefitinib shows antagonistic pattern at high concentrations of vemurafenib and low concentrations of gefitinib, while it results in a synergistic effect at high concentrations of both components (see SI). Further, the experiments revealed a highly synergistic combination between gefitinib and vismodegib, a SMO inhibitor of the hedgehog pathway, and this combination scores high also in our MOP solution because of this synergy along the high additive effect and nonselective synergy of vemurafenib and gefitinib.

Discussion

We introduced a multiobjective optimization framework for designing safe and effective drug combinations by simultaneous optimization of therapeutic and nonselective effects. The MOP approach requires only the measurements of single drugs and a subset of their pairwise combinations, and it enables optimization of also higher-order combinatorial treatments. The nonselectivity was modeled here by the mean effect over all the cell lines in the NCI ALMANAC data set, yet any other control data can be used for the purpose of estimating toxic side effects. The exact MOP solutions from Pareto-optimization provide the decision maker with a targeted set of patient-specific combinations, illustrated and quantified by their therapeutic and nonselective effects. We further defined Bliss multiexcess, a multiobjective generalization of the traditional Bliss excess, as a new measure of selective synergy.

Our results confirmed the previously suggested combinatorial inhibition of the MAPK/ERK pathway as an effective and safe treatment for malignant melanoma with BRAF V600E mutation. However, we note that the approved combination of vemurafenib with cobimetinib [28, 29], or with any MEK inhibitor, was not tested in NCI ALMANAC, but our mechanistic-agnostic model pairs the EGFR inhibitor gefitinib with vemurafenib instead. Several higher order combinations were also found to be Pareto optimal. Of these, an aromatase inhibitor anastrozole is typically used to treat estrogen-responsive breast cancer, and it is therefore quite surprising to see it selective for melanoma. On the other hand, already the unsupervised hierarchical clustering of drugs without any interaction information paired vemurafenib with estramustine, an estrogen receptor binder. There is increasing evidence that hormone levels play a role in growth and progression of melanoma, although the underlying conditions are still unclear [30]. These examples demonstrate how our MOP solutions identify among a massive number of potential combinations those with optimal efficacy and safety profiles for further pre-clinical and clinical testing.

1 Materials and Methods

Cell line

Melanoma cell line Malme-3M (ATCC®HTB-64™) was purchased from ATCC. It was maintained at 37C with 5% CO2 in a humidified incubator in IMDM (Cat#12440046) supplemented with 20% fetal bovine serum (Cat#10270-106) and 1% Penicillin-Streptomycin (Cat#15140-122). All the reagents were purchased from ThermoFisher Scientific for the in-house assays.

Drug combination assays

The drug combination screening approach described previously [31] was adopted on MALME-3M cell line. Seven different concentrations in log3-fold dilution ranging from $10\,\text{nM}$ to $10\,000\,\text{nM}$ of vemurafenib and gefitinib, vemurafenib and anastrozole, and gefitinib and anastrozole were combined with each other in 8x8 matrix formats in duplicate. For the triple combination, $500\,\text{nM}$ of vemurafenib or $500\,\text{nM}$ gefitinib or $1000\,\text{nM}$ of anastrozole was added to the alternate drug combination matrix (in one copy). Exactly as described above, another set of combination was prepared for vemurafenib, gefitinib and vismodegib. For triple combination, $1000\,\text{nM}$ of vismodegib was used whereas the concentrations of vemurafenib and gefitinib were the same as mentioned above (Supplementary file FILE.XLSX). The compounds were plated to black clear bottom 384-well plates (Corning #3764) using an Echo 550 Liquid Handler (Labcyte). $100\,\mu\text{M}$ benzethonium chloride (BzCl2) and 0.1% dimethyl sulfoxide (DMSO) were used as positive and negative controls respectively. All subsequent liquid handling was performed using MultiFlo FX multi-mode dispenser (BioTek). The pre-dispensed compounds were dissolved in $5\,\mu\text{l}$ of culture media containing cytotoxicity measurement reagents CellTox Green (Promega) and left in plate shaker at RT for 30 min. Twenty microliter cell suspension (75000cells/ ml) was dispensed in the drugged plates. After 72 h incubation, cytotoxicity (fluorescence, 485/520 nm excitation/emission filters) was recorded using PheraStar plate reader (BMG Labtech). Subsequently, $25\,\mu\text{l}$ per well of CellTiter-Glo (Promega) reagent was added, and after 10 min of incubation at room temperature, luminescence (cell viability) was measured using PheraStar plate reader (BMG Labtech).

Code Availability

A Jupyter notebook for multiobjective optimization of drug sensitivity data and for producing the figures in this article is available at http://tiny.cc/combimop

2 Acknowledgements

The authors thank Kaisa Miettinen (University of Jyväskylä) for comments concerning multiobjective optimization, Alexander Ianevski and Anil Kumar (FIMM) for sharing their Excel spreadsheet of the NCI ALMANAC data, and Laura Turunen (FIMM HTB unit) for the drugs plates in the in-house experiments. This work was supported by the Academy of Finland [grants 292611, 310507, 313267, 326238 to TA; grant 313270 to VM], the Cancer Society of Finland [TA] and the Sigrid Jusélius Foundation [TA].

References

[1] Liye He, Evgeny Kulesskiy, Jani Saarela, Laura Turunen, Krister Wennerberg, Tero Aittokallio, and Jing Tang. Methods for high-throughput drug combination screening and synergy scoring. *Methods in molecular biology (Clifton, N.J.)*, 1711:351–398, 2018. 29344898[pmid].

[2] Alina Malyutina, Muntasir Mamun Majumder, Wenyu Wang, Alberto Pessia, Caroline A. Heckman, and Jing Tang. Drug combination sensitivity scoring facilitates the discovery of synergistic and efficacious drug combinations in cancer. *PLOS Computational Biology*, 15(5):1–19, 05 2019.

[3] Bissan Al-Lazikani, Udai Banerji, and Paul Workman. Combinatorial drug therapy for cancer in the post-genomic era. *Nature Biotechnology*, 30(7):679–692, 2012.

[4] Anna H.C. Vlot, Nata Aniceto, Michael P. Menden, Gudrun Ulrich-Merzenich, and Andreas Bender. Applying synergy metrics to combination screening data: agreements, disagreements and pitfalls. *Drug Discovery Today*, 24(12):2286 – 2298, 2019.

[5] Jing Tang, Krister Wennerberg, and Tero Aittokallio. What is synergy? the Saariselkä agreement revisited. *Frontiers in Pharmacology*, 6:181, 2015.

[6] Michael P. Menden, Dennis Wang, Mike J. Mason, Bence Szalai, Krishna C. Bulusu, Yuanfang Guan, Thomas Yu, Jaewoo Kang, Minji Jeon, Russ Wolfinger, Tin Nguyen, Mikhail Zaslavskiy, Jordi Abante, Barbara Schmitz Abecassis, Nanne Aben, Delasa Aghamirzaie, Tero Aittokallio, Farida S. Akhtari, Bissan Al-lazikani, Tanvir Alam, Amin Allam, Chad Allen, Mariana Pelicano de Almeida, Doaa Altarawy, Vinicius Alves, Alicia Amadoz, Benedict Anchang, Albert A. Antolin, Jeremy R. Ash, Victoria Romeo Aznar, Wail Ba-alawi, Moeen Bagheri, Vladimir Bajic, Gordon Ball, Pedro J. Ballester, Delora Baptista, Christopher Bare, Mathilde Bateson, Andreas Bender, Denis Bertrand, Bhagya Wijayawardena, Keith A. Boroevich, Evert Bosdriesz, Salim Bougouffa, Gergana Bounova, Thomas Brouwer, Barbara Bryant, Manuel Calaza, Alberto Calderone, Stefano Calza, Stephen Capuzzi, Jose Carbonell-Caballero, Daniel Carlin, Hannah Carter, Luisa Castagnoli, Remzi Celebi, Gianni Cesareni, Hyeokyoon Chang, Guocai Chen, Haoran Chen, Huiyuan Chen, Lijun Cheng, Ariel Chernomoretz, Davide Chicco, Kwang-Hyun Cho, Sunghwan Cho, Daeseon Choi, Jaejoon Choi, Kwanghun Choi, Minsoo Choi, Martine De Cock, Elizabeth Coker, Isidro Cortes-Ciriano, Miklós Cserzö, Cankut Cubuk, Christina Curtis, Dries Van Daele, Cuong C. Dang, Tjeerd Dijkstra, Joaquin Dopazo, Sorin Draghici, Anastasios Drosou, Michel Dumontier, Friederike Ehrhart, Fatma-Elzahraa Eid, Mahmoud ElHefnawi, Haitham Elmarakeby, Bo van Engelen, Hatice Billur Engin, Iwan de Esch, Chris Evelo, Andre O. Falcao, Sherif Farag, Carlos Fernandez-Lozano, Kathleen Fisch, Asmund Flobak, Chiara Fornari, Amir B. K. Foroushani, Donatien Chedom Fotso, Denis Fourches, Stephen Friend, Arnoldo Frigessi, Feng Gao, Xiaoting Gao, Jeffrey M. Gerold, Pierre Gestraud, Samik Ghosh, Jussi Gillberg, Antonia Godoy-Lorite, Lizzy Godynyuk, Adam Godzik, Anna Goldenberg, David Gomez-Cabrero, Mehmet Gonen, Chris de Graaf, Harry Gray, Maxim Grechkin, Roger Guimera, Emre Guney, Benjamin Haibe-Kains, Younghyun Han, Takeshi Hase, Di He, Liye He, Lenwood S. Heath, Kristoffer H. Hellton, Manuela Helmer-Citterich, Marta R. Hidalgo, Daniel Hidru, Steven M. Hill, Sepp Hochreiter, Seungpyo Hong, Eivind Hovig, Ya-Chih Hsueh, Zhiyuan Hu, Justin K. Huang, R. Stephanie Huang, László Hunyady, Jinseub Hwang, Tae Hyun Hwang, Woochang Hwang, Yongdeuk Hwang, Olexandr Isayev, Oliver Bear Don't Walk, John Jack, Samad Jahandideh, Jiadong Ji, Yousang Jo, Piotr J. Kamola, Georgi K. Kanev, Loukia Karacosta, Mostafa Karimi, Samuel Kaski, Marat Kazanov, Abdullah M. Khamis, Suleiman Ali Khan, Narsis A. Kiani, Allen Kim, Jinhan Kim, Juntae Kim, Kiseong Kim, Kyung Kim, Sunkyu Kim, Yongsoo Kim, Yunseong Kim, Paul D. W. Kirk, Hiroaki Kitano, Gunter Klambauer, David Knowles, Melissa Ko, Alvaro Kohn-Luque, Albert J. Kooistra, Melaine A. Kuenemann, Martin Kuiper, Christoph Kurz, Mijin Kwon, Twan van Laarhoven, Astrid Laegreid, Simone Lederer, Heewon Lee, Jeon Lee, Yun Woo Lee, Eemeli Lepp_aho, Richard Lewis, Jing Li, Lang Li, James Liley, Weng Khong Lim, Chieh Lin, Yiyi Liu, Yosvany Lopez, Joshua Low, Artem Lysenko, Daniel Machado, Neel Madhukar, Dries De Maeyer, Ana Belen Malpartida, Hiroshi Mamitsuka, Francesco Marabita, Kathleen Marchal, Pekka Marttinen, Daniel Mason, Alireza Mazaheri, Arfa Mehmood, Ali Mehreen, Magali Michaut, Ryan A. Miller, Costas Mitsopoulos, Dezso Modos, Marijke Van Moerbeke, Keagan Moo, Alison Motsinger-Reif, Rajiv Movva, Sebastian Muraru, Eugene Muratov, Mushthofa Mushthofa, Niranjan Nagarajan, Sigve Nakken, Aritro Nath, Pierre Neuvial, Richard Newton, Zheng Ning, Carlos De Niz, Baldo Oliva, Catharina Olsen, Antonio Palmeri, Bhawan Panesar, Stavros Papadopoulos, Jaesub Park, Seonyeong Park, Sungjoon Park, Yudi Pawitan, Daniele Peluso, Sriram Pendyala, Jian Peng, Livia Perfetto, Stefano Pirro, Sylvia Plevritis, Regina Politi, Hoifung Poon, Eduard Porta, Isak Prellner, Kristina Preuer, Miguel Angel Pujana, Ricardo Ramnarine, John E. Reid, Fabien Reyal, Sylvia Richardson, Camir Ricketts, Linda Rieswijk, Miguel Rocha, Carmen Rodriguez-Gonzalvez, Kyle Roell, Daniel Rotroff, Julian R. de Ruiter, Ploy Rukawa, Benjamin Sadacca, Zhaleh Safikhani, Fita Safitri, Marta Sales-Pardo, Sebastian Sauer, Moritz Schlichting, Jose A. Seoane, Jordi Serra, Ming-Mei Shang, Alok Sharma, Hari Sharma, Yang Shen, Motoki Shiga, Moonshik Shin, Ziv Shkedy, Kevin Shopsowitz, Sam Sinai, Dylan Skola, Petr Smirnov, Izel Fourie Soerensen, Peter Soerensen, Je-Hoon Song, Sang Ok Song, Othman Soufan, Andreas Spitzmueller, Boris Steipe, Chayaporn Suphavilai, Sergio Pulido Tamayo, David Tamborero, Jing Tang, Zia-ur-Rehman Tanoli, Marc Tarres-Deulofeu, Jesper Tegner, Liv Thommesen, Seyed Ali Madani Tonekaboni, Hong Tran, Ewoud De Troyer, Amy Truong, Tatsuhiko Tsunoda, Gábor Turu, Guang-Yo Tzeng, Lieven Verbeke, Santiago Videla, and AstraZeneca-Sanger Drug Combination DREAM Consortium. Community assessment to advance computational prediction of cancer drug combinations in a pharmacogenomic screen. *Nature Communications*, 10(1):2674, 2019.

[7] Samira Jaeger, Ana Igea, Rodrigo Arroyo, Victor Alcalde, Begoña Canovas, Modesto Orozco, Angel R. Nebreda, and Patrick Aloy. Quantification of pathway cross-talk reveals novel synergistic drug combinations for breast cancer. *Cancer Research*, 77(2):459–469, 2017.

[8] Krishna C. Bulusu, Rajarshi Guha, Daniel J. Mason, Richard P.I. Lewis, Eugene Muratov, Yasaman Kalantar Motamedi, Murat Cokol, and Andreas Bender. Modelling of compound combination effects and applications to efficacy and toxicity: state-of-the-art, challenges and perspectives. *Drug Discovery Today*, 21(2):225 – 238, 2016.

[9] Kaisa Miettinen. *Nonlinear Multiobjective Optimization*, volume 12 of *International Series in Operations Research & Management Science*. Kluwer Academic Publishers, Boston, USA, 1999.

[10] Jürgen Branke, Kalyanmoy Deb, Kaisa Miettinen, and Roman Słowiński. *Multiobjective Optimization: Interactive and Evolutionary Approaches*. Springer-Verlag, Berlin, Heidelberg, 2008.

[11] H.W. Hamacher and K.-H. Küfer. Inverse radiation therapy planning — a multiple objective optimization approach. *Discrete Applied Mathematics*, 118(1):145 – 161, 2002. Special Issue devoted to the ALIO-EURO Workshop on Applied Combinatorial Optimization.

[12] David L. Craft, Theodore S. Hong, Helen A. Shih, and Thomas R. Bortfeld. Improved planning time and plan quality through multicriteria optimization for intensity-modulated radiotherapy. *International Journal of Radiation Oncology*Biology*Physics*, 82(1):e83 – e90, 2012.

[13] K. Matlock, N. Berlow, C. Keller, and R. Pal. Combination therapy design for maximizing sensitivity and minimizing toxicity. *BMC Bioinformatics*, 18(Suppl 4):116, Mar 2017.

[14] Noah Berlow, Lara E. Davis, Emma L. Cantor, Bernard Séguin, Charles Keller, and Ranadip Pal. A new approach for prediction of tumor sensitivity to targeted drugs based on functional data. *BMC Bioinformatics*, 14(1):239, 2013.

[15] S. L. Holbeck, R. Camalier, J. A. Crowell, J. P. Govindharajulu, M. Hollingshead, L. W. Anderson, E. Polley, L. Rubinstein, A. Srivastava, D. Wilsker, J. M. Collins, and J. H. Doroshow. The National Cancer Institute ALMANAC: A Comprehensive Screening Resource for the Detection of Anticancer Drug Pairs with Enhanced Therapeutic Activity. *Cancer Res.*, 77(13):3564–3576, 07 2017.

[16] C. I. Bliss. The toxicity of poisons applied jointly. *Annals of Applied Biology*, 26(3):585–615, 1939.

[17] D. Y. Baeder, N. Yu G., Hozé, J. Rolff, and R. R. Regoes. Antimicrobial combinations: Bliss independence and Loewe additivity derived from mechanistic multi-hit models. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 371(1695):20150294, may 2016.

[18] Avichai Tendler, Anat Zimmer, Avi Mayo, and Uri Alon. Noise-precision tradeoff in predicting combinations of mutations and drugs. *PLOS Computational Biology*, 15(5):1–17, 05 2019.

[19] Kevin Wood, Satoshi Nishida, Eduardo D. Sontag, and Philippe Cluzel. Mechanism-independent method for predicting response to multidrug combinations in bacteria. *Proceedings of the National Academy of Sciences*, 109(30):12254–12259, 2012.

[20] Kevin B. Wood. Pairwise interactions and the battle against combinatorics in multidrug therapies. *Proceedings of the National Academy of Sciences*, 113(37):10231–10233, 2016.

[21] Anat Zimmer, Itay Katzir, Erez Dekel, Avraham E. Mayo, and Uri Alon. Prediction of multidimensional drug dose responses based on measurements of drug pairs. *Proceedings of the National Academy of Sciences*, 113(37):10442–10447, 2016.

[22] Anat Zimmer, Avichai Tendler, Itay Katzir, Avi Mayo, and Uri Alon. Prediction of drug cocktail effects when the number of measurements is limited. *PLOS Biology*, 15(10):1–16, 10 2017.

[23] Michael T. M. Emmerich and André H. Deutz. A tutorial on multiobjective optimization: fundamentals and evolutionary methods. *Natural Computing*, 17(3):585–609, Sep 2018.

[24] Arne Van Hoeck, Niels H. Tjoonk, Ruben van Boxtel, and Edwin Cuppen. Portrait of a cancer: mutational signature analyses for cancer diagnostics. *BMC Cancer*, 19(1):457, 2019.

[25] James Tsai, John T. Lee, Weiru Wang, Jiazhong Zhang, Hanna Cho, Shumeye Mamo, Ryan Bremer, Sam Gillette, Jun Kong, Nikolas K. Haass, Katrin Sproesser, Ling Li, Keiran S. M. Smalley, Daniel Fong, Yong-Liang Zhu, Adhirai Marimuthu, Hoa Nguyen, Billy Lam, Jennifer Liu, Ivana Cheung, Julie Rice, Yoshihisa Suzuki, Catherine Luu, Calvin Settachatgul, Rafe Shellooe, John Cantwell, Sung-Hou Kim, Joseph Schlessinger, Kam Y. J. Zhang, Brian L. West, Ben Powell, Gaston Habets, Chao Zhang, Prabha N. Ibrahim, Peter Hirth, Dean R. Artis, Meenhard Herlyn, and Gideon Bollag. Discovery of a selective inhibitor of oncogenic b-raf kinase with potent antimelanoma activity. *Proceedings of the National Academy of Sciences*, 105(8):3041–3046, 2008.

[26] R. Fisher and J. Larkin. Vemurafenib: a new treatment for BRAF-V600 mutated advanced melanoma. *Cancer Manag Res*, 4:243–252, 2012.

[27] A. Kim and M. S. Cohen. The discovery of vemurafenib for the treatment of BRAF-mutated metastatic melanoma. *Expert Opin Drug Discov*, 11(9):907–916, Sep 2016.

[28] James Larkin, Paolo A. Ascierto, Brigitte Dr, Victoria Atkinson, Gabriella Liszkay, Michele Maio, Mario Mandalev Demidov, Daniil Stroyakovskiy, Luc Thomas, Luis de la Cruz-Merino, Caroline Dutriaux, Claus Garbe, Mika A. Sovak, Ilsung Chang, Nicholas Choong, Stephen P. Hack, Grant A. McArthur, and Antoni Ribas. Combined vemurafenib and cobimetinib in braf-mutated melanoma. *New England Journal of Medicine*, 371(20):1867–1876, 2014. PMID: 25265494.

[29] Antoni Ribas, Adil Daud, Anna C. Pavlick, Rene Gonzalez, Karl D. Lewis, Omid Hamid, Thomas F. Gajewski, Igor Puzanov, Matthew Wongchenko, Isabelle Rooney, Jessie J. Hsu, Yibing Yan, Erica Park, and Grant A. McArthur. Extended 5-year follow-up results of a phase ib study (brim7) of vemurafenib and cobimetinib in braf-mutant melanoma. *Clinical Cancer Research*, 26(1):46–53, 2020.

[30] Monica Marzagalli, Marina Montagnani Marelli, Lavinia Casati, Fabrizio Fontana, Roberta Manuela Moretti, and Patrizia Limonta. Estrogen receptor β in melanoma: From molecular insights to potential clinical utility. *Frontiers in Endocrinology*, 7:140, 2016.

[31] Prson Gautam, Leena Karhinen, Agnieszka Szwajda, Sawan Kumar Jha, Bhagwan Yadav, Tero Aittokallio, and Krister Wennerberg. Identification of selective cytotoxic and synthetic lethal drug responses in triple negative breast cancer cells. *Molecular Cancer*, 15(1):34, 2016.

Figure 1: Selectivity analysis of the 104 individual drug compounds in NCI ALMANAC. (**A**) Data consists of drug sensitivity measurements of the NCI-60 cell lines at various concentrations c_i, c_j of drug pairs (here vemurafenib and gefitinib, respectively). Maximal single compound activity, Q_{min}, of a drug is obtained by fitting Hill curves to the dose-response data. (**B**) Hierarchical clustering of cell lines and drug compounds according to their Q_{min}. Highly effective but nonselective drugs are clustered together and show up as dark horizontal bands in the heatmap. Leukemias (red color columns) cluster strongly together as they are more vulnerable to a large variety of treatments than the other cancer types. Melanomas (violet columns) form one major cluster, and the rest of the melanoma cell lines are associated with other cancer types. In particular, all but one melanoma cell line stand out from most of the other cell lines because of their susceptibility to vemurafenib (light blue box). The drug activity profile of vemurafenib is clustered together most closely with estramustine, an estradiol derivative with a nitrogen mustard moiety, and the anthracycline neoplastics epirubicin and idarubicin. (**C**) Multiobjective optimization of single drug compounds acting on Malme-3M cell line. Vemurafenib is clearly the optimal single-compound treatment because it is both effective and selective. Bortezomib and valrubicin, near the rightmost tip of the convex hull, are the least optimal drugs. Mithramycin leads to highest therapeutic effect but it is also highly nonselective, far more than the common cytotoxic treatment cisplatin. EGFR inhibitor gefitinib alone is not a Pareto optimal solution. The multi-coloured dots show that a substantially better efficacy can be achieved by combining drugs. For the combination of vemurafenib and gefitinib, a higher therapeutic effect than predicted by Bliss independence model emerges, yet the boost in efficacy comes with an increase in nonselective effect as well.

Figure 2: **(A)** Multiobjective optimization (MOP) solutions for monotherapies and two-drug combinations. Monotherapies are represented by big monochrome dots, and multicoloured dots indicate combinations of selected compounds (see color legend). The small light blue dots represent all the other drugs alone and in two-drug combinations. Anastrozole alone has next to no effect on any cell line, yet in combination with vemurafenib and gefitinib, a selective effect emerges. The triplet combination of these three drugs is shown for comparison, and it yields a significant advance over all pairs of the drugs. Pareto fronts for $M = 1, 2$ are shown. **(B)** MOP for drug triplets and their subcombinations. Due to high number of possible combinations, only monotherapies of drugs indicated in the legend and drug combinations involving vemurafenib, gefitinib and anastrozole are shown. The triplet of vemurafenib and gefitinib with sirolimus is on the convex hull, indicating a strong selective effect, yet it comes with a significant cost in nonselective effect. The Pareto front for $M = 4$ coincides with the one for triplets within this (and actually much wider) window of nonselective effects, indicating that none of the fourth-order combinations perform better than drug triplets and their subcombinations.

Figure 3: Therapeutic dominance (bars, left y-axis) and selective synergy measured by the Bliss multiexcess (red curve, right y-axis) of Pareto optimal combinations as a function of nonselective effect. The numbered combinations correspond to Pareto optimal combinations highlighted in Fig. 2 B, from tretinoin (1) monotherapy to vemurarfenib+gefitinib+sirolimus (13) triplet. Vemurafenib monotherapy (2) is dominant at low mean effect. Vemurafenib + gefitinib (9) duplet has high a dominance but very low selective synergy, indicating that the combination is optimal because of its strong therapeutic effect, not due to selective drug interaction. The combination vemurafenib+gefitinib+sirolimus (13) has the highest selective synergy among shown combinations. The combinations vemurafenib+anastrozole (3) and vemurafenib+gefitinib+anastrozole (10) exhibit both considerable dominance and local maxima in selective synergy.