TITLE: Evolution in *Sinocyclocheilus* cavefish is marked by rate

shifts, reversals and origin of novel traits

Ting-Ru Mao†,[1], Ye-Wei Liu†, [1], Madhava Meegaskumbura†,‡, [1,3], Jian Yang[2], Gajaba

Ellepola[1,3], Gayani Senevirathne[4], Cheng-Hai Fu[1], Joshua B. Gross[5], Marcio R. Pie[6]

[1]Guangxi Key Laboratory for Forest Ecology and Conservation, College of Forestry, Guangxi

University, Nanning, Guangxi, P.R.C.

[2]Key Laboratory of Environment Change and Resource Use, Beibu Gulf, Nanning Normal

University, Nanning, Guangxi, P.R.C.

[3]Deparment of Molecular Biology and Biotechnology & Department of Zoology, Faculty of

Science, University of Peradeniya, Peradeniya, Sri Lanka

[4]Department of Organismal Biology & Anatomy, University of Chicago, Chicago, IL, USA

[5]Department of Biological Sciences, University of Cincinnati, Cincinnati, Ohio, USA

[6]Departamento de Zoologia, Universidade Federal do Paraná, Curitiba, Paraná, Brazil 81531-980

† - equal contribution

‡ - corresponding author - madhava_m@mac.com

ABSTRACT

Epitomized by the well-studied *Astyanax mexicanus*, cavefishes provide important model

organisms to understand adaptations in response to divergent natural selection. However, the

spectacular *Sinocyclocheilus* diversification of China, the most diverse cavefish clade in the

world harboring nearly 75 species, demonstrate evolutionary convergence for many traits, yet

remain poorly understood in terms of their morphological evolution. Here, using a broad sample

of 49 species representative of this diversification, we analyze patterns of *Sinocylocheilus*

evolution in a phylogenetic context. We categorized species into morphs based on eye-related

condition: Blind, Micro-eyed (small-eyed), and Normal-eyed and we also considered three

habitat types (Troglodytic – cave-restricted; Troglophilic – cave-associated; Surface – outside of

caves). Geometric morphometric analyses show Normal-eyed morphs with fusiform shapes

being segregated from Blind/Micro-eyed (Eye-regressed) morphs with deeper bodies along the

first principal component ("PC") axis. The second PC axis accounts for shape complexity related

to the presence of horns. Ancestral character reconstructions of morphs suggest at least three

independent origins of Blind morphs, each with different levels of modification in relation to the

typical morphology of ancestral Normal-eyed morphs. Interestingly, only some Blind or Micro-

eyed morphs bear horns and they are restricted to a single clade (Clade B) and arising from a

Troglodytic ancestral species. Our geophylogeny shows an east-to-west diversification spanning

the Pliocene and the Pleistocene, with Troglodytic species dominating karstic subterranean

habitats of the plains whereas predominantly Surface species inhabit streams and pools in hills to

the west (perhaps due to the scarcity of caves). Integration of morphology, phylogeny and

geography suggests *Sinocyclocheilus* are pre-adapted for cave dwelling. Analyses of

evolutionary rates suggest that lineages leading to Blind morphs were characterized by

significant rate shifts, such as a slowdown in body size evolution and a 3.3 to 12.5 fold increase

in the evolutionary rate of eye regression. Furthermore, body size and eye size have undergone

reversals, but horns have not, a trait that seem to require substantial evolutionary time to form.

These results, compared to the *Astyanax* model system, indicate *Sinocyclocheilus* fishes

demonstrate extraordinary morphological diversity and variation, offering an invaluable model

system to explore evolutionary novelty.

KEYWORDS: Phylomorphospace, Evolutionary Convergence, Blind fish, Troglodytism,

Geophylogeny, Pre-adaptation

INTRODUCTION

Due to the absence of light, stable mean temperatures, absence of primary productivity, and

paucity of dissolved oxygen, subterranean habitats are among the most challenging environments

for life on earth (Ginet and Decou, 1977, Camacho, 1992). From surface-dwelling ancestral

species, cavefish have secondarily adapted to live in cave systems, often demonstrating a

remarkable array of morphological and behavioral adaptations (Soares and Niemiller, 2013;

Yoshizawa, 2015). These involve enhanced sensation, and also dispensing of traits that incur a

developmental or energetic cost. Cavefish species can be divided into two forms: troglophiles are

closely associated with caves, but do not entirely depend on them, and troglobites are obligate

cave dwellers (Dowling et al., 2002; Jingcheng and Weicheng, 2015). Troglobite fish *may* harbor

special adaptations, such as complete eye loss, loss of pigmentation, changes in cranial

symmetry, proliferation of neuromast sensory organs, development of horns, and in some species

flat, hollow heads (Strecker et al., 2004; Zhao, 2006; Gross et al., 2008). Despite the ~200

cavefish species from across the world, large diversification of cavefishes is rare. However, one

extensive diversification occurs in *Sinocyclocheilus*, a monophyletic group of cyprinid fishes

endemic to China, which allows a robust analysis of trait evolution relative to troglomorphism in

a phylogenetic context.

The specialized traits cavefish bear have led them to be developed as models of evolution

especially with respect to adaptations to novel environments and evolutionary convergence

(Culver et al., 1995; Dowling et al., 2002; Jeffery, 2001; Li et al. 2008; Strecker et al., 2004;

Yang et al.， 2016). A lion's share of knowledge on evolution and development in cavefishes

has come from *Astyanax mexicanus* (Mexican tetra), a species with both surface-dwelling

(pigmented and eyed) and cave-dwelling morphs (depigmented and blind), which can readily

interbreed (Borowsky, 2008). In contrast to this well-studied model system, *Sinocyclocheilus*

species not only include blind and normal-eyed morphs (Lan et al., 2013), but demonstrate a

continuum from blind to normal-eyed species. Indeed, members of the *Sinocyclocheilus* genus

display remarkable morphological evolution with divergent cave-dwelling, cave-associated, and

surface-dwelling species.

Sinocyclocheilus species are thought to have shared a common ancestor in the late Miocene,

undergoing a spectacular diversification spanning the Pliocene and Pleistocene across the

southwestern parts of China's 620,000 km^2 of karst habitats (Huang et al., 2008), with nearly 75

extant species (Jiang et al., 2019). This resulted in an adaptive diversification into subterranean

refugia traversing the intersection of the Guizhou, Guangxi and Yunnan provinces around the

time of the uplifting of Tibetan/Guizhou plateau (Li et al., 2008).

One of the most striking forms of cave adaptation in *Sinocyclocheilus* is variation in eye

morphology, categorized often into three morphs (Zhao and Zhang, 2009), ranging from Normal-

eyed to Micro-eyed (small-eyed) to Blind species. Of all Chinese hypogean fishes, 56 species

show troglomorphism such as reduction and/or loss of eyes, pigmentation, and the gas bladder.

Presence of a horn-like structure and hyper-development of the dorsal protuberance (similar to a

humpback whale) are two additional unique characters to certain Chinese hypogean species

(Romero et al., 2009). These dramatic adaptations to cave life are reflected in the unique

morphology of these fish.

The morphology of *Sinocyclocheilus* is most likely attributed to their habitat and local

adaptations, however the precise function of certain morphologies (e.g., their horns and humps)

remains unknown (Ma and Zhao, 2012). For instance, many blind species are obligate cave

dwellers that have the ability to navigate along cave walls, cave-bottoms and within narrow

passages (Yoshizawa, 2015). Yet, others are open water species that navigate in the manner of

typical fish. There are also intermediate forms between these two principal morphs (Zhao and

Zhang, 2009). However, the morphology of these fishes is so extreme that substantial variation

in morphology is evident even within blind, intermediate and the open water species.

The pattern of body shape evolution in *Sinocyclocheilus* is a conundrum, which has not been

addressed in an evolutionary context. Here, we explore key patterns of morphological evolution

in these fishes, and demonstrate that the evolution in *Sinocyclocheilus* has been associated with

significant rate changes and trait reversals across their phylogenetic history.

MATERIALS & METHODS

Phylogeny estimation

We compiled sequence data from GenBank for two mtDNA fragments (*NADH4* and *cytb*) of

39 *Sinocyclocheilus* species, and the outgroup species *Linichthys laticeps* (Cyprinidae). In

addition, we generated sequence data for the *cytb* gene fragment of ten additional

Sinocyclocheilus species (Table S1). For these species, total genomic DNA was extracted using

the DNeasy Blood and Tissue Kit (Qiagen Inc., Valencia, CA) following the manufacturer's

protocols. DNA was amplified in 25-μL volume reactions: 3 mM $MgCl_2$, 0.4 mM of dNTP, 1X

buffer, 0.06 U of Taq DNA Polymerase, 2 mM of each primer. Thermocycling conditions

included an initial step at 94 °C for 3 min, followed by 35 cycles at 45 s at 94 °C, 1 min at 46-50

°C and 45 s at 48-56 °C, and a final step at 72 °C for 5 min. PCR products were electrophoresed

in a 1.5% agarose gel, stained with ethidium bromide and visualized under UV light.

Successfully amplified products were purified using MicroconTM Centrifugal Filter Units

(Millipore, Billerica, MA, U.S.A.). Sequencing reactions were carried out in 10 µl solutions

including the following final concentrations: 5 ng/µl of template DNA, 0.5 µl of Big DyeTM

(Applied Biosystems Inc., Foster City, CA, U.S.A.), 0.2 µM of each primer and 0.1X of reaction

buffer. The final product was purified using SephadexTM G-50 (GE Healthcare Bio-Sciences

AB, Uppsala, Sweden) for sequencing. Forward and reverse strands were reconciled using

Staden v.1.6.0 (Staden 1996). Sequences from both genes were concatenated and aligned

unambiguously using CLUSTALW (Thompson et al., 2003), as implemented in MEGA v. 6.0

(Tamura et al. 2013), for a total alignment length of 2155 bp. We used JMODELTEST v.2.1.10

(Santorum et al., 2014) to determine the best models of evolution for each fragment, which were

implemented in BEAST v.1.10.4 (Drummond and Rambaut, 2007) as a partitioned analysis to

estimate the phylogenetic relationships and relative divergence times within *Sinocyclocheilus*.

We calibrated the tree using the relative time period which diversification of *Sinocyclocheilus*

initiated around 11.31 Mya, a reference point obtained from (Chen et al., 2018). We used a strict

molecular clock and a calibrated Yule tree prior, as well as a GTR+I+G for each partition and

ran the analysis for 20 million generations using the Cipres Science Gateway Server (Miller et

al., 2010). Convergence was assessed by inspecting the log-output file in TRACER v.1.6

(Drummond et al., 2012), and by ensuring ESS values were above 200. The first 10% of the trees

were discarded, and the post burn-in trees were used to infer the maximum clade credibility tree

using TREEANNOTATOR v.1.4.4 (Rambaut and Drummond，2019). The maximum clade

credibility tree, as well as a set of 1000 post burn-in topologies, were retained for further

analyses (see below).

Morphometric data acquisition and analyses

We assembled a database of images of referenced specimens of *Sinocyclocheilus* from

scaled photographs, which was complemented by species that we photographed (Table S2). The

final dataset included 90 images (54 from species descriptions and catalogues and 36 from

photographs by the authors) for 50 species, which included 70% of the total number of described

species for the genus. These images were used for geometric morphometrics analyses, which

were based on 15 landmarks (See Fig. 1A) and 180 sliding semi-landmarks, obtained using

tpsDig v. 2.16 (Rohlf, 2010). Semi-landmarks were collected as curves outlining the body. These

data were subsequently reduced to equidistant landmarks, and defined as semi-landmarks using

tpsUtil v. 1.46 (Rohlf, 2010). We then slid the landmarks using the bending energy method

(Gunz & Mitteroecker, 2013) implemented in GEOMORPH v.3.2.0 (Adams & Otárola-Castillo,

2013). The landmark coordinates were aligned using a generalized Procrustes superimposition

analysis (Adams *et al*., 2013), and a principal component analysis (PCA) was used to evaluate

shape variation within the sample.

Multiple images for the same species were used to obtain the landmarks and the mean of

their Procrustes coordinates were calculated to be used in later analyses. We also obtained

traditional linear measurements, namely standard length (SL), eye diameter (ED) and

standardized eye diameter (sED, calculated as the ratio between ED and SL).

Morphological and habitat evolution

Since shape variation in *Sinocyclocheilus* cavefishes occurs mostly in the anterior end of the

fish and as one of the major features leading to this is eye-related, we considered the absence or

the size of the eye (when present) as a proxy for the categorization of morphs. Since the eye size

has an allometric association with body size, we used the standardized eye diameter (sED) in

placing them into three morphological categories: Blind (eye absent); Micro-eyed (0.0 – 3.0mm);

Normal-eyed (< 3.0mm) respectively. For ease of discussion, we considered Blind and Micro-

eyed together as Regressed-eyed; Micro-eyed and Normal-eyed together as Eyed species.

Sinocyclocheilus were also categorized based on their habitat as Troglodytic, Troglophilic,

and Surface species. Troglodytic species live in an obligatory association with caves, and are not

sampled outside of caves. Caves, as meant here represent roofed-caves, submerged caves, and

subterranean waterways that form windows intermittently with the surface. Troglophilic species

live in a close association with caves and are sampled both in the vicinity of cave entrances and

within caves. Finally, Surface species are found in habitats even when a cave is not found in

close proximity and live in normal streams ponds and lakes as typical fish do, but they could

venture into caves (underground water bodies) during unfavorable periods, when water is only

available in caves. It should be noted here that this categorization is strictly habitat based and not

morphology based (for instance, there are Normal-eyed species that are Troglodytic,

Troglophilic or Surface). These habitat associations are based on published literature (Zhao and

Zhang, 2009; Romero et al., 2009) and personal observations as outlined in Table S1.

To infer the number and timing of evolutionary shifts within eye-related morphs, horn

distribution and the habitat type, we used stochastic character mapping (Nielsen, 2002;

Huelsenbeck et al. 2003), as implemented in the make.simmap function in PHYTOOLS. On each of

the 1000 post burn-in trees obtained from BEAST, we used stochastic character mapping to

generate 100 potential histories. This approach therefore considers uncertainty both in the

evolutionary history of the traits as well as in the inferred topology of the phylogeny.

The landmark coordinates obtained were aligned using a generalized Procrustes

superimposition analysis (Adams et al., 2013), and a principal component analysis (PCA) was

used to explore shape variation within the sample. In addition, we described the eye-related

morphological variation in *Sinocyclocheilus* by estimating ancestral states of SL and ED and

visualizing them using traitgrams (Evans et al., 2009) as implemented in the phenogram function

in PHYTOOLS 0.7.20 (Revell, 2012) using the maximum clade credibility tree. We also visualized

the evolution of both traits simultaneously using a phylomorphospace (a projection of the tree

into morphospace, *sensu* Sidlauskas, 2008) using the phylomorphospace function in PHYTOOLS.

Evolutionary rate variation in eye related morphs

We tested whether the evolutionary rates of the studied continuous traits (SL, ED, sED) are

significantly different in different morphs. We used 100 potential trait histories from stochastic

character mapping and then fit two alternative models of evolution on each studied trait, one that

fixes the rate of evolution to be identical between morphs against an alternative model in which

the morphs have separate rates. We calculated the Akaike Information Criterion for small sample

size (AICc) from the maximum likelihood estimate on each tree using the brownie.lite function

in PHYTOOLS. Finally, if a multi-rate model provided a better fit to the data, we calculated model-

averaged estimates of evolutionary rates for each morph. Unless otherwise indicated, all analyses

were conducted using R v. 3.6.0 (R Core Team, 2019).

Geophylogeny analyses

We could not carry out a formal biogeographical analysis, given that their high endemism

and the complex pattern of underground connections between caves limits the establishment of

reasonable biogeographical areas. However, we assessed the geographical structuring

of *Sinocyclocheilus* diversification by building a geophylogeny on GenGIS v. 2.5.3 (Parks et al.,

2013) based on the maximum credibility tree.

RESULTS

The maximum credibility tree of *Sinocyclocheilus* is shown in Fig. 2 together with the

reconstruction of ancestral states for eye-related morphs; we consider four major clades

(A,B,C,D), as previously reported by other authors (Zhao and Zhang, 2009). Despite the inherent

uncertainty in ancestral state reconstructions, it is clear that Blind species evolved at least three

times in *Sinocyclocheilus*. Two of these events involved single species evolving from Normal-

eyed ancestors, namely *S. xunlensis* and *S. anophthalmus*. On the other hand, the third lineage of

blind *Sinocyclocheilus,* Clade B, includes several closely related species of Blind, Micro-eyed

and a few Normal-eyed species, with two cases of reversal from either Micro-eyed or Blind to

Normal-eyed morphs, namely *S. zhenfengensis* and *S. brevibarbatus* (Fig. 2. All four clades

contain Regressed-eyed species and comparatively, clade Blind contains the most cases.

Interestingly, clade B originated around the time of the beginning of the aridification process in

China in the late Pliocene, whereas the other two transitions to blind species were much more

recent (Fig. 2.

Interestingly, the evolution of body size and eye diameter seem to have often involved

reversals, with little correspondence between body size (Fig. 3A) or eye diameter (Fig. 3B) and

their corresponding morphs, except for the case of blind species for which eye diameter is

necessarily zero. On the other hand, morphs are clearly distinguished when eye diameter and

body size are visualized simultaneously (Fig. 3C), which suggests that the evolution of different

morphs is achieved by altering the relationships between body size and eye diameter. Habitat

associations traced on the phylomorphospace (Fig. 3D) indicates that species with regressed eyes

and small-to-medium body sizes are obligate cave dwellers. However, normal eyed species can

be Troglodytic, Troglophilic or surface dwellers regardless of their body size. Interestingly all

horned species are obligate cave dwellers while all cave species are not horned (Fig. 3E).

A more precise description of the overall changes associated with different morphs can be

visualized in the projections build from the geometric morphometrics analyses (Fig. 1B). The

first PC, which accounted for approximately 32% of the variance in the dataset (see Table S3),

tended to distinguish the slender Normal-eyed species on the left and Micro/Blind species on the

right, which were characterized by changes in shape and widening of the anterior dorsal area

between mouth and beginning of the dorsal fin of their body, resulting in a shift from the

fusiform shape of the Normal-eyed forms to a more "boxy" form of the Micro-eyed and Blind

forms. The second PC, which explained approximately 17% of the variance in the dataset,

emphasized the differences in the type of dorsoventral broadening of the mid-section between

morphs, with a shortening of the tail region (Fig. 1B). The variation in this axis is very high

among the Micro-eyed and Blind forms when compared to the Normal morphs.

The multiple-rate model of evolution provided the best fit to the data for all three

quantitative traits (ΔAIC=2.7, 10.7 and 7.9 respectively for ED, sED and SL; Table 1), indicating

that the evolution of different *Sinocyclocheilus* morphs was associated with significant changes

in their evolutionary rates. However, there were intriguing differences between traits in their

rates (Table 2). The rates of evolution of eye diameter and standardized eye diameter were

similar between Normal-eyed and Micro-Eyed species, but increased between 3.3 to 12.56 times

during shifts towards Blind species.

Geophylogeny represents the phylogeny overlaid across the geographic location of each

species, where phylogenetic clustering is evident across the landscape. Considering the

distribution of *Sinocyclocheilus,* we mainly see a pattern where the basal, Normal-eyed morphs

are placed in the east, a substantial portion of Blind/Micro-eyed (Regressed-eyed) species are in

the center, and Normal-eyed morphs are predominant towards the western mountains (Fig. 4).

DISCUSSION

Habitat utilization in context of eye-morphology

Integrating evolution of eye size and habitat manifests interesting and previously

unrecognized evolutionary patterns in the evolution of *Sinocyclocheilus*. The Eye-size based

ancestral reconstruction suggests the base of the phylogeny is most likely an Eyed species (i.e.

Normal- or Micro-), but habitat reconstructions places, with high probability, Troglodytic species

at the base (Fig. S1). This suggests an ancestral Eyed-species evolved a Troglodytic habit before

they became blind. This may be an example of preadaptation in *Sinocyclocheilus*, i.e., the

advancement of a functional change with little or no evolutionary modification (Ardila, 2016). In

Astyanax cavefish, surface-dwelling forms are scotophilic, they prefer to remain away from

direct light suggesting that scotophilia may be preadaptive for colonizing the dark, cave

environment (Espinasa et al., 2001). In *Sinocyclocheilus*, since a basal (eyed) species

demonstrated preference for the cave habitat, this preadaptation to darkness may hint towards

why certain species tend to become cave-dwellers while others do not. This pattern is supported

by two principal lines of evidence. First, most of the basal species are eyed, and Troglodytic

(except for one species with an unusual eye-related polymorphic condition that we discuss

below). Second, the most westward group (Clade D; Normal-eyed Surface fish), re-colonized

caves whenever cave habitats were available within that area, suggesting a strong predisposition

for cave-dwelling across all *Sinocyclocheilus*. In other words, when caves were present,

members of *Sinocylocheilus,* irrespective of eye-related condition, preferred the cave habitat.

The preference for caves may not be a preference for darkness, but in fact a preference for

depth, in search of water for survival. In a karstic environment where drying of surface running

water is common, a preference for such deeper habitats may have provided an evolutionary

advantage. In the presence of an array of subterranean waterways, such a predisposition would

have given rise to the eye-regressed forms living close to, or associated with, caves that are

characteristic of the genus.

Furthermore, apart from the Troglodytic and Troglophilic species of Clade D, some of the

putative Surface species of Clade D are often observed at the entrances of caves or at windows to

subterranean rivers (Zhao and Zhang, 2009). Hence, with more intensive ecological studies,

some species recognized as Surface species may indeed be Troglophilic species, bolstering the

notion that *Sinocyclochelus* are predisposed to seek deeper waters of the karstic caves.

Resource utilization plays a key survival role in harsh environments (Culver and Pipan,

2009). Some of the Troglophilic, eyed-species are nocturnal, emerging from submerged caves,

presumably to feed at night to reduce competition from other non-cave inhabiting fish species

(personal observations). Some species like *S. altishoulderus, S. donglanensis* (Romero et al.，

2009), *S. bamaensis* (Su et al., 2003), *S. malacopterus* (Chen et al., 2017) and *S. longibarbatus*

(personal observation, video evidence as Supplementary information); are known to come out of

caves during the high water season, presumably to feed and breed. This explains dependence on

the cave as a diurnal refugium, from where these species can exploit the surface habitats at night.

Strategies such as this, where multiple resources are utilized simultaneously, points to the

adaptability of some *Sinocyclocheilus* species, resulting in their evolutionary success in a harsh

and changing environment. Hence, the cave entrances are possibly an important ecotone that is

important in *Sinocyclochelius* diversification and conservation.

Season and time of day seem to be important factors in determining habitat utilization

patterns, but this level of resolution in habitat data is not currently available for a majority of the

species to carry out a comprehensive habitat analysis across the diversification – indeed, many

species are known only from one or a few specimens (Zhao and Zhang, 2009).

Adaptations in the light of geophylogeny

In combination with the data analyzed, basal *Sinocyclocheilus* (Clade A) are Normal-Eyed,

predominantly cave dwelling and non-horned species from the Eastern region of their

distribution. As pointed out, this suggests that the earliest ancestors of *Sinocyclocheilus* species

where Normal-eyed and but still lived in close association with caves. The ancestral

reconstructions suggest that the affinity to caves would have evolved early and is present in most

Sinocyclocheilus. The clade comprising basal species are from the east of the *Sinocyclocheilus*

distribution, i.e. the He Jiang and Gui Jiang river basin in northeastern Guangxi, and hence, it

seems that the diversification of these fish occurred from East to West (Fig. 4).

Within this predominantly Normal-eyed clade (Clade A), there is an exception,

Sinocyclocheilus guanyangensis, a species that we coded as Micro-eyed, has Normal-Eyed,

Micro-eyed and effectively Blind morphs within the same population – polymorphic for this

trait. But these blind morphs have their eyes completely covered by skin and the Micro-eye is not

itself degenerate. This kind of condition has been noted in several other taxa also (*S. xunlensis*

and *S. flexuodorsalis* – not available for our analysis), but is uncommon. This suggests a degree

of polymorphism for this trait, suggesting that the earliest ancestors of *Sinocyclocheilus* may

have been able to lose or gain eyes relatively easily as an adaptation to local conditions, this

ability appears several times within this cave-driven diversification.

The major adaptation for cave dwelling evolves predominantly in the expansive karstic area

in northwestern Guangxi (associated with the Liu Jiang basin and Hongshui river basin joining

the main Xijiang River system from the North), in Clade B, the southeastern corner of Guizhou

province (upper reaches of Hongshui River) and the northeastern plateau of Yunnan province.

This region can be considered the center for novel adaptations for *Sinocyclocheilus,* where these

fishes express their full morphological diversity, blindness, micro-eyedness, and their remarkable

horns. In the shape-related analyses, these species cluster on the right of morphospace (Fig. 1B).

The deeper caves and extensive subterranean river system associated with the Guangxi plains

(Zhao and Zhang, 2009) would have facilitated this extensive adaptive diversification (Fig. 4).

The karstic northwestern region that the Guangxi-dominated Clade (Clade B) experiences

drought conditions during much of the year, and one of the major sources of rain for the region is

through storms sweeping from the southeast that are strong enough to persist through the vast

plains of Guangxi, mainly from April to August (Zhao and Zhan, 2009). During unfavorable

periods, these fishes seem to have found refuge in the subterranean caves. The morphologically

most diverse clade being present in the region where the climatic conditions are most

unfavorable for surface fish, reinforces the notion that *Sinocyclocheilus* species adapted to life in

caves as climatic refuges (Zhao and Zhang, 2009).

The distribution of Clade C, characterized by mostly Normal-eyed but Troglodytic species

largely overlaps Clade B. In Clade C, the single Blind species (*S. xunlensis*) and the two Micro-

Eyed species (*S. cyphotergous* and *S. multipunctatus*) are shown within a narrow geographic area

on the Liu Jiang and Hongshui river system (Fig. 4).

Species in the clade that is found in the west (Clade D), predominantly on the hilly terrain of

Yunnan plateau, are predominantly Normal-eyed Surface species lacking horns (Fig. 1, Fig. 4).

However, wherever there are cave habitats and subterranean river systems, some of these

putative surface species have become facultative or obligate cave species. The obligate cave

species found within the region, *S. anophthalmus*, is blind. However, this Blind species stands

clustered with the Normal-eyed morphs in the morphospace, signifying that the shape of the

species has not extensively changed, possibly due to recent (Pleistocene) invasion of the cave

habitat from a Normal-eyed ancestor (Fig. 2) – time since becoming blind is not been long

enough for change into the box-like shape of the Blind species of Clade B.

Horn distributions show several peculiar trends. In most *Sinocyclocheilus* species, a

prominent hump is present (He et al., 2013). However, this hump is markedly low in the Normal-

eyed surface inhabiting species of the Yunnan clade (Fig. 4). For the species that bears a horn,

the horn represents the region in which the dorso-frontal hump is present, and always occurs

before the hump begins, at the boundary of the edge of the dorsal skull. The exception to this is

S. cyphotergous (Clade C), where a horn like structure is placed on top of the hump. *S.*

cyphotergous, a species found in clade C, is phylogenetically separate from other horned species,

suggesting that the origins of the "horn" for this species is evolutionarily different from the other

horned species (Fig. S2). Though the function of the horn remains unknown (protection of head,

anchoring in strong current and protection of head has been suggested; Zhao and Zhang, 2009),

functionally this structure may be similar to other species, if it is actually anchoring in strong

current is the main function.

The rates of evolution of various traits show some incongruent (non-allometric), but

interesting patterns that can be explained in the context to adaptations to a Troglodytic condition.

The rates of evolution in eye diameter are similar between Normal- and Micro-Eyed species but,

increases dramatically (3.3-12.56 times) with shifts towards blind forms. However, body size

evolution for these morpho groups shows a reversed pattern, with a 0.03 decrease in body size

evolution in the Blind morphs compared to the eyed-morphs. These patterns in rate variation

suggest that the evolution of Blind morphs to a Troglodytic habitat were simultaneously

associated with an increase in the rate of evolution of the eye-size degeneration and a decrease in

the rate of body size evolution. The smaller body size resulting from a sluggish rate of change

will facilitate both navigation within constricted spaces and sustenance on a limited supply of

resources expected to be experienced in subterranean habitats.

Much of our collective knowledge of the patterns and mechanisms of regressive evolution

come from studies of animals that have colonized the subterranean biome. Within this group, a

several studies have focused on the Mexican tetra, *Astyanax mexicanus* (Jeffery, 2009). This

natural animal model system comprises multiple cave-adapted morphs and a surface-dwelling

morph that resides in near the caves themselves (Gross, 2012). Since the discovery of *Astyanax*

cavefish in 1936, countless studies have provided insight to the developmental and genetic bases

for cave-associated traits (Hubbs and Innes, 1936). Indeed, much of this insight has emerged

from the interbreeding studies of conspecific cave and surface morphs (reviewed in Wilkens,

2016). However, several aspects of regressive evolution and troglomorphic adaptation remain

unresolved. Owing to several of the differences with *Astyanax*, we argue that *Sinocyclocheilus* is

well-positioned to provide important new insights to broader patterns of diversification and

adaptation in cave-dwelling organisms.

For instance, *Astyanax* represents a single species with 30 different interfertile populations

from a relatively small geographic location (Wilkens and Strecker, 2017). In contrast,

Sinocyclochelius harbors about 75 species (49 in our study) inhabiting diverse geographic

biomes across a much larger geographic area. Moreover, while *Astyanax* cavefish converge on

similar phenotypes (regressed vision and pigmentation), they are not regarded as having

numerous morphological novelties. In contrast, *Sinocyclocheilus* species have recurrently

evolved a unique "horn" (Soares et al., 2019) from several eyeless species. Similarly, the larger

number of *Sinocyclocheilus* species allows keener resolution for understanding broad

phylogenetic processes, such as trait reversals and directions of diversification. Although a

reversal from an eyeless to an eyed form has been reported for one cave population in *Astyanax*

(Caballo Moro; Krishnan and Rohner, 2017), this phenomenon appears to be much less common

than in *Sinocyclocheilus*. Additionally, a clear polarity of diversification is lacking in *Astyanax*

cavefish, rather ancient stocks of surface-dwelling forms appear to have recurrently invaded

caves to the east (i.e., Sierra de El Abra caves), with more recent invasions having occurred in

the northern (Sierra de Colmena) and the western caves (Sierra de Guatemala; Bradic et al.,

2012). However, the well-characterized gene flow between the cave and surface waters obscures

the ability to understand clear boundaries between different cave groups. Further, most *Astyanax*

cave populations are believed to have diverged over the course of the last ~200 – 500 Ky

(Herman et al., 2018). By contrast, *Sinocyclocheilus* species are much older, and therefore one

can determine how longer-term processes unfold in these cave-dwelling animals. Thus, despite

clear phylogenetic differences between *Astyanax* and *Sinocyclocheilus*, both genera have the

ability to provide complementary and critical insights to the processes underlying cave evolution

and diversification.

The integration of morphology, phylogeny, rate analyses, dating and distribution show not

only several remarkable patterns of evolution, but also interesting exceptions to these patterns

that signifies the diversification of *Sinocyclocheilus* as a unique model system to study

evolutionary novelty.

ACKNOWLEDGEMENTS

We thank the following institutions and individuals: funding from Guangxi University

Special Talent Recruitment Grant to MM; funding from National Natural Science Foundation of

China (31860600 and Guangxi Natural Science Foundation (2017GXNSFFA198010) JY;

Shipeng Zhou, Bing Chen, Dan Sun, Jayampathi Herath and Amrapali Rajput for assistance in

the field; Ethical review approval from Guangxi University.

REFERENCES

Adams, D. C., and E. Otárola-Castillo. 2013. geomorph: an r package for the collection and

analysis of geometric morphometric shape data. Methods Ecol Evol 4:393–399.

Adams, D. C., F. J. Rohlf, and D. E. Slice. 2013. A field comes of age: geometric morphometrics

in the 21st century. Hystrix, the Italian Journal of Mammalogy 24:7–14.

Ardila, A. 2016. The Evolutionary Concept of "Preadaptation" Applied to Cognitive

Neurosciences. Front. Neurosci. 10:103.

Borowsky, R. 2008. Breeding *Astyanax mexicanus* through Natural Spawning. Cold Spring

Harbor Protocols 2008:pdb.prot5091-pdb.prot5091.

Bradic, M., P. Beerli, F. J. García-de León, S. Esquivel-Bobadilla, and R. L. Borowsky. 2012.

Gene flow and population structure in the Mexican blind cavefish complex (*Astyanax*

mexicanus). BMC Evol Biol 12:9.

Camacho , A. I. ed. 1992. The Natural History of Biospeology. Monographs of the National

Museum of Natural Sciences. . National Museum of Natural Sciences., Madrid.

Chen, Y.-Y., R. Li, C.-Q. Li, W.-X. Li, H.-F. Yang, H. Xiao, and S.-Y. Chen. 2018. Testing the

validity of two putative sympatric species from *Sinocyclocheilus* (Cypriniformes:

Cyprinidae) based on mitochondrial cytochrome b sequences. Zootaxa 4476:130.

Chen, Z. M., J. Luo, H. Xiao, and J. X. Yang. 2010. Subterranian fishes of China. Pp. 397–

414 *in* Biology of Subterranean Fishes. E. Trajano, M. E. Bichuette, B.G. Kapoor (eds）.

Science Publishers, USA.

Culver, D. C., T. C. Kane, and D. W. Fong. 1995. Adaptation and natural selection in caves: the

evolution of Gammarus minus. Harvard University Press, Cambridge, Mass.

Culver, D. C., and T. Pipan. 2009. The biology of caves and other subterranean habitats. Oxford

University Press, New York.

Dowling, T. E., D. P. Martasian, and W. R. Jeffery. 2002. Evidence for Multiple Genetic Forms

with Similar Eyeless Phenotypes in the Blind Cavefish, *Astyanax mexicanus*. Molecular

Biology and Evolution 19:446–455.

Drummond, A. J., and A. Rambaut. 2007. BEAST: Bayesian evolutionary analysis by sampling

trees. BMC Evol Biol 7:214.

Drummond, A. J., M. A. Suchard, D. Xie, and A. Rambaut. 2012. Bayesian Phylogenetics with

BEAUti and the BEAST 1.7. Molecular Biology and Evolution 29:1969–1973.

Espinasa, L., P. Rivas-Manzano, and H. E. Pérez. 2001. A New Blind Cave Fish Population of

Genus *Astyanax*: Geography, Morphology and Behavior. Environmental Biology of Fishes

62:339–344.

Evans, M. E. K., S. A. Smith, R. S. Flynn, and M. J. Donoghue. 2009. Climate, Niche Evolution,

and Diversification of the "Bird-Cage" Evening Primroses (*Oenothera* , Sections *Anogra*

and *Kleinia*). The American Naturalist 173:225–240.

Ginet, R., and V. Decu. 1977. Initiation à la biologie et à l'écologie souterraines. J.-P. Delarge,

Paris.

Gross, J. B. 2012. The complex origin of *Astyanax* cavefish. BMC Evol Biol 12:105.

Gross, J. B., M. Protas, M. Conrad, P. E. Scheid, O. Vidal, W. R. Jeffery, R. Borowsky, and C. J.

Tabin. 2008. Synteny and candidate gene prediction using an anchored linkage map of

Astyanax mexicanus. Proceedings of the National Academy of Sciences 105:20106–20111.

Gunz, P., and P. Mitteroecker. 2013. Semilandmarks: a method for quantifying curves and

surfaces. Hystrix, the Italian Journal of Mammalogy 24(1):103-109.

He, Y., X. Y. Chen, T. Q. Xiao, and J. X. Yang. 2013. Three-dimensional morphology of the

Sinocyclocheilus hyalinus (Cypriniformes: Cyprinidae) horn based on synchrotron X-ray

microtomography . Zoological Research 34(4−5):128–134.

Herman, A., Y. Brandvain, J. Weagley, W. R. Jeffery, A. C. Keene, T. J. Y. Kono, H. Bilandžija,

R. Borowsky, L. Espinasa, K. O'Quin, C. P. Ornelas-García, M. Yoshizawa, B. Carlson, E.

Maldonado, J. B. Gross, R. A. Cartwright, N. Rohner, W. C. Warren, and S. E. McGaugh.

2018. The role of gene flow in rapid and repeated evolution of cave-related traits in Mexican

tetra, *Astyanax mexicanus*. Mol Ecol 27:4397–4416.

Huang, Q., Y. Cai, and X. Xing. 2008. Rocky Desertification, Antidesertification, and

Sustainable Development in the Karst Mountain Region of Southwest China. AMBIO: A

Journal of the Human Environment 37:390–392.

Hubbs, C. L., and W. T. Innes . 1936. The first known blind fish of the family Characidae: A

new genus from Mexico. Occasional papers of the Museum of Zoology, University of

Michigan 342:1–10.

Huelsenbeck, J. P., R. Nielsen, and J. P. Bollback. 2003. Stochastic Mapping of Morphological

Characters. Systematic Biology 52:131–158.

Jeffery, W. R. 2001. Cavefish as a Model System in Evolutionary Developmental Biology.

Developmental Biology 231:1–12.

Jeffery, W. R. 2009. Regressive evolution in *Astyanax* cavefish. Annual Review of Genetics

43:25–47.

Jiang, W. S., J. Li, X. Z. Lei, Z. R. Wen, Y. Z. Han, J. X. Yang, and J. B. Chang. 2019.

Sinocyclocheilus sanxiaensis, a new blind fish from the Three Gorges of Yangtze River

provides insights into speciation of Chinese cavefish. Zoological research 40(6):552–557.

Jingcheng, R., and Y. Weicheng. 2015. A Review of Progress in Chinese Troglofauna Research.

Journal of Resources and Ecology 6:237–246.

Krishnan, J., and N. Rohner. 2017. Cavefish and the basis for eye loss. Phil. Trans. R. Soc. B

372:20150487.

Lan, J. H., X. Gan, T. J. Wu, and J. Yang. 2013. Cave fishes of Guangxi, China. Science Press,

Beijing.

Li, Z., B. Guo, J. Li, S. He, and Y. Chen. 2008. Bayesian mixed models and divergence time

estimation of Chinese cavefishes (Cyprinidae: *Sinocyclocheilus*). Sci. Bull. 53:2342–2352.

Lunghi, E., Y. Zhao, X. Sun, and Y. Zhao. 2019. Morphometrics of eight Chinese cavefish

species. Sci Data 6:233.

Ma, L. and Y.H. Zhao. 2012. Cavefish of China. in: White, W.B.; and D.C. Cuvier, editors.

Encyclopedia of Caves. Pp. 107—125 Elsevier. ISBN 9780123838322

Miller, M. A., W. Pfeiffer, and T. Schwartz. 2010. Creating the CIPRES Science Gateway for

inference of large phylogenetic trees. Pp. 1–8 *in*. New Orleans, LA.

Nielsen, R. 2002. Mapping Mutations on Phylogenies. Systematic Biology 51:729–739.

Parks, D. H., T. Mankowski, S. Zangooei, M. S. Porter, D. G. Armanini, D. J. Baird, M. G. I.

Langille, and R. G. Beiko. 2013. GenGIS 2: Geospatial Analysis of Traditional and Genetic

Biodiversity, with New Gradient Algorithms and an Extensible Plugin Framework. PLoS

ONE 8:e69885.

R Development Core Team. 2019. a language and environment for statistical computing:

reference index. R Foundation for Statistical Computing, Vienna.

Rambaut, A., and A. J. Drummond. 2013. TreeAnnotator v1.10.4. Available as part of the

BEAST package at http://beast.bio.ed.ac.uk. .

Revell, L. J. 2012. phytools: an R package for phylogenetic comparative biology (and other

things): phytools: R package. Methods in Ecology and Evolution 3:217–223.

Rohlf, F. J. 2010. tpsDig 2.16 edn. at http://life.bio.sunysb.edu/morph/.

Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish

86:211–278.

Santorum, J. M., D. Darriba, G. L. Taboada, and D. Posada. 2014. jmodeltest.org: selection of

nucleotide substitution models on the cloud. Bioinformatics 30:1310–1311.

Sidlauskas, B. 2008. Continuous and arrested morphological diversification in sister clades of

characiform fishes: A phylomorphospace approach. Evolution 62:3135–3156.

Soares, D., and M. L. Niemiller. 2013. Sensory Adaptations of Fishes to Subterranean

Environments. BioScience 63:274–283.

Soares, D., M. Pluviose, and Y. Zhao. 2019. Ontogenetic development of the horn and hump of

the Chinese cavefish *Sinocyclocheilus furcodorsalis* (Cypriniformes: Cyprinidae). Environ

Biol Fish 102:741–746.

Staden, R. 1996. The staden sequence analysis package. Mol Biotechnol 5:233–241.

Strecker, U., V. H. Faúndez, and H. Wilkens. 2004. Phylogeography of surface and cave

Astyanax (Teleostei) from Central and North America based on cytochrome b sequence data.

Molecular Phylogenetics and Evolution 33:469–481.

Su, R., J. Yang, and G. Cui. 2003. Taxonomic review of the genus *Sinocrossocheilus wu*

(Teleostei: Cyprinidae), with a description of four new species. Zoological Studies 42:420–

430.

Tamura, K., G. Stecher, D. Peterson, A. Filipski, and S. Kumar. 2013. MEGA6: Molecular

Evolutionary Genetics Analysis Version 6.0. Molecular Biology and Evolution 30:2725–2729.

Thompson, J. D., T. J. Gibson, and D. G. Higgins. 2003. Multiple Sequence Alignment Using

ClustalW and ClustalX. Current Protocols in Bioinformatics 00: 2.3.1-2.3.22.

Wilkens, H. 2016. Genetics and hybridization in surface and cave *Astyanax* (Teleostei): a

comparison of regressive and constructive traits. Biol. J. Linn. Soc. 118:911–928.

Wilkens, H., and U. Strecker. 2017. Surface and cave populations of Mexican *Astyanax*. Pp. 37–

52 *in* Evolution in the Dark . Springer, Berlin, Heidelberg.

Yang, J., X. Chen, J. Bai, D. Fang, Y. Qiu, W. Jiang, H. Yuan, C. Bian, J. Lu, S. He, X. Pan, Y.

Zhang, X. Wang, X. You, Y. Wang, Y. Sun, D. Mao, Y. Liu, G. Fan, H. Zhang, X. Chen, X.

Zhang, L. Zheng, J. Wang, and L. Cheng. 2016. The *Sinocyclocheilus* cavefish genome

provides insights into cave adaptation. BMC Biology 14:1.

Yoshizawa, M. 2015. Behaviors of cavefish offer insight into developmental evolution: Evo-

Devo of cave fish behaviour. Mol. Reprod. Dev. 82:268–280.

Zhao, Y. 2006. Cavefishes: concept, diversity and research progress. Biodiversity Science

14:451.

Zhao, Y., and C. Zhang. 2009. Endemic Fishes of *Sinocyclocheilus* (Cypriniformes: Cyprinidae)

in China -Species diversity, cave adaptation, systematics and zoogeography. Science Press,

Beijing.

Main Figures

Fig. 1. (A) A specimen of *Sinocyclocheilus altishoulderus* indicating the position of 15 landmarks (red: larger points indicated by numbers 1-15) and 180 semi-landmarks (red: smaller point) used for the calculation of Procrustes coordinates and traditional linear measurements (SL: standard Length, ED: eye diameter and sED: standardized eye diameter) used in the geometric morphometric analyses. (B) **PCA showing the variation in body shape of the genus *Sinocyclocheilus* traced with eye morphology and habitat occupation.** PC1 and PC2 accounts for 32% and 17% of the variance respectively. A shift from the fusiform shape of the Normal-eyed surface forms to a more "boxy" form of the Micro-eyed and Blind forms is evident.

Fig. 2. Ancestral state reconstruction of morphology (Blind, Mico-eyed & Normal-eyed) on a time calibrated phylogeny. Posterior probabilities of node support also shown on the tree. **Maximum-likelihood reconstructions for the ancestral state of the eye-trait morphology (Blind, Micro and Normal-eyed morphs) on a time calibrated phylogeny.** A, B, C and D are four major clades. Posterior probabilities of node support values of 100% are indicated by *. Key events of *Sinocyclocheilus* evolution includes; at least three independent evolutionary events for Blind morphs; Blind, Micro-eyed and a few Normal-eyed species in clade B, with two cases of reversal from either Micro-eyed or Blind to Normal-eyed species.

Fig. 3. **Temporal patterns of body size and eye diameter in *Sinocyclocheilus* from the perspective of their eye trait related morphs.** The traitgrams suggests that the evolution of different morphs is attained by altering the allometric relationships between body size and eye

diameter. A. Traitgram of body size ; B. traitgram of eye size; C. Eye related morphs traced on

the phylomorphospace indicating clear separation of the three morphs in the morphospace.

(D)Habitat associations traced on the phylomorphospace showing species having eye diameter

<3mm and small to medium body sizes are obligate cave dwellers whereas species with eye

diameter >3mm can be Troglodytic, Troglophilic or Surface dwellers regardless of their body

size   (E) Horn related morphs traced on the phylomorphospace indicating the presence of a horn

in smaller fish with reduced eye size. Horned species are Troglodytes.

Fig. 4. Geophylogeny, the phylogeny laid across the geographic distribution of the species

considered in the analysis, with eye-related morphology, habitats, and horn-existence of the

species traced. A pattern where basal, Normal-eyed, Troglodytic species are placed in the east,

predominantly Blind/Micro-eyed/Normal-Eyed, Troglodytic species in the center and Normal-

eyed, Surface dwelling species towards the western mountains is evident, indicating an East to

west dispersion of the genus *Sinocyclocheilus* across South and South Western China. Eye

specializations mostly occurred in Clade B, and horn evolution occurred exclusively in Clade B,

within the Central range of the *Sinocyclocheilus* distribution.

Main Tables:

Table 1. Model fit and estimated Brownian rate parameters for three traits (ED, sED and SL) in eye related morphs of *Sinocyclocheilus*. Multiple-rate models of evolution providing the best fit to the data for all three quantitative traits with ΔAIC >2.

Morph

Trait	Single rate model		Multiple rate model		ΔAIC
	AICc	AICweights	AICc	AICweights	
ED	253.9337	0.19991263	251.1600	0.800088	2.773681
sED	246.1558	0.004509138	235.3616	0.995491	10.794260
SL	538.3145	0.018806041	530.4053	0.981194	7.909184

Table 2. Model averaged rate parameters for the measured traits in eye related morphs of *Sinocyclocheilus*. Normal-eyed and Micro-Eyed species indicate similar evolutionary rates with marked shifts towards Blind species.

Trait	Model Averaged Rate		
	Blind	Micro-eyed	Normal-eyed
SL	45.3506	1597.9230	1275.5080
ED	11.3935	2.9984	3.8872
sED	24.9441	1.6925	2.2773

Fig.1

Fig.2.

Fig. 3.

621
622
623

624
625 **Fig.4.**

Supplementary Information:

Fig. S1. Maximum-likelihood reconstructions for the ancestral states of the habitat occupation

(Troglodytic, Troglophilic and Surface) on the phylogeny of the genus *Sinocyclocheilus*.

Fig. S2. Maximum-likelihood reconstructions for the ancestral state of the horn related trait

(presence/absence of horn) on the phylogeny of the genus *Sinocyclocheilus*.

Table S1. Species information and GenBank accession numbers of two mtDNA fragments

(*NADH4* and *cytb*) of 39 *Sinocyclocheilus* species. The information of the outgroup species

Linichthys laticeps (Cyprinidae) is also indicated along with calculated standard length (SL), eye

diameter (ED), standard eye diameter (sED), discrete trait categories related to eye morphology

(Blind, Micro-eyed, Normal eyed), presence or absence of a horn and habitat occupation

(Troglodytic, Troglophilic and surface) for all species used in the analysis. Accession numbers

indicated as XXXX will be accessible upon the acceptance of the manuscript.

Species name	SL	ED	SED	CYTB	NADH4	Morph	Habitat	Horn
Linichthys laticeps	110.20	4.91	4.45	AY854739	AY854796	Normal-eyed	Surface	Absent
S. altishoulderus	125.10± 16.74	3.07±0. 35	2.46	AY854724	AY854781	Micro-eyed	Troglodytic	Absent
S. anatirostris	88.40	0.00	0	AY854708	AY854765	Blind	Troglodytic	Present
S. angustiporus	99.57±19. 79	5.05±0. 77	5.07	AY854702	AY854759	Normal-eyed	Surface	Absent
S. anophthalmus	99.80	0.00	0	AY854715	AY854772	Blind	Troglodytic	Absent
S. anshuiensis	87.40	0.00	0	NC_027169	NC_039769	Blind	Troglodytic	Present
S. bicornutus	98.80	1.39	1.41	AY854730	AY854787	Micro-eyed	Troglodytic	Present
S. brevibarbatus	137.55±18 .17	4.50±0. 84	3.27	XXXX	-	Normal-eyed	Troglophilic	Absent

Species								
S. brevis	92.91±6.39	3.55±0.14	3.83	XXXX	-	Normal-eyed	Troglodytic	Absent
S. guanyangensis	148.92±7.39	1.90±1.14	1.27	AY854711	AY854768	Micro-eyed	Troglodytic	Absent
S. cyphotergous	106.95±13.08	1.34±0.61	1.25	AB196440	-	Micro-eyed	Troglodytic	Absent
S. donglanensis	173.37±15.94	7.09±0.69	4.09	AY854709	AY854766	Normal-eyed	Troglodytic	Absent
S. furcodorsalis	91.29±16.36	0.00	0	AY854694	AY854751	Blind	Troglodytic	Present
S. grahami	112.60	6.64	5.9	XXXX	-	Normal-eyed	Surface	Absent
S. guilinensis	120.68±15.04	4.74±0.33	3.93	XXXX	-	Normal-eyed	Troglodytic	Absent
S. guishanensis	64.90	3.48	5.35	AY854722	AY854779	Normal-eyed	Surface	Absent
S. huangtianensis	67.32±20.48	4.03±0.72	5.99	XXXX	-	Normal-eyed	Troglodytic	Absent
S. huaningensis	92.20	5.30	5.75	AY854718	AY854775	Normal-eyed	Surface	Absent
S. huanjiangensis	80.79±17.68	3.99±0.45	4.94	XXXX	-	Normal-eyed	Troglodytic	Absent
S. huizeensis	85.60	4.61	5.39	NC_044072	NC_039769	Normal-eyed	Troglophilic	Absent
S. hyalinus	59.80±28.43	0.00	0	AY854721	AY854778	Blind	Troglodytic	Present
S. jii	111.43±17.21	4.97±0.47	4.46	AY854727	AY854784	Normal-eyed	Troglodytic	Absent
S. jiuxuensis	125.10	2.84	2.27	AY854736	AY854793	Micro-eyed	Troglodytic	Absent
S. lateristritus	120.00	6.52	5.43	AY854703	AY854760	Normal-eyed	Surface	Absent
S. lingyunensis	77.69±9.35	2.75±0.60	3.53	AY854691	AY854748	Normal-eyed	Troglodytic	Absent
S. longibarbatus	90.63±1.82	4.84±0.24	5.34	AY854714	AY854771	Normal-eyed	Troglophilic	Absent
S. macrocephalus	202.30	9.42	4.65	AY854683	AY854740	Normal-eyed	Surface	Absent
S. macrolepis	93.30	4.06	4.35	AY854729	AY854786	Normal-eyed	Troglophilic	Absent
S. macrophthalmus	73.16±11.50	3.96±0.71	5.41	AY854735	AY854792	Normal-eyed	Troglodytic	Absent
S. maculatus	82.70	2.84	3.43	EU366193	EU366183	Normal-eyed	Surface	Absent

S. maitianheensis	90.00	3.96	4.4	AY854710	AY854767	Normal-eyed	Surface	Absent
S. malacopterus	129.20	4.65	3.6	AY854697	AY854754	Normal-eyed	Surface	Absent
S. mashanensis	107.36±5.72	3.13±0.34	2.91	XXXX	-	Micro-eyed	Troglodytic	Absent
S. microphthalmus	134.89±10.51	1.89±0.38	1.4	AY854687	AY854744	Micro-eyed	Troglodytic	Absent
S. multipunctatus	188.60	4.11	2.18	AY854712	AY854769	Micro-eyed	Troglophilic	Absent
S. oxycephalus	103.10	3.80	3.69	AY854685	AY854742	Normal-eyed	Surface	Absent
S. purpureus	97.60	3.61	3.7	EU366194	EU366177	Normal-eyed	Surface	Absent
S. qiubeiensis	103.30	3.84	3.72	EU366195	EU366181	Normal-eyed	Surface	Absent
S. qujingensis	99.90	4.91	4.92	AY854719	AY854776	Normal-eyed	Surface	Absent
S. rhinocerous	78.20	1.55	1.98	AY854720	AY854777	Micro-eyed	Troglodytic	Present
S. ronganensis	112.72	5.92	5.25	NC_032385	NC_039769	Normal-eyed	Troglodytic	Absent
S. tianeensis	102.34±12.43	0.00	0	AY854717	AY854774	Blind	Troglodytic	Present
S. tianlinensis	87.96±27.97	0.00	0	XXXX	-	Blind	Troglodytic	Present
S. tingi	109.00	6.42	5.89	AY854701	AY854758	Normal-eyed	Surface	Absent
S. wumengshanensis	92.80	4.52	4.87	NC_039769	NC_039769	Normal-eyed	Surface	Absent
S. xunlensis	69.01±21.44	0.00	0	EU366187	EU366184	Blind	Troglodytic	Absent
S. yangzongensis	144.50	5.56	3.85	AY854725	AY854782	Normal-eyed	Surface	Absent
S. yimenensis	92.50	3.31	3.58	EU366192	EU366179	Normal-eyed	Troglophilic	Absent
S. yishanensis	105.34±8.15	5.23±0.23	4.97	XXXX	-	Normal-eyed	Troglodytic	Absent
S. zhenfengensis	56.78	2.41	4.24	MK610342	MK610347	Normal-eyed	Troglodytic	Absent

659   Table S2. Information of digitized images used in the morphometric geometric analysis. Table

660   indicates the voucher number of the specimen used for the analysis and the reference from which

661   the image was obtained. Images photographed during the current study are also indicated with

662   voucher numbers stated as GXUXXX (GXU: Guangxi University, China).

Species name	Reference	Voucher #
Linichthys laticeps	Zhang, E., and F. Fang. 2005. Linichthys: A New Genus of Chinese Cyprinid Fishes (Teleostei: Cypriniformes). Copeia 2005:61–67.	*IHB 78X6242*
S. altishoulderus	Guangxi University (this study).	*GXU001, GXU002, GXU003*
S. anatirostris	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*IHB 84VII255,*
S. angustiporus	Guangxi University (this study).	*GXU007, GXU008 , GXU009*
S. anophthalmus	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*KIZ 865949*
S. anshuiensis	Xi, G., W. Tie-Jun, W. Mu-Lan, and Y. Jian. 2013. A new blind barbine species, *Sinocyclocheilus anshuiensis* sp. nov.(Cypriniformes: Cyprinidae) from Guangxi, China. Kunming Institute of Zoology, Chinese Academy of Sciences.	*12070276*
S. bicornutus	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*IHB 12209043-9o5o241*
S. brevibarbatus	Guangxi University (this study).	*GXU010, GXU011, GXU012*
S. brevis	Guangxi University (this study).	*GXU013,*

Species	Reference	Voucher
	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*GXU014* *IHB12209033-87087496*
S. cyphotergous	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278. Huang, J., A. Gluesenkamp, D. Fenolio, Z. Wu, and Y. Zhao. 2017. Neotype designation and redescription of *Sinocyclocheilus cyphotergous* (Dai) 1988, a rare and bizarre cavefish species distributed in China (Cypriniformes: Cyprinidae). Environ Biol Fish 100:1483–1488.	*IHB 12209040* *ASIZB 204678*
S. donglanensis	Guangxi University (this study).	*GXU015, GXU016, GXU017*
S. furcodorsalis	Guangxi University (this study).	*GXU018, GXU019, GXU020*
S. grahami	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*ASIZB 03496*
S. guanyangensis	Guangxi University (this study).	*GXU004,GXU005, GXU006*
S. guilinensis	Guangxi University (this study).	*GXU021, GXU022, GXU023*
S. guishanensis	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean	*Li980514005*

	fishes of China. Environ Biol Fish 86:211–278.	
S. huangtianensis	Guangxi University (this study).	*GXU024, GXU025, GXU026*
S. huaningensis	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*ASIZB79228*
S. huanjiangensis	Guangxi University (this study).	*GXU027, GXU028, GXU029*
S. huizeensis	Cheng, C., P. Xiao-Fu, C. Xiaoyong, J. Li, L. Ma, and J. Yang. 2015. A new species of the genus *Sinocyclocheilus*(Teleostei:Cypriniformes), from Jinshajiang Drainage, Yunnan, China. Cave Research 1:1–4.	*KIZ2013001246*
S. hyalinus	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278. You, H., C. Xiaoyong, X. Ti-Qao, and Y. Jun-Xing. 2013. Three-dimensional morphology of the *Sinocyclocheilus hyalinus* (Cypriniformes : Cyprinidae) horn based on synchrotron X-ray microtomography. Zoological research 34:128–134.	*KIZ916001* *Photograph in life*
S. jii	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*ASIZB62726*

	Baradi, W., Z. Yahui, Z. Chunguang, D. Rongji, and A. Abdul. 2013. Anatomical Studies of the Olfactory Epithelium of Two Cave Fishes *Sinocyclocheilus jii* and *S. furcodorsalis* (Cypriniformes: Cyprinidae) from China. Pakistan Journal of Zoology 45:1091 –1101.	*Photograph in life*
S. jiuxuensis	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*ASIZB102260*
S. lateristritus	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*IHB12209036-865027*
S. lingyunensis	Guangxi University (this study). Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*GXU030* *ASIZB 73038*
S. longibarbatus	Guangxi University (this study).	*GXU031, GXU032, GXU033*
S. macrocephalus	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*IHB12209012-662001*
S. macrolepis	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*IHB12209035-87IV457*
S. macrophthalmus	Guangxi University (this study).	*GXU034,GXU035,GXU036*
S. maculatus	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*Li870808001*
S. maitianheensis	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*IHB12209039-874001*
S. malacopterus	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean	*KIZ775831*

	fishes of China. Environ Biol Fish 86:211–278.	
S. mashanensis	Guangxi University (this study).	*GXU037, GXU038, GXU039*
S. microphthalmus	Guangxi University (this study).	*GXU040, GXU041, GXU042*
S. multipunctatus	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*ASIZB73000*
S. oxycephalus	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*IHB12209013-652047*
S. purpureus	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*IHB12209015-731004*
S. qiubeiensis	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*Li990527002*
S. qujingensis	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*ASIZB78790*
S. rhinocerous	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*ASIZB93907*
S. ronganensis	FuGuang, L., H. Jie, L. Xia, L. Tong, and W. YanHong. 2016. *Sinocyclocheilus ronganensis* Luo, Huang et Wen sp. nov., a new species belonging to *Sinocyclocheilus* Fang from Guangxi (Cypriniformes: Cyprinidae). Guangxi Academy of Agricultural Sciences 47:650–655.	*20151114001*
S. tianeensis	Guangxi University (this study).	*GXU043, GXU044, GXU045*

S. tianlinensis	Guangxi University (this study).	*GXU052, GXU053, GXU054*
S. tingi	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*ASIZB60227*
S. wumengshanensis	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*KIZ82100006*
S. xunlensis	Guangxi University (this study).	*GXU046, GXU047, GXU048*
S. yangzongensis	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*KIZ6351069*
S. yimenensis	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*Li030509009*
S. yishanensis	Romero, A., Y. Zhao, and X. Chen. 2009. The Hypogean fishes of China. Environ Biol Fish 86:211–278.	*GXU049, GXU050, GXU051*
S. zhenfengensis	Liu, T., H. Q. Deng, L. Ma, N. Xiao, and J. Zhou. 2018. *Sinocyclocheilus zhenfengensis* , a new cyprinid species (Pisces: Teleostei) from Guizhou Province, Southwest China. J Appl Ichthyol 34:945–953.	*GZNU20120701 001*

666   Table S3. Calculated Principal Component values (PC1, PC2 and PC3) of all the specimens used

667   in the current analysis

Spname	PC1val	PC2val	PC3cal
Linichthys laticeps	0.1069618	-0.006044939	-0.032010383
S. altishoulderus	-0.01804998	-0.010081536	-0.046423134

S. anatirostris	-0.01766386	0.046175863	0.051169448
S. angustiporus	0.04334953	-0.002675304	0.034186136
S. anophthalmus	0.02908059	0.031606384	0.00917709
S. anshuiensis	-0.06639039	-0.000250595	-0.004873197
S. bicornutus	-0.07667729	0.060886565	0.012973894
S. brevibarbatus	-0.05048176	-0.011045867	0.006805293
S. brevis	-0.0142754	-0.011768784	-0.001763996
S. guanyangensis	-4.38E-05	0.015156895	-0.025200906
S. cyphotergous	-0.1220022	-0.139796596	0.008525168
S. donglanensis	0.004198408	0.013428914	-0.021746042
S. furcodorsalis	-0.050741	-0.000898602	-0.044412032
S. grahami	0.02231859	0.026236004	0.022098623
S. guilinensis	0.01963493	-0.0033727	-0.023009295
S. guishanensis	0.07886827	0.00257981	-0.059167855
S. huangtianensis	0.05562456	-0.011114954	0.021111628
S. huaningensis	0.04117411	-0.037450418	0.031061221
S. huanjiangensis	0.009958768	-0.021868616	0.007063138
S. huizeensis	-0.003779219	0.010286192	0.031012201
S. hyalinus	-0.03974731	0.030183657	0.039671171
S. jii	0.02155773	-0.019701057	-0.005284772
S. jiuxuensis	-0.06337799	0.004932247	-0.008080491
S. lateristritus	0.01609448	-0.006156199	0.021918263
S. lingyunensis	0.01666848	0.015544267	0.002003128
S. longibarbatus	0.0238826	-0.037134912	0.007645544
S. macrocephalus	0.0223646	-0.03930768	0.034945546
S. macrolepis	0.002586613	0.0056233	-0.005281302

S. macrophthalmus	0.01921085	-0.004423179	0.011957836
S. maculatus	-0.01164108	-0.003338214	-0.043022304
S. maitianheensis	0.04058493	-0.015399253	-0.004022456
S. malacopterus	0.03369873	0.016341474	0.002649988
S. mashanensis	-0.03299028	-0.033529198	-0.005547116
S. microphthalmus	-0.02936266	0.061223556	-0.040563771
S. multipunctatus	-0.05126049	0.014657848	-0.010821531
S. oxycephalus	0.04140438	0.030989746	-0.036348165
S. purpureus	0.02288238	-0.045035385	-0.018448807
S. qiubeiensis	0.03248934	-0.003880679	0.012191068
S. qujingensis	0.000644406	0.033555511	0.046291334
S. rhinocerous	-0.08985807	0.03649387	-0.020510521
S. ronganensis	0.05067357	-0.004282673	-0.02013066
S. tianeensis	-0.06234116	0.02827463	-0.030564848
S. tianlinensis	-0.01905521	0.012948686	-0.00406653
S. tingi	0.01508119	-0.019031787	-0.002178747
S. wumengshanensis	-0.000171791	-0.027155405	0.003238896
S. xunlensis	0.008420042	0.022462118	0.047598226
S. yangzongensis	0.02622031	-0.003887167	-0.006000758
S. yimenensis	-0.01762049	0.008265106	0.019258774
S. yishanensis	0.007505367	-0.029610044	-0.01952727
S. zhenfengensis	0.01938822	-0.002301361	0.003044235

669
670
671
672

673 **Fig.S1.**

Habitat Associations
- ■ Troglodytic
- ■ Troglophilic
- □ Surface

674
675

676　**Fig.S2.**