Title: Resistance, resilience, and functional redundancy of freshwater bacterioplankton communities facing a gradient of agricultural stressors in a mesocosm experiment

Running title: Bacterioplankton responses to pesticides

Authors: Naíla Barbosa da Costa[1,2], Vincent Fugère[2,3,4,5], Marie-Pier Hébert[2,6], Charles C.Y. Xu[3,6,7], Rowan D.H. Barrett[3,6,7], Beatrix E. Beisner[2,4], Graham Bell[3,6], Viviane Yargeau[8], Gregor F. Fussmann[2,3,6], Andrew Gonzalez[3,6], B. Jesse Shapiro[1,2,3,9,10]†

[1] Département des sciences biologiques, Université de Montréal, Montreal, Canada

[2] Groupe de Recherche Interuniversitaire en Limnologie et environnement aquatique (GRIL)

[3] Québec Centre for Biodiversity Science (QCBS)

[4] Département des sciences biologiques, Université du Québec à Montréal, Montréal, Canada

[5] Département des sciences de l'environnement, Université du Québec à Trois-Rivières, Trois-Rivières, Canada

[6] Department of Biology, McGill University, Montreal, Canada

[7] Redpath Museum, McGill University, Montreal, Canada

[8] Department of Chemical Engineering, McGill University, Montreal, Canada

[9] Department of Microbiology and Immunology, McGill University, Montreal, Canada

[10] McGill Genome Centre, McGill University, Montreal, Canada

† Corresponding author: jesse.shapiro@mcgill.ca

Abstract

Agricultural pollution with fertilizers and pesticides is a common disturbance to

freshwater biodiversity. Bacterioplankton communities are at the base of aquatic food

webs, but their responses to these potentially interacting stressors are rarely explored.

To test the extent of resistance and resilience in bacterioplankton communities faced

with agricultural stressors, we exposed freshwater mesocosms to single and combined

gradients of two commonly used pesticides: the herbicide glyphosate (0-15 mg/L) and

the neonicotinoid insecticide imidacloprid (0-60 μg/L), in high or low nutrient

backgrounds. Over the 43-day experiment, we tracked variation in bacterial density with

flow cytometry, carbon substrate use with Biolog EcoPlates, and taxonomic diversity and

composition with environmental 16S rRNA gene amplicon sequencing. We show that

only glyphosate (at the highest dose, 15 mg/L), but not imidacloprid, nutrients, or their

interactions measurably changed community structure, favoring members of the

Proteobacteria including the genus *Agrobacterium*. However, no change in carbon

substrate use was detected throughout, suggesting functional redundancy despite

taxonomic changes. We further show that communities are resilient at broad, but not

fine taxonomic levels: 24 days after glyphosate application the precise amplicon

sequence variants do not return, and tend to be replaced by phylogenetically close taxa.

We conclude that high doses of glyphosate – but still within commonly acceptable

regulatory guidelines – alter freshwater bacterioplankton by favoring a subset of higher

taxonomic units (i.e. genus to phylum) that transiently thrive in the presence of

glyphosate. Longer-term impacts of glyphosate at finer taxonomic resolution merit

further investigation.

Introduction

Agricultural expansion and intensification are major drivers of global environmental

change in both terrestrial and aquatic ecosystems (Song et al., 2018; Springmann et al.,

2018; Tilman et al., 2001). Chemicals derived from agricultural landscapes, such as

fertilizers and pesticides, are among the main sources of freshwater pollution

(Vörösmarty et al., 2010), leading to eutrophication (Carpenter et al., 1998; Keatley,

Bennett, Macdonald, Taranu, & Gregory-Eaves, 2011) and biodiversity loss (DeLorenzo,

Scott, & Ross, 2001; Relyea, 2009; Stehle & Schulz, 2015). Anthropogenic climate

change may intensify these effects as variation in precipitation patterns and increased

temperatures affect agrochemicals fate, transport, and behavior in surface and

groundwater (Bloomfield, Williams, Gooddy, Cape, & Guha, 2006; Jeppesen et al.,

2009). Agricultural runoff to waterbodies particularly increases after storms, acting as a

pulse perturbation (Cedergreen & Rasmussen, 2017) while bringing a mixture of

nutrients, herbicides and insecticides that may interact to affect aquatic microbial taxa

(Flood & Burkholder, 2018) and communities (Lozano & Pratt, 1994; Starr, Bargu, Maiti,

& DeLaune, 2017). The impact of agricultural contaminants may depend on whether

they are applied alone or in combination (Altenburger, Backhaus, Boedeker, Faust, &

Scholze, 2013), and the effects of combinations may be difficult to predict based upon

data from single contaminants, possibly due to complex interactions within diverse

bacterial communities (Romero, Acuña, & Sabater, 2020).

Agricultural activity has a major impact on bacterioplankton (Kraemer et al., 2020) and,

as a consequence, on the ecosystem processes they provide; e.g. decomposition of

organic matter (Piggott, Niyogi, Townsend, & Matthaei, 2015) and nutrient cycling

(Romero et al., 2020). Altering these processes may have broad consequences for

aquatic ecosystem productivity, food webs, and the human activities that depend upon

them (Carpenter, Stanley, & Vander Zanden, 2011).

Nutrient pollution is among the most important stressors affecting biodiversity in lakes

(Birk et al., 2020). It promotes eutrophication (Smith, Joye, & Howarth, 2006), which can

increase bacterial biomass, reduce phytoplankton diversity, and trigger harmful algal

blooms (Paerl, Otten, & Kudela, 2018; Smith & Schindler, 2009). While few studies have

addressed individual and combined effects of fertilizers with herbicides or insecticides

on phytoplankton and zooplankton communities (Baker, Mudge, Thompson, Houlahan,

& Kidd, 2016; Chará-Serna, Epele, Morrissey, & Richardson, 2019; Geyer, Smith, &

Rettig, 2016), analogous assessments of bacterioplankton are more scarce. Yet, similar

to other planktonic communities, bacteria may also be directly or indirectly (e.g. through

trophic effects) affected by the individual or combined effects of these stressors, despite

not being their intended targets (Muturi, Donthu, Fields, Moise, & Kim, 2017).

The herbicide glyphosate, mainly formulated commercially as Roundup, and the

neonicotinoid insecticide imidacloprid (available in different commercial formulations) are

among the most commonly used pesticides worldwide (Benbrook, 2016; Simon-Delso et

al., 2015), despite restrictions on their use in different jurisdictions. In North America and

the European Union, common benchmarks to protect aquatic life range from 800 to

26,600 μg/L of glyphosate for long-term (chronic) exposure, and between 27,000 to

49,000 μg/L for short-term (acute) exposure (CCME, 2012; EFSA, 2016; EPA, 2019). In

contrast, lower concentrations of imidacloprid are considered safe for aquatic

invertebrates, ranging from 0.009-0.385 μg/L (CCME, 2007; EFSA, 2014; EPA, 2019)

(Table S1). Most of these criteria were developed based on toxicity tests on individual

eukaryotic organisms, and it remains unclear how bacterial communities respond to

these concentrations considered "safe for aquatic life" and what consequences their

responses might have on the ecosystem functions they provide.

Glyphosate is a broad-spectrum synthetic phosphonate herbicide used for weed control.

It acts by inhibiting the enzyme enolpyruvylshikimate-3-phosphate synthase (EPSPS)

involved in the biosynthesis of aromatic amino acids essential to plants, its target group,

but also to many fungi and bacteria (Pollegioni, Schonbrunn, & Siehl, 2011). However,

some microorganisms are resistant to glyphosate either by expressing an insensitive

form of the target enzyme (Funke, Han, Healy-Fried, Fischer, & Schönbrunn, 2006;

Healy-Fried, Funke, Priestman, Han, & Scho, 2007) or by metabolizing the molecule and

using it as a phosphorus source (Hove-Jensen, Zechel, & Jochimsen, 2014).

Glyphosate could therefore select for resistant species within bacterial communities

(Muturi et al., 2017). Moreover, as it may prevent the growth of some phytoplankton

species (Smedbol, Lucotte, Labrecque, Lepage, & Juneau, 2017), bacterioplankton

could be affected indirectly, for example by reduced competition with phytoplankton.

Unlike glyphosate, imidacloprid is an insecticide commonly used as a seed-coating

agent intended to control sapling damage from piercing-sucking insects (CCME, 2007;

Jeschke & Nauen, 2008). It acts on insect nervous systems (Roberts & Hutson, 1999)

and can be toxic to many aquatic invertebrates, especially insects and crustaceans

(Morrissey et al., 2015). Although it is not known to inhibit bacteria directly, it could affect

them indirectly via trophic effects on their predators or grazers. If imidacloprid reduces

total zooplankton biomass, for example, a reduction in predation pressure could promote

an increase in bacterioplankton biomass. Ecosystem functions provided by

bacterioplankton, such as carbon use, could subsequently be affected, as has been

observed in experiments with other insecticides (Thompson et al., 2016).

Bacterioplankton are important drivers of energy and nutrient cycling in freshwater

ecosystems (Falkowski, Fenchel, & Delong, 2008; Konopka, 2009), and more

observations are needed to understand how they respond to anthropogenic

disturbances (Allison & Martiny, 2008). They may respond with detectable changes in

species composition (Allison & Martiny, 2008) that could be permanent, thereby

providing a measure of the historical impact of anthropogenic activities on ecosystem

health (Kraemer et al., 2020). Alternatively, community composition could be resistant or

resilient to changes (Shade et al., 2012). Even if disturbances alter community

composition, ecosystem processes may remain stable if pre- and post-disturbance

communities are functionally redundant (Allison & Martiny, 2008).

Functional redundancy is thought to be common in microbial communities, as most

metabolic pathways controlling biogeochemical cycles are encoded by several different

phylogenetic groups. Certain functions, such as photosynthesis and methanogenesis,

are however phylogenetically restricted (Falkowski et al., 2008). It is likely that

communities are partially redundant for general functions like respiration or biomass

production, but non-redundant for more specific functions encoded by unique taxa

(Louca et al., 2018). The prevalence of, and reasons for microbial community

resistance, resilience, and functional redundancy are still debated (Allison & Martiny,

2008; Shade et al., 2012), particularly in response to novel anthropogenic disturbances

which increasingly involve combinations of stressors.

In this study, we experimentally tested the effects of pulse applications of glyphosate

and imidacloprid, under low (mesotrophic) or high (eutrophic) nutrient conditions, on

bacterioplankton community density, taxonomic composition and richness, and functions

related to carbon substrate use. To do so, we filled 1,000 L mesocosms with water and

planktonic organisms from a pristine lake located on a mountaintop of a protected area

with no history of agricultural activity. Using a regression design, we applied gradients of

pesticide concentrations (Fig. 1), spanning ranges observed in surface runoff and

freshwater systems (Hénault-Ethier et al., 2017; Morrissey et al., 2015; van Bruggen et

al., 2018). Highest doses applied are considered harmful to eukaryotic organisms upon

which nationwide water quality guidelines are based (Table S1). To quantify individual

and interactive effects of agricultural stressors, we applied these pesticides alone and in

combination, and in the presence or absence of nutrient enrichment simulating fertilizer

pollution. Pesticides were applied as pulse perturbations to mimic how these

contaminants reach natural freshwater ecosystems from agricultural fields, while nutrient

enrichments were applied as press treatments to mimic mesotrophic and eutrophic

conditions.

We hypothesize that glyphosate will change bacterial community composition and

reduce richness, as many taxa depend on the target enzyme (EPSPS) to synthesize

aromatic amino acids, while other species encode a resistant allele of EPSPS (Funke et

al., 2006; Healy-Fried et al., 2007; Rainio et al., 2021) or are able to metabolize

glyphosate (Hove-Jensen et al., 2014). While imidacloprid is less likely to directly impact

bacteria, we hypothesize that it can exert indirect effects due to its potential toxicity to

aquatic invertebrates (Chará-Serna et al., 2019), releasing grazing pressure on bacterial

communities and increasing their density. When applied in combination with glyphosate,

imidacloprid may therefore delay or mask the effects of glyphosate on bacterioplankton

community structure. Similarly, fertilizers might also increase microbial productivity and

mask negative effects of glyphosate, as it does with other contaminants (Alexander,

Luis, Culp, Baird, & Cessna, 2013). We also expect some positive effects of glyphosate

on bacterial density, as it may serve as a source of phosphorus for some species

(Hébert, Fugère, & Gonzalez, 2018; Lu et al., 2020). Finally, we predict that functional

diversity will be less prone to changes than taxonomic composition, as bacterial

communities tend to be functionally redundant (Louca et al., 2018). We thus expect to

detect changes in bacterial community composition and species richness at lower

pesticide doses, and changes in functional diversity only at higher doses, or not at all.

Materials and methods

Experimental design and sampling

We conducted a mesocosm experiment at the Large Experimental Array of Ponds

(LEAP) platform at McGill University's Gault Nature Reserve (45°32'N, 73°08'W), a

protected area with no history of agricultural pollution (Beauséjour, Handa, Lechowicz,

Gilbert, & Vellend, 2015) in Quebec, Canada. The pond mesocosms at LEAP are

connected to a reservoir that receives water from the upstream Lake Hertel through a 1

km pipe by gravity. On May 11th, 2016 (99 days prior to the start of the experiment), 100

ponds were simultaneously filled with 1,000 L of lake water, to acclimate communities to

the mesocosm setting. When filling ponds we used a coarse sieve to prevent fish

introduction. To maximize initial homogeneity among communities (before treatments),

and because this study focuses on planktonic microbes, no sediment substrate was

added to the ponds. Tadpoles and large debris such as leaves and pollen were

periodically removed with a net before the experiment commenced. Additional lake

water was added on a biweekly basis (~10% of total volume) between May and August

to ensure a continuous input of lake bacterioplankton, tracking seasonal changes in the

source lake community, and to homogenize communities across ponds. The experiment

reported here used 48 of these pond mesocosms from August 17th (day 1) to September

28[th] (day 43), and it is part of a collaborative experiment that also assessed responses

of zooplankton in the same set of ponds (Hébert et al., 2021) and phytoplankton

responses in a subset of these ponds for a longer period of time (Fugère et al., 2020).

Throughout our experiment, phosphorus (P) and nitrogen (N) were simultaneously

added biweekly to simulate nutrient enrichment at a constant rate, starting on August

10[th], 7 days before the first sampling day to ensure communities would have passed

their exponential growth phase before the first pulse of pesticides was applied. Our

nutrient treatment included two levels, with target concentrations of 15 µg P/L (hereafter

referred as low-nutrient treatment) typical of mesotrophic Lake Hertel (Thibodeau,

Walsh, & Beisner, 2015) and 60 µg P/L (high-nutrient treatment; eutrophic conditions).

Nutrient solutions were made using nitrate (KNO_3) and phosphate (KH_2PO_4 and K_2PO_4)

preserving the same N:P molar ratio (33:1) found in Lake Hertel; the target

concentrations were therefore 231 µg N/L and 924 µg N/L for low- and high- nutrient

treatments respectively. Over the course of the experiment, the average total P (TP)

concentration measured in the source lake was 20.4 µg/L (standard error, SE=±1.3) and

the average TP achieved in ponds with no pesticide addition was 13.6 µg/L (SE=±0.71)

and 36.7 µg/L (SE=±10.8) respectively for low- and high-nutrient treatment. The average

total N (TN) concentration was 556.9 µg/L (SE=±60.7) at Lake Hertel, 407.8 µg/L

(SE=±32.7) and 789.0 µg/L (SE=±177.6) respectively in control ponds with low and high

nutrient inputs.

Within each nutrient treatment, ponds received varying amounts of the herbicide

glyphosate or the insecticide imidacloprid, separately or in combination, in a regression

design with seven levels of pesticide concentration plus controls with no pesticide

addition (Fig. 1A). The seven levels of target concentration were: 0.04, 0.1, 0.3, 0.7, 2,

5.5 and 15 mg/L for glyphosate and 0.15, 0.4, 1, 3, 8, 22 and 60 μg/L for imidacloprid.

There was no replication for each combination of nutrient and pesticide concentration,

which is compensated by the wide gradient of pesticides concentration established in

the regression design (Fig. 1A). Glyphosate was added in the form of Roundup Super

Concentrate (Monsanto©) and target concentrations calculated based on its glyphosate

acid content, while imidacloprid was added in the form of a solution prepared with pure

imidacloprid powder (Sigma-Aldrich, Oakville, Canada) dissolved in ultrapure water.

Treatment ponds received two pulses of pesticides (at days 6 and 34 of the experiment)

while nutrients were applied biweekly to maintain a press treatment. The target

concentrations of glyphosate and imidacloprid were well correlated with the measured

concentrations in the ponds (Fig. S1A-B) with the exception of a few ponds receiving the

highest imidacloprid dose which reached lower concentrations than intended, especially

after the second pulse (Fig. S1C). These ponds nonetheless reached higher

concentrations than ponds lower on the imidacloprid gradients (i.e., a clear gradient was

established).

Bacterioplankton communities were sampled at six different timepoints (Fig. 1B): one

before pesticide application (day 1); three between pulse 1 and pulse 2 applications

(days 7, 15, and 30); and two timepoints after the second pulse (days 35 and 43).

Pesticide quantification was performed immediately after each pulse application (days 6

and 34) and at two time points between them (days 14 and 29) while nutrients were

quantified on the same days as bacterioplankton except for days one and 35 (Fig. 1B).

Water samples for nutrient and microbial community analyses were collected from each

mesocosm with integrated samplers (made of 2.5 cm-wide PVC tubing) and stored in

dark clean 1L Nalgene (Thermo Scientific) bottles triple-washed with pond water. To

avoid cross contamination, we sampled each pond with a separate sampler and bottle.

We kept bottles in coolers while sampling and then moved them to an on-site laboratory,

where they were stored at 4 °C until processing, for no longer than 4 hours. Water

samples for pesticide quantification were collected immediately after pesticide

application (days 6 and 34) in a subset of ponds englobing each gradient and in a

smaller subset between the pulses (days 14 and 29). They were stored in clear Nalgene

bottles (1 L), acidified to a pH < 3 with sulfuric acid and frozen at -20 ºC until analysis.

Nutrient and pesticide quantification

Quantification of TP and TN from unfiltered water samples were processed at the GRIL

(Interuniversity Group in Limnology) analytical laboratory at the Université du Québec à

Montréal following standard protocols as outlined by McComb (2002). Duplicate

subsamples (40 mL) of water sampled from each pond were stored in acid-washed

glass tubes and kept at 4 ºC until nutrient concentrations were quantified. TN

concentration was determined using the alkaline persulfate digestion method coupled

with a cadmium reactor (Patton & Kryskalla, 2003) in a continuous flow analyzer (OI

Analytical, College Station, TX, USA). TP was estimated based on optical density in a

spectrophotometer (Biocrom Ultrospec 2100pro, Holliston, MA, USA) after persulfate

digestion through the molybdenum blue method (Wetzel & Likens, 2000). Glyphosate

and imidacloprid concentrations were quantified through liquid chromatography coupled

to mass spectrometry using an Accela 600-Orbitrap LTQ XL (LC–HRMS, Thermo

Scientific). The method consisted of heated electrospray ionization (HESI) in negative

mode for glyphosate, acquisition in full scan mode (50-300 m/z) at high resolution

(FTMS = 30,000 m/Dz) and the same LC-HRMS system but using positive HESI mode

for imidacloprid (mass range 50-700m/z). Limits of detection were 1.23 and 1.44 µg/L for

glyphosate and imidacloprid respectively, while quantification thresholds were

respectively 4.06 µg/L, and 4.81 µg/L. Samples falling below limits of detection were pre-

concentrated with a factor of 40X (10 mL samples were reconstituted to 250 µL) and

their final concentration were back-calculated according to the concentration factor.

Estimating bacterial density through flow cytometry

To estimate the density of bacterial cells, we fixed 1 mL of the 1 L sampled pond water

with glutaraldehyde (1% final concentration) and flash froze this subsample in liquid

nitrogen (Gasol & Del Giorgio, 2000; Ruiz-González et al., 2018). We stored samples at

-80 ºC until they were processed via a BD Accuri C6 flow cytometer (BD Biosciences,

San Jose, CA, USA). After samples were thawed at room temperature (18-20 °C), we

prepared dilutions (1:25) with Tris-EDTA buffer (Tris-HCl 10 mM; EDTA 1 mM; pH 8)

and aliquoted in two duplicate tubes. Samples were then stained with Syto13 Green-

Fluorescent Nucleic Acid Stain (0.1 v/v in DMSO; ThermoFisher S7575) and incubated

in the dark at room temperature (18-20 °C) for 10 min. To validate the equipment

calibration, we ran BD TruCount Absolute Count Tubes (BD Biosciences) each day,

prior to sample processing. Samples were run until reaching 20,000 events, at a rate of

100-1,000 events/s in slow fluidics (14 µL/min). Events within a predefined gate on a 90°

light side scatter (SSC-H) versus green fluorescence (FL1-H) cytogram were used for

cell counts estimation. This inclusive gate was defined to maximize cell counts accuracy

by excluding background noise and large debris. Bacterial density was estimated based

on cell counts detected within the gate, flow volume, and sample dilution. We calculated

the average bacterial density for each pair of analytical duplicates with a coefficient of

variation (CV, i.e., ratio between the standard deviation and average of the duplicate

values) less than 0.08. If the CV was greater than or equal to 0.08, the sample was run

a third time, and the outlying value was discarded before taking the mean of the two

remaining samples.

Carbon substrate utilization patterns

We used Biolog EcoPlate® assays (Hayward, CA, USA) to infer community-level

utilization of dissolved organic carbon by microbes. For all treatments (Fig. 1A) and at

each of the six sampled timepoints (Fig. 1B), we added 125 µL of unfiltered pond water

to each well of the EcoPlates. Each plate contains, in triplicates, 31 different organic

carbon substrates and water controls. These substrates can be grouped into five main

guilds (amines/amides, amino acids, carbohydrates, carboxylic acetic acids and

polymers), as summarized in Table S2. We measured the optical density at 590 nm in

each well as a proxy for microbial carbon substrate use, since it causes a concomitant

reduction of the redox-sensitive tetrazolium dye, whose color intensity is measurable at

this wavelength. Plates were incubated in the dark at room temperature (18-20 °C) and

well absorbance was measured daily until an asymptote was reached (Ruiz-González et

al., 2018; Ruiz-González, Niño-García, Lapierre, & del Giorgio, 2015). For each daily

measurement, an average well color development (AWCD) was calculated. To correct

for variation in inoculum density we selected substrate absorbance values of the plate

measurements with AWCD closest to 0.5 (usually after 3-8 days of incubation) as

suggested in Garland (2001). We then calculated the blank-corrected median

absorbance of each substrate at each sampled timepoint for analyses.

DNA extraction, 16S rRNA gene amplification and sequencing

We selected a subset of ponds for DNA extraction and subsequent analyses (outlined in

bold in Fig. 1A) to assess bacterioplankton community responses at the extremes and

the middle of the experimental gradient. From each timepoint and nutrient treatment, we

chose two control ponds (beginning of the gradient, no pesticide addition), ponds with

the third lowest concentration (middle of the gradient) of each or both pesticides (1 μg/L

imidacloprid and/or 0.3 mg/L glyphosate), and ponds with the highest concentration (end

of the gradient) used in the experiment for each or both pesticides (60 μg/L imidacloprid

and/or 15 mg/L glyphosate). We selected ponds with high concentrations of pesticides

to maximize the chance of detecting a response from the bacterial community. That

said, we still kept concentrations that fall below available regulatory acceptable

concentrations for glyphosate in North America (Table S1), allowing us to ask whether

changes in bacterial communities can be detected at concentrations considered safe for

aquatic eukaryotes in a region where glyphosate is extensively used (Benbrook, 2016;

Simon-Delso et al., 2015). In total, we sampled 16 of the 48 experimental ponds at six

timepoints, yielding a total of 96 samples for 16S rRNA amplicon sequencing (Fig. 1B).

After sampling 1 L of pond water as described above, we immediately filtered 250 mL

through a 0.22 μm pore size Millipore hydrophilic polyethersulfone membrane of 47 mm

diameter (Sigma-Aldrich, St. Louis, USA) and stored filters at -80 ºC until DNA

extraction. We extracted and purified total genomic DNA from frozen filters using the

PowerWater DNA Isolation Kit (MoBio Technologies Inc., Vancouver, Canada) following

the manufacturer's protocol, that includes a 5-min vortex agitation of the filter with beads

and lysis buffer to enhance cell lysis. We quantified genomic DNA with a Qubit 2.0

fluorometer (ThermoFisher, Waltham, MA, USA) and used 10 ng to prepare amplicon

libraries for paired-end sequencing (2 x 250 bp) on two Illumina MiSeq (Illumina, San

Diego, CA, USA) runs. We performed a two-step polymerase chain reaction (PCR)

targeting the V4 region of the 16S rRNA gene, with primers U515_F and E786_R, as

described in Preheim et al. (2013). Further details on PCR reactions, library preparation

and amplicon sequencing, including positive controls (mock communities) and negative

controls are described in the Supplementary Material.

Sequence data processing

We used idemp (https://github.com/yhwu/idemp) to demultiplex barcoded fastq files from

the sequencing data, and cutadapt to remove remaining Illumina adapters (Martin,

2011). The DADA2 package (Callahan et al., 2016) in R was used to filter and trim

reads, using the default filtering parameters with a maximum expected error (maxEE)

score of two. Reads were trimmed on the left to remove primers and those shorter than

200 or 150 bp were discarded, respectively, for forward and reverse reads. DADA2 was

also used to infer amplicon sequence variants (ASVs), remove chimeras and finally

obtain a matrix of ASV counts in each sample for each MiSeq run independently. We

used the default parameters of the "learning error rates" function with the multithread

option enabled. The number of raw reads and non-chimeric reads obtained from each

sample are summarized in Table S3 (average raw reads per sample: 43,159;

SE=2,245). Excluding mock communities, extraction blanks and PCR controls, we

obtained 1,787,412 raw reads in the first run and 4,702,355 in the second run, of which

we retained, respectively, 1,565,021 and 4,188,644 non-chimeric reads. PCR negative

controls and extraction blanks produced 214 non-chimeric reads in total; these were

excluded from downstream analyses as we only included samples with a minimum of

6,000 reads. Of the 30 expected sequences from the custom mock community (Preheim

et al., 2013), DADA2 found 25 exact sequence matches, producing 5 false negatives

and 7 false positives (for a total of 32 sequences). In the ATCC mock, 23 of the 24

expected sequences were found, with only 1 false negative but 10 false positives (for a

total of 33 sequences). We concatenated DADA2 abundance matrices from each MiSeq

run and then used TaxAss (Rohwer, Hamilton, Newton, & McMahon, 2018) to assign

ASV taxonomy with a database specifically curated for freshwater bacterioplankton,

FreshTrain (Newton, Jones, Eiler, Mcmahon, & Bertilsson, 2011), and GreenGenes

(DeSantis et al., 2006), with a minimum bootstrap support of 80% and 50%,

respectively. After performing a multiple sequence alignment with the R package

DECIPHER (Wright, 2016), we constructed a maximum likelihood phylogenetic tree

using the phangorn package following recommendations made by Callahan (2016). For

subsequent analyses, we imported the ASV abundance matrix together with taxonomic

assignments and environmental data as an object in the phyloseq package (McMurdie &

Holmes, 2013) in R. We removed sequence data identified as mitochondria or

chloroplast DNA and normalized read counts using the DESeq2 package (Love, Huber,

& Anders, 2014), which performs a variance stabilizing transformation without discarding

reads or samples (McMurdie & Holmes, 2014), which is important in the context of high

read depth variation, as observed among our samples (Table S3). As normalizations

such as the DESeq2 method tend to reduce the importance of dominant taxa while

inflating the importance of rare taxa (McKnight et al., 2019), for comparison with

DESeq2, we additionally normalized the abundance matrix in two ways: (1) by

calculating relative abundances (proportions) of each ASV, and (2) by rarefying to

10,000 reads (948 ASVs and 7 samples were consequently removed). These two

alternative normalizations are presented in the Supplementary Materials, and are

generally concordant with the DESeq2 results in the main text. For most compositional

analyses in the main text, we calculated the estimated absolute abundance (EAA) of

ASVs per sample by multiplying the DESeq2 normalized ASVs relative abundance by

the total bacterial cell counts found in the sample through flow cytometry (Zhang et al.,

2017).

Statistical analyses

To assess resistance and resilience to experimental treatments, we compared changes

in bacterial community density, microbial carbon substrate use, as well as

bacterioplankton community taxonomic structure (richness and composition), as

explained in detail below. We conducted all statistical analyses in R version 3.5.1 (R

Core Team, 2008). As we tested hypotheses of different treatment effects at different

timepoints, we applied a Bonferroni correction for multiple hypothesis testing.

Treatment effects on bacterioplankton density

Time series of bacterial density were analyzed with a generalized additive mixed model

(GAMM) with the mgcv R package (Wood, 2017) to quantify the singular and interactive

effects of nutrient and of each pesticide treatment on bacterioplankton density as a

function of time while accounting for nonlinear relationships. Glyphosate and

imidacloprid target concentrations were rescaled (from 0 to 1) to match the scale of the

nutrient treatment factor (binary) and we tested for their effect individually or in

combination. Individual mesocosms (ponds) were included as a random effect (random

smooth) to account for non-independence among measurements from the same pond

over time. Model validation was performed by investigating residual distributions,

comparing fitted and observed values and checking if basis dimensions (k) of smooth

terms were not too low. The model fit (adjusted R^2) and further details on predictors

used in the model, including their statistical significance, are provided in Table S4.

Treatment effects on carbon substrate use

We quantified treatment effects on the number of carbon substrates used at each pond

and timepoint with a GAMM with the same terms as the GAMM described above for

modeling bacterial density. More details are provided in Table S4. To assess the effects

of the treatments on carbon substrate utilization patterns by microbial communities over

time, we built principal response curves (PRCs) (Auber, Travers-Trolet, Villanueva, &

Ernande, 2017). PRCs are a special case of partial redundancy analysis (pRDA) in

which time and treatments, expressed as ordered factors, are used as explanatory

variables, while community composition is the multivariate response. Time is considered

as a covariable (or conditioning variable) whose effect is partialled out, and changes in

community composition with the treatments over time are always expressed as

deviations from the control pond at each timepoint. PRCs also assess the contribution of

each species to the treatment effect through the taxa weight (also known as species

score) usually displayed in the right y-axis of a PRC diagram (Van den Brink, den

Besten, bij de Vaate, & ter Braak, 2009). The significance of the first PRC axis was

inferred by permuting the treatment label of each pond while keeping the temporal order,

using the permute R package (Simpson, 2019) followed by a permutation test with the

vegan R package (Oksanen et al., 2018). Before performing PRCs we transformed the

community matrix (containing carbon substrate use data) using the Hellinger

transformation (Legendre & Gallagher, 2001). PRC of community carbon utilization

patterns was performed for the 31 substrates individually and grouped into five guilds

(Table S2).

Treatment effects on bacterioplankton community taxonomic structure

To infer the impact of treatments on bacterioplankton taxonomic diversity over time, we

calculated alpha diversity as richness (number of observed ASVs) and as the exponent

of the Shannon index (or Hill numbers (Jost, 2006)) of each sample after rarefying the

ASV abundance matrix to 10,000 reads without replacement and modelled their

response to pesticide and nutrient treatments using GAMMs. Model equations, their fit

(adjusted R^2) and statistics of significant terms are reported in Table S5. In this analysis,

pesticides treatments were considered factors (low vs. high) because 16S rRNA reads

data were only available for a subset of concentrations (Fig. 1A). Pesticides and nutrient

treatments were coded as ordered factors and models were validated after investigation

of residual distributions, comparison of fitted and observed values and checking if the

basis dimension (k) of smooth terms was sufficiently large.

To assess differences in community composition, we calculated weighted UniFrac

distances (Lozupone & Knight, 2005) and Jensen-Shannon divergence (JSD) among

the subset of samples selected for DNA analyses and represented them in principal

coordinate analysis (PCoA) bidimensional plots. These two metrics are complementary

as the first is weighted for phylogenetic branch lengths unique to a particular treatment,

and the second assesses changes in community composition at the finest possible

resolution, tracking ASVs regardless of their phylogenetic relatedness. We performed a

series of permutational analyses of variance (PERMANOVA) based on weighted

UniFrac distances and JSD to test the effect of treatments (as factors) on community

composition at four sampled timepoints separately: at day 1 (before any treatment was

applied), day 7 (immediately after the first pulse), day 15 (11 days after the first pulse),

day 30 (immediately before the second pulse) and day 43 (last day of the experiment,

after the second pulse). We also performed an analysis of multivariate homogeneity

(PERMDISP) to test for homoscedasticity in groups dispersions (Anderson, 2006)

because the PERMANOVA may be sensitive to non-homogeneous dispersions within

groups and thus mistake it as among-group variation (Anderson, 2001). A significant

PERMDISP ($p<0.05$) indicates different within-group dispersions and thus should be

used in combination with visual inspection of the ordination plots to interpret the

PERMANOVA results.

Using EAA after read depth normalization with DESeq2, we further visualized

bacterioplankton community temporal shifts with PRCs, asking if the extent of

community turnover varied across phylogenetic levels. Separate models were built for

ASVs grouped at various phylogenetic levels, from phylum to genus. For each PRC

model, we evaluated the proportion of variance (inertia) explained by the conditional

variable (time) and the constrained variable (treatments), as well as the proportion of

explained variance per axis (the eigenvalue of each RDA axis divided by the sum of all

eigenvalues). We used these values to decide which PRC model, if at the phylum, class,

order, family, genus or ASV level, best explained the variation in the data, and we tested

for the significance of the first PRC axis through a permutation test with the permute and

vegan packages in R (Oksanen et al., 2018; Simpson, 2019). Taxa weights representing

the affinity of the most responsive taxa with the treatment response curve are displayed

the right y-axis of each PRC diagram. Before performing each PRC we transformed the

community matrix using the Hellinger transformation (Legendre & Gallagher, 2001).

The abundance of the three genera with the highest PRC taxa weights were modeled

with GAMMs to explore how treatments impacted their (potentially non-linear)

abundances over time, and to provide further validation of the treatment effects detected

by PRCs. The GAMM response variable was the log-transformed (log (1+x), where x is

the variable) EAA of each of the three genera, after reads had been rarefied to 10,000

reads per sample without replacement. We opted for using rarefied data instead of

DESeq2 normalization which is intended for community analyses (Weiss et al., 2017)

and the GAMMs focused on specific taxa of interest. Modeled abundances were

visualized with the R package itsadug (Van Rij, Wieling, Baayen, & van Rijn, 2020).

Results

Bacterial cell density is weakly affected by glyphosate while microbial community carbon

substrate utilization is resistant to all stressors

Overall bacterial cell density showed a strong but non-linear increase over time across

all ponds (GAMM, effect of time: F=17.5, $p<0.001$, Table S4; Fig. 2). The time-

independent effect of nutrients on bacterial cell density was weak but positive (GAMM,

t=4.1, $p<0.001$), and, over time, glyphosate had a weak positive effect on bacterial

density (GAMM, factor-smooth interaction between time and glyphosate: F=6.6,

$p<0.001$) (Table S4). The interactive effect of nutrients and glyphosate was also weak,

and not significant after Bonferroni correction for multiple testing (GAMM, F=5.7,

uncorrected $p=0.018$, Table S4). Overall, these results indicate that, despite increasing

over time across ponds, bacterioplankton densities also slightly increased in response to

nutrient and glyphosate addition.

The number of carbon substrates used by the microbial community diminished slightly

over time (GAMM, F=6.0, p<0.001, Table S4). However, neither glyphosate,

imidacloprid, nutrients, nor their interactions had significant effects on carbon substrate

utilization as assayed by EcoPlates (Table S4). In addition, the PRC analysis did not

reveal any significant treatment effects on microbial utilization of any of the 31 unique

carbon substrates when considered separately (Fig. 3A; permutation test for the first

constrained eigenvalue, F=12.28 p=0.295) or when grouped into guilds (Fig. 3B;

F=34.46 p=0.355). To simplify visualization and facilitate comparison with treatments

selected for community taxonomic characterization, the PRCs in Fig. 3 included the

same ponds as those used for DNA analyses. PRCs including all ponds in the tested

gradient showed similar results (Fig. S2A and Fig. S2B). We conclude that, despite

slight changes in the number of substrates being used over time, none of the treatments

significantly affected microbial community-level carbon utilization profiles.

Bacterioplankton community structure responses

Glyphosate has a minor time-independent effect on community diversity and a

major effect on community composition over time

We calculated two metrics of bacterioplankton community alpha diversity in each

sample: taxon richness, estimated as the logarithm of the total number of observed

ASVs after rarefying (Fig. 4A), and the exponent of the Shannon index, which combines

information about ASV richness and evenness (Fig. 4B). No significant time-dependent

effect of any treatment was detected, although ponds with high glyphosate concentration

(15 mg/L) had a lower Shannon diversity when averaged across all timepoints (GAMM,

t=-3.51, p=0.001, Table S5), and the same was observed for ASV richness but with a

non-significant effect after multiple test correction (GAMM, t=-2.89, uncorrected

p=0.006, Table S5). Overall, the effect of glyphosate on bacterioplankton alpha diversity

was relatively weak and not influenced by time (Table S5).

We also tracked changes in bacterioplankton community composition, in two ways: with

weighted UniFrac distance and JSD, both calculated after normalizing read depth per

sample with DESeq2 (or with alternative normalizations described below). We display

these changes in community composition using PCoA, with a separate plot for each

timepoint of the experiment (Fig. 5). Glyphosate explained a significant proportion of the

variation in both metrics of community composition, with R^2 ranging from 0.29 to 0.58,

depending on the time following glyphosate application (PERMANOVA, p<0.007 for both

metrics at all tested timepoints after pesticide pulses, except for weighted UniFrac

distance at day 30, Tables S6 and S7). Nutrients and imidacloprid did not significantly

affect community composition, alone or in combination with other treatments (Tables S6

and S7). Although nutrients appear to have a slight effect on community composition on

day 15 (uncorrected p=0.027 for weighted UniFrac and JSD, Table S6 and Table S7)

and on day 30 (uncorrected p=0.055 for weighted UniFrac, Table S6, and uncorrected

p=0.013 for JSD, Table S7), the effect is not significant after Bonferroni correction, and

the explained variance is never as high as it is for glyphosate on the same day (R^2=0.12

for both weighted UniFrac and for JSD at both days, Table S6 and Table S7). We

conclude that glyphosate was the dominant driver of compositional changes as it

produced a significant and consistent effect on bacterioplankton communities,

independent of other stressors, on days 7, 15, 30 and 43 according to JSD, and on days

7, 15 and 43 according to weighted UniFrac distance.

Alternative read depth normalization methods (ASV relative abundance and rarefied

data; see Methods) produced qualitatively similar results, showing the predominant

effect of glyphosate on community composition (Fig. S3), with a slight delay in the effect

of the first glyphosate pulse compared to the DESeq2 normalization (Fig. 5). The effect

of glyphosate on bacterioplankton community composition is detected regardless of the

data normalization (Table S8), but is more apparent using DESeq2 (compare Fig. 5 to

Fig. S3). This might be because DESeq2 involves a log transformation which reduces

the weight of highly abundant community members (McKnight et al., 2019). If less

abundant taxa are more responsive to glyphosate, this could explain why this effect is

more apparent with DESeq2 normalization.

Bacterioplankton communities recover over time at broad phylogenetic scales

from the first glyphosate pulse

On day 30 (24 days after the first pesticide pulse and before the second pulse), the

bacterioplankton community composition in ponds that had been affected (on day 15) by

a high dose of glyphosate (15 mg/L) appeared to recover according to weighted UniFrac

(Fig. 5A), but not when using JSD applied to ASVs (Fig. 5B). Using weighted UniFrac,

the effect of glyphosate was visibly weaker on day 30 (Fig. 5A) and at the limit of

significance after Bonferroni correction (PERMANOVA, R^2=0.29, uncorrected p=0.007,

Table S6), but still significant using JSD (PERMANOVA, R^2=0.34, p=0.001, Table S7).

Viewed together, our series of ordinations show that detection of community recovery

depends upon whether phylogenetic information is taken into account. Recovery was

apparent when phylogenetic distance among ASVs was calculated (as measured by

UniFrac distance, on day 30, control and high-glyphosate communities approach each

other, Fig. 5A) but undetected at the ASV level, independent of phylogeny (as measured

by JSD, differences between control and high-glyphosate communities keep significant

on day 30, Fig. 5B). As such, the community appears to be resilient at a broad

phylogenetic level, but not at the finer ASV level, indicating that glyphosate-sensitive

ASVs are replaced with phylogenetically-close relatives.

To further assess how resilience varied at different phylogenetic scales, we used PRCs

to track community changes at the phylum and ASV levels (Fig. 6). Given that nutrient

inputs were not major drivers of community composition (Tables S6 and S7), we built

PRCs by combining ponds with the same pesticide treatment, irrespective of nutrient

load. This facilitated the visualization of pesticide effects, while capturing the same

effects as PRCs considering all experimental treatments separately (compare Fig. 6A

and Fig. S4). We further compared PRCs at different phylogenetic scales, from class to

genus level (Fig. S5). PRCs captured a significant amount of the variation in community

responses to pesticide treatments over time (phylum level: F=31.22, class: F=34.28,

order: F=26.19, family: F=21.30, genus: F=20.6, ASV: F=10.61, all p=0.001; Table S9),

with greater variation explained at broader taxonomic levels compared to finer levels.

The variance explained by the first PRC axis decreased from 47.7% at the phylum level

to 22.1% at the ASV level (Table S9). At the broadest taxonomic scale (phylum),

communities showed a clear response to high (15 mg/L) but not low (0.3 mg/L)

concentrations of glyphosate, followed by a recovery before the second pulse (Fig. 6A).

Notably, no recovery was observed at the ASV level (Fig. 6B), consistent with the

community composition analysis (Fig. 5). Imidacloprid had no detectable effect at any

concentration, whereas the highest concentration of glyphosate caused the greatest

effect on bacterioplankton communities. Similar response and recovery patterns were

also observed down to the genus level, with progressively weaker recovery at finer

taxonomic scales (Fig. S5). Community composition showed recovery 24 days after the

first pulse of glyphosate, but failed to recover after the second pulse (Fig. 5A and Fig.

6A). While this does not exclude the possibility of an eventual recovery, the duration of

our experiment (which ended nine days after the second pulse) was likely insufficient to

permit subsequent recovery. These results further support that high concentrations of

glyphosate led to long-lasting community shifts at the ASV or genus level, whereas

community resilience can be achieved at broader phylogenetic scales.

Dynamics of the taxa most responsive to treatments

The phylum Proteobacteria was the most positively affected by glyphosate (Fig 6A;

Table S10), with relative abundance over 60% in the high glyphosate treatment (15

mg/L) and ~50% or less in other treatments and controls (Table S11). Bacteroidetes,

Actinobacteria and Cyanobacteria were the most negatively affected phyla (Fig. 6A;

Table S10, Table S11). Of the ten ASVs with the highest absolute taxa weights, all

belonged to the phylum Proteobacteria (Fig. 6B, Table S12) and, except for sp283 and

sp2111, they were all positively affected by glyphosate. An ASV assigned to the genus

Agrobacterium was among the ASVs that responded most positively to high glyphosate

treatment (Fig. 6B; Table S12). The GAMM showed that ASVs assigned to the genus

Agrobacterium increased in EAA over time in ponds receiving high doses of glyphosate

(GAMM, factor-smooth interaction between time and high glyphosate treatment:

F=19.49, *p*<0.001, Table S13), or receiving both high glyphosate and imidacloprid

(GAMM, factor-smooth interaction between time and treatment with high concentrations

of both glyphosate and imidacloprid: F=20.66, *p*<0.001, Table S13). A linear time-

independent effect of glyphosate was also detected in experimental ponds treated with

the highest concentrations of both pesticides together (GAMM, t=7.50, *p*<0.001, Table

S13) or glyphosate alone (GAMM, t=6.25, *p*<0.001, Table S13). The modeled

Agrobacterium abundance (Fig. S6A) shows a similar 'response followed by recovery'

pattern over time as the overall community response at the phylum level (Fig. 6A),

suggesting that the positive effect of glyphosate on Proteobacteria may be driven by

Agrobacterium.

The other two most positively affected genera (*Flavobacterium* and *Azospirillum*, Fig.

S5D) increased in abundance in response to the combination of glyphosate at 15 mg/L

and imidacloprid at 60 μg/L (GAMM, factor-smooth interaction between time and

treatment with high concentrations of both glyphosate and imidacloprid on

Flavobacterium: F=17.35, *p*<0.001, and on *Azospirillum*: F=6.27 p=0.001, Table S13) or

glyphosate alone at 15 mg/L (GAMM, factor-smooth interaction between time and high

glyphosate treatment on *Flavobacterium*: F=3.63, *p*=0.031, not significant after

Bonferroni correction; and on *Azospirillum*: F=5.41, *p*=0.002, Table S13), but the effects

were not as strong as detected for *Agrobacterium* (Table S13). In contrast to the

recovery pattern observed in *Agrobacterium* exposed to both the independent and

combined highest concentrations of glyphosate (Fig. S6A), the modeled abundance of

Flavobacterium (Fig. S6B) and *Azospirillum* (Fig. S6C) followed distinct patterns in these

two treatments. *Flavobacterium* responded weakly to high doses of both pesticides,

mainly after the second pulse, whereas *Azospirillum* recovered partially after responding

to the first pulse, but only in ponds treated with the highest concentrations of both

pesticides. Despite the overall strong effect of glyphosate on Proteobacteria, these

results highlight how different bacterioplankton taxa (including *Agrobacterium* and

Azospirillum – both Alphaproteobacteria) can show subtly different responses and

recovery patterns to pesticides.

648
Discussion

Context and summary of the experiment

The herbicide glyphosate has been shown to affect aquatic microbial community

structure in a variety of natural environments and experimental setups (Berman et al.,

2020; Lu et al., 2020; Muturi et al., 2017; Stachowski-Haberkorn et al., 2008). Likewise,

the insecticide imidacloprid may disrupt aquatic food webs (Yamamuro et al., 2019), with

potential, yet poorly explored consequences for bacterioplankton. The interactive effects

of these pesticides on bacterioplankton – and how they might vary depending on

fertilizer use and lake trophic status – are relevant because such agrochemical mixtures

are common in agriculturally impacted watersheds. Here, we tested how individual and

combined gradients of glyphosate and imidacloprid affected bacterioplankton

communities in aquatic mesocosms receiving different nutrient inputs. Although they are

incomplete representations of natural ecosystems, mesocosm experiments allow us to

manipulate and replicate the exposure of complex lake bacterial communities to

agricultural chemical pollutants commonly found in freshwaters (Alexander, Luiker,

Finley, & Culp, 2016). The current experiment is limited to the response of

bacterioplankton communities derived from a pristine lake. Future studies focusing on

biofilms and sediments could complement our results, as many contaminants

accumulate in lake sediments and may affect the biofilm community structure

(Fernandes et al., 2019; Khadra, Planas, Girard, & Amyot, 2018; Romero et al., 2020).

Glyphosate as a driver of community structure

Our data support the prediction that glyphosate would affect bacterioplankton

community structure, which occurred at the highest tested concentration (15 mg/L).

Contrary to expectation, no evident interaction between glyphosate and imidacloprid or

nutrient load was detected in determining either bacterial density or community

structure. High doses of glyphosate resulted in a weak time-independent reduction of

bacterioplankton alpha diversity, and a more pronounced change in community

composition over time. As hypothesized, glyphosate and nutrient treatments slightly

increased bacterial density, suggesting a mild fertilizing effect of glyphosate consistent

with it being a potential phosphorus source (Hove-Jensen et al., 2014; Lu et al., 2020).

Most bacterioplankton from a pristine source environment (Lake Hertel) are thus able to

cope with concentrations of imidacloprid as high as 60 µg/L and of glyphosate as high

as 0.3 mg/L, but they may be sensitive to glyphosate concentrations exceeding 15 mg/L.

The regulatory criteria intended for eukaryotes (below 60 µg/L for imidacloprid; Table

S1) were sufficient to preserve bacterioplankton diversity in the experimental conditions

at LEAP. On the other hand, the threshold of 15 mg/L for glyphosate deserves further

attention from regulatory agencies, as this concentration impacted bacterioplankton

composition, which is known to affect lake health and freshwater quality (Kraemer et al.,

2020).

Although the highest targeted imidacloprid concentration was not always achieved in all

ponds (Fig. S1), this cannot entirely explain its lack of detectable effect on

bacterioplankton. Community composition of ponds receiving measured concentrations

of imidacloprid as high as 15 µg/L or more did not deviate from controls, confirming a

true lack of effect at least up to that concentration. Alternatively, the absence of a

detectable response might be due in part to rapid degradation of imidacloprid in water,

which fell below the limit of detection between pulses (Fig. S1). The absence of a

bacterioplankton response is also consistent with the weak or undetectable response of

zooplankton biomass to imidacloprid pulses in the same experiment (Hébert et al.,

2021). The invertebrate community in the experimental ponds was mainly composed of

the zooplanktonic groups Cladocera, Copepoda and Rotifera, and only copepods

declined over time after pulse 2, with no resulting effect in total zooplankton biomass

(Hébert et al., 2021). Overall, these results indicate that the concentrations of

imidacloprid applied in this experiment were not sufficient to strongly alter either

zooplankton or bacterioplankton biomass or community structure.

Our results suggest that two properties of ecological stability – resistance and resilience

– are at play in lake bacterioplankton: functions related to microbial carbon substrate

use are resistant to imidacloprid, glyphosate and their interactions in different nutrient

backgrounds, while bacterioplankton community composition is resilient following

disturbance caused by a glyphosate pulse at 15 mg/L. The recovery of bacterioplankton

community composition was only evident when grouping ASVs at higher (more

inclusive) taxonomic or phylogenetic levels. Glyphosate thus drove a turnover of

bacterioplankton ASVs which, even after the recovery, are different from the ASVs

initially found in the undisturbed community.

Proteobacteria are major responders to glyphosate

Glyphosate treatments had a strong positive effect on the phylum Proteobacteria,

previously found to be favoured by high concentrations of glyphosate in rhizosphere-

(Newman et al., 2016) and phytoplankton-associated communities (Wang, Lin, Li, Lin, &

Lin, 2017). Multiple species of Proteobacteria can use glyphosate as a source of

phosphorus by breaking its C-P bond (Hove-Jensen et al., 2014). We identified

Agrobacterium, a genus of *Rhizobiaceae* containing species known to degrade

glyphosate (Hove-Jensen et al., 2014), as being highly favored in the glyphosate

treatment at 15 mg/L. The abundance of ASVs assigned to this genus peaked after each

pulse and decreased before the second pulse, coinciding with the community recovery

observed 24 days after the first perturbation. The ability to degrade glyphosate may be

widespread in the family *Rhizobiaceae* (Liu, McLean, Sookdeo, & Cannon, 1991), and

Agrobacterium have also been found to encode glyphosate-resistant EPSPS genes

(Funke et al., 2006). In fact, this genus was used to create glyphosate-resistant crops,

i.e. the so-called 'Roundup-ready technology' (Funke et al., 2006). While glyphosate

may be a stressor for the microbial community at large (e.g.*,* phytoplankton (Fugère et

al., 2020)), it may be a resource for some members such as *Agrobacterium*, who could

potentially detoxify the environment and thus facilitate community recovery after a pulse

perturbation. Further genomic and metagenomic analyses of our experimental samples

could reveal whether these ecological dynamics are underlain by evolutionary

adaptation, and whether community resistance and resilience can be explained by the

initial presence of resistant bacteria in the community, or to *de novo* mutations or gene

transfer events.

Glyphosate could have driven changes in the bacterial community via direct

mechanisms (e.g. by affecting species with a sensitive EPSPS, its target enzyme) or

indirect mechanisms (e.g. effects on other trophic levels that cascaded down to bacteria

via predation or other interactions). In a previous study of the same experiment

described here that focused on the responses of eukaryotic phytoplankton, we found

that glyphosate treatment reduced the diversity of phytoplankton, but did not significantly

change phytoplankton community composition (Fugère et al., 2020). Although a reduced

phytoplankton diversity could indirectly affect bacterioplankton community composition,

a direct effect of glyphosate on bacteria seems more plausible as the taxa favored by

the treatment (mainly Proteobacteria) have been previously shown to be directly

affected in a similar way (Janßen et al., 2019; Wang et al., 2017). Indeed, bacterial

degradation of glyphosate likely released bioavailable phosphorus, stimulating

phytoplankton growth (Fugère et al., 2020). Further studies will be needed to

disentangle how the effects of pesticides cascade through food webs, and how trophic

structure influences their effects.

Functional redundancy in carbon utilization potential

Despite the marked changes in taxonomic composition driven by glyphosate, microbial

communities did not change their carbon substrate use throughout the experiment,

providing evidence for functional redundancy in carbon utilization potential. This was an

expected result, as broad-scale ecosystem functions such as respiration and dissolved

organic carbon consumption are weakly coupled with species composition (Girvan,

Campbell, Killham, Prosser, & Glover, 2005; Langenheder, Lindström, & Tranvik, 2006;

Peter et al., 2011), allowing these functions to remain unaffected by fluctuations in

microbial community composition (Louca et al., 2018). While less diverse communities

(in terms of species richness) may lack functional redundancy, more diverse

communities are expected to encode more redundant functions (Konopka, 2009). We

can thus surmise that the freshwater bacterioplankton communities studied here were

sufficiently diverse to be functionally redundant for carbon utilization in the face of

disturbance. The weak and time-independent effect of high concentrations of glyphosate

on alpha diversity was insufficient to alter community carbon substrate use. However,

our experiment was conducted with communities originating from a pristine lake in a

nature reserve, and this result might not be generalized to freshwaters historically

impacted by other forms of anthropogenic stress. For example, land use intensity is

negatively correlated with bacterioplankton richness in lakes across Eastern Canada

(Kraemer et al., 2020). It remains to be seen whether such impacted lakes are less

functionally redundant, and thus possibly more susceptible to impaired ecosystem

functioning. Lastly, although bacterioplankton respiration accounts for a large fraction of

organic carbon processing in freshwaters (Berggren, Lapierre, & del Giorgio, 2012), the

carbon substrate use we measured could also be due in part to fungal activity which

could be compensating or masking changes in bacterioplankton activity. There was no

macroscopically observable fungal growth in the plates, yet microscopic fungi likely

contributed a fraction of the inoculum used to initiate the plates.

The phylogenetic depth of glyphosate resistance: Methodological considerations

The inference of bacterioplankton ASVs in this study allowed a relatively fine-scale

taxonomic resolution of community changes in response to a pulse perturbation of

glyphosate. Notably, the recovery of bacterioplankton composition was detectable at

broader taxonomic units (e.g. phylum in particular) but not at the ASV level. This implies

that the taxonomic resolution of traits under selection during recovery from a glyphosate

pulse is relatively coarse (Martiny, Jones, Lennon, & Martiny, 2015). This result could

also be explained if ASVs are too fine-scale as a measure of diversity, and mostly reflect

sequencing or base calling errors rather than true biological diversity. We deem this

unlikely, first because the ASV inference algorithm includes a model-based approach to

correct for amplicon sequencing errors (Callahan et al., 2016), and second because

ASV detection methods are usually more accurate than OTU-clustering methods based

on sequence similarity thresholds of usually 97% (Caruso, Song, Asquith, & Karstens,

2019). For example, we only found 7 to 10 false-positive ASVs (Methods), but dozens to

hundreds of false positive are detected by even state-of-the-art (distribution-based)

sequence clustering-based methods to identify operational taxonomic units, when

applied to the same or similar mock communities as used here (Tromas et al., 2017).

Although we cannot exclude the impact of possible false ASVs on our results, we expect

them to be relatively minimal and evenly distributed across all timepoints (Callahan,

McMurdie, & Holmes, 2017). In other words, there is no reason to believe that

sequencing errors should be non-randomly distributed over time or across experimental

treatments. Moreover, PRC analyses show a steady decline from the phylum level to the

genus level in both the response to, and recovery from, high concentrations of

glyphosate. Therefore, even without considering the ASV level, there is still a discernible

pattern of greater community resilience at broader taxonomic scales. This suggests that

the traits (and underlying genes) required for survival or growth in the presence of

glyphosate are relatively deeply conserved. Higher-resolution genomic or metagenomic

analyses could be used to confirm this result, and pinpoint the genes involved in

resistance.

Ecosystem resistance, resilience and stability

Our study provides evidence of ecosystem stability in terms of carbon substrate use

maintained by microbial communities when faced by a perturbation by two of the most

commonly used pesticides in the world, separately or in combination. We also showed

resistance to a wide gradient of imidacloprid contamination, and resilience to high doses

of glyphosate in bacterioplankton communities that have no known history of contact

with the herbicide. Finally, whether a stressed community is considered resilient

depends on the phylogenetic depth of the traits required to deal with the stress (Martiny

et al., 2015). Our results provide an example of how resilience to stressors can be a

feature of deeper phylogenetic groups, but not finer-scale groupings (ASVs), which

could be involved in adaptation to other stressors or niches.

Acknowledgments

We are grateful to D. Maneli, C. Normandin, A. Arkilanian and T. Jagadeesh for their

assistance in the field, to J. Marleau, C. Girard, O.M. Pérez-Carrascal and N. Tromas for

their assistance during laboratory analyses, to K. Velghe for nutrient analyses and to

M.A.P. Castro for developing the LC–MS method for pesticides quantification. We thank

the three anonymous reviewers for comments on a previous version of this manuscript.

Funding information

This study was supported by a Canada Research Chair and NSERC Discovery Grant to

B.J.S. N.B.C. was funded by FRQNT and NSERC-CREATE/GRIL fellowships. V.F. was

supported by an NSERC postdoctoral fellowship. M-P.H. was funded by NSERC and

NSERC-CREATE/GRIL. C.C.Y.X. was funded by a Vanier Canada Graduate

Scholarship. R.D.H.B. was supported by a Canada Research Chair. LEAP was built and

operated with funds from a CFI Leaders Opportunity Fund, NSERC Discovery Grant and

the Liber Ero Chair to A.G.

842   References

Alexander, A. C., Luiker, E., Finley, M., & Culp, J. M. (2016). Mesocosm and Field
Toxicity Testing in the Marine Context. In *Marine Ecotoxicology* (pp. 239–256).
Elsevier Inc. https://doi.org/10.1016/b978-0-12-803371-5.00008-4

Alexander, A. C., Luis, A. T., Culp, J. M., Baird, D. J., & Cessna, A. J. (2013). Can
nutrients mask community responses to insecticide mixtures? *Ecotoxicology*, *22*(7),
1085–1100. https://doi.org/10.1007/s10646-013-1096-3

Allison, S. D., & Martiny, J. B. H. (2008). Resistance, resilience, and redundancy in
microbial communities. *Proceedings of the National Academy of Sciences of the*
USA, *105*, 11512–11519. https://doi.org/10.1073/pnas.0801925105

Altenburger, R., Backhaus, T., Boedeker, W., Faust, M., & Scholze, M. (2013).
Simplifying complexity: Mixture toxicity assessment in the last 20 years.
Environmental Toxicology and Chemistry, *32*(8), 1685–1687.
https://doi.org/10.1002/etc.2294

Anderson, M. J. (2001). A new method for non-parametric multivariate analysis of
variance, *26*, 32–46.

Anderson, M. J. (2006). Distance-based tests for homogeneity of multivariate
dispersions. *Biometrics*, *62*(1), 245–253. https://doi.org/10.1111/j.1541-
0420.2005.00440.x

Auber, A., Travers-Trolet, M., Villanueva, M.-C., & Ernande, B. (2017). A new application
of principal response curves for summarizing abrupt and cyclic shifts of
communities over space. *Ecosphere*, *8*(12), e02023.
https://doi.org/10.1002/ecs2.2023

Baker, L. F., Mudge, J. F., Thompson, D. G., Houlahan, J. E., & Kidd, K. A. (2016). The
combined influence of two agricultural contaminants on natural communities of
phytoplankton and zooplankton. *Ecotoxicology*, *25*(5), 1021–1032.
https://doi.org/10.1007/s10646-016-1659-1

Beauséjour, R., Handa, I. T., Lechowicz, M. J., Gilbert, B., & Vellend, M. (2015).
Historical anthropogenic disturbances influence patterns of non-native earthworm
and plant invasions in a temperate primary forest. *Biological Invasions*, *17*(4),
1267–1281. https://doi.org/10.1007/s10530-014-0794-y

Benbrook, C. M. (2016). Trends in glyphosate herbicide use in the United States and
globally. *Environmental Sciences Europe*, *28*(1), 1–15.
https://doi.org/10.1186/s12302-016-0070-0

Berggren, M., Lapierre, J. F., & del Giorgio, P. A. (2012). Magnitude and regulation of
bacterioplankton respiratory quotient across freshwater environmental gradients.
The ISME Journal, *6*(5), 984–993. https://doi.org/10.1038/ismej.2011.157

Berman, M. C., Llames, M. E., Minotti, P., Fermani, P., Quiroga, M. V., Ferraro, M. A., …
Zagarese, H. E. (2020). Field evidence supports former experimental claims on the
stimulatory effect of glyphosate on picocyanobacteria communities. *Science of the*
Total Environment, *701*, 134601. https://doi.org/10.1016/j.scitotenv.2019.134601
Birk, S., Chapman, D., Carvalho, L., Spears, B. M., Andersen, H. E., Argillier, C., …
Hering, D. (2020). Impacts of multiple stressors on freshwater biota across spatial
scales and ecosystems. *Nature Ecology & Evolution*, *4*(8), 1060–1068.
https://doi.org/10.1038/s41559-020-1216-4
Bloomfield, J. P., Williams, R. J., Gooddy, D. C., Cape, J. N., & Guha, P. (2006).
Impacts of climate change on the fate and behaviour of pesticides in surface and
groundwater-a UK perspective. *Science of the Total Environment*, *369*(1–3), 163–
177. https://doi.org/10.1016/j.scitotenv.2006.05.019
Callahan, B., McMurdie, P., Rosen, M., Han, A. W., Johnson, A. J. A., & Holmes, S. P.
(2016). DADA2: High-resolution sample inference from Illumina amplicon data.
Nature Methods, *13*(7), 581–583. https://doi.org/10.1038/nmeth.3869
Callahan, Ben J., Sankaran, K., Fukuyama, J. A., McMurdie, P. J., & Holmes, S. P.
(2016). Bioconductor workflow for microbiome data analysis: From raw reads to
community analyses. *F1000Research*, *5*(1492), 1–49.
https://doi.org/10.12688/f1000research.8986.2
Callahan, Benjamin J, McMurdie, P. J., & Holmes, S. P. (2017). Exact sequence
variants should replace operational taxonomic units in marker-gene data analysis.
The ISME Journal, 1–5. https://doi.org/10.1038/ismej.2017.119
Carpenter, S. R., Caraco, N. F., Correll, D. L., Howarth, R. W., Sharpley, A. N., & Smith,
V. H. (1998). Nonpoint pollution of surface waters with phosphorus and nitrogen.
Ecological Application, *8*(January 1998), 559–568. https://doi.org/10.1890/1051-
0761(1998)008[0559:NPOSWW]2.0.CO;2
Carpenter, S. R., Stanley, E. H., & Vander Zanden, M. J. (2011). State of the world's
freshwater ecosystems: Physical, chemical, and biological changes. *Annual Review*
of Environment and Resources, *36*, 75–99. https://doi.org/10.1146/annurev-environ-
021810-094524
Caruso, V., Song, X., Asquith, M., & Karstens, L. (2019). Performance of Microbiome
Sequence Inference Methods in Environments with Varying Biomass. *MSystems*,
4(1), 1–19. https://doi.org/10.1128/msystems.00163-18
CCME. (2007). *Canadian Water Quality Guidelines for Protection of Aquatic Life:*
Imidacloprid. Scientific Supporting Document. Winnipeg, Manitoba.
CCME. (2012). *Canadian Water Quality Guidelines for the Protection of Aquatic Life:*
Glyphosate. Scientific Supporting Document. Winnipeg, Manitoba.
Cedergreen, N., & Rasmussen, J. J. (2017). Low dose effects of pesticides in the
aquatic environment. In *ACS Symposium Series* (Vol. 1249, pp. 167–187).
https://doi.org/10.1021/bk-2017-1249.ch012
Chará-Serna, A. M., Epele, L. B., Morrissey, C. A., & Richardson, J. S. (2019). Nutrients
and sediment modify the impacts of a neonicotinoid insecticide on freshwater
community structure and ecosystem functioning. *Science of the Total Environment*,
692, 1291–1303. https://doi.org/10.1016/j.scitotenv.2019.06.301
DeLorenzo, M. E., Scott, G. I., & Ross, P. E. (2001). Toxicity of pesticides to aquatic

microorganisms: A review. *Environmental Toxicology and Chemistry*, *20*(1), 84–98. https://doi.org/10.1002/etc.5620200108

DeSantis, T. Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E. L., Keller, K., … Andersen, G. L. (2006). Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Applied and Environmental Microbiology*, *72*(7), 5069–5072. https://doi.org/10.1128/AEM.03006-05

EFSA. (2014). *Conclusion on the peer review of the pesticide risk assessment for aquatic organisms for the active substance imidacloprid. EFSA Journal* (Vol. 12). https://doi.org/10.2903/j.efsa.2013.3066

EFSA. (2016). *Conclusion on the peer review of the pesticide risk assessment of the active substance glyphosate. EFSA Journal* (Vol. 13). https://doi.org/10.2903/j.efsa.2015.4302

EPA. (2019). Aquatic Life Benchmarks and Ecological Risk Assessments for Registered Pesticides. Retrieved from https://www.epa.gov/pesticide-science-and-assessing-pesticide-risks/aquatic-life-benchmarks-and-ecological-risk#ref_4

Falkowski, P. G., Fenchel, T., & Delong, E. F. (2008). The microbial engines that drive earth's biogeochemical cycles. *Science*, *320*(5879), 1034–1039. https://doi.org/10.1126/science.1153213

Fernandes, G., Carolina, V., Camotti, M., Gerónimo, E. De, Labanowski, J., Damian, O., … Rheinheimer, D. (2019). Indiscriminate use of glyphosate impregnates river epilithic biofilms in southern Brazil. *Science of the Total Environment*, *651*, 1377–1387. https://doi.org/10.1016/j.scitotenv.2018.09.292

Flood, S. L., & Burkholder, J. A. M. (2018). Imbalanced nutrient regimes increase Prymnesium parvum resilience to herbicide exposure. *Harmful Algae*, *75*, 57–74. https://doi.org/10.1016/j.hal.2018.04.006

Fugère, V., Hébert, M., Barbosa da Costa, N., Xu, C. C. Y., Barrett, R. D. H., Beisner, B. E., … Gonzalez, A. (2020). Community rescue in experimental phytoplankton communities facing severe herbicide pollution. *Nature Ecology & Evolution*, *4*, 578–588. https://doi.org/10.1038/s41559-020-1134-5

Funke, T., Han, H., Healy-Fried, M. L., Fischer, M., & Schönbrunn, E. (2006). Molecular basis for the herbicide resistance of Roundup Ready crops. *Proceedings of the National Academy of Sciences of the USA*, *103*(35), 13010–13015. https://doi.org/10.1073/pnas.0603638103

Garland, J. L., Mills, A. L., & Young, J. S. (2001). Relative effectiveness of kinetic analysis vs single point readings for classifying environmental samples based on community-level physiological profiles (CLPP). *Soil Biology and Biochemistry*, *33*(7–8), 1059–1066. https://doi.org/10.1016/S0038-0717(01)00011-6

Gasol, J. M., & Del Giorgio, P. A. (2000). Using flow cytometry for counting natural planktonic bacteria and understanding the structure of planktonic bacterial communities. *Scientia Marina*, *64*(2), 197–224. https://doi.org/10.3989/scimar.2000.64n2197

Geyer, R. L., Smith, G. R., & Rettig, J. E. (2016). Effects of Roundup formulations, nutrient addition, and Western mosquitofish (Gambusia affinis) on aquatic communities. *Environmental Science and Pollution Research*, *23*(12), 11729–11739. https://doi.org/10.1007/s11356-016-6381-2

Girvan, M., Campbell, C., Killham, K., Prosser, J., & Glover, L. (2005). Bacterial diversity
promotes community stability and functional resilience after perturbation.
Environmental Microbiology, *7*(3), 301–313. https://doi.org/10.1111/j.1462-
2920.2004.00695.x
Healy-Fried, M. L., Funke, T., Priestman, M. A., Han, H., & Scho, E. (2007). Structural
Basis of Glyphosate Tolerance Resulting from Mutations of Pro 101 in Escherichia
coli. *The Journal of Biological Chemistry*, *282*(45), 32949–32955.
https://doi.org/10.1074/jbc.M705624200
Hébert, M.-P., Fugère, V., Beisner, B. E., Barbosa da Costa, N., Barrett, R. D. H., Bell,
G., … Fussmann, G. F. (2021). Widespread agrochemicals differentially affect
zooplankton biomass and community structure. *Ecological Applications*.
https://doi.org/10.1002/eap.2423
Hébert, M.-P., Fugère, V., & Gonzalez, A. (2018). The overlooked impact of rising
glyphosate use on phosphorus loading in agricultural watersheds. *Frontiers in*
Ecology and the Environment, *3*, 1–9. https://doi.org/10.1002/fee.1985
Hénault-Ethier, L., Lucotte, M., Moingt, M., Paquet, S., Maccario, S., Smedbol, É., …
Labrecque, M. (2017). Herbaceous or Salix miyabeana 'SX64' narrow buffer strips
as a means to minimize glyphosate and aminomethylphosphonic acid leaching from
row crop fields. *Science of the Total Environment*, *598*, 1177–1186.
https://doi.org/10.1016/j.scitotenv.2017.04.104
Hove-Jensen, B., Zechel, D. L., & Jochimsen, B. (2014). Utilization of glyphosate as
phosphate source: biochemistry and genetics of bacterial carbon-phosphorus lyase.
Microbiology and Molecular Biology Reviews : MMBR, *78*(1), 176–197.
https://doi.org/10.1128/MMBR.00040-13
Janßen, R., Skeff, W., Werner, J., Wirth, M. A., Kreikemeyer, B., Schulz-Bull, D., &
Labrenz, M. (2019). A Glyphosate Pulse to Brackish Long-Term Microcosms Has a
Greater Impact on the Microbial Diversity and Abundance of Planktonic Than of
Biofilm Assemblages. *Frontiers in Marine Science*, *6*(December), 1–17.
https://doi.org/10.3389/fmars.2019.00758
Jeppesen, E., Kronvang, B., Meerhoff, M., Søndergaard, M., Hansen, K. M., Andersen,
H. E., … Olesen, J. E. (2009). Climate Change Effects on Runoff, Catchment
Phosphorus Loading and Lake Ecological State, and Potential Adaptations. *Journal*
of Environment Quality, *38*(5), 1930. https://doi.org/10.2134/jeq2008.0113
Jeschke, P., & Nauen, R. (2008). Neonicotinoids – from zero to hero in insecticide
chemistry. *Pest Management Science*, *64*, 1100–1106.
https://doi.org/10.1002/ps.1631
Jost, L. (2006). Entropy and diversity. *Oikos*, *113*(2), 363–375.
Keatley, B. E., Bennett, E. M., Macdonald, G. K., Taranu, Z. E., & Gregory-Eaves, I.
(2011). Land-Use Legacies Are Important Determinants of Lake Eutrophication in
the Anthropocene. *PLOS One*, *6*(1), e15913.
https://doi.org/10.1371/journal.pone.0015913
Khadra, M., Planas, D., Girard, C., & Amyot, M. (2018). Age matters: Submersion period
shapes community composition of lake biofilms under glyphosate stress. *Facets*,
3(1), 934–951. https://doi.org/10.1139/facets-2018-0019
Konopka, A. (2009). What is microbial community ecology? *The ISME Journal*, *3*(11),

1223–1230. https://doi.org/10.1038/ismej.2009.88

Kraemer, S. A., Barbosa da Costa, N., Shapiro, B. J., Fradette, M., Huot, Y., & Walsh, D.
A. (2020). A large-scale assessment of lakes reveals a pervasive signal of land use
on bacterial communities. *The ISME Journal.* https://doi.org/10.1038/s41396-020-
0733-0

Langenheder, S., Lindström, E. S., & Tranvik, L. J. (2006). Structure and function of
bacterial communities emerging from different sources under identical conditions.
Applied and Environmental Microbiology, *72*(1), 212–220.
https://doi.org/10.1128/AEM.72.1.212-220.2006

Legendre, P., & Gallagher, E. D. (2001). Ecologically meaningful transformations for
ordination of species data. *Oecologia*, *129*, 271–280.
https://doi.org/10.1007/s004420100716

Liu, C. M., McLean, P. A., Sookdeo, C. C., & Cannon, F. C. (1991). Degradation of the
herbicide glyphosate by members of the family Rhizobiaceae. *Applied and*
Environmental Microbiology, *57*(6), 1799–1804.

Louca, S., Polz, M. F., Mazel, F., Albright, M. B. N., Huber, J. A., Connor, M. I. O., …
Parfrey, L. W. (2018). Function and functional redundancy in microbial systems.
Nature Ecology & Evolution, *2*, 936–943. https://doi.org/10.1038/s41559-018-0519-
1

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(550).
https://doi.org/10.1186/s13059-014-0550-8

Lozano, R. B., & Pratt, J. R. (1994). Interaction of toxicants and communities: The role
of nutrients. *Environmental Toxicology and Chemistry*, *13*(3), 361–368.
https://doi.org/10.1002/etc.5620130302

Lozupone, C., & Knight, R. (2005). UniFrac☐: a New Phylogenetic Method for
Comparing Microbial Communities. *Applied and Environmental Microbiology*,
71(12), 8228–8235. https://doi.org/10.1128/AEM.71.12.8228

Lu, T., Xu, N., Zhang, Q., Zhang, Z., Debognies, A., Zhou, Z., … Qian, H. (2020).
Understanding the influence of glyphosate on the structure and function of
freshwater microbial community in a microcosm. *Environmental Pollution*, *260*,
114012. https://doi.org/10.1016/j.envpol.2020.114012

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput
sequencing reads. *EMBnet.Journal*, *17*, 10–12.

Martiny, J. B. H., Jones, S. E., Lennon, J. T., & Martiny, A. C. (2015). Microbiomes in
light of traits: a phylogenetic perspective. *Science*, *350*(6261), aac9323.
https://doi.org/10.1126/science.aac9323

McComb, A. J. (2002). *Limnological Analyses. Lakes and Reservoirs: Research and*
Management (Vol. 7). https://doi.org/10.1046/j.1440-1770.2002.01722.x

McKnight, D. T., Huerlimann, R., Bower, D. S., Schwarzkopf, L., Alford, R. A., & Zenger,
K. R. (2019). Methods for normalizing microbiome data: An ecological perspective.
Methods in Ecology and Evolution, *10*(3), 389–400. https://doi.org/10.1111/2041-
210X.13115

McMurdie, P. J., & Holmes, S. (2013). phyloseq☐: An R Package for Reproducible
Interactive Analysis and Graphics of Microbiome Census Data. *PLOS One*, *8*(4),

e61217. https://doi.org/10.1371/journal.pone.0061217

McMurdie, P. J., & Holmes, S. (2014). Waste not, want not◻: why rarefying Microbiome
data is inadmissible. *PLOS Computational Biology, 10*(4), e1003531.
https://doi.org/10.1371/journal.pcbi.1003531

Morrissey, C. A., Mineau, P., Devries, J. H., Sanchez-Bayo, F., Liess, M., Cavallaro, M.
C., & Liber, K. (2015). Neonicotinoid contamination of global surface waters and
associated risk to aquatic invertebrates: A review. *Environment International, 74*,
291–303. https://doi.org/10.1016/j.envint.2014.10.024

Muturi, E. J., Donthu, R. K., Fields, C. J., Moise, I. K., & Kim, C. (2017). Effect of
pesticides on microbial communities in container aquatic habitats. *Scientific
Reports*, (November 2016), 1–10. https://doi.org/10.1038/srep44565

Newman, M. M., Hoilett, N., Lorenz, N., Dick, R. P., Liles, M. R., Ramsier, C., &
Kloepper, J. W. (2016). Glyphosate effects on soil rhizosphere-associated bacterial
communities. *Science of the Total Environment, 543*, 155–160.
https://doi.org/10.1016/j.scitotenv.2015.11.008

Newton, R. J., Jones, S. E., Eiler, A., Mcmahon, K. D., & Bertilsson, S. (2011). A Guide
to the Natural History of Freshwater Lake Bacteria. *Microbiology and Molecular
Biology Reviews, 75*(1), 14–49. https://doi.org/10.1128/MMBR.00028-10

Oksanen, A. J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., Hara, R. B. O.,
… Wagner, H. (2018). Vegan: community ecology package. R package version 2.4-
6. https://doi.org/ISBN 0-387-95457-0

Paerl, H. W., Otten, T. G., & Kudela, R. (2018). Mitigating the Expansion of Harmful
Algal Blooms Across the Freshwater-to-Marine Continuum. *Environmental Science
and Technology, 52*(10), 5519–5529. https://doi.org/10.1021/acs.est.7b05950

Patton, C. J., & Kryskalla, J. R. (2003). *Methods of Analysis by the U.S. Geological
Survey National Water Quality Laboratory—Evaluation of Alkaline Persulfate
Digestion as an Alternative to Kjeldahl Digestion for Determination of Total and
Dissolved Nitrogen and Phosphorus in Water. USGS.*

Peter, H., Beier, S., Bertilsson, S., Lindström, E. S., Langenheder, S., & Tranvik, L. J.
(2011). Function-specific response to depletion of microbial diversity. *The ISME
Journal, 5*(2), 351–361. https://doi.org/10.1038/ismej.2010.119

Piggott, J. J., Niyogi, D. K., Townsend, C. R., & Matthaei, C. D. (2015). Multiple
stressors and stream ecosystem functioning: Climate warming and agricultural
stressors interact to affect processing of organic matter. *Journal of Applied Ecology,
52*(5), 1126–1134. https://doi.org/10.1111/1365-2664.12480

Pollegioni, L., Schonbrunn, E., & Siehl, D. (2011). Molecular basis of glyphosate
resistance: Different approaches through protein engineering. *The FEBS Journal,
278*(16), 2753–2766. https://doi.org/10.1111/j.1742-4658.2011.08214.x

Preheim, S. P., Perrotta, A. R., Martin-platero, A. M., Gupta, A., & Alm, E. J. (2013).
Distribution-Based Clustering◻: Using Ecology To Refine the Operational
Taxonomic Unit. *Applied and Environmental Microbiology, 79*(21), 6593–6603.
https://doi.org/10.1128/AEM.00342-13

R Core Team. (2008). R: A Language and Environment for Statistical Computing. In *R
Foundation for Statistical Computing* (Vol. 1, pp. 1–2630). R Foundation for
Statistical Computing. https://doi.org/10.1007/978-3-540-74686-7

Rainio, M. J., Ruuskanen, S., Helander, M., Saikkonen, K., Saloniemi, I., & Puigbò, P. (2021). Adaptation of bacteria to glyphosate: a microevolutionary perspective of the enzyme 5-enolpyruvylshikimate-3-phosphate synthase. *Environmental Microbiology Reports*, *00*. https://doi.org/10.1111/1758-2229.12931

Relyea, R. A. (2009). A cocktail of contaminants: How mixtures of pesticides at low concentrations affect aquatic communities. *Oecologia*, *159*(2), 363–376. https://doi.org/10.1007/s00442-008-1213-9

Roberts, T., & Hutson, D. (1999). *Metabolic Pathways of Agrochemicals. Part 2: Insecticides and Fungicides*. (T. R. Roberts, D. H. Hutson, P. W. Lee, P. H. Nicholls, & J. R. Plimmer, Eds.), *The Royal Society of Chemistry*. Cambridge: The Royal Society of Chemistry. https://doi.org/10.1017/CBO9781107415324.004

Rohwer, R. R., Hamilton, J. J., Newton, R. J., & McMahon, K. D. (2018). TaxAss: Leveraging a Custom Freshwater Database Achieves Fine-Scale Taxonomic Resolution. *MSphere*, *3*(5), 1–14. https://doi.org/10.1128/msphere.00327-18

Romero, F., Acuña, V., & Sabater, S. (2020). Multiple stressors determine community structure and estimated function of river biofilm bacteria. *Applied and Environmental Microbiology*, *86*(12), 1–13. https://doi.org/10.1128/AEM.00291-20

Ruiz-González, C., Archambault, E., Laforest-Lapointe, I., Del Giorgio, P. A., Kembel, S. W., Messier, C., … Beisner, B. E. (2018). Soils associated to different tree communities do not elicit predictable responses in lake bacterial community structure and function. *FEMS Microbiology Ecology*, *94*(8), 1–15. https://doi.org/10.1093/femsec/fiy115

Ruiz-González, C., Niño-García, J. P., Lapierre, J.-F., & del Giorgio, P. A. (2015). The quality of organic matter shapes the functional biogeography of bacterioplankton across boreal freshwater ecosystems. *Global Ecology and Biogeography*, *24*, 1487–1498. https://doi.org/10.1111/geb.12356

Shade, A., Peter, H., Allison, S. D., Baho, D. L., Berga, M., Bürgmann, H., … Handelsman, J. (2012). Fundamentals of microbial community resistance and resilience. *Frontiers in Microbiology*, *3*(DEC), 1–19. https://doi.org/10.3389/fmicb.2012.00417

Simon-Delso, N., Amaral-Rogers, V., Belzunces, L. P., Bonmatin, J. M., Chagnon, M., Downs, C., … Wiemers, M. (2015). Systemic insecticides (Neonicotinoids and fipronil): Trends, uses, mode of action and metabolites. *Environmental Science and Pollution Research*, *22*(1), 5–34. https://doi.org/10.1007/s11356-014-3470-y

Simpson, M. G. L. (2019). permute: Functions for Generating Restricted Permutations of Data. R package version 0.9-5. Retrieved from https://cran.r-project.org/package=permute

Smedbol, É., Lucotte, M., Labrecque, M., Lepage, L., & Juneau, P. (2017). Phytoplankton growth and PSII efficiency sensitivity to a glyphosate-based herbicide (Factor 540®). *Aquatic Toxicology, 192*(September), 265–273. https://doi.org/10.1016/j.aquatox.2017.09.021

Smith, V. H., Joye, S. B., & Howarth, R. W. (2006). Eutrophication of freshwater and marine ecosystems. *Limnology and Oceanography*, *51*(1), 351–355.

Smith, V. H., & Schindler, D. W. (2009). Eutrophication science: where do we go from here? *Trends in Ecology and Evolution*, *24*(4), 201–207.

https://doi.org/10.1016/j.tree.2008.11.009

Song, X. P., Hansen, M. C., Stehman, S. V., Potapov, P. V., Tyukavina, A., Vermote, E.
F., & Townshend, J. R. (2018). Global land change from 1982 to 2016. *Nature*,
560(7720), 639–643. https://doi.org/10.1038/s41586-018-0411-9

Springmann, M., Clark, M., Mason-D'Croz, D., Wiebe, K., Bodirsky, B. L., Lassaletta, L.,
… Willett, W. (2018). Options for keeping the food system within environmental
limits. *Nature*, *562*(7728), 519–525. https://doi.org/10.1038/s41586-018-0594-0

Stachowski-Haberkorn, S., Becker, B., Marie, D., Haberkorn, H., Coroller, L., & de la
Broise, D. (2008). Impact of Roundup on the marine microbial community, as shown
by an in situ microcosm experiment. *Aquatic Toxicology*, *89*(4), 232–241.
https://doi.org/10.1016/j.aquatox.2008.07.004

Starr, A. V., Bargu, S., Maiti, K., & DeLaune, R. D. (2017). The Effect of Atrazine on
Louisiana Gulf Coast Estuarine Phytoplankton. *Archives of Environmental
Contamination and Toxicology*, *72*(2), 178–188. https://doi.org/10.1007/s00244-
016-0335-z

Stehle, S., & Schulz, R. (2015). Agricultural insecticides threaten surface waters at the
global scale. *Proceedings of the National Academy of Sciences of the USA*,
112(18), 5750–5755. https://doi.org/10.1073/pnas.1500232112

Thibodeau, G., Walsh, D. A., & Beisner, B. E. (2015). Rapid eco-evolutionary responses
in perturbed phytoplankton communities. *Proceedings of the Royal Society B*, *282*,
20151215. https://doi.org/10.1098/rspb.2015.1215

Thompson, M. S. A., Bankier, C., Bell, T., Dumbrell, A. J., Gray, C., Ledger, M. E., …
Woodward, G. (2016). Gene-to-ecosystem impacts of a catastrophic pesticide spill:
testing a multilevel bioassessment approach in a river ecosystem. *Freshwater
Biology*, *61*(12), 2037–2050. https://doi.org/10.1111/fwb.12676

Tilman, D., Fargione, J., Wolff, B., D'Antonio, C., Dobson, A., Howarth, R., …
Swackhamer, D. (2001). Forecasting agriculturally driven global environmental
change. *Science*, *292*(5515), 281–284. https://doi.org/10.1126/science.1057544

Tromas, N., Fortin, N., Bedrani, L., Terrat, Y., Cardoso, P., Bird, D., … Shapiro, B. J.
(2017). *Characterising and predicting cyanobacterial blooms in an 8-year amplicon
sequencing time course*. *The ISME Journal* (Vol. 11).
https://doi.org/10.1038/ismej.2017.58

van Bruggen, A. H. C., He, M. M., Shin, K., Mai, V., Jeong, K. C., Finchk, M. R., &
Morris, J. J. (2018). Environmental and health effects of the herbicide glyphosate.
Science of The Total Environment, *617*, 255–268.
https://doi.org/10.1016/j.scitotenv.2017.10.309

Van den Brink, P. J., den Besten, P. J., bij de Vaate, A., & ter Braak, C. J. F. (2009).
Principal response curves technique for the analysis of multivariate biomonitoring
time series. *Environmental Monitoring and Assessment*, *152*, 271–281.
https://doi.org/10.1007/s10661-008-0314-6

Van Rij, J., Wieling, M., Baayen, H. R., & van Rijn, H. (2020). Interpreting Time Series
and Autocorrelated Data Using GAMMs Author.

Vörösmarty, C. J., McIntyre, P. B., Gessner, M. O., Dudgeon, D., Prusevich, A., Green,
P., … Davies, P. M. (2010). Global threats to human water security and river
biodiversity. *Nature*, *467*(7315), 555–561. https://doi.org/10.1038/nature09440

Wang, C., Lin, X., Li, L., Lin, L., & Lin, S. (2017). Glyphosate Shapes a Dinoflagellate-
Associated Bacterial Community While Supporting Algal Growth as Sole
Phosphorus Source. *Frontiers in Microbiology*, *8*(2530).
https://doi.org/10.3389/fmicb.2017.02530
Weiss, S., Xu, Z. Z., Peddada, S., Amir, A., Bittinger, K., Gonzalez, A., … Knight, R.
(2017). Normalization and microbial differential abundance strategies depend upon
data characteristics. *Microbiome*, *5*(1), 27. https://doi.org/10.1186/s40168-017-
0237-y
Wetzel, R. G., & Likens, G. E. (2000). *Limnological Analyses. Springer Science &*
Business Media (3rd ed., Vol. 7). New York, New York, USA.
https://doi.org/10.1046/j.1440-1770.2002.01722.x
Wood, S. (2017). *Generalized Additive Models* (2nd ed.). New York: Chapman and
Hall/CRC. Retrieved from https://doi.org/10.1201/9781315370279
Wright, E. S. (2016). Using DECIPHER v2.0 to analyze big biological sequence data in
R. *The R Journal*, *8*(1), 352–359. https://doi.org/10.32614/rj-2016-025
Yamamuro, M., Komuro, T., Kamiya, H., Kato, T., Hasegawa, H., & Kameda, Y. (2019).
Neonicotinoids disrupt aquatic food webs and decrease fishery yields. *Science*,
366(6465), 620–623. https://doi.org/10.1126/science.aax3442
Zhang, Z., Qu, Y., Li, S., Feng, K., Wang, S., Cai, W., … Deng, Y. (2017). Soil bacterial
quantification approaches coupling with relative abundances reflecting the changes
of taxa. *Scientific Reports*, *7*(1), 1–11. https://doi.org/10.1038/s41598-017-05260-w
1215
1216
1217

1218   **Data accessibility**

1219   Sequence data corresponding to raw 16S rRNA reads and metadata, as well as carbon

1220   substrate utilization dataset based on Biolog EcoPlates assessment, are available on

1221   https://figshare.com/projects/MEC-LEAP/78297. Sequences of 16S rRNA reads are also

1222   available at NCBI SRA (BioProject ID PRJNA664121).

1223

Author contributions

N.B.C., V.F., M.-P.H., R.D.H.B., B.E.B., G.B., G.F.F., B.J.S. and A.G. designed the

study. N.B.C., V.F. and M.-P.H. collected the data. N.B.C., C.C.Y.X. and V. Y.

contributed to the development of laboratory methods. N.B.C. and V.F. analysed data.

N.B.C. made the figures and drafted the manuscript. N.B.C. and B.J.S. wrote the first

manuscript draft and all authors contributed significantly to data interpretation and

commented on manuscript drafts.

1231

Figure captions

Fig. 1 Experimental design and sampling timeline. A) In total, 48 mesocosms

(ponds) at the Large Experimental Array of Ponds (LEAP) at the Gault Nature Reserve

were filled with 1,000 L of pristine lake water and received two pulses of the pesticides

glyphosate and imidacloprid, alone or in combination, at two different nutrient

enrichment scenarios. Each box represents an experimental pond and those outlined in

bold indicate ponds sampled for DNA extraction and 16S rRNA gene amplicon

sequencing. B) The experiment lasted 43 days and pesticides were applied on days 6

(pulse 1) and 34 (pulse 2). Dates of sampling for each variable are indicated with points.

Nutrients were added every two weeks at a constant dose, starting seven days before

the first sampling day.

1243

1244 **Fig. 2 Bacterial cell density dynamics during the experiment.** Total bacterial density

1245 is plotted over time in ponds with low- or high- nutrient enrichment. Dashed vertical lines

1246 indicate days of pesticide pulses application. Ponds with both glyphosate and

1247 imidacloprid follow the same gradient pattern as treatments with either of these

1248 pesticides applied alone.

1249

Fig. 3 Microbial community carbon substrate utilization. Principal response curves

(PRCs) of selected experimental treatments show no significant difference between

controls and pesticide treatments when microbial communities are described according

to (A) their ability to metabolize 31 different carbon substrates when analysed

individually or (B) when grouped into guilds. Weights of each tested compound or guild

are shown along the Y axis (right). Dashed vertical lines indicate days of pesticides

pulses application. For ease of comparison, the PRCs were calculated based on the

subset of samples for which DNA was extracted. The PRC displayed in (A) explains

15.1% of the variation while the one displayed in (B) explains 42.2%, suggesting that

grouping substrates into guilds improves the explanatory power of the PRC.

1260

1261   **Fig. 4 Bacterioplankton alpha diversity variation across experimental treatments**

1262   **over time**, calculated as (A) the (log transformed) observed number of ASVs per

1263   sample and as (B) the exponent of Shannon index. Dashed vertical lines indicate days

1264   of pesticides pulses application.

1265

Fig. 5 PCoA ordinations of bacterioplankton community composition in response

to experimental treatments, based on (A) weighted UniFrac distance or (B) Jensen-

Shannon divergence calculated on ASV estimated absolute abundances after a DESeq2

normalization. Dashed vertical lines indicate days of pesticides pulses application. Each

sampling day is plotted in a separate panel to facilitate visualization of treatment effects

on community composition, mainly driven by high glyphosate (15 mg/L).

1272

Fig. 6 PRCs showing the effect of pesticide treatments over time relative to

control ponds at (A) the phylum level or (B) the ASV level. Dashed vertical lines

indicate days of pesticides pulses application. Only phyla with weights >0.2 and ASVs

with weight >0.1 are plotted on the right Y axis to facilitate visualization. The finest

available annotated level of taxonomic assignment of each ASV is indicated in panel B.

Low- and high-nutrient treatments were grouped together for clarity, but follow the same

pattern when considered separately (Fig. S4). See Fig. S5 for PRCs at other taxonomic

levels. These analyses were based on ASV estimated absolute abundances after a

DESeq2 normalization.

1282

Supplementary figures captions

Fig. S1 Experimental gradient established for (A) glyphosate and (B) imidacloprid

concentrations between two application pulses (at days 6 and 34) and (C) the

correlation between target and measured concentrations at each pulse. The top row of

figure C shows results for glyphosate, and the bottom two rows for imidacloprid, after

pulse 1 (left column) and pulse 2 (right column) respectively.

1289

1290 **Fig. S2** PRC plots show no effect of experimental treatments on community metabolic

1291 profiles when considering (A) the 31 different compounds individually (F=32.6 p=0.69) or

1292 (B) grouped according to functional guilds (F=79.2 p=0.86). The PRC axis shown in A

1293 explains 13.4% of total variance and in B 43.1%.

1294

1295 **Fig. S3** PCoA ordinations based on (A, B) weighted UniFrac distance or (B, D) Jensen-

1296 Shannon divergence exploring different normalization approaches: (A, C) calculation of

1297   reads relative abundance and (B, D) rarefying to a threshold of 10,000 reads per

1298   sample. Each sampling day is plotted separately to facilitate visualization of treatment

1299   effects on community composition.

1300

Fig. S4 PRCs show how high glyphosate treatments affected community composition at

(A) phylum and (B) ASV levels. Low- and high-nutrient treatments show the same

pattern, and, for this reason, they were grouped in Fig. 4, to facilitate data visualization.

The finest level of taxonomic assignment based on FreshTrain and GreenGenes

database is shown following ASV names in panel B. Only taxa with weights higher than

0.2 are shown in A and higher than 0.095 are shown in B.

1307

1308   **Fig. S5** PRCs show how high glyphosate treatments (15 mg/L) affected community

1309   composition at different taxonomic levels: (A) class, (B) order, (C) family/lineage, (D)

1310   genus/clade. Taxonomic assignment based on FreshTrain and GreenGenes databases.

1311   Low and high nutrient treatments were grouped as they follow the same pattern. Only

1312   taxa with weights higher than 0.2 are shown.

1313

Fig. S6 Summed effects of GAMMs on abundance of three genera most positively

affected by the glyphosate treatments: (A) Agrobacterium, (B) Flavobacterium and (C)

Azospirillum. Shades indicate a confidence interval of 95%. Abundance of each genus is

the estimated absolute abundance of all ASVs assigned to Agrobacterium,

Flavobacterium or Azospirillum after normalization by rarefying each sample to 10,000

reads without replacement.

Supplementary tables captions

Table S1 Regulatory acceptable concentrations (RACs) of glyphosate and imidacloprid

residues in freshwater according to regulatory agencies in Canada (CCME, Canadian

Council of Ministers of the Environment), Europe (EFSA, European Food Safety

Agency) and in the USA (EPA, Environmental Protection Agency). Chronic (long-term)

and acute (short-term) exposure RACs are specified when available.

Table S2 Carbon substrates present in Biolog EcoPlates and their respective grouping

(guild)

Table S3 Barcode sequences of the reverse primer used in step 2 PCR, and total read

counts obtained after sample demultiplexing. The number of non-chimeric reads

obtained after filtering, denoising, merging paired ends and removing chimeras with

DADA2, is also shown.

Table S4 Summarized results of the generalized additive mixed models (GAMMs) for

bacterial density and number of carbon substrate used as a response variables. A

Gaussian residual distribution was used for both models. For each response variable we

report the sample size (n), adjusted R^2, the predictors used in the model, the parameter

estimate and respective standard error (SE) of parametric effects or the effective

degrees of freedom (EDF) of smooth terms, the corresponding test statistics (*t* value for

parametric and F for smooth terms) and the *p*-value. Smooths terms are described as

mgcv syntax: 's()' functions are thin plate regression splines and 'ti()' tensor product

1343   interactions, pond represents the random variable of the mixed model and 'bs='fs''

1344   specifies the underline base function as a random smooth. Following a Bonferroni

1345   multiple testing correction for 9 tests, we only considered significant variables with

1346   unadjusted p-value <0.005 (shown in bold).

1347

Table S5 Summarized results of the generalized additive mixed models (GAMMs) for

alpha diversity: observed ASV and exponential Shannon. Gaussian residual distributions

were used in all models. For each response variable we report the sample size (n),

adjusted R2, the predictors and factors used in the model, the parameter estimate and

respective standard error (SE) of parametric effects or the effective degrees of freedom

(EDF) of smooth terms, the corresponding test statistics (t value for parametric and F for

smooth terms) and the p-value. Smooths terms are described as mgcv syntax: 's()'

functions are thin plate regression splines and 'ti()' tensor product interactions, pond

represents the random variable of the mixed model and 'bs='fs'' specifies the underline

base function as a random smooth. Following a Bonferroni multiple testing correction for

16 tests, we only considered significant variables with p-value <0.003, shown in bold.

1359

Table S6 PERMANOVA for different explanatory variables (and their interaction) in

models with the weighted UniFrac distances among communities as the response. The

same model was tested at five different time points and an asterisk indicates p-values

that are significant after a Bonferroni correction for 7 hypothesis tests (i.e. p-values

<0.007 are considered significant, shown in bold). A PERMDISP was performed to

confirm homogeneity of groups dispersions and significant p-values (<0.05) point out to

predictors whose significance in the PERMANOVA output should be carefully analysed

as they may be a result of within-group variation rather than among-group variation.

df=degrees of freedom

1369

Table S7 PERMANOVA for different explanatory variables (and their interaction) in

models with the Jensen-Shannon divergence among communities as the response. The

same model was tested at five different time points and an asterisk indicates p-values

that are significant after a Bonferroni correction for 7 hypothesis tests (i.e. only p-values

<0.007 are considered significant, shown in bold). A PERMDISP was performed to

confirm homogeneity of groups dispersions and significant p-values (<0.05) point out to

predictors whose significance in the PERMANOVA output should be carefully analysed

as they may be a result of within-group variation rather than among-group variation.

df=degrees of freedom

1379

Table S8 PERMANOVA for glyphosate as the explanatory variable in models with

weighted UniFrac distance or Jensen-Shannon divergence among communities as the

response variable after data transformation by ASV relative abundance calculation

(unrarefied) or by rarefying samples to 10,000 reads. The same model was tested at five

different time points and an asterisk indicates p-values that are significant after a

conservative Bonferroni correction for 7 hypothesis tests (i.e. only p-value<0.007 are

considered significant). A PERMDISP was performed to confirm homogeneity of groups

dispersions and significant p-values (<0.05) point out to predictors whose significance in

the PERMANOVA output should be carefully analysed as they may be a result of within-

group variation rather than among-group variation.

1390

1391   **Table S9** Percent of variance explained by the two first PRC axes, and by time or

1392   treatment when nutrient treatments are grouped as replicates (see Fig. 6 and Fig. S5). F

1393   statistic and p-value of permutation test for first constrained eigenvalue is also shown,

1394   and an asterisk denote significant p-values.

1395

1396   **Table S10** All bacterioplankton taxa weights for the PRC model at the phylum level,

1397   ranked from largest (positive effects of glyphosate treatment) to smallest (negative

1398   effects of glyphosate treatment).

1399

1400   **Table S11** Relative abundance of the main affected phyla by treatment. Percentage was

1401   calculated after normalization with DESeq2 or by rarefying samples to 10,000 reads

1402   each and the respective standard error is indicated in parenthesis.

1403

1404   **Table S12** ASVs with the highest PRC taxa weights, and their respective weight in the

1405   first RDA axis, the ratio between this value and the maximum taxa weight of the PRC

1406   model, and their taxonomy assignment from TaxAss using FreshTrain and GreenGenes

1407   databases.

1408

Table S13 Summarized results of the generalized additive mixed models (GAMMs) for

abundance of the three genera most positively impacted by the experimental treatments.

Gaussian residual distributions were used in all models. For each response variable we

report the sample size (n), adjusted R2, the predictors and factors used in the model,

the parameter estimate and respective standard error (SE) of parametric effects or the

effective degrees of freedom (EDF) of smooth terms, the corresponding test statistics (t

value for parametric and F for smooth terms) and the p-value. Smooths terms are

described as mgcv syntax: 's()' functions are thin plate regression splines and 't()' tensor

product interactions, pond represents the random variable of the mixed model and

'bs='fs'' specifies the underline base function as a random smooth. Following a

Bonferroni multiple testing correction for 16 tests, we only considered significant

variables with p-value <0.003, shown in bold.

A

Glyphosate (mg/L)

| **0** | 0.04 | 0.1 | **0.3** | 0.7 | 2.0 | 5.5 | **15** |

Imidacloprid (μg/L)

| **0** | 0.15 | 0.4 | **1.0** | 3.0 | 8.0 | 22 | **60** |

Glyphosate (mg/L) and imidacloprid (μg/L)

| 0/0 | 0.04/0.15 | 0.1/0.4 | **0.3/1.0** | 0.7/3.0 | 2.0/8.0 | 5.5/22 | **15/60** |

Replicated in low and high nutrient backgrounds

B

Figure showing sampling schedule: Pulse 1 at day 6, Pulse 2 at day 34. Rows: Microbial carbon substrate use; Bacterioplankton (DNA and density); Total nitrogen; Total phosphorus; Imidacloprid; Glyphosate. Day of experiment on x-axis (0–44).

A Weighted-UniFrac distance

B Jensen-Shannon divergence

Pesticide treatment
- Control
- Glyphosate (0.3 mg/L)
- Glyphosate (15 mg/L)
- Imidacloprid (1 μg/L)
- Imidacloprid (60 μg/L)
- Glyphosate (0.3 mg/L) and imidacloprid (1 μg/L)
- Glyphosate (15 mg/L) and imidacloprid (60 μg/L)

Nutrient concentration
- Low
- High