***Jchain*-driven cre enables specific genetic manipulation**

and timestamping of plasma cell in their niche.

Xu AQ[1], Barbosa RR[1], Calado DP[1,2,3*]

1. Immunity and Cancer, The Francis Crick Institute, 1 Midland Road, London NW1 1AT, UK.

2. Peter Gorer Department of Immunobiology, School of Immunology & Microbial Sciences,

King's College London, London, UK.

3. Lead contact.

* Corresponding author: Dinis.Calado@crick.ac.uk

Keywords

Jchain, *Blimp1*, *Irf4*, plasma cells, B cells, cre-loxP

Highlights

• *Jchain*-driven cre-mediated genetic manipulation is effective only in plasma cells

• *Jchain* expression occurs in plasma cells across immunoglobulin isotypes

• Genetic manipulation using Jchain-driven cre targets plasma cells already in their niche

• *Jchain*-driven cre allows tracing of long-lived plasma cells

Summary

Plasma cells are essential for long-term protection from infection and underlie vaccination success.

However, studies of gene function in plasma cells are hindered by technical limitations. Here we report

an engineered allele that allows to trace and genetically manipulate plasma cells.

Abstract

Plasma cells produce antibodies that are essential for protection from infection and for vaccination

success. Plasma cells are also at the origin of currently incurable hematological malignancies, most

notably multiple myeloma. However, the study of gene function in plasma cells has mostly been

performed using *in vitro* cultures and cell transfer systems that remove plasma cells from their

microenvironment. Further, existing systems to genetically manipulate plasma cells are not specific and

target multiple cell lineages including B cells, making it difficult to discriminate gene function in plasma

cell formation and/or maintenance. Here we characterized a genetically engineered allele: *Igjcre$^{ERT2(GFP)}$*

that expresses GFP and cre^{ERT2} under the activity of the endogenous *Jchain* promoter. We found at the

single cell level that *Jchain* expression occurred in plasma cells across immunoglobulin isotypes. Also

using the *Igjcre$^{ERT2(GFP)}$* we performed highly specific cre-loxP genetic manipulation of plasma cells

residing in their natural microenvironment *in vivo* in the mouse and traced long-lived plasma cells.

Introduction

Plasma cells produce antibodies that are crucial for immune protection against infection and for vaccination success (Nutt et al., 2015). Upon activation, B cells terminally differentiate into plasma cells, a process initiated by the downregulation of the B cell transcription factor PAX5 (Kallies et al., 2007). This event allows the expression of multiple factors normally repressed by PAX5, including *Xbp1* and *Jchain* (Castro and Flajnik, 2014; Nutt et al., 2015; Rinkenberger et al., 1996; Shaffer et al., 2004). PAX5 downregulation is also followed by the expression of the transcription factors IRF4 and BLIMP1 that are essential for the establishment of the plasma cell program (Kallies et al., 2007; Klein et al., 2006; Shapiro-Shelef et al., 2003). Beyond physiology, multiple cancers have a plasma cell as the cell of origin, including multiple myeloma that is the second most frequent hematological malignancy overall and for which a cure remains to be found (Palumbo and Anderson, 2011). As consequence, the study of gene function in plasma cell homeostasis is a subject of intense investigation. However, because of the lack of tools many studies make use of *in vitro* and cell transfer systems that remove plasma cells from their microenvironment. Further, genetic manipulation of plasma cells is currently not specific, targeting other cellular populations including B cells. These strategies are limited in their ability to study gene function in plasma cells in their microenvironment and confound plasma cell formation and maintenance.

We found that amongst plasma cell associated genes, *Jchain* (*Igj*) had the highest level and most specific expression in plasma cell populations. JCHAIN is a small polypeptide that is required to multimerize IgM and IgA and that is necessary for the transport of these Ig classes across the mucosal epithelium in a poly-Ig receptor mediated process (Brandtzaeg and Prydz, 1984; Castro and Flajnik, 2014; Max and Korsmeyer, 1985). Here, we characterized a GFP-tagged cre^{ERT2} allele: *Igjcre$^{ERT2(GFP)}$* at the *Jchain* endogenous locus. At the single cell level *Jchain* expression occurred in plasma cells across immunoglobulin isotypes, including IgG1. Using the *Igjcre$^{ERT2(GFP)}$* we performed highly specific cre-loxP genetic manipulation of plasma cells residing in their natural microenvironment *in vivo* and traced long-lived plasma cells. *Igjcre$^{ERT2(GFP)}$* is a validated allele for plasma cell studies, allowing the mechanistic understanding of gene function in plasma cells *in vivo* and in their niche.

Results

Jchain transcripts are highly enriched in plasma cell populations

B-to-plasma cell differentiation is a process that involves a complex network of factors (**Fig. 1A**; (Nutt et al., 2015)). We investigated the level and specificity of the expression of genes associated with plasma cells (*Xbp1*, *Jchain*, *Scd1*, *Irf4*, and *Prdm1*) through the analysis of a publicly available RNA sequencing dataset (ImmGen, (Heng et al., 2008)). We first determined the cell populations with the highest transcript level for each factor (**Fig. 1B**). *Xbp1* and *Irf4* were primarily expressed in plasma cell populations, however the expression in bone marrow plasma cells (B_PC_BM) was less than 2-fold greater than that of non-plasma cell populations (**Fig. 1B**). The expression of *Sdc1* and *Prdm1* was not specific to plasma cell populations (**Fig. 1C**). Peritoneal cavity macrophages (MF_226+II+480lo_PC) expressed more *Sdc1* than bone marrow plasma cells (B_PC_BM), and a subset of FOXP3$^+$ cells (Treg_4_FP3+_Nrplo_Co) expressed higher levels of *Prdm1* than that observed in splenic plasmablasts (B_PB_Sp) and bone marrow plasma cells (B_PC_BM; **Fig. 1C**). In contrast, *Jchain* had the highest level of transcript expression in plasma cell populations compared to non-plasma cells and was the most plasma cell specific amongst all factors, with a 40-fold enrichment over germinal center (GC) B cells (B_GC_CB_Sp; **Fig. 1D**). We concluded that the *Jchain* was a suitable candidate for genetic engineering and generation of plasma cell specific tools.

Jchain is expressed in a small fraction of GC B cells and in most plasma cells

We searched alleles produced by the EUCOMMTools consortium and identified a genetically engineered *Jchain* allele produced by the Wellcome Trust Sanger Institute: MGI:5633773, hereafter termed: *Igjcre*$^{ERT2(GFP)}$. The genetically engineered *Jchain* allele contained an *FRT* site between exons 1 and 2 followed by an engrailed 2 splice acceptor sequence and an *EGFP.2A.cre*^{ERT2} expression cassette (**Fig. 2A**). In this design the expression of the EGFP (GFP) and of cre^{ERT2} is linked by a self-cleaving 2A peptide under the transcriptional control of the *Jchain* promoter (**Fig. 2A**). To determine cells with GFP expression we initially analyzed the spleen from mice heterozygous for the *Igjcre*$^{ERT2(GFP)}$ allele that had been immunized 12 days earlier **(Fig. 2B)**. B220 is expressed on the surface of B cells and downregulated during plasma cell differentiation (Pracht et al., 2017). We therefore defined three

cell populations based on the levels of GFP fluorescence and B220 surface expression: $GFP^{low}B220^{high}$,

$GFP^{int}B220^{int}$, $GFP^{high}B220^{low}$, and a population negative for both markers ($GFP^{neg}B220^{neg}$; **Fig. 2C**).

We next determined the fraction of cells within these populations expressing surface CD138, that is a

commonly used marker to define plasma cells by flow-cytometry (Pracht et al., 2017). The

$GFP^{neg}B220^{neg}$ population did not contain $CD138^+$ cells, however, the fraction of $CD138^+$ cells

increased in the remaining populations in agreement with the reduction of B220 during plasma cell

differentiation, with the $GFP^{int}B220^{int}$ and $GFP^{high}B220^{low}$ populations being mostly composed of

$CD138^+$ cells (**Fig. 2D and E**). Identical results were observed when defining plasma cells using an

additional marker, the CXCR4 chemokine receptor that facilitates homing of plasma cells to the bone

marrow (**Fig. 2D and E**; (Hargreaves et al., 2001)). We investigated the cellular composition of the

$GFP^{low}B220^{high}$, that contained the fewest $CD138^+$ cells (1 to 20%; **Fig. 2E**). We found that virtually all

$CD138^{neg}$ cells within $GFP^{low}B220^{high}$ expressed the B cell marker CD19 (**Fig. 2F**). Also and in

agreement with *Jchain* gene expression analysis (**Fig. 1D**) these cells were mostly GC B cells

($CD38^{low}FAS^{high}$; **Fig. 2F and G**). We also performed analyses in the spleen and bone marrow of 12-

day immunized mice heterozygous for the $Igjcre^{ERT2(GFP)}$ allele in which mature and B cell precursors

as well as non-B cell populations were first defined using surface markers and the fraction of GFP

expressing cells within those populations determined (**Fig. S1**). We found that a small fraction of B1b

cells (0 to 6%) in the spleen and most plasma cells ($CD138^+CXCR4^+$, 60 to 90%) in the spleen and

bone marrow expressed GFP (**Fig. S1**). B1b cells are prone to differentiate into plasma cells and are a

source of IgM antibodies upon T cell independent responses (Alugupalli et al., 2004). Collectively these

data confirmed at the single cell level the analysis of gene expression using bulk populations (**Fig. 1D**)

and suggested that *Jchain* expression is enriched in plasma cells.

***Jchain* expression correlates with IRF4 and BLIMP1 co-expression**

IRF4 and BLIMP1 transcription factors play an essential role in plasma cell differentiation (Kallies et

al., 2007; Klein et al., 2006; Shapiro-Shelef et al., 2003). We therefore analyzed the spleen of 12-day

immunized mice heterozygous for the $Igjcre^{ERT2(GFP)}$ allele and determined whether IRF4 and BLIMP1

were expressed in the populations defined by varied GFP and B220 surface expression (**Fig. 2C, 3A**

141 and **B**). The GFP^{neg}B220^{neg} cells were virtually devoid of BLIMP1 and IRF4 expression (**Fig. 3C**). In

142 contrast the GFP^{low}B220^{high} cell population contained ~7% (median) of BLIMP1$^+$IRF4$^+$ cells (**Fig. 3D**

143 and **E**). This suggested that *Jchain* expression can precede that of IRF4 and BLIMP1, something

144 previous suggested from *in vitro* cultures of *Blimp1* deficient B cells (Kallies et al., 2007). Still,

145 increased *Jchain* expression strongly correlated with that of IRF4 and BLIMP1 as the vast majority of

146 cells within the GFP^{int}B220^{int} and GFP^{high}B220^{low} populations were BLIMP1$^+$IRF4$^+$ (**Fig. 3D** and **E**).

147 These data confirmed *in vivo* the observations of others *in vitro* (Kallies et al., 2007) and identified

148 population subsets in which *Jchain* expression precedes that of IRF4 and BLIMP1. The data also

149 demonstrated that increased levels of *Jchain* promoter-driven GFP expression correlates highly with

150 cells expressing critical plasma cell transcription factors supporting the observations made using cell

151 surface markers (**Fig. 2**).

153 ***Igjcre$^{ERT2(GFP)}$* mediated cre recombination is effective only in plasma cells**

154 We next sought to determine whether the *Igjcre$^{ERT2(GFP)}$* could be used to perform genetic manipulation

155 of plasma cells. We generated compound mutant mice carrying the *Igjcre$^{ERT2(GFP)}$* allele and a Rosa 26

156 allele in which RFP expression is conditional to cre-mediated recombination of a loxP-STOP-loxP

157 cassette (*R26^{lslRFP}*; (Luche et al., 2007)). In the *Igjcre$^{ERT2(GFP)}$* cre is fused to an estrogen binding domain

158 (ERT2) that sequesters cre in the cytoplasm through the binding to HSP90 (Feil et al., 2009) . Addition

159 of tamoxifen to the system displaces the cre^{ERT2}-HSP90 complex effectively allowing nuclear import of

160 cre^{ERT2} and its access to loxP flanked DNA sequences (Feil et al., 2009). We first performed an *in vitro*

161 experiment using a classical plasmablast (B220^{low}CD138$^+$) differentiation assay in which B cells

162 purified from mice carrying the *Igjcre$^{ERT2(GFP)}$* and *R26^{lslRFP}* alleles were cultured with LPS in the

163 presence or absence of 4-OH tamoxifen (Andersson et al., 1972). GFP expression was highly enriched

164 in plasmablasts compared to B cells and RFP expression was only observed upon addition of 4-OH

165 tamoxifen to the culture, and that occurred virtually only in plasmablasts (**Fig. S2**). This data suggests

166 that cre^{ERT2} was mostly induced in plasmablasts and was effectively retained in the cytoplasm in the

167 absence of 4-OH tamoxifen. Similar observations were made in mice carrying the *Igjcre$^{ERT2(GFP)}$* and

168 *R26^{lslRFP}* alleles *in vivo* in that absence of tamoxifen administration RFP expressing cells were not

detected, confirming the *in vitro* results and supporting that the *Igjcre^ERT2(GFP)* allele was not "leaky" in

the control of cre activity (**Fig. S1B**, **C** and **E**). We next immunized mice carrying *Igjcre^ERT2(GFP)* and

R26^lslRFP alleles and administered tamoxifen on days 7, 8, 9 and 10 followed by analysis of spleen and

bone marrow at day 12 (**Fig. 4A**). Analyses of the populations defined by varied GFP and B220 surface

expression (**Fig. 2 and 4B**) revealed that a small fraction of GFP^{low}B220^{high} cells were positive for RFP

in the spleen (median ~2%) and in the bone marrow (median ~1.2%; **Fig. 4C**). In contrast, cre mediated

recombination and as consequence RFP expression occurred in ~37% and ~31% of GFP^{int}B220^{int} in the

spleen and bone marrow, respectively, whereas the vast majority of cells within the GFP^{high}B220^{low}

population had undergone cre mediated recombination and were RFP positive (~76% in the spleen, and

~88% in the bone marrow, median; **Fig. 4C and D**). These data showed that *Igjcre^ERT2(GFP)* mediated cre

recombination was only effective in plasma cell populations and validated *Igjcre^ERT2(GFP)* as a tool to

specifically perform genetic manipulation of plasma cells.

Genetic manipulation using *Igjcre^ERT2(GFP)* in plasma cells occurs across immunoglobulin isotypes

IgG1 does not multimerize, and due to differences in its secretory tail to that of IgA and IgM, JCHAIN

does not associate with IgG1 (Johansen et al., 2000). It is currently assumed that *Jchain* expression

occurs in all plasma cells regardless of isotype, however, this has not been demonstrated at the single

cell level (Castro and Flajnik, 2014; Johansen et al., 2000; Mather et al., 1981). We performed

experiments that investigated whether *Igjcre^ERT2(GFP)* allele GFP expression (as proxy for *Jchain*), and

cre-mediated loxP recombination occurred in plasma cells across immunoglobulin isotypes. For that

we analyzed the spleen, mLN, peyer's patches of younger (15 weeks) and older (30 weeks) mice

carrying the *Igjcre^ERT2(GFP)* and *R26^lslRFP* alleles. These mice were immunized and administered with

tamoxifen on days 7, 8, 9 and 10 followed by analysis at day 12 (**Fig. 5A**). We first gated on total

plasma cells (B220^{low}CD138^{+}), and then on total GFP^{+} cells (RFP^{+} and RFP^{neg}, i.e. *Jchain* expressing)

and after that on GFP^{+}RFP^{+} cells (i.e. cre recombined) and determined the proportions of IgA, IgM and

IgG1 expressing cells within the plasma cell population and subpopulations using intracellular stain

(**Fig. 5B-D**). Analysis of spleens of 15-week-old mice revealed a slightly increased representation of

IgA^{+} cells within the GFP^{+}RFP^{+} plasma cells only when compared to total plasma cells (**Fig. 5E**). A

similar observation was made when analyzing the peyer's patches of 15-week-old mice (**Fig. 5E**).

However, for 15-week-old mice we did not observe differences in the representation of IgA in any of

the other tissues analyzed, and in none of the tissues for IgM and IgG1 (**Fig. 5E**). Analysis of 30-week-

old mice showed a slightly increased representation of IgM^+ cells within the GFP^+RFP^+ plasma cells

only when compared to total plasma cells, and for IgA^+ in mesenteric lymph node (mLN; **Fig. 5F**).

Nevertheless, for 30-week-old mice we did not observe a significant difference in the representation of

IgM or IgA in any of the other tissues analyzed, and in none of the tissues for IgG1 (**Fig. 5F**). Taken

together, these data suggested that *Jchain* expression is not overly represented in IgA or IgM expressing

plasma cells compared to $IgG1^+$ plasma cells. As consequence, only minor differences were observed

for *Igjcre*[ERT2(GFP)] mediated cre-loxP recombination across immunoglobulin isotypes indicating that the

Igjcre[ERT2(GFP)] is useful for genetic manipulation also of plasma cells expressing IgG1.

Igjcre[ERT2(GFP)] **allows lifespan studies and to trace long-lived plasma cells**

The lifespan of plasma cells has been a matter of debate and investigation for over 70 years (Manz and

Radbruch, 2002). This knowledge is important as it directly relates to long-term protection from

infection and could provide valuable information to improve vaccines. Multiple analyses using

primarily BrdU labeling pulse-chase experiments have generated conflicting results with some studies

suggesting a 3-week lifespan for a plasma cell and others more than 3 months (Ho et al., 1986; Manz

et al., 1997). These experiments have also showed that the half-life of plasma cells differs between

spleen and the bone marrow (Sze et al., 2000). We investigated the suitability of the *Igjcre*[ERT2(GFP)] allele

to determine the lifespan of plasma cells in the spleen and bone marrow. We immunized mice carrying

Igjcre[ERT2(GFP)] and *R26*[lslRFP] alleles at two time-points spaced by a period of 21 days (**Fig. 6A**, (Calado

et al., 2010)). Thirty days after the secondary immunization (day 51) we administered tamoxifen for 5

consecutive days to genetically label plasma cells (**Fig. 6A**). Next we determined genetically labelled

GFP^+RFP^+ plasma cells present in the spleen and bone marrow at day 5, and after 2 and 5 months after

the last tamoxifen administration (**Fig. 6B** and **C**). Compared to day 5, in which ~70% (median) of

$CD138^+CXCR4^+$ plasma cells in spleen and bone marrow were GFP^+RFP^+, that fraction was

significantly reduced in spleen and bone marrow at two months and more so at five months after the

last tamoxifen injection (**Fig. 6B** and **C**). As expected, with the generation of new plasma cells over

time, the opposite was true for the fraction of GFP^{+}RFP^{neg} plasma cells (**Fig. 6B** and **C**). Remarkably,

by five months the reduction in the fraction of GFP^{+}RFP^{+} plasma cells was more pronounced in the

spleen compared to the bone marrow (**Fig. 6B** and **C**). Still the data indicated that ~6.4% of plasma

cells in the spleen and ~16.4% in the bone marrow lived for at least 5 months (**Fig. 6B** and **C**). We next

calculated the half-life of plasma cells in the spleen and bone marrow. Consistent with the sharper

decrease of genetically labelled GFP^{+}RFP^{+} plasma cells in the spleen over time, splenic GFP+RFP+

plasma cells displayed a calculated half-life of ~49.5 days, whereas that of GFP^{+}RFP^{+} plasma cells in

the bone marrow was ~86.4 days. Notably the calculated half-life of genetically labelled plasma cells

in the bone marrow was consistent with BrdU labeling pulse-chase experiments suggesting that a

fraction of plasma cells survives for at least 3 months (Manz et al., 1997). These data demonstrated the

suitability of *Igjcre$^{ERT2(GFP)}$* in time-stamping plasma cells using the expression of a fluorescent marker

and to identify long-lived plasma cells.

Discussion

Plasma cells are an essential component of the adaptive immune system. However, compared to other B cell lineage populations, the molecular analysis of gene function *in vivo* as lagged behind. One underlying cause is the lack of tools for specific genetic manipulation, something available for B cells since 1997, more than 20 years ago (Rickert et al., 1997), and recently for B cells at multiple stages of development (Boross et al., 2009; Casola et al., 2006; Croker et al., 2004; Crouch et al., 2007; de Boer et al., 2003; Dogan et al., 2009; Duber et al., 2009; Georgiades et al., 2002; Hobeika et al., 2006; Kraus et al., 2004; Kwon et al., 2008; Moriyama et al., 2014; Robbiani et al., 2008; Schweighoffer et al., 2013; Shinnakasu et al., 2016; Weber et al., 2019; Yasuda et al., 2013).

Here we used a systematic analysis of developmental factors for which expression is increased during plasma cell differentiation (*Xbp1*, *Jchain*, *Scd1*, *Irf4*, and *Prdm1*) to identify suitable candidates for the generation of plasma cell-specific tools. *Blimp1*, the critical transcription factor for the establishment of the plasma cell program, displayed an expression pattern unspecific to plasma cells. This was unsurprising given the knowledge that *Blimp1* is expressed by other hematopoietic and non-hematopoietic cells (John and Garrett-Sinha, 2009), including germ cells (Ohinata et al., 2005). Amongst the analyzed factors we found *Jchain* to be the one with the highest RNA expression level in plasma cells and the most plasma cell-specific expression pattern. We therefore considered *Jchain* to be the most suitable candidate for the generation of plasma cell-specific tools.

We searched alleles generated by the EUCOMMTools consortium and identified a *Jchain* allele: *Igjcre^ERT2(GFP)^* and tested its functionally. Through a series of experiments, we characterized this allele and demonstrated its suitability for specific genetic manipulation of plasma cells. Briefly, although *Jchain* was expressed in a small subset of B cells (primarily GC B cells) and preceded BLIMP1 and IRF4 expression, *Igjcre^ERT2(GFP)^* mediated cre recombination was only effective in plasma cells. In the *Igjcre^ERT2(GFP)^* allele GFP and cre^ERT2^ are linked by a self-cleaving 2A peptide which ensures a near equitable co-expression of the proteins (Szymczak et al., 2004). As consequence, it is likely that effectiveness of cre recombination is relatable to the level of cre^ERT2^ because plasma cells expressed the

highest level of *Jchain*, as suggested by the GFP expression. Of note, the *Igjcre$^{ERT2(GFP)}$* allele analyzed

in this work retained the puromycin selection cassette that is flanked by Rox recombination sites. This

knowledge should be taken in consideration in future experiments involving compound mutant mice

that make use of Dre recombinase (Anastassiadis et al., 2009). On occasion it has been observed that

retention of the selection cassette reduces gene expression of the targeted locus (Meyers et al., 1998;

Nagy et al., 1998). As consequence removal of the selection may potentially increase cre^{ERT2} expression,

including in B cells, and reduce the specificity of *Igjcre$^{ERT2(GFP)}$* cre-mediated recombination to plasma

cells. Lastly in the *Igjcre$^{ERT2(GFP)}$* allele the expression of *Jchain* is interrupted by a GFP-2A-creERT2

cassette and induction of cre-mediated recombination deletes *Jchain* exon 2 that is loxP-flanked. It was

previously shown that heterozygous deletion of *Jchain* displayed an intermediate phenotype to that of

knockout mice (Lycke et al., 1999). This knowledge must be taken into consideration when performing

Igjcre$^{ERT2(GFP)}$ cre-mediated genetic manipulation, including the need to use the *Igjcre$^{ERT2(GFP)}$* allele

without the targeted manipulation as control, and the utility of the *Igjcre$^{ERT2(GFP)}$* allele in homozygosity.

Future iterations of the *Igjcre$^{ERT2(GFP)}$* in which JCHAIN protein expression is preserved should be

considered.

Previous analysis of *in vitro* cultures of *Blimp1* deficient B cells showed that initiation of *Jchain*

expression does not require BLIMP1 (Kallies et al., 2007). However, *Jchain* expression is higher in

BLIMP1 expressing cells (Kallies et al., 2007). We used the *Igjcre$^{ERT2(GFP)}$* allele and identified a

population of cells *in vivo* marked by low GFP expression and high surface expression of B220

(GFP^{low}B220^{hi}) where only ~7% of cells expressed BLIMP1. It is thus possible that the GFP^{low}B220^{hi}

population identified in mice carrying the *Igjcre$^{ERT2(GFP)}$* allele contains BLIMP1^{neg} plasma cell

precursor cells and future in-depth analysis may provide information on the mechanisms underlying the

very first steps of plasma cell differentiation.

It is generally assumed that *Jchain* expression occurs in all plasma cells regardless of their isotypes

(Castro and Flajnik, 2014; Johansen et al., 2000; Mather et al., 1981). The analysis performed in this

work using the *Igjcre$^{ERT2(GFP)}$* allele confirmed at the single cell level that *Jchain* expression occurs in

plasma cells across immunoglobulin isotypes. However, given that in mice carrying the *Igjcre*[ERT2(GFP)]

allele not all plasma cells expressed GFP, it did not support the concept *Jchain* is expressed by all

plasma cells. It is, however, necessary to consider that these analysis measure GFP expression that is

under the activity of the *Jchain* promoter, rather than JCHAIN directly.

We also demonstrated that the *Igjcre*[ERT2(GFP)] allele is a useful tool to study plasma cell life-span and to

identify long-lived plasma cells. Until now, most studies aiming to determine plasma cell longevity

have relied on BrdU or EdU labeling pulse-chase experiments (Ho et al., 1986; Lemke et al., 2016;

Manz et al., 1997). There are key differences between the genetic and the BrdU/EdU time-stamping

methods. The former labels existing plasma cells independently of when these cells were generated and

even if labelled cells were to undergo cell cycle the genetic mark would be maintained. On other hand,

BrdU or EdU labelling pulse-chase experiments are strictly dependent on cell cycle and therefore more

accurately mark the time at which plasma cells were generated, however, if division occurs in the

plasma cell pool the label may be lost. Still, the calculated half-life of plasma cells using the

Igjcre[ERT2(GFP)] allele was similar to that of some EdU or BrdU labelling studies (Lemke et al., 2016;

Manz et al., 1997). Where the *Igjcre*[ERT2(GFP)] allele will supersede BrdU labelling technologically is on

the ability to retrieve time-stamped live plasma cells at multiple time-points for downstream analysis.

That property can easily be combined with further genetic manipulation to investigate gene function,

paving the way for in-depth mechanistic understanding of physiology and pathology of plasma cells in

in vivo and in their niche.

Acknowledgments

We thank the members of the Immunity and Cancer laboratory (FCI, London) for critical discussions and review of the manuscript; the FCI BRF and Flow-cytometry platforms for expert advice and technical support.

This work was supported by The Francis Crick Institute which receives its core funding from Cancer Research UK (FC001057), the UK Medical Research Council (FC001057), and the Wellcome Trust (FC001057) to D.P.C., an MRC career development award MR/J008060/1 to D.P.C.

The authors declare no competing financial interests.

Author contributions: Conceptualization, A.Q.X., R.R.B, D.P.C.; Methodology, A.Q.X., R.R.B, D.P.C.; Investigation, A.Q.X., R.R.B, D.P.C.; Resources, D.P.C.; Writing – Original Draft, A.Q.X., and D.P.C.; Supervision D.P.C.; Funding Acquisition, D.P.C.

References

Alugupalli, K.R., J.M. Leong, R.T. Woodland, M. Muramatsu, T. Honjo, and R.M. Gerstein. 2004. B1b lymphocytes confer T cell-independent long-lasting immunity. *Immunity* 21:379-390.

Anastassiadis, K., J. Fu, C. Patsch, S. Hu, S. Weidlich, K. Duerschke, F. Buchholz, F. Edenhofer, and A.F. Stewart. 2009. Dre recombinase, like Cre, is a highly efficient site-specific recombinase in E. coli, mammalian cells and mice. *Dis Model Mech* 2:508-515.

Andersson, J., O. Sjoberg, and G. Moller. 1972. Induction of immunoglobulin and antibody synthesis in vitro by lipopolysaccharides. *Eur J Immunol* 2:349-353.

Bonami, R.H., W.T. Wolfle, J.W. Thomas, and P.L. Kendall. 2014. NFATc2 (NFAT1) assists BCR-mediated anergy in anti-insulin B cells. *Mol Immunol* 62:321-328.

Boross, P., C. Breukel, P.F. van Loo, J. van der Kaa, J.W. Claassens, H. Bujard, K. Schonig, and J.S. Verbeek. 2009. Highly B lymphocyte-specific tamoxifen inducible transgene expression of CreER T2 by using the LC-1 locus BAC vector. *Genesis* 47:729-735.

Brandtzaeg, P., and H. Prydz. 1984. Direct evidence for an integrated function of J chain and secretory component in epithelial transport of immunoglobulins. *Nature* 311:71-73.

Calado, D.P., B. Zhang, L. Srinivasan, Y. Sasaki, J. Seagal, C. Unitt, S. Rodig, J. Kutok, A. Tarakhovsky, M. Schmidt-Supprian, and K. Rajewsky. 2010. Constitutive canonical NF-kappaB activation cooperates with disruption of BLIMP1 in the pathogenesis of activated B cell-like diffuse large cell lymphoma. *Cancer Cell* 18:580-589.

Casola, S., G. Cattoretti, N. Uyttersprot, S.B. Koralov, J. Seagal, Z. Hao, A. Waisman, A. Egert, D. Ghitza, and K. Rajewsky. 2006. Tracking germinal center B cells expressing germ-line immunoglobulin gamma1 transcripts by conditional gene targeting. *Proc Natl Acad Sci U S A* 103:7396-7401.

Castro, C.D., and M.F. Flajnik. 2014. Putting J chain back on the map: how might its expression define plasma cell development? *J Immunol* 193:3248-3255.

Croker, B.A., D. Metcalf, L. Robb, W. Wei, S. Mifsud, L. DiRago, L.A. Cluse, K.D. Sutherland, L. Hartley, E. Williams, J.G. Zhang, D.J. Hilton, N.A. Nicola, W.S. Alexander, and A.W. Roberts. 2004. SOCS3 is a critical physiological negative regulator of G-CSF signaling and emergency granulopoiesis. *Immunity* 20:153-165.

Crouch, E.E., Z. Li, M. Takizawa, S. Fichtner-Feigl, P. Gourzi, C. Montano, L. Feigenbaum, P. Wilson, S. Janz, F.N. Papavasiliou, and R. Casellas. 2007. Regulation of AID expression in the immune response. *J Exp Med* 204:1145-1156.

de Boer, J., A. Williams, G. Skavdis, N. Harker, M. Coles, M. Tolaini, T. Norton, K. Williams, K. Roderick, A.J. Potocnik, and D. Kioussis. 2003. Transgenic mice with hematopoietic and lymphoid specific expression of Cre. *Eur J Immunol* 33:314-325.

Dogan, I., B. Bertocci, V. Vilmont, F. Delbos, J. Megret, S. Storck, C.A. Reynaud, and J.C. Weill. 2009. Multiple layers of B cell memory with different effector functions. *Nat Immunol* 10:1292-1299.

Duber, S., M. Hafner, M. Krey, S. Lienenklaus, B. Roy, E. Hobeika, M. Reth, T. Buch, A. Waisman, K. Kretschmer, and S. Weiss. 2009. Induction of B-cell development in adult mice reveals the ability of bone marrow to produce B-1a cells. *Blood* 114:4960-4967.

Feil, S., N. Valtcheva, and R. Feil. 2009. Inducible Cre mice. *Methods Mol Biol* 530:343-363.

Georgiades, P., S. Ogilvy, H. Duval, D.R. Licence, D.S. Charnock-Jones, S.K. Smith, and C.G. Print. 2002. VavCre transgenic mice: a tool for mutagenesis in hematopoietic and endothelial lineages. *Genesis* 34:251-256.

Hargreaves, D.C., P.L. Hyman, T.T. Lu, V.N. Ngo, A. Bidgol, G. Suzuki, Y.R. Zou, D.R. Littman, and J.G. Cyster. 2001. A coordinated change in chemokine responsiveness guides plasma cell movements. *J Exp Med* 194:45-56.

Heng, T.S., M.W. Painter, and C. Immunological Genome Project. 2008. The Immunological Genome Project: networks of gene expression in immune cells. *Nat Immunol* 9:1091-1094.

Ho, F., J.E. Lortan, I.C. MacLennan, and M. Khan. 1986. Distinct short-lived and long-lived antibody-producing cell populations. *Eur J Immunol* 16:1297-1301.

Hobeika, E., S. Thiemann, B. Storch, H. Jumaa, P.J. Nielsen, R. Pelanda, and M. Reth. 2006. Testing gene function early in the B cell lineage in mb1-cre mice. *Proc Natl Acad Sci U S A* 103:13789-13794.

Johansen, F.E., R. Braathen, and P. Brandtzaeg. 2000. Role of J chain in secretory immunoglobulin formation. *Scand J Immunol* 52:240-248.

John, S.A., and L.A. Garrett-Sinha. 2009. Blimp1: a conserved transcriptional repressor critical for differentiation of many tissues. *Exp Cell Res* 315:1077-1084.

Kallies, A., J. Hasbold, K. Fairfax, C. Pridans, D. Emslie, B.S. McKenzie, A.M. Lew, L.M. Corcoran, P.D. Hodgkin, D.M. Tarlinton, and S.L. Nutt. 2007. Initiation of plasma-cell differentiation is independent of the transcription factor Blimp-1. *Immunity* 26:555-566.

Klein, U., S. Casola, G. Cattoretti, Q. Shen, M. Lia, T. Mo, T. Ludwig, K. Rajewsky, and R. Dalla-Favera. 2006. Transcription factor IRF4 controls plasma cell differentiation and class-switch recombination. *Nat Immunol* 7:773-782.

Kraus, M., M.B. Alimzhanov, N. Rajewsky, and K. Rajewsky. 2004. Survival of resting mature B lymphocytes depends on BCR signaling via the Igalpha/beta heterodimer. *Cell* 117:787-800.

Kwon, K., C. Hutter, Q. Sun, I. Bilic, C. Cobaleda, S. Malin, and M. Busslinger. 2008. Instructive role of the transcription factor E2A in early B lymphopoiesis and germinal center B cell development. *Immunity* 28:751-762.

Lemke, A., M. Kraft, K. Roth, R. Riedel, D. Lammerding, and A.E. Hauser. 2016. Long-lived plasma cells are generated in mucosal immune responses and contribute to the bone marrow plasma cell pool in mice. *Mucosal Immunol* 9:83-97.

Luche, H., O. Weber, T. Nageswara Rao, C. Blum, and H.J. Fehling. 2007. Faithful activation of an extra-bright red fluorescent protein in "knock-in" Cre-reporter mice ideally suited for lineage tracing studies. *Eur J Immunol* 37:43-53.

Lycke, N., L. Erlandsson, L. Ekman, K. Schon, and T. Leanderson. 1999. Lack of J chain inhibits the transport of gut IgA and abrogates the development of intestinal antitoxic protection. *J Immunol* 163:913-919.

Manz, R.A., and A. Radbruch. 2002. Plasma cells for a lifetime? *Eur J Immunol* 32:923-927.

Manz, R.A., A. Thiel, and A. Radbruch. 1997. Lifetime of plasma cells in the bone marrow. *Nature* 388:133-134.

Mather, E.L., F.W. Alt, A.L. Bothwell, D. Baltimore, and M.E. Koshland. 1981. Expression of J chain RNA in cell lines representing different stages of B lymphocyte differentiation. *Cell* 23:369-378.

Max, E.E., and S.J. Korsmeyer. 1985. Human J chain gene. Structure and expression in B lymphoid cells. *J Exp Med* 161:832-849.

Meyers, E.N., M. Lewandoski, and G.R. Martin. 1998. An Fgf8 mutant allelic series generated by Cre- and Flp-mediated recombination. *Nat Genet* 18:136-141.

Moriyama, S., N. Takahashi, J.A. Green, S. Hori, M. Kubo, J.G. Cyster, and T. Okada. 2014. Sphingosine-1-phosphate receptor 2 is critical for follicular helper T cell retention in germinal centers. *J Exp Med* 211:1297-1305.

Nagy, A., C. Moens, E. Ivanyi, J. Pawling, M. Gertsenstein, A.K. Hadjantonakis, M. Pirity, and J. Rossant. 1998. Dissecting the role of N-myc in development using a single targeting vector to generate a series of alleles. *Curr Biol* 8:661-664.

Nutt, S.L., P.D. Hodgkin, D.M. Tarlinton, and L.M. Corcoran. 2015. The generation of antibody-secreting plasma cells. *Nat Rev Immunol* 15:160-171.

Ohinata, Y., B. Payer, D. O'Carroll, K. Ancelin, Y. Ono, M. Sano, S.C. Barton, T. Obukhanych, M. Nussenzweig, A. Tarakhovsky, M. Saitou, and M.A. Surani. 2005. Blimp1 is a critical determinant of the germ cell lineage in mice. *Nature* 436:207-213.

Palumbo, A., and K. Anderson. 2011. Multiple myeloma. *N Engl J Med* 364:1046-1060.

Pracht, K., J. Meinzinger, P. Daum, S.R. Schulz, D. Reimer, M. Hauke, E. Roth, D. Mielenz, C. Berek, J. Corte-Real, H.M. Jack, and W. Schuh. 2017. A new staining protocol for detection of murine antibody-secreting plasma cell subsets by flow cytometry. *Eur J Immunol* 47:1389-1392.

Rickert, R.C., J. Roes, and K. Rajewsky. 1997. B lymphocyte-specific, Cre-mediated mutagenesis in mice. *Nucleic Acids Res* 25:1317-1318.

Rinkenberger, J.L., J.J. Wallin, K.W. Johnson, and M.E. Koshland. 1996. An interleukin-2 signal relieves BSAP (Pax5)-mediated repression of the immunoglobulin J chain gene. *Immunity* 5:377-386.

Robbiani, D.F., A. Bothmer, E. Callen, B. Reina-San-Martin, Y. Dorsett, S. Difilippantonio, D.J.
Bolland, H.T. Chen, A.E. Corcoran, A. Nussenzweig, and M.C. Nussenzweig. 2008. AID is
required for the chromosomal breaks in c-myc that lead to c-myc/IgH translocations. *Cell*
135:1028-1038.
Schweighoffer, E., L. Vanes, J. Nys, D. Cantrell, S. McCleary, N. Smithers, and V.L. Tybulewicz. 2013.
The BAFF receptor transduces survival signals by co-opting the B cell receptor signaling
pathway. *Immunity* 38:475-488.
Shaffer, A.L., M. Shapiro-Shelef, N.N. Iwakoshi, A.H. Lee, S.B. Qian, H. Zhao, X. Yu, L. Yang, B.K.
Tan, A. Rosenwald, E.M. Hurt, E. Petroulakis, N. Sonenberg, J.W. Yewdell, K. Calame, L.H.
Glimcher, and L.M. Staudt. 2004. XBP1, downstream of Blimp-1, expands the secretory
apparatus and other organelles, and increases protein synthesis in plasma cell differentiation.
Immunity 21:81-93.
Shapiro-Shelef, M., K.I. Lin, L.J. McHeyzer-Williams, J. Liao, M.G. McHeyzer-Williams, and K.
Calame. 2003. Blimp-1 is required for the formation of immunoglobulin secreting plasma cells
and pre-plasma memory B cells. *Immunity* 19:607-620.
Shinnakasu, R., T. Inoue, K. Kometani, S. Moriyama, Y. Adachi, M. Nakayama, Y. Takahashi, H.
Fukuyama, T. Okada, and T. Kurosaki. 2016. Regulated selection of germinal-center cells into
the memory B cell compartment. *Nat Immunol* 17:861-869.
Sze, D.M., K.M. Toellner, C. Garcia de Vinuesa, D.R. Taylor, and I.C. MacLennan. 2000. Intrinsic
constraint on plasmablast growth and extrinsic limits of plasma cell survival. *J Exp Med*
192:813-821.
Szymczak, A.L., C.J. Workman, Y. Wang, K.M. Vignali, S. Dilioglou, E.F. Vanin, and D.A. Vignali.
2004. Correction of multi-gene deficiency in vivo using a single 'self-cleaving' 2A peptide-
based retroviral vector. *Nat Biotechnol* 22:589-594.
Weber, T., J. Seagal, W. Winkler, T. Wirtz, V.T. Chu, and K. Rajewsky. 2019. A novel allele for
inducible Cre expression in germinal center B cells. *Eur J Immunol* 49:192-194.
Yasuda, T., T. Wirtz, B. Zhang, T. Wunderlich, M. Schmidt-Supprian, T. Sommermann, and K.
Rajewsky. 2013. Studying Epstein-Barr virus pathologies and immune surveillance by
reconstructing EBV infection in mice. *Cold Spring Harb Symp Quant Biol* 78:259-263.

502
503
504

Figure legends

Figure 1. *Jchain* transcripts are highly enriched in plasma cell populations.

(A) Schematic of the network of factors associated with plasma cell differentiation. Upwards arrows indicate increased expression compared to the precursor population. **(B-D)** Differential gene expression analysis using RNA sequencing data (ImmGen, (Heng et al., 2008)). (B) Analysis of *Xbp1* and *Irf4*, which encode for XBP1 and IRF4, respectively. (C) Analysis of *Sdc1* and *Prdm1*, which encode for CD138 and BLIMP1, respectively. (D) Analysis for *Jchain* (*Igj*) that encodes for JCHAIN. "x" indicates fold change. Expression Value Normalized by DESeq2. http://rstats.immgen.org/Skyline/skyline.html.

Figure 2. *Jchain* is expressed in a small fraction of GC B cells and in most Plasma cells.

(A) Schematic of *Jchain* targeted allele Wtsi MGI:5633773. Rectangular boxes indicate exons, and exon number is on top; pink triangle indicates a FRT sequence; EGFPcre^{ERT2} cassette contains a splice acceptor site (SA)-led EGFP-2A-creERT2 expression cassette followed by a poly-A tail inserted in the intron between exons 1 and 2; white triangle indicates a ROX sequence; orange rectangle indicates a promoter driven puromycin resistance cassette; green triangle indicates loxP sequence. **(B)** Schematic of experimental procedure protocol. Mice carrying the *Igjcre$^{ERT2(GFP)}$* allele were immunized with sheep red blood cells (SRBC) intravenously (i.v.) on day 0 and spleens of mice were analyzed at day 12 post-immunization. **(C)** Gating strategy of populations by flow-cytometry according to the expression of GFP and B220 in mice carrying the *Igjcre$^{ERT2(GFP)}$* allele and wild-type B6 mice for negative control of GFP expression. **(D)** Gating strategy by flow-cytometry for plasma cells within the GFP^{hi}B220^{low} population using CD138$^+$ and CD138$^+$CXCR4$^+$ markers. **(E)** Cumulative data for CD138$^+$ and CD138$^+$CXCR4$^+$ plasma cells analyzed as in (D). Top: fraction of CD138$^+$ plasma cells; Bottom: fraction of CD138$^+$CXCR4$^+$ plasma cells within the four populations defined by flow-cytometry according to the expression of GFP and B220 in mice carrying the *Igjcre$^{ERT2(GFP)}$* allele. **(F)** Gating strategy by flow-cytometry for total CD138$^+$ and CD138$^+$CXCR4$^+$ plasma cells within GFP^{low}B220^{hi} population. The CD138^{neg} cell fraction within the GFP^{low}B220^{hi} population was analyzed for the CD19 B cell marker and stained for CD38 and FAS to determine germinal center (GC) B cells. **(G)** Cumulative

data for the frequency of GC B cells within the CD138^{neg}GFP^{low}B220^{hi} population. Each symbol (E, G)

represents an individual mouse; small horizontal lines indicate median, minimum and maximum values.

***=p≤0.001, ****=p≤0.0001 (unpaired Student's *t*-test). Data are representative of three independent

experiments (E, G).

Figure 3. *Jchain* precedes IRF4 and BLIMP1 expression.

(A) Schematic of experimental procedure protocol. Mice carrying the *Igjcre*$^{ERT2(GFP)}$ allele were

immunized with sheep red blood cells (SRBC) intravenously (i.v.) on day 0 and spleens of mice were

analyzed at day 12 post-immunization. **(B)** Gating strategy of populations by flow-cytometry according

to the expression of GFP and B220 in mice carrying the *Igjcre*$^{ERT2(GFP)}$ allele. Wild-type B6 mice for

negative control of GFP expression. **(C)** Gating strategy for IRF4 and BLIMP1 expression by flow-

cytometry within the GFP^{neg}B220^{neg} population. **(D)** Gating strategy for IRF4 and BLIMP1 expression

by flow-cytometry within GFP^{low}B220^{hi}, GFP^{int}B220^{int} and GFP^{hi}B220^{low} populations, defined as in (B).

(E) Cumulative data for the frequency of BLIMP1^{+}IRF4^{+} cells within the four populations defined as

in (B). Each symbol in (E) represents an individual mouse; small horizontal lines indicate median,

minimum and maximum values. ns=not significant, ***=p≤0.001, ****=p≤0.0001 (unpaired Student's

t-test). Data are representative of three independent experiments (E).

Figure 4. *Igjcre*$^{ERT2(GFP)}$ mediated cre recombination is effective only in plasma cells.

(A) Schematic of experimental procedure protocol. Mice carrying the *Igjcre*$^{ERT2(GFP)}$ and *R26*^{lslRFP} alleles

were immunized with sheep red blood cells (SRBC) intravenously (i.v.) on day 0 and spleens (top) and

bone marrow (bottom) of mice were analyzed at day 12 post-immunization. A group of mice received

tamoxifen treatment for four consecutive days from day 7 to day 10 after immunization. **(B)** Gating

strategy of populations by flow-cytometry in spleen (top) and bone marrow (bottom) according to the

expression of GFP and B220 in mice carrying the *Igjcre*$^{ERT2(GFP)}$ allele. Wild-type B6 mice for negative

control of GFP expression. **(C)** Gating strategy for GFP and RFP expression by flow-cytometry in the

four populations defined as in (B). Top: spleen; bottom: bone marrow. **(D)** Cumulative data for the

frequency and number of GFP$^+$RFP$^+$ cells within GFP^{neg}B220^{neg}, GFP^{low}B220^{hi}, GFP^{int}B220^{int} and

GFP^{hi}B220^{low} populations, defined as in (B). Top: spleen; bottom: bone marrow. Each symbol (D)

represents an individual mouse; small horizontal lines indicate median, minimum and maximum values.

ns=not significant, *=p≤0.05, **=p≤0.01, ***=p≤0.001, ****=p≤0.0001 (unpaired Student's *t*-test).

Data are representative of three independent experiments (D).

Figure 5. Genetic manipulation using *Igjcre*$^{ERT2(GFP)}$ occurs across immunoglobulin isotypes.

(A) Schematic of experimental procedure protocol. Mice carrying the *Igjcre*$^{ERT2(GFP)}$ and *R26*^{lslRFP} alleles

were immunized with sheep red blood cells (SRBC) intravenously (i.v.) on day 0 and spleens,

mesenteric lymph nodes (mLN), peyer's patches, and bone marrows of mice were analyzed at day 12

post-immunization. Mice received tamoxifen treatment for four consecutive days from day 7 to day 10

after immunization. **(B)** Gating strategy by flow-cytometry for intracellular and extracellular expression

of IgG1, IgM and IgA within total B220^{low}CD138$^+$ plasma cells. Analysis in the spleen is provided as

example. **(C)** Gating strategy by flow-cytometry for intracellular and extracellular expression of IgG1,

IgM and IgA within B220^{low}CD138$^+$GFP$^+$ (RFP$^+$ and RFP^{neg}) plasma cells. Analysis in the spleen is

provided as example. **(D)** Gating strategy by flow-cytometry for intracellular and extracellular

expression of IgG1, IgM and IgA within B220^{low}CD138$^+$GFP$^+$RFP$^+$ plasma cells. Analysis in the spleen

is provided as example. **(E)** Cumulative data for the fractions of IgA, IgM or IgG1 expressing cells

within total plasma cells (PC) (black, B220^{low}CD138$^+$), GFP$^+$ (RFP$^+$ and RFP^{neg}) plasma cells (green,

B220^{low}CD138$^+$GFP$^+$), and GFP$^+$RFP$^+$ plasma cells (orange, B220^{low}CD138$^+$GFP$^+$RFP$^+$) at 15 weeks

of age. **(F)** Cumulative data for the fractions of IgA, IgM or IgG1 expressing cells within total plasma

cells (PC) (black, B220^{low}CD138$^+$), GFP$^+$ (RFP$^+$ and RFP^{neg}) plasma cells (green,

B220^{low}CD138$^+$GFP$^+$), and GFP$^+$RFP$^+$ plasma cells (orange, B220^{low}CD138$^+$GFP$^+$RFP$^+$) at 30 weeks

of age. Each symbol (E, F) represents an individual mouse; small horizontal lines indicate median,

minimum and maximum values. ns=not significant, *=p≤0.05 (2way ANOVA). Data are representative

of three independent experiments (E, F).

Figure 6. *Igjcre[ERT2(GFP)]* allows the study of plasma cell lifespan.

(A) Schematic of experimental procedure protocol. Mice carrying the *Igjcre[ERT2(GFP)]* and *R26[lslRFP]* alleles

were immunized twice with sheep red blood cells (SRBC) intravenously (i.v.) on day 0 and day 21.

Mice received tamoxifen treatment for five consecutive days from day 51 to 55 after the first

immunization. Spleens and bone marrow of mice were analyzed at the five-day, two-month and five-

month timepoints after the last tamoxifen administration. **(B)** Cumulative data for the fraction of

GFP^+RFP^+ cells (left) and GFP^+RFP^{neg} cells (right) within $CD138^+CXCR4^+$ plasma cells in the spleen.

(C) Cumulative data for the fraction of GFP^+RFP^+ cells (left) and GFP^+RFP^{neg} cells (right) within

$CD138^+CXCR4^+$ plasma cells in the bone marrow. **(D)** Graphical representation of R^2 and half-life of

$GFP^+RFP^+CD138^+CXCR4^+$ plasma cells in the spleen using the data presented in (B). Graphing of best-

fit curve was performed using the GraphPad Prism 8 software. **(E)** Graphical representation of R^2 and

half-life of $GFP^+RFP^+CD138^+CXCR4^+$ plasma cells in the bone marrow using the data presented in

(B). Graphing of best-fit curve was performed using the GraphPad Prism 8 software. Each symbol (B,

C) represents an individual mouse; small horizontal lines indicate median, minimum and maximum

values. ns=not significant, *=p≤0.05 **=p≤0.01, ***=p≤0.001, ****=p≤0.0001 (unpaired Student's *t*-

test). Data are representative of three independent experiments (B, C).

Supplemental Figure Legends

Figure S1. *Jchain* expression in multiple stages of B cell development and other cell lineages

(A) Gating strategy by flow-cytometry for the B cell lineage in the spleen. Mice carrying the

Igjcre[ERT2(GFP)] and *R26[lslRFP]* alleles were immunized with sheep red blood cells (SRBC) intravenously

(i.v.) at day 0 and spleens were analyzed at day 12. Mice did not receive tamoxifen treatment. Cells

were pre-gated on live singlets and for the expression of CD138 and CXCR4 markers. $CD138^+CXCR4^+$

cells were defined as plasma cells (PC). Non-PCs positive for the CD19 and B220 markers were defined

as B2 cells, whereas those cells displaying a $CD19^+B220^{low}$ were defined as B1 cells. B1 cells were

further characterized as B1a ($CD5^{hi}CD23^{neg}$) and B1b ($CD5^{low}CD23^{neg}$) cells. B2 cells were further

delineated into marginal zone (MZ) B cells ($CD21^{hi}CD23^{int}$), T1 B cells ($CD21^{low}CD23^{low}$), T2 B cells

($CD21^{int}IgM^{hi}CD23^{hi}$) and Follicular B cells (Fo, $CD21^{int}IgM^{int}CD23^{hi}$. Non MZ or T1 cells were gated

for germinal center (GC) B cells (CD38^{low}FAS^{high}) and these were further delineated into dark zone GC

B cells (CXCR4^{high}CD86^{low}) and light zone GC B cells (CXCR4^{low}CD86^{high}). (Bonami et al., 2014) **(B)**

Cumulative data for fractions of GFP$^+$RFP^{neg} and GFP$^+$RFP$^+$ cells within the B cell lineage in the spleen.

Analyzed as in (A). **(C)** Cumulative data for fractions of GFP$^+$RFP^{neg} cells and GFP$^+$RFP$^+$ cells in

additional B cell lineage and non-B cell lineage populations in the spleen. Analyzed as in (A). **(D)**

Gating strategy by flow-cytometry for the B cell lineage in the bone marrow. Mice carrying the

Igjcre$^{ERT2(GFP)}$ and *R26^{lslRFP}* alleles were immunized with sheep red blood cells (SRBC) intravenously

(i.v.) at day 0 and spleens were analyzed at day 12. Mice did not receive tamoxifen treatment. Cells

were pre-gated on live singlets and gated for CD138 and B220 expression. CD138$^+$B220^{low} cells were

defined as plasma cells (PCs) Non-PCs positive for the CD19 and B220 markers were further examined

and further delineated to immature B cells (IgM$^+$CD43^{neg}), Pre-B cells (fraction D; IgM^{neg}CD43^{neg}) and

Pro-B cells (IgM^{neg}CD43$^+$). Pro B cells were further sub-divided into Hardy fraction A (BP1^{neg}CD24^{neg}),

fraction B (BP1^{neg}CD24$^+$), fraction C (BP1$^+$CD24^{neg}) and fraction C' (BP1$^+$CD24$^+$). **(E)** Cumulative

data for fractions of GFP$^+$RFP^{neg} cells and GFP$^+$RFP$^+$ cells in the B cell lineage and in plasma cells in

the bone marrow. Analyzed as in (D). Each symbol (B, C, E) represents an individual mouse; small

horizontal lines indicate median, minimum and maximum values. Data are representative of three

independent experiments (B, C, E).

Figure S2. *Igjcre$^{ERT2(GFP)}$* mediated recombination in vitro.

(**A**) Gating strategy by flow-cytometry for GFP and RFP expression within CD19$^+$B220$^+$ B cells and

B220^{low}CD138$^+$ plasmablasts. Splenic B cells from unmanipulated mice carrying the *Igjcre$^{ERT2(GFP)}$* and

R26^{lslRFP} alleles were enriched using CD43 negative depletion (MACS) and culture *in vitro* in the

presence of LPS and 4-hydroxytamoxifen (4-OH-TAM). Cells were analyzed 48h, 72h, and 96h time-

points for GFP and RFP expression by flow-cytometry. Data from 96h analysis of a B cell culture with

LPS and 1μM 4-OH-TAM is shown as example. (**B**) Cumulative data for fractions of GFP^{neg}RFP^{neg},

GFP$^+$RFP^{neg}, GFP$^+$RFP$^+$ cells within B cells. Analyzed as in (A). (**C**) Cumulative data for fractions of

GFP^{neg}RFP^{neg}, GFP$^+$RFP^{neg}, GFP$^+$RFP$^+$ cells within plasmablasts. Analyzed as in (A).

Materials and Methods

Mice: The *Igjcre^ERT2(GFP)* allele was purchased from EMMA in agreement with the Wellcome Trust Sanger Institute (Jchain targeted allele Wtsi MGI:5633773, genebank https://www.i-dcc.org/imits/targ_rep/alleles/43805/escell-clone-genbank-file) and mice were rederived at the The Francis Crick Institute. The allele contains a splice acceptor site (SA), an *EGFP-2A-creERT2* expression cassette and a poly-A tail in the intron between exons 1 and 2 under the *Jchain* promoter. In addition, exon 2 is loxP-flanked and the allele also contains a Rox-flanked puromycin resistance cassette. These mice were crossed to carry a *Rosa26^lsl RFP* cre recombination reporter allele (R26^lsl.RFP) allele that expresses a non-toxic tandem-dimer red fluorescent protein upon cre-mediated deletion of a floxed STOP cassette (Luche et al., 2007). Mice were maintained on the C57BL/6 background and bred at The Francis Crick Institute biological resources facility under specific pathogen-free conditions. Animal experiments were carried out in accordance with national and institutional guidelines for animal care and were approved by The Francis Crick Institute biological resources facility strategic oversight committee (incorporating the Animal Welfare and Ethical Review Body) and by the Home Office, UK. All animal care and procedures followed guidelines of the UK Home Office according to the Animals (Scientific Procedures) Act 1986 and were approved by Biological Research Facility at the Francis Crick Institute. The age of mice ranged between 15-30 weeks as specified.

Immunization and *in vivo* induction of cre activity: Mice were injected intravenously with 1 x 10^9 sheep red blood cells (SRBCs, TCS Biosciences Ltd) in PBS. For the induction of cre activity, 4mg tamoxifen (SIGMA T5648) dissolved in sunflower seed oil were administered by oral gavage to mice once per day for multiple days depending on experimental design.

***In vitro* B cell culture and induction cre activity**: Splenic cells were harvested and B cells were isolated using CD43 (Ly-48) MicroBeads, mouse (Miltenyi Biotec). The purity of B cells was determined by flow-cytometry (>95%). B cells were cultured in 96-well round-bottom plates (Falcon) in B cell media in B cell media (DMEM high glucose/Glutamax from ThermoFisher Scientific

supplemented with 10% fetal bovine serum F7524 from Sigma, 100 U/mL Penicillin and 100 µg/mL

Streptomycin from Life Technologies, 10 mM HEPES Buffer Solution from Life Technologies, 100

µM MEM Non-essential Amino Acids from ThermoFisher Scientific, 1 mM sodium pyruvate from Life

Technologies and 50 uM β-mercaptoethanol from Sigma) at 1 million/mL concentration (200,000

cells/well) with 10ug/mL LPS and varied concentrations of (Z) 4-hydroxytamoxifen (Sigma-Aldrich).

Analysis was performed by flow-cytometry at 48, 72 or 96 hours of culture.

Antibodies: Rat Anti-Mouse Blimp-1, Clone 6D3 (BD Biosciences); Rat Anti-Mouse CD16/CD32,

Clone 2.4G2 (BD Biosciences); Rat Anti-Mouse CD19, Clone 1D3 (BD Biosciences); Rat Anti-Mouse

CD23, Clone B3B4 (BD Biosciences); Rat Anti-Mouse CD38, Clone 90 (BD Biosciences); Hamster

Anti-Mouse CD95, Clone Jo2 (BD Biosciences); Rat Anti-Mouse IgG1, Clone A85-1 (BD

Biosciences); Rat Anti-Mouse CD138, Clone 281-2 (BD Biosciences); Rat Anti-Mouse CD45R/B220,

Clone RA3-6B2 (BioLegend); Hamster Anti-Mouse CD11c, Clone N418 (BioLegend); Rat Anti-

Mouse CD19, Clone 6D5 (BioLegend); Rat Anti-Mouse CD21/35, Clone 7E9 (BioLegend); Rat Anti-

Mouse CD43, Clone 1B11 (BioLegend); Rat Anti-Mouse CD86, Clone GL1 (BioLegend); Rat Anti-

Mouse BP1, Clone 6C3 (eBioscience); Rat Anti-Mouse CD5, Clone 53-7.3 (eBioscience); Rat Anti-

Mouse CXCR4, Clone 2B11 (eBioscience); Rat Anti-Mouse IgA, Clone 11-44-2 (eBioscience); Rat

Anti-Mouse IgM, Clone II/41 (eBioscience); Rat Anti-mouse IRF4, Clone 3E4 (eBioscience).

Flow cytometry: Single cell suspensions were stained with antibodies. We used Zombie NIR Fixable

Viability Kit (BioLegend) for live/dead discrimination. For intracellular staining we fixed cells using

the BD CytoFix/Cytoperm (BD Biosciences) kit as per manufacturer instructions. Samples were

acquired on a BD LSRFortessa analyser using FACSDiva software (BD) and analysed on FlowJo

software.

Quantification and statistical analysis: Data were analyzed with unpaired two-tailed Student's t test

or 2way ANOVA; a p-value of 0.05 or less was considered significant. Prism (v7 and v8, GraphPad)

was used for statistical analysis. A single asterisk (*) in the graphs of figures represents a p-value ≤0.05,

double asterisks (∗∗) a p-value ≤0.01, triple asterisks (∗∗∗) a p-value ≤0.001, quadruple asterisks (∗∗∗∗) a

p-value ≤0.0001, and "ns" stands for not statistically significant, i.e., a p-value >0.05.

Online supplemental material: Figure S1 displays the analysis of *Jchain* expression across cell

lineages in the spleen and bone marrow. These data also demonstrates that *Igjcre*$^{ERT2(GFP)}$ cre-mediated

recombination does not occur in the absence of tamoxifen treatment. Figure S1 shows experiments

using a classical LPS induced B to plasmablast differentiation and the occurrence of *Igjcre*$^{ERT2(GFP)}$ cre-

mediated recombination upon addition of increasing concentrations of 4-hydroxytamoxifen (4-OH-

TAM).

B cell lineage

Xu A.Q. et al.

Graphical Abstract

Figure 1:

Jchain transcripts are highly enriched in plasma cell populations

Xu A.Q. et al.

Figure 2.

Jchain is expressed in a small fraction of GC B cells and in most plasma cells

Xu A.Q. et al.

Figure 3:

Jchain expression correlates with IRF4 and BLIMP1 co-expression

Xu A.Q. et al.

Figure 4.

Igjcre^ERT2(GFP) mediated cre recombination is effective only in plasma cells

Xu A.Q. et al.

Figure 5.

Genetic manipulation using *Igjcre^ERT2(GFP)* occurs across immunoglobulin isotypes

Xu A.Q. et al.

Figure 6.

Igjcre^ERT2(GFP) allows lifespan studies and to trace long-lived plasma cells

Xu A.Q. et al.

Figure S1.

Jchain expression in multiple stages of B cell development and other cell lineages

Xu A.Q. et al.

Figure S2.

Igjcre[ERT2(GFP)] mediated recombination *in vitro*

Xu A.Q. et al.