Bioinformatic characterization of angiotensin-converting enzyme 2, the entry receptor for SARS-

CoV-2

Harlan Barker and Seppo Parkkila#

Faculty of Medicine and Health Technology, Tampere University and Fimlab Ltd, Tampere

University Hospital, 33520 Tampere, Finland

Running title: Bioinformatic characterization of angiotensin-converting enzyme 2

#Correspondence to:

Professor Seppo Parkkila, M.D., Ph.D.

Faculty of Medicine and Health Technology

Tampere University

Arvo Ylpön katu 34, 33520 Tampere, Finland

e-mail: seppo.parkkila@tuni.fi

ABSTRACT

The World Health Organization declared the COVID-19 epidemic a public health emergency of

international concern on March 11th, 2020, and the pandemic is rapidly spreading worldwide.

COVID-19 is caused by a novel coronavirus SARS-CoV-2, which enters human target cells via

angiotensin converting enzyme 2 (ACE2). We used a number of bioinformatics tools to

computationally characterize ACE2 by determining its cell-specific expression in trachea, lung, and

small intestine, derive its putative functions, and predict transcriptional regulation. The small

intestine expressed higher levels of ACE2 than any other organ. The large intestine, kidney and

testis showed moderate signals, whereas the signal was weak in the lung. Single cell RNA-Seq data

from trachea indicated positive signals along the respiratory tract in key protective cell types

including club, goblet, proliferating, and ciliary epithelial cells; while in lung the ratio of ACE2-

expressing cells was low in all cell types ($<2.6\%$), but was highest in vascular endothelial and

goblet cells. Gene ontology analysis suggested that, besides its classical role in renin-angiotensin

system, ACE2 may be functionally associated with angiogenesis/blood vessel morphogenesis.

Using a novel tool for the prediction of transcription factor binding sites we identified several

putative binding sites within two tissue-specific promoters of the *ACE2* gene. Our results also

confirmed that age and gender play no significant role in the regulation of ACE2 mRNA expression

in the lung.

IMPORTANCE

Vaccines and new medicines are urgently needed to prevent spread of COVID-19 pandemic, reduce

the symptoms, shorten the duration of disease, prevent virus spread in the body, and most

importantly to save lives. One of the key drug targets could be angiotensin-converting enzyme 2

(ACE2), which is a crucial receptor for the corona virus (SARS-CoV-2). It is known that SARS

coronavirus infections lead to worse outcome in the elderly and in males. Therefore, one aim of the

present study was to investigate whether age or sex could contribute to the regulation of ACE2

expression. We also decided to explore the transcriptional regulation of *ACE2* gene expression.

Since data on ACE2 distribution is still conflicting, we aimed to get a more comprehensive view of

the cell types expressing the receptor of SARS-CoV-2. Finally, we studied the coexpression of

ACE2 with other genes and explored its putative functions using gene ontology enrichment

analysis.

## INTRODUCTION

A zinc metalloenzyme, angiotensin-converting enzyme (ACE) was discovered 64 years ago and first named as a hypertension-converting enzyme [1]. Classically, ACE is well known for its roles in the regulation of arterial pressure through conversion of angiotensin I to active angiotensin II and cleavage of bradykinin and neurotensin [2]. As a zinc metalloenzyme, ACE belongs to a large cluster of zinc-binding proteins. The first zinc metalloenzyme, carbonic anhydrase was discovered in 1932 by Meldrum and Roughton [3] and thereafter thousands of such metalloenzymes have been reported in different species of all phyla [4, 5].

Angiotensin-converting enzyme 2 (ACE2) was first discovered in 2000 when a novel homologue of ACE was cloned [2, 6, 7]. Although ACE and ACE2 share significant sequence similarity in their catalytic domains, they appear to act on different peptide substrates of angiotensins [8, 9]. Previous studies identified ACE2 as a functional receptor for severe acute respiratory syndrome corona virus 1 (SARS-CoV-1) which led to an outbreak of SARS infection in 2003 [10]. ACE2 is also a crucial receptor for the novel corona virus (SARS-CoV-2), which has caused a large global outbreak of COVID-19 infection with rapidly growing numbers of patients (5,488,825 confirmed cases as of May 27th, 2020, https://www.who.int/emergencies/diseases/novel-coronavirus-2019). A recent report suggested that soluble ACE2 fused to the Fc portion of immunoglobulin can neutralize SARS-CoV-2 *in vitro* [11]. This result was further confirmed by showing that human recombinant soluble ACE2 reduced SARS-CoV-2 infection on cultured Vero-E6 cells in a dose dependent manner [12]. Therefore, ACE2 also holds promise for treating patients with coronavirus infection.

The structural key for target cell infection by coronavirus is the viral spike (S) protein of SARS-CoV. ACE2 acts as a locking device for the virus, whereby the binding of the surface unit S1 facilitates viral attachment to the surface of target cells [13]. The cellular serine protease (TMPRSS2) promotes SARS-CoV entry via a dual mechanism. It cleaves both the SARS-CoV S protein and the virus receptor, ACE2, promoting both the viral uptake and the viral and cellular

membrane fusion events [13-15]. The critical residues contributing to the receptor-spike protein

interaction were first determined for SARS-CoV-1 [16] and recently in three independent studies

for SARS-CoV-2 [17-19]. It has been proposed by biolayer interferometry studies that the receptor-

binding domains of SARS-CoV-1 and SARS-CoV-2 S proteins bind with comparable affinities to

human ACE2 [20]. In contrast, a modelling study suggested that binding of SARS-CoV-2 is

stronger [21], which was convincingly confirmed by structural and biochemical data [17, 18].

The clinical characteristics of COVID-19 infection have recently been described based on data from

1,099 patients from mainland China [22]. It was found that the clinical characteristics of COVID-19

mimic those of SARS-CoV-1 infection. The most dominant symptoms include fever, cough,

fatigue, and sputum production, whereas gastrointestinal symptoms are less common. In laboratory

parameters, lymphopenia was detected in 83.2% of patients on admission. According to another

recent survey of 278 patients with pneumonia caused by SARS-CoV-2, fever was the most common

symptom, followed by cough [23]. Bilateral pneumonia has been detected by computed tomography

scans in 67.0% of patients [24]. A recent study from Wuhan, China listed the most common clinical

complications determined in critically ill COVID-19 patients [25]. The complications during

clinical worsening included acute respiratory distress syndrome and respiratory failure, sepsis, acute

cardiac injury, and heart failure.

Data on the localization of virus receptors can provide insight into mechanisms of virus entry, tissue

tropism, and pathogenesis of the disease. Therefore, it is of particular interest to correlate COVID-

19 symptoms with the distribution pattern of ACE2. The first studies performed by northern

blotting indicated that ACE2 is located in the human heart, kidney, and testis [2]. By

immunohistochemistry, the expression of the ACE2 protein was identified in the human lung

alveolar epithelial cells (type I and II pneumocytes), enterocytes of the small intestine, the brush

border of the renal proximal tubules, and the endothelial cells of arteries and veins and arterial

smooth muscle cells in several organs [26]. It was proposed that this distribution pattern of ACE2

could explain the tissue tropism of SARS-CoV-1 for the lung, small intestine, and kidney [27]. On

the other hand, the symptoms of COVID-19, in contrast to SARS-CoV-1 infection, are not

associated to the same extent with the gastrointestinal tract in spite of the high expression of ACE2

in the intestinal enterocytes [28]. In COVID-19, diarrhea has been reported in just 3.8% of patients,

in contrast to 40-70% in SARS-CoV-1 infection [22, 29]. A recent preprint report indicated diarrhea

in 18.1% of 254 COVID-19 patients [30].

There are conflicting reports on the expression of ACE2 in the upper respiratory tract [29].

Hamming and coworkers found that only the basal layer of nonkeratinized airway squamous

epithelium shows positive signal [26], whereas Sims and colleagues demonstrated ACE2 expression

on the luminal surface of ciliated cells in freshly excised human nasal and tracheobronchial tissue

[31]. Ren and coworkers showed weak ACE2-positive signal in the epithelial cells of trachea and

main bronchus [32]. Although lymphopenia is a typical feature of SARS [22, 29], ACE2 is not

highly expressed on T or B cells or macrophages in the spleen or lymphoid organs [26].

It is known that both SARS-CoV and SARS-CoV-2 infections lead to worse outcome in the elderly

[29, 33]. Therefore, one aim of the present study was to investigate whether age could contribute to

the regulation of ACE2 expression. We also decided to explore the transcriptional regulation of

*ACE2* gene expression using a novel computational tool recently developed by the first author of

this article. Notably, data on ACE2 distribution is still conflicting, and thus we aimed to get a more

comprehensive view of the cell types expressing the receptor of SARS-CoV-2. Finally, we studied

the coexpression of ACE2 with other genes and explored its putative functions using a gene

ontology enrichment analysis.

RESULTS

ACE2 is weakly expressed in the lung

The first aim of our study was to investigate different human tissues using publicly available

datasets for the distribution of ACE2 mRNA and protein. In the FANTOM5 dataset, the highest

values for ACE2 mRNA, ranked according to signal intensity, were seen for the small intestine,

dura mater, colon, testis, thalamus, and rectum (Fig. 1).

Figure 2 shows the expression of ACE2 protein in selected human tissues. Representative example

images of the ACE2 immunostaining were prepared from tissue specimens of the Human Protein

Atlas database (https://www.proteinatlas.org/). The results indicate a strong signal for ACE2 protein

in the brush border of small intestinal enterocytes. In the kidney, prominent immunostaining

reactions were present in the epithelial cells of proximal convoluted tubules and Bowman´s capsule.

The seminiferous tubules and interstitial cells of testis also demonstrated strong immunostaining.

No immunoreactions for ACE2 were observed in the lung specimens. Very weak signal, associated

with apical membranes, was detected in sporadic ciliary cells of a nasopharyngeal mucosa sample.

Although the evaluation of immunostaining reaction is generally considered semiquantitative at

most, the results seem to correlate fairly well with the corresponding mRNA expression levels.

Single cell RNA-Seq analysis indicates cell-specific expression for ACE2 mRNA

The respiratory tract is the main target region that is affected by COVID-19 infection. Bulk RNA-

Seq data from lung specimens showed low expression levels for ACE2 (Fig. 1). Therefore, we

performed an analysis of single cell RNA-Seq using both human lung and mouse trachea datasets,

representing the breadth of the lower respiratory tract. Figures 3 and 4 show the expression of

ACE2 mRNA in identified cell types of lung and trachea, respectively. In lung, ACE-2 expressing

cells are generally uncommon with no cell type having a ratio of ACE2-expressing cells greater

than 2.6%. The cell types with the greatest proportion of ACE2 expression are those of arterial

vascular endothelial cells (2.55%), goblet cells (2.02%), and venous vascular endothelial cells

(1.33%). In trachea, the highest ratio of ACE2-expressing cells included the club cells (16.62%),

goblet cells (13.84%), and ciliary epithelial cells (6.63%).

Since both the airways and intestine contain goblet cells, SARS-CoV-1 affects gastrointestinal tract,

and bulk RNA-Seq data shows high expression in small intestine and colon, we decided to analyze

another single cell RNA-Seq dataset covering mouse intestinal epithelial cells. Figure 5 indicates

the highest levels of ACE2 mRNA signal in the absorptive enterocytes (44.09%), whereas the

intestinal goblet cells (1.35%) remain mostly negative.

ACE2 mRNA expression levels are unrelated to age and gender in the lung

Since both age and gender may contribute to onset and severity of COVID-19 symptoms we aimed

to investigate the effect of these variables on the expression levels of ACE2 mRNA. Figure 6

indicates that some tissues showed a slight trend to lower expression in older age categories.

Among all tested tissues, statistically significant differences between the age categories were seen

in the tibial nerve ($p=8.58 \times 10^{-6}$), minor salivary gland (p=0.002), aorta (p=0.003), whole blood

(p=0.005), transverse colon (p=0.010), hypothalamus (p=0.039), and sun exposed skin (p=0.046).

Importantly, the lung specimens showed no significant difference of ACE2 mRNA expression

between different age categories (p=0.681). Complete data on ACE mRNA expression levels in

different age categories are shown in Supplementary Table 1. To make a binary comparison of

expression by age, samples were divided into groups of ≤45 and >45 years of age. In comparison of

these younger and older age groups, significant differences in expression were found in tibial nerve

($p=2.47 \times 10^{-7}$), whole blood ($p=3.21 \times 10^{-4}$), minor salivary gland ($p=4.89 \times 10^{-4}$), sun exposed

skin (p=0.003), transverse colon (p=0.022), testis (p=0.025), esophageal muscle layer (p=0.040),

and subcutaneous adipose tissue (p=0.045).

ACE2 mRNA levels largely overlapped between male and female sexes as shown in Figure 7. In

the lung, no statistically significant difference was observed in the expression levels between the

male and female subjects (p=0.908). Statistically significant differences were observed in the

adipose tissue (p=0.0001), whole blood (p=0.0002), amygdala (p=0.0006), transverse colon

(p=0.0008), muscle layer of esophagus (p=0.002), left ventricle of heart (p=0.005), Epstein-Barr

virus-transformed lymphocytes (p=0.015), and esophagus-gastroesophageal junction (p=0.024).

Notably, there was no clear sex-specific trend pointing to one direction in all these cases. ACE

mRNA expression levels in all studied tissues sorted according to subjects´ gender are shown in

Supplementary Table 2.

Proximal promoter contains putative TFBSs for ileum, colon, and kidney expression

TFBS analysis of the *ACE2* intestinal transcript promoter (ENST00000252519) revealed several

candidate binding sites which occur in a cluster extending from 400 bp upstream of the transcription

start site; CDX2, HNF1A, FOXA1, SOX4, TP63, HNF4A, DUX4, FOXA2, NR2F6, and SOX11

(Fig. 7A). These predicted sites overlap an evolutionarily conserved region in mammals and are

proximal to several ATAC-Seq peaks. In several tissues these TFs are found to be highly positively

correlated (>0.7) with expression of ACE2: CDX2 (colon, terminal ileum), HNF1A (colon, kidney,

terminal ileum), FOXA1 (cervix, colon, terminal ileum), HNF4A (colon, terminal ileum), FOXA2

(colon, kidney), NR2F6 (colon, kidney, terminal ileum), and SOX11 (kidney). In addition, two of

the TFs are highly negatively correlated with ACE2 expression DUX4 (kidney) and FOXA1

(kidney). Full prediction results are included as Supplementary Table 3 and TF correlations by

tissue are present in Supplementary Table 4.

Analysis of the *ACE2* lung transcript promoter (ENST00000427411) produced putative TFBS

predictions for ESRRA, HNF4A, CDX2, CEBPA, ESRRB, MEF2B, TCF7, TCF7L2, JUN, and

LEF1 (Fig. 7B). The predicted TFBSs clustered within 200 base pairs of the TSS, and overlap with

evolutionarily conserved regions, TFBS metaclusters, and ATAC-Seq peaks. The TFs

corresponding to predicted TFBSs, which are positively correlated (>0.7) with ACE2 expression,

are ESRRA (terminal ileum, colon), HNF4A (terminal ileum, colon), CDX2 (colon, terminal

ileum), CEBPA (colon, terminal ileum), ESRRB (cervix), TCF7L2 (testis). Those TFBSs with TFs

which strongly (<-0.7) negatively correlate with ACE2 are ESRRA (kidney) and TCFL72 (kidney).

The lung-specific transcript TSS aligns with the p3@ACE2 FANTOM5 dataset CAGE peak, which

indicates that the expression of this transcript is much lower than the intestinal transcript, which

corresponds with p1@ACE2 and p2@ACE2 FANTOM5 CAGE peaks. Common between the two

tissue-specific transcripts, are predictions for CDX2 and HNF-family transcription factors.

ACE2 mRNA expression correlates with metalloproteases and transporter genes

Coexpression analysis identified numerous genes in ileum, testis, colon, and kidney which are

highly correlated (>0.8) with ACE2 (Table 1; Supplementary Table 5). In particular, in the ileum

there are a number of genes with correlation values greater than 0.95. In contrast, analysis of the

lung shows a maximum correlation of expression of 0.6275. The genes with which ACE2 mRNA

expression shows the highest levels of coexpression code for metalloprotease and transporter

proteins. Selected tissue-specific ACE2-correlated genes, determined with bulk RNA-Seq data, are

presented with their expression levels within the scRNA-Seq trachea and intestinal datasets in

Figure 9.

ACE2 is associated with vascular growth

GO enrichment analysis of ACE2 mRNA expression in all tissues produced 22 terms which were

enriched in BP, CC, HP, KEGG, and WP ontologies (Table 2). A total of 12 of these terms were

related to blood vessel growth, including the three most strongly enriched terms, 'angiogenesis

[GO:0001525]', 'blood vessel morphogenesis [GO:0048514]', and 'vasculature development

[GO:0072358]'. Full GO enrichment results for relevant tissues (lung, small intestine, kidney,

colon, and testis) are included as Supplementary Table 6.

DISCUSSION

The predominant pathological features of COVID-19 infection largely mimic those previously

reported for SARS-CoV-1 infection. They include dry cough, persistent fever, progressive dyspnea,

and in some cases acute exacerbation of lung function with bilateral pneumonia [31]. Major lung

lesions include several pathological signs, such as diffuse alveolar damage, inflammatory exudation

in the alveoli and interstitial tissue, hyperplasia of fibrous tissue, and eventually lung fibrosis [34-

36]. It has been shown by fluorescence *in situ* hybridization technique that SARS-CoV-1 RNA

locates to the alveolar pneumocytes and alveolar space [37, 38]. Mossel and colleagues

demonstrated that SARS-CoV-1 replicates in type 2 (AT2) pneumocytes, but not in type 1 (AT1)

cells [39]. Considering all of these facts, it is not surprising that most histopathological analyses

have been focused on distal parts of the respiratory airways, while the regions other than the

alveolus have been less systematically studied.

To understand better the pathogenesis of COVID-19 we need to know where ACE2, the receptor for

SARS-CoV, is located within the human respiratory tract and elsewhere. Overall, different studies

including ours have convincingly shown that several organs, such as the small intestine, colon,

kidney, and testis, express higher levels of ACE2 than the lung and other parts of the respiratory

tract. Our analysis of ACE2 expression in the human lung show low levels of expression in all cell

types, with arterial vascular endothelial cells achieving the highest overall ratio of just ~2.5%. The

present results based on mouse tracheal dataset suggested that ACE2 mRNA is predominantly

expressed in the club cells, goblet cells, and ciliated epithelial cells, and at significantly higher

frequency than found in the lung. The mouse dataset used in our study contained no secretory3

cells, which Lukassen and colleagues recently reported to express the highest levels of ACE2

mRNA along the human respiratory tract [40]. Another study reported positive expression in the

type AT2 pneumocytes [41], which is in line with the results of Lukassen et al. [40], but only a few

cells appeared positive. A third study based on single cell expression data demonstrated the

strongest positive signal in the lung AT2 cells, while other cells including AT1 cells, club cells,

ciliated cells, and macrophages showed weaker expression [42]. A fourth single cell expression

analysis using Gene Expression Omnibus (GEO) database recently demonstrated ACE2-positive

signal in 1% of AT2 cells and in 2% of respiratory tract epithelial cells [43]. This correlates with

our own findings for lung. For comparison about 30% of ileal epithelial cells were ACE2-positive,

and 44% of enterocytes in small intestine of mouse. Immunohistochemical analysis of mouse

tissues has shown positive signal in the club cells, AT2 cells, endothelial cells, and smooth muscle

cells [44]. In spite of the obvious discrepancies between different datasets, that highlights the need

for large numbers of thoroughly characterized cells for single cell RNA-Seq analyses, we can now

make some conclusions of the expression of ACE2 mRNA in the respiratory tract. First, ACE2 is

positively though weakly expressed in the AT2 cells of the lung and less so in AT1 cells. Second,

ACE2 also shows a weak positive signal, but at significantly higher proportions of cells, in several

other cell types of the trachea, including goblet cells, club cells, and ciliated cells. Third, based on

the findings of Lukassen et al. [40] secretory3 cells, a transient cell type of the bronchial tree, may

express the highest levels of ACE2. These ACE2-positive cell types may represent the main host

cells for SARS-CoV-2 along the whole respiratory tract. However, the median percentage of ACE-

expressing secretory3 cells in the study was less than 6%, significantly less than that observed in

club (16.62%), goblet (13.84%), and ciliated (6.63%) cells of trachea we have identified in the

GSE103354 dataset.

Goblet cells, ciliated epithelial cells, and club cells are considered important cell types for the

protection of airway mucosa. Lukassen and coworkers [40] described secretory3 cells as

intermediate cells between goblet, ciliated, and club cells. If SARS-coronaviruses predominantly

attack these cells, locating along the airway segments including the trachea, bronchi, and

bronchioles until the last segment that is the respiratory bronchioles, it would be obvious that

physiological protective mechanisms are severely affected. Defective mucosal protection and

inefficient removal of pathogens due to viral infection may contribute to onset of severe bilateral pneumonia that is common for SARS-diseases [45]. This pathogenic mechanism is supported by previous findings, showing that early disease is manifested as a bronchiolar disease with respiratory epithelial cell necrosis, loss of cilia, squamous cell metaplasia, and intrabronchiolar fibrin deposits [31]. In fact, it has been suggested that early diffuse damage as a result of SARS-CoV-1 infection may actually initiate at the level of the respiratory bronchioles [46, 47].

Our findings confirm that the respiratory tract tissues have quite limited expression levels of ACE2 compared to several other tissues that show much more prominent signal. Because ACE2 is highly expressed in the intestine [28], as also confirmed by our bioinformatics study, it would be obvious to predict that both SARS-CoV-1 and -2 infections cause significant gastrointestinal pathology and symptoms including diarrhea. Interestingly, the patients with COVID-19 have reported less gastrointestinal symptoms than the SARS-CoV-1-infected patients [22, 29]. The pathophysiological basis for this phenomenon is not understood at this point, and thus further investigations on this topic are warranted.

When we initiated the present study, we hypothesized that understanding better the transcriptional regulation of the *ACE2* gene might help to explain the peculiar distribution pattern of ACE2 in tissues. Since upregulation of ACE2 would reflect an increased number of SARS-coronavirus receptors on cell surfaces, it could possibly help us to understand the mechanisms why certain patients (males more than females, old more than young, smokers more than non-smokers) are more susceptible for the most detrimental effects of the COVID-19 infection. In our study, the signals for ACE2 mRNA in the lung specimens did not vary much in different age groups nor did they show significant differences between males and females, which is in line with previous findings [40]. Therefore, different expression levels of lung ACE2 may not explain the variable outcome of the disease concerning age groups and genders.

To investigate the transcriptional regulation of *ACE2* gene we made predictions for the binding sites

of transcription factors within the proximal promoter region of the intestine-specific and lung-

specific human *ACE2* transcript promoters. Our findings introduced several putative binding sites in

the *ACE2* promoter for known transcription factors, which showed high levels of coexpression with

ACE2 in several tissues including the ileum, colon, and kidney. The identified transcription factors

could represent potential candidate target molecules which regulate ACE2 expression. Two of our

predictions, for HNF1A and HNF1B have been previously identified experimentally to drive ACE2

expression in pancreatic islet cells and insulinoma cells, respectively [48]. Later work by the same

group has shown that our prediction of FOXA binding sites in the ACE2 promoter are also likely

correct [49]. It is of interest that ACE2 might be regulated by oxygen status. Zhang and coworkers

previously demonstrated that ACE2 mRNA and protein levels increased during the early stages of

hypoxia and decreased to near-baseline levels at later stages after hypoxia inducible factor (HIF)-1α

accumulation [50]. Based on these findings *ACE2* has been listed as a HIF1α-target gene [51],

although it does not follow the typical HIF1α regulated expression pattern, nor is there any

predicted HIF1α binding site in our analyses. However, HNF1B has been identified as upregulated

in hypoxia in kidney, independent of HIF1α [52], and in hypoxic embryonic stem cells HIF1α has

been shown to increase expression of transcription factors TCF7/LEF1 (predicted to bind the

promoter of the lung-specific ACE2 transcript) through Wnt/β-catenin signaling [53].

The regulation of ACE2 expression remains an enigma and there may be multiple factors involved.

There has been concern that the use of ACE inhibitors and angiotensin receptor blockers could

increase the expression of ACE2 and increase patient susceptibility to viral host cell entry [54, 55].

Previous studies have suggested that both ACE inhibitor and angiotensin II receptor type I

antagonist therapies increase ACE2 mRNA expression in the rat heart [56]. There has also been

some evidence in humans showing increased expression of ACE2 in the heart, brain, and even in

urine after treatment with angiotensin receptor blockers [54]. Since these drugs are widely used for

treatment of hypertension and heart failure, it would be important to determine in COVID-19

patients whether these medications have any significant effects on symptoms or outcome of the

disease.

Gene ontology investigations revealed interesting novel data on potential physiological roles of

ACE2. The five most significant gene ontology terms included angiogenesis, blood vessel

morphogenesis, vasculature development, cardiovascular system development, and blood vessel

development. Our study of lung showed that the arterial and venous vascular endothelial cell types

had the highest ratios of ACE2-expressing cells, first and third highest, respectively. In another

study, ACE2 expression was previously detected in blood vessels [26], and a recent study showed

that SARS-CoV-2 is capable of directly infecting blood vessel cells [12]. Endothelial ACE2

expression may be linked to clotting and multi-organ dysfunction reported in many patients with

COVID-19. Based on the present finding angiogenesis/blood vessel morphogenesis may be

considered a putative function for ACE2 in addition to its classical role as the key angiotensin-(1-7)

forming enzyme [57].

Conclusions: Our bioinformatics study confirmed the low expression of ACE2 in the respiratory

tract. In lung it was lowest of all, while significantly higher in the trachea. Bulk RNA-Seq analyses

indicated the highest expression levels in the small intestine, colon, testis, and kidney. In the human

lung scRNA-Seq dataset, the strongest positive signals for ACE2 mRNA were observed in vascular

endothelial cells, goblet cells, ciliated cells, and AT2 and AT1 pneumocytes. In the mouse trachea

dataset, positive signals were most common in club cells, goblet cells and ciliated epithelial cells.

The results suggest that SARS-CoV infection may target the cell types that are important for the

protection of airway mucosa and their damage may lead to deterioration of epithelial cell function,

finally leading to a more severe lung disease with accumulation of alveolar exudate and

inflammatory cells and lung edema, the signs of pneumonia recently described in the lung

specimens of two patients with COVID-19 infection [58]. Gene ontology analysis based on

expression in all tissues suggested that ACE2 is involved in angiogenesis/blood vessel

morphogenesis processes in addition to its classical function in renin-angiotensin system.

METHODS

ACE2 mRNA expression

From the FANTOM5 project [59], cap analysis of gene expression (CAGE) sequencing of cDNA

has been performed in 1,839 human samples from 875 different primary cells, tissues, and cell

lines. Expression of transcription start sites (TSSs) was extracted and combined for all genes in all

samples as tags per million (TPM). From this compiled set, *ACE2* gene expression was extracted

and presented as barplot using the Matplotlib [60] and Seaborn [61] Python libraries. Similarly,

human gene expression data (as TPM) was extracted from the GTEx database, along with metadata

on the samples. *ACE2* gene expression values were separated by tissue and compared among 10-

year interval age groups to determine if the values showed any differences throughout the lifecycle.

Boxplots for tissues of relevance were generated using Matplotlib and Seaborn libraries.

Coexpression and gene ontology enrichment analysis

In each of the tissues present in the GTEx dataset, expression values for ACE2 were compared with

expression of all other genes by Spearman correlation analysis using the SciPy [62] Python library

to identify those genes with concordant expression patterns. Bonferroni correction was used to

derive an adjusted p-value threshold of 9.158E-07. For each tissue, those genes which both satisfied

the Bonferroni-adjusted p-value threshold and had a correlation of expression of 0.50 or greater

were analyzed using the Gprofiler gene ontology (GO) enrichment analysis [63] Python library to

identify possible enriched terms in biological process (BP), molecular function (MF), cellular

component (CC), human phenotype (HP), KEGG pathway, and WikiPathways (WP) ontologies.

ACE2 protein expression

Immunohistochemical localization of human ACE2 was evaluated from immunostained specimens

provided by Protein Expression Atlas (https://www.proteinatlas.org/). The images of the Figure 2

represent duodenum from 77-years-old female, kidney from 36-years-old male, testis from 38-

years-old male, lung from 61-years-old female, and nasopharyngeal mucosa from 78-years-old

female. According to Protein Expression Atlas the immunostainings were performed with the rabbit

anti-human polyclonal antibody (HPA000288; Sigma Aldrich, St. Louis, MO) raised against 111 N-

terminal amino acids of ACE2 and diluted 1:250 for the staining.

Promoter Analysis

Analysis of *ACE2* promoter regions was performed using the TFBSfootprinter tool

(https://github.com/thirtysix/TFBS_footprinting) which uses transcription-relevant data from

several major databases to enhance prediction of putative TFBSs, including: ATAC-Seq data from

ENCODE [64], transcription start sites and expression data from FANTOM5 [65], expression

quantitative trail loci from GTEx [66], TFBS metacluster data from GTRD [67], TFBS binding

profile data from JASPAR [68], and sequence and conservation data from Ensembl [69]. Detailed

description of this novel tool is under preparation (Barker et al. manuscript under preparation).

Previous studies identified two distinct tissue-specific transcription start sites (TSS) for intestine

and lung expression [48], which correspond to primary protein-coding Ensembl transcripts

ENST00000252519 and ENST00000427411, respectively. These two transcripts were targeted for

transcription factor binding site (TFBS) analysis; input parameters of 1,000 base pairs (bp)

upstream and 200 bp downstream, relative to the TSS.

Single-Cell RNA-Seq

Single-cell expression datasets were identified for relevant tissues/cells of lung (human) [70],

trachea (mouse) [71], and small intestine (mouse) [72]. Using a modified workflow described

previously in [73], for each dataset the samples were filtered by Gaussian fit of read count,

expressed gene count, and number of cells in which a gene is expressed. Counts were normalized by

cell, log transformed, principle component analysis performed with 15 components, and k-nearest

neighbors computed using SCANPY [74], and then the full data set normalized with R package

'scran' [75]. Batch correction by individual and sample region was performed with SCANPY using

the ComBat function. The top 1,000 genes with highly differential expression were identified for

cluster analysis which was performed with Uniform Manifold Approximation and Projection

(UMAP) and force directed graph models. The top 100 marker genes were identified as those with

higher expression unique to each cluster by Welch t-test in SCANPY. Expression of the *ACE2* gene

was mapped onto cluster figures to determine overlap with previously identified cell types or cell

type marker genes identified in the literature. Cell type was mapped by expression of known marker

genes of cell types expressed in the lung and small intestine, as defined by *de novo* prediction in the

original articles.

Statistics

Comparisons of ACE2 expression values in different tissues and between groups delineated by age

or sex, were carried out by one-way ANOVA using the stats package in the SciPy [62] Python

library. Only groups with 20 or more observations and a 2-sided chi squared probability of

normality of $<=0.1$ (due to the robustness of ANOVA to non-normal distributions) were used for

comparison. Correlation of gene expression values was calculated by two-sided Spearman rank-

order analysis, where a Bonferroni-corrected p-value threshold was computed using $\alpha=0.05$/number

of comparisons. Gene ontology enrichment analyses performed using the GProfiler tool utilize a

custom algorithm for multiple testing of dependent results, which corresponds to an experiment-

wide threshold of α=0.05. TFBSfootprinter analysis of the *ACE2* promoter limits results for

individual TFBSs whose score satisfies a genome-wide threshold of α=0.01.

REFERENCES

1. Skeggs LT, Jr., Kahn JR and Shumway NP (1956) The preparation and function of the hypertensin-converting enzyme. J Exp Med 103:295-9. doi: 10.1084/jem.103.3.295

2. Donoghue M, Hsieh F, Baronas E, Godbout K, Gosselin M, Stagliano N, Donovan M, Woolf B, Robison K, Jeyaseelan R, Breitbart RE and Acton S (2000) A novel angiotensin-converting enzyme-related carboxypeptidase (ACE2) converts angiotensin I to angiotensin 1-9. Circ Res 87:E1-9. doi: 10.1161/01.res.87.5.e1

3. Meldrum NU and Roughton FJW (1932) Some properties of carbonic anhydrase, the $CO_2$ enzyme present in blood. J. Physiol. 75:15-16.

4. Andreini C, Banci L, Bertini I and Rosato A (2006) Zinc through the three domains of life. J Proteome Res 5:3173-8. doi: 10.1021/pr0603699

5. Andreini C, Banci L, Bertini I and Rosato A (2006) Counting the zinc-proteins encoded in the human genome. J Proteome Res 5:196-201. doi: 10.1021/pr050361j

6. Tipnis SR, Hooper NM, Hyde R, Karran E, Christie G and Turner AJ (2000) A human homolog of angiotensin-converting enzyme. Cloning and functional expression as a captopril-insensitive carboxypeptidase. J Biol Chem 275:33238-43. doi: 10.1074/jbc.M002615200

7. Douglas GC, O'Bryan MK, Hedger MP, Lee DK, Yarski MA, Smith AI and Lew RA (2004) The novel angiotensin-converting enzyme (ACE) homolog, ACE2, is selectively expressed by adult Leydig cells of the testis. Endocrinology 145:4703-11. doi: 10.1210/en.2004-0443

8. Imai Y, Kuba K and Penninger JM (2008) The discovery of angiotensin-converting enzyme 2 and its role in acute lung injury in mice. Exp Physiol 93:543-8. doi: 10.1113/expphysiol.2007.040048

9. Cheng H, Wang Y and Wang GQ (2020) Organ-protective effect of angiotensin-converting enzyme 2 and its effect on the prognosis of COVID-19. J Med Virol. doi: 10.1002/jmv.25785

10. Li W, Moore MJ, Vasilieva N, Sui J, Wong SK, Berne MA, Somasundaran M, Sullivan JL, Luzuriaga K, Greenough TC, Choe H and Farzan M (2003) Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus. Nature 426:450-4. doi: 10.1038/nature02145

11. Lei C, Qian K, Li T, Zhang S, Fu W, Ding M and Hu S (2020) Neutralization of SARS-CoV-2 spike pseudotyped virus by recombinant ACE2-Ig. Nat Commun 11:2070. doi: 10.1038/s41467-020-16048-4

12. Monteil V, Kwon H, Prado P, Hagelkruys A, Wimmer RA, Stahl M, Leopoldi A, Garreta E, del Pozo CH, Prosper F, Romero JP, Wirnsberger G, Zhang H, Slutsky AS, Conder R, Montserrat N, Mirazimi A and Penninger JM (2020) Inhibition of SARS-CoV-2 infections in engineered human tissues using clinical-grade soluble human ACE2. Cell:in press. doi: 10.1016/j.cell.2020.04.004

13. Hoffmann M, Kleine-Weber H, Schroeder S, Kruger N, Herrler T, Erichsen S, Schiergens TS, Herrler G, Wu NH, Nitsche A, Muller MA, Drosten C and Pohlmann S (2020) SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. Cell 181:271-280 e8. doi: 10.1016/j.cell.2020.02.052

14. Shulla A, Heald-Sargent T, Subramanya G, Zhao J, Perlman S and Gallagher T (2011) A transmembrane serine protease is linked to the severe acute respiratory syndrome coronavirus receptor and activates virus entry. J Virol 85:873-82. doi: 10.1128/JVI.02062-10

15. Heurich A, Hofmann-Winkler H, Gierer S, Liepold T, Jahn O and Pohlmann S (2014) TMPRSS2 and ADAM17 cleave ACE2 differentially and only proteolysis by TMPRSS2 augments entry driven by the severe acute respiratory syndrome coronavirus spike protein. J Virol 88:1293-307. doi: 10.1128/JVI.02202-13

16. Li F, Li W, Farzan M and Harrison SC (2005) Structure of SARS coronavirus spike receptor-binding domain complexed with receptor. Science 309:1864-8. doi: 10.1126/science.1116480

17. Wang Q, Zhang Y, Wu L, Niu S, Song C, Zhang Z, Lu G, Qiao C, Hu Y, Yuen KY, Wang Q, Zhou H, Yan J and Qi J (2020) Structural and functional basis of SARS-CoV-2 entry by using human ACE2. Cell in press. doi: 10.1016/j.cell.2020.03.045

18. Shang J, Ye G, Shi K, Wan Y, Luo C, Aihara H, Geng Q, Auerbach A and Li F (2020) Structural basis of receptor recognition by SARS-CoV-2. Nature 581:221-224. doi: 10.1038/s41586-020-2179-y

19. Lan J, Ge J, Yu J, Shan S, Zhou H, Fan S, Zhang Q, Shi X, Wang Q, Zhang L and Wang X (2020)
Structure of the SARS-CoV-2 spike receptor-binding domain bound to the ACE2 receptor. Nature 581:215-
220. doi: 10.1038/s41586-020-2180-5
20. Walls AC, Park YJ, Tortorici MA, Wall A, McGuire AT and Veesler D (2020) Structure, Function,
and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. Cell 181:281-292 e6. doi: 10.1016/j.cell.2020.02.058
21. Chen Y, Guo Y, Pan Y and Zhao ZJ (2020) Structure analysis of the receptor binding of 2019-
nCoV. Biochem Biophys Res Commun 525:135-140. doi: 10.1016/j.bbrc.2020.02.071
22. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, Liu L, Shan H, Lei CL, Hui DSC, Du B, Li LJ, Zeng
G, Yuen KY, Chen RC, Tang CL, Wang T, Chen PY, Xiang J, Li SY, Wang JL, Liang ZJ, Peng YX, Wei L, Liu Y, Hu
YH, Peng P, Wang JM, Liu JY, Chen Z, Li G, Zheng ZJ, Qiu SQ, Luo J, Ye CJ, Zhu SY, Zhong NS and China
Medical Treatment Expert Group for C (2020) Clinical Characteristics of Coronavirus Disease 2019 in China.
N Engl J Med 382:1708-1720. doi: 10.1056/NEJMoa2002032
23. Lai CC, Shih TP, Ko WC, Tang HJ and Hsueh PR (2020) Severe acute respiratory syndrome
coronavirus 2 (SARS-CoV-2) and coronavirus disease-2019 (COVID-19): The epidemic and the challenges. Int
J Antimicrob Agents 55:105924. doi: 10.1016/j.ijantimicag.2020.105924
24. Qian GQ, Yang NB, Ding F, Ma AHY, Wang ZY, Shen YF, Shi CW, Lian X, Chu JG, Chen L, Wang
ZY, Ren DW, Li GX, Chen XQ, Shen HJ and Chen XM (2020) Epidemiologic and Clinical Characteristics of 91
Hospitalized Patients with COVID-19 in Zhejiang, China: A retrospective, multi-centre case series. QJM:in
press. doi: 10.1093/qjmed/hcaa089
25. Chen T, Wu D, Chen H, Yan W, Yang D, Chen G, Ma K, Xu D, Yu H, Wang H, Wang T, Guo W,
Chen J, Ding C, Zhang X, Huang J, Han M, Li S, Luo X, Zhao J and Ning Q (2020) Clinical characteristics of 113
deceased patients with coronavirus disease 2019: retrospective study. BMJ 368:m1091. doi:
10.1136/bmj.m1091
26. Hamming I, Timens W, Bulthuis ML, Lely AT, Navis G and van Goor H (2004) Tissue
distribution of ACE2 protein, the functional receptor for SARS coronavirus. A first step in understanding
SARS pathogenesis. J Pathol 203:631-7. doi: 10.1002/path.1570
27. Ding Y, He L, Zhang Q, Huang Z, Che X, Hou J, Wang H, Shen H, Qiu L, Li Z, Geng J, Cai J, Han
H, Li X, Kang W, Weng D, Liang P and Jiang S (2004) Organ distribution of severe acute respiratory syndrome
(SARS) associated coronavirus (SARS-CoV) in SARS patients: implications for pathogenesis and virus
transmission pathways. J Pathol 203:622-30. doi: 10.1002/path.1560
28. To KF and Lo AW (2004) Exploring the pathogenesis of severe acute respiratory syndrome
(SARS): the tissue distribution of the coronavirus (SARS-CoV) and its putative receptor, angiotensin-
converting enzyme 2 (ACE2). J Pathol 203:740-3. doi: 10.1002/path.1597
29. Chen J and Subbarao K (2007) The Immunobiology of SARS*. Annu Rev Immunol 25:443-72.
doi: 10.1146/annurev.immunol.25.022106.141706
30. Zhou Z, Zhao N, Shu Y, Han S, Chen B and Shu X (2020) Effect of gastrointestinal symptoms
on patients infected with COVID-19. Gastroenterology:in press. doi: 10.1053/j.gastro.2020.03.020
31. Sims AC, Baric RS, Yount B, Burkett SE, Collins PL and Pickles RJ (2005) Severe acute
respiratory syndrome coronavirus infection of human ciliated airway epithelia: role of ciliated cells in viral
spread in the conducting airways of the lungs. J Virol 79:15511-24. doi: 10.1128/JVI.79.24.15511-
15524.2005
32. Ren X, Glende J, Al-Falah M, de Vries V, Schwegmann-Wessels C, Qu X, Tan L, Tschernig T,
Deng H, Naim HY and Herrler G (2006) Analysis of ACE2 in polarized epithelial cells: surface expression and
function as receptor for severe acute respiratory syndrome-associated coronavirus. J Gen Virol 87:1691-5.
doi: 10.1099/vir.0.81749-0
33. Lai CC, Liu YH, Wang CY, Wang YH, Hsueh SC, Yen MY, Ko WC and Hsueh PR (2020)
Asymptomatic carrier state, acute respiratory disease, and pneumonia due to severe acute respiratory
syndrome coronavirus 2 (SARS-CoV-2): Facts and myths. J Microbiol Immunol Infect:in press. doi:
10.1016/j.jmii.2020.02.012
34. Cheung OY, Chan JW, Ng CK and Koo CK (2004) The spectrum of pathological changes in
severe acute respiratory syndrome (SARS). Histopathology 45:119-24. doi: 10.1111/j.1365-
2559.2004.01926.x

35.        Ksiazek TG, Erdman D, Goldsmith CS, Zaki SR, Peret T, Emery S, Tong S, Urbani C, Comer JA,
Lim W, Rollin PE, Dowell SF, Ling AE, Humphrey CD, Shieh WJ, Guarner J, Paddock CD, Rota P, Fields B,
DeRisi J, Yang JY, Cox N, Hughes JM, LeDuc JW, Bellini WJ, Anderson LJ and Group SW (2003) A novel
coronavirus associated with severe acute respiratory syndrome. N Engl J Med 348:1953-66. doi:
10.1056/NEJMoa030781
36.        Kuiken T, Fouchier RA, Schutten M, Rimmelzwaan GF, van Amerongen G, van Riel D, Laman
JD, de Jong T, van Doornum G, Lim W, Ling AE, Chan PK, Tam JS, Zambon MC, Gopal R, Drosten C, van der
Werf S, Escriou N, Manuguerra JC, Stohr K, Peiris JS and Osterhaus AD (2003) Newly discovered coronavirus
as the primary cause of severe acute respiratory syndrome. Lancet 362:263-70. doi: 10.1016/S0140-
6736(03)13967-0
37.        To KF, Tong JH, Chan PK, Au FW, Chim SS, Chan KC, Cheung JL, Liu EY, Tse GM, Lo AW, Lo YM
and Ng HK (2004) Tissue and cellular tropism of the coronavirus associated with severe acute respiratory
syndrome: an in-situ hybridization study of fatal cases. J Pathol 202:157-63. doi: 10.1002/path.1510
38.        Chow KC, Hsiao CH, Lin TY, Chen CL and Chiou SH (2004) Detection of severe acute
respiratory syndrome-associated coronavirus in pneumocytes of the lung. Am J Clin Pathol 121:574-80. doi:
10.1309/C0ED-U0RA-QBTX-BHCE
39.        Mossel EC, Wang J, Jeffers S, Edeen KE, Wang S, Cosgrove GP, Funk CJ, Manzer R, Miura TA,
Pearson LD, Holmes KV and Mason RJ (2008) SARS-CoV replicates in primary human alveolar type II cell
cultures but not in type I-like cells. Virology 372:127-35. doi: 10.1016/j.virol.2007.09.045
40.        Lukassen S, Lorenz Chua R, Trefzer T, Kahn NC, Schneider MA, Muley T, Winter H, Meister M,
Veith C, Boots AW, Hennig BP, Kreuter M, Conrad C and Eils R (2020) SARS-CoV-2 receptor ACE2 and
TMPRSS2 are primarily expressed in bronchial transient secretory cells. EMBO J:in press. doi:
10.15252/embj.20105114
41.        Guo J, Wei X, Li Q, Li L, Yang Z, Shi Y, Qin Y, Zhang X, Wang X, Zhi X and Meng D (2020) Single-
cell RNA analysis on ACE2 expression provides insight into SARS-CoV-2 blood entry and heart injury.
medRxiv:2020.03.31.20047621. doi: 10.1101/2020.03.31.20047621
42.        Qi F, Qian S, Zhang S and Zhang Z (2020) Single cell RNA sequencing of 13 human tissues
identify cell types and receptors of human coronaviruses. Biochem Biophys Res Commun 526:135-140. doi:
10.1016/j.bbrc.2020.03.044
43.        Zou X, Chen K, Zou J, Han P, Hao J and Han Z (2020) Single-cell RNA-seq data analysis on the
receptor ACE2 expression reveals the potential risk of different human organs vulnerable to 2019-nCoV
infection. Front Med 14:185-192. doi: 10.1007/s11684-020-0754-0
44.        Wiener RS, Cao YX, Hinds A, Ramirez MI and Williams MC (2007) Angiotensin converting
enzyme 2 is primarily epithelial and is developmentally regulated in the mouse lung. J Cell Biochem
101:1278-91. doi: 10.1002/jcb.21248
45.        Shi H, Han X, Jiang N, Cao Y, Alwalid O, Gu J, Fan Y and Zheng C (2020) Radiological findings
from 81 patients with COVID-19 pneumonia in Wuhan, China: a descriptive study. Lancet Infect Dis 20:425-
434. doi: 10.1016/S1473-3099(20)30086-4
46.        Franks TJ, Chong PY, Chui P, Galvin JR, Lourens RM, Reid AH, Selbs E, McEvoy CP, Hayden CD,
Fukuoka J, Taubenberger JK and Travis WD (2003) Lung pathology of severe acute respiratory syndrome
(SARS): a study of 8 autopsy cases from Singapore. Hum Pathol 34:743-8. doi: 10.1016/s0046-
8177(03)00367-8
47.        Nicholls JM, Poon LL, Lee KC, Ng WF, Lai ST, Leung CY, Chu CM, Hui PK, Mak KL, Lim W, Yan
KW, Chan KH, Tsang NC, Guan Y, Yuen KY and Peiris JS (2003) Lung pathology of fatal severe acute
respiratory syndrome. Lancet 361:1773-8. doi: 10.1016/s0140-6736(03)13413-7
48.        Pedersen KB, Chhabra KH, Nguyen VK, Xia H and Lazartigues E (2013) The transcription factor
HNF1alpha induces expression of angiotensin-converting enzyme 2 (ACE2) in pancreatic islets from
evolutionarily conserved promoter motifs. Biochim Biophys Acta 1829:1225-35. doi:
10.1016/j.bbagrm.2013.09.007
49.        Pedersen KB, Chodavarapu H and Lazartigues E (2017) Forkhead Box Transcription Factors of
the FOXA Class Are Required for Basal Transcription of Angiotensin-Converting Enzyme 2. J Endocr Soc
1:370-384. doi: 10.1210/js.2016-1071

50.        Zhang R, Wu Y, Zhao M, Liu C, Zhou L, Shen S, Liao S, Yang K, Li Q and Wan H (2009) Role of
HIF-1alpha in the regulation ACE and ACE2 expression in hypoxic human pulmonary artery smooth muscle
cells. Am J Physiol Lung Cell Mol Physiol 297:L631-40. doi: 10.1152/ajplung.90415.2008
51.        Slemc L and Kunej T (2016) Transcription factor HIF1A: downstream targets, associated
pathways, polymorphic hypoxia response element (HRE) sites, and initiative for standardization of
reporting in scientific literature. Tumour Biol 37:14851-14861. doi: 10.1007/s13277-016-5331-4
52.        Faguer S, Mayeur N, Casemayou A, Pageaud AL, Courtellemont C, Cartery C, Fournie GJ,
Schanstra JP, Tack I, Bascands JL and Chauveau D (2013) Hnf-1beta transcription factor is an early hif-
1alpha-independent marker of epithelial hypoxia and controls renal repair. PLoS One 8:e63585. doi:
10.1371/journal.pone.0063585
53.        Mazumdar J, O'Brien WT, Johnson RS, LaManna JC, Chavez JC, Klein PS and Simon MC (2010)
O2 regulates stem cells through Wnt/beta-catenin signalling. Nat Cell Biol 12:1007-13. doi:
10.1038/ncb2102
54.        Patel AB and Verma A (2020) COVID-19 and Angiotensin-Converting Enzyme Inhibitors and
Angiotensin Receptor Blockers: What Is the Evidence? JAMA:in press. doi: 10.1001/jama.2020.4812
55.        Magrone T, Magrone M and Jirillo E (2020) Focus on Receptors for Coronaviruses with
Special Reference to Angiotensin-converting Enzyme 2 as a Potential Drug Target - A Perspective. Endocr
Metab Immune Disord Drug Targets:in press. doi: 10.2174/1871530320666200427112902
56.        Ferrario CM, Jessup J, Chappell MC, Averill DB, Brosnihan KB, Tallant EA, Diz DI and Gallagher
PE (2005) Effect of angiotensin-converting enzyme inhibition and angiotensin II receptor blockers on
cardiac angiotensin-converting enzyme 2. Circulation 111:2605-10. doi:
10.1161/CIRCULATIONAHA.104.510461
57.        Santos RAS, Sampaio WO, Alzamora AC, Motta-Santos D, Alenina N, Bader M and
Campagnole-Santos MJ (2018) The ACE2/Angiotensin-(1-7)/MAS Axis of the Renin-Angiotensin System:
Focus on Angiotensin-(1-7). Physiol Rev 98:505-553. doi: 10.1152/physrev.00023.2016
58.        Tian S, Hu W, Niu L, Liu H, Xu H and Xiao SY (2020) Pulmonary Pathology of Early-Phase 2019
Novel Coronavirus (COVID-19) Pneumonia in Two Patients With Lung Cancer. J Thorac Oncol 15:700-704.
doi: 10.1016/j.jtho.2020.02.010
59.        Consortium F, the RP, Clst, Forrest AR, Kawaji H, Rehli M, Baillie JK, de Hoon MJ, Haberle V,
Lassmann T, Kulakovskiy IV, Lizio M, Itoh M, Andersson R, Mungall CJ, Meehan TF, Schmeier S, Bertin N,
Jorgensen M, Dimont E, Arner E, Schmidl C, Schaefer U, Medvedeva YA, Plessy C, Vitezic M, Severin J,
Semple C, Ishizu Y, Young RS, Francescatto M, Alam I, Albanese D, Altschuler GM, Arakawa T, Archer JA,
Arner P, Babina M, Rennie S, Balwierz PJ, Beckhouse AG, Pradhan-Bhatt S, Blake JA, Blumenthal A, Bodega
B, Bonetti A, Briggs J, Brombacher F, Burroughs AM, Califano A, Cannistraci CV, Carbajo D, Chen Y, Chierici
M, Ciani Y, Clevers HC, Dalla E, Davis CA, Detmar M, Diehl AD, Dohi T, Drablos F, Edge AS, Edinger M, Ekwall
K, Endoh M, Enomoto H, Fagiolini M, Fairbairn L, Fang H, Farach-Carson MC, Faulkner GJ, Favorov AV, Fisher
ME, Frith MC, Fujita R, Fukuda S, Furlanello C, Furino M, Furusawa J, Geijtenbeek TB, Gibson AP, Gingeras T,
Goldowitz D, Gough J, Guhl S, Guler R, Gustincich S, Ha TJ, Hamaguchi M, Hara M, Harbers M, Harshbarger
J, Hasegawa A, Hasegawa Y, Hashimoto T, Herlyn M, Hitchens KJ, Ho Sui SJ, Hofmann OM, Hoof I, Hori F,
Huminiecki L, Iida K, Ikawa T, Jankovic BR, Jia H, Joshi A, Jurman G, Kaczkowski B, Kai C, Kaida K, Kaiho A,
Kajiyama K, Kanamori-Katayama M, Kasianov AS, Kasukawa T, Katayama S, Kato S, Kawaguchi S, Kawamoto
H, Kawamura YI, Kawashima T, Kempfle JS, Kenna TJ, Kere J, Khachigian LM, Kitamura T, Klinken SP, Knox AJ,
Kojima M, Kojima S, Kondo N, Koseki H, Koyasu S, Krampitz S, Kubosaki A, Kwon AT, Laros JF, Lee W,
Lennartsson A, Li K, Lilje B, Lipovich L, Mackay-Sim A, Manabe R, Mar JC, Marchand B, Mathelier A, Mejhert
N, Meynert A, Mizuno Y, de Lima Morais DA, Morikawa H, Morimoto M, Moro K, Motakis E, Motohashi H,
Mummery CL, Murata M, Nagao-Sato S, Nakachi Y, Nakahara F, Nakamura T, Nakamura Y, Nakazato K, van
Nimwegen E, Ninomiya N, Nishiyori H, Noma S, Noma S, Noazaki T, Ogishima S, Ohkura N, Ohimiya H, Ohno
H, Ohshima M, Okada-Hatakeyama M, Okazaki Y, Orlando V, Ovchinnikov DA, Pain A, Passier R, Patrikakis
M, Persson H, Piazza S, Prendergast JG, Rackham OJ, Ramilowski JA, Rashid M, Ravasi T, Rizzu P, Roncador
M, Roy S, Rye MB, Saijyo E, Sajantila A, Saka A, Sakaguchi S, Sakai M, Sato H, Savvi S, Saxena A, Schneider C,
Schultes EA, Schulze-Tanzil GG, Schwegmann A, Sengstag T, Sheng G, Shimoji H, Shimoni Y, Shin JW, Simon
C, Sugiyama D, Sugiyama T, Suzuki M, Suzuki N, Swoboda RK, t Hoen PA, Tagami M, Takahashi N, Takai J,

Tanaka H, Tatsukawa H, Tatum Z, Thompson M, Toyodo H, Toyoda T, Valen E, van de Wetering M, van den
Berg LM, Verado R, Vijayan D, Vorontsov IE, Wasserman WW, Watanabe S, Wells CA, Winteringham LN,
Wolvetang E, Wood EJ, Yamaguchi Y, Yamamoto M, Yoneda M, Yonekura Y, Yoshida S, Zabierowski SE,
Zhang PG, Zhao X, Zucchelli S, Summers KM, Suzuki H, Daub CO, Kawai J, Heutink P, Hide W, Freeman TC,
Lenhard B, Bajic VB, Taylor MS, Makeev VJ, Sandelin A, Hume DA, Carninci P and Hayashizaki Y (2014) A
promoter-level mammalian expression atlas. Nature 507:462-70. doi: 10.1038/nature13182
60.        Hunter JD (2007) Matplotlib: A 2D Graphics Environment. Computing in Science &
Engineering 9:90-95. doi: 10.1109/MCSE.2007.55
61.        Waskom M, Botwinnik O, Ostblom J, Lukauskas S, Hobson P, MaozGelbart, Gemperline DC,
Augspurger T, Halchenko Y and Cole JB (2020) Mwaskom/Seaborn: V0.10.0 (January 2020). Zenodo. doi:
10.5281/zenodo.3629446
62.        Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, Burovski E,
Peterson P, Weckesser W, Bright J, van der Walt SJ, Brett M, Wilson J, Millman KJ, Mayorov N, Nelson ARJ,
Jones E, Kern R, Larson E, Carey CJ, Polat I, Feng Y, Moore EW, VanderPlas J, Laxalde D, Perktold J, Cimrman
R, Henriksen I, Quintero EA, Harris CR, Archibald AM, Ribeiro AH, Pedregosa F, van Mulbregt P and SciPy C
(2020) SciPy 1.0: fundamental algorithms for scientific computing in Python. Nat Methods 17:261-272. doi:
10.1038/s41592-019-0686-2
63.        Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H and Vilo J (2019) g:Profiler: a
web server for functional enrichment analysis and conversions of gene lists (2019 update). Nucleic Acids
Res 47:W191-W198. doi: 10.1093/nar/gkz369
64.        Consortium EP (2012) An integrated encyclopedia of DNA elements in the human genome.
Nature 489:57-74. doi: 10.1038/nature11247
65.        Lizio M, Harshbarger J, Shimoji H, Severin J, Kasukawa T, Sahin S, Abugessaisa I, Fukuda S,
Hori F, Ishikawa-Kato S, Mungall CJ, Arner E, Baillie JK, Bertin N, Bono H, de Hoon M, Diehl AD, Dimont E,
Freeman TC, Fujieda K, Hide W, Kaliyaperumal R, Katayama T, Lassmann T, Meehan TF, Nishikata K, Ono H,
Rehli M, Sandelin A, Schultes EA, t Hoen PA, Tatum Z, Thompson M, Toyoda T, Wright DW, Daub CO, Itoh
M, Carninci P, Hayashizaki Y, Forrest AR, Kawaji H and consortium F (2015) Gateways to the FANTOM5
promoter level mammalian expression atlas. Genome Biol 16:22. doi: 10.1186/s13059-014-0560-6
66.        Consortium GT (2015) Human genomics. The Genotype-Tissue Expression (GTEx) pilot
analysis: multitissue gene regulation in humans. Science 348:648-60. doi: 10.1126/science.1262110
67.        Yevshin I, Sharipov R, Valeev T, Kel A and Kolpakov F (2017) GTRD: a database of
transcription factor binding sites identified by ChIP-seq experiments. Nucleic Acids Res 45:D61-D67. doi:
10.1093/nar/gkw951
68.        Vlieghe D, Sandelin A, De Bleser PJ, Vleminckx K, Wasserman WW, van Roy F and Lenhard B
(2006) A new generation of JASPAR, the open-access repository for transcription factor binding site profiles.
Nucleic Acids Res 34:D95-7. doi: 10.1093/nar/gkj115
69.        Yates AD, Achuthan P, Akanni W, Allen J, Allen J, Alvarez-Jarreta J, Amode MR, Armean IM,
Azov AG, Bennett R, Bhai J, Billis K, Boddu S, Marugan JC, Cummins C, Davidson C, Dodiya K, Fatima R, Gall
A, Giron CG, Gil L, Grego T, Haggerty L, Haskell E, Hourlier T, Izuogu OG, Janacek SH, Juettemann T, Kay M,
Lavidas I, Le T, Lemos D, Martinez JG, Maurel T, McDowall M, McMahon A, Mohanan S, Moore B, Nuhn M,
Oheh DN, Parker A, Parton A, Patricio M, Sakthivel MP, Abdul Salam AI, Schmitt BM, Schuilenburg H,
Sheppard D, Sycheva M, Szuba M, Taylor K, Thormann A, Threadgold G, Vullo A, Walts B, Winterbottom A,
Zadissa A, Chakiachvili M, Flint B, Frankish A, Hunt SE, G II, Kostadima M, Langridge N, Loveland JE, Martin
FJ, Morales J, Mudge JM, Muffato M, Perry E, Ruffier M, Trevanion SJ, Cunningham F, Howe KL, Zerbino DR
and Flicek P (2020) Ensembl 2020. Nucleic Acids Res 48:D682-D688. doi: 10.1093/nar/gkz966
70.        Adams TS, Schupp JC, Poli S, Ayaub EA, Neumark N, Ahangari F, Chu SG, Raby BA, DeIuliis G,
Januszyk M, Duan Q, Arnett HA, Siddiqui A, Washko GR, Homer R, Yan X, Rosas IO and Kaminski N (2019)
Single Cell RNA-seq reveals ectopic and aberrant lung resident cell populations in Idiopathic Pulmonary
Fibrosis. bioRxiv:759902. doi: 10.1101/759902
71.        Montoro DT, Haber AL, Biton M, Vinarsky V, Lin B, Birket SE, Yuan F, Chen S, Leung HM,
Villoria J, Rogel N, Burgin G, Tsankov AM, Waghray A, Slyper M, Waldman J, Nguyen L, Dionne D,
Rozenblatt-Rosen O, Tata PR, Mou H, Shivaraju M, Bihler H, Mense M, Tearney GJ, Rowe SM, Engelhardt JF,

Regev A and Rajagopal J (2018) A revised airway epithelial hierarchy includes CFTR-expressing ionocytes.
Nature 560:319-324. doi: 10.1038/s41586-018-0393-7
72.        Haber AL, Biton M, Rogel N, Herbst RH, Shekhar K, Smillie C, Burgin G, Delorey TM, Howitt
MR, Katz Y, Tirosh I, Beyaz S, Dionne D, Zhang M, Raychowdhury R, Garrett WS, Rozenblatt-Rosen O, Shi
HN, Yilmaz O, Xavier RJ and Regev A (2017) A single-cell survey of the small intestinal epithelium. Nature
551:333-339. doi: 10.1038/nature24489
73.        Luecken MD and Theis FJ (2019) Current best practices in single-cell RNA-seq analysis: a
tutorial. Mol Syst Biol 15:e8746. doi: 10.15252/msb.20188746
74.        Wolf FA, Angerer P and Theis FJ (2018) SCANPY: large-scale single-cell gene expression data
analysis. Genome Biol 19:15. doi: 10.1186/s13059-017-1382-0
75.        Lun AT, Bach K and Marioni JC (2016) Pooling across cells to normalize single-cell RNA
sequencing data with many zero counts. Genome Biol 17:75. doi: 10.1186/s13059-016-0947-7
694

**FIGURES**

Figure 1. Expression of ACE2 mRNA in selected human tissues. Expression values as TPM have

been extracted from the FANTOM5 dataset.

Figure 2. Immunohistochemical localization of ACE2 protein in selected human tissues. In the

duodenum (A), the protein is most strongly localized to the apical plasma membrane of absorptive

enterocytes (arrows). The goblet cells (arrowheads) show weaker apical staining. Intracellular

staining is confined to the absorptive enterocytes. In the kidney (B), ACE2 shows prominent apical

staining in the epithelial cells of the proximal convoluted tubules (arrows) and Bowman´s capsule

epithelium (arrowheads). The distal convoluted tubules are negative (asterisk). The testis specimen

(C) shows strong immunostaining in the seminiferous tubules (arrows) and interstitial cells

(arrowheads). The lung sample (D) is negative. In the nasopharyngeal mucosa (E), ACE2 signal is

very weak and only occasional epithelial cells show weak signals (arrows). Immunostained

specimens were taken from the Protein Expression Atlas (https://www.proteinatlas.org/).

Figure 3. Single cell RNA-Seq analysis of different cell types from the lung (human), derived from

data from GEO dataset GSE136831 [70]. ACE2 mRNA expression as normalized, batch-corrected

counts is shown for comparison in upper panel. The force directed layout plot was computed and

visualized in ScanPy [74]. For each cell type the ratio of cells expressing ACE2 is presented in

addition to a stacked barplot of the relative cell type frequencies in the whole dataset. Alveolar type

I (ATI), alveolar type II (ATII), pulmonary neuroendocrine cells (PNEC), smooth muscle cells

(SMC), vascular endothelia (VE).

## ACE2

## Cells Expressing ACE2

## Cell Type

## Cell Type Ratio

Figure 4. Single cell RNA-Seq analysis of different cell types from the respiratory tract (mouse

tracheal epithelium), derived from data from GEO dataset GSE103354 [71]. ACE2 mRNA

expression as normalized, batch-corrected counts is shown for comparison in upper panel. The force

directed layout plot was computed and visualized in ScanPy [74]. For each cell type the ratio of

cells expressing ACE2 is presented in addition to a stacked barplot of the relative cell type

frequencies in the whole dataset.

Figure 5. Single cell RNA-Seq analysis of mouse intestinal epithelial cells. Data is from GEO

dataset GSE92332 [72]. ACE2 mRNA expression as normalized, batch-corrected counts is shown

for comparison in upper panel. The force directed layout plot was computed and visualized in

ScanPy [74]. For each cell type the ratio of cells expressing ACE2 is presented in addition to a

stacked barplot of the relative cell type frequencies in the whole dataset.

## ACE2

## Cells Expressing ACE2

## Cell Type

## Cell Type Ratio

Figure 6. Effect of age on ACE2 mRNA expression levels. Data is extracted from the GTEx dataset

as TPM. In these organs, ANOVA revealed significant differences between age categories in tibial

nerve (p=8.58 x 10$^{-6}$), minor salivary gland (p=0.002), and whole blood (p=0.005). In other tissues,

the differences did not reach statistical significance. The highest TPM values are seen the small

intestine, testis, and kidney.

Figure 7. Effect of gender on ACE2 mRNA expression levels. Data is extracted from the GTEx

dataset as TPM. The expression levels in males and females overlap in all tissue categories.

Statistically significant differences studied by ANOVA analysis were determined in esophagus-

gastroesophageal junction (p=0.024) and whole blood (p=0.0002). The ACE2 mRNA expression

levels in testis specimens are shown here for comparison. 1=male, 2=female.

Figure 8. Prediction of transcription factor binding sites in the human *ACE2* gene promoter regions

of Ensembl transcripts ENST00000252519 and ENST00000427411 using TFBSfootprinter. A)

Promoter region of the intestine specific ACE2 transcript. The results show putative binding sites

for several transcription factors which have a strong correlation of expression with ACE2 in colon,

kidney, and ileum. The predicted binding sites overlap regions of conservation in mammal species

(Ensembl GERP), and cluster within 400 base pairs (bp) of the transcription start site. B) Promoter

region of the lung specific ACE2 transcript. The predicted binding sites cluster within 200 bp of the

TSS and overlap regions of conservation in mammal species, ATAC-Seq peaks (ENCODE), and

TFBS metaclusters (GTRD).

A

B

762     Figure 9. Expression of genes most highly correlated with ACE2 in single cell datasets of trachea

763     and intestinal epithelia. Trachea expression data is taken from GSE103354 [71] and intestinal

764     epithelia data is derived from GSE92332 [72]. Visualized in ScanPy [74].

Table 1. Genes associated with ACE2 mRNA expression in selected human tissues. Derived from GTEx bulk RNA-Seq data [66].

| Tissue | Correlated_gene | Correlation | p-value | HGNC | UniProt | Description | Panther protein class |
|---|---|---|---|---|---|---|---|
| terminal ileum | *ENPEP* | 0.9653 | 8.29E-110 | 3355 | Q07075 | Glutamyl aminopeptidase | metalloprotease (PC00153) |
| terminal ileum | *MEP1B* | 0.9653 | 8.35E-110 | 7020 | Q16820 | Meprin A subunit beta | metalloprotease (PC00153) |
| terminal ileum | *SLC15A1* | 0.9621 | 2.19E-106 | 10920 | P46059 | Solute carrier family 15 member 1 | transporter (PC00227) |
| terminal ileum | *MEP1A* | 0.9619 | 4.4E-106 | 7015 | Q16819 | Meprin A subunit alpha | metalloprotease (PC00153) |
| terminal ileum | *SLC3A1* | 0.9608 | 5.63E-105 | 11025 | Q07837 | Neutral and basic amino acid transport protein rBAT | amylase (PC00048) |
| terminal ileum | *APOB* | 0.9587 | 6.45E-103 | 603 | P04114 | Apolipoprotein B-100 | |
| terminal ileum | *SLC6A19* | 0.9581 | 2.33E-102 | 27960 | Q695T7 | Sodium-dependent neutral amino acid transporter B(0)AT1 | primary active transporter (PC00068) |
| terminal ileum | *ANPEP* | 0.9562 | 1.16E-100 | 500 | P15144 | Aminopeptidase N | metalloprotease (PC00153) |
| terminal ileum | *MTTP* | 0.9559 | 2.45E-100 | 7467 | P55157 | Microsomal triglyceride transfer protein large subunit | transporter (PC00227) |
| terminal ileum | *TUBAL3* | 0.9554 | 6.24E-100 | 23534 | A6NHL2 | Tubulin alpha chain-like 3 | tubulin (PC00228) |
| testis | *PLP1* | 0.9092 | 1.07E-138 | 9086 | P60201 | Myelin proteolipid protein | myelin protein (PC00161) |
| colon-transverse | *MEP1A* | 0.9056 | 1.52E-152 | 7015 | Q16819 | Meprin A subunit alpha | metalloprotease (PC00153) |
| kidney | *TINAG* | 0.9012 | 7.07E-32 | 14599 | Q9UJW2 | Tubulointerstitial nephritis antigen | cysteine protease (PC00081) |
| colon-transverse | *GDA* | 0.8964 | 8.19E-145 | 4212 | Q9Y2T3 | Guanine deaminase | deaminase (PC00088) |
| colon-transverse | *MGAM2* | 0.8963 | 9E-145 | 28101 | Q2M2H8 | Probable maltase-glucoamylase 2 | glucosidase (PC00108) |
| colon-transverse | *TMEM236* | 0.894 | 6.66E-143 | 23473 | Q5W0B7 | Transmembrane protein 236 | |
| colon-transverse | *GDPD2* | 0.8935 | 1.58E-142 | 25974 | Q9HCC8 | Glycerophosphoinositol inositolphosphodiesterase GDPD2 | |
| colon-transverse | *PLS1* | 0.8932 | 2.81E-142 | 9090 | Q14651 | Plastin-1 | non-motor actin binding protein (PC00165) |
| colon-transverse | *HHLA2* | 0.893 | 3.59E-142 | 4905 | Q9UM44 | HERV-H LTR-associating protein 2 | immunoglobulin receptor superfamily (PC00124) |
| colon-transverse | *SLC3A1* | 0.8928 | 5.27E-142 | 11025 | Q07837 | Neutral and basic amino acid transport protein rBAT | amylase (PC00048) |
| colon-transverse | *COL17A1* | 0.8914 | 6.12E-141 | 2194 | Q9UMD9 | Collagen alpha-1(XVII) chain | extracellular matrix structural protein (PC00103) |
| testis | *NDFIP1* | 0.8911 | 2.91E-125 | 17592 | Q9BT67 | NEDD4 family-interacting protein 1 | |
| testis | *SPARC* | 0.8908 | 4.67E-125 | 11219 | P09486 | SPARC | extracellular matrix glycoprotein (PC00100) |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| **colon-transverse** | *TINAG* | 0.8905 | 2.94E-140 | 14599 | Q9UJW2 | Tubulointerstitial nephritis antigen | cysteine protease (PC00081) |
| **testis** | *SCP2* | 0.8886 | 1.34E-123 | 10606 | P22307 | Non-specific lipid-transfer protein | transfer/carrier protein (PC00219) |
| **testis** | *SMARCA1* | 0.8852 | 2.44E-121 | 11097 | P28370 | Probable global transcription activator SNF2L1 | DNA helicase (PC00011) |
| **testis** | *ATP2B1* | 0.8829 | 6.45E-120 | 814 | P20020 | Plasma membrane calcium-transporting ATPase 1 | primary active transporter (PC00068) |
| **testis** | *VSIG1* | 0.8815 | 4.6E-119 | 28675 | Q86XK7 | V-set and immunoglobulin domain-containing protein 1 | |
| **testis** | *HTATSF1* | 0.8809 | 1.09E-118 | 5276 | O43719 | HIV Tat-specific factor 1 | RNA splicing factor (PC00148) |
| **testis** | *ACO1* | 0.8808 | 1.26E-118 | 117 | P21399 | Cytoplasmic aconitate hydratase | RNA binding protein (PC00031) |
| **testis** | *ENPP5* | 0.8802 | 3.01E-118 | 13717 | Q9UJA9 | Ectonucleotide pyrophosphatase/phosphodiesterase family member 5 | nucleotide phosphatase (PC00173) |
| **kidney** | *METTL7B* | 0.8705 | 2.78E-27 | 28276 | Q6UX53 | Methyltransferase-like protein 7B | methyltransferase (PC00155) |
| **kidney** | *ANPEP* | 0.8683 | 5.3E-27 | 500 | P15144 | Aminopeptidase N | metalloprotease (PC00153) |
| **kidney** | *LRP2* | 0.8671 | 7.59E-27 | 6694 | P98164 | Low-density lipoprotein receptor-related protein 2 | |
| **kidney** | *SLC13A1* | 0.8667 | 8.5E-27 | 10916 | Q9BZW2 | Solute carrier family 13 member 1 | secondary carrier transporter (PC00258) |
| **kidney** | *UGT3A1* | 0.8608 | 4.52E-26 | 26625 | Q6NUS8 | UDP-glucuronosyltransferase 3A1 | |
| **kidney** | *SLC27A2* | 0.8535 | 3.26E-25 | 10996 | O14975 | Very long-chain acyl-CoA synthetase | secondary carrier transporter (PC00258) |
| **kidney** | *TMEM27* | 0.8526 | 4.1E-25 | 29437 | Q9HBJ8 | Collectrin | |
| **kidney** | *CLRN3* | 0.8443 | 3.36E-24 | 20795 | Q8NCR9 | Clarin-3 | |
| **kidney** | *ACP5* | 0.8367 | 2.05E-23 | 124 | P13686 | Tartrate-resistant acid phosphatase type 5 | |
| **lung** | *PARM1* | 0.6275 | 1.32E-64 | 24536 | Q6UWI2 | Prostate androgen-regulated mucin-like protein 1 | |
| **lung** | *CNR1* | 0.6055 | 4.21E-59 | 2159 | P21554 | Cannabinoid receptor 1 | G-protein coupled receptor (PC00021) |
| **lung** | *NUDT16* | 0.5918 | 6.52E-56 | 26442 | Q96DE0 | U8 snoRNA-decapping enzyme | |
| **lung** | *DUOX1* | 0.5817 | 1.23E-53 | 3062 | Q9NRD9 | Dual oxidase 1 | oxidase (PC00175) |
| **lung** | *DUOXA1* | 0.5739 | 6.2E-52 | 26507 | Q1HG43 | Dual oxidase maturation factor 1 | |
| **lung** | *FMN1* | 0.571 | 2.53E-51 | 3768 | Q68DA7 | Formin-1 | |
| **lung** | *TMEM164* | 0.5692 | 6.05E-51 | 26217 | Q5U3C3 | Transmembrane protein 164 | |
| **lung** | *HSD17B4* | 0.5673 | 1.54E-50 | 5213 | P51659 | Peroxisomal multifunctional enzyme type 2 | |
| **lung** | *MGST1* | 0.5631 | 1.16E-49 | 7061 | P10620 | Microsomal glutathione S-transferase 1 | |
| **lung** | *SHMT1* | 0.5594 | 6.77E-49 | 10850 | P34896 | Serine hydroxymethyltransferase, cytosolic | methyltransferase (PC00155) |

Table 2. Gene ontology annotation results for the processes associated with genes strongly coexpressed (≥0.5) with ACE2 across all tissues in GTEx dataset. Calculated with GProfiler Python library [63].

| Source | Native | Process | p-value |
|--------|--------|---------|---------|
| GO:BP | GO:0001525 | angiogenesis | 1.797E-08 |
| GO:BP | GO:0048514 | blood vessel morphogenesis | 4.611E-08 |
| GO:BP | GO:0001944 | vasculature development | 1.893E-07 |
| GO:BP | GO:0072358 | cardiovascular system development | 2.365E-07 |
| GO:BP | GO:0001568 | blood vessel development | 4.746E-07 |
| GO:BP | GO:0035239 | tube morphogenesis | 5.907E-07 |
| GO:BP | GO:0072359 | circulatory system development | 6.641E-07 |
| GO:BP | GO:0048646 | anatomical structure formation involved in morphogenesis | 1.885E-05 |
| GO:BP | GO:0035295 | tube development | 2.900E-05 |
| GO:BP | GO:1901342 | regulation of vasculature development | 0.025 |
| GO:BP | GO:0008217 | regulation of blood pressure | 0.035 |
| GO:BP | GO:0045765 | regulation of angiogenesis | 0.009 |
| GO:CC | GO:0071944 | cell periphery | 0.001 |
| GO:CC | GO:0009986 | cell surface | 0.002 |
| GO:CC | GO:0005886 | plasma membrane | 0.013 |
| GO:CC | GO:0046930 | pore complex | 0.003 |
| HP | HP:0005381 | recurrent meningococcal disease | 0.004 |
| HP | HP:0005430 | recurrent Neisserial infections | 0.007 |
| HP | HP:0100601 | eclampsia | 0.021 |
| KEGG | KEGG:04610 | complement and coagulation cascades | 0.002 |
| KEGG | KEGG:04923 | regulation of lipolysis in adipocytes | 0.037 |
| WP | WP:WP558 | complement and coagulation cascades | 0.011 |