1

4 Correcting False Memories: The Effect of Mnemonic Generalization

5 on Original Memory Traces

7 Nathan M. Muncy[1] and C. Brock Kirwan[1,2*]

11 [1] Department of Psychology, Brigham Young University, Provo, Utah, USA and [2] Neuroscience
12 Center, Brigham Young University, Provo, Utah, USA
13

15 * Corresponding author

16 E-mail: kirwan@byu.edu (CBK)

Abstract

False memories are a common occurrence but the impact of misremembering on the original memory trace is ill-described. While the original memory may be rewritten, it is also possible for a second false memory to exist concurrently with the original, and if a false memory exists concurrently then recovery of the original information should be possible. This study investigates first, whether false recognition overwrites the original memory representation using a mnemonic discrimination task, and second, which neural processes are involved in recovering the original memory following a false memory. Thirty-five healthy, young adults performed multiple recognition memory tests, where the design of the experiment induced participants to make memory errors in the first recognition memory test and then allowed us to determine whether the memory error would be corrected in the second test session. FMRI signal associated with the encoding and retrieval processes during the experiment were investigated in order to determine the important regions for false memory correction. We found that false memories do not overwrite the original trace in all instances, as recovery of the original information was possible. Critically, we determined that recovery of the original information was dependent on higher-order processes during the formation of the false memory during the first test, and not on processing at the time of encoding or the second test episode.

KEYWORDS

episodic memory, discrimination, generalization, functional magnetic resonance imaging

1   INTRODUCTION

The field of study on mnemonic discrimination in episodic memory has largely focused on a single aspect—the capacity of the participant to correctly identify lure stimuli. In animal models, where pattern separation processes can be targeted, previous work has demonstrated that the successful identification of lures is dependent on the orthogonalization processes of the dentate gyrus in the hippocampus as well as an intact hippocampal trisynaptic pathway (Gilbert et al., 1998, 2001; Gilbert & Kesner, 2006; Gold & Kesner, 2005; Hunsaker et al., 2008; Kesner, 2007a, 2007b; Leutgeb et al., 2007). Human studies have implicated the same processes for successful mnemonic discrimination performance: participants with healthy hippocampi perform well on tests of mnemonic discrimination while participants with hippocampal atrophy perform poorly and participants with hippocampal lesions perform worst of all (Bakker et al., 2008; Kirwan et al., 2012; Mueller et al., 2011; Small et al., 2011). Additionally, hippocampal CA3/DG volumes in humans are known to positively correlate with performance on tests of mnemonic discrimination (Doxey & Kirwan, 2015; Grady & Ryan, 2017; Leal & Yassa, 2014; Marshall et al., 2016; Reagh et al., 2014; Roberts et al., 2014; Sheppard et al., 2016). That said, while mnemonic discrimination processes appear to be pattern separation dependent, higher order processes such as semantic labeling, visual spatial processing, attention, and decisiveness (Goebel & Vincze, 2007; Hunsaker & Kesner, 2013; Lacy et al., 2011; Leal & Yassa, 2014; Pidgeon & Morcom, 2016; Reagh et al., 2016; Reagh & Yassa, 2014; Rolls & Kesner, 2016; Steemers et al., 2016; Wais et al., 2017) integrate with hippocampal pattern separation output to coordinate the overall behavior. But while the roles of the medial temporal lobe structures in mnemonic discrimination are well understood, higher-order processes are relatively understudied in this context.

In addition to these gaps in the field of mnemonic discrimination, the failure in detecting

lures is also relatively unaddressed. The detection of lure stimuli in a recognition memory test is

the result of both encoding and retrieval processes: the lure's representation must be successfully

orthogonalized from the target's, and the target must be successfully retrieved. At the point of

successful lure identification, then, two distinct representations exist (Target, Lure) between

which the participant is able to discriminate. This process is sometimes known as "recall to

reject" (J. Kim & Yassa, 2013; Kirwan & Stark, 2007; Lacy et al., 2011; van den Honert et al.,

2016). A failure in lure detection, then, could have multiple antecedents: a failure to encode the

lure's salient details, a failure to retrieve the salient target information, and/or lure interference

may lead to mnemonic generalization rather than discrimination. Such failures may result from

underperforming lower-order hippocampal processes, but may also be the result of deficient

attention, semantic labelling, and/or decision making. Further, and perhaps more interestingly,

the consequence of incorrectly identifying the Lure as Target (a False Alarm, FA) is unknown: as

studies of mnemonic discrimination typically use a single study/test presentation, the effect of

the FA response on the original memory trace has not been established. It is possible that either

the lure representation could overwrite the target memory trace, or that a second representation

could exist concurrently with the original. If the latter is true and multiple memory traces result

from a FA response, then the two memory traces (one of the Target, and one of the Lure) may be

dissociable in subsequent testing. And while certain studies, particularly in the false-memory

literature, have specifically targeted the FA responses (rather than record it as a task failure in the

mnemonic discrimination task) and the underlying mechanism at both the encoding and test

phase of the trial (Cabeza, 2013; Hanczakowski & Mazzoni, 2011; H. Kim, 2011; Kubota et al.,

2006; Moritz et al., 2006; Okado & Stark, 2005), the differential processing that could

potentially be involved with correcting versus perpetuating FA behavior has not been

investigated.

In order to address these gaps in the literature, we developed a paradigm to investigate

both the impact of a false alarm on the original memory trace as well as the relative contributions

of hippocampal and cortical processes in promoting and recovering from a failed lure detection.

We hypothesized that (a) FAs would result in multiple memory traces that exist concurrently,

and as such, we predicted that participants would be able to distinguish target from lure when

both were presented simultaneously. Further, we hypothesized that subsequent target-trace

recovery would be predicted by (b) increased BOLD signal in CA3/DG during the orienting task

and false alarm, and (c) increased BOLD signal in a priori regions associated with encoding and

retrieval. Finally, we performed exploratory whole-brain analyses with the prediction that (d) we

would observe regions associated with higher-order cognitive processes, such as attention and

content processing, that were involved in subsequent target-trace recovery. In short, we predicted

that not all FA responses are equal, but that some would occur during better processing of the

Lure stimulus which would then precede a retrieval of the Target memory trace during a

subsequent test.

2   METHODS

2.1   Participants

Thirty-five participants (15 female, mean age = 23.1, SD = 2.2) were recruited from the

University and local community and were compensated for their participation with either

payment or a 3D-printed, quarter-scale model of their brain. The university's Institutional

Review Board approved the research, and all participants gave written informed consent prior to

participation. Inclusion criteria consisted of speaking English as a native language, being right-

hand dominant, and having normal or corrected-to-normal vision. Exclusion criteria consisted of

a history of psychologic disorder or brain injury, head dimensions that were too large for the 32-

channel coil, the existence of non-surgical metal in the participant's body (including non-

removable piercings), and color blindness. Participants were excluded from MRI analyses

separately for excess motion within a particular phase of the experiment as noted in the MRI

Data Preprocessing section below.

2.2   Mnemonic Discrimination Task

Stimuli for the task consisted of 200 pairs of perceptually and semantically similar

Target-Lure pairs that were randomly selected from those hosted at

https://github.com/celstark/MST. Each stimulus depicted an everyday object in the center of the

image with a white background, and one member of each set was randomly selected to be a

Target while the other was selected to be a Lure for each participant. Previous to this study,

approximately 900 Target-Lure pairs were rated for similarity by 35 individuals (who were

independent of the current study) on a 7-point Likert scale where a response of "1" indicated

"extremely dissimilar" and "7" indicated "extremely similar". Only items that received an

averaged similarity rating of 6.0 or higher were used in this task in order to ensure a sufficiently

large number of FA responses for functional analyses.

The behavioral task consisted of three phases (Study, Test1, and Test2), all of which were

performed while the participant was scanned using fMRI. Stimuli were displayed by means of an

MR-compatible LCD monitor placed at the head of the scanner and viewed using a mirror

mounted on the head coil. Responses were collected via a fiber optic button box. E-prime (v 2.0)

was used to control stimulus display and record behavioral responses. While in the scanner but

prior to engaging in the task, participants viewed a task instruction video; for consistency, the

same researcher (NMM) screened and debriefed all participants, responded to any questions

about the task, and operated the scanner. Three different instruction videos were shown, one

immediately prior to each respective phase of the task, in which explicit directions and examples

were given to the participant.

Study phase: In the initial encoding phase, participants were shown 200 images of

everyday objects and were instructed to indicate via button press whether each object is typically

encountered indoors or outdoors (Figure 0, left). Each stimulus was preceded by a fixation cross

for 500 ms, then presented for 3500 ms across one continuous block. For all phases of the

experiment, 16 blank trials were added in a pseudo-random fashion to create jitter in stimulus

timing, with care taken to ensure that each block contained an equal number of blank trials

(when applicable). These blank trials, in addition to the inter-stimulus fixation crosses, served as

the baseline condition in the single subject regression analysis (see MRI Data Pre-Processing).

Test1 phase: Participants were shown 100 Target stimuli (items identical to the original

stimuli in the Study phase), and 100 Lure stimuli (items which were similar to, but differed

visually from, the original items) one at a time in the center of the screen. Participants were

instructed to indicate via button press whether each stimulus was the "same" (an exact repeat, or

Target) or "different" (a related Lure) to the study item (Figure 0, middle). Stimuli were

presented in a randomized order for both type (Target/Lure) and for level of Target-Lure

similarity. The Test1 phase was conducted in two blocks of 100 stimuli each (50 Target, 50

Lure), where each stimulus was presented for 3500 ms and preceded by a fixation cross for 500

ms.

Test2 phase: In this phase, all stimuli from the Study phase (Targets), accompanied with

the similar item (Lure), were presented side-by-side to the participant in a two-alternative forced-

choice format. The stimuli were counterbalanced for whether the Target appeared on the Left or

Right (Figure 0, right). Participants were asked to indicate which of the two items they had seen

during the Study phase (Left or Right) and their confidence in their decision (Very or Somewhat)

via one of four potential response options (left very confident; left somewhat confident; right

somewhat confident; right very confident). This phase was conducted in four blocks of 50

stimuli in an attempt to combat participant fatigue. Each stimulus was presented for 5000 ms and

preceded by a fixation cross for 500 ms. During all phases of this experiment, participants were

instructed to respond while the stimulus was presented on the screen, and no feedback was given

relating to participant performance.

163
Figure 0. Mnemonic discrimination task. Left, during the study phase the participants made
indoor/outdoor judgements about a series of objects. Middle, during the first test phase,
participants were presented with either the same (Target) stimulus or one that was similar (Lure)
and were asked to decide whether the stimulus was the same or similar. Right, during the second
test phase, participants were presented with both Target and Lure stimuli simultaneously and
asked to indicate which item was originally presented during the Study phase. Additionally,
participants indicated the level of confidence (Very or Somewhat) in their decision and made
both their target and confidence decision with a single button press. For instance, if they decided
that the left stimulus was the Target and they were very confident, they pressed "1" whereas if
they were only somewhat confident the right stimulus was the Target, they pressed "3".

2.3 Statistical Analyses of Behavioral Data

Participant responses during the Study phase of the experiment were not analyzed for

indoor-outdoor accuracy as this question was merely meant to orient participants' attention to the

stimuli. During the second phase of the experiment, Test1, participants saw either Target or Lure

stimuli and made a forced-dichotomous judgment of whether the item was "Same" or

"Different". Correctly identifying the Target and Lure is termed Hit and Correct Rejection (CR),

respectively, and the incorrect identification is termed Miss and False Alarm (FA), respectively.

Here, we operationalize a false memory as the FA response to the lure stimulus. The third phase

of the experiment, Test2, was a two-alternative forced-choice where both Target and Lure

stimuli were presented and the participant was asked to identify the Target; consequently, only

Hits and FAs were possible.

Behavioral analyses were conducted on response proportions, considering the number of

items to which each participant responded. D-prime (d') scores were calculated for both Test1

and Test2 phases as an indicator of each participant's sensitivity to the stimuli. Test1 d' scores

were calculated the standard way (Equation 1; Wickens, 2002) where Z = z-score, $pHit$ =

proportion of Hits to the number of Targets, and pFA = proportion of FAs to the number of

Lures. As the Test2 phase presented both stimuli simultaneously, a correction was added to the d'

calculation (Equation 2; Wickelgren, 1968).

$$d'_{T1} = \ z(pHit) - \ z(pFA) \qquad\qquad d'_{T2} = \frac{z(pHit) - z(pFA)}{\sqrt{2}} \qquad\qquad \text{Equations (1, 2)}$$

Further, one-sample proportion testing using a continuity correction were used assess

whether participants were more likely to respond one way versus another (e.g., if participants

were more likely to Hit or Miss when presented with a Target, or more likely to respond Very

versus Somewhat confident during CRs). Outliers were detected according to the $1.5 \times IQR$

method. Paired and one-sample *t*-testing was used to compare the d' scores to each other and

against 0, respectively, utilizing an FDR correction to account for multiple comparisons. When

the exclusion of outliers differentially impacted the statistics, tests including and excluding

outliers are reported, otherwise reporting represents the exclusion of outliers.

2.4 MRI Acquisition

Functional and structural images were acquired on a Siemens TIM Trio 3T MRI scanner

utilizing the 32-channel head coil. Participants contributed a T1-weighted scan, a high-resolution

T2-weighted scan, and a series of functional T2*-weighted scans as described below. Standard-

resolution structural images were acquired using a T1-weighted magnetization-prepared rapid

acquisition with gradient echo (MP-RAGE) sequence with the following parameters: TR = 1900

ms, TE = 2.26 ms, flip angle = 9°, FoV = 218 × 250, voxel size = 0.97 × 0.97 × 1 mm, slices =

176 interleaved. High-resolution structural images were acquired using a T2-weighted sequence

with the following parameters: TR = 8020 ms, TE = 80 ms, flip angle = 150°, FoV = 150 × 150

mm, voxel size = 0.4 × 0.4 × 2 mm, slices = 30 interleaved. The T2 data were acquired

perpendicular to the long axis of the hippocampus. High-resolution multi-band echo-planar

images were acquired with a T2*-weighted pulse sequence with the following parameters: Multi-

band factor = 8, TR = 875 ms, TE = 43.6 ms, flip angle = 55°, FoV = 180 × 180 mm, voxel size

= 1.8 mm^3, slices = 72 interleaved. This sequence was aligned parallel with the long axis of the

hippocampus, and the first 11 TRs were discarded in order to allow for T1 equilibration.

2.5 MRI Data Pre-Processing

The pre-processing and visualization of structural and functional MRI data were

accomplished with the following software: dcm2nii (Li et al., 2016), Convert 3D Medical Image

Processing Tool (c3d) and ITK-SNAP (Yushkevich et al., 2006), Analysis of Functional

NeuroImages (AFNI; version 18.3.15; Cox, 2012), Advanced Normalization Tools software

(ANTs; Avants et al., 2008, 2011), Automatic Segmentation of Hippocampal Subfields (ASHS;

Yushkevich et al., 2015), and Multi-image Analysis GUI (Mango; Lancaster et al., 2012).

T1-weighted structural files were converted into 3D NIfTI files, rotated into functional

space (referencing the volume with the minimum calculated outlier voxels) using a rigid

transformation with a Local Pearson Correlation cost function. The rotated scan was then skull-

stripped and warped into MNI space via a non-linear diffeomorphic transformation.

High-resolution T2-weighted files were converted to 3D NIfTI files, and both T1w and

T2w files, in native space, were used to segment hippocampal subregions via ASHS, referencing

the UPenn atlas (Yushkevich et al., 2015). Hippocampal subregion masks from each participant

were then used to construct template priors via Joint Label Fusion (Figure 1; H. Wang et al.,

2013), and study-specific masks for the subiculum, CA1, and the combined CA2-CA3-DG

regions where constructed and resampled into functional dimensions. Voxels associated with

overlapping subregions were excluded to combat partial-voluming effects when down-sampling,

and the CA2, CA3, and DG masks were combined into a single mask due to functional resolution

constraints.

236
Figure 1. Hippocampal subregion masks. Left, a participant's T2-weighted scan segmented for
the medial temporal lobe via ASHS. Middle, template medial temporal lobe masks constructed in
MNI space via Joint Label Fusion. Right, the resampled masks, where overlapping labels were
excluded.

T2*-weighted DICOMs were first converted to 4D NIfTI files, and the volume from the

experiment with the smallest percentage of outlier voxels was extracted to serve as a volume

registration base. The calculations for moving each volume into the same space as the volume

registration base were then performed, using cubic polynomial interpolations, and the T1-

weighted file was also aligned with the volume registration base using a rigid transformation that

utilized a Local Pearson Correlation cost function. This rotated structural scan was then used to

produce normalization calculations via a non-linear transformation into MNI space. Movement

of the functional data into template space was then accomplished by concatenating the volume

registration calculations with the MNI transformation calculations, thereby moving all volumes

from native space to MNI space via a single transformation, resulting in less partial-voluming

and blurring of the functional data than if each transformation was applied separately. A series of

masks were then generated in order to remove volumes with missing data as well as determine

the intersection between functional and structural data, and the functional data were then scaled

by the mean signal. Finally, centered (demeaned) motion files for six degrees of freedom were

generated for each run of the experiment.

Behavioral data during each of the three phases (Study, Test1, and Test2) were coded

according to behavioral responses during that and subsequent phases of the experiment. Trials

from the Study phase were coded according to subsequent Test1 performance (i.e., Study

predicting Test1 activation or SpT1): subsequent Hit, subsequent FA, subsequent Miss, or

subsequent CR. Additionally, Study-phase trials were coded according to behavioral responses in

Test1 and Test2 (SpT1pT2). For example, a stimulus presented during Study might subsequently

receive a FA in Test1 and a Hit in Test2 (subsequent FA-Hit). Coding Study trials in this way

allows us to potentially localize differences during the encoding process that result in Test1 FAs

that are perpetuated into Test2 (subsequent FA-FA) compared to FAs that are corrected in Test2

(subsequent FA-Hit). It should be noted, however, that a low number of certain behaviors

(particularly Test1 Misses preceding Test2 behaviors) necessitated collapsing across CR and

Miss response bins in the deconvolution phase in order to avoid producing vectors for some

participants consisting entirely of zeros, vectors which are incompatible with deconvolution.

Such behavioral bins were not included in group-level analyses. Test1 trials were coded

according to four outcomes (Hit, FA, CR, and Miss). Responses during Test1 that preceded

Test2 (T1pT2) produced 4×2 possible outcomes since Test2 had two possible behaviors (Hit,

FA). Test2 responses that followed Test1 (T2fT1) also produced 4×2 possible outcomes, and

Test2 had two outcomes. Dividing all Test2 behavioral data according to confidence interval as

well as response type resulted in bins with trial counts that were too small for analyses, and as

such the confidence interval was dropped from functional analyses. Finally, trials where

participants failed to respond were coded separately from each of the above.

Vectors coding for the above behavioral outcomes were used in single-subject regression

models (Generalized Least Square) in which the canonical hemodynamic response was

convolved with a boxcar function with a duration equal to stimulus presentation. Modelling

included polynomial regressors accounting for scanner drift and scan run and regressors coding

for movement. The baseline condition consisted of collapsing across inter-stimulus intervals and

jittered blank stimuli. Volumes consisting of movement greater than 0.3° and/or more than 0.3

mm in any direction relative to the previous volume were discarded along with the previous

volume. Each deconvolution was assessed for the number of volumes that necessitated removal

and any participant with more than 10% of volumes dropped *and* 1.5 × IQR of volumes dropped

were excluded from subsequent analyses: one subject was removed from the Study phase, two

subjects were removed from the Test1 phase, and three subjects were removed from the Test2

phase. As such 34 participants were used to investigate Study performance, 33 for Test1, and 32

for Test2 performance.

Region of interest (ROI) specific masks were constructed to test *a priori* hypotheses. In

addition to the hippocampal subregion masks described above, clusters associated with memory

tasks were constructed by using the key words "Memory Encoding" and "Memory Retrieval" as

search terms in the NeuroSynth.org algorithm, which conducts a meta-analysis of PubMed

articles reporting the same key words and constructs a set of clusters derived from the reported

coordinates in said papers. The search term "Memory Encoding" yielded 124 articles, while

"Memory Retrieval" yielded 183. Association clusters (corrected for meta-analysis multiple

comparisons at the p = .01 level) were downloaded and thresholded at k=80 for the encoding

clusters, and k=100 for the retrieval clusters, which produced a set of binary masks that were

then resampled into functional dimensions. The "encoding" set contained four clusters,

consisting of the left hippocampus, right collateral sulcus, left inferotemporal gyrus, and right

amygdala (Figure 2). The "retrieval" set contained 10 clusters, consisting of the left retrosplenial,

angular gyrus, hippocampus, medial PFC, temporal-parietal junction, dorsal medial PFC, and the

right hippocampus, parietal-occipital sulcus, and posterior hippocampus (Figure 3). NeuroSynth

masks were then rotated and resampled into functional space and used to extract region-of-

interest (ROI) specific mean β-coefficients, using "encoding" masks to investigate volumes from

the Study phase, and "retrieval" masks for the Test1 and Test2 phases. Two-factor repeated

measures ANOVAs were used to investigate an ROI × behavior interaction for hippocampal

subregion and NeuroSynth masks, individually, where a Greenhouse-Geisser (GG) correction

was conducted for violations of sphericity. Any significant interaction surviving FDR multiple-

comparison adjustments were investigated via a one-way repeated measures ANOVA to

determine which ROIs had differential activation for the various behaviors, and subsequent

pairwise t-testing to determine how the β-coefficients for the behaviors differed within the ROI.

313
314   **Figure 2. NeuroSynth encoding masks.** A left hippocampus, B left inferotemporal gyrus, C
315   right collateral sulcus, D right posterior amygdala.

Figure 3. NeuroSynth retrieval masks. A bilateral posterior cingulate, B left posterior parietal, C left hippocampus, D left middle frontal gyrus, E left medial prefrontal, F left temporoparietal junction, G left dorsal medial prefrontal, H right hippocampus, I right parieto-occipital sulcus, and J right posterior hippocampus.

In preparation for exploratory group-level analyses, a gray matter mask was constructed in template space using Atropos priors (Tustison et al., 2014), which was multiplied by the intersection mask to produce a gray matter, intersection inclusion mask. Exploratory group-level analyses were then conducted on all voxels within this inclusion mask for each phase of the experiment using the Equitable Thresholding and Clustering method (ETAC; Cox, 2019) using the p-values 0.01, 0.005, and 0.001, blurs of 4, 6, and 8mm, nearest neighbors 1, and two-sided thresholding. Any surviving clusters were used to extract mean β-coefficients from participant data. All scripts used in this experiment can be found at https://github.com/nmuncy/STT.

3 RESULTS

3.1 Behavioral Analyses and Descriptive Statistics

Analyses of Test1 performance yield a mean d'_{T1} score of 0.66, indicating that successfully distinguishing between Targets and Lures was difficult, as the design of the experiment intended. However, participants were successful at detecting Targets and Lures in Test1, as the d'_{T1} differed significantly from zero ($t(34) = 11.3$, $p_{FDR} < .0001$, 95% CI [0.54, 0.78]). Response proportions were tested against chance (0.5) for each stimulus type; participants were more likely to Hit than Miss on Targets (estimate = .79, $\chi^2 = 32.18$, $p_{FDR} < .0001$, 95% CI [.69, .86]) while being equally likely to CR and FA on Lures (estimate = .43, $\chi^2 = 1.43$, $p_{FDR} = .41$, 95% CI [.33, .54]; Figure 1, left). On average, participants had 76 Hits, 54 FAs, 42 CRs, 20 Misses, and 6 non-responses in this phase. The 54 FA responses are of particular interest to the study: we designed the difficulty of the experiment to elicit approximately 60 FA responses in Test1 as two responses (Hit, FA) were possible in Test2, so subdividing Test1 FAs bins according to subsequent Test2 responses would yield sufficient trials to test statistically and model the HRF.

Behavioral Responses

Comparison

Figure 1. Boxplots of response proportions. Performance at chance results in a proportion of 0.5. Test1 (T1), Test2 (T2), and Test1 preceding Test2 (T1 p T2). "Hit-Very Rate" = likelihood of responding Very Confident versus Somewhat Confident during Hits. "T1 Hit p T2 Hit Rate" = proportion of Test1 Hits that precede a Test2 Hit versus Test2 FA. Outliers are included.

Analyses of Test2 performance yielded a d'$_{T2}$ of 0.45, which also differed significantly

from zero ($t(34) = 9.12$, $p_{FDR} < .0001$, 95% CI [.35, .55]), and indicated that participants were

more likely to Hit than FA (estimate = .62, $\chi^2 = 11.43$, $p_{FDR} = .002$, 95% CI [.55, .69]; Figure 1,

middle). In terms of confidence, participants were equally likely to indicate Very and Somewhat

confident during Test2 Hits (estimate = .53, $\chi^2 = 0.2$, $p_{FDR} = .7$, 95% CI [.43, .62]) and during

Test2 FAs (estimate = .38, $\chi^2 = 4.1$, $p_{FDR} = .09$, 95% CI [.27, .49]). Finally, testing the d' scores

for Test1 versus Test2 indicated that participants performed better during Test1 ($t(34) = 4.37$,

$p_{FDR} = .001$, 95% CI [.11, .31]). On average, participants had 121 Hits, 73 FAs, and 5 non-

responses.

Proportion testing of Test1 Hits and FAs which precede Test2 responses (T1pT2) reveal

that a Test1 Hit was more likely to result in a subsequent Test2 Hit than a FA (estimate = .72, χ^2

= 13.8, $p_{FDR} < .001$, 95% CI [.60, .82]), but that Test1 FAs were equally likely to result in either

a subsequent Test2 Hit or FA (estimate = .5, $\chi^2 \approx 0$, $p_{FDR} \approx 1$, 95% CI [.37, .64]; Figure 1, right).

These results, when viewed with respect to the Test2 results (d' which differs from 0 and a

tendency to detect the target), suggest that recovery from a Test1 FA is possible but not

guaranteed. It should be noted, though, that Test2 Target detection (particularly following a

Test1 Hit) may be driving some of the behavioral findings. Analyses of the BOLD response

during Test1 FAs that precede Hits versus FAs in Test2 time will help determine why some

Test1 FAs are perpetuated into Test2 while others are corrected, and if participants are indeed

performing differently from chance.

Test1 responses that preceded Test2 confidence responses consist of permutations of

Test1 behaviors, Test2 behaviors, and Test2 confidence responses. It was originally thought that

Test1-Test2 FA-Hit responses would yield higher confidence responses than FA-FA, which

could show interesting activation in the BOLD response. Paired t-testing revealed, however, an

equal number of "Very Confident" responses for both Hits and FAs in Test2 following a Test1

FA ($t(34) = 0.1$, $p = .9$, 95% CI [-2.3, 2.5]), indeed, participants were equally likely to select

"Very" and "Somewhat" during Test2 Hits and FAs following Test1 FAs (estimate = .4, $\chi^2 = $

0.7, $p_{FDR} = .53$, 95% CI [.23, .61], estimate = .4, $\chi^2 = 0.8$, $p_{FDR} = .53$, 95% CI [.22, .59],

respectively). Finally, parsing the data in such a fashion yielded bins that were too small to

reasonably model the hemodynamic response (e.g. Test1 FA preceding Test2 Hit-Very = 11 ±6

responses on average). As such, all subsequent analyses were collapsed across confidence

ratings.

3.2 Functional MRI Analyses

To address our research hypotheses, we conducted *a priori* analyses on both hippocampal

subregions and NeuroSynth memory regions, separately, as well as *a posteriori* exploratory,

whole-brain analyses. First, analyses investigated the effects of encoding on subsequent test

performance. The first analysis examined signal during the Study phase coded for subsequent

Test1 behavior, termed "SpT1" or "Study preceding Test1" (Figure 2, blue), while the second

analysis investigated the Study phase coded instead for trials which resulted in either a Test1 FA

that was then corrected (Test2 Hit) or perpetuated (Test2 FA), termed "SpT1pT2" (Figure 2,

purple). One participant was excluded from these analyses due to excessive movement during the

Study phase (n=34). Next, two analyses investigated the Test1 phase of the experiment. One

analysis investigated signal during Test1 associated with the various Test1 behaviors, termed

"T1", and the next analysis investigated Test1 FAs according to whether or not they would be

corrected in Test2, termed "T1pT2" (Figure 2, red). Two participants were excluded from these

analyses due to excessive movement in the Test1 phase (n=33). The final set of analyses

involved signal from the Test2 phase. First, we investigated signal during Test2 associated with

Test2 behaviors, termed "T2". The final analysis coded trials according to whether a Test2 Hit or

FA followed a Test1 FA, here termed "T2fT1" or "Test2 following Test1" (Figure 2, green).

Three participants were excluded from the last set of analyses due to excessive movement during

Test2 (n=32). In this way, the signal from each respective phase of the experiment is

investigated, but trials are coded for the current phase (T1 and T2), for future responses to

current stimuli (SpT1, SpT1pT2, T1pT2), or with respect to previous responses (T2fT1), thereby

allowing for the investigation of potential encoding and retrieval effects that differentially impact

the memory trace and attempted recovery thereof.

404
Figure 2. Schematic of analyses performed. The experiment consists of three phases, Study,
Test1, Test2. Study trials were coded for which stimuli preceded Test1 responses (SpT1, blue) as
well as the resulting response in Test2 following a Test1 response (SpT1pT2, purple). Test1
trials were analyzed both for the behavior which occurred in Test1 (not colored) as well as for
responses which preceded a Test2 response (T1pT2, red). Similarly, Test2 trials were analyzed
for Test2 behaviors (not colored) as well as for which Test1 behavior the Test2 response
followed (T2fT1).

3.2.1 Study trials preceding Test1 responses (SpT1). Study stimulus responses were

coded for subsequent Test1 behavior. For example, one stimulus in Study could elicit a Hit in

Test1 while a different stimulus could elicit a FA response.

For *a priori* analyses, a two-factor repeated measures MANOVA, where hippocampal

subregion ROIs (Left and Right Subiculum, CA1, and CA2-CA3-DG) served as within-subject

factor A and subsequent Test1 behavior (Hit, FA, CR) served as within-subject factor B, failed to

detect any significant interaction between the factors ($F(10, 330) = .89$, $\eta_G^2 < .001$, GGε = .45,

p(GG)$_{FDR}$ = .47). A similar analysis using NeuroSynth ROIs associated with encoding (Left

hippocampus, inferotemporal gyrus, and right amygdala, and collateral sulcus; see Figure 2) also

failed to detect a significant interaction between the factors ($F(6, 198) = 1.4$, $\eta_G^2 < .001$, GGε =

.38, p(GG)$_{FDR}$ = .24).

Exploratory whole-brain analyses revealed seven clusters (Figure 3, Table 1) which

significantly differed in Study volumes for subsequent Test1 Hits versus FAs (two subsequent

behaviors were investigated given that ETAC only supports t-test equivalent analyses). Three

clusters were more active preceding a Hit than a FA (left and right visual stream [Figure 3, A,

B], right intraparietal sulcus [C]) which are consistent with visual and attention networks. Four

clusters demonstrated the opposite pattern of activation (left and right angular gyrus [D, E], left

precuneus [F], and left dorsal medial PFC [G]), where the angular gyri and precuneus regions are

consistent with the default mode network.

Figure 3. SpT1 whole-brain results. Regions which demonstrated differential signal during SpT1. Hits versus FAs during an exploratory whole-brain analysis. A left visual stream, B right visual stream, C right intraparietal sulcus, D left angular gyrus, E right angular gyrus, F left precuneus, G left dorsal medial PFC.

Table 1

Phase	ROI	K	X	Y	Z	T(33)	Sig	LB	UB
SpT1	LVS	1649	-42.6	-74.3	-1.2	6.8	***	0.04	0.08
	RVS	2054	42.1	-72.4	-2.4	7.1	***	0.06	0.10
	RIPS	561	30.1	-75.2	33.7	5.3	***	0.04	0.09
	LAG	825	-47.8	-64.7	39.1	-6.1	***	-0.11	-0.05

	ROI	K	X	Y	Z	T(32)	Sig	LB	UB
	RAG	56	53.6	-64	30.6	-5.1	***	-0.14	-0.06
	LPPCU	1334	-6.3	-60.4	27.1	-6.7	***	-0.11	-0.06
	LDMPFC	269	-4.7	34.6	39.6	-7.7	***	-0.08	-0.05
T1	ROI	K	X	Y	Z	T(32)	Sig	LB	UB
	RPUT	46	25	-9	-22	4.51	***	0.066	0.174
	RMPFC	984	6	-3	-2	6.27	***	0.056	0.11
	RAG	917	52	-14	-40	7.37	***	0.053	0.093
	RAMTG	646	59	22	-40	6.03	***	0.05	0.101
	RPCU	174	4	-41	0	6.48	***	0.068	0.13
	LDMPFC	18	-3	-48	5	-5.23	***	-0.22	-0.097
	LINS	103	-33	-3	38	-6.2	***	-0.141	-0.071
	RIPS	321	28	-40	-18	-6.26	***	-0.11	-0.056
	RIFG	47	47	-25	-43	-6.64	***	-0.149	-0.079
T1pT2	ROI	K	X	Y	Z	T(32)	Sig	LB	UB
	LINS	98	31	-26	-3	4.7	***	0.05	0.12
	RSMG	19	-63	30	41	6.8	***	0.08	0.15
T2fT1	ROI	K	X	Y	Z	T(31)	Sig	LB	UB
	RAG	193	45	-72	18	4.95	***	0.05	0.12
	RPCU	813	2	-56	16	4.45	***	0.05	0.13

Cluster sizes and test statistics for the exploratory whole-brain analyses. ROI = region of interest, K = cluster size, X-Z = coordinate (peak), T(xx) = t-test(df), Sig = significant, *** = p < .0001, LB = 95% confidence interval lower bound, UB = upper bound. See text for ROI abbreviations.

3.2.2 Study trials preceding Test1 FAs preceding Test2 responses (SpT1pT2). Study trials were coded for whether or not an encoding stimulus which resulted in a Test1 FA was then corrected in Test2. For example, while two stimuli both may have resulted in a Test1 FA, the participant may have successfully identified one stimulus as the Target but failed to do so for the other stimulus in Test2. Such coding allowed us to investigate the potential role of encoding in recovering from interference in Test2. On average there were an equal number of encoding trials (26) that resulted in a T1 FA that were and were not corrected in Test2.

For the a priori analyses, individual two-factor repeated measures MANOVAs were used to investigate the interaction of ROI and behaviors for hippocampal subregion as well as NeuroSynth ROIs associated with encoding. ROIs were used as factor A in the model and factor B consisted of whether a FA would be corrected (Test1 FA, Test2 Hit) or perpetuated (Test1 FA, Test2 FA). No ROI × behavior interaction was detected in this phase of the experiment for either the hippocampal or NeuroSynth ROIs ($F(5, 165) = 1.4$, $\eta_G^2 < .001$, GGε = .72, $p = .23$; $F(3, 99) =$

2.7, $\eta_G^2 = .001$, GGε = .58, p(GG)$_{FDR}$ = .09, respectively). Further, exploratory whole-brain

analyses failed to detect any regions during the encoding phase that discriminated for whether a

subsequent Test1 FA would be corrected or perpetuated in Test2. Taken together, our data

suggest that while the neural activation during encoding trials had a significant impact on

subsequent Test1 performance, the activation during the encoding phase did not predict a

recovery from interference, as indicated by a Test1 FA preceding a Test2 Hit.

3.2.3 Test1 responses. Test1 trials were coded for the relative success of target and lure

detection, using the labels Hit, FA, CR, and Miss. A two-factor repeated measures MANOVA

detected a significant interaction of hippocampal subregions (left and right subiculum, CA1, and

CA2-CA3-DG) and Test1 behaviors (Hit, FA, CR; $F(10, 320) = 2.7$, $\eta_G^2 = .001$, GGε = .58,

p(GG)$_{FDR}$ = .03). One-way ANOVA testing revealed that left CA2-CA3-DG and CA1 were

driving this effect ($F(2, 64) = 8.28$, $\eta_G^2 = .01$, GGε = .96, p(GG) < .001; $F(2, 64) = 3.64$, $\eta_G^2 <$

.01, GGε = .86, p(GG) < .05, respectively). Further, post-hoc pairwise t-testing indicated that the

left CA2-CA3-DG was significantly more active during Hits than FAs and CRs ($t(32) = 2.16$, p <

.05, 95% CI [.004, .12]; $t(32) = 4.47$, p < .001, 95% CI [.06, .17], respectively) and that the left

CA1 was also more active during Hits than CRs ($t(32) = 3.06$, p < .01, 95% CI [.01, .06]; Figure

4).

Figure 4. Test1 hippocampal subfield results. The CA2-CA3-DG region is more active during Hits than FAs or CRs. The CA1 is more active during Hits than CRs.

Using the same analysis as above to investigate NeuroSynth masks associated with

retrieval (left angular gyrus, dorsal medial PFC, hippocampus, middle frontal gyrus, medial PFC,

retrosplenial cortex, temporo-parietal junction, and right hippocampus, parieto-occipital sulcus,

and posterior hippocampus; see Figure 3) revealed a significant mask × behavior interaction was

detected ($F(18, 576) = 13.15$, $\eta_G^2 = .01$, GG$\varepsilon = .36$, p(GG)$_{FDR} < .0001$). Subsequent one-factor

ANOVA testing revealed that five regions (Table 2, T1) were driving this effect: the left

temporo-parietal junction (LTPJ), ventral medial PFC (LVMPFC), dorsal medial PFC

(LDMPFC), middle frontal gyrus (LMFG), and angular gyrus (LAG). Post-hoc t-testing was

conducted for these regions in order to determine for which behaviors a difference in BOLD

signals occurred (Table 3, T1), which revealed two patterns of activity (Figure 5). First, the LTPJ

and LVMPFC were associated with greater BOLD signal during Hits than CRs; with such a

pattern of activity it appears that these two regions are involved in recognition aspects of

memory retrieval. Second, activity in the LDMPFC, LMFG, and LAG regions during CRs

demonstrated greater BOLD signal than either Hits or FAs, consistent with novelty detection in

memory systems. In sum, the effect found in these regions is driven largely by the differential

BOLD response during CRs versus Hits and FAs (Table 3, T1), and the effects appear to be

associated with recognition memory and novelty detection.

492
493 Table 2

	ROI	F(2,64)	η_G^2	GGe	Sig
	LTPJ	6.9	0.006	0.82	**
	LVMPFC	7.3	0.016	0.81	**
T1	LDMPFC	22.5	0.029	0.72	***
	LMFG	10.9	0.011	0.80	***
	LAG	13.9	0.012	0.81	***
	ROI	F(2,64)	η_G^2	GGe	Sig
	LDMPFC	20.4	0.025	0.7	***
T1pT2	LAG	17.9	0.02	0.9	***
	LMFG	9.6	0.01	0.86	***
	ROI	F(2,62)	η_G^2	GGe	Sig
	LAG	16.2	0.023	0.96	***
	LDMPFC	12.2	0.019	0.95	***
T2fT1	LMFG	10.7	0.017	0.91	**
	LPCU	11.8	0.019	0.88	***
	LTPJ	15.1	0.022	0.87	***
	RPOS	13.9	0.023	0.87	***

494 NeuroSynth ROI ANOVA statistics. T1 = Test1,T1pT2 = Test1 preceding Test2, T2fT1 = Test2
495 following Test1. GGe = Greenhouse-Geisser epsilon. ** = p < .01, *** = p < .001.

496
497
498 Table 3

	Contrast	Hit-FA				Hit-CR				FA-CR			
	ROI	T(32)	Sig	LB	UB	T(32)	Sig	LB	UB	T(32)	Sig	LB	UB
	LTPJ	3.65	***	0.03	0.10	3.37	**	0.03	0.13	0.72	n.s.	-0.04	0.07
T1	LVMPFC	2.22	*	0.00	0.08	3.60	**	0.04	0.15	1.87	n.s.	0.00	0.12
	LDMPFC	-0.21	n.s.	-0.04	0.03	-5.62	***	-0.21	-0.10	-4.72	***	-0.22	-0.09
	LMFG	-0.14	n.s.	-0.02	0.02	-3.98	***	-0.09	-0.03	-3.39	**	-0.09	-0.02
	LAG	-1.22	n.s.	-0.02	0.01	-4.44	***	-0.07	-0.03	-3.64	***	-0.07	-0.02
	Contrast	FpH-FpF				FpH-CpH				FpF-CpH			
	ROI	T(32)	Sig	LB	UB	T(32)	Sig	LB	UB	T(32)	Sig	LB	UB
T1pT2	LDMPFC	2.10	*	0.00	0.08	-4.28	***	-0.23	-0.08	-5.13	***	-0.27	-0.12
	LAG	0.99	n.s.	-0.01	0.03	-4.36	***	-0.08	-0.03	-5.21	***	-0.09	-0.04

	Contrast	HfF-FfF				HfF-HfC				FfF-HfC			
	LMFG	0.22	n.s.	-0.02	0.03	-3.49	**	-0.09	-0.02	-3.45	**	-0.10	-0.02
	Contrast	HfF-FfF				HfF-HfC				FfF-HfC			
	ROI	T(31)	Sig	LB	UB	T(31)	Sig	LB	UB	T(31)	Sig	LB	UB
T2fT1	LAG	-1.00	n.s.	-0.08	0.03	-5.84	***	-0.14	-0.07	-3.33	**	-0.13	-0.03
	LDMPFC	-0.07	n.s.	-0.03	0.03	-5.42	***	-0.10	-0.04	-4.49	***	-0.10	-0.04
	LMFG	0.42	n.s.	-0.03	0.04	-4.14	***	-0.11	-0.04	-3.64	***	-0.13	-0.04
	LPCU	0.50	n.s.	-0.04	0.06	-4.91	***	-0.17	-0.07	-4.24	***	-0.19	-0.07
	LTPJ	0.86	n.s.	-0.03	0.07	-3.87	***	-0.15	-0.05	-4.21	***	-0.17	-0.06
	RPOS	-0.40	n.s.	-0.06	0.04	-5.02	***	-0.19	-0.08	-3.75	***	-0.20	-0.06

499   NeuroSynth ROI post-hoc t-tests corresponding to Table 2. * = p < .05, n.s. = p > .05.

501
Figure 5. Test1 NeuroSynth results. Five regions (left temporo-parietal junction [LTPJ],
ventral medial prefrontal cortex [LVMPFC], dorsal medial prefrontal cortex [LDMPFC], middle
frontal gyrus [LMFG], angular gyrus [LAG]) differentiate between Hits, FAs, and CRs. The
LTPJ and LVMPFC demonstrate significantly greater activity during Hits than CRs, while the
LDMPFC, LMFG, and LAG show the opposite pattern of activity.

Exploratory analyses detected nine clusters (Figure 6) which demonstrated differential

activation according to whether a Hit versus a FA was occurring (Table 1, Figure 6). Five

regions had a greater BOLD response during Hits all of which were located in the right

hemisphere: the putamen (RPUT, A), medial PFC (RMPFC, B), angular gyrus (RAG, C),

anterior middle temporal gyrus (RAMTG, D), and precuneus (RPCU, E). These findings are

largely consistent with known patterns of activity in the default mode network associated with

successful retrieval during episodic memory tasks (Figure 6, B-E). Four regions demonstrated

the opposite pattern of BOLD response: left dorsal medial PFC (LDMPFC, F), insular cortex

(LINS, G), right intraparietal sulcus (RIPS, H), and inferior frontal gyrus (RIFG, I). These

regions have been associated with dorsal and ventral networks which attend to self-referential

and saliency processing, respectively, in episodic memory tasks (Figure 6, F-H) as well as

attention networks (Figure 6, I).

520
Figure 6. Test1 whole-brain results. Regions which demonstrated differential signal during
Test1 Hits versus FAs during an exploratory whole-brain analysis. A right putamen, B right
medial PFC, C right angular gyrus, D right anterior middle temporal gyrus, E right precuneus, F
left dorsal medial PFC, G left insula, H right intraparietal sulcus, I right inferior frontal gyrus.

3.2.4 Test1 FA responses preceding Test2 responses (T1pT2). Test1 FA responses

were coded according to whether or not they would be corrected in Test2 (Hit versus FA),

resulting in the same number of trials per bin as in Section 3.2.2. This was done in order to

investigate any potential processes occurring during the FA that may allow for subsequent

recovery from such interference. Additionally, Test1 CR trials which preceded a Hit were

included in the analysis in order to investigate processes which may differ during Test1 Lure

trials (FA versus CR) that precede the same Test2 performance (Hit).

A two-factor repeated measures MANOVA failed to detect a significant interaction of

hippocampal subregions and beta-coefficients ($F(10, 320) = 0.97$, $\eta_G^2 < .001$, GGε = .69,

$p(GG)_{FDR} = .45$). However, an investigation into NeuroSynth regions associated with retrieval

showed a significant interaction ($F(18, 576) = 7.28$, $\eta_G^2 = .008$, GGε = .42, $p(GG)_{FDR} < .0001$).

Subsequent one-factor ANOVA testing revealed that three ROIs were driving this effect (Table

2, T1pT2; Figure 7): the left dorsal medial PFC (LDMPFC), angular gyrus (LAG), and middle

frontal gyrus (LMFG). The LDMPFC differed significantly in signal across all behaviors, and of

particular interest had stronger signal during a Test1 FA that preceded a Test2 Hit than a Test2

FA (Table 3, T1pT2). The LAG and LMFG did not differ among the Test1 FA responses but

showed stronger signal for Test1 CR preceding a Hit than either Test1 FA response. These

results suggest that while all three regions are particularly important for Test1 performance in

terms of CRs versus FAs, only the LDMPFC had differential activity during Test1 FAs that

predicted subsequent Test2 performance.

546
Figure 7. T1pT2 NeuroSynth results. Three regions (left dorsal medial prefrontal cortex
[LDMPFC], angular gyrus [LAG], middle frontal gyrus [LMFG]) demonstrated differential
activity during Test1 behaviors (FA, CR) preceding Test2 Hits and FAs. FpH = FA preceding
Hit, FpF = FA preceding FA, CpH = CR preceding Hit.

551   Exploratory whole-brain analyses of Test1 FA trials that precede a Test2 Hit versus FA

552   implicated two regions as being differentially engaged (Figure 8): the left insular cortex (LINS)

553   and right supramarginal gryus (RSMG). Both regions had greater BOLD signal preceding a Hit

554   than a FA (Figure 8, Table 3).

Figure 8. T1pT2 whole-brain results. Clusters which demonstrate differential signal during Test1 FAs that precede a Test2 Hit versus FA. A left insular cortex, B right supramarginal gyrus. FpH = FA preceding Hit, FpF = FA preceding FA.

3.2.5 Test2 responses. A two-factor repeated measures MANOVA failed to detect a

significant interaction of hippocampal subregions and Test2 behaviors (Hit, FA; $F(5, 155) =$

1.04, $\eta_G^2 < .001$, GG$\varepsilon = .75$, p(GG)$_{FDR} = .39$) or among NeuroSynth ROIs associated with

retrieval ($F(9, 279) = .19$, $\eta_G^2 < .001$, GG$\varepsilon = .51$, p(GG)$_{FDR} = .99$). Further, and unexpectedly, an

exploratory whole-brain analysis failed to detect any regions which differentially engaged in Hits

versus FAs.

3.2.6 Test2 responses following Test1 FAs (T2fT1). Test2 responses (Hit, FA)

following a Test1 FA were coded in order to investigate any potential processes during the

second testing phase involved in overcoming interference resulting from a Test1 FA. Further,

Test2 Hits following a Test1 CRs were also investigated in order to determine whether any

processes were differentially involved in Test2 Hits with respect to whether the Hit was preceded

by an incorrect or correct Test1 Lure identification (FA, CR). On average there were 27 Hits

following a CR.

A two-factor repeated measures MANOVA failed to detect a significant interaction of

hippocampal subregions and behavioral outcomes during the behavioral coding described above

($F(10, 310) = 1.1$, $\eta_G^2 = .001$, GG$\varepsilon = .53$, p(GG)$_{FDR} = .39$). An investigation in NeuroSynth

masks associated with Retrieval, however, showed a significant interaction in the same analysis

($F(18, 558) = 4.68$, $\eta_G^2 = .004$, GG$\varepsilon = .48$, p(GG)$_{FDR} < .0001$), and subsequent one-way ANOVA

analyses indicated that six regions were driving this effect (left angular gyrus, dorsal medial

PFC, middle frontal gyrus, precuneus, temporoparietal junction, and right parietooccipital sulcus;

Table 3, T2fT1). Greater signal was detected during Test2 in all regions for Hits following a

Test1 CR than for Hits following a Test1 FA (Figure 9), while no difference was detected in the

same ROIs for Hits versus FAs following a Test1 FA (Table 3, T2fT1). As such, we do not

detect any compensatory processes in Test2 in these ROIs associated with overcoming

interference resulting from a Test1 FA, as would be implicated by differential signal in a region

during Test2 Hits versus FAs following a Test1 FA. Further, we find the difference between

Test2 Hits according to whether they were preceded by a Test1 FA or CR to be interesting; as

Test2 presents both the Target and Lure simultaneously, increased activity during a Hit following

a CR but not a FA may be indicative of processes associated with selecting between two salient

memory representations. Indeed, in Test2 Hits following a Test1 CR, the participant (a) encoded

the Target sufficiently to be able to (b) correctly identify the Lure. When both are presented

simultaneously and the participant is tasked with identifying the Target, the participant may have

to resolve interference resulting from two competing representations. Finally, exploratory whole-

brain analyses did not detect any regions that differed significantly between their BOLD signal

associated with Test2 Hits versus FAs following a Test1 FA.

Figure 9. T2fT1 NeuroSynth results. Six regions (left angular gyrus [LAG], dorsal medial
prefrontal cortex [LDMPFC], middle frontal gyrus [LMFG], precuneus [LPCU], temporoparietal
junction [LTPJ], right parietooccipital sulcus [RPOS]) demonstrated differential activity during
Test2 behaviors (Hit, FA) following Test1 responses (FA, CR). HfF = Hit following FA, FfF =
FA following FA, HfC = Hit following CR.

4   DISCUSSION

This experiment set out to investigate the impact of a false memory response on the

original memory representation, and to determine what, if any, neural processes were associated

with recovering the original memory trace. To this end, we employed a Study-Test1-Test2

paradigm within an MRI scanner, in which participants were trained on a series of images

(Study), tasked with identifying either Targets or similar Lures presented individually (Test1),

and then tasked with identifying the Target when both Targets and Lures were presented

simultaneously (Test2). False Alarm (FA) responses in Test1 were operationalized as false

memories and were used to investigate whether participants could correctly identify the Target

(Hit) in Test2 following a Test1 FA. Briefly, we hypothesized (a) that a FA would write a

memory trace of the Lure without necessarily overwriting the Target's trace such that the two

representations would be dissociable, and that (b) the CA3/DG regions of the hippocampus as

well as (c) encoding and retrieval nodes and (d) higher-order executive processes would be

integral to the formation of a second, dissociable representation.

As the paradigm anticipated, participants made a large number of FA responses in Test1,

and proportion testing revealed that participants were equally likely to correct (Hit) and

perpetuate (FA) the mistake in Test2 (Figure 1). That is, a FA does not unambiguously overwrite

the Target memory trace, as would be indicated by a greater likelihood to perpetuate the FA

response in Test2, nor does the FA necessarily write multiple representations from which the

Target is dissociable. Given that participants were indeed successful in Test2 (Section 3.1), the

behavioral data is ambiguous as to the effect of the FA on the Target trace, and so from the

behavioral analyses alone we are unable to reject the null of hypothesis (a). The functional data,

however, shed some light on why some FA responses are corrected and others perpetuated.

Functional analyses both replicate and reveal several aspects of memory. First, consistent

with literature on subsequent memory and forgetting (Cabeza et al., 2008; H. Kim, 2011; Shrager

et al., 2008), seven regions were differentially involved during the encoding phase according to

test performance (SpT1). Three regions, the bilateral visual stream and right intraparietal sulcus

(Figure 3, A-C), showed greater BOLD signal preceding a Test1 Hit versus a Test1 FA,

indicative of increased visual processing and attention, respectively. Conversely, three regions,

the bilateral angular gyri and left precuneus (Figure 3, D-F), had the opposite pattern of activity,

which is consistent with increased default mode network activity during encoding preceding an

erroneous response during test. The final region, the dorsal medial PFC region (Figure 3, G), also

demonstrated greater BOLD signal preceding a FA versus Hit and appears to be associated with

recognition memory (Phillips et al., 2009; Suzuki et al., 2009); as our stimulus set consists of

everyday objects, increased familiarity with certain objects may result in top-down interference

thereby resulting in poorer encoding. As such, suppressed signal in this region during encoding

trials for stimuli that precede a Hit may be indicative of decreased familiarity with the stimulus

which allowed for greater encoding and subsequent successful target identification.

Second, in addition to confirming that hippocampal subfields and NeuroSynth retrieval

clusters demonstrated differential signal during this retrieval task (Figure 4, Figure 5), multiple

regions were detected in an exploratory analysis of the Test1 phase of the experiment that were

consistent with the retrieval literature: four regions in the right hemisphere, consisting of the

medial PFC, angular gyrus, anterior middle temporal gyrus, and precuneus (Figure 6, B-E), all of

which are again consistent with the default mode network, were differentially engaged in the task

such that they demonstrated greater BOLD signal during Hits relative to FAs. These findings

replicate previous results (H. Kim, 2010; Sestieri et al., 2011) which have demonstrated that the

default mode network is involved in self-referential processing, an aspect of which is episodic

memory retrieval. Additionally, regions associated with saliency processing and attention

networks (left anterior insula, right intraparietal sulcus, right inferior frontal gyrus; Figure 6, G-I)

had greater signal during FA responses relative to Hits. It appears that such patterns of activation

are the result of the unsuccessful attempt to overcome significant interference resulting from the

highly similar Lures used in this task. Indeed, the right intraparietal sulcus is particularly

interesting as it is implicated in post-retrieval processes, where hippocampal information is

integrated with executive orchestration (Cabeza et al., 2008; Gilmore et al., 2015; Hao et al.,

2018; Sestieri et al., 2011; Uncapher & Wagner, 2009).

Third, and importantly, differential signal was detected during Test1 FAs according to

whether or not the FA was perpetuated into Test2 (T1pT2). Three regions, the left dorsomedial

PFC, left anterior insula, and right supramarginal gyrus (Table 3, Figure 7, Figure 8),

demonstrated significantly greater BOLD signal during Test1 FAs the preceded a Test2 Hit

versus a Test2 FA. The dorsomedial PFC region was derived from a NeuroSynth search term of

"Memory Retrieval" and is often implicated in the retrieval of episodic memory while the left

anterior insular cortex and right supramarginal gyrus are themselves associated with saliency and

bottom-up attention networks (Cabeza et al., 2008; Euston et al., 2012; Menon & Uddin, 2010;

Seeley et al., 2007; Shirer et al., 2012). These findings, then, suggest that increased activity in

these regions during a Test1 FA may be associated with better retrieval of the original memory

trace (LDMPFC) and attention to the relevant details of the stimulus (LINS, RSMG) which,

while subthreshold for Lure detection, nevertheless led to the formation of a Lure representation

such that the Target and Lure memory traces were subsequently dissociable in Test2.

Correspondingly, no dissociable pattern of activity was detected during the encoding phase that

predicted which Test1 FAs would be corrected in Test2 (SpT1pT2), nor is any such pattern

detected in Test2 when comparing Hits to FAs for stimuli following a Test1 FA (T2fT1) as

would be expected if such resolution of Target-Lure interference was only a function of the

quality of Target encoding (SpT1pT2) or retrieval processes used to dissociate Target and Lure

representations (T2fT1). Instead, our data suggest that the capacity to recover the Target trace

following a FA is entirely dependent on processes occurring *during* the FA response, and not

before or after, where greater attention and retrieval processes result in a dissociable Lure trace.

As such, and with respect to the behavioral data, the rejection of hypothesis (a) may or may not

be warranted; it appears that multiple representations *can* occur during a FA but we do not

demonstrated that multiple representations *will* occur, as the probability appears to dependent on

retrieval and attention processes during Test1. We do, however, reject the null of hypotheses (c)

and (d).

Some results of this study were surprising. First, participants were equally likely to Hit or

FA in Test2 to stimuli that had previously resulted in a Test1 FA (Figure 1, right), which is to

say that their Test2 behavior for these trials was not significantly different from chance. While

the d' score of Test2 differed significantly from 0, this was likely driven by the large number of

Test2 Hits following a Test1 Hit. Analyses of confidence ratings indicate that participants may

have been less confident in the Test2 FA responses (Figure 1, middle); note, though, that while

the confidence intervals of the proportion test do not include 0.5, indicating a difference from

chance, the FDR-corrected p-value is not significant at the predetermined α. Having confidence

intervals that do not include 0.5 gives some indication that participants were not responding with

FAs and Hits in Test2 at random. Regardless of the responses at varying confidences, however,

an equal likelihood to Hit or FA in Test2 following a Test1 Hit undermines subsequent fMRI

analyses in that any differential signal detected may simply be the capitalization on noise and

confirmation bias. That said, such a result may very well be a function of the experimental

design itself: a large number of difficult Target-Lure pairs were utilized in order to elicit a

sufficient number of Test1 FA responses such that dividing the number of Test1 FA responses by

subsequent behavior would still have sufficient trials to model the HRF, resulting in a task that

lasted 1.5 hours. Such an ambiguous result may very well therefore be the result of participant

fatigue in a long, arduous task. It is interesting, however, that the response proportion did not

approach 0, which would be indicative of a greater tendency for the Test1 FA to overwrite the

original representation. It is our interpretation, then, that the participants were indeed engaged in

the task and not performing at random given the Test2 d' and a decreased confidence in Test2-

FA responses. As such, a false alarm may or may not overwrite the original representation,

according to processes occurring during the initial false alarm (Section 3.2.4). Somewhat

relatedly, another minor limitation is the fact that participants performed better during the Test1

phase than Test2, even though it has been documented that a two-alternative forced-choice task

is easier than a Yes/No (Jou et al., 2016). We interpret this fact as evidence of participant fatigue.

A second unexpected result was the lack of findings within the hippocampal subfields in

all analyses save those of Test1 (Section 3.2.3). While we demonstrate the differential roles of

left CA1 and CA3-CA3-DG in this task with respect to Hits and CRs, in which the hippocampus

appeared to be biased towards Target detection in this task, the lack of subfield findings in other

analyses of the task was surprising to us give the wealth of literature that has well established the

roles of these regions in similar tasks (Bakker et al., 2008; J. Kim & Yassa, 2013; Kirwan &

Stark, 2007; Lacy et al., 2011; Rolls & Kesner, 2016). Accordingly, we cannot reject the null of

hypothesis (b) as we did not detect differential subregion processing during the FA that predicted

subsequent performance. These null results are perhaps a type II or false negative error which

resulted from extracting a single parameter estimate for the entire ROI, thereby collapsing across

a significant amount of noise as well as signal, possibly washing out the effect. Indeed, it has

been noted that the long-axis of the hippocampus contains a functional gradient and discrete

domains (Strange et al., 2014), so collapsing across all variance along the anterior-posterior axis

of the hippocampal subregions may be inappropriate. Similarly, our contrast of Test1 FA signal

coded for subsequent Test2 performance may have simply failed to detect a difference between

those two conditions in terms of subregion BOLD signal, but this does not necessarily imply that

the hippocampus is not differentially processing the lures such that subsequent behavior results

in a Hit versus FA. Rather, the processes involved in such a computation could be rather similar

to one another and so a comparison of the mean subfield BOLD signal associated with each

process may again be simply too insensitive.

Third, we were surprised by the lack of significant differences for the fMRI data of the

Test2 phase. The behavioral data indicate that successful Target detection was possible, and that

participants were more likely to Hit in Test2 following a Test1 Hit, so performance was not at

chance. The lack of findings may not be unwarranted, however, as there are very few studies

reporting positive results from two-alternative, forced choice, recognition memory paradigms

using fMRI. One notable exception by Olsen and colleagues (2009) is somewhat similar to this

experiment in terms of participants and power. Olsen et al. (2009) demonstrated that the medial

temporal region was active while maintaining the memory of a set of faces during a delay period

prior to a two-alternative forced choice test, an area similar to where we expected activation to

occur. The lack of findings in the present experiment may be due to differences between the

experiments in fMRI processing methods. One stark difference is in the methods modelling both

the noise and auto-correlation function at the group-analysis step employed in this experiment, as

this improvement is more conservative than previous random-field based models in terms of

false discovery rates but was only recently developed (Cox, 2018, 2019; Eklund et al., 2016).

Furthermore, while significant differences were not detected within the fMRI data at the group

level, one should not assume that "[i]f the result is not statistically significant, then it proves that

no effect or difference exists" (Chen et al., 2017, p. 955). Indeed, as Chen et al. (2017) points

out, null results may very well be the result of sub-threshold signal and/or a lack of differential

signal between our chosen contrasts at our resolution.

In conclusion, our data suggest that a false alarm does not necessarily overwrite the

original memory trace of the target, but that subsequent detection of the target is possible,

suggesting that a false alarm results in multiple memory representations in some instances.

Further, the successful recovery of the target trace following a false alarm appears to be

dependent not on the initial encoding trial or on retrieval processes following the false alarm, but

on processes which occurred during the false alarm itself. Differential activation was detected in

the left dorsal medial prefrontal cortex, left insular cortex, and right supramarginal gyrus during

the false alarm that predicted whether or not the target would subsequently be retrieved, regions

which are associated with episodic memory, saliency, and bottom-up attention, respectively.

References

Avants, B. B., Epstein, C. L., Grossman, M., & Gee, J. C. (2008). Symmetric diffeomorphic image registration with cross-correlation: Evaluating automated labeling of elderly and neurodegenerative brain. *Medical Image Analysis*, *12*(1), 26–41.

Avants, B. B., Tustison, N. J., Song, G., Cook, P. A., Klein, A., & Gee, J. C. (2011). A reproducible evaluation of ANTs similarity metric performance in brain image registration. *NeuroImage*, *54*(3), 2033–2044.

Bakker, A., Kirwan, C. B., Miller, M., & Stark, C. E. (2008). Pattern separation in the human hippocampal CA3 and dentate gyrus. *Science*, *319*. https://doi.org/10.1126/science.1152882

Cabeza, R. (2013). Introduction to the special issue on functional neuroimaging of episodic memory. *Neuropsychologia*, *51*(12), 2319–2321.

Cabeza, R., Ciaramelli, E., Olson, I. R., & Moscovitch, M. (2008). The parietal cortex and episodic memory: An attentional account. *Nature Reviews Neuroscience*, *9*(8), 613.

Chen, G., Taylor, P. A., & Cox, R. W. (2017). Is the statistic value all we should care about in neuroimaging? *NeuroImage*, *147*, 952–959. https://doi.org/10.1016/j.neuroimage.2016.09.066

Cox, R. W. (2012). AFNI: What a long strange trip it's been. *NeuroImage*, *62*(2), 743–747.

Cox, R. W. (2018). Equitable thresholding and clustering. *BioRxiv*, 295931.

Cox, R. W. (2019). Equitable thresholding and clustering (ETAC): A novel method for fMRI clustering in AFNI. *Brain Connectivity*. https://doi.org/10.1089/brain.2019.0666

Doxey, C. R., & Kirwan, C. B. (2015). Structural and functional correlates of behavioral pattern separation in the hippocampus and medial temporal lobe. *Hippocampus*, *25*(4), 524–533.

Eklund, A., Nichols, T. E., & Knutsson, H. (2016). Cluster failure: Why fMRI inferences for

    spatial extent have inflated false-positive rates. *Proceedings of the National Academy of*

    Sciences, *113*(28), 7900–7905. https://doi.org/10.1073/pnas.1602413113

Euston, D. R., Gruber, A. J., & McNaughton, B. L. (2012). The role of medial prefrontal cortex

    in memory and decision making. *Neuron*, *76*(6), 1057–1070.

    https://doi.org/10.1016/j.neuron.2012.12.002

Gilbert, P. E., & Kesner, R. P. (2006). The role of the dorsal CA3 hippocampal subregion in

    spatial working memory and pattern separation. *Behavioural Brain Research*, *169*(1),

    142–149.

Gilbert, P. E., Kesner, R. P., & DeCoteau, W. E. (1998). Memory for spatial location: Role of the

    hippocampus in mediating spatial pattern separation. *Journal of Neuroscience*, *18*(2),

    804–810.

Gilbert, P. E., Kesner, R. P., & Lee, I. (2001). Dissociating hippocampal subregions: A double

    dissociation between dentate gyrus and CA1. *Hippocampus*, *11*(6), 626–636.

Gilmore, A. W., Nelson, S. M., & McDermott, K. B. (2015). A parietal memory network

    revealed by multiple MRI methods. *Trends in Cognitive Sciences*, *19*(9), 534–543.

    https://doi.org/10.1016/j.tics.2015.07.004

Goebel, P. M., & Vincze, M. (2007). Implicit modeling of object topology with guidance from

    temporal view attention. *Proceedings of the International Cognitive Vision Workshop,*

    ICVW.

Gold, A. E., & Kesner, R. P. (2005). The role of the CA3 subregion of the dorsal hippocampus in

    spatial pattern completion in the rat. *Hippocampus*, *15*(6), 808–814.

Grady, C. L., & Ryan, J. D. (2017). Age-related differences in the human hippocampus: Behavioral, structural and functional measures. In *The Hippocampus from Cells to Systems* (pp. 167–208). Springer.

Hanczakowski, M., & Mazzoni, G. (2011). Both differences in encoding processes and monitoring at retrieval reduce false alarms when distinctive information is studied. *Memory*, *19*(3), 280–289.

Hao, X., Wang, X., Song, Y., Kong, X., & Liu, J. (2018). Dual roles of the hippocampus and intraparietal sulcus in network integration and segregation support scene recognition. *Brain Structure & Function*, *223*(3), 1473–1485. https://doi.org/10.1007/s00429-017-1564-2

Hunsaker, M. R., & Kesner, R. P. (2013). The operation of pattern separation and pattern completion processes associated with different attributes or domains of memory. *Neuroscience & Biobehavioral Reviews*, *37*(1), 36–58.

Hunsaker, M. R., Rosenberg, J. S., & Kesner, R. P. (2008). The role of the dentate gyrus, CA3a, b, and CA3c for detecting spatial and environmental novelty. *Hippocampus*, *18*(10), 1064–1073.

Jou, J., Flores, S., Cortes, H. M., & Leka, B. G. (2016). The effects of weak versus strong relational judgments on response bias in two-alternative-forced-choice recognition: Is the test criterion-free? *Acta Psychologica*, *167*, 30–44. https://doi.org/10.1016/j.actpsy.2016.03.014

Kesner, R. P. (2007a). A behavioral analysis of dentate gyrus function. *Progress in Brain Research*, *163*, 567–576.

Kesner, R. P. (2007b). Behavioral functions of the CA3 subregion of the hippocampus. *Learning & Memory*, *14*(11), 771–781.

Kim, H. (2010). Dissociating the roles of the default-mode, dorsal, and ventral networks in episodic memory retrieval. *NeuroImage*, *50*(4), 1648–1657. https://doi.org/10.1016/j.neuroimage.2010.01.051

Kim, H. (2011). Neural activity that predicts subsequent memory and forgetting: A meta-analysis of 74 fMRI studies. *NeuroImage*, *54*(3), 2446–2461.

Kim, J., & Yassa, M. A. (2013). Assessing recollection and familiarity of similar lures in a behavioral pattern separation task. *Hippocampus*, *23*(4), 287–294.

Kirwan, C. B., Hartshorn, A., Stark, S. M., Goodrich-Hunsaker, N. J., Hopkins, R. O., & Stark, C. E. (2012). Pattern separation deficits following damage to the hippocampus. *Neuropsychologia*, *50*(10), 2408–2414.

Kirwan, C. B., & Stark, C. E. (2007). Overcoming interference: An fMRI investigation of pattern separation in the medial temporal lobe. *Learn Mem*, *14*. https://doi.org/10.1101/lm.663507

Kubota, Y., Toichi, M., Shimizu, M., Mason, R. A., Findling, R. L., Yamamoto, K., & Calabrese, J. R. (2006). Prefrontal hemodynamic activity predicts false memory—A near-infrared spectroscopy study. *Neuroimage*, *31*(4), 1783–1789.

Lacy, J. W., Yassa, M. A., Stark, S. M., Muftuler, L. T., & Stark, C. E. (2011). Distinct pattern separation related transfer functions in human CA3/dentate and CA1 revealed using high-resolution fMRI and variable mnemonic similarity. *Learning & Memory*, *18*(1), 15–18.

Lancaster, J. L., Laird, A. R., Eickhoff, S. B., Martinez, M. J., Fox, P. M., & Fox, P. T. (2012). Automated regional behavioral analysis for human brain images. *Frontiers in Neuroinformatics*, *6*, 23.

Leal, S. L., & Yassa, M. A. (2014). Effects of aging on mnemonic discrimination of emotional information. *Behavioral Neuroscience*, *128*(5), 539.

Leutgeb, J. K., Leutgeb, S., Moser, M.-B., & Moser, E. I. (2007). Pattern separation in the dentate gyrus and CA3 of the hippocampus. *Science*, *315*(5814), 961–966.

Li, X., Morgan, P. S., Ashburner, J., Smith, J., & Rorden, C. (2016). The first step for neuroimaging data analysis: DICOM to NIfTI conversion. *Journal of Neuroscience Methods*, *264*, 47–56.

Marshall, A. C., Cooper, N. R., & Geeraert, N. (2016). The impact of experienced stress on aged spatial discrimination: Cortical overreliance as a result of hippocampal impairment. *Hippocampus*, *26*(3), 329–340.

Menon, V., & Uddin, L. Q. (2010). Saliency, switching, attention and control: A network model of insula function. *Brain Structure and Function*, *214*(5–6), 655–667.

Moritz, S., Gläscher, J., Sommer, T., Büchel, C., & Braus, D. F. (2006). Neural correlates of memory confidence. *Neuroimage*, *33*(4), 1188–1193.

Mueller, S. G., Chao, L. L., Berman, B., & Weiner, M. W. (2011). Evidence for functional specialization of hippocampal subfields detected by MR subfield volumetry on high resolution images at 4T. *NeuroImage*, *56*(3), 851–857.

Okado, Y., & Stark, C. E. (2005). Neural activity during encoding predicts false memories created by misinformation. *Learning & Memory*, *12*(1), 3–11.

Olsen, R. K., Nichols, E. A., Chen, J., Hunt, J. F., Glover, G. H., Gabrieli, J. D. E., & Wagner, A.
D. (2009). Performance-related sustained and anticipatory activity in human medial
temporal lobe during delayed match-to-sample. *Journal of Neuroscience*, *29*(38), 11880–
11890. https://doi.org/10.1523/JNEUROSCI.2245-09.2009

Philippi, C. L., Tranel, D., Duff, M., & Rudrauf, D. (2015). Damage to the default mode network
disrupts autobiographical memory retrieval. *Social Cognitive and Affective Neuroscience*,
10(3), 318–326. https://doi.org/10.1093/scan/nsu070

Phillips, J. S., Velanova, K., Wolk, D. A., & Wheeler, M. E. (2009). Left posterior parietal
cortex participates in both task preparation and episodic retrieval. *NeuroImage*, *46*(4),
1209–1221. https://doi.org/10.1016/j.neuroimage.2009.02.044

Pidgeon, L. M., & Morcom, A. M. (2016). Cortical pattern separation and item-specific memory
encoding. *Neuropsychologia*, *85*, 256–271.

Reagh, Z. M., Ho, H. D., Leal, S. L., Noche, J. A., Chun, A., Murray, E. A., & Yassa, M. A.
(2016). Greater loss of object than spatial mnemonic discrimination in aged adults.
Hippocampus, *26*(4), 417–422.

Reagh, Z. M., Roberts, J. M., Ly, M., DiProspero, N., Murray, E., & Yassa, M. A. (2014).
Spatial discrimination deficits as a function of mnemonic interference in aged adults with
and without memory impairment. *Hippocampus*, *24*(3), 303–314.

Reagh, Z. M., & Yassa, M. A. (2014). Object and spatial mnemonic interference differentially
engage lateral and medial entorhinal cortex in humans. *Proceedings of the National
Academy of Sciences*, *111*(40), E4264–E4273.

Roberts, J. M., Ly, M., Murray, E., & Yassa, M. A. (2014). Temporal discrimination deficits as a
function of lag interference in older adults. *Hippocampus*, *24*(10), 1189–1196.

Rolls, E. T., & Kesner, R. P. (2016). *Pattern separation and pattern completion in the hippocampal system. Introduction to the Special Issue*. Elsevier.

Seeley, W. W., Menon, V., Schatzberg, A. F., Keller, J., Glover, G. H., Kenna, H., Reiss, A. L., & Greicius, M. D. (2007). Dissociable intrinsic connectivity networks for salience processing and executive control. *Journal of Neuroscience*, *27*(9), 2349–2356. https://doi.org/10.1523/JNEUROSCI.5587-06.2007

Sestieri, C., Corbetta, M., Romani, G. L., & Shulman, G. L. (2011). Episodic memory retrieval, parietal cortex, and the default mode network: Functional and topographic analyses. *Journal of Neuroscience*, *31*(12), 4407–4420. https://doi.org/10.1523/JNEUROSCI.3335-10.2011

Sheppard, D. P., Graves, L. V., Holden, H. M., Delano-Wood, L., Bondi, M. W., & Gilbert, P. E. (2016). Spatial pattern separation differences in older adult carriers and non-carriers for the apolipoprotein E epsilon 4 allele. *Neurobiology of Learning and Memory*, *129*, 113–119.

Shirer, W. R., Ryali, S., Rykhlevskaia, E., Menon, V., & Greicius, M. D. (2012). Decoding subject-driven cognitive states with whole-brain connectivity patterns. *Cerebral Cortex*, *22*(1), 158–165. https://doi.org/10.1093/cercor/bhr099

Shrager, Y., Kirwan, C. B., & Squire, L. R. (2008). Activity in both hippocampus and perirhinal cortex predicts the memory strength of subsequently remembered information. *Neuron*, *59*(4), 547–553.

Small, S. A., Schobel, S. A., Buxton, R. B., Witter, M. P., & Barnes, C. A. (2011). A pathophysiological framework of hippocampal dysfunction in ageing and disease. *Nature Reviews Neuroscience*, *12*(10), 585–601.

Steemers, B., Vicente-Grabovetsky, A., Barry, C., Smulders, P., Schröder, T. N., Burgess, N., &

    Doeller, C. F. (2016). Hippocampal attractor dynamics predict memory-based decision

    making. *Current Biology*, *26*(13), 1750–1757.

Strange, B. A., Witter, M. P., Lein, E. S., & Moser, E. I. (2014). Functional organization of the

    hippocampal longitudinal axis. *Nature Reviews. Neuroscience*, *15*(10), 655–669.

    https://doi.org/10.1038/nrn3785

Suzuki, C., Tsukiura, T., Mochizuki-Kawai, H., Shigemune, Y., & Iijima, T. (2009). Prefrontal

    and medial temporal contributions to episodic memory-based reasoning. *Neuroscience*

    Research, *63*(3), 177–183. https://doi.org/10.1016/j.neures.2008.11.010

Tustison, N. J., Cook, P. A., Klein, A., Song, G., Das, S. R., Duda, J. T., Kandel, B. M., van

    Strien, N., Stone, J. R., Gee, J. C., & Avants, B. B. (2014). Large-scale evaluation of

    ANTs and FreeSurfer cortical thickness measurements. *NeuroImage*, *99*(0), 166–179.

    http://dx.doi.org/10.1016/j.neuroimage.2014.05.044

Uncapher, M. R., & Wagner, A. D. (2009). Posterior parietal cortex and episodic encoding:

    Insights from fMRI subsequent memory effects and dual-attention theory. *Neurobiology*

    of Learning and Memory, *91*(2), 139–154.

van den Honert, R. N., McCarthy, G., & Johnson, M. K. (2016). Reactivation during encoding

    supports the later discrimination of similar episodic memories. *Hippocampus*, *26*(9),

    1168–1178.

Wais, P. E., Jahanikia, S., Steiner, D., Stark, C. E. L., & Gazzaley, A. (2017). Retrieval of high-

    fidelity memory arises from distributed cortical networks. *NeuroImage*, *149*, 178–189.

    https://doi.org/10.1016/j.neuroimage.2017.01.062

Wang, H., Suh, J. W., Das, S. R., Pluta, J., Craige, C., & Yushkevich, P. A. (2013). Multi-atlas segmentation with joint label fusion. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, *35*(3), 611–623. https://doi.org/10.1109/TPAMI.2012.143

Wickelgren, W. A. (1968). Unidimensional strength theory and component analysis of noise in absolute and comparative judgments. *Journal of Mathematical Psychology*, *5*(1), 102–122.

Wickens, T. D. (2002). *Elementary signal detection theory*. Oxford University Press, USA.

Yushkevich, P. A., Piven, J., Hazlett, H. C., Smith, R. G., Ho, S., Gee, J. C., & Gerig, G. (2006). User-guided 3D active contour segmentation of anatomical structures: Significantly improved efficiency and reliability. *Neuroimage*, *31*(3), 1116–1128.

Yushkevich, P. A., Pluta, J. B., Wang, H., Xie, L., Ding, S.-L., Gertje, E. C., Mancuso, L., Kliot, D., Das, S. R., & Wolk, D. A. (2015). Automated volumetry and regional thickness analysis of hippocampal subfields and medial temporal cortical structures in mild cognitive impairment. *Human Brain Mapping*, *36*(1), 258–287. https://doi.org/10.1002/hbm.22627