Mapping the genetic and environmental aetiology of autistic traits in Sweden and

the UK

Zoe E. Reed[1,2*], Henrik Larsson[4,5], Claire M.A. Haworth[1,2,3,6,7], Dheeraj Rai[2,3,8],

Sebastian Lundström[9], Angelica Ronald[10], Abraham Reichenberg[11], Paul

Lichtenstein[12], Oliver S.P. Davis[1,2,3,7]

* Corresponding author

1) MRC Integrative Epidemiology Unit, University of Bristol, UK

2) Department of Population Health Sciences, Bristol Medical School, University

of Bristol, UK

3) National Institute for Health Research Biomedical Research Centre, University

Hospitals Bristol NHS Foundation Trust and the University of Bristol, Bristol, UK

4) School of Medical Sciences, Örebro University, Sweden

5) Department of Medical Epidemiology and Biostatistics, Karolinska Institutet,

Sweden

6) School of Psychological Science, University of Bristol, UK

7) The Alan Turing Institute, London, UK

8) Avon and Wiltshire Partnership NHS Mental Health Trust

9) Gillberg Neuropsychiatry Centre, Centre for Ethics, Law and Mental Health,

University of Gothenburg, Sweden

10) Department of Psychological Sciences, Birkbeck, University of London, UK

11) Department of Psychiatry and Seaver Autism Center for Research and

Treatment, Ichan School of Medicine at Mount Sinai, New York, USA

12) Department of Medical Epidemiology and Biostatistics, Karolinska Institutet,

Sweden

Abstract

Autistic traits are influenced by both genetic and environmental factors, and are

known to vary geographically in prevalence. But to what extent does their

aetiology also vary from place to place? We applied a novel spatial approach to

data from two large twin studies, the Child and Adolescent Twin Study in

Sweden (CATSS) and the Twins Early Development Study (TEDS) in the UK, to

explore how the influence of nature and nurture on autistic traits varies from

place to place. We present maps of gene- and environment- by geography

interactions that suggest, for example, higher heritability and lower non-shared

environmental influence in more densely populated areas. We hope this

systematic approach to aetiological interactions will inspire research to identify

previously unknown environmental influences on the aetiology of autistic traits.

Background

Autism spectrum disorder (ASD) is a neurodevelopmental condition that manifests in childhood. ASD is generally characterised by persistent difficulties with social communication and repetitive behaviours. Reported prevalence of ASD varies, but in developed countries the prevalence is estimated to be between 1-1.5% [1-4] and this recorded prevalence has increased over the past few decades [1,3,4]. Factors such as diagnostic criteria, age, time of study and location of study may all contribute to this heterogeneity in prevalence estimates. ASD has a significant impact on child development, often including language difficulties and other co-occurring conditions which may persist into adulthood [5].

The aetiology of ASD reflects both genetic and environmental influences. Twin studies suggest that genetic differences between people explain around 80% of the population variance for ASD [6]. Most studies suggest that the remaining variance is explained by variation in the non-shared environment. That is, environmental influences that do not contribute to similarity within families. Similarly, a recent study of over 3.5 million twin and sibling pairs in Sweden found that 83% of the variance is explained by genetic differences and 17% by non-shared environmental influences [7]. Another study across 5 different countries (Sweden, Finland, Denmark, Western Australia and Israel) estimated heritability for ASD to be around 80%, using data from whole populations, although there was variation between countries [8].

ASD is known to vary in prevalence across geographical regions. For example, spatial analyses of ASD have revealed areas of increased prevalence in Salt Lake

County in Utah [9], in northern Taiwan [10] and in areas of California [11,12]. ASD also

appears more common in those born in New England compared to those born in

the south east of the United States (US) [13]. Similarly, a study of Greater Glasgow

in Scotland identified variation in prevalence across the city [14]. Several studies

have also suggested differences in prevalence between urban and rural areas,

where living in or growing up in an urban environment is associated with

greater risk of ASD compared to rural environments [10,15-18]. Possible reasons for

geographical variation in prevalence of ASD across these areas include regional

diagnostic bias, differences in the access to health services or diagnostic

resources, different levels of parental awareness, air pollution exposure during

pregnancy, green space in an area and local trends in socioeconomic status

(SES).

If prevalence of autistic traits varies from place to place, is the same true of the

aetiology? For example, does variation in the environment explain variation in

autistic traits in some areas more than others? Similarly, does the environment

in some areas draw out genetic differences between children in their propensity

for developing autistic traits? We previously developed a spatial approach to

twin model-fitting called spACE that showed genetic and environmental

influences vary spatially within a country in response to geographically-

distributed environments [19]. This approach has the potential to highlight gene-

environment (G×E) and environment-environment (E×E) interactions for

outcomes such as ASD traits. G×E and E×E represent variation in the aetiological

genetic influences on a trait depending on environmental exposure. For example,

genetic risk of a mental health disorder may be drawn out by a stressful

environment, genetic risk of asthma may be apparent only in polluted

environments, or genetic risk of hay fever may only reveal itself in pollen-rich

areas. The spACE approach allows us to investigate this, mapping geographical

patterns of nature and nurture without requiring the measurement of specific

genetic variants or specific environmental characteristics. This systematic

geographical approach may facilitate the discovery of novel specific genetic and

environmental influences.

Here we apply the spACE approach to data on autistic traits in Sweden and the

UK. Autistic traits and diagnostic categories of ASD show substantial aetiological

overlap [20,21], with genetic correlations from bivariate twin models of 0.52-0.89

and SNP based genetic correlations of 0.27-0.30. The heritability of ASD traits

does not change as a function of severity [22–24], and genetic links have been

identified between extreme and sub-threshold variation in ASD [22,24], so to

maximise power we have focussed on trait measures rather than diagnoses.

It seems likely that environments previously identified as important for the

development of autistic traits will also influence aetiology. For example, given

previous research on the social stress of urban compared to rural upbringing [25],

we hypothesise that urban-rural differences will be apparent in the aetiology of

autistic traits. However, more importantly, we hope that by systematically

mapping geographical differences in aetiology we will facilitate identification of

new environments and shed light on the mechanisms by which they act.

Methods

The Swedish Twin Registry and CATSS

The Swedish twin registry, established in the 1950s, currently includes over 194,000 twins [26]. Phenotypic information on the twins comes from a variety of sources such as medical registers and questionnaires and is regularly updated. Several sub-studies of the registry have been established, including the longitudinal Child and Adolescent Twin Study in Sweden (CATSS) [27]. CATSS was launched in 2004 to investigate childhood-onset neurodevelopmental problems such as ADHD and ASD in childhood and adolescence, for all twins turning 9 or 12 years since 2004. Parents of all Swedish twins aged 9 and 12 years old were asked to participate in a telephone interview to collect information on various health-related issues. By the time data on autistic traits were obtained in 2013, 8,610 parents had responded to this request, accounting for 17,220 twins. The CATSS-9/12 study obtained ethical approval from the Karolinska Institute Ethical Review Board: Dnr 03-672 and 2010/507-31/1, CATSS-9 – clinical 2010/1099-31/3 CATSS-15 Dnr: 2009/1599-32/5, CATSS-15/DOGSS Dnr: 03-672 and 2010/1356/31/1, and CATSS-18 Dnr: 2010/1410/31/1.

For ASD traits, 16,677 participants had data available (including 8,307 complete pairs and 63 incomplete pairs of twins). Interviews were carried out when the twins were around the age of 9 or 12 years and 51% were male.

CATSS measures of autistic traits

The Autism-Tics, ADHD and other Comorbidities (A-TAC) inventory, based on the Diagnostic and Statistical Manual of Mental Disorders (DSM)-IV criteria, was used in the

telephone interview with parents to collect information on a range of

neurodevelopmental problems. This inventory has previously been validated in both

clinically diagnosed children and the general population [28–32]. The inventory includes 17

items that assess ASD symptoms, where respondents can answer 'yes/1', 'yes, to some

extent/0.5', and 'no/0'. Following the standard approach, we created a symptom score

for each individual by summing these item scores. Further details can be found in

previous publications [32].

In our sample the median score was 0.00 (interquartile range [IQR]=1.00) for autistic

symptoms, where, in previous validation studies a low and high cut-off of 4.5 and 8.5 for

ASD have been established for broad screening and for use as a clinical proxy,

respectively. This indicates that most people, in this general population sample, score

well below these cut-offs. As expected, the distribution of this symptom score was zero-

skewed, as shown in the histogram in **supplementary figure 1**.

CATSS location data

To conduct the spACE analysis, we assigned a geographical location to each family. In

CATSS we matched each twin pair to a Small Areas for Market Statistics (SAMS)

location, for the most recent location data we had available up to 2009, using data from

Statistics Sweden (http://www.scb.se/en/) and assigned coordinates based on the

centroid of the SAMS location. There are approximately 9,200 SAMS in Sweden,

subdivisions of 290 municipalities. The average population within each SAMS is 1,000

people and therefore the area covered by each SAMS varies by population density.

To provide context for the results for Sweden, it is useful to understand a little about its

geography. **Figure 1** shows a map of Sweden and some general indicators of the

country's geography; the **supplementary materials** contain a detailed description. In

summary, Sweden is split into a more rural north and central area, known as the

lowlands and the more populated areas in a belt from Gothenburg in the west to

Stockholm in the east and the very south near Malmö. Much of Sweden is covered by

forest and lakes. The capital, Stockholm, is in the east with a mix of tourist-centred and

residential areas and a number of islands. Gothenburg, the second largest city with a

port, also has varied areas like Stockholm and an archipelago. Further inland is rural

Värmland. South-west Sweden is a coastal and lowland area and is the third most

populated area in Sweden. The main city in this area is industrialised and multicultural

Malmö. South-east Sweden is heavily forested with some large lakes and a number of

large towns. Sweden's two largest islands are also found here, Öland and Gotland, which

are popular summer destinations, due to their warmer climate. They both have fairly

rural landscapes with small towns and villages. The Bothnian coast is the most

populated area in the north, with some large towns along the coastline. Central Sweden

is a sparsely populated, rural, lowland area covered in forests, with numerous lakes and

mountains along the Norwegian border. Further north is Swedish Lapland, a very

remote area with a mountainous, rural landscape.

Figure 1. Map of Sweden with top 10 cities by population

Sweden is shown in dark grey with the surrounding countries in a lighter shade of grey. The top 10 most

populated cities are shown on the map with orange circles, the area of which reflects the population of the

city.

 The Twins Early Development Study

The Twins Early Development Study (TEDS) contacted parents of twins born in England

and Wales between January 1994 and December 1996 [33]. 16,810 pairs of twins were

initially recruited, and currently there are over 10,000 twin pairs still enrolled in TEDS.

The participants are demographically representative of the UK population of a similar

age, with the majority identifying themselves as white British and with English as their

first language. TEDS has collected wide-ranging data on cognitive and behavioural

development, using approaches that include questionnaire booklets, telephone testing

and web-based tests. The twins, their parents and teachers have all participated in data

collection. Ethical approval for TEDS research is provided by the Institute of Psychiatry,

Psychology and Neuroscience Ethics Committee, King's College London.

Full phenotypic data for autistic traits were available for 11,594 TEDS participants

(including 5,796 complete pairs and 62 incomplete pairs of twins). For these twins the

mean age was 11.30 (SD=0.72) and 48% were male.

TEDS measures of autistic traits

Parents in TEDS completed the Childhood Autism Spectrum Test (CAST) when the twins

were age 12 years. The CAST consists of 30 items, scored 1 for yes or 0 for no [34]. For

participants included in our analyses, the median score for ASD symptoms was 4.0

(IQR=4.84). The CAST score considered indicative of ASD is 15.

    TEDS location data

    We assigned each twin pair geographical coordinates based on the centroids of their

    postcodes at age 12. There are over 1.5 million postcode units in the UK, covering, on

    average, 15 properties. As with SAMS, the area covered by each postcode varies

    depending on population density.

    To provide context for the results for the UK, **Figure 2** displays a map and some general

    indicators of the country's geography; the **supplementary materials** include a detailed

    description. The UK is split into England, Wales, Scotland and Northern Ireland

    (although we do not describe Northern Ireland here because it was not included in the

    TEDS recruitment area and few participants have moved there since recruitment).

    Generally, the south of the UK has a milder climate compared to the north and has more

    low-lying land. The UK is a mix of some very urban, previously (or still) industrial areas

    and more rural traditional countryside areas. London, a diverse, multicultural city in the

    south-east, is the capital, with its own distinct boroughs. The south-east of England has

    many commuter areas and is surrounded by coastline. This area is historically rich and

    has a mix of industrial and countryside areas and a number of seaside towns. The south

    of the UK is fairly rural, has many historical sites and also many coastal towns as well as

    the New Forest. More inland are the areas of Oxfordshire, with the city of Oxford and

    the picturesque, rural Cotswolds. In the west are the areas of cosmopolitan Bristol, spa-

    city Bath and rural Somerset, with the wooded Mendips, the Quantock and Exmoor

    National Park, on the Bristol channel. South-west England consists of pre-industrial

    Devon and Cornwall, popular Summer destinations with many seaside and fishing

    towns and plenty of farmland, and Dartmoor National Park. East Anglia is an area of

    flatland, wetlands and coastal areas.

The west Midlands are a mix of lowlands and hilly areas with the industrial city of

Birmingham (England's second largest city) and the Peak district. The east midlands has

a number of large urbanised cities and is an old coal mining area, but rural areas can

still be found. North-west England has the large cities of Manchester and Liverpool and

the seaside resort of Blackpool, but also the unspoilt, mountainous Isle of Man. The

scenic Lake district is also found in the north as well as the varied area of Yorkshire,

with urbanised, coastal and rural areas. Wales is split into the more populated and

coastal south, hilly, rural Mid-Wales and mountainous north Wales. Scotland is split into

the Highlands in the north and the Lowlands in the south. Southern Scotland is home to

the main cities of cosmopolitan and medieval Edinburgh and urban Glasgow and this

area has coastal towns, forests and agricultural land. Central Scotland is more varied

with large lochs and forests in the west, rural and industrialised areas and fishing

villages in the east and peaks in the north. Argyll in the west is a remote area,

transitioning between lowland and highland and with numerous islands. North east

Scotland has a number of industrial cities and port towns, although further north

becomes mountainous. The Highlands are a very remote but unspoilt area, with forests,

lochs, mountains and rugged coastline. Scotland has a number of island clusters, the

Inner and Outer Hebrides in the west and the Orkney and Shetland islands in the north.

Figure 2. Map of the UK with top 10 cities by population

The recruitment area (England and Wales) for the Twins Early Development Study (TEDS) is shown in dark

grey, with the rest of the UK (Scotland and Northern Ireland [N.I]) in a lighter shade of grey. Other countries

are shown in the lightest shade of grey. The top 10 most populated cities are shown on the map with orange

circles, the area of which reflects the population of the city.

Statistical analyses

ACE models and maps in CATSS and TEDS

In twin analysis, within-pair similarity of monozygotic (MZ) and dizygotic (DZ) twins is compared to estimate parameters for additive genetic (A), shared environmental (C) and non-shared environmental (E) influences on a trait. In this context, the shared environment refers to influences other than DNA similarity that make children growing up in the same family more similar to each other, whilst the non-shared environment refers to influences that do not contribute to similarity within families. Although tempting, it is not possible to assign specific environments to one or the other environmental component, because most environments themselves show both shared and non-shared (and often genetic) influences. We can estimate the contribution of genetic and environmental influences because of the different ways these influences are shared in MZ and DZ twin pairs. For MZ twins, who share 100% of their segregating alleles, A influences correlate 1, whereas for DZ twins they correlate 0.5 because DZ twins share, on average, 50% of their segregating alleles. For both MZ and DZ twins growing up in the same family the shared environmental correlation is 1. In contrast, the non-shared environment is uncorrelated and contributes to differences between twins [35]. In this study, we applied a version of the spACE analysis method [19] to explore how A, C and E for ASD traits vary geographically. To do this, we fit full information maximum likelihood structural equation models to twin data in R (version 3.3.1) using the OpenMx package (version 2.9.4), calculating A, C and E at many different target locations across an area. The contribution of each twin pair to a model is weighted by a function of the inverse Euclidean distance of the twin pair from the target location. In this study we built on our previous work by applying the weights within the structural

equation modelling framework, rather than by calculating weighted correlation

matrices and using those as input (although for normally distributed measures the

results are the same with either approach). In twin analysis it is possible to model non-

additive genetic effects (D) instead of shared environmental effects (C), and D influences

are sometimes found with ASD. However, the D component is highly correlated with the

A component, which means confidence intervals are wide and the tendency of variance

to swap between these two components makes it difficult to compare results across

locations. Because of this, we have fitted ACE models, although in this case, A should be

considered broad-sense heritability, including both additive and non-additive genetic

influences.

For target locations in Sweden we used the centroid of each unique SAMS that included

at least one twin pair. Because UK postcodes give more precise locations than Swedish

SAMS, we instead selected UK target locations representative of local population density

to preserve participant anonymity. All twin pairs contributed to the results at each

location, but contributions were weighted according to the distance of each twin pair

from the target location:

$$w_i(x) = \frac{1}{d(x, x_i)^p}$$

where x represents the target location, x_i represents the location of a twin pair, d is the

Euclidean distance between x and x_i, and p is the power parameter that controls the rate

of drop-off of a twin pair's influence over distance (0.5 for these analyses). We included

sex as a covariate in all the models (accounting for on average 2.59% of the variance),

and age in the TEDS data (where it accounted for on average 0.34% of the variance).

Further detail on the spACE approach can be found in the original article [19]. We plotted

maps to visualise the results (**figures 3 and 4**). In the maps each target location is

coloured according to the value of the estimate at that location compared to the full

range of values across the map. Low values appear blue and high values appear red,

with increasing brightness of the colour representing increasing distance from the

mean. To avoid outliers having a large effect on the distribution of colours in the maps,

we assigned the highest 4% of values to the brightest red and the lowest 4% of values to

the brightest blue before assigning colour values to equal ranges between the two. The

histograms show the distribution of results and the corresponding colours.

We estimated 95% confidence intervals for A, C and E at each target location and using

the CATSS data we performed sensitivity analyses for how A, C and E estimates vary

based on the historical residential location used for the twin pairs. To do this we

repeated analyses based on participants' locations at different ages and we combined

the resulting maps into a video (**supplementary video 1**). Changes across time may

allow identification of critical developmental periods when the geographical

environment is particularly influential; for example, if clear patterns are seen when

participant locations for the analysis are based on their location at a specific age.

Sex limitation models

While some previous studies have identified no aetiological sex differences for ASD,

others have. For example, one study using the Missouri twin study and a continuous

measure of autistic traits, found no sex differences [36], and neither did previous work in

TEDS[24], but modest sex differences were found in previous work with the Swedish Twin

Study[37]. To maximise power, in the main text we report results that equate the

aetiological influences for males and females. But for the Swedish data, where we

replicated quantitative sex differences in aetiology, we conducted further separate

analyses for males and females, and we include these in supplementary materials

(**supplementary figures 4 and 5**).

Data availability

The data used in this study are available to researchers directly from CATSS and TEDS.

Procedures for accessing the data are described at https://ki.se/en/meb/the-child-and-

adolescent-twin-study-in-sweden-catss (CATSS) and

https://www.teds.ac.uk/researchers/teds-data-access-policy (TEDS).

Code availability

Code that implements the spACE model described here is available in the scripts

directory at https://github.com/DynamicGenetics/spACEjs/.

Results

Mapping the aetiology of autistic traits in Sweden

We plotted the results from each of the 4,199 locations on a map (**Figure 3**, an

interactive version is available at https://dynamicgenetics.github.io/spACEjs/), where

red points represent locations where results fall above the population mean, and blue

points represent locations where they fall below. The brighter the points, the further

they are from the population mean. This is shown in the histograms below the maps,

where the colours of the bars match the points in the map above. Because we modelled

raw variance after standardising data to mean 0 and SD 1 at the population level (i.e. we

did not standardize the A, C and E estimates at each location to add up to one) genetic

and environmental influences are not reciprocal at each location, so it is possible for a

location to show both strong genetic and environmental influences. The consequence of

this is that each map stands alone: differences in genetic influence really do imply

differences in the genetic component, and not just a reflection of differences in a

reciprocal environmental component. Maps with A and E constrained to add up to one

in each location (i.e. proportional) are shown in **supplementary figure 2** and show

similar results to those for the raw variance. For comparison, we have also plotted

results of the weighted means of scaled autistic trait scores at the same locations in

supplementary figure 3 and we observe geographical variation for mean autistic trait

scores, reflecting the expected variation in the prevalence of ASD.

Figure 3. Mapping genetic (A) and non-shared environmental (E) influences on autistic

traits in Sweden suggests that genetic variation is more influential in more densely-

populated areas

Geographical variation in genetic (A) and non-shared (E) influences on childhood ASD traits in Sweden

(results are overlaid on an outline of the SAMS areas). The contributions of A and E range from low (blue) to

high (red). The histograms below show the distribution of the estimates, coloured in the same way as the

points on the map. The estimates are not standardised and are therefore not constrained to add up to one.

Shared environment (C) estimates were approximately zero across the whole map, so they are not shown

here. An interactive version of this map is available at https://dynamicgenetics.github.io/spACEjs/.

The results suggest that the amount of variation in autistic traits explained by genetic

influences (A) is generally greater in urban areas and lesser in the sparsely populated

north and more rural southern belt. The non-shared environment (E) frequently shows

the opposite pattern, with the variation explained generally less in and around the

capital and more in Southern and Northern rural areas. However, we also observe

greater contribution of E in the areas around the cities of Gothenburg and Malmö.

Variation in A and E can also be seen within local areas, such as around Stockholm,

where there are both low and high values for A, suggesting genetic influences are

moderated by other factors beyond urbanicity. The histograms for the raw variance

indicate that the variance explained by genetic influences ranges from 0.55 to 0.91, with

most values around the mean of 0.65 (SD=0.02). The variance explained by E ranges

from 0.25 to 0.46, again with most values around the mean of 0.34 (SD=0.02). Variation

in autistic traits explained by C was approximately zero over the whole of Sweden.

Confidence intervals for estimates at each location are provided in **supplementary**

table 1. Supplementary video 1 shows that the overall patterns for variation in A and

E remained similar irrespective of the historical location used for each twin pair.

Sex limitation models for ASD traits

Population-level sex limitation model results for autistic traits are shown in

supplementary table 2. We used nested sub-models to test for sex differences in

aetiology. The common effects model is the most parsimonious model that adequately

fits the data, indicating quantitative, but not qualitative, sex differences. However,

because this is a large and well-powered sample, the parameter estimates for males and

females that are "significantly" different at an alpha of 0.05 are actually within 1% of

each other. To maximise power, we have presented the maps for males and females

combined in the main text, but maps for males and females separately are shown in

supplementary figures 4 and 5.

Mapping the aetiology of autistic traits in the UK

Figure 4 maps genetic and environmental influences on autistic traits in the UK at 6,758

locations chosen to represent sample density across the UK (an interactive version of

this map is available at https://dynamicgenetics.github.io/spACEjs/). Again, this is a

map of the raw variance, so A, C and E are not constrained to add up to one. However,

maps with A, C and E constrained to add to one at each location are shown in

supplementary figure 6, with very similar results.

Figure 4. Similar to findings in Sweden, genetic influences for autistic traits in the UK

appear more influential in more densely populated areas, although patterns of non-

shared environmental influences follow a north-south divide

Geographical variation in genetic (A) and non-shared (E) influences on childhood autistic traits in the UK.

The contributions of A and E range from low (blue) to high (red). The histograms below show the

distribution of the estimates, coloured in the same way as the points on the map. The A and E estimates are

not standardised and are therefore not constrained to add up to one at each location. An interactive version

of this map is available at https://dynamicgenetics.github.io/spACEjs/.

The raw results for A are consistent with those from the CATSS sample in Sweden

where we observed higher heritability in more densely populated areas. The mean of A

is slightly higher in the UK than in Sweden: 0.76 (SD=0.01) compared to 0.65 (SD=0.02).

For non-shared environment (E) the patterns are less similar across countries, as are

the mean values 0.23 (SD=0.01) in the UK, compared to 0.34 (SD=0.02) in Sweden.

London, the capital city, and the surrounding south-east of the UK show greater

influence of E compared to the north and some regions in the mid-west of England and

Wales. In contrast, Sweden's capital, Stockholm, and the surrounding areas show lower

estimates of E. Again, C is approximately zero for autistic traits across all regions. As

before, local variation in A and E is apparent within large cities such as London. As the

histograms show, A is fairly normally distributed between 0.69 and 0.84 across regions.

E ranges more narrowly from 0.21 to 0.29 in a bimodal distribution with a positive

skew. Confidence intervals for estimates at each location are provided in

supplementary table 3.

Discussion

In this study we looked at how genetic and environmental influences on symptoms of

ASD vary geographically in Sweden and the UK. Our results are consistent with previous

population-level estimates of genetic and environmental influences, and demonstrate

geographical variation in genetic and non-shared environmental influences on autistic

traits in Sweden and the UK.

These geographical differences in genetic and environmental influences on autistic

traits are indicative of gene-environment and environment-environment interactions

where the interacting environmental variable varies by location. Where we find areas of

increased genetic or environmental influences for autistic traits this means that the

environment in these areas draws out genetic or environmental influence, in the same

way that the presence of airborne pollen would reveal individual differences in genetic

risk for hay fever. By studying this in a systematic way, rather than relying on a specific

measured environment, we can use our results to develop novel hypotheses about

currently unknown environmental influences.

Our findings complement previous research that has focused on geographical

prevalence differences in ASD [9-14,38]. Similarly, alongside aetiological differences, we

observe geographical variation in mean autistic trait scores. These mean differences

may be linked to aetiological differences. For example, areas of greater prevalence could

represent regions where the environment triggers genetic predisposition to ASD traits.

This provides a basis for future research into specific geographically distributed

environments that draw out or mitigate genetic or environmental risk, which could in

turn be useful for population health measures seeking to reduce the impact of ASD.

From our results we can hypothesise about what these factors could be. For example,

we find that there is generally higher heritability in more densely populated areas of

Sweden, such as Gothenburg, Malmö, and Stockholm, and in a band running from

Gothenburg to Stockholm where the majority of the population live. We find lower

heritability in the southern highlands and northern regions, which are less populated.

This may suggest that urban environments draw out genetic differences in

predisposition to autistic traits between people of the same ancestral background.

These geographically distributed environments might include psychosocial factors such

as the stress of urban living or income inequality, or aspects of the physical

environment such as air pollution. This explanation fits with neuroscience literature

that suggests that living in an urban environment is associated with specific neural

correlates in response to stress, which may influence the onset of related mental health

disorders [25]. The literature on prevalence suggests that other potentially important

factors may include geographical differences in access to healthcare, diagnostic bias and

parental awareness, socio-economic status, neighbourhood deprivation, infrastructure

of the area, or access to green space. However, factors such as rater effects or access to

healthcare are less likely to play a role in this aetiological variation as we have used

data from structured interviews in population representative samples, and

environmental influences on prevalence are not necessarily the same as environmental

influences on aetiology. For non-shared environmental influences urban-rural

differences are confined to areas in and around Stockholm, the Swedish capital.

Therefore, it may be that there are environments related specifically to living in or

around the capital that result in decreased non-shared environmental influences

compared to other areas in Sweden.

We see similar patterns for genetic influences on in the UK, with higher heritability

estimates in city areas such as central and south London, Birmingham, Bristol,

Manchester, Newcastle. Estimates are generally lower in East Anglia, the south west,

Wales and other less densely populated areas. Again, as in Sweden, non-shared

environmental influence shows a more complex pattern in the UK.

Whilst we see similarities in patterns of aetiology between Sweden and the UK for

autistic traits, there are also substantial differences. There are several possible reasons

for this. For example, it could be due to differences in the measurement of ASD

symptoms in the cohorts, or it could be due to environmental differences between the

two countries, for example differences in the level of awareness of ASD and therefore

possible differential reporting in ASD symptoms, or differences in the physical or social

environments, which may vary between countries in the same way as they do within

each country. It will be important to investigate this in other countries to explore these

international similarities and differences further.

When interpreting these results there are a few important points to consider. First, in

some areas the effective sample size is lower than others, for example in densely

populated areas the proximity of some twin pairs relative to others can weight their

influence relatively highly. However, across all areas we have taken care to maintain

effective sample sizes in the thousands for both identical and fraternal twin pairs, so

estimates remain reasonably precise. Second, due to how the weighting of participants'

contributions to the analyses works, i.e. participants contribute more to analysis the

closer they are to the target location, this results in smoothing over the estimates for A,

C and E. The amount by which results over the area are smoothed depends on the

tuning parameter used in the weighting. There is a trade-off when selecting the tuning

parameter between smoothing over noise and detecting real variation, or between

accurately estimating variance components and accurately localising them. Here, we

have chosen the tuning parameter to result in some smoothing towards the population

mean, but this may mean that some larger localised variation remains undetected. In

interpreting the maps, it is important to take into account both the pattern of results

shown on the map, and the range of estimates shown by the histogram, while bearing in

mind that the effect sizes are smoothed towards the population mean. Third, in common

with the previous literature, we find that an ADE model is often a slightly better fit to

the data, but here we have fitted ACE models and generally presented results for A and

E alone because the high correlation between A and D brings noise to spatial analysis

due to switching between the two across locations [39]. Instead, we interpret A here as a

broad genetic component, without the usual connotation of additivity. Fourth, as with

any statistical analysis, it is important to consider the assumptions of the model. For

twin modelling, these include random mating within the population, that MZ and DZ

twins share their environments to the same extent (at least where those environments

are not genetically influenced), and that twins are representative of the general

population for the traits studied [35]. These assumptions have generally been found to be

reasonable [40], although there is some evidence to suggest that there is assortative

mating for ASD, for example a study in Sweden that found phenotypic correlations of

0.48 for ASD [41]. This would have the effect of inflating the shared environmental

influences, which we find to be approximately zero across locations. For our

geographical analyses we do not assume that there is no gene-environment interaction

or correlation, because we are explicitly modelling them as our main point of interest.

Our systematic analysis shows geographical variation in genetic and non-shared

environmental influences for symptoms of ASD in both Sweden and the UK. These

results will inform further studies of measured geographically distributed

environments, beyond those already identified as influencing prevalence in the

literature. For example, by correlating the spatial distribution of these environments

with the spatial distribution of the aetiological estimates or by using formal continuous

moderator models. Identifying these environments and understanding how they draw

out or mask genetic predisposition may lead to population health and social policy

innovation to support people with ASD.

References

1.   Lundström, S., Reichenberg, A., Anckarsäter, H., Lichtenstein, P. & Gillberg, C.

Autism phenotype versus registered diagnosis in Swedish children: Prevalence

trends over 10 years in general population samples. *BMJ* **350**, h1961 (2015).

2.   Brugha, T. S. *et al.* Epidemiology of autism spectrum disorders in adults in the

community in England. *Arch. Gen. Psychiatry* **68**, 459–466 (2011).

3.   Idring, S. *et al.* Changes in Prevalence of Autism Spectrum Disorders in 2001–

2011: Findings from the Stockholm Youth Cohort. *J. Autism Dev. Disord.* **45**, 1766–

1773 (2015).

4.   Lyall, K. *et al.* The Changing Epidemiology of Autism Spectrum Disorders. *Annu.*

Rev. Public Health **38**, 81–102 (2017).

5.   Seltzer, M. M., Shattuck, P., Abbeduto, L. & Greenberg, J. S. Trajectory of

development in adolescents and adults with autism. *Ment. Retard. Dev. Disabil.*

Res. Rev. **10**, 234–247 (2004).

6.   Tick, B., Bolton, P., Happé, F., Rutter, M. & Rijsdijk, F. Heritability of autism

spectrum disorders: A meta-analysis of twin studies. *J. Child Psychol. Psychiatry*

Allied Discip. **57**, 585–595 (2016).

7.   Sandin, S. *et al.* The Heritability of Autism Spectrum Disorder. *JAMA* **318**, 1182–

1184 (2017).

8.   Bai, D. *et al.* Association of Genetic and Environmental Factors With Autism in a 5-

Country Cohort. *JAMA Psychiatry* 1–9 (2019).

doi:10.1001/jamapsychiatry.2019.1411

9.   Bakian, A. V., Bilder, D. A., Coon, H. & McMahon, W. M. Spatial Relative Risk

Patterns of Autism Spectrum Disorders in Utah. *J. Autism Dev. Disord.* **45**, 988–

1000 (2015).

10.   Chen, C. Y., Liu, C. Y., Su, W. C., Huang, S. L. & Lin, K. M. Urbanicity-related variation

in help-seeking and services utilization among preschool-age children with

autism in Taiwan. *J. Autism Dev. Disord.* **38**, 489–497 (2008).

11.   Mazumdar, S., King, M., Liu, K. Y., Zerubavel, N. & Bearman, P. The spatial

structure of autism in California, 1993-2001. *Heal. Place* **16**, 539–546 (2010).

12.   Mazumdar, S., Winter, A., Liu, K. Y. & Bearman, P. Spatial clusters of autism births

and diagnoses point to contextual drivers ofincreased prevalence. *Soc. Sci. Med.*

95, 87–96 (2013).

13.   Hoffman, K. *et al.* Geographic Patterns of Autism Spectrum Disorder Among

Children of Participants in Nurses ' Health Study II. *Am. J. Epidemiol.* **186**, 834–

842 (2017).

14.   Campbell, M., Reynolds, L., Cunningham, J., Minnis, H. & Gillberg, C. Autism in

Glasgow: Cumulative incidence and the effects of referral age, deprivation and

geographical location. *Child. Care. Health Dev.* **39**, 688–694 (2011).

15.   Lauritsen, M. B., Pedersen, C. B. & Mortensen, P. B. Effects of familial risk factors

and place of birth on the risk of autism: A nationwide register-based study. *J. Child*

Psychol. Psychiatry **46**, 963–971 (2005).

16. Lauritsen, M. B. *et al.* Urbanicity and autism spectrum disorders. *J. Autism Dev.*

Disord. **44**, 394–404 (2014).

17. Wu, J. & Jackson, L. Inverse relationship between urban green space and

childhood autism in California elementary school districts. *Environ. Int.* **107**, 140–

146 (2017).

18. Vassos, E., Agerbo, E., Mors, O. & Bøcker Pedersen, C. Urban-rural differences in

incidence rates of psychiatric disorders in Denmark. *Br. J. Psychiatry* **208**, 435–

440 (2016).

19. Davis, O. S. P., Haworth, C. M. A., Lewis, C. M. & Plomin, R. Visual analysis of

geocoded twin data puts nature and nurture on the map. *Mol. Psychiatry* **17**, 867–

874 (2012).

20. Colvert, E. *et al.* Heritability of autism spectrum disorder in a UK population-

based twin sample. *JAMA Psychiatry* **72**, 415–423 (2015).

21. Robinson, E. B. *et al.* Genetic risk for autism spectrum disorders and

neuropsychiatric variation in the general population. *Nat. Genet.* **48**, 552–555

(2016).

22. Robinson, E. B. *et al.* Evidence that autistic traits show the same etiology in the

general population and at the quantitative extremes (5%, 2.5%, and 1%). *Arch.*

Gen. Psychiatry **68**, 1113–1121 (2011).

23.   Lundström, S. *et al.* Autism spectrum disorders and autisticlike traits: Similar

etiology in the extreme end and the normal variation. *Arch. Gen. Psychiatry* **69**,

46–52 (2012).

24.   Ronald, A. *et al.* Genetic heterogeneity between the three components of the

autism spectrum: A twin study. *J. Am. Acad. Child Adolesc. Psychiatry* **45**, 691–699

(2006).

25.   Lederbogen, F. *et al.* City living and urban upbringing affect neural social stress

processing in humans. *Nature* **474**, 498–501 (2011).

26.   Magnusson, P. K. E. *et al.* The Swedish Twin Registry: Establishment of a Biobank

and Other Recent Developments. *Twin Res. Hum. Genet.* **16**, 317–329 (2012).

27.   Anckarsäter, H. *et al.* The Child and Adolescent Twin Study in Sweden (CATSS).

Twin Res. Hum. Genet. **14**, 495–508 (2011).

28.   Hansson, S. L. *et al.* Psychiatric telephone interview with parents for screening of

childhood autism - Tics, attention-deficit hyperactivity disorder and other

comorbidities (A-TAC): Preliminary reliability and validity. *Br. J. Psychiatry* **187**,

262–267 (2005).

29.   Anckarsäter, H. *et al.* Child neurodevelopmental and behavioural problems are

intercorrelated and dimensionally distributed in the general population. *The Open*

Psychiatry Journal **2**, 5–11 (2008).

30.   Mårland, C. *et al.* The Autism-Tics, ADHD and other Comorbidities inventory (A-
TAC): Previous and predictive validity. *BMC Psychiatry* **17**, 403 (2017).

31.   Larson, T. *et al.* Predictive properties of the A-TAC inventory when screening for
childhood-onset neurodevelopmental problems in a population-based sample.
BMC Psychiatry **13**, 233 (2013).

32.   Larson, T. *et al.* The Autism-Tics, AD/HD and other Comorbidities inventory (A-
TAC): further validation of a telephone interview for epidemiological research.
BMC Psychiatry **10**, 1 (2010).

33.   Haworth, C. M. A., Davis, O. S. P. & Plomin, R. Twins Early Development Study
(TEDS): A Genetically Sensitive Investigation of Cognitive and Behavioural
Development From Childhood to Young Adulthood. *Twin Res. Hum. Genet.* **16**,
117–125 (2013).

34.   Scott, F. J., Baron-Cohen, S., Bolton, P. & Brayne, C. The CAST (Childhood Asperger
Syndrome Test): preliminary development of a UK screen for mainstream
primary-school-age children. *Autism* **6**, 9–31 (2002).

35.   Rijsdijk, F. V & Sham, P. C. Analytic approaches to twin data using structural
equation models. *Brief. Bioinform.* **3**, 119–133 (2002).

36.   Constantino, J. N. & Todd, R. D. Autistic Traits in the General Population. *Arch. Gen.*
Psychiatry **60**, 524 (2003).

37.   Ronald, A., Larsson, H., Anckarsäter, H. & Lichtenstein, P. A twin study of autism

symptoms in Sweden. *Mol. Psychiatry* **16**, 1039–1047 (2011).

38.   Hoffman, K., Kalkbrenner, A. E., Vieira, V. M. & Daniels, J. L. The spatial distribution

of known predictors of autism spectrum disorders impacts geographic variability

in prevalence in central North Carolina. *Environ. Heal.* **11**, 80 (2012).

39.   Rietveld, M. J. H., Posthuma, D., Dolan, C. V. & Boomsma, D. I. ADHD: Sibling

interaction or dominance: An evaluation of statistical power. *Behav. Genet.* **33**,

247–255 (2003).

40.   Evans, D. M. & Martin, N. G. The validity of twin studies. *Genescreen* **1**, 77 (2000).

41.   Nordsletten, A. E. *et al.* Patterns of nonrandom mating within and across 11 major

psychiatric disorders. *JAMA Psychiatry* **73**, 354–361 (2016).

Acknowledgements

We gratefully acknowledge the ongoing contribution of the participants in the TEDS and

CATSS and their research teams. This work was supported in part by the UK Medical

Research Council Integrative Epidemiology Unit at the University of Bristol (Grant ref:

MC_UU_12013/1). ZR is supported by a Wellcome Trust PhD studentship (Grant ref:

109104/Z/15/Z). OSPD and CMAH are funded by the Alan Turing Institute under the

EPSRC grant EP/N510129/1. This study was also supported by the National Institute

for Health Research (NIHR) Biomedical Research Centre at the University Hospitals

Bristol NHS Foundation Trust and the University of Bristol (BRC-1215-2011). The

CATSS study is supported by the Swedish Foundation for International Cooperation in

Research and Higher Education (STINT), the Swedish Council for Health, Working Life

and Welfare, the Söderström-Königska Foundation, and the Swedish Research Council

(Medicine and SIMSAM). TEDS is funded by Medical Research Council grant

MR/M021475/1 to Robert Plomin. AbR is supported by the Beatrice and Samuel L

Seaver Foundation.

Author contributions

Conceptualization: ZER, OSPD; Methodology: ZER and OSPD; Software: ZER and OSPD;

Formal Analysis: ZER; Resources: OSPD, CMAH, ARo and PL; Data Curation: ZER;

Writing—Original Draft: ZER; Writing—Review and Editing: OSPD, HL, CMAH, DR, SL,

AnR, AbR and PL; Visualization: ZER and OSPD; Supervision: OSPD; Project

Administration: OSPD; Funding Acquisition: OSPD.

Competing interests

Angelica Ronald has received a consultancy fee for writing for the National Childbirth

Trust and receives an annual honorarium as joint editor of the Journal of Child

Psychology and Psychiatry. H. Larsson has served as a speaker for Evolan Pharma and

Shire and has received research grants from Shire; all outside the submitted work.