Relaxation of purifying selection suggests low effective population size in eusocial Hymenoptera and pollinating bees

Arthur Weyna[1], Jonathan Romiguier[1]

[1]Institut des Sciences de l'Evolution (UMR 5554), University of Montpellier, CNRS

Corresponding author: jonathan.romiguier@umontpellier.fr

ABSTRACT

Eusociality is a rare life history strategy that evolved repeatedly in Hymenoptera. At the population genetics level, inbreeding and low effective population size have been suggested as both evolutionary causes and consequences of social life. In this study, we tested these hypotheses by estimating the relative rate of non-synonymous substitution in 169 species to investigate the variation in natural selection efficiency and effective population size throughout the Hymenoptera tree of life. We show that relaxed selection is strongly associated with eusociality. This suggests that the division of reproductive labour decreases effective population size in ants, bees and wasps. Interestingly, we also report a striking and widespread relaxation of selection in both social and non social bees, which indicates that these keystone pollinator species generally feature low effective population sizes. We suggest that a particularly high inbreeding rate in bees might increase the benefits of kin selection, which would explain why most independent origins of eusociality in the tree of life occurred in this taxon. The particularly high load of deleterious mutations we report in the genome of these crucial ecosystem engineer species suggest new concerns about their ongoing population decline.

INTRODUCTION

The intensity of the genetic drift experienced by a population depends on its effective population size Ne (Wright 1931). Deleterious mutations reach fixation with a higher probability in small populations, which undergo more drift, than in large populations in which purifying selection is more efficient. Ne is usually defined for any observed population as the theoretical census size an ideal Wright-Fisher population should have to show the level of drift displayed by the observed population (Wang et al. 2016). While different definitions of Ne exist depending on the field, it generally correlates negatively with any process breaking the assumption of panmixia which underlies the Wright-Fisher model (i.e. population structure, homogamy, inbreeding…). Building on this knowledge, it has been proposed that basic traits influencing the reproductive output and mating choices of organisms, such as life-history traits, should correlate with their genome-wide deleterious substitutions rates. Several examples confirming these predictions have been uncovered in the last two decades: species generation time, longevity or body mass is positively correlated with the genome-wide dN/dS (ratio of the non-synonymous substitution rate to the synonymous substitution rate) (Nikolaev et al. 2007; Popadin et al. 2007; Romiguier et al. 2014; Figuet et al. 2016; Botero-Castro et al. 2017, Rolland et al. 2020). However, most known examples are clustered within a few Vertebrate taxa: mainly mammals, birds or reptiles. To date, only few examples of such patterns have been found in invertebrates, which cast doubt on the existence of a general

relationship between life history strategies and the efficiency of natural selection in Metazoa. Various reasons might explain the difficulty to demonstrate such relationships in invertebrates: There is relatively less genomic data available than in mammals or birds, and most taxa are expected to feature larger and less variable long-term Ne. Additionally, gathering life-history data in a large number of non-model invertebrates can be difficult as they have generally received less attention than Vertebrates.

Among invertebrates, Hymenoptera display a particularly wide diversity of life-history strategies. Notably, they exhibit extreme lifestyles that can be predicted to strongly influence their reproductive output, and thus their long-term Ne. First, many species within this clade are parasites of plants (phytophageous) or other Arthropodes (parasitoids) (Mayhew 2016), which could shape their demography, as hosts population structure and size can influence the parasite's (Mazé-Guilmo et al. 2016). Second, Hymenoptera contain a large number of pollinators, such as bees, that are involved in keystone insect-plant mutualisms and strictly depends on a limited floral resource likely to be scattered in time and space. Finally, eusociality, which is a rare lifestyle in Animals, is relatively common in the Hymenoptera order with at least 9 independant apparitions (Hughes et al. 2008). Eusociality is characterized by the differentiation of reproductive and non-reproductive castes of individuals within a colony, which cooperate for resource acquisition and brood care (Crespi and Yanega 1995). Because reproduction is typically monopolized by few long-lived reproductive individuals (Keller and Genoud 1997), a decrease of long-term Ne, and of the efficiency of natural selection, is often believed to be a general consequence of eusociality (Bromham and Leys 2005; Romiguier et al. 2014; Settepani et al. 2016). Without excluding the previous hypothesis, low Ne and inbreeding has also been raised as a potential *prerequesite*, facilitating evolution towards eusociality by inflating within-group relatedness and the benefit of kin selection (Hamilton 1972; Husseneder et al. 1999; Hughes et al. 2008; Tabadkani et al. 2012). Ancestral bottlenecks associated with such inbreeding episodes could thus be a typical feature of taxa in which eusociality frequently evolves.

Some early studies have tried to detect Ne decreases in eusocial lineages by comparing their heterozygosity (Berkelhamer 1983; Owen 1985; Reeve et al. 1985) with that of solitary lineages. These studies did not, however, allow a consensus to be reached, maybe in part because of the limiting size of the genetic data set available at the time. A more recent study showed that the genome-wide genetic diversity of four eusocial lineages was closer to that of mammals and birds(Romiguier et al. 2014) than to that of four solitary insects, suggesting a reduced Ne in eusocial lineages. This study's results are to be taken with care however, as the number of compared taxa is low, and as none of the solitary reference taxa are Hymenoptera, while almost all compared eusocial taxa belong to this group. Indeed, Ne could be inherently reduced in Hymenoptera, irrespective of eusociality, due to their haploid males (Wang et al. 2016). A study accommodating this problem by comparing genetic data for eusocial and solitary sister species in Hymenoptera and other taxa did not reveal any significant effect of eusociality on molecular evolution (Bromham and Leys 2005). This represents the strongest argument to date against a systematic reduction of Ne in eusocial lineages, as pair comparisons allow to correct for the effects of ploïdy and phylogeny, and as implicated pairs are numerous (25) and distributed over most known eusocial lineages. However, the small size of the genetic dataset (1 to 3 genes) in this study might have made it statistically difficult to detect any effect.

Here we estimated the efficiency of purifying selection in a phylogenomic dataset of 3256 genes in 169 Hymenoptera species (Peters et al. 2017), including 5 independent origins of eusociality (Formicidae : 3 species; Polistinae/Vespinae wasps : 3 species; Stenogastrinae wasps : 1 species; Corbiculate bees : 3 species; Halictidae : 2 species), with the goal of disentangling the links between effective population size, deleterious mutation load, and causes and consequences of eusociality.

RESULTS

dN/dS distribution across the Hymenoptera phylogeny
We estimated dN/dS in 3241 gene alignments of 169 Hymenoptera species using the mapNH program (Romiguier et al. 2012, https://github.com/BioPP/testnh) from the *testnh* program suite (Dutheil and Boussau 2008; Guéguen and Duret 2018). Average genomic dN/dS values are displayed along the Hymenoptera tree on Figure 1. We used this tree obtained by Peters et al. (2017) and its topology thorough all analyses. The largest and smallest mean ratios were inferred for *Eucera nigrescens* (0.0745) and *Macrocentrus marginator* (0.0318). As expected for conserved coding regions, the distribution of genomic dN/dS ratios is close to 0 (overall average of 0.0424), indicative of the large prevalence of purifying selection. We observed above average dN/dS ratios in all 5 available eusocial clades: Formicidae (0.0463 ± 0.0019, 3 species), Polistinae/Vespinae wasps (0.0492 ± 0.0023, 3 species), Stenogastrinae (0.0447, 1 species only), the *Apis/Bombus/Tetragonula* clade (0.0511 ± 0.0057), and Halictidae (0.0613 ± 0.0039; 2 species). These last two clades of bees do not clearly stand out however, as most bees in the dataset (Antophila, species characterized by pollen feeding of larvae: Apidae, Megachilidae, Halictidae, Colettidae, Andrenidae, and Melittidae) show high dN/dS ratios (0.0522 ± 0.0073, 41 species) with no clear dependence on their social organization. Finally, only two purely solitary taxa displayed comparable dN/dS ratios: Siricoidea (0.0466 ± 0.0027, 3 species) and Cynipoidea (0.0447 ± 0.0059, 5 species). We further used simple linear modelling to try and relate variation in dN/dS ratios to life history traits. When needed, phylogenetic independent contrasts where used to account for phylogenetic relationships (Felsenstein 1985).

Effect of body size and parasitism on relative protein evolution rates
As in birds and mammals (Figuet et al. 2016; Botero-Castro et al. 2017), we found a significant (df=131, F=5.2647, p-val = 0.0234) and positive (slope = 1.844e-04) effect of body size on dN/dS ratio in Hymenoptera. This effect only explains a very small part of the observed variation (R^2 = 0.03864) but remains significant when taking phylogeny into account (df=130, F=4.3121, p-val = 0.03981, slope = 1.542e-04, R^2 = 0.0321). When testing for a difference in dN/dS ratios between parasitic (parasitoid or parasites) and free-living Hymenoptera, we found a significant effect (df= 167, F= 46.327, p-val = 1.715e-10, R^2 = 0.2172), but which completely disappears when taking phylogeny into account (df= 166, F= 1.2111, p-val = 0.2727, R^2 = 0.007243). We thus interpret this as being a confounding effect of sampling disequilibrium, as groups with elevated ratios completely lack parasite (with the exception of the cuckoo bumblebee *Bombus rupestris*). We further tried and test for an association between dN/dS ratios of reproductive strategy and diet specialization within parasitoids using life-history and host range descriptors found in the litterature (Jervis et al. 2003; Traynor and Mayhew 2005a; Traynor and Mayhew 2005b; Mayhew 2016), and summarized in table S1. However these descriptors were very seldom available for the species contained in the present phylogenomic dataset, forcing us to use genus-level averaging for both traits and dN/dS ratios. We detected no siginificant associations between average dN/dS ratios and life-history in parasitoids at the genus-level.

Eusociality is associated to accelerated relative protein evolution rates
The social status of a terminal branch (eusocial or solitary species at tip) significantly explains dN/dS variations (df = 167; F = 15.403; p-val = 0.00012, R^2=0.0844) and this effect gets clearer when accounting for body size in the model (df = 130; F =17.7127; p-val = 4.757e-05, R^2=0.1153). Depending on the exact definition of eusocial branches, this effect can be robust to phylogenetic corrections (df=129, F=9.2684, p-val = 0.002827, R^2=0.0648) or not (df = 129, F=0.17, p-val=0.68083, R^2=0.0012). More specifically, the effect is still significant only when considering terminal branches leading to *Halictus* and *Lassioglossum* genera as eusocial. We argue that these

branches are likely to reflect eusocial evolutionary dynamics, because *Halictus* and *Lassioglossum* are known to originate from an eusocial ancestor (Cardinal and Danforth 2011) in spite of the secondary loss of eusociality in the specific species of our dataset. Table 1 presents results obtained when constructing full models accounting for all mentioned effects jointly, as well as results obtained when considering *Halictus* and *Lassioglossum* genera as solitary.

Antophila bees displays highly accelerated relative protein evolution rates
High dN/dS ratios in Antophila bees is by far the main pattern observed in our results. Treating appartenance to Antophila as a covariable allows to significantly explain (df = 167; F = 175.84; p-val < $2.2.10^{-16}$) more than half the observed variation (R^2 = 0.5129). Despite Antophila being only one monophyletic group, this effect is still present when accounting for phylogeny and adult size (df = 139; F = 6.4645; p-val < 0.01219). As five eusocial species in the dataset belong to Antophila and were susceptible to drive the eusociality effect, we also validated the presence of an effect of eusociality when accounting for the Antophila effect (see table 1).

Controlling for branch lengths, biased gene conversion or species sampling
Short terminal branches are known to bias dN/dS estimations upward because they yield more inaccurate estimations of this parameter, which real value is often close to its zero boundary at a genomic scale. There is some association between dN/dS ratios and branch length in this study (df = 167; F = 4.0492; p-val = 0.0458, R^2=0.02367) but it can not be responsible for the Antophila and eusociality effect as taking branch length into account does not nullifie any effects (see Table 1).

As eusocial Hymenoptera are known to have high recombination rates (Wilfert et al. 2007; Sirviö et al. 2011; Wallberg et al. 2015), which in turn are known to inflate dN/dS when associated to biased gene conversion in Vertebrates (Duret and Galtier 2009; Lartillot 2013; Galtier et al. 2018), we further validated these results using dN/dS estimated considering GC-conservative substitutions only, and which should therefore be impervious to the effects of biased gene conversion (Galtier et al. 2018). Average corrected genomic dN/dS values are displayed along the Hymenoptera tree on figure S1. All previous effects were recovered using these dN/dS ratios, with the exception of the adult size effect which is not significant when taking the phylogeny into account (table S2).

Four of the 12 available eusocial Hymenoptera (*Apis mellifera* and the three available ants), are species with published and annotated genomes (Consortium and The Honeybee Genome Sequencing Consortium 2006; Bonasio et al. 2010; Nygaard et al. 2011), and were used by Peters et al. (2017) as reference species for the identification of 1-1 orthologous genes, along with only one solitary reference species *Nasonia vitripennis (Werren et al. 2010)* This translated into a better power for gene prediction by Peters et al. (2017) in eusocial species, and thus into an over-representation of these eusocial species (mean number of alignments available per species: 2732.40 ± 88.09) as compared to solitary species (2276.7 ± 90.74) that is significant (T = 3.0567, df = 9.3549, p-value = 0.01305).To control for a potential bias originating from varying precision in estimations, we validated all our results on a balanced subsampled dataset containing 134 alignments, each of them containing data for the same 88 species (most represented half of the species). Average corrected genomic dN/dS values are displayed along the Hymenoptera tree on figure S2. Both the effect of eusociality and of Antophila were recovered with this dataset (table S3).

Eusocial taxa and Antophila bees experienced a relaxation of purifying selection
To ensure that previous results stem from a relaxation of selection and not from strong positive selection, we applied the Hyphy RELAX procedure (Pond et al. 2005; Wertheim et al. 2015) on each available alignment separately. This procedure allows to formally test for selection relaxation by modelling the distribution of dN/dS ratios along a phylogeny and by comparing the distribution

fitted on a focal group of branches (eusocial taxa and Antophila alternatively) to the distribution fitted for the rest of the tree.

Out of 3236 realized tests, 1743 (53.9%) detected relaxed selection on eusocial branches and 184 (5.7%) detected intensified selection. Genes under relaxation of selection thus represent 90% of the genes for which a difference of selection efficiency between eusocial branches and focal branches could be detected. Results of a gene ontology enrichment analysis conducted with genes under intensified selection in eusocial species as focal genes are presented in table S3. Using a conservative bonferroni correction for multiple testing in this procedure still leads to the detection of selection relaxation in 751 genes and of selection intensification in 28 genes. These results also hold if the more balanced subsampled dataset is used, as out of 134 alignments, 68 genes supported a relaxation of selection and 16 genes supported an intensification of selection. Moreover, the detected effect of eusociality doesn't seem to be driven by any over-representation of bees within eusocial species. The average number of eusocial bee sequences available for genes with relaxed selection (2.427 +- 0.018) is not different than within genes without relaxed selection (2.463 +- 0.024) (F=2.11; pval=0.146).

These verifications are needed because bees experience an even stronger relaxation of selection. If this was apparent from simple modelling of genomic dN/dS ratios, it is made even more obvious by the application of the RELAX procedure with Antophila branches as focal branches. Out of 3239 realized tests, 2000 (61.74%) detected relaxed selection on eusocial branches, while 294 detected an intensification of selection (9.07%). Using a conservative bonferroni correction for multiple testing in this procedure still leads to the detection of selection relaxation in 1210 genes and of selection intensification in 66 genes.

DISCUSSION

Molecular consequences of eusociality
We observed a significant higher accumulation of non-synonymous substitutions in eusocial genomes. This increase was associated with the detection of relaxed selection on eusocial branches for more than half of the available alignments. This result supports the hypothesis of a relaxation of selection associated with eusociality through demographic effects, as any other selective effects would be expected to have more heterogenous and localised effects.

Because eusocial Hymenoptera are known to display very high recombination rates (Wilfert et al. 2007; Sirviö et al. 2011; Wallberg et al. 2015), these results could have been imputed to biased gene conversion, which is known to increase dN/dS by promoting the fixation of any G/C alleles (including deleterious alleles). Biases in dN/dS estimation due to gBGC can be excluded however, as our results are not qualitatively changed by the use of dN/dS ratios computed using only GC-conservative substitutions. This is confirmed by RELAX analyses which would be expected to have detected gBGC effects as directional/adaptive selection, because both signals are very similar (Rousselle et al. 2019).

We rather suggest that the high dN/dS ratios we report are primarily due to *Ne* reduction and purifying selection relaxation in eusocial lineages. Long-lived reproductive female with delayed sexual maturity, as well as a biased sex-ratio and monopolization of the reproductive labour by few individuals, are typical features of eusocial species, which are bound to reduce effective population size. The hypothesis of a life-history effect matches well with the observation of a higher *dN/dS* in the highly eusocial formicoids ants *Acromyrmex echinatior* and *Camponotus floridanus* than in *Harpegnatos saltator,* which possesses a less complex social organization (Hölldobler and Wilson 1990). Ants, which display a variety of complexity levels in their social organisation, could

represent an ideal model for a more quantitative approach (Bourke, 1999), allowing to test for an effect of eusociality complexification on selection efficiency. We observed a clear effect of adult body size on relative non-synonymous substitution rates, an observation already made in vertebrates and usually interpreted as the effect of life history strategies on demographic parameters (Ne) and natural selection efficiency (dN/dS). In this study, we show that the emergence of a new level of organismic complexity (eusocial life history strategy) has a similar effect. One logical next step will be to investigate the existence of such associations within eusocial organisms, along a gradient of complexity in colonial organisation.

However, the hypothesis of such an effect of the level of sociality on natural selection efficiency matches somewhat badly with the observation of a smaller increase of dN/dS ratios in highly eusocial and long-lived ants than in less highly eusocial wasps. This hypothesis alone also fails in explaining why the dN/dS ratio shows no additional increase in social bees when compared to solitary ones.

Ecological and molecular predisposition to eusociality in bees
A high genomic dN/dS ratios in all social and solitary bees is a clear pattern in our results. Interestingly, Antophila (bees) is the taxa with the highest number of independent origins of eusociality in the tree of life (Hughes et al. 2008). This suggests that high dN/dS might be not only a consequence of group-living, but might also be associated to pre-requisites facilitating evolution toward eusociality. Several ecological characteristics have been suggested as facilitating the emergence of eusociality in the litterature, such as communal behaviors in nesting, the ability to defend such communal nests, parental care, progressive provisioning of food to the larvae or co-occurrence of nesting/feeding sites (Wilson 2008). Interestingly, some of these ecological characteristics can be linked to Ne, such as communal behaviors in nesting and parental care, that theoretically increase within-group relatedness through inbreeding, thus reducing Ne and increasing dN/dS. While the idea is controversial, it has been suggested that inbreeding itself might directly favor the evolution of eusociality by strengthening the advantages of within-group altruism (Hamilton 1972; Hussedener et al. 1999, Tabadkani et al., 2012, but see Nowak et al. 2010). Whether it be direct or indirect, inbreeding is thus clearly believed to be associated with the evolution of eusociality, but few genomic evidences supporting such a link have been observed so far. By showing a striking increase in dN/dS ratio in all Antophila bees - the taxa concentrating more than half of the origins of eusociality in the tree of life - our results are the first genomic insight supporting the idea that high inbreeding rates might have preceded and/or favoured evolution towards eusociality.

The fact that the increase of dN/dS precedes eusociality in bees, the hymenopteran group where eusociality has appeared the most frequently, also leaves one to wonder whether the relaxation of selection itself could have had a role in eusociality emergence. This might be seen as an argument supporting the idea that analogously to some hypotheses for the transition from unicellularity to multicellularity (Lynch and Conery 2003; Lynch 2007), eusociality can be seen as a complex trait that can evolve only in taxa with low Ne, where selection is not strong enough to maintain simpler organisations.

Besides many independent transitions toward eusociality, Antophila are also characterized by their pollen-collecting behaviors. This dependence to large amounts of pollen to feed their larvae is believed to be a potential constraint on *Ne*, and this particularly in specialist species, an argument which has been made previously (Zayed and Packer 2007). Pollen is a resource which is scattered in space and time, and which can require a large energetic investment to come by and exploit (through progressive provisioning), thus constraining the very fecundity of female individuals. If pollen-collecting is linked to Ne variation in Hymenoptera, it might help explaining our results, where dN/dS are similarly high between social and solitary pollen-collecting species. Indeed, group-living

might enhance the productivity of pollen collecting and metabolizing, thus effectively compensating the decrease of Ne linked to eusociality in Antophila. A global survey of Hymenoptera fecundity would be necessary to test this hypothesis, even if such tests would be quite limited by the fact that Antophila is the only pollen-collecting taxa in Hymenoptera.

Besides their implication regarding the evolution of eusociality, our results have important consequences for the conservation field. Indeed, pollination has been found to rely heavily on wild and domesticated bees, which ensure the majority of animal-mediated pollination of wild and domesticated plants in most ecosystems (Winfree 2010). Our finding of particularly high deleterious substitution rates within this group raises the additional concern that bee species might be especially sensitive to any further population decline, which are already known as particularly alarming (Powney et al. 2019; Murray et al. 2009; Arbetman et al. 2017).

Conclusion

This study brings new and genomic evidence supporting the existence of an association between eusociality and reduced selection efficiency, through reduced effective population size. Interestingly, the observation of a widespread reduction of selection efficiency in eusocial and solitary bees suggests that the picture might not be as clear-cut as thought previously. It is indeed possible that this reduction in effective population size is caused in part by ecological innovations that are not strictly eusocial characteristics, or that these reductions themselves have favored the complexification of social organisations. Besides the new questions raised by this study, this study brings the first genomic evidence suggesting that, as Vertebrates, invertebrates are constrained in their molecular evolutionary dynamics by their ecology and reproductive strategy.

MATERIALS AND METHODS

Genetic data

Data was downloaded from the authors' online repository (http://dx.doi.org/10.17632/trbj94zm2n.2). It originally contained nucleotide and amino-acids multi-sample alignments for 3256 protein coding genes predicted to be 1-1 orthologs in 174 species (see Peters et al. 2017 for details about the production of these alignments), 5 of which are outgroups to the Hymenoptera (2 Coleoptera, 1 Megaloptera, 1 Neuroptera and 1 Raphidioptera), and 12 of which are eusocial species. The latter belong to 5 independent eusocial clades: corbiculate bees (*Tetragonula carbonaria*, *Bombus rupestris* and *Apis mellifera*), ants (*Acromyrmex echinatior*, *Camponotus floridanus* and *Harpegnathos saltator*), Polistinae/Vespinae wasps (*Vespa crabro*, *Vespula germanica* and *Polistes dominula*), Stenogastrinae wasps (*Parischnogaster nigricans*) and Halictine bees (*Lasioglossum xanthopus* and Halictus). *Lassioglossum xanthopus* and *Halictus quadricinctus* were considered as eusocial despite not being specifically reported as so in the litterature, because they belong to genera which were reported as descending from an eusocial ancestor (Cardinal and Danforth 2011).The data also contained the trees inferred using this data by the original authors. We used the dated chronogram inferred by the authors using amino-acid data throughout this study. This tree corresponds to their main results and is contained in the file dated_tree_aa_inde_2_used_in_Fig1.tre available on the authors' online repository.

Data cleaning

Each amino-acid alignment was first checked for potential false homology using HmmCleaner (Philippe et al. 2017; Di Franco et al. 2019) with default settings. The resulting maskings were then reported on corresponding nucleotide sequences using the reportMaskAA2NT program from the MASCE program suite (Ranwez et al. 2011). At this point, we discarded individual sequences containing less than 50% of informative site within one alignment.

dN/dS ratios estimation
Cleaned alignments were then used, along with the tree topology inferred by Peters et al. (2017) and the mapNH binary (Romiguier et al 2012, https://github.com/BioPP/testnh), to estimate synonymous and non-synonymous substitution rates along the branches of the Hymenoptera tree. MapNH allows a fast estimation of those rates by using tree-wide parameters obtained *a priori* by fitting a homogeneous model (YN98) to the data with the help of paml (Yang 1997), to parsimoniously map observed substitutions to the supplied topology. Estimated substitution counts for specific branches, obtained separately for each alignments, can then be summed to obtain genome-wide substitution rates. We used this method to obtain dN/dS ratios of terminal branches, susceptible to carry information about the long-term drift regime of extant lineages. 15 alignments did not contain enough data to allow correct convergence of the homogeneous model needed by mapNH.

Controlling for biased gene conversion
We produced a corrected dN/dS using only GC conservative substitutions to estimate dN/dS. This was achieved using a custom version of mapNH developed in our lab (Rousselle et al. 2019) which categorizes mapped substitutions into GC-conservative (GC->GC or AT->AT) and GC-modifying (AT->GC or GC->AT) substitutions, and uses only the former to compute dN/dS ratios. Ratios obtained this way show more sampling variance, as they are obtained from smaller substitution counts. This translates in higher genomic dN/dS, as this paramter is usually close to its zero bound in exons. These rates are however supposedly impervious to gBGC.

Controlling for sampling bias
Four Hymenoptera (*Apis mellifera* and the three ants), which represent nearly half the eusocial species considered, are species with published genomes. This translates into a better power for gene prediction and thus, into an over-representation of these species in the dataset. Imprecisions in dN/dS ratios estimations is in turn known to yield higher values, because the real value of this ratio in functional sequences is often close to its zero boundary. We thus applied an additional sub-sampling procedure, designed to correct for any potential bias in our estimations that could stem from variation in the quantity of information available for each species. We applied every analysis mentioned before to a reduced but complete dataset containing data only for the most represented half of the species (88 species), and only alignments containing information for each of these species (135 alignments).

Linear modelling of dN/dS ratios
Estimated rates, corrected rates and rates obtained from the subsampled dataset were then modelled through simple linear models using the R software environment, using adult size, social status (eusocial or solitary) and membership to Antophila as covariables. We also used this statistical setting to evaluate the effect of branch length. Short branches are known to bias dN/dS estimations upward because they yield more inaccurate and thus generally higher estimations of this parameter. The phylogenetic setting was taken into account by using phylogenetic independent contrast (Felsenstein 1981)).

To try and further uncover the potential links between dN/dS ratios and life-history within Hymenoptera, we also attempted to correlate dN/dS ratios with major descriptors of parasitic type within parasitoid Hymenoptera. These descriptors were gathered from databases designed to describe the reproductive strategy of parasitoids (Traynor & Mayhew 2005a, Traynor & Mayhew 2005b, Jervis & Ferns 2011; Mayhew 2016) and are summarized in table S1. We conducted the analysis at the genus level using genus-averaged dN/dS ratios and descriptors. This was necessary because the species-level concordance between databases was too low (only 6 species in common between the genomic database and the parasitoid life-history database). We used Pearson's linear

correlation coefficient for continuous descriptors and Kruskal-Wallis tests for discrete descriptors.

RELAX analyses

We used the RELAX procedure (Wertheim et al. 2015) from the HyPhy program suite (Pond et al. 2005) to test for the presence of a systematic relaxation of selection on branches belonging to eusocial groups (thereafter called "eusocial branches"), that is all branches descending from the ancestral node of one of the eusocial clade present in the dataset. Hyphy allows, for a specific sequence alignment, to model the distribution of dN/dS ratios along the branches of a tree. The RELAX procedure consists first in defining focal and background branches, associated with one focal and one background distribution of dN/dS ratios. It then consists in comparing a model where the two mentioned distribution are identical (null model, no differences between branch sets) to a model where the focal distribution is a power transform of the background distribution ($\omega_f = \omega_b^k$). Relaxation of selection is inferred when the second model appears superior based on a log-ratio test (differences between branch sets), and when the focal distribution is narrower than the background distribution (k parameter estimated to be less than 1). Indeed, strong selection is thought to produce both low (close to 0) and high (greater than 1) dN/dS ratios, while neutrality should produce rates close to 1. This test thus correctly takes into account the fundamental two-sided nature of dN/dS ratios. 20 alignments did not contain enough data to allow models necessary to the HyPhy RELAX procedure to be fitted with eusocial branches as background branches, and 17 alignments didn't allow the procedure with Antophila branches as background branches.

AKNOWLEDGEMENTS

We thanks Nicolas Galtier for advices during the writing of the manuscript and Laurent Keller for useful discussions.

REFERENCES

Arbetman MP, Gleiser G, Morales CL, Williams P, Aizen MA. 2017. Global decline of bumblebees is phylogenetically structured and inversely related to species range size and pathogen incidence. Proceedings of the Royal Society B: Biological Sciences [Internet] 284. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5543210/

Berkelhamer RC. 1983. Intraspecific genetic variation and haplodiploidy, eusociality, and polygyny in the Hymenoptera. Evolution 37:540–545.

Bonasio R, Zhang G, Ye C, Mutti NS, Fang X, Qin N, Donahue G, Yang P, Li Q, Li C, et al. 2010. Genomic comparison of the ants *Camponotus floridanus* and *Harpegnathos saltator*. Science 329:1068–1071.

Botero-Castro F, Figuet E, Tilak M-K, Nabholz B, Galtier N. 2017. Avian genomes revisited: hidden genes uncovered and the rates versus traits paradox in birds. Mol. Biol. Evol. 34:3123–3131.

Bromham L, Leys R. 2005. Sociality and the rate of molecular evolution. Mol. Biol. Evol. 22:1393–1402.

Cardinal S, Danforth BN. 2011. The Antiquity and Evolutionary History of Social Behavior in Bees. PLoS ONE [Internet] 6:e21086. Available from: http://dx.doi.org/10.1371/journal.pone.0021086

Consortium THGS, The Honeybee Genome Sequencing Consortium. 2006. Insights into social insects from the genome of the honeybee Apis mellifera. Nature [Internet] 443:931–949. Available from: http://dx.doi.org/10.1038/nature05260

Crespi BJ, Yanega D. 1995. The definition of eusociality. Behavioral Ecology [Internet] 6:109–115. Available from: http://dx.doi.org/10.1093/beheco/6.1.109

Di Franco A, Poujol R, Baurain D, Philippe H. 2019. Evaluating the usefulness of alignment filtering methods to reduce the impact of errors on evolutionary inferences. BMC Evol. Biol. 19:21.

Duret L, Galtier N. 2009. Biased gene conversion and the evolution of mammalian genomic landscapes. Annu. Rev. Genomics Hum. Genet. 10:285–311.

Dutheil J, Boussau B. 2008. Non-homogeneous models of sequence evolution in the Bio++ suite of libraries and programs. BMC Evol. Biol. 8:255.

Felsenstein J. 1985. Phylogenies and the Comparative Method. The American Naturalist [Internet] 125:1–15. Available from: http://dx.doi.org/10.1086/284325

Figuet E, Nabholz B, Bonneau M, Mas Carrio E, Nadachowska-Brzyska K, Ellegren H, Galtier N. 2016. Life History Traits, Protein Evolution, and the Nearly Neutral Theory in Amniotes. Mol. Biol. Evol. 33:1517–1527.

Galtier N, Roux C, Rousselle M, Romiguier J, Figuet E, Glémin S, Bierne N, Duret L. 2018. Codon usage bias in animals: disentangling the effects of natural selection, effective population size, and GC-Biased gene conversion. Mol. Biol. Evol. 35:1092–1103.

Guéguen L, Duret L. 2018. Unbiased estimate of synonymous and nonsynonymous substitution rates with nonstationary base composition. Molecular Biology and Evolution [Internet] 35:734–742. Available from: http://dx.doi.org/10.1093/molbev/msx308

Hamilton WD. 1972. Altruism and related phenomena, mainly in Social Insects. Annual Review of Ecology and Systematics [Internet] 3:193–232. Available from: http://dx.doi.org/10.1146/annurev.es.03.110172.001205

Hölldobler B, Wilson EO. 1990. The Ants. Available from: http://dx.doi.org/10.1007/978-3-662-10306-7

Hughes WOH, Oldroyd BP, Beekman M, Ratnieks FLW. 2008. Ancestral monogamy shows kin selection is key to the evolution of eusociality. Science [Internet] 320:1213–1216. Available from: http://dx.doi.org/10.1126/science.1156108

Husseneder, Husseneder, Epplen JT, Brandl, Epplen, Kaib. 1999. Within-colony relatedness in a termite species: genetic roads to eusociality? Behaviour [Internet] 136:1045–1063. Available from: http://dx.doi.org/10.1163/156853999501702

Jervis MA, Ferns PN, Heimpel GE. 2003. Body size and the timing of egg production in parasitoid wasps: a comparative analysis. Functional Ecology [Internet] 17:375–383. Available from: http://dx.doi.org/10.1046/j.1365-2435.2003.00742.x

Keller L, Genoud M. 1997. Extraordinary lifespans in ants: a test of evolutionary theories of ageing. Nature [Internet] 389:958–960. Available from: http://dx.doi.org/10.1038/40130

Lartillot N. 2013. Phylogenetic patterns of GC-biased gene conversion in placental mammals and the evolutionary dynamics of recombination landscapes. Mol. Biol. Evol. 30:489–502.

Lynch M. 2007. The frailty of adaptive hypotheses for the origins of organismal complexity. Proc. Natl. Acad. Sci. U. S. A. 104 Suppl 1:8597–8604.

Lynch M, Conery JS. 2003. The origins of genome complexity. Science [Internet] 302:1401–1404. Available from: http://dx.doi.org/10.1126/science.1089370

Mayhew PJ. 2016. Comparing parasitoid life histories. Entomologia Experimentalis et Applicata [Internet] 159:147–162. Available from: http://dx.doi.org/10.1111/eea.12411

Mazé-Guilmo E, Blanchet S, McCoy KD, Loot G. 2016. Host dispersal as the driver of parasite genetic structure: a paradigm lost? Ecol. Lett. 19:336–347.

Murray TE, Kuhlmann M, Potts SG. 2009. Conservation ecology of bees: populations, species and communities. Apidologie [Internet] 40:211–236. Available from: http://dx.doi.org/10.1051/apido/2009015

Nikolaev SI, Montoya-Burgos JI, Popadin K, Parand L, Margulies EH, National Institutes of Health Intramural Sequencing Center Comparative Sequencing Program, Antonarakis SE. 2007. Life-history traits drive the evolutionary rates of mammalian coding and noncoding genomic elements. Proc. Natl. Acad. Sci. U. S. A. 104:20443–20448.

Nygaard S, Zhang G, Schiøtt M, Li C, Wurm Y, Hu H, Zhou J, Ji L, Qiu F, Rasmussen M, et al. 2011. The genome of the leaf-cutting ant *Acromyrmex echinatior* suggests key adaptations to advanced social life and fungus farming. Genome Res. 21:1339–1348.

Owen RE. 1985. Difficulties with the interpretation of patterns of genetic variation in the eusocial Hymenoptera. Evolution [Internet] 39:201–205. Available from: http://dx.doi.org/10.1111/j.1558-5646.1985.tb04093.x

Peters RS, Krogmann L, Mayer C, Donath A, Gunkel S, Meusemann K, Kozlov A, Podsiadlowski L, Petersen M, Lanfear R, et al. 2017. Evolutionary history of the Hymenoptera. Curr. Biol. 27:1013–1018.

Philippe H, de Vienne DM, Ranwez V, Roure B, Baurain D, Delsuc F. 2017. Pitfalls in supermatrix phylogenomics. European Journal of Taxonomy [Internet]. Available from: http://dx.doi.org/10.5852/ejt.2017.283

Pond SLK, Frost SDW, Muse SV. 2005. HyPhy: hypothesis testing using phylogenies. Bioinformatics 21:676–679.

Popadin K, Polishchuk LV, Mamirova L, Knorre D, Gunbin K. 2007. Accumulation of slightly deleterious mutations in mitochondrial protein-coding genes of large versus small mammals. Proc. Natl. Acad. Sci. U. S. A. 104:13390–13395.

Powney GD, Carvell C, Edwards M, Morris RKA, Roy HE, Woodcock BA, Isaac NJB. 2019. Widespread losses of pollinating insects in Britain. Nat. Commun. 10:1–6.

Rahnenfuhrer J, Alexa A. 2019. topGO: Enrichment Analysis for Gene Ontology. Bioconductor. Published online.

Ranwez V, Harispe S, Delsuc F, Douzery EJP. 2011. MACSE: Multiple Alignment of Coding SEquences accounting for frameshifts and stop codons. PLoS One 6:e22594.

Reeve HK, Reeve JS, Pfennig DW. 1985. Eusociality and genetic variability: a re-evaluation. Evolution 39:200–201.

Rolland J, Schluter D, Romiguier J. 2020. Vulnerability to fishing and life-history traits correlate with the load of deleterious mutations in teleosts. Mol. Biol. Evol. https://doi.org/10.1093/molbev/msaa067

Romiguier J, Figuet E, Galtier N, Douzery EJ, Boussau B, Dutheil JY, Ranwez V. 2012. Fast and robust characterization of time-heterogeneous sequence evolutionary processes using substitution mapping. *PLoS One*, *7*(3).

Romiguier J, Lourenco J, Gayral P, Faivre N, Weinert LA, Ravel S, Ballenghien M, Cahais V, Bernard A, Loire E, et al. 2014. Population genomics of eusocial insects: the costs of a vertebrate-like effective population size. Journal of Evolutionary Biology [Internet] 27:593–603. Available from: http://dx.doi.org/10.1111/jeb.12331

Rousselle M, Laverré A, Figuet E, Nabholz B, Galtier N. 2019. Influence of recombination and GC-biased gene conversion on the adaptive and nonadaptive substitution rate in mammals versus birds. Mol. Biol. Evol. 36:458–471.

Settepani V, Bechsgaard J, Bilde T. 2016. Phylogenetic analysis suggests that sociality is associated with reduced effectiveness of selection. Ecol. Evol. 6:469–477.

Sirviö A, Johnston JS, Wenseleers T, Pamilo P. 2011. A high recombination rate in eusocial Hymenoptera: evidence from the common wasp Vespula vulgaris. BMC Genet. 12:95.

Tabadkani SM, Nozari J, Lihoreau M. 2012. Inbreeding and the evolution of sociality in arthropods. Naturwissenschaften 99:779–788.

Traynor RE, Mayhew PJ. 2005a. A comparative study of body size and clutch size across the parasitoid Hymenoptera. Oikos [Internet] 109:305–316. Available from: http://dx.doi.org/10.1111/j.0030-1299.2005.13666.x

Traynor RE, Mayhew PJ. 2005b. Host range in solitary versus gregarious parasitoids: a laboratory experiment. Entomologia Experimentalis et Applicata [Internet] 117:41–49. Available from: http://dx.doi.org/10.1111/j.1570-7458.2005.00331.x

Wallberg A, Glémin S, Webster MT. 2015. Extreme recombination frequencies shape genome variation and evolution in the honeybee, *Apis mellifera*. PLOS Genetics [Internet] 11:e1005189. Available from: http://dx.doi.org/10.1371/journal.pgen.1005189

Wang J, Santiago E, Caballero A. 2016. Prediction and estimation of effective population size. Heredity [Internet] 117:193–206. Available from: http://dx.doi.org/10.1038/hdy.2016.43

Werren JH, Richards S, Desjardins CA, Niehuis O, Gadau J, Colbourne JK, Nasonia Genome Working Group, Werren JH, Richards S, Desjardins CA, et al. 2010. Functional and evolutionary insights from the genomes of three parasitoid Nasonia species. Science 327:343–348.

Wertheim JO, Murrell B, Smith MD, Kosakovsky Pond SL, Scheffler K. 2015. RELAX: Detecting Relaxed Selection in a Phylogenetic Framework. Molecular Biology and Evolution [Internet] 32:820–832. Available from: http://dx.doi.org/10.1093/molbev/msu400

Wilfert L, Gadau J, Schmid-Hempel P. 2007. Variation in genomic recombination rates among animal taxa and the case of social insects. Heredity 98:189–197.

Wilson EO. 2008. One giant leap: how insects achieved altruism and colonial life. BioScience [Internet] 58:17–25. Available from: http://dx.doi.org/10.1641/b580106

Winfree R. 2010. The conservation and restoration of wild bees. Ann. N. Y. Acad. Sci. 1195:169–197.

Wright S. 1931. Evolution in Mendelian Populations. Genetics 16:97–159.

Yang Z. 1997. PAML: a program package for phylogenetic analysis by maximum likelihood. Bioinformatics [Internet] 13:555–556. Available from: http://dx.doi.org/10.1093/bioinformatics/13.5.555

Zayed A, Packer L. 2007. The population genetics of a solitary oligolectic sweat bee, *Lasioglossum* (Sphecodogastra) *oenotherae* (Hymenoptera: Halictidae). Heredity 99:397–405.

		Halictus and *Lasioglossum* as solitary	*Halictus* and *Lasioglossum* as eusocial

		R^2	F	p-value	R^2	F	p-value
Original Data residual df = 128 R^2=0.5526	branch length	0.0199	5.4338	**0.021313**	0.0199	5.7165	**0.0182621**
	adult size	0.0349	9.4942	**0.002524**	0.0349	9.9881	**0.0019667**
	Antophila	0.4462	121.36	**< 2.2e-16**	0.4462	127.6745	**< 2.2e-16**
	Eusociality	0.0281	7.6648	**0.006468**	0.0514	14.7227	**0.0001947**
Phylogenetic Independent Contrasts residual df = 127 R^2=0.1688	branch length	0.0389	5.5767	**0.01972**	0.0389	5.9517	**0.016084**
	adult size	0.0269	3.8587	**0.05167**	0.0269	4.1182	**0.044514**
	Antophila	0.0455	6.5271	**0.01180**	0.0455	6.9660	**0.009348**
	Eusociality	0.00145	0.2089	0.64841	0.0573	8.7634	**0.003668**

Table 1: Results of simple linear modelling of dN/dS ratios. Displayed results are those obtained when simultaneously accounting for all effects inside a multiple linear model.

Figure 1: Genomic dN/dS ratios for 169 Hymenoptera species. dN/dS ratios estimated on terminal branches using 3241 genes are represented on the chronogram inferred by Peters *et al.* (2017). Green rectangles around labels indicate eusocial taxa.

Figure S1: Corrected genomic dN/dS ratios for 169 Hymenoptera species. dN/dS ratios estimated on terminal branches using 3241 genes and GC conservative substitutions are represented on the chronogram inferred by Peters *et al.* (2017). Green rectangles around labels indicate eusocial taxa. Note that the color scale differs from that of figure 1, corrected dN/dS being generally higher due to smaller sample size (less substitutions overall).

Figure S2: Genomic dN/dS ratios for 88 Hymenoptera species. dN/dS ratios estimated on terminal branches using 134 genes with data for each of the displayed species are represented on the chronogram inferred by Peters *et al.* (2017). Green rectangles around labels indicate eusocial taxa.

Variable		Description	Number of genus with information	correlation	p-val	source
	Number of mature eggs at emergence	Maximal observed number during dissections of emerged females	11	-0.048	0.889	Jervis & Ferns (2011) Mayhew (2016)

continuous	Maximal number of mature eggs	Maximal observed number during dissections	10	0.146	0.688	Jervis & Ferns (2011) Mayhew (2016)
	Ovigenic index	Ratio of the last two variables	10	0.114	0.711	Jervis & Ferns (2011) Mayhew (2016)
	Oviposition rate	Average number of eggs laid by one female in one day	10	0.176	0.627	Jervis & Ferns (2011)
	Egg length	Propagule size	3	-0.5	0.667	Traynor & Mayhew (2005a)
	Clutch size	Average number of eggs in one clutch	5	-0.707	0.182	Traynor & Mayhew (2005a)
	Fecundity	Average number of eggs laid by one female until death (continuous)	14	0.415	0.14	Jervis & Ferns (2011) Traynor & Mayhew (2005a)
	Developpement time : egg -> adult	Proxy of developmental speed	11	-0.2	0.55	Jervis & Ferns (2011)
	Time as an egg	Proxy of developmental speed	4	0.2	0.8	Traynor & Mayhew (2005a)
	Time as a pupa	Proxy of developmental speed	5	-0.3	0.624	Jervis & Ferns (2011) Traynor & Mayhew (2005a)
	Time as an adult	Proxy of developmental speed	14	0.226	0.436	Jervis & Ferns (2011) Traynor & Mayhew (2005a)
	Host body length	Loose proxy of ressource abundance	8	0.119	0.779	Traynor & Mayhew (2005b)
	Number of potential host species	Proxy of the level of specialization	10	-0.234	0.515	Traynor & Mayhew (2005b)
discrete	Egg hydropisy	High or low nutritional ressources in eggs	11	0.375	0.54	Jervis & Ferns (2011)
	Gregarious behavior	One or more larvae on one individual host	12	1.444	0.229	Jervis & Ferns (2011)
	Idiobiont/Koïnobiont	Parasitoid types	13	1.371	0.242	Jervis & Ferns (2011)
	Ecto/Endoparasitism	Parasitoid types	18	0.022	0.882	Jervis & Ferns (2011) Traynor & Mayhew (2005b)
	Hyperparasitism	Host is free or a parasite itself	12	0.185	0.667	Jervis & Ferns (2011)
	Host consumption behavior	Is Adult able to feed on host too ?	12	0.26	0.61	Jervis & Ferns (2011)
	Host developpement stage at infection	On which developmental stage is the host when eggs are laid.	10	3.836	0.28	Traynor & Mayhew (2005b)
	Host hiding behavior	How well is the host dissimulated	7	2.4	0.121	Traynor & Mayhew (2005b)

Table S1: Life-history and specialisation descriptors for parasitoids. Tested variables and their description are displayed along with the value of the statistic obtained for each correlation test with dN/dS ratios. Correlation tests are Spearman tests for continuous variables and Kruskal-Wallis tests for discrete variables.

		Halictus and *Lassioglossum* as solitary			*Halictus* and *Lassioglossum* as eusocial		
		R^2	F	p-value	R^2	F	p-value
Original Data	branch length	0.0149	3.7726	0.05430	0.015	3.9185	**0.049903**
	adult size	0.0241	6.0739	**0.01505**	0.0241	6.3089	**0.013257**

residual df = 128 R²=0.5526	Antophila	0.4451	112.2325	**< 2e-16**	0.4452	116.5747	**< 2.2e-16**
	Eusociality	0.008	2.0219	0.15748	0.027	7.0524	**0.008922**
Phylogenetic Independent Contrasts residual df = 127 R²=0.1688	branch length	0.0347	4.8040	**0.03022**	0.0347	5.0455	**0.02642**
	adult size	0.0085	1.1793	0.27956	0.0085	1.2386	0.26785
	Antophila	0.0383	5.3012	**0.02294**	0.0383	5.5676	**0.01982**
	Eusociality	0.0006	0.0897	0.76502	0.0446	6.4785	**0.01212**

Table S2: Results of simple linear modelling of corrected dN/dS ratios. Displayed results are obtained when using dN/dS ratios obtained using only GC conservative substitutions, and simultaneously accounting for all effects inside a multiple linear model.

		Halictus and *Lassioglossum* as solitary			*Halictus* and *Lassioglossum* as eusocial		
		R^2	F	p-value	R^2	F	p-value
Original Data residual df = 128 R²=0.5526	branch length	0.0159	2.1251	0.149378	0.0159	2.2648	0.1368421
	adult size	0.0654	8.7625	**0.004194**	0.0654	9.3386	**0.0031745**
	Antophila	0.3325	44.541	**4.882e-09**	0.3325	47.4704	**1.982e-09**
	Eusociality	0.0637	8.5353	**0.004686**	0.096	13.6993	**0.0004239**
Phylogenetic Independent Contrasts residual df = 127 R²=0.1688	branch length	0.0159	2.2860	0.13511	0.0282	2.3519	0.12970
	adult size	0.0654	2.2963	0.13425	0.0283	2.3625	0.12886
	Antophila	0.3325	5.2339	**0.02522**	0.0645	5.3847	**0.02328**
	Eusociality	0.0637	2.3086	0.13323	0.0523	4.3627	**0.04042**

Table S3: Results of simple linear modelling of dN/dS ratios. Displayed results are obtained when using dN/dS ratios obtained using the subsampled dataset (88 species and 134 genes), and simultaneously accounting for all effects inside a multiple linear model.

domain	GO ID	Term	p-val
biological process	GO:0043623	cellular protein complex assembly	0.00011
	GO:0016043	cellular component organization	0.00011
	GO:0043604	amide biosynthetic process	0.00012

molecular function	GO:0003723	RNA binding	0.00018
	GO:0008092	cytoskeletal protein binding	0.00030
	GO:0003735	structural constituent of ribosome	0.00083
	GO:0005488	binding	0.00109
	GO:0051020	GTPase binding	0.00113
	GO:0005085	guanyl-nucleotide exchange factor activity	0.00350
	GO:0017069	snRNA binding	0.00376
	GO:0019899	enzyme binding	0.00491
	GO:0005198	structural molecule activity	0.00500
	GO:0030246	carbohydrate binding	0.01089
	GO:0019904	protein domain specific binding	0.01286
	GO:0008536	Ran GTPase binding	0.02269
	GO:0003924	GTPase activity	0.02838
	GO:0017016	Ras GTPase binding	0.03243
	GO:0031267	small GTPase binding	0.03243

Table S4: Go terms enriched in genes supporting an intensification of selection in eusocial Hymenoptera. P-values are those of a Fisher hypergeometric test used for significance in the GO enrichment analysis, as implemented in the **R** package *topGO (Rahnenfuhrer and Alexa 2019)*