# Actin binding domain of Rng2 strongly inhibits actin movement on myosin II HMM through structural changes of actin filaments

**Yuuki Hayakawa[1, 6], Masak Takaine[2, 6, #], Taiga Imai[3], Masafumi Yamada[4], Keiko Hirose[4], Kiyotaka Tokuraku[3], Kien Xuan Ngo[5], Noriyuki Kodera[5], Osamu Numata[2], Kentaro Nakano[2,*], Taro Q.P. Uyeda[1, 4,*]**

[1] Department of Physics, Faculty of Science and Engineering, Graduate School of Waseda University, Shinjuku, Tokyo 169-8555, Japan

[2] Department of Biology, Degree Programs in Life and Earth Sciences, Graduate School of Science and Technology, University of Tsukuba, Tsukuba, Ibaraki 305-8572, Japan

[3] Department of Applied Sciences, Muroran Institute of Technology, Muroran, Hokkaido 050-8585, Japan

[4] Biomedical Research Institute, National Institute of Advanced Industrial Science and Technology, Tsukuba, Ibaraki 305-8565, Japan

[5] WPI Nano Life Science Institute, Kanazawa University, Kanazawa, Ishikawa 920-1192, Japan,

[6] These authors contributed equally.

[#] Present addresses: Gunma University Initiative for Advanced Research (GIAR) and Institute for Molecular and Cellular Regulation (IMCR), Gunma University, Maebashi 371-8512, Japan

* Correspondence: t-uyeda@waseda.jp and knakano@biol.tsukuba.ac.jp

# Abstract

The contraction of contractile rings (CRs) depends on sliding between actin filaments and myosin II filaments. The rate of contraction in the fission yeast *Schizosaccharomyces pombe* is less than 1/120 of the velocity of acto-myosin II movement *in vitro*, but the mechanism of inhibition has not been described. Here, we found that the calponin-homology actin binding domain of fission yeast IQGAP Rng2 (Rng2CHD) strongly inhibits the motility of actin filaments on skeletal muscle myosin II fragments *in vitro*, even at a low ratio of bound Rng2CHD to actin protomers, reducing the sliding velocity to half when the binding ratio was 1/75. Rng2CHD also induced structural changes of actin filaments and reduced the affinity between actin filaments and subfragment 1 (S1) of muscle myosin II carrying ADP. Intriguingly, actin-activated ATPase of S1 was only mildly inhibited, even by high concentrations of Rng2CHD. Moreover, the motility of actin filaments by myosin V was not inhibited by Rng2CHD. We propose a new regulatory mechanism for acto-myosin II movement that involves Rng2CHD-induced structural changes of actin filaments.

# Introduction

Actin exists in all eukaryotic cells and performs a wide variety of functions. One of these functions is the contraction of contractile rings (CRs). The CR is a ring-like structure that appears transiently on the equatorial plane underneath the cell membrane during cytokinesis of animal cells and many unicellular eukaryotes. CR mainly consists of actin filaments and myosin II filaments. It is thought that a CR contracts by sliding between the two filament systems due to actomyosin movement (Mabuchi & Okuno, 1977; Satterwhite & Pollard, 1992). However, how contraction is regulated remains ambiguous. For example, the circumferential contraction rate of the CR of the fission yeast *Schizosaccharomyces pombe*, a model organism for cytokinesis studies, is 0.5 μm/min (Pelham & Chang, 2002), while actin filaments on myosin II derived from *S. pombe* move at a rate of 30 μm/min *in vitro* (Lord & Pollard, 2004). This 60-fold difference in speed is most likely an underestimate of the extent of inhibition, as explained next. CRs have actin filaments of opposite polarities, as is expected for contraction in a manner similar to muscle sarcomeres or stress fibers, but they are not as regularly organized as in sarcomeres or stress fibers (Kamasaki *et al*, 2007). Nonetheless, to simplify the argument, if we assume that CRs consist of contraction units, each of which contains myosin II filaments and anti-parallel actin filaments, and there are N of those contractile units in series in a CR, then the maximum contraction rate would be the sliding velocity *in vitro* multiplied by a factor of 2N. Therefore, it is estimated that the contraction rate of the CR in *S.*

*pombe* is inhibited to less than 0.5/(30x2)=1/120 of the velocity of acto-myosin II movement *in vitro*, even when assuming the extreme case of N=1. In the cellular slime mold *Dictyostelium discoideum*, the contractile velocity of the CR is 0.18 μm/s (Zang *et al*, 1997), whereas the sliding velocity of actin filaments on myosin II derived from *D. discoideum* is 1.4 μm/s *in vitro* (Kron & Spudich, 1986). These results suggest that contraction of the CR is regulated in an inhibitory manner in cells with or without a cell wall. Two distinct mechanisms may contribute to this significant inhibition. First, the increase in membrane surface tension due to membrane deformation and high viscoelasticity of the cytoplasm may create high loads that slow down the movement speed of myosin II, as was demonstrated for skeletal muscle contraction (Hill, 1938). The other possibility is that myosin II motor activity is negatively regulated in CRs. In this study, we experimentally examined the latter hypothesis. Several actin-binding proteins (ABPs), such as formin, tropomyosin and Rng2, play important roles in the formation of CRs in *S. pombe* (Eng *et al*, 1998), and one of those ABPs may be involved in the inhibitory regulation of the contraction rate of CRs. In particular, Rng2, an IQ motif containing GTPase activating protein (IQGAP) with a calponin-homology actin binding domain (CHD) at the N-terminus (Fig. 1), is a prime candidate for this regulator. It is thought that Rng2 crosslinks and bundles actin filaments together to form CRs, because CRs were not formed in Rng2 knockout *S. pombe* cells even though actin filaments were assembled at their division sites (Eng *et al*, 1998, Takaine *et al*, 2009). Additionally, temperature-sensitive Rng2 mutant

cells showed normal actin filament rings at the permissive temperature but the distribution of myosin II on actin filaments was abnormal (Takaine *et al*, 2009). This result suggests that Rng2 regulates the interaction between actin filaments and myosin II filaments.

In this study, we performed *in vitro* motility assays in which actin filaments move on myosin II HMM to examine if Rng2 inhibits actomyosin motility. We found that CHD fragments of Rng2 (Rng2CHD) bound on actin filaments strongly inhibited acto-myosin II motility even when binding was sparse, without significantly inhibiting ATPase activity. Furthermore, the affinity of actin filaments sparsely decorated by Rng2CHD displayed a reduced affinity for the motor domain of myosin II carrying ADP, demonstrating that this is a novel mode of regulation of acto-myosin II motility.

# Results

**Rng2CHD strongly inhibits sliding of actin filaments on myosin II HMM but not on myosin V HMM *in vitro***

To examine if Rng2 inhibits actomyosin motility, we performed *in vitro* motility assays in which actin filaments move on a glass surface coated with heavy meromyosin (HMM) of skeletal muscle myosin II (Kron & Spudich, 1986). The movement of actin filaments was strongly inhibited by Rng2CHD, the actin-binding domain of Rng2, and was completely stopped in the presence of 200 nM Rng2CHD (Fig. 2A). A trajectory analysis showed that, under all the inhibition conditions tested, the front end of the actin filament was followed by the remainder of the filament (Fig. 2B). Buckling of the filaments, indicative of local inhibition of movement, was rarely observed. This indicates that movement is more or less uniformly inhibited along the entire length of the filaments.

We also performed *in vitro* motility assays in which actin filaments moved on recombinant myosin V HMM that was expressed in insect cells. In striking contrast with myosin II, up to 1 μM Rng2CHD did not inhibit the sliding of actin filaments on myosin V HMM (Fig. 2A).

**Rng2CHD on actin filaments inhibits acto-myosin II movement *in vitro* even when binding is sparse**

We estimated the binding ratio of Rng2CHD to actin protomers when the movement of

actin filaments on myosin II HMM was inhibited. The concentrations of Rng2CHD that caused 50%, 75% and 95% reduction of movement speed were estimated from the velocity curve to be 12.4 nM, 31.6 nM and 71.4 nM, respectively (Fig. 2A). In parallel, we performed co-sedimentation assays of actin filaments with Rng2CHD, and the dissociation constant ($K_d$) between Rng2CHD and actin protomers was calculated (Fig. 3A, 3C). $K_d$ was determined to be 0.92 μM by the following fitting function: $[Rng2CHD_{bound}] = [Actin_{total}][Rng2CHD_{free}]/[Rng2CHD_{free}] + K_d)$ (Eq. 1 in Materials and Methods). In *in vitro* motility assays, where the concentration of actin protomers is much lower than that of Rng2CHD, the following approximation holds: $[Rng2CHD_{bound}]/[Actin_{free}] \cong [Rng2CHD_{total}]/K_d$ (Eq. 3 in Materials and Methods). Using this approximation, the binding ratio of Rng2CHD to actin protomers that caused 50%, 75% and 95% reduction of acto-myosin II movement speed was estimated to be 1/75, 1/30 and 1/13, respectively (Table 1). In other words, Rng2CHD was sparsely bound to actin filaments (1.3% occupancy) when the speed was reduced to half, and 7.7% occupancy even when it inhibited 95% of movement.

To directly confirm that sparsely bound Rng2CHD potently inhibits actomyosin movements, we prepared Rng2CHD fused with GFP to its N-terminus via a 16-residue linker (GFP-Rng2CHD), and determined the binding ratio of GFP-Rng2CHD to actin protomers from the fluorescence intensity of GFP. GFP-Rng2CHD strongly inhibited actin filament movement on

myosin II HMM in a manner similar to Rng2CHD (Fig. 4A). However, the dissociation constant

determined by the co-sedimentation assay was 3.37 μM, which was three-fold higher than that of

Rng2CHD (Fig. 3B, 3D). The binding ratio of GFP-Rng2CHD to actin protomers estimated from $K_d$

was also higher than that of Rng2CHD when GFP-Rng2CHD and Rng2CHD caused the same

degree of movement inhibition (Table 1), implying that the mode of actin binding is somewhat

different between Rng2CHD and GFP-Rng2CHD. Nonetheless, we went on to measure the

fluorescence intensity of bound GFP-Rng2CHD per unit length of actin filaments. The fluorescence

intensity increased depending on the GFP-Rng2CHD concentration in the buffer, and was saturated

in the presence of 5,000 nM GFP-Rng2CHD (Fig. 4B, 4C). The co-sedimentation assay showed that

one molecule of GFP-Rng2CHD binds to one molecule of an actin protomer in the presence of 5,000

nM GFP-Rng2CHD (Fig. 3D). Therefore, we regarded this saturated fluorescence intensity as a

one-to-one binding state, and used it as the reference to calculate how much GFP-Rng2CHD binds to

actin protomers when movements were inhibited (Fig. 4D). Although this directly measured binding

ratio was higher than the value estimated from $K_d$, it was unambiguously demonstrated that sparsely

bound GFP-Rng2CHD on actin filaments potently inhibits acto-myosin II sliding.

**Rng2CHD locally kinks or distorts actin filaments**

Rng2CHD reduced the rate of acto-myosin II movement to 50% at a very low binding

ratio, estimated to be 1/75 from $K_d$ (Fig. 2A and Table 1). Thus, we inferred that sparsely bound

Rng2CHD somehow induces global structural changes in actin filaments and inhibits acto-myosin II

movement. To examine whether sparsely bound Rng2CHD actually changes the structure of actin

filaments, we observed actin filaments in the presence of Rng2CHD using negative stain electron

microscopy. In the absence of Rng2CHD, double helical filaments typical of actin filaments were

observed (Fig. 5A), but the helical structures became irregular and the filaments often looked locally

distorted in the presence of Rng2CHD (Fig. 5B-5D). In some portions of the filaments, the two actin

protofilaments appeared to be separated so that a dark straight line was observed in between

(indicated by yellow brackets), suggesting that the interactions between neighboring actin protomers

are altered. At a higher concentration (1 µM) of Rng2CHD, the filaments were frequently kinked or

even fragmented. The binding ratios of Rng2CHD to actin protomers were estimated to be 1/25 and

1/5 at 40 nM and 200 nM of Rng2CHD, respectively, based on $K_d$ (Eq. 2 in Materials and Methods).

These results indicate that when low concentrations of Rng2CHD was allowed to interact with actin

filaments, structural changes are induced even in actin protomers not in direct contact with

Rng2CHD molecules.

**Steady state actin-activated S1 ATPase is only weakly inhibited by Rng2CHD**

To gain insight into the mechanism by which structural changes of actin filaments induced

by Rng2CHD inhibits actomyosin motility, we investigated the effects of Rng2CHD on actin-activated ATPase activity of myosin II subfragment-1 (S1). Actin-activated S1 ATPase was moderately inhibited (approximately 50%) by the highest concentration of Rng2CD tested (5 µM; Fig. 6). In the presence of 0.33 µM, 0.82 µM and 1.91 µM of Rng2CHD and 23.8 µM of actin filaments, the binding ratios of Rng2CHD to actin were calculated to be 1/75, 1/30, and 1/13, respectively (Eq. 2 in Materials and Methods). These binding ratios caused a 50%, 75% and 95% reduction in the speed of actomyosin movement (Fig. 2A and Table 1). However, the inhibition of actin-activated S1 ATPase activity was much weaker and disproportional to the inhibition of movement (Fig. 6). This indicates that even when actomyosin movement was strongly inhibited or halted by Rng2CHD, the myosin II ATPase cycle was running, and was only partially inhibited.

**Rng2CHD inhibits the steady-state binding of S1 to actin filaments in the presence of ADP, but not in the presence of ATP**

We examined the possibility that Rng2CHD might affect the affinity between actin filaments and myosin II when it inhibits acto-myosin II motility. First, we performed a co-sedimentation assay of actin filaments and S1, and found that Rng2CHD did not significantly inhibit steady-state binding of S1 to actin filaments in the presence of ATP as in *in vitro* motility assays (Fig. 7A, 7C). However, Rng2CHD weakly but statistically significantly inhibited the binding

of S1• ADP to actin filaments in the presence of ADP in the buffer (Fig. 7B, 7C).

A cosedimentation assay using myosin II filaments showed that Rng2CHD did not bind to myosin II under the conditions employed in the *in vitro* motility assays (Supplementary Figure 1).

**Rng2CHD decreases the fluorescence of HMM-GFP along actin filaments in the presence of ATP**

We previously reported that when HMM of *D. discoideum* myosin II fused with GFP was allowed to interact with actin filaments in the presence of a very low concentration of ATP, HMM-GFP forms clusters along actin filaments (Tokuraku *et al*, 2009; Hirakawa *et al*, 2017). This was interpreted to represent local polymorphism of actin filaments, such that some sections of the filaments have a higher affinity for HMM than other parts of the filament, and HMM-GFP preferentially repeats transient binding to those sections. In the presence of a high concentration of ATP, no fluorescent clusters were observed, and in the absence of ATP, HMM-GFP uniformly bound along the entire filaments (Tokuraku *et al*, 2009).

. We speculated that, for the detectable fluorescent clusters to form, HMM needs to bind to actin filaments repetitively and transiently but the dwell time of each binding event must be long enough to allow visualization. Based on this hypothesis, we observed how Rng2CHD affects the formation of myosin II HMM-GFP clusters along actin filaments in the presence of a very low

concentration of ATP using fluorescence microscopy. Numerous fluorescent spots, each representing a HMM-GFP cluster, were observed along actin filaments in the presence of 0.5 µM ATP and the absence of Rng2CHD. In the presence of 1 nM Rng2CHD, the number and fluorescent intensity of fluorescent spots were significantly reduced, and fluorescent spots were apparently absent in the presence of 10 nM Rng2CHD (Fig. 8A). In contrast, Rng2CHD did not affect the binding of HMM-GFP to actin filaments in the nucleotide-free state (Fig. 8B).

**Rng2CHD significantly decreases the number of S1 molecules transiently bound to actin filaments in the presence of ATP**

We also directly observed transient binding of S1 molecules to actin filaments in the presence of ATP using high-speed atomic force microscopy (HS-AFM). At a scan speed of 0.5 s per field of view, transient binding of S1 to actin filaments was rarely observed in the presence of 500 µM ATP alone, but was frequently observed in the presence of 50 µM ATP and 1 mM ADP. S1 molecules were easily identified based on their size and shape, while Rng2CHD molecules were too small to visualize under our experimental condition. We analyzed images scanned between 1 and 2 min after the addition of S1 or ATP, and visually counted the number of transiently bound S1 molecules (Fig. 9A). The binding dwell time of S1 molecules on the top of the filament was shorter than those bound along the sides of the filaments. Therefore, we separately counted the S1 molecules

bound on the top and along the sides of the filaments. The number of S1 molecules that transiently

bound to actin filaments was significantly decreased when 12.4 nM Rng2CHD, the concentration

that caused 50% inhibition of motility, was added before the addition of S1 (Fig. 9B, 8C). It was thus

directly confirmed that sparsely bound Rng2CHD affected the binding of myosin II S1 to actin

filaments in the presence of ATP and ADP. In contrast, when S1 was strongly bound to actin

filaments in the nucleotide-free state in advance, and then 50 $\mu$M ATP, 1 mM ADP and 12.4 nM

Rng2CHD were added, the number of S1 molecules transiently bound to actin filaments was

unaffected by Rng2CHD (Fig. 9C). Thus, there is apparent hysteresis in the impact of Rng2CHD on

S1 binding to actin filaments, depending on whether S1 or Rng2CHD actin filaments first encounter.

# Discussion

**Structural changes of actin filaments induced by sparsely bound Rng2CHD inhibit acto-myosin II movement**

Rng2CHD, the actin-binding domain of Rng2, strongly inhibited acto-myosin II motility along the entire length of the filament, when actin filaments were estimated to be only sparsely decorated by Rng2CHD (Fig. 2A and Table 1). Furthermore, fluorescence measurements unambiguously showed that sparsely bound GFP-Rng2CHD strongly inhibited acto-myosin II movement (Fig. 4A, 4D). Since binding of Rng2CHD or GFP-Rng2CHD was sparse, the inhibition of motility could not be due to steric hindrance or direct competition for a binding site on actin molecules. We thus inferred that sparsely bound Rng2CHD and GFP-Rng2CHD induced some structural changes in actin filaments, and these inhibited the productive interaction between actin filaments and myosin II. Previous studies advocated that structural changes of actin filaments modulate affinities for ABPs. For example, cooperative structural changes of actin filaments induced by myosin II and cofilin contribute to mutually exclusive cooperative binding of myosin II and cofilin to actin filaments (Ngo *et al*, 2016). Meanwhile, in this study, electron microscopy observations showed that Rng2CHD significantly changed the structure of actin filaments even when Rng2CHD was only sparsely bound to actin filaments (Fig. 5). The images of actin filaments suggest that Rng2CHD somehow reduces, or at least changes, the interaction between the two protofilaments

even when the binding ratio of Rng2CHD to actin protomers was only 1/25. Thus, it was actually

confirmed that Rng2CHD alters wide areas of the structure of actin filaments even when its binding

ratio to actin protomers is very low. At the saturating concentration of Rng2CHD (1 μM), actin

filaments became kinked, distorted or fragmented (Fig. 5D), which can also be explained by altered

interactions between neighboring actin protomers.

We consider two possible mechanisms how sparsely bound Rng2CHD inhibits

acto-myosin II movements. The first mechanism proposes that one or two actin protomers in direct

contact with the bound Rng2CHD molecule undergo structural changes, and those affected actin

protomers bind persistently to myosin II motors even in the presence of ATP, acting as a potent break.

The second mechanism assumes that a bound Rng2CHD molecule changes the structure of multiple

actin protomers, and the affected actin protomers become unable to interact with myosin II

productively. The first mechanism predicts that Rng2CHD should increase the amount of

co-sedimented S1 in the presence of ATP. However, this was not the case (Fig. 7). Furthermore, the

buckling of the moving actin filaments on HMM-coated surfaces, indicative of local inhibition of the

movement, was rarely observed in the presence of various concentrations of Rng2CHD (Fig. 2B). In

the presence of 5 nM Rng2CHD, movement speed was inhibited by 40%, and the binding ratio to

actin protomers was estimated as 1/185 based on $K_d$ (Eq. 3 in Materials and Methods). In other

words, on average, one molecule of Rng2CHD is bound per 500 nm of actin filament. A buckling of

this length would have been detectable with a spatial resolution of about 380 nm calculated from the numerical aperture and the fluorescence wavelength of this observation. Those two reasons led us to reject the first mechanism and conclude that force generation by myosin II is inhibited in broad sections of actin filaments not in direct contact with Rng2CHD. The velocity of acto-myosin II movement would decrease as the proportion of the inhibited section within the filaments increases in a Rng2CHD concentration-dependent manner. This idea is supported by the fact that the number of fluorescent spots of HMM-GFP was significantly reduced in a Rng2CHD concentration-dependent manner in the presence of a very low concentration of ATP (Fig. 8A).

Hereafter, we resolve the inhibition process by two aspects, and discuss their mechanisms next. The first is the mechanism by which sparsely bound Rng2CHD causes global structural changes in actin filaments. The second aspect is the mechanism by which structural changes of actin filaments inhibit actin motility driven by myosin II.

**The mechanism by which Rng2CHD causes global structural changes in actin filaments**

We now consider two hypotheses for the mechanism by which one molecule of Rng2CHD changes the structure of multiple actin protomers. The first is a cooperative structural change, in which one molecule of Rng2CHD bound to one actin protomer changes the structure of multiple neighboring actin protomers in the same filament. Such cooperative propagation of conformational

changes has been reported for many ABPs. For example, cofilin binds to actin filaments

cooperatively to form clusters along the filaments, and the half helical pitch of actin filaments in the

cluster is reduced by 25% (McGough *et al*, 1997). Notably, this structural change was propagated to

the neighboring cofilin-unbound bare region (Galkin *et al*, 2001, Ngo *et al*, 2015). Moreover, a

single molecule of gelsolin bound to the barbed end of an actin filament changes the structure of all

actin protomers in the filament (Orlova *et al*, 1995). Other ABPs such as myosin II (Oosawa *et al*,

1973), tropomyosin (Khaitlina *et al*, 2017), α-actinin (Craig-Schmidt *et al*, 1981) and formin (Papp

*et al*, 2006) also cause cooperative structural changes of actin filaments. The other hypothesis is the

memory effect, in which Rng2CHD molecules repeat transient binding to different actin protomers,

and the actin protomers remain in the altered conformation for a certain period of time after

dissociation of Rng2CHD. If there is a memory effect, Rng2CHD can alter the structure of entire

actin filaments even when its binding ratio to actin protomers is low. We speculate that either or both

of the two mechanisms, cooperative structural change and memory effect, underlie the global

structural changes in actin filaments induced by sparsely bound Rng2CHD.

**The mechanism by which structural changes of actin filaments inhibit acto-myosin II motility**

Rng2CHD inhibited steady-state binding of S1•ADP to actin filaments (Fig. 7B, 7C). In

other words, Rng2CHD reduced the affinity between S1•ADP and actin filaments. We also observed

transient binding of S1 in the presence of ATP using a fluorescent microscope and HS-AFM. Fluorescent microscopy demonstrated that Rng2CHD significantly decreased the region along actin filaments where HMM-GFP fluorescence was observed in the presence of 0.5 µM ATP (Fig. 8A). Moreover, HS-AFM directly showed that Rng2CHD significantly reduced the number of imaged S1 molecules transiently bound to actin filaments in the presence of 50 µM ATP and 1 mM ADP (Fig. 9).

First, we consider the results of high-speed AFM observation. Unlike when excess ADP coexists in the buffer, S1 molecules transiently bound to actin filaments were hardly observed in the presence of 500 µM ATP alone. Therefore, in the presence of 50 µM ATP and 1 mM ADP, HS-AFM presumably detected S1•ADP bound to actin filaments before the low concentration of ATP in the presence of excess ADP slowly disrupted the binding. The decrease in the number of detected S1 molecules caused by Rng2CHD can be interpreted in the following two ways: (1) the number of transient binding events of S1 decreased, or (2) the duration of each binding event was shortened. Hypothesis (1) predicts that actin-activated S1 ATPase is also very strongly inhibited by Rng2CHD, which was not the case. We thus concluded that the unstable binding of S1•ADP to actin filaments caused by Rng2CHD shortened the duration of transient binding of S1 to actin filaments, and decreased the efficiency of detection of transient binding by HS-AFM. Rng2CHD decreased the fluorescence of HMM-GFP along actin filaments, a result that is consistent with this idea (Fig. 8A).

Based on this conclusion, we propose two possible mechanisms of acto-myosin II movement inhibition caused by Rng2CHD, in the framework of the swinging lever arm model (Huxley, 1969; Cooke *et al*, 1984; Tokunaga *et al*, 1991; Uyeda *et al*, 1996) tightly coupled with the acto-myosin ATPase cycle (Lymn & Taylor, 1971) (Fig. 10). The first mechanism proposes that phosphate release from myosin II•ADP•Pi is promoted normally even by actin filaments structurally altered by Rng2CHD, but without the lever arm swing. Consequently, myosin II•ADP, which does not have the authentic post-power stroke structure, reduces its affinity to actin filaments. The second mechanism assumes that although the lever arm swing occurs, the myosin II motor domain slips at the contact surface with actin filaments, or myosin II•ADP dissociates from actin filaments because of the low affinity between myosin II•ADP and structurally altered actin filaments. This would lead to a failure of myosin II•ADP to maintain the tension, generated by the swing of the lever arm, to drive the movement of the actin filaments.

It is noteworthy that Rng2CHD did not inhibit actin filament movement on myosin V HMM, which is well established to move actin filaments processively with a lever arm swing (Forkey *et al*, 2003, Kodera *et al*, 2010). In line with this finding, two actin mutations, M47A (Kubota *et al*, 2009) and G146V (Noguchi *et al*, 2012), both of which likely perturb conformational changes of actin filaments, inhibit actin movements on myosin II, but not on myosin V. We suggest that the movement by myosin II, which is a fast and non-processive motor, requires some

unidentified structural property of actin filaments which is dispensable for the slow and processive

myosin V involving the lever arm swing. Further studies are needed to understand the mechanism by

which actin filaments structurally altered by Rng2CHD fails to move on myosin II.

Another interesting feature of the inhibition of acto-myosin II movement by Rng2CHD is

that there is hysteresis in the inhibition. The number of S1 molecules transiently bound to actin

filaments in the presence of ATP and ADP was significantly reduced when S1 was added after

Rng2CHD was allowed to interact with actin filaments (Fig. 9C). In contrast, when S1 was strongly

and densely bound to actin filaments in the nucleotide-free state and then Rng2CHD was added

together with ATP and ADP, Rng2CHD hardly changed the number of S1 molecules transiently

bound to actin filaments (Fig. 9C). These results showed apparent hysteresis in the impact of

Rng2CHD on S1 binding to actin filaments. We speculate that the dense rigor binding of S1 to actin

filaments or the subsequent active interaction of S1 with actin induced by the addition of ATP

changed the structure of actin filaments in one way, which hindered the subsequent inhibitory

interaction of Rng2CHD with the altered actin filaments. This explanation is apparently at odds with

the fact that Rng2CHD inhibited actomyosin movement in *in vitro* motility assays, in which myosin

II HMM was bound to actin filaments in the rigor state before the addition of ATP and Rng2CHD.

This discrepancy may be attributed to differences in experimental conditions, such as the density of

the bound myosin heads along the actin filaments before the addition of Rng2CHD and ATP, or

whether S1 molecules were present in the buffer containing ATP and Rng2CHD.

Phosphorylation of the myosin light chain (Fujime, 1983, Sellers *et al*, 1985; Griffith *et al*, 1987) and calcium regulation via tropomyosin and troponin (Ebashi & Kodama, 1965, Ebashi & Kodama, 1966) are two widely known major regulatory mechanisms of acto-myosin II movements. Additionally, it has been reported that caldesmon and calponin inhibit the movement of actin filaments on smooth muscle myosin II (Shirinsky *et al*, 1992). These ABPs are similar to Rng2CHD in that they potently inhibit acto-myosin II movements even with sparse binding to actin filaments (Shirinsky *et al*, 1992). Furthermore, calponin is homologous to Rng2CHD. However, these two ABPs are clearly different from Rng2CHD in their degree of cooperativity of motility inhibition (Shirinsky *et al*, 1992) and ATPase inhibition (Chalovich *et al*, 1987, Kolakowski *et al*, 1997). In addition, some filaments move even in the presence of concentrations of caldesmon and calponin that stop the movement of most actin filaments (Shirinsky *et al*, 1992), which is in sharp contrast to Rng2CHD that completely stops the movement of all actin filaments (Fig. 2A). It is unknown whether the inhibition of acto-myosin movement caused by these ABPs share the same mechanism with Rng2CHD, but Rng2CHD induces the highest degree of cooperative inhibition, making Rng2CHD suitable for elucidating the mechanism of acto-myosin II movement inhibition.

Further investigations are warranted to reveal the structural aspects of the actomyosin movement inhibition induced by Rng2CHD, and to clarify the relationship between the structural

changes of actin filaments and acto-myosin II motility.

# Materials and Methods

## Protein purification

Actin was purified from rabbit skeletal muscle acetone powder (Spudich & Watt, 1971; Pardee & Spudich, 1982). Myosin II HMM and myosin II S1 were prepared by digestion of rabbit skeletal muscle myosin with papain and α-chymotrypsin, respectively (Margossian & Lowey, 1982). HMM-GFP was expressed in *D. discoideum* and purified as described previously (Tokuraku *et al*, 2009). The HMM version of human myosin V with a FLAG-tag at the N-terminus and a c-myc tag at the C-terminus was coexpressed with calmodulin in insect cells and purified using a method described previously (Watanabe *et al*, 2006).

The gene encoding Rng2CHD (Takaine *et al*, 2009) was inserted at the *Hind*III and *Pst*I sites of the pCold-TEV vector (Ngo *et al*, 2015), which had a TEV protease recognition sequence between the 6×His sequence and the multiple cloning site of pColdI (Takara Bio). The amino acid sequence of His-TEV Rng2CHD was MNHKVHHHHHHIEGRHM<u>ENLYFQG</u>TLEGSEFKLDVNVGL…(Rng2CHD)…LPNFKA, where the underline shows the TEV recognition sequence.

Rng2CHD was expressed in BL21 *Escherichia coli* (Takara) according to the instructions provided by the manufacturer of pColdI. The cells were lysed by sonication in 2 mM 2-mercaptoethanol, 0.3% Triton X-100, 0.1 mM phenylmethylsulfonylfluoride, 400 mM NaCl, 10

mM imidazole (pH 7.4) and 20 mM Hepes (pH 7.4) on ice. The homogenate was clarified by centrifugation and mixed with Ni Sepharose 6 Fast Flow (GE Healthcare). After extensive washing, the peak fractions eluted by 7 mM 2-mercaptoethanol, 400 mM imidazole (pH 7.4) and 10 mM Hepes (pH 7.4) were combined and supplemented with His-tagged TEV protease at a 1/10 molar amount of proteins in the pooled fractions to separate the His-tag from Rng2CHD at the cleavage site for TEV protease. After dialysis against 50 mM KCl, 0.1 mM DTT and 10 mM Hepes (pH 7.4) overnight at 4°C, the protein solution was clarified by centrifugation and passed through Ni Sepharose 6 Fast Flow in a column to remove the released His-tag, His-TEV protease and uncleaved His-tagged Rng2CHD. This was followed by concentration with a centrifugal concentrator (Amicon Ultra-15 3 k device, Merck Millipore), and after supplemented with 10 mM DTT, aliquots were snap-frozen in liquid nitrogen and stored at -80°C.

To fuse GFP to the N-terminus of Rng2CHD, the GFP gene with a S65T mutation was inserted at the *Kpn*I and *BamH*I sites of pCold-TEV-Rng2CHD. A Gly-based 16 amino acid-residue linker sequence was inserted between the GFP gene and the Rng2CHD gene, so that the expressed GFP would not spatially inhibit the binding of Rng2CHD to the actin filament. The resultant amino acid sequence of GFP-Rng2CHD was MNHKVHHHHHHIEGRHM<u>ENLYFQG</u>TMSKGE…(GFP)…MDELY<u>GGSEFGSSGSSGSSKL</u>DV NVGL…(Rng2CHD)…LPNFKA, where the underline shows the TEV recognition sequence and the

double underline shows the linker sequence. GFP-Rng2CHD was expressed in Rosetta DE3 *E. coli*

(Merck Millipore), and purified basically in the same way as Rng2CHD, except that the protein was

further purified by anion exchange chromatography with an Econo-pac High Q cartridge

(BIO-RAD) before it was concentrated.

### *In vitro* motility assays

G-actin was polymerized in F-buffer (0.1 M KCl, 2 mM $MgCl_2$, 1 mM DTT, 1 mM ATP

and 10 mM Hepes pH 7.4) for 1 h at 22°C, and was diluted to 1 μM with NF buffer (25 mM KCl, 2

mM $MgCl_2$, 10 mM DTT and 20 mM Hepes pH 7.4). Diluted actin filaments were incubated with 1

μM rhodamine phalloidin (Invitrogen) or 1 μM phalloidin (Wako) for 1 h at 22°C.

*In vitro* actomyosin motility assays in which actin filaments move on myosin II were

performed according to the method of Kron and Spudich (Kron & Spudich, 1986). Rhodamine

phalloidin-stabilized actin filaments were bound to myosin II HMM on a nitrocellulose-coated glass

surface in a flow chamber filled with NF buffer containing 10 mg/ml BSA. Movement of actin

filaments was started by injecting 15 μl of MA buffer (25 mM KCl, 2 mM $MgCl_2$, 0.5 mM ATP, 10

mM DTT, 3 mg/ml glucose, 1.2 μM glucose oxidase, 0.15 μM catalase, 10 mg/ml BSA and 20 mM

Hepes pH 7.4) into the flow chamber. Various concentrations of Rng2CHD (5, 10, 50, 100 and 200

nM) were present in this buffer. Fluorescence of rhodamine phalloidin was imaged with an EMCCD

camera (iXon X3, Andor) on a fluorescent microscope (IX-71; Olympus) equipped with a Plan Apo

100X, 0.9 NA objective lens (Nikon) at a frame rate of 0.5 fps.

The acto-myosin II motility assay in the presence of GFP-Rng2CHD was performed

basically as above, with several modifications. Rhodamine phalloidin stabilized and unlabeled

phalloidin stabilized actin filaments were mixed at a 1:1 concentration in order to observe GFP

fluorescence without the interference of rhodamine fluorescence. Fluorescence micrographs were

taken with an EMCCD camera (iXon X3) on a TIRF microscope (IX-71) equipped with a UApo N

100X, 1.49 NA objective lens (Olympus). Laser light of 538 nm for exciting rhodamine and 488 nm

for exciting GFP were irradiated alternately at 2 s intervals, and the time-lapse imaging of moving

actin filaments and GFP-Rng2CHD dynamics was performed semi-simultaneously.

The images were processed with Image J (Schneider *et al*, 2012). Movements of actin

filaments were tracked by MTrackJ, a plugin for Image J (Meijering *et al*, 2012) and the velocities

were calculated. The fluorescence intensity of GFP was quantified by Image J as follows. Five

filaments near the center of the images were selected for each condition, and the fluorescence

intensity in five frames was measured. The light intensity at five points near the filament was

averaged and subtracted from the measured values along the filament as the background for each

filament. The values were normalized by the unit length of each filament.

For *in vitro* motility assays in which actin filaments move on myc-tagged myosin V

HMM, HMM molecules were immobilized on a glass surface via anti-c-myc antibody. MA2 buffer

(20 mM KCl, 4 mM MgCl$_2$, 2 mM ATP, 120 μM calmodulin, 1 mM EGTA, 1 mM DTT, 0.5%

methylcellulose (v/v), 1 mg/ml BSA and 25 mM imidazole pH 7.4) was used instead of MA buffer.

Mouse calmodulin was expressed in *E. coli* and purified as described previously (Shishido *et al*,

2009). The fluorescence images were captured with a camera (ORCA-Flash 2.8; Hamamatsu

Photonics) on a fluorescent microscope (IX-70; Olympus) equipped with a Plan-Fluor 100X, 1.3 NA

objective lens (Nikon) at a frame rate of 0.5 fps.

**Measurement of dissociation constant**

G-actin was polymerized in F-buffer for 1 h at 22°C and incubated with phalloidin at a

1:1 molar ratio for 1 h at 22°C. Phalloidin-stabilized actin filaments that were diluted to 3 μM and

various concentrations of Rng2CHD or GFP-Rng2CHD (1, 2, 3, 4, 5, 6 nM) were incubated together

in SA buffer (25 mM KCl, 2 mM MgCl$_2$, 0.5 mM ATP, 10 mM DTT and 20 mM Hepes pH 7.4) for 5

min at 22°C, and then centrifuged at 278,800 g for 10 min at 22°C. The supernatants and pellets

were subjected to SDS-PAGE. Images of Coomassie brilliant blue (CBB)-stained gels were read into

Image J and the concentration of Rng2CHD in each fraction was quantified by densitometry. The

dissociation constant ($K_d$) for Rng2CHD to actin filaments was calculated by fitting plots of

[Rng2CHD bound to actin filaments] versus [Rng2CHD free] with the following equation:

$$[Rng2CHD_{bound}] = [Actin_{total}] [Rng2CHD_{free}] / ([Rng2CHD_{free}+K_d]) \qquad [1]$$

$K_d$ between actin filaments and GFP-Rng2CHD was calculated in the same way.

**Estimation of binding ratio of Rng2CHD and GFP-Rng2CHD to actin filament from $K_d$**

$K_d$ between Rng2CHD and actin filaments is given by:

$$K_d = [Actin_{free}] [Rng2CHD_{free}] / [Rng2CHD_{bound}] \qquad [2]$$

The concentration of actin filaments is extremely low in solutions in which Rng2CHD in

the buffer interacts with actin filaments immobilized on the substrate, such as *in vitro* actomyosin

motility assays and observations of binding with high-speed AFM and fluorescence microscopy, so

that $[Rng2CHD_{free}]$ can be approximated by the concentration of total Rng2CHD

($[Rng2CHD_{total}]$). Therefore, the following approximation holds from Eq. 2:

$$[Rng2CHD_{bound}] / [Actin_{free}] \simeq [Rng2CHD_{total}] / K_d \qquad [3]$$

The binding ratios of Rng2CHD and GFP-Rng2CHD to the dilute actin protomers were

estimated with this approximation using the value of $K_d$.

**Electron microscopy**

G-actin was polymerized in F-buffer for 1 h at room temperature, and 1 μM actin

filaments were mixed with various concentrations of Rng2CHD in EM buffer (25 mM KCl, 4 mM

$MgCl_2$, 1 mM DTT, 0.1 mM ATP and 10 mM imidazole pH 7.4) at room temperature. A small

volume of each sample was placed on a carbon-coated grid at 30 s (40 nM Rng2CHD), 2 min (200

nM Rng2CHD) or 4 min (1 µM Rng2CHD) after mixing. The samples were negatively stained with

1% uranyl acetate, and observed in a transmission electron microscope (Tecnai F20; FEI). The

electron micrographs were recorded with a Gatan ORIUS 831 CCD camera, adjusted for contrast

and Gaussian-filtered using Adobe Photoshop.

**Myosin II S1 ATPase measurements**

Actin-activated S1 ATPase was measured according to the method of Kodama *et al*.

using malachite green (Kodama *et al*, 1986). G-actin was polymerized in F-buffer for 1 h at 22°C.

The solution was centrifuged at 278,800 g for 10 min at 22°C and actin filaments in the pellet were

resuspended in NF buffer. This procedure was repeated once more to minimize the amount of carried

over phosphate. Actin filaments diluted to 23.8 µM was mixed with various concentrations of

Rng2CHD (0, 0.33, 0.82, 1.91, 5 µM) in NF buffer containing 2 mM ATP, and was incubated for 10

min at 25°C. The reaction was started by the addition of 50 nM S1, and phosphate released at 0, 2, 4,

6 and 8 min was measured.

**Co-sedimentation assay of actin filaments and S1 with Rng2CHD**

For the co-sedimentation assays in the presence of ATP, G-actin was polymerized in

F-buffer for 1 h at 22°C and was then incubated with phalloidin at a 1:1 molar ratio for 1 h at 22°C.

For the co-sedimentation assays in the presence of ADP, the solution of actin filaments was centrifuged at 278,800 g for 10 min at 22°C, and the pelleted actin filaments were resuspended in NF buffer. This procedure was repeated once again to minimize the amount of carried over ATP before the addition of phalloidin. The following two samples were prepared for both experiments: (1) 3 μM actin filaments were incubated with 2 μM S1 for 5 min at 22°C; (2) 3 μM actin filaments were incubated with Rng2CHD for 10 min at 22°C, and then incubated with subsequently added S1 for 5 min at 22°C. Each sample was prepared in S-ATP buffer (25 mM KCl, 2 mM MgCl$_2$, 2 mM ATP, 10 mM DTT, 10 mM phosphoenolpyruvate, 10 units / ml pyruvate kinase and 20 mM Hepes pH 7.4) or S-ADP buffer (25 mM KCl, 2 mM MgCl$_2$, 2 mM ADP, 10 mM DTT and 20 mM Hepes pH 7.4). After incubation, each sample was centrifuged at 278,800 g for 10 min at 22°C. The supernatants and pellets were subjected to SDS-PAGE. The images of the CBB-stained gels were read into Image J and the concentration of S1 in each fraction was quantified by densitometry.

**Fluorescence microscope-based binding assay**

Binding of HMM-GFP to actin filaments was observed as follows. G-actin was polymerized in NF-buffer (50 mM KCl, 2 mM MgCl$_2$, 0.5 mM EGTA, 1 mM DTT, 20 mM PIPES, pH 6.5) for 2 h at 22°C. Actin filaments and Alexa 647 phalloidin (Invitrogen) were mixed at a molar ratio of 20:1 and incubated overnight on ice. The surface of coverslips was covered with a

positively charged lipid bilayer and was used to construct flow chambers as described previously (Ngo *et al*, 2015), except that the weight ratio of 1,2-dipalmitoyl-sn-glycero-3-phosphocholine (DPPC; Avanti Polar Lipids, Alabaster, AL) and 1,2-dipalmitoyl-3-trimethylammonium-propane (DPTAP; Avanti Polar Lipids) was 17:3 (Hirakawa *et al*., 2017). Alexa 647 phalloidin-stabilized actin filaments diluted in NF-ATP buffer (NF buffer containing 0.5 μM ATP) were introduced into the flow chamber to loosely bind on the positively charged lipid monolayer. HMM-GFP and Rng2CHD diluted in NF-ATP buffer were then introduced to the flow chamber. Alternatively, NF-ATP buffer in the above procedures was replaced with NF buffer for the assays in the nucleotide-free state. Fluorescence of Alexa 647 and GFP was imaged with a fluorescence microscope (ECLIPSE E600, Nikon, Tokyo, Japan) equipped with an ARUGUS-HiSCA system (Hamamatsu Photonics, Hamamatsu, Japan). Images were captured using a 100x objective lens (CFI Plan Apo Lambda 100x Oil, NA 1.45; Nikon).

**High-speed atomic force microscopy**

We used a laboratory-built high-speed atomic microscope (HS-AFM) as described previously (Ando *et al*, 2013). The cantilevers and the sample stage, positively charged lipid bilayers on a mica surface, were prepared according to the method of Ngo *et al*. (Ngo *et al*, 2015). The free oscillation amplitude of the cantilever was set at 2.0 nm, and the amplitude set point for feedback

control was set to 1.6 -1.9 nm during observations.

G-actin was polymerized in F-buffer for 1 h at 22°C, and then the buffer on the sample stage was replaced by 2 μl of 20 μM actin filaments in AFM buffer (25 mM KCl, 2 mM MgCl$_2$, 50 μM ATP, 1 mM ADP, 10 mM DTT, 20 mM Hepes pH 7.4). After incubation for 10 min at 22°C, the surface of the sample stage was rinsed with 20 μl of AFM buffer to remove free actin filaments. Subsequently, the surface of the sample stage was immersed in 60 μl of AFM buffer in the observation chamber of the HS-AFM. The observations of the transient binding of myosin II S1 to actin filaments in the presence of ATP and ADP were performed under the following three conditions: (1) S1 diluted in AFM buffer was added to the observation chamber to a final concentration of 500 nM; (2) 12.4 nM Rng2CHD was allowed to interact with actin filaments in the observation chamber for 15 min at 22°C, and then 20 μM S1 in AFM buffer was added to the observation chamber to a final concentration of 500 nM; (3) 500 nM S1 was allowed to interact with actin filaments in the observation chamber filled with NF buffer (25 mM KCl, 2 mM MgCl$_2$, 10 mM DTT, 20 mM Hepes pH 7.4) for 10 min at 22°C. Subsequently, 1/40 vol of a mixture of ATP, ADP and Rng2CHD in AFM buffer was added to the observation chamber to a final concentration of 50 μM, 1 mM and 12.4 nM, respectively.

AFM images were obtained at a scan speed of 0.5 s per field of view, and were then visualized by Kodec4, our laboratory-built software (Ngo *et al*, 2015). Images scanned between 1

min and 2 min after the addition of S1 or ATP were analyzed, and the number of transiently bound

S1 molecules on actin filaments was visually counted.

**Acknowledgments**

We thank Dr. Toshio Ando for technical support for HS-AFM experiments, and Dr. Atsuko

H. Iwane and Dr. Toshio Yanagida for the gift of the virus to express HMM of myosin V. This work

was supported in part by Bio-SPMs Collaborative Research of WPI Nano Life Science Institute,

Kanazawa University, and Grants-in-aid from the Ministry of Education, Culture, Sports, Science

and Technology to KT (No. 24370069 and 24117008), K Nakano (No. 22019004), MT (No.

24770177) and TU (No. 24370069 and 24117008).

# Figures

**Figure 1 Domain structure of Rng2.**

Domain structure of human IQGAP1 and fission yeast Rng2 and Rng2CHD. Numbers written above

the domains show the percentage of amino acid sequence homology (modified from Takaine *et al*,

2009).

**Figure 2 Rng2CHD strongly inhibits movement of actin filaments on myosin II, but not on myosin V.**

**(A)** Movement velocity of actin filaments on myosin II HMM (black plots) or myosin V HMM (red plots) attached to glass surfaces in the presence of various concentrations of Rng2CHD. The

concentration of ATP was 0.5 mM for myosin II HMM, and 2 mM for myosin V HMM. Ten

smoothly moving filaments were chosen for each condition, and approximately 10 consecutive

measurements were made for each filament. Data are expressed as the mean ± SD. **(B)** Trajectories

of moving actin filaments on myosin II HMM in the presence of 0, 5 and 50 nM Rng2CHD. Eleven

consecutive images with 0.2 s intervals are coded in rainbow colors from red to blue, and overlaid.

**Figure 3 Measurement of $K_d$ between actin filaments and Rng2CHD or GFP-Rng2CHD.**

**(A,B)** Co-sedimentation assay of actin filaments with Rng2CHD (A) or GFP-Rng2CHD (B). **(C,D)**

For each condition, the concentration of bound Rng2CHD (C) or GFP-Rng2CHD (D) was plotted

against the concentration of the free fraction and fitted with the following equation: $[Rng2CHD_{bound}]$

$= [Actin_{total}]\ [Rng2CHD_{free}]\ /\ ([Rng2CHD_{free}+K_d])$. $K_d$ between Rng2CHD and actin protomers was

calculated as an average value of two trials.

**Figure 4 Sparsely bound GFP-Rng2CHD on actin filaments inhibits movement of actin**

**filaments on myosin II.**

**(A)** Movement velocity of actin filaments on myosin II HMM-coated glass surfaces in the presence

of various concentration of GFP-Rng2CHD (black plots). Ten smoothly moving filaments were

chosen for each condition, and approximately 10 consecutive measurements were made for each

filament. Data are expressed as the mean ± SD. The movement velocity in the presence of

Rng2CHD is also shown for the reference (red plots). **(B)** Fluorescence micrographs of

GFP-Rng2CHD bound to actin filaments. Actin filaments stabilized by non-fluorescent phalloidin

and those labeled with rhodamine phalloidin were present at a molar ratio of 1: 1. Green: GFP

fluorescence. Red: rhodamine fluorescence. Non-fluorescent phalloidin-stabilized actin filaments

were visualized by bound GFP-Rng2CHD (green filaments). **(C)** Fluorescence intensity of

GFP-Rng2CHD on non-fluorescent phalloidin-stabilized actin filaments. The fluorescence intensity

along five filaments was measured for five frames. Data are expressed as the mean ± SD of 25

measurements. Bars: 10 μm. **(D)** Binding ratio of GFP-Rng2CHD to actin protomers calculated from

measured fluorescence intensity that caused indicated degrees of movement inhibition.

**Figure 5 Rng2CHD deforms the helical structures of actin filaments.**

Electron micrographs of negatively stained actin filaments in the presence of 0 nM (A), 40 nM (B),

200 nM (C) and 1 μM (D) Rng2CHD. The boxed regions in each image are magnified to the right.

In the presence of Rng2CHD, the filaments often appear to be slightly curved or bent. In some parts,

the two actin protofilaments look separated (yellow brackets). At a high concentration of Rnd2CHD,

the filaments are highly deformed and show discontinuities (D).

**Figure 6 Actin-activated S1 ATPase in the presence of Rng2CHD.**

The orange (0.33 μM Rng2CHD), green (0.82 μM Rng2CHD) and blue (1.91 μM Rng2CHD) plots

were measured in the presence of Rng2CHD concentrations that were expected to bind to actin

protomers at binding ratios which caused a 50%, 75% and 95% reduction of acto-myosin II

movement speed in *in vitro* motility assays, respectively. Data are expressed as the mean ± SD of

three independent experiments.

**Figure 7 Rng2CHD inhibits the steady-state binding of S1 to actin filaments in the presence of**

**ADP, but not in the presence of ATP.**

**(A,B)** Co-sedimentation assay of S1 and actin filaments in the presence of Rng2CHD and 2 mM ATP

(A) or 2 mM ADP (B). **(C)** Fraction of S1 dissociated from actin filaments was compared with and

without Rng2CHD. Rng2CHD made a significant difference in the presence of ADP (Student's t-test,

p<0.01), but not in the presence of ATP. Data are expressed as the mean ± SD of three independent

experiments.

**Figure 8 Rng2CHD decreases the fluorescence of HMM-GFP along actin filaments in the**

**presence of ATP.**

Fluorescence micrographs of actin filaments labeled lightly with Alexa647 phalloidin and

HMM-GFP in the presence of 0.5 μM ATP (A) and in the nucleotide-free state (B). In each panel, the

top row shows the fluorescence of Alexa647, and the bottom row shows that of GFP.

**Figure 9 Rng2CHD significantly decreases the number of S1 molecules bound along actin filaments in the presence of ATP.**

**(A, B)** HS-AFM images of actin filaments interacting with S1 in the presence of 50 µM ATP and 1 mM ADP. The images were scanned at about 2 min after the addition of 500 nM S1. S1 molecules bound on the top and along the side of the filaments are indicated by yellow and white arrowheads, respectively. In (A), 500 nM S1 was added to actin filaments. In (B), in contrast, actin filaments were incubated with 12.4 nM Rng2CHD for 15 min, then 500 nM S1 was added. Bars: 50 nm. **(C)** Number of S1 molecules bound on the top or along the sides of the filaments in the presence of 50 µM ATP and 1 mM ADP. The values were normalized by the total length of the measured filaments

and time. White bars: 500 nM S1 was added to actin filaments. Red bars: Actin filaments were

incubated with 12.4 nM Rng2CHD for 15 min, then 500 nM S1 was added. Blue bars: Actin

filaments were incubated with 500 nM S1 in the nucleotide-free state for 10 min, then 50 μM ATP, 1

mM ADP and 12.4 nM Rng2CHD were added. The number of bound S1 molecules was counted in

the images scanned between 1 and 2 min after the addition of S1 (white and red bars) or ATP (blue

bars).

**Figure 10 Lever arm swing in the myosin II motor domain that is tightly coupled with the**

**ATPase cycle.**

**Supplemental Figure 1 Co-sedimentation of Rng2CHD with skeletal muscle myosin II filaments.**

Skeletal muscle myosin II filaments, Rng2CHD and a mixture of the two proteins were centrifuged at 30,000 rpm for 10 min at 22°C in a Beckman TLA100.1 rotor, and the supernatant and pellet fractions were analyzed by SDS-PAGE. The concentrations of myosin II and Rng2CHD were 2 μM, and the buffer was 10 mM Hepes (pH 7.4), 25 mM KCl, 4 mM MgCl$_2$, 0.2 mM ATP and 1 mM DTT.

**Table 1 Estimated binding ratio of Rng2CHD and GFP-Rng2CHD to actin filaments.**

| Inhibition rate of movement | 50% | 75% | 80% | 90% | 95% | 100% |
|---|---|---|---|---|---|---|
| $[\text{Rng2CHD}_{bound}]/[Actin_{total}]$ | 1/75 | 1/30 | 1/28 | 1/21 | 1/13 | 1/6 |
| $[\text{GFP} - \text{Rng2CHD}_{bound}]/[Actin_{total}]$ | 1/23 | 1/15 | 1/14 | 1/8 | 1/5 | 1/4 |

The estimated binding ratio of Rng2CHD and GFP-Rng2CHD to actin protomers that caused various degrees of movement inhibition. These values were estimated based on the following approximation: $[Rng2CHD_{total}]/K_d \cong [Rng2CHD_{bound}]/[Actin_{free}]$.

# 5. References

Ando T, Uchihashi T & Kodera N (2013) High-Speed AFM and Applications to Biomolecular

    Systems. *Annu. Rev. Biophys.* **42:** 393–414

Chalovich JM, Cornelius P & Benson CE (1987) Caldesmon inhibits skeletal actomyosin

    subfragment-1 ATPase activity and the binding of myosin subfragment-1 to actin. *J. Biol.*

    *Chem.* **262:** 5711–5716

Cooke R, Crowder MS, Wendt CH, Barnett VA & Thomas DD (1984) Muscle Cross-Bridges: Do

    They Rotate? *Adv. Exp. Med. Biol.* **170:** 413–427

Craig-Schmidt MC, Robson RM, Goll DE & Stromer MH (1981) Effect of alpha-actinin on actin

    structure. *Biochim. Biophys. Acta - Protein Struct.* **670:** 9–16

Ebashi S & Kodama A (1965) A new protein factor promoting aggregation of tropomyosin. *J.*

    *Biochem.* **58:** 107–108

Ebashi S & Kodama A (1966) Interacton of Troponin with F-actin in the Presence of

    Tropomyosin. *J. Biochem.* **59:** 425–426

Eng K, Naqvi NI, Wong KCY & Balasubramanian MK (1998) Rng2p, a protein required for

    cytokinesis in fission yeast, is a component of the actomyosin ring and the spindle pole body.

    *Curr. Biol.* **8:** 611–621

Forkey JN, Quinlan ME, Shaw MA, Corrie JET & Goldman YE (2003) Three-dimensional

structural dynamics of myosin V by single-molecule fluorescence polarization. *Nature* **422:** 399–404

Fujime S (1983) Phosphorylation of Myosin Light Chain Modulates the In Vitro Movement of Fibers Composed of Actin and Myosin from Skeletal Muscle. *J. Biochem.* **94:** 1539–1545

Galkin VE, Orlova A, Lukoyanova N, Wriggersd W & Egelman EH (2001) Actin depolymerizing factor stabilizes an existing state of F-actin and can change the tilt of F-actin subunits. *J. Cell Biol.* **153:** 75–86

Griffith LM, Downs SM & Spudich JA (1987) Myosin light chain kinase and myosin light chain phosphatase from Dictyostelium: Effects of reversible phosphorylation on myosin structure and function. *J. Cell Biol.* **104:** 1309–1323

Hill AV (1938) The heat of shortening and the dynamic constants of muscle. *Proc. R. Soc. London. Ser. B - Biol. Sci.* **126:** 136–195

Hirakawa R, Nishikawa Y, Uyeda TQP & Tokuraku K (2017) Unidirectional growth of heavy meromyosin clusters along actin filaments revealed by real-time fluorescence microscopy. *Cytoskelet.* **74:** 482–489

Huxley H (1969) The Mechanism of Muscular Contraction. *Science* **164:** 1356–1366

Kamasaki T, Osumi M & Mabuchi I (2007) Three-dimensional arrangement of F-actin in the contractile ring of fission yeast. *J. Cell Biol.* **178:** 765–771

Khaitlina S, Tsaplina O & Hinssen H (2017) Cooperative effects of tropomyosin on the dynamics of the actin filament. *FEBS Lett.* **591:** 1884–1891

Kodama T, Fukui K & Kometani K (1986) The initial phosphate burst in ATP hydrolysis by myosin and subfragment-1 as studied by a modified malachite green method for determination of inorganic phosphate. *J. Biochem.* **99:** 1465–1472

Kodera N, Yamamoto D, Ishikawa R & Ando T (2010) Video imaging of walking myosin v by high-speed atomic force microscopy. *Nature* **468:** 72–76

Kolakowski J, Karkucinska A & Dabrowska R (1997) Calponin inhibits actin-activated MgATPase of myosin subfragment 1 (S1) without displacing S1 from its binding site on actin. *Eur. J. Biochem.* **243:** 624–629

Kron SJ & Spudich JA (1986) Fluorescent actin filaments move on myosin fixed to a glass surface. *Proc. Natl. Acad. Sci. U. S. A.* **83:** 6272–6276

Kubota H, Mikhailenko S V., Okabe H, Taguchi H & Ishiwata S (2009) D-loop of actin differently regulates the motor function of myosins II and V. *J. Biol. Chem.* **284:** 35251–35258

Lord M & Pollard TD (2004) UCS protein Rng3p activates actin filament gliding by fission yeast myosin-II. *J. Cell Biol.* **167:** 315–325

Lymn RW & Taylor EW (1971) Mechanism of Adenosine Triphosphate Hydrolysis by

Actomyosin. *Biochemistry* **10:** 4617–4624

Mabuchi I & Okuno M (1977) The effect of myosin antibody on the division of starfish

blastomeres. *J. Cell Biol.* **74:** 251–263

Margossian SS & Lowey S (1982) Preparation of myosin and its subfragments. *Meth.Enzymol.*

**85:** 55–71

McGough A, Pope B, Chiu W & Weeds A (1997) Cofilin changes the twist of F-actin:

Implications for actin filament dynamics and cellular function. *J. Cell Biol.* **138:** 771–781

Meijering E, Dzyubachyk O & Smal I (2012) Methods for Cell and Particle Tracking. *Methods*

*Enzymol.* **504:** 183–200

Ngo KX, Kodera N, Katayama E, Ando T & Uyeda TQP (2015) Cofilin-induced unidirectional

cooperative conformational changes in actin filaments revealed by high-speed atomic force

microscopy. *Elife* **4:** 1–19

Ngo KX, Umeki N, Kijima ST, Kodera N, Ueno H, Furutani-Umezu N, Nakajima J, Noguchi

TQP, Nagasaki A, Tokuraku K & Uyeda TQP (2016) Allosteric regulation by cooperative

conformational changes of actin filaments drives mutually exclusive binding with cofilin and

myosin. *Sci. Rep.* **6:** 1–11

Noguchi TQP, Komori T, Umeki N, Demizu N, Ito K, Iwane AH, Tokuraku K, Yanagida T &

Uyeda TQP (2012) G146V mutation at the hinge region of actin reveals a myosin

class-specific requirement of actin conformations for motility. *J. Biol. Chem.* **287:** 24339–24345

Oosawa F, Fujime S, Ishiwata S & Mihashi K (1973) Dynamic Property of F-Actin and Thin Filament. *Cold Spring Harb. Symp. Quant. Biol.* **37:** 277–285

Orlova A, Prochniewicz E & Egelman EH (1995) Structural Dynamics of F-Actin: II. Cooperativity in Structural Transitions. *J. Mol. Biol.* **245:** 598–607

Papp G, Bugyi B, Ujfalusi Z, Barkó S, Hild G, Somogyi B & Nyitrai M (2006) Conformational changes in actin filaments induced by formin binding to the barbed end. *Biophys. J.* **91:** 2564–2572

Pardee JD & Spudich JA (1982) Purification of muscle actin. *Methods Enzymol.* **85:** 164–181

Pelham RJ & Chang F (2002) Actin dynamics in the contractile ring during cytokinesis in fission yeast. *Nature* **419:** 82–86

Satterwhite LL & Pollard TD (1992) Cytokinesis. *Cell Biol.* **4:** 43–52

Schneider CA, Rasband WS & Eliceiri KW (2012) NIH Image to ImageJ: 25 years of Image Analysis HHS Public Access. *Nat. Methods* **9:** 671–675

Sellers JR, Spudich JA & Sheetz MP (1985) Light chain phosphorylation regulates the movement of smooth muscle myosin on actin filaments. *J. Cell Biol.* **101:** 1897–1902

Shirinsky VP, Biryukov KG, Hettasch JM & Sellers JR (1992) Inhibition of the relative movement

of actin and myosin by caldesmon and calponin. *J. Biol. Chem.* **267:** 15886–15892

Shishido H, Yamada MD, Kondo K & Maruta S (2009) Photocontrol of calmodulin interaction

with target peptides using azobenzene derivative. *J. Biochem.* **146:** 581–590

Spudich JA & Watt S (1971) The Regulation of Rabbit Skeletal Muscle Contraction. *J. Biol.*

*Chem* **246:** 4866–4871

Takaine M, Numata O & Nakano K (2009) Fission yeast IQGAP arranges actin filaments into

the cytokinetic contractile ring. *EMBO J.* **28:** 3117–3131

Tokunaga M, Sutoh K & Wakabayashi T (1991) Structure and Structural Change of the Myosin

Head. *Adv. Biophys.* **27:** 157–167

Tokuraku K, Kurogi R, Toya R & Uyeda TQP (2009) Novel Mode of Cooperative Binding

between Myosin and Mg2+-actin Filaments in the Presence of Low Concentrations of ATP.

*J. Mol. Biol.* **386:** 149–162

Uyeda TQP, Abramson PD & Spudich JA (1996) The neck region of the myosin motor domain

acts as a lever arm to generate movement. *Proc. Natl. Acad. Sci.* **93:** 4459–4464

Watanabe S, Ikebe R & Ikebe M (2006) Drosophila myosin VIIA is a high duty ratio motor with a

unique kinetic mechanism. *J. Biol. Chem.* **281:** 7151–7160

Zang JH, Cavet G, Sabry JH, Wagner P, Moores SL & Spudich JA (1997) On the role of

myosin-II in cytokinesis: division of *Dictyostelium* cells under adhesive and nonadhesive

conditions. *Mol. Biol. Cell* **8:** 2617–2629