1 **Genome diversification via genetic exchanges between temperate and**
2 **virulent bacteriophages**

4 Jorge A. Moura de Sousa[1,*], Eugen Pfeifer[1], Marie Touchon[1] & Eduardo P.C. Rocha[1,*]
5
6 [1]Microbial Evolutionary Genomics, Institut Pasteur, CNRS, UMR3525, Paris 75015, France

7 * corresponding authors: jorge-andre.sousa@pasteur.fr, erocha@pasteur.fr

10   ABSTRACT
11
12

Bacteriophages (henceforth phages) evolve by mutation and recombination. Temperate

phages are known to evolve rapidly by genetic exchanges. In contrast, recombination events

between virulent and between temperate and virulent phages have rarely been reported. A

gene flow barrier between these two large distinct groups of phages could affect the ability of

phages to acquire novel functions. Here, we show that genomes of temperate and virulent

phages are very distinct but often have a few almost identical genes. These cases are due to

recent genetic exchanges of both phage-like and bacterial-like genes. These exchanges were

probably mediated by phage recombinases with low homology requirements and mechanisms

of non-homologous end joining, since both were over-represented in recombinant phages and

their hosts. When assessing the impact of gene flow across the two populations of phages we

realized that temperate phages have narrower host ranges than virulent phages. This

suggests, and we find some evidence, that gene flow between temperate phages infecting

distant bacterial hosts might be mediated by recombination with the broader host virulent

phages. Hence, gene flow across phages with distinct lifestyles could drastically increase the

gene repertoire available for phage evolution, including the transfer of functional innovations

across taxa. These results also have implications for bacterial evolution because of the impact

of phage predation in bacterial population dynamics and the contribution of temperate phages

to the evolution of bacterial gene repertoires.

SIGNIFICANCE STATEMENT
33

Many   microbial   communities   are   intensely   predated   by   temperate   and   virulent

bacteriophages. The former also contribute to bacterial gene repertoires. Gene transfers

between bacteriophages favor their rapid adaptation, but are thought to occur rarely between

virulent and temperate bacteriophages because their genomes are very different in terms of

gene repertoires and genetic organization. We found that genetic exchanges occur frequently

between these types of bacteriophages, involve different phage and bacterial functions, and

are likely due to multiple DNA repair processes. Since we show that virulent bacteriophages

have broader host ranges, they can shuttle genes between temperate bacteriophages present

in taxa that are beyond the typical range of the latter. This enhances phages diversification

and has multiple impacts on bacterial evolution.

Introduction

Bacterial viruses (bacteriophages or phages) are ubiquitous across environments, where they are often the most abundant biological entities and shape bacterial population dynamics. Most known phages have double stranded DNA genomes and can be either virulent or temperate (1). Virulent phages are restricted to lytic infections, where rapid viral replication ends in progeny release and bacterial death. Temperate phages can follow either a lytic or a lysogenic cycle. In the latter, viral DNA maintains itself in the host as a prophage. Most prophage genes are silent until a signal activates their lytic cycle. Yet, some genes are expressed and they may provide adaptive phenotypes to their hosts (2, 3). Half of the bacterial genomes have at least one prophage and some have up to 20 prophages (4), showing the important contribution of temperate phages to bacterial gene repertoires. The relative frequency of temperate and virulent phages vary across bacterial taxa and environments. In the mammalian gut, for instance, temperate phages seem to be more prevalent (5), while most intracellular and many environmental bacteria lack prophages (6), and are thus presumably rarely infected by temperate phages. Phages also drive horizontal gene transfer among bacteria by transduction (3), which may disseminate virulence factors (7) and antibiotic resistance (8).

Phage genomes vary considerable in size, from a few dozen to hundreds of genes (9). The largest genomes tend to be found in virulent phages and seem to have gene repertoire dynamics like those of bacteria, where a core genome is complemented by very diverse sets of accessory genes (10, 11). There is very little information on the mechanisms of gene accretion in virulent phages, even if some families of virulent phages have remarkable diversity in their genomes' size, suggesting the existence of mechanisms of gene exchange (12). The genomic plasticity of temperate lambdoid phages has been much more extensively studied (9, 13, 14). Their analyses reveal that pairs of phages tend to have patches of regions of very similar sequences within genomes that are often very dissimilar. This genome mosaicism is facilitated by the modular organization of phage genomes, where groups of

functionally related genes are encoded, and potentially exchanged, together (15). It is also

facilitated by the key role of recombination in the production of phage genome concatemers

to be packaged into the virion (16). The molecular mechanisms of recombination underlying

the mosaicism are not completely understood. They may involve phage-encoded

recombinases, which are more permissive to differences between sequences than bacterial

RecA-mediated recombination (17). They may also involve homology-free mechanisms such

as non-homologous end joining or some type of illegitimate recombination (18, 19). In any

case, past studies revealed pairs of (typically temperate) phages with a few almost identical

genes within extremely divergent genomes.

The evolutionary dynamics of phage genomes reflect their distinct lifestyles. A recent study

showed that temperate and virulent phages have different "evolutionary modes" representing

the relative importance of genetic exchanges in their evolution (20). Indeed, there are many

reports of extensive mosaicism in temperate phages. These exchanges can occur between

phages during co-infections, between prophages in the genome, or between prophages and

infecting temperate phages (14, 17). Given the aforementioned prevalence of prophages in

bacteria, and that prophages tend to reside in bacterial genomes for many generations, this

may explain the very high rates of recombination of temperate phages. Virulent phages tend

to have more conserved core genomes and only a few highly variable regions (21, 22). They

might not exchange DNA outside restricted phage taxonomic groups (23, 24). Recombination

between virulent phages requires co-infection by two phages, which is probably rare.

Nevertheless, some cases have been studied in detail. Evolution of virulent phages of

Lactococcus lactis from a dairy production line showed that recombination brings more

polymorphism than point mutations to their genomes (25), and studies on T4 and T7 phages

revealed events of recombination and gene accretion (26, 27). Much less is known regarding

genetic exchanges between temperate and virulent phages, even if they are assumed to be

rare. For instance, a recent study found no evidence of recent exchanges between a set of 84

virulent and temperate phages of *Eschericha coli* (17), and a broader analysis suggested that

temperate phages rarely have genes in common with virulent ones (20). However, there are

some reports of virulent phages acquiring genes from temperate phages or their prophages

(28, 29). To the best of our knowledge, no study has systematically traced genetic exchanges

between temperate and virulent phages, even if their existence could have important

consequences. From a fundamental point of view, gene exchanges between the two types of

phages vastly increase the gene repertoire available for the evolution of each type of phage.

This could be particularly important for temperate phages infecting bacterial clades where they

are rare and for virulent phages if they are segregated in distinct families with little gene flux

between them (as previously proposed (23, 24)). Furthermore, if virulent and temperate

phages have different host ranges, gene exchanges can expand the taxonomic range of gene

flux in the most restricted types of phages (via recombination with phages with broader host

ranges). These exchanges could also affect the adaptation of bacteria, because phage

predation shapes microbial communities and temperate phages contributes to bacterial gene

repertoires. These exchanges could also have consequences for the safety of phage therapy.

In this work, we searched for the existence, prevalence and mechanisms underlying genetic

exchanges between temperate and virulent phages. Genetic exchanges between cellular

organisms are usually detected from variations in the density of polymorphism in genomes or

the phylogenetic congruence of core genes (30). This is impossible when analyzing divergent

phages because they have no core genes that could be used to build reliable phylogenies and

many genes lack homologs in other phages. Furthermore, the observed extensive mosaicism

of temperate phages implies that most genes have different phylogenies. Therefore, we

searched instead for strong mosaicism, i.e. highly similar genes within highly dissimilar

genomes. We were able to identify recent events of recombination between virulent and

temperate phages, although our conservative approach will miss very ancient recombination

events, or those between similar genomes. These findings impact our understanding of how

virulent and temperate phages evolve, especially because our results suggest that virulent

phages can facilitate the transfer of adaptive traits across more distant taxa.

Results

The network of similarities between temperate and virulent phages

We analysed a dataset of 2487 complete bacteriophage (phage) sequences available from NCBI (as of May of 2019). We used PHACTS (31) to predict their lifestyle and identified 1166 virulent and 1336 temperate phages. This method classifies some phages with high confidence and others with less confidence (see Methods). The analysis presented in the main text of this article focuses on this complete dataset, but restricting the study to the ca. 54% of phages classed by PHACTS with high confidence provides similar qualitative results (Supplementary Data). We searched for protein similarity across all phages in the dataset with at least 15 predicted proteins (2387 phages, ~96% of the dataset), identified reciprocal best hits between pairs of phages and computed the corresponding fraction of homologs weighted by sequence identity (weighted Genome Repertoire Relatedness, wGRR, see Methods). This resulted in 2,847,691 pairwise wGRR values, among which ca. 91% were null because there was no detectable homology using our procedure. The remaining 9% of non-null pairwise wGRR values involve almost all phages included in the analysis (2386 phages, >99%). The histograms of non-null wGRR values (Fig 1A) revealed higher values for the comparisons between pairs of phages with similar lifestyles than for the comparisons involving virulent and temperate phages. Very few (6%) of the comparisons between phages with different lifestyles show some level of homology, and even these tend to have lower wGRR than the ones observed for pairs of phages of similar lifestyles (i.e., temperate-temperate or virulent-virulent). The analysis of the phages classed with high confidence in terms of lifestyle showed an even greater difference in the distributions between pairs of phages of similar or opposite lifestyle (Fig S1A). This may indicate that the few high wGRR values between temperate and virulent phages correspond to erroneous assignments of lifestyles, or that confident lifestyle assignment becomes difficult when there are homologous genes across temperate and virulent phages. In any case, this analysis suggests the existence of relations of homology

between many temperate and virulent phages, even if these tend to result in small wGRR

values.

Network based representations are frequently used to describe evolutionary relationships

between phages (1, 24). We used the gene repertoire relatedness to build a graph where

phages are nodes and edges represent wGRR. This results in a network that is highly

connected, given that 99% of the phages have some level of homology with at least one other

phage (Fig S2). We used the Louvain method for community detection to cluster phages by

their wGRR, which resulted in 34 different clusters with at least 3 phages, 5 with 2 phages,

and 3 singletons (Fig S3). In order to compare the effects of the phage lifestyle, family (the

characterized virion morphology, see Methods) and host phyla in the separation of phages in

different clusters of wGRR, we used Shannon information entropy to quantify how

homogeneous each cluster is, regarding each of these three traits (Fig 1B, Fig S1B). As a

control, we randomly re-assigned all the labels (host phyla, family and lifestyle) to the phage

set. Although randomized clusters are more heterogeneous than any of the three traits being

tested, there was a clear and significant difference in terms of cluster homogeneity between

the lifestyle and the other two traits. Clusters are very homogeneous in terms of host phyla

and phage family, as revealed by average entropies close to zero. In contrast, they are much

less homogeneous in terms of lifestyle, where almost all clusters included both temperate and

virulent phages. These results show that there is genetic similarity between many temperate

and virulent phages independently of phage or bacterial taxonomic considerations.

Fig 1. Results of the analysis of similarity between phages. A) Histograms of the wGRR values (with wGRR>0). B) Shannon's

Entropy values for each cluster identified with the Louvain community detection method in the wGRR matrix. Results are given

for the three phage traits (N=34 for each trait, one per cluster). Boxplots represent the distribution of the concatenation of 100

repetitions of randomized cluster label re-assignments (N=3400 for each trait). All distributions are significantly less

heterogeneous than their random counterparts (all p<0.0001, 2-sample Kolmogorov Smirnov test). The observed clusters are

significantly more heterogeneous in phage lifestyle than in the other variables (both p<0.001, Tukey Honest Significant Difference

(HSD) Test).

Recombination drives genetic relatedness between temperate and virulent phage

The observed similarities between temperate and virulent phages could be explained by

common ancestry as well as by recent gene flux. When the wGRR is small, the two processes

translate into different patterns: common ancestry results in many homologs of low similarity

along the genomes, whereas recent genetic exchanges result in a small number of highly

similar homologs within very dissimilar genomes. To distinguish between these two scenarios,

one can use the contrast between the fraction of homologous genes (above a low minimal

similarity threshold of 35% identity) and the fraction of genes that are very similar (more than

80% identity). Distant common ancestry results in a sizeable fraction of the pairs of genomes

with homologs but no single very similar gene, whereas recent gene flow between distantly

related phages results in a few highly similar homologs and a low wGRR. This procedure does

not allow to identify very ancient events of recombination that result in a few homologs of low

sequence similarity. It also misses events of recombination between very closely related

genomes or when a large fraction of the genome is exchanged. Since we showed above that

the most confident virulent and temperate genomes have low wGRR values, these two cases

should be rare regarding comparisons between temperate and virulent phages (Fig 1A).

Genomes that engage rarely in recombination, as is thought to be the case between virulent

and temperate phages, should show a rapid decrease of the fraction of very similar genes in

function of the fraction of detectable homologs. Indeed, comparisons between temperate and

virulent (Fig 2A) and between virulent phages (Fig 2B) show a few nearly identical genes

(>80% identity) within genomes otherwise lacking homologues. These cases are best

explained by recent genetic exchanges. In contrast, when genomes exchange genes very

frequently, as is the case of many temperate phages, the comparisons show mosaics of highly

similar and highly dissimilar (eventually non-homologous) genes. This results in a linear

distribution where almost identical genes co-exist in very different genomes (Fig 2C).

The comparisons between temperate and virulent phages suggest a frequency of

recombination that is lower between the two populations of phages than between temperate

phages. This is in agreement with the network-based analyses above, where relations of

homology between temperate and virulent phages were frequent but limited to a small number

of genes (low wGRR). We then built a linear model based on the relationship between the

fraction of very similar proteins (>80% identity) and the fraction of homologous proteins (see

Methods) between temperate and virulent genomes (Fig 2A, blue line). We limited this dataset

to pairs of phages with at least 50% of homologous proteins, where the influence of outliers

that are associated with recombination is expected to be weaker. The linear model fitted to

this dataset represents a conservative model of the null hypothesis that the relationship

between these two parameters is mostly due to ancestry. It fits well the major group of

comparisons between virulent and temperate phages across almost all the range of the

regression (Fig 2A). We computed the negative residuals of the linear model to identify

significant negative deviations to the main trend. These represent cases where genomes have

an unexpectedly high number of very similar genes given the overall level of homology. This

threshold on the value of negative residuals was used across all datasets to identify putative

recent events of recombination (Fig 2, see Methods). For the case of exchanges between

temperate and virulent or between virulent phages, the overwhelming majority of exchanges

were found in very dissimilar phages (wGRR below 0.2, Fig 3A and Fig 3B). However, the

comparisons implicating pairs of temperate phages have a notably different distribution of

genetic similarity (Fig 3B), with a large fraction showing intermediate (>0.25) wGRR values.

This is again suggestive of a higher gene flux in temperate phages, in agreement with other

works showing their high mosaicism (1, 9).

The analysis of sequence similarity between phages connected in the wGRR network shows

two distinct patterns: genomes that are more likely to be ancestrally related tend to have large

regions of low sequence similarity, whereas the others show homology limited to one or a few

very similar genes (Fig 2, Fig S4). This suggests that our method discriminates recent

recombination from shared ancestry (see examples of pairs of phages more likely to be

ancestrally related in Fig 2 and Fig S4). 48% (528) of the virulent phages and 66% (852) of

temperate phages in our dataset were classified as recombinant. Considering only

recombination between phages of different lifestyle, 36% of the virulent phages were inferred

to recombine with temperate phages, whilst 37% of temperate phages were found to

recombine with their virulent counterparts.

The network restricted to pairs of recombinant phages (Fig 3C) shows once again clusters

that are less homogeneous in terms of lifestyle than in terms of host phyla or phage family

(Fig S5A). This means that there is more recombination inferred between phages of different

lifestyle than between phages of different families or between those that infect different

bacterial clades. Furthermore, it suggests that the coincidence of virulent and temperate

phages in the same clusters of the graph is partly driven by recombination. We found

qualitatively similar conclusions when restricting the analysis to phages with confidently

assigned lifestyles, albeit in this case there was a non-significant difference in heterogeneity

between phage lifestyle and the phage family (Fig S5B). Overall, these results show that

recombination between temperate and virulent phages is frequent in our dataset.

Fig 2. Identification of putative recombinant phage pairs. A-C) Scatterplots of pairs of phage genomes with recombinant proteins indicated as colored points (otherwise grey). The linear regression model (blue lines) was inferred for the temperate-virulent dataset in A and applied to all datasets. The genomic maps in front of each panel show representative examples of recombinant (top) or non-recombinant (ancestrally related) phage pairs (bottom). Color codes for points and genes: virulent (red), temperate (green), virulent-temperate (blue). Colors in the blocks linking the phages indicate sequence similarity between homologs.

Fig 3. Identification of putative recombinant phage pairs in the wGRR network. A) Scatterplot of pairs of temperate-virulent phages in terms of wGRR and the fraction of high sequence-identity homologous genes. B) Histogram of wGRR values for the subset of recombinant phage pairs. C) Network of recombinant phages. Each node represents a phage genome and each edge a relationship of homology (with wGRR below 0.5).

Functional classification of genes recombined between temperate and virulent phage

We investigated the genes exchanged between phages with different lifestyles to understand if recombination affects specific functions. These genes were the ones encoding highly similar proteins (at least 80% identity) present in dissimilar phage genomes (wGRR lower than 0.5). We found an average of 6 recombinant genes per phage genome. These genes were only slightly smaller than the other genes (191 vs 207 amino acids, Fig S6). To inquire on their role, we clustered the Prokaryotic Virus Orthologous Groups (pVOGs) profiles database into functional classes and used it to annotate phage-related functions (see Methods). Some functions are over (e.g., proteins involved in packaging, injection and assembly) or under (e.g., structural proteins) represented in the proteins exchanged between temperate and virulent phages. Yet, these genes matched a very large number of different protein profiles (>100), showing that genes identified as recombinant are not restricted to a few functional categories of slow evolving proteins that might spuriously be identified as recombinant. Hence, recombination between phages with different lifestyles can result in the exchange of a large range of functions (Table 1, Table S1, File S2).

Temperate phages sometimes carry genes that were acquired from bacterial genomes. We thus searched if recombinant genes were associated with some specific bacterial functions. This would suggest that recombination between temperate and virulent phages disseminates bacterial traits. We removed from the bacteria Non-supervised Orthologous Groups (bactNOG) profiles database the functions associated with phages by removing profiles matching pVOG profiles with a VQ higher than 80%. We then used the remaining bactNOG profiles to annotate the recombinant genes (see Methods). The vast majority of these genes

have no defined function. Some of the remaining might be typical phages genes (like putative structural components) that were missed by the filter using the pVOG profiles. Yet, other genes matched profiles whose functions are related to carbohydrate or nucleotide transport and metabolism, DNA related functions, or energy production and conservation (Table 2, Table S2, File S2). The category of energy production corresponds to the majority of genes of known function. They are found in phages infecting marine bacteria and correspond to proteins implicated in photosynthetic activity. Genes with these types of functions were previously found to be beneficial in virulent phages that infect cyanobacteria (32). Other notable functions include methylases, transporters and single-strand binding proteins. Hence, our results suggest that a multiplicity of phage and bacterial functions can be exchanged between temperate and virulent phages by recombination.

Table 1. Bacteriophage-related functions of recombinant proteins resulting from genetic exchanges between temperate and virulent phages. pVOG profiles matching the phage protein dataset (first column). Number of recombinant proteins (between temperate and virulent phages, second column). Enrichment (+) or depletion (-) of the functional category in recombinant proteins relative to all phage proteins (Fisher-exact test, significant differences for each functional category shown as asterisks (*) and adjusted for multiple comparisons using the Benjamin-Hochberg method).

pVOG Category	Matched profiles	Matched recombinant proteins in temperate-virulent phages	P-value (sign)
Unknown	201	821	**** (+)
Structure	26	96	**** (-)
DNA metabolism, Regulation & Recombination	25	73	**** (-)
Packaging, Injection & Assembly	21	115	* (+)
Others	3	5	**** (-)
Lysis	5	10	* (-)
Structure & Lysis	6	21	*** (+)
Other combinations	5	14	n.s.

Table 2. Bacterial-related functions of recombinant proteins resulting from genetic exchanges between temperate and virulent phages. Profiles matching the phage protein dataset and number of phage recombinant proteins matched by them.

bactNOG Category	Matched profiles	Matched recombinant proteins in temperate-virulent phages
Function unknown	87	212
Replication, recombination and repair	7	27
Transcription	6	19
Energy production and conversion	3	68
Carbohydrate transport and metabolism	1	2
Inorganic ion transport and metabolism	1	6
Cell wall/membrane/envelope biogenesis	1	1
Nucleotide transport and metabolism	1	11
Transcription + Signal transduction mechanisms	1	1

Recombinant phages are enriched for recombinases

Recombination between distantly related phages cannot usually be achieved by the cellular recombinases, because these tolerate few mismatches (33). However, some phage recombinases, e.g. lambda Red, allow recombination between more dissimilar sequences (17). To test if these recombinases could facilitate recombination between temperate and virulent phages, we searched for the Erf, Sak, Sak4, RecT, RecA and Gp2.5 families in the

genomes of phages (see Methods). We found that ca. 42% of the phages with recombinant

genes encode at least one of these recombinases, while only 26% of the remaining phages

encode them (Tables S3-4). This suggests that these recombinases may play an important

role in the recombination between distant phages.

There is no known mechanism that allows for genetic exchanges between virions. Instead,

the DNA of different phages can only recombine within bacterial hosts. It is therefore possible

that phage-like recombinases encoded in the genomes of bacteria play a role in the genetic

exchanges between phages. We searched for these recombinases in the bacterial hosts (see

Methods). For this, we removed RecA recombinases from the analysis, since they are

encoded by the vast majority of bacterial genomes (34). We found that hosts of phages with

recombinant genes are 29% more likely to encode recombinases in their genomes than the

hosts of the remaining phages. These host recombinases are concentrated (ca. 90% of the

total) in prophages and absent from the rest of the bacterial genomes (Tables S5-6). This

suggests that recombinases from prophages facilitate recombination between other phages

within bacterial cells. In agreement with this view, the genomes of bacterial species that host

phages with recombinant proteins tend to have more prophages (Fig S7-8).

Since it has been proposed that recombination between phages could also result from non-

homologous recombination (18), we searched the genomes for the gene encoding the Ku

protein. Ku is a key marker of the non-homologous end-joining (NHEJ) pathway (35). This

analysis showed that genomes of bacterial hosts of phages with recombinant proteins are

significantly more likely to encode NHEJ than the others (Fig S9, see Methods). In the Mu

phage, a Ku homolog has been shown to promote NHEJ using a host ligase (36). Interestingly,

we found that 10 phages in our dataset (6 temperate and 4 virulent) also code for the Ku

protein. All of these phages have recombinant proteins. Even if their number is small, the over-

representation of phages carrying Ku within those with recombinant genes is significant

(p=0.003, Fisher exact test). Either encoded in phages or bacteria, such a mechanism could

facilitate genetic exchanges between phages lacking homologs.

Broader host range of virulent phages facilitates gene flux across
clades of temperate phages
360

Defining the host range of phages is a notoriously difficult problem. However, if recombinant

genes are observed between phages that have distinct bacterial hosts, it is reasonable to

assume that either their host ranges overlap or that their ancestors must have at some point

infected the same bacteria. Because the recombined genes are very similar, this

recombination event must not have been too ancient (although translation of sequence

divergence into coalescence time is impossible with this data). Consequently, recombination

events can contribute to the characterisation of the phages' host range. To assess this

hypothesis, we computed both the patristic distance (based on the 16S rDNA, Fig 4A) and the

differences in 3-mer genomic signatures between the bacterial hosts (identified at the species

level, see Methods) of all pairs of phages with recombinant genes (Fig 4B, Fig S10-11). In all

datasets, most cases of recombination occurred within very closely related bacteria. However,

all three datasets (recombination between temperate, between virulent and between

temperate and virulent) significantly differ in their host range (as measured by this approach).

Importantly, pairs of virulent phages with recombinant genes are more frequently associated

with distant (or dissimilar) hosts, compared to those of temperate phages. This suggests that

virulent phages are able to infect (and undergo recombination in) a wider range of hosts

species than temperate phages. Interestingly, recombination between temperate and virulent

phages is associated with intermediate distances between bacterial hosts, which likely reflects

the contribution of broader host (virulent) and narrower host (temperate) phages.

381

382

383 **Fig 4. Recombination between phages indicates a broader host range of virulent phages. A)** Boxplots of the non-null

384 patristic distances (computed from the 16S rDNA gene tree) between the hosts' species of each recombinant phage pair. **B)**

385 Distributions of the differences in 3-mer genomic signatures between the hosts's species of each phage pair with recombinant

386 genes. *** p=0.001, Tukey HSD for all pairs. T-T: temperate-temperate, T-V: temperate-virulent, V-V: virulent-virulent.

387

388

389 If virulent phages have broader host ranges and recombine with temperate phages in

390 phylogenetically distant bacteria, they can shuttle genes between groups of temperate phages

391 that are sexually isolated (because they infect distantly related bacterial clades). To

392 systematize these observations, we built a simplified network with only recombinant

393 temperate-virulent phage pairs where each phage infects a different host genus (Fig 5A). This

394 network reveals several exchanges between virulent and temperate phages from distinct

395 genuses. In some cases, these exchanges concern distinct genes (Fig 5B), whereas in others

396 they concern the exact same gene (Fig 5C). In these examples, temperate phages share little

397 or no homology beyond genes homologous to the virulent phage (Fig S12), suggesting that

398   these exchanges were mediated by the virulent phage. These results suggest that

399   recombination between temperate and virulent phages, in tandem with the broader host range

400   of virulent phages, increases the frequency of genomic exchanges between temperate phages

401   in phylogenetically distant hosts.

Fig 5. Genetic exchanges between virulent and temperate phages with different host genus. A) Simplified network restricted to pairs of temperate-virulent phages with distinct bacterial host genuses (i.e., edges shown are only those that link phages with different hosts' genuses). Each node color represents a different bacterial host genus, and the node sizes identify either temperate (large nodes) or virulent (small nodes) phages. Examples from panels B and C are highlighted in ellipses. B) Examples of virulent phages (red) having recombined with different recombinant proteins exchanged with temperate phages infecting (green) of distinct bacterial genuses (green). C) Analogous to B, but representing genes that were exchanged with the virulent phage sharing the exact same protein with both temperate phages infecting distinct bacterial genuses. Colors in the blocks linking the phages indicate sequence similarity between homologs. The genomic maps between each of the two temperate phages for each of the cases are shown in Fig S12.

Discussion

Our study provides a first systematic quantification of genetic exchanges between temperate and virulent phages, highlighting their potential to drive phage diversification. We had to develop a specific method to identify these exchanges, since the lack of species phylogenies and core genes in phages precludes the use of standard approaches. A recent characterization of the evolution of phage genomes suggested that there are very few similarities between temperate and virulent phages (20). Interestingly, the phages in these few cases showed a pattern of rapid gene acquisition and loss. Moreover, this study found that these genomes have very few homologous genes which is consistent with the exchange of a small number of genes between temperate and virulent phages. Our method focuses on these types of cases, by analysing the mosaicism in individual (pairwise) genomes and by contrasting low global genome similarity, based on wGRR, with high individual protein similarity. However, our approach does have some caveats. First, incorrect classification of lifestyle (virulent or temperate) leads to spurious inference of inter-lifestyle recombination events. We controlled for this uncertainty by redoing the key analyses using the subset of phages with a lifestyle assigned with high confidence, which revealed similar qualitative patterns, often with a stronger statistical signal (see Supplementary Data). Second, the method is incapable of identifying certain types of recombination events: the ancient ones (proteins that are now less than 80% identical), those covering a significant fraction of the phage genome (raising the value of wGRR due to a high global level of similarity, as in (37)), and those between closely related phages (where wGRR is high). Even if this last limitation is unlikely to be important, since virulent and temperate phages are typically quite distinct, this means that we have probably underestimated the number of gene exchanges between phages with different lifestyles. Third, the distinction between recombination and ancestry is particularly challenging in the analysis of temperate phages because of their pervasive mosaicism. This leads to less confident identification of specific recombination events between

temperate phages. Nevertheless, and even with these caveats, a large fraction of the pairs of

phages with recombinant genes show clear traces of recent genetic exchanges.

When does recombination between temperate and virulent phages occur? We envisage three

major opportunities for such events. First, temperate and virulent phages may recombine

when co-infecting a bacterial host or when one of them is already present in the cell in a state

of pseudo-lysogeny. We do not know the frequency of such co-infections, but the existence of

recombination between virulent phages suggests that co-infections occur in nature (except if

all these occur indirectly through temperate phages). Second, virulent phages may recombine

with prophages present in the infected bacteria. Given the prevalence of bacteria with active

or defective prophages, such events are probably much more likely than those of co-infection.

Accordingly, the few previous descriptions of recombination between virulent phages and

temperate phages involved recombination with prophages (28, 38, 39). Virulent phages could

acquire genes from prophages by restricting chromosome DNA, as found in T2 and T4 phages

decades ago (40, 41). Such exchanges could also occur if prophages are induced by the

infection of virulent phage. Finally, prophages could acquire genes from virulent phages when

the genomes of the latter are cleaved by the hosts' defence systems, such as restriction

modification systems or CRISPR-Cas systems, which produce recombinogenic linear double

stranded DNA (42). A quantification of the relevance of each of these situations requires more

genome sequences, and experimental data, than the one available at this stage. Indeed, our

approach does not allow to identify the direction of the gene exchanges (from temperate to

virulent or vice versa), and this is key to know the relevance of the above scenarios.

Which mechanisms mediate gene flux between temperate and virulent phages? Given the

differences between these types of phages, genetic exchanges between them are likely to

occur through mechanisms with less stringent homology requirements than bacterial RecA

mediated homologous recombination. Two schools of thought have proposed that

recombination between very divergent temperate lambdoid phages could result from either

relaxed homologous recombination or illegitimate recombination (9, 17). These mechanisms

are not mutually exclusive, and both were proposed to act on the observed recombination

between virulent phages and prophages in *Lactococcus* (43). We observe that phages with

recombinant genes tend to encode recombinases known to tolerate more mismatches than

bacterial RecA-mediated recombination. Similarly, many bacterial genomes encode such

recombinases in their prophages. These recombinases might facilitate homologous

recombination between divergent phages. It is possible that alternative mechanisms not

involving homologous recombination also favour gene flux across temperate and virulent

phages. Accordingly, we observed that phages with recombinant genes are more likely to be

hosted by bacteria encoding Ku, the key protein of the NHEJ pathway. This mechanism results

in the ligation of two linear double stranded DNA molecules even if they may lack sequence

similarity (44). Recent works have shown that NHEJ-like activity could occur even in the

absence of Ku, *e.g.* it was found to recombine non-homologous DNA in *E. coli* (45) and the

T4 ligase performs NHEJ-like functions with high efficiency in the absence of Ku (46, 47). The

poor understanding of these Ku-independent processes precludes assessing their role in

phage recombination, but they could contribute to explain gene flux between unrelated

phages. In summary, the presence of prophages in many bacterial genomes provides ample

opportunity for recombination with infecting virulent phages, and these genetic exchanges

could occur by a multitude of molecular processes.

Recombination between temperate and virulent phages allows transfer of functional

innovations from one group to the other. This has the potential to facilitate phage adaptation

to novel challenges. When investigating the functions of exchanged genes, we found proteins

involved in the production of virions, but also bacteriocins or RNA polymerases. Some

exchanged proteins are recombinases and their transfer could disseminate the potential for

further genetic exchanges. Moreover, a recent work found evidence that anti-CRISPR

proteins, used by phage to evade their hosts' CRISPR defences, can be exchanged between

temperate and virulent phages (48).

Our findings suggest that the impact of recombination in phage functional and morphological

diversification can be enhanced by the differences in host range between temperate and

virulent phages. We observed that virulent phages have broader host ranges, which suggests

that temperate phages could be more constrained in their host range by the additional genetic

interactions they establish with their hosts when deciding between lysis and lysogeny and

during the lysogenic cycle. These processes require some integration of the phage genome

on the cell's genetic network, e.g. the development of genetic regulation for induction under

the control of the host SOS response. This integration adds constraints to the genome of

temperate phages and may contribute to their narrower host ranges. There is some published

evidence on differences in host ranges between temperate and virulent phages. For example,

virulent coliphages isolated from the faeces of toddlers infect a broader range of gut strains

than temperate ones (49). The implications of our findings are that groups of temperate

phages that might be sexually isolated (because they infect phylogenetically distant hosts)

can exchange genes indirectly through recombination with broader host range virulent

phages. Conversely, it was observed that virulent-to-virulent genetic exchanges are rare

outside restricted taxonomic groups (23, 24). The genetic flux between virulent phages, and

also between virulent and temperate phages, might then increase when they infect hosts

whose genomes have many prophages.

The expression of prophage genes can change bacterial phenotypes. This so-called lysogenic

conversion has been shown to have an important impact on pathogenicity, growth and stress

responses (50, 51). We analysed the functions of genes exchanged between temperate and

virulent phages that are not typically found in phages and tend to be associated with bacteria.

Although the permanence of genes with bacterial functions might be, for most cases, transient

in virulent genomes, there are examples of such genes that have been acquired and

repurposed by those phages. For instance, bacterial photosynthetic genes encoded by

genomes of virulent phages were shown to increase viral progeny (32, 52). These genes are

often referred to as phage auxiliary metabolic genes (53). Amongst other genes with bacterial

functions, they were identified as being exchanged between temperate and virulent phages in

our data, which opens the possibility that they were acquired by the latter through this

mechanism. Interestingly, photosynthetic genes were previously shown to be exchanged

between *Prochlorococcus* and *Synechococcus* through viral intermediates (54), suggesting

that gene flow with and between phages plays an active role in the evolution of cyanobacterial

photosynthesis. Thus, in addition to the acquisition of genes from bacterial genomes (55) and

other prophages (56), exchanges between virulent and temperate phages could enhance the

genetic repertoire of prophages with functions adaptive to the bacterial host.

Given the pervasiveness of prophages in bacterial genomes, recombination between

temperate and virulent phages might be more frequent than expected based on their low

global sequence similarity, and perhaps even more than what we report here. This could have

implications for the use of phage therapy, because it suggests that there may be exchanges

between temperate and virulent phages during therapy and the former may favour transfer to

and between bacterial genomes. The assessment of safety in therapies should account for

this factor. At this stage, the rates of exchange between the two populations of phages seem

low to dramatically change the flux of genetic information between bacteria at the very short

time scale of an infection. Furthermore, bacteria themselves can exchange DNA through many

other means (transduction, conjugation, transformation, etc). In any case, the frequency of

recombination events between temperate and virulent phages, coupled with the ability of

virulent phages to shuttle genes between temperate phages that are segregated in distinct

clades, can contribute to the high rates of diversification of phage genomes, as well as to the

expansion of gene repertoires of lysogenic bacteria.

Methods

Data

We retrieved the complete genomes of 13513 bacteria and 2502 phages from NCBI non-redundant RefSeq database (ftp://ftp.ncbi.nlm.nih.gov/genomes/refseq/, last accessed in May 2019). The lifestyle of these phages was predicted using PHACTS v.0.3 (31). Predictions were considered as confident if the average probability score of the predicted lifestyle was at least two standard deviations (SD) away from the average probability score of the other lifestyle, which leads a precision rate in lifestyle identification of 99% (31). Using these criteria, we classified as confident 54% of the phages into 571 virulent and 780 temperate phages.

We retrieved information (when available) on the phage hosts from the Genbank files of the phages or from the Virus-Host DB (57) (https://www.genome.jp/virushostdb/). A few phages (~2%) have no identified host, since they were collected from environmental (soil, sea, etc.) or faeces samples. In this dataset, the phage hosts belong to 332 species and 145 genera. Most of these species (69%) have at least one complete genome sequenced and available in NCBI RefSeq. In the absence of a complete genome of the same species (31%, indicated as "manually annotated" in File S1), we used another genome from the same genus as a proxy for the species. Some analysis are repeated by excluding these phages with manually annotated (non-confident) host species, in order to provide a conservative control for these assignments.

Prophages (integrated temperate phages) were predicted using VirSorter v.1.0.3 (58) with the RefSeqABVir database in these bacterial genomes (corresponding to the host species/genus of the phages). The least confident predictions, i.e., categories 3 and 6, which may be prophage remnants or erroneous assignments, were excluded from the analyses. We also retrieved the phage family from the Genbank files of each phage. Most of them are Caudovirales (94%) and belong to the 3 main phage families, i.e., Siphoviridae (1291), Myoviridae (632) and Pododviridae (383). The complete dataset of phage genomes can be found in File S1.

Protein similarity and weighted gene repertoire between bacteriophage genomes

We searched for sequence similarity between all proteins of all phages using mmseqs2 (59)

(Nature Biotechnology release, August 2017) with the sensitivity parameter set at 7.5. The

results were converted to the blast format for analysis and we kept for analysis the hits

respecting the following thresholds: e-value lower than 0.0001, at least 35% identity, and a

coverage of at least 50% of the proteins. The hits were used to compute the bi-directional best

hits between pair of phages, which were used to compute a score of gene repertoire

relatedness weighted by sequence identity:

586
$$wGRR = \frac{\sum_i^p id(A_i, B_i)}{\min(A, B)}$$

where A_i and B_i is the pair *i* of homologous proteins present in *A* and *B*, id(A_i,B_i) is the percent

sequence identity of their alignment, and min(*A*,*B*) is the number of proteins of the genome

encoding the fewest proteins (*A* or *B*). wGRR is the fraction of bi-directional best hits between

two genomes weighted by the sequence identity of the homologs. It varies between zero (no

bi-directional best hits) and one (all genes of the smallest genome have an identical homolog

in the largest genome). wGRR integrates information on the frequency of homologs and

sequence identity. Hence, when the smallest genome is 100 proteins long, a wGRR of 0.03

can result from three pairs of identical homologs or six pairs of homologs with 50% identity.

Similarity networks, community detection and calculation of entropy

The phage network was built based on the filtered wGRR values, using the *networkx* and

graphviz Phyton (v2.7) packages, and the *neato* algorithm. The Louvain community clusters

(60) were calculated using the *best_partition* function from the *community* package in Python

(v2.7). For each cluster (considering only those with at least 3 nodes), we first calculated their

total number of nodes, and then the number of nodes corresponding to each category: nodes

in either one of two lifestyles (temperate or virulent), nodes with a given phage family and

nodes with a given described bacterial host phyla. The Shannon entropy of a cluster (X) with

nodes classed according to a given variable that takes t different values ($y_{1,X}$... $y_{t,X}$) was

calculated as:

$$Entropy(X) = -\sum_{i \in \{1,..,t\}} f(y_{i,X}) * \log_t f(y_{i,X})$$

Where $f(y_{i,X})$ is the relative frequency of the nodes classed i in the cluster. As an example,

consider a cluster composed of 20 nodes, where 10 of them have a temperate lifestyle and

10 have a virulent lifestyle. Because there are two labels (temperate or virulent), $t = 2$. If the

frequencies of the two types of phages were identical, close to the expectation under a random

distribution, the two frequencies would be 0.5 and the entropy would amount to 1, which is the

maximum and corresponds to a highly heterogeneous cluster. When all phages are from one

single type, the entropy is equal to 0, corresponding to an homogeneous cluster. To be able

to compare each trait to the expectation of random cluster compositions, we randomly re-

assigned the labels (host phyla, phage family and lifestyle) to the phages and computed the

Shannon entropy as described above. The results (shown as boxplots in the figures)

summarise the distribution of the 100 random experiments.

Identification of pairs of phage genomes with recombinant genes

In order to separate events of recombination and from homology due to shared ancestry, we

used the relationship between the fraction of homologous proteins (over the average number

of proteins for each phage pair) and the fraction of homologous proteins with very high identity

(at least 80% identity, also over the average number of proteins for each phage pair). We used

the *ols* function from the *statsmodels* package in Python (v2.7) to generate a linear model

describing the expectation of ancestry between phage genomes. To characterize common

ancestry as robustly as possible, we restricted the dataset to temperate-virulent pairs of

phages, as this dataset is expected to contain the lowest frequency of recombination events.

Further, and in order to reduce the influence of the outliers associated with recombination (and

thus maximize the fit of the linear model to cases of common ancestry), we analysed only the

pairs of phage with at least 50% of the proteins in their (average) genomes classified as

homologous (meaning at least 35% identity and 50% coverage) and a minimum of one highly

similar homologous protein (at least 80% identity). Using this dataset, we applied residual

analysis (with the *fit* and *outlier_test* function from *statsmodels*) to identify the first significant

residual value. This was defined as the least distant residual from the linear model that is

classed as outside the confidence interval, with an adjusted pvalue<0.05 using the

Benjamin/Hochberg method – *fdr_bh*. The value of this residual is assumed as the significance

threshold beyond which a residual is classified as a putative recombination event (since it

significantly departs from the expectation of common ancestry defined by the linear model).

Thus, the linear model and the residual threshold are applied to the entire range of the data

across the three datasets (temperate-virulent, temperate-temperate and virulent-virulent

pairs). The pairs of phages whose residuals are larger than the minimum significant residual

threshold (and whose wGRR are below 0.5 to avoid high similarity caused by recent

divergence of the lineages) are classed as pairs of putatively recombined phages. The

recombinant genes are identified as the proteins with at least 80% identity between the pairs

of phages showing evidence of recombination.

Functional annotation of proteins

We used HMMER v3.1b2 (61) (default parameters) to search for genes matching the

prokaryotic Virus Orthologous Groups (pVOGS) (Version May 2016 (62)) database of hmm

profiles (filters used were e-value<1e-5 and profile coverage>60%). Only pVOGs with a viral

quotient (VQ) above 0.8 were used (7751 out of 9518 pVOG profiles in total). The pVOG

profiles were classed into seven functional categories: a) structure, b) lysis, c) packaging,

maturation/Assembly and DNA injection, d) DNA metabolism, recombination and regulation,

e) others and f) unknown by two approaches. First a profile-profile comparison was done using

the HHsuite 2.0.9 (63) with phage-specific profiles from the PFAM (64) and TIGRFAM (65)

database (taken from (66)). Applying a threshold of p-value $<10^{-5}$ resulted in 711 profiles that

cluster in 261 groups using the Louvain algorithm. The annotations of the PFAM and

TIGFRAM profiles were used to assign one of the functional categories to a group. The

remaining profiles were manually assigned considering the piled-up annotations of the pVOGs

(File S3). The identification of bacterial functions was performed based the EggNOG database

of hmm profiles (bactNOG Version (67)). In order to minimize the number of bactNOG entries

that derive from prophages in bacterial genomes, we used hhsearch (HHSuite version 3.2

(68)) to remove from bactNOG the profiles matching pVOG profiles. The bactNOG profiles

with matches in pVOGs with VQ above 0.8 (p-value <0.0001) were discarded. This resulted

in a reduction of 34% of the bactNOG profiles' dataset. The remaining 135814 bactNOG

profiles were used to class the dataset of putative recombinant genes (filters used were e-

value<1e-5 and profile coverage>60%) in broad functional categories.

Detection of phage-like recombinases in phage and bacterial genomes

The families of recombinases of phage were described in (69), for which we built profiles or

recovered them from PFAM. To build the profiles, we retrieved the homologs given in the

reference above and aligned them using default options with MAFFT (v7.407) (70). The

alignments were used to build the profiles using hmmbuild from HMMER (default options).

Our novel profiles are given in supplementary material. We used HMMER to search for

homologs of Sak (PF04098, --cut_ga), Sak4 (Sak4 from phage T7, minimum score = 20), Erf

(PF04404, --cut_ga), RecT (PF03837, --cut_ga) and gp2.5 (gp2.5 from phage T7, minimum

score = 20). An additional profile, RecA (PF00154, --cut_ga) was searched only in phage

genomes. Hosts were retrieved from the GenBank file of each phage, and all the genomes in

the database belonging to that host's species were used for the calculation. E.g., if the host

species of a given phage is described as *E. coli*, we calculated the mean number of

recombinases in all *E. coli* genomes. The values shown on the table are the mean of these

means for phages with recombinant genes and the remaining phages. If recombinases were

found within the coordinates of a prophage, they were associated with prophage regions. Note

that a particular phage host can be included multiple times in the distribution, if subsequent

phages have a similar host, and can even result in the host being included in the dataset of

recombinant and non-recombinant phages simultaneously. Although this results in repeated

sampling, it represents the likelihood that hosts of recombinant or non-recombinant phages

encode for recombinases. Moreover, this is unbiased for either class of phages, since the

hosts of both recombinant and non-recombinant phages are subject to this repeated sampling

process.

Detection of NHEJ in bacterial genomes

We used HMMER to search for homologs of the Ku protein (PF02735.16) in the dataset of

bacterial genomes, using the -cut_ga parameter. We did not require the presence of a

neighboring ligase to consider the system complete, because such a ligase is often absent

(71). The set of bacterial genomes assigned as host of a given phage was asserted as

described in the section above. The fraction of hosts that encode for NHEJ was calculated as

the number of genomes that where at least one homolog of Ku was found, divided by the total

number of genomes considered for a given phage. The use of the presence/absence if Ku as

the key variable, and not the number of Ku genes, is due to three reasons: 1) the numbers

were always low (most often 0 or 1), 2) some genomes encoding multiple NHEJ systems

express them during different times, 3) some bacteria seem to encode heterodimeric Ku where

two genes are necessary for the process (72).

Calculation of the patristic distance between bacterial hosts

We used the 16S rRNA of the bacterial genomes identified in the RefSeq annotations,

corresponding to the species level of the identified phage hosts. We selected the first entry of

each genome and aligned them using the secondary structure models with the program

SSU_Align version 0.1.1 (73). Poorly aligned positions were eliminated with the ssu-mask.

The alignment was trimmed with trimAl version 1.2 (74) using the option –noallgaps to delete

only the gap proteins but not the regions that are poorly conserved. The 16S rRNA

phylogenetic tree was inferred using maximum likelihood with IQTREE version 1.6.5 (75)

under the best-fit model automatically selected by ModelFinder (76), and with the options –

wbtl (to conserve all optimal trees and their branch lengths) and –bb 1000 to run the ultrafast

bootstrap option with 1000 replicates. The patristic distances amongst the taxa in the 16S

trees were calculated from the tree (weighted by the edge distances) with the *dendropy*

package in Python (77), using the functions *phylogenetic_distance_matrix* and

patristic_distance, with the default parameters.

Calculation of the genetic distance between bacterial hosts

We calculated the trinucleotide composition, or 3-mer genomic signature, of each bacterial

genome from the phage host species, using the relative abundance value of each of the 64

possible trinucleotides, as in (78). This is defined as the observed trinucleotide frequency

divided by the expected trinucleotide frequency (the product of the mononucleotide

frequencies), or

726
$$x_{ijk} = \frac{f_{ijk}}{f_i f_j f_k}$$

where f_i, f_j and f_k represent the frequency of the nucleotides i, j and k, respectively (with i, j,

$k \in$ A, C, T, G). This allows to quantify the deviation of the observed frequency of trinucleotides

to the one expected given the nucleotide composition of the genomes, which are known to

differ between phages and bacteria (79). The genetic distance between two hosts was then

calculated using the average absolute difference between the 3-mer genomic composition of

each, as

733
$$\delta(a,b) = \frac{1}{64} \sum_{i \in \{ACGT\}} \sum_{j \in \{ACGT\}} \sum_{k \in \{ACGT\}} |x_{ijk,a} - x_{ijk,b}|$$

where $x_{ijk,a}$ and $x_{ijk,b}$ are the relative abundances of the trinucleotide *ijk* in each of the

bacterial hosts. Note that multiple genomes might be available for each host. To use all the

available information when multiple genomes are available for a species, we used the grand

mean of all pairwise comparisons. As an example, if one host is *Escherichia coli* and the

another is *Staphylococcus aureus*, we calculated the differences in genomic signature

between each pairwise combinations of all genomes of *Escherichia coli* and all genomes of

Staphylococcus aureus. We then calculate the average of all the pairwise calculations of the

genetic distance as δ_T, with

$$\delta_T(a, b) = \frac{1}{N}\sum_a\sum_b \delta(a, b)$$

where *a* and *b* are individual strain genomes of the first and second host, respectively, and *N*

is the total pairwise calculations between the strains of the first and second hosts.

ACKNOWLEDGEMENTS

We acknowledge support by the PRESTIGE programme (PRESTIGE-2017-1-0012), the ANR

grants (ANR-16-CE15-0022 and ANR-16-CE16-0029), the Fondation pour la Recherche

Médicale (Equipe FRM EQU201903007835), and the Laboratoire d'Excellence IBEID (ANR-

10-LABX-62-IBEID). We thank Mireille Ansaldi, Marta Lourenço and the members of the

Microbial Evolutionary Genomics laboratory by discussions and comments on earlier versions

of this manuscript. This work used the computational and storage services (TARS cluster)

provided by the IT department at Institut Pasteur, Paris.

AUTHOR CONTRIBUTIONS

Designed research: JMS, EPCR. Analyzed the data: JMS. Visualization: JMS. Provided

materials and methods: EP, JMS, MT. Drafted the paper: JMS, EPCR. Revised the

manuscript: all authors.

REFERENCES

1.   Dion MB, Oechslin F, Moineau S (2020) Phage diversity, genomics and phylogeny.
767   *Nat Rev Microbiol* 437:356–14.

768   2.   Harrison E, Brockhurst MA (2017) Ecological and Evolutionary Benefits of Temperate
769   Phage: What Does or Doesn't Kill You Makes You Stronger. *Bioessays*
770   197(12):1700112.

771   3.   Touchon M, Moura de Sousa JA, Rocha EP (2017) Embracing the enemy: the
772   diversification of microbial gene repertoires by phage-mediated horizontal gene
773   transfer. *Curr Opin Microbiol* 38:66–73.

774   4.   Touchon M, Bernheim A, Rocha EP (2016) Genetic and life-history traits associated
775   with the distribution of prophages in bacteria. *ISME J* 10(11):2744–2754.

776   5.   Sausset R, Petit MA, Gaboriau-Routhiau V, De Paepe M (2020) New insights into
777   intestinal phages. *Mucosal Immunol* 16(11):760–11.

778   6.   Canchaya C, Fournous G, Brüssow H (2004) The impact of prophages on bacterial
779   chromosomes. *Mol Microbiol* 53(1):9–18.

780   7.   Penadés JR, Chen J, Quiles-Puchalt N, Carpena N, Novick RP (2015) Bacteriophage-
781   mediated spread of bacterial virulence genes. *Curr Opin Microbiol* 23:171–178.

782   8.   Fillol-Salom A, et al. (2019) Bacteriophages benefit from generalized transduction.
783   *PLoS Pathog* 15(7):e1007888.

784   9.   Hatfull GF, Hendrix RW (2011) Bacteriophages and their genomes. *Curr Opin Virol*
785   1(4):298–303.

786   10.   Arbiol C, Comeau AM, Kutateladze M, Adamia R, Krisch HM (2010) Mobile
787   Regulatory Cassettes Mediate Modular Shuffling in T4-Type Phage Genomes.
788   *Genome Biol Evol* 2:140–152.

789   11.   Devoto AE, et al. (2019) Megaphages infect Prevotella and variants are widespread in
790   gut microbiomes. *Nat Microbiol* 4(4):1–11.

791   12.   Yuan Y, Gao M (2017) Jumbo Bacteriophages: An Overview. *Front Microbiol* 8:403.

792   13.   Botstein D (1980) A theory of modular evolution for bacteriophages. *Ann N Y Acad*
793   *Sci* 354:484–490.

794   14.   Hendrix RW, Lawrence JG, Hatfull GF, Casjens S (2000) The origins and ongoing
795   evolution of viruses. *Trends Microbiol* 8(11):504–508.

796   15.   Campbell A, Botstein D (1983) *Evolution of the lambdoid phages* (Cold Spring Harbor
797   Laboratory, Cold Spring Harbor, NY).

798   16.   Smith GR (1983) *General Recombination* (Cold Spring Harbor, NY: Cold Spring
799   Harbor Laboratory).

800   17.   De Paepe M, et al. (2014) Temperate Phages Acquire DNA from Defective
801   Prophages by Relaxed Homologous Recombination: The Role of Rad52-Like

Recombinases. *PLoS Genet* 10(3):e1004181–15.

18. Morris P, Marinelli LJ, Jacobs-Sera D, Hendrix RW, Hatfull GF (2008) Genomic characterization of mycobacteriophage Giles: evidence for phage acquisition of host DNA by illegitimate recombination. *J Bacteriol* 190(6):2172–2182.

19. d'Adda di Fagagna F, Weller GR, Doherty AJ, Jackson SP (2003) The Gam protein of bacteriophage Mu is an orthologue of eukaryotic Ku. *EMBO Rep* 4(1):47–52.

20. Mavrich TN, Hatfull GF (2017) Bacteriophage evolution differs by host, lifestyle and genome. *Nat Microbiol* 2:1–9.

21. Filée J, Tétart F, Suttle CA, Krisch HM (2005) Marine T4-type bacteriophages, a ubiquitous component of the dark matter of the biosphere. *Proc Natl Acad Sci USA* 102(35):12471–12476.

22. Ignacio-Espinoza JC, Sullivan MB (2012) Phylogenomics of T4 cyanophages: lateral gene transfer in the "core" and origins of host genes. *Environ Microbiol* 14(8):2113–2126.

23. Chopin A, Bolotin A, Sorokin A, Ehrlich SD, Chopin M (2001) Analysis of six prophages in Lactococcus lactis IL1403: different genetic structure of temperate and virulent phage populations. *Nucleic Acids Res* 29(3):644–651.

24. Lima-Mendez G, Van Helden J, Toussaint A, Leplae R (2008) Reticulate Representation of Evolutionary and Functional Relationships between Phage Genomes. 25(4):762–777.

25. Kupczok A, et al. (2018) Rates of Mutation and Recombination in Siphoviridae Phage Genome Evolution over Three Decades. *Mol Biol Evol* 35(5):1147–1159.

26. Filée J, Bapteste E, Susko E, Krisch HM (2006) A selective barrier to horizontal gene transfer in the T4-type bacteriophages that has preserved a core genome with the viral replication and structural genes. *Mol Biol Evol* 23(9):1688–1696.

27. Dekel-Bird NP, et al. (2013) Diversity and evolutionary relationships of T7-like podoviruses infecting marine cyanobacteria. *Environ Microbiol* 15(5):1476–1491.

28. Bouchard JD, Moineau S (2000) Homologous Recombination between a Lactococcal Bacteriophage and the Chromosome of Its Host Strain. *Virology* 270(1):65–75.

29. Garneau JE, Tremblay DM, Moineau S (2008) Characterization of 1706, a virulent phage from Lactococcus lactis with similarities to prophages from other Firmicutes. *Virology* 373(2):298–309.

30. Martin DP, Lemey P, Posada D (2011) Analysing recombination in nucleotide sequences. *Mol Ecol Resour* 11(6):943–955.

31. McNair K, Bailey BA, Edwards RA (2012) PHACTS, a computational approach to classifying the lifestyle of phages. *Bioinformatics* 28(5):614–618.

32. Mann NH, Cook A, Millard A, Bailey S, Clokie M (2003) Bacterial photosynthesis genes in a virus. *Nature* 424(6950):741–741.

33. Shen P, Huang HV (1986) Homologous recombination in Escherichia coli: dependence on substrate length and homology. *Genetics* 112(3):441–457.

34. Rocha EPC, Cornet E, Michel B (2005) Comparative and evolutionary analysis of the
bacterial homologous recombination systems. *PLoS Genet* 1(2):e15.

35. Aravind L, Koonin EV (2001) Prokaryotic homologs of the eukaryotic DNA-end-
binding protein Ku, novel domains in the Ku protein and prediction of a prokaryotic
double-strand break repair system. *Genome Res* 11(8):1365–1374.

36. Bhattacharyya S, et al. (2018) Phage Mu Gam protein promotes NHEJ in concert with
Escherichia coli ligase. *Proc Natl Acad Sci USA* 115(50):E11614–E11622.

37. Latino L, Essoh C, Blouin Y, Vu Thien H, Pourcel C (2014) A novel Pseudomonas
aeruginosa Bacteriophage, Ab31, a Chimera Formed from Temperate Phage PAJU2
and P. putida Lytic Phage AF: Characteristics and Mechanism of Bacterial
Resistance. *PLoS ONE* 9(4):e93777–16.

38. Durmaz E, Klaenhammer TR (2000) Genetic analysis of chromosomal regions of
Lactococcus lactis acquired by recombinant lytic phages. *Appl Environ Microbiol*
66(3):895–903.

39. Schuch R, Fischetti VA (2006) Detailed genomic analysis of the Wbeta and gamma
phages infecting Bacillus anthracis: implications for evolution of environmental fitness
and antibiotic resistance. *J Bacteriol* 188(8):3037–3051.

40. Hershey AD (1953) Nucleic acid economy in bacteria infected with bacteriophage T2.
J Gen Physiol 37(1):1–23.

41. Wiberg JS (1966) Mutants of bacteriophage T4 unable to cause breakdown of host
DNA. *Proc Natl Acad Sci USA* 55(3):614–621.

42. Arber W (2000) Genetic variation: molecular mechanisms and impact on microbial
evolution. *FEMS Microbiol Rev* 24(1):1–7.

43. Labrie SJ, Moineau S (2007) Abortive Infection Mechanisms and Prophage
Sequences Significantly Influence the Genetic Makeup of Emerging Lytic Lactococcal
Phages. *J Bacteriol* 189(4):1482–1487.

44. Shuman S, Glickman MS (2007) Bacterial DNA repair by non-homologous end
joining. *Nat Rev Microbiol* 5(11):852–861.

45. Chayot R, Montagne B, Mazel D, Ricchetti M (2010) An end-joining repair mechanism
in Escherichia coli. *Proc Natl Acad Sci USA* 107(5):2141–2146.

46. Su T, et al. (2019) The phage T4 DNA ligase mediates bacterial chromosome DSBs
repair as single component non-homologous end joining. *Synth Syst Biotechnol*
4(2):107–112.

47. Wu DY, Wallace RB (1989) Specificity of the nick-closing activity of bacteriophage T4
DNA ligase. *Gene* 76(2):245–254.

48. Hynes AP, et al. (2018) Widespread anti-CRISPR proteins in virulent bacteriophages
inhibit a range of Cas9 proteins. *Nat Commun* 9(1):2919–10.

49. Mathieu A, et al. (2020) Virulent coliphages in 1-year-old children fecal samples are
fewer, but more infectious than temperate coliphages. *Nat Commun*:1–12.

50. Brüssow H, Canchaya C, Hardt W-D (2004) Phages and the evolution of bacterial

pathogens: from genomic rearrangements to lysogenic conversion. *Microbiol Mol Biol*
Rev 68(3):560–602– table of contents.

51.   Obeng N, Pratama AA, van Elsas JD (2016) The Significance of Mutualistic Phages
for Bacterial Ecology and Evolution. *Trends Microbiol* 24(6):1–10.

52.   Lindell D, et al. (2004) Transfer of photosynthesis genes to and from Prochlorococcus
viruses. *Proc Natl Acad Sci USA* 101(30):11013–11018.

53.   Thompson LR, et al. (2011) Phage auxiliary metabolic genes and the redirection of
cyanobacterial host carbon metabolism. *Proc Natl Acad Sci USA* 108(39):E757–64.

54.   Zeidner G, et al. (2005) Potential photosynthesis gene recombination between
Prochlorococcus and Synechococcus via viral intermediates. *Environ Microbiol*
7(10):1505–1513.

55.   Mata M, Delstanche M, Robert-Baudouy J (1978) Isolation of specialized transducing
bacteriophages carrying the structural genes of the hexuronate system in Escherichia
coli K-12: exu region. *J Bacteriol* 133(2):549–557.

56.   Mirold S, Rabsch W, Tschäpe H, Hardt WD (2001) Transfer of the Salmonella type III
effector sopE between unrelated phage families. *J Mol Biol* 312(1):7–16.

57.   Mihara T, et al. (2016) Linking Virus Genomes with Host Taxonomy. *Viruses* 8(3):66.

58.   Roux S, Enault F, Hurwitz BL, Sullivan MB (2015) VirSorter: mining viral signal from
microbial genomic data. *PeerJ* 3(348):e985.

59.   Steinegger M, Söding J (2017) MMseqs2 enables sensitive protein sequence
searching for the analysis of massive data sets. *Nat Biotechnol* 35(11):1026–1028.

60.   Blondel VD, Guillaume J-L, Lambiotte R, Lefebvre E (2008) Fast unfolding of
communities in large networks. *J Stat Mech* 2008(10):P10008.

61.   Eddy SR (2011) Accelerated Profile HMM Searches. *PLoS Comput Biol*
7(10):e1002195.

62.   Grazziotin AL, Koonin EV, Kristensen DM (2017) Prokaryotic Virus Orthologous
Groups (pVOGs): a resource for comparative genomics and protein family annotation.
Nucleic Acids Res 45(D1):D491–D498.

63.   Söding J (2005) Protein homology detection by HMM-HMM comparison.
Bioinformatics 21(7):951–960.

64.   El-Gebali S, et al. (2019) The Pfam protein families database in 2019. *Nucleic Acids*
Res 47(D1):D427–D432.

65.   Haft DH, Selengut JD, White O (2003) The TIGRFAMs database of protein families.
Nucleic Acids Res 31(1):371–373.

66.   Fouts DE (2006) Phage_Finder: automated identification and classification of
prophage regions in complete bacterial genome sequences. *Nucleic Acids Res*
34(20):5839–5851.

67.   Huerta-Cepas J, et al. (2019) eggNOG 5.0: a hierarchical, functionally and
phylogenetically annotated orthology resource based on 5090 organisms and 2502

viruses. *Nucleic Acids Res* 47(D1):D309–D314.

68.   Steinegger M, et al. (2019) HH-suite3 for fast remote homology detection and deep
protein annotation. *BMC Bioinformatics* 20(1):473–15.

69.   Lopes A, Amarir-Bouhram J, Faure G, Petit M-A, Guerois R (2010) Detection of novel
recombinases in bacteriophage genomes unveils Rad52, Rad51 and Gp2.5 remote
homologs. *Nucleic Acids Res* 38(12):3952–3962.

70.   Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version
7: improvements in performance and usability. *Mol Biol Evol* 30(4):772–780.

71.   Bernheim A, Bikard D, Touchon M, Rocha EPC (2019) A matter of background: DNA
repair pathways as a possible cause for the sparse distribution of CRISPR-Cas
systems in bacteria. *Philos Trans R Soc Lond, B, Biol Sci* 374(1772):20180088.

72.   Bertrand C, Thibessard A, Bruand C, Lecointe F, Leblond P (2019) Bacterial NHEJ: a
never ending story. *Mol Microbiol* 111(5):1139–1151.

73.   Nawrocki EP, Kolbe DL, Eddy SR (2009) Infernal 1.0: inference of RNA alignments.
Bioinformatics 25(10):1335–1337.

74.   Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T (2009) trimAl: a tool for
automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*
25(15):1972–1973.

75.   Nguyen L-T, Schmidt HA, Haeseler von A, Minh BQ (2015) IQ-TREE: a fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol*
Evol 32(1):268–274.

76.   Kalyaanamoorthy S, Minh BQ, Wong TKF, Haeseler von A, Jermiin LS (2017)
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods*
14(6):587–589.

77.   Sukumaran J, Holder MT (2010) DendroPy: a Python library for phylogenetic
computing. *Bioinformatics* 26(12):1569–1571.

78.   Suzuki H, Yano H, Brown CJ, Top EM (2010) Predicting plasmid promiscuity based
on genomic signature. *J Bacteriol* 192(22):6045–6055.

79.   Rocha EPC, Danchin A (2002) Base composition bias might result from competition
for metabolic resources. *Trends Genet* 18(6):291–294.

A

B

A

○ Temperate phages
· Virulent phages

B

NC021782 (*Salmonella*)

NC024122 (*Salmonella*)

Identity>80%
60%<Identity<80%
40%<Identity<60%
Identity<40%

NC023589 (*Shigella*)

20 kb

NC029029 (*Brevibacillus*)

NC028841 (*Paenibacillus*)

NC041867 (*Paenibacillus*)

5 kb

C

NC023574 (*Lactococcus*)

NC029058 (*Lactobacillus*)

NC020197 (*Streptococcus*)

20 kb

NC031025 (*Klebsiella*)

NC029070 (*Cronobacter*)

NC031068 (*Pectobacterium*)

5 kb