From Value to Saliency: Neural Computations of Subjective Value under Uncertainty in Combat Veterans

Authors: Ruonan Jia[1,2], Lital Ruderman[2], Charles Gordon[3,4], Daniel Ehrlich[1], Mark Horvath[3,4], Serena Mirchandani[3,4], Clara DeFontes[3,4], Steven Southwick[3,4], John H. Krystal[3,4,5,6], Ilan Harpaz-Rotem[3,4,*], Ifat Levy[1,2,3,5,6,*]

1. Interdepartmental Neuroscience Program, Yale University; 2. Department of Comparative Medicine, Yale University; 3. National Center for PTSD, West Haven VA Medical Center; 4. Department of Psychiatry, Yale University; 5. Department of Neuroscience, Yale University; 6. Department of Psychology, Yale University

* These authors contributed equally.

Abstract

Military personnel engaged in combat are vulnerable to Posttraumatic Stress Disorder (PTSD), following traumatic experiences in the battlefield. Prior research has mostly employed fear-related paradigms to unravel neural underpinnings of fear dysregulation in individuals with PTSD. The ability to acquire and update fear responses depends critically on the individual's ability to cope with uncertainty, yet the role of individual uncertainty attitudes in the development of trauma-related psychopathology has hardly been examined. Here, we investigated the association between PTSD-related alterations and the subjective valuation of uncertain outcomes during decision-making. We used a monetary gambling paradigm inspired by behavioral economics in conjunction with fMRI and explored neural markers of both vulnerability and resilience to PTSD in a group of combat veterans. Behaviorally, PTSD symptom severity was associated with increased aversion to uncertainty. Neurally, activity in the ventromedial prefrontal cortex (vmPFC) during valuation of uncertain options was associated with PTSD symptoms, an effect which was specifically

driven by numbing symptoms. Moreover, the neural encoding of the subjective value of those uncertain options was markedly different in the brains of veterans diagnosed with PTSD, compared to veterans who experienced trauma but did not develop PTSD. Most notably, veterans with PTSD exhibited enhanced representations of the saliency of rewards and punishments in the neural valuation system, especially in ventral striatum, compared with trauma-exposed controls. Our results point to a link between the function of the valuation system under uncertainty and the development and maintenance of PTSD symptoms, and stress the significance of studying reward processes in PTSD.

Introduction

Following a life-threatening experience, some individuals develop Posttraumatic Stress Disorder (PTSD) symptoms, which include re-experiencing the traumatic event, avoidance of trauma reminders, emotional numbing, and exaggerated arousal and reactivity. These symptoms can be socially and vocationally disabling. While medications and psychotherapy help some individuals, many people with PTSD remain symptomatic following treatment (Foa, Keane, and Friedman 2000). A better understanding of the neural basis of PTSD is crucial, as it can inform new approaches to individualized treatment.

Soldiers in combat face highly uncertain life-threatening events, which are uncontrollable (van der Kolk et al. 1985), and that may result in serious injury to themselves or death of teammates. An individual's attitude towards uncertainty and their capacity to handle uncertainty may therefore affect one's ability to cope with potentially traumatic events. The notion of uncertainty was incorporated in studies of fear-learning attempting to unravel the behavioral and neural mechanisms of PTSD (Brown et al. 2018; Homan et al. 2019). Participants in these studies encountered probabilistic deliveries of adverse outcomes (e.g. electric shocks), and their ability to predict these outcomes was measured (e.g. by their skin conductance responses). In a separate line of work, using a behavioral economic framework, our group showed increased aversion to ambiguity (an uncertain situation where outcome probabilities are not known) in combat veterans with PTSD compared to trauma-exposed veterans without PTSD when choosing between potential monetary

losses (Ruderman et al. 2016). This aversion to uncertainty demonstrated in situations unrelated to the trauma, may also contribute to the exaggerated behavior in fear conditioning paradigms, and to the development and maintenance of PTSD symptoms. Further understanding of this aversion to uncertainty may also provide evidence for targeting uncertainty in behavioral interventions, to improve the daily decision-making and well-being of individuals with PTSD.

As uncertainty attitudes affect the subjective value of options, it is plausible that neural computations of subjective value are altered in the brains of individuals who developed PTSD following trauma exposure. Such changes may be particularly related to numbing symptoms, which include losing interest in significant activities, having difficulty experiencing happiness or love, and feeling distant from others (Harpaz-Rotem et al. 2014). A few studies indeed showed that PTSD symptoms were associated with changes in neural processing of uncertain monetary outcomes in several brain regions, including medial prefrontal cortex (mPFC) and striatum (Sailer et al. 2008; Elman et al. 2009).

 In the general population, a network of brain regions was implicated in valuation and decision making, including the ventromedial prefrontal cortex (vmPFC), anterior cingulate cortex (ACC), posterior cingulate cortex (PCC), dorsolateral prefrontal cortex (dlPFC), ventral striatum, amygdala, and thalamus (Kable and Glimcher 2009; Gleichgerrcht et al. 2010). While the neural encoding of the subjective value of rewards was extensively studied in the general population, it is still unclear how subjective values of punishments are represented in the brain. On the one hand, there is evidence for a unified valuation system, which encodes the value of both rewards and punishments (Tom et al. 2007; Fujiwara et al. 2009; Bartra, McGuire, and Kable 2013; Kahnt et al. 2014). On the other hand, there is evidence for different brain circuits processing outcomes from the two valence domains (O'Doherty et al. 2001; Yacubian et al. 2006; Kim et al. 2014; Monosov and Hikosaka 2012; Fiorillo 2013). Moreover, while we know that the subjective values of ambiguous gains are encoded in the valuation system (Levy et al. 2010; Blankenstein and van Duijvenvoorde 2019), we are not aware of any examination of the neural encoding of subjective value of ambiguous losses. Here we combined our simple economic task with functional MRI and computational

modeling to examine the neural encoding of subjective value under risk and ambiguity, and the alterations in this encoding in individuals exposed to trauma. We compared combat veterans with PTSD to those who did not develop PTSD symptoms (trauma-exposed controls), and were thus able to investigate both the psychopathology of PTSD and the resilience to PTSD. We find that veterans with PTSD encode the subjective values of uncertain monetary gains and losses in a U-shape manner, with increased activation for both increased gains and increased losses (compatible with saliency encoding). Conversely, trauma-exposed controls encode the same type of subjective values monotonically, with increased activation for increased gains, and decreased activation for increased losses (compatible with value encoding). Our results suggest that this shift from value-encoding to saliency-encoding, especially of ambiguous monetary losses, could be a neural marker for PTSD symptom severity.

Results

In an fMRI experiment, combat veterans with current PTSD diagnoses and those who never developed PTSD completed a gambling task under four decision conditions on two separate days. Participants chose between a sure monetary outcome (either gaining or losing money) and an uncertain outcome (either risky or ambiguous, under gains or losses) (Fig 1). We estimated the uncertainty attitudes of each participant through a behavioral model, and aimed to understand the influence of PTSD symptom severity on both the behavioral attitudes and the neural mechanisms of valuation.

Figure 1. Study design

A: Timeline of the study. Participants went through a screening session and two scanning sessions on three different days. The screening session determined participants' eligibility based on PTSD diagnosis, combat exposure, and exclusion of other neurological disorders. Eligible participants were scanned on two separate days on a decision making task. Measure labels: SCID: Structured Clinical Interview for DSM-4, CAPS: Clinician Administered PTSD Scale, PCL5: , PTSD Checklist for DSM-5 , BDI: Beck Depression Inventory, STAI-1: State Anxiety, STAI-2: Trait Anxiety, DES: Dissociative Experiences Scale, CES: Combat Exposure Scale, CTQ: Childhood Trauma Questionnaire, KBIT: Kaufman Brief Intelligence Test, BIS/BAS: Behavioral Avoidance/Inhibition Scale, BIS-11: Barratt Impulsiveness Scale, DOSPERT: Domain-Specific Risk-Taking Scale. B: Task design: participants chose between a lottery and a sure outcome under four conditions: risky gains, ambiguous gains, risky losses, and ambiguous losses. Lotteries are shown as examples. C: Levels of risk (0.25, 0.5, and 0.75), ambiguity (0.74, 0.5, and 0.24), and monetary outcomes (20 monetary gains and 20

monetary losses) of the lottery. D: On each trial, participants had 6 seconds to view the options, and made a choice following a green response cue. They had a time limit of 3.5 seconds to register the choice, after which they would immediately see a confirmation with the yellow square representing the side they chose. The lottery was not played out during the scan to avoid learning. The inter-trial-interval (ITI) was jittered among 4, 6, and 8 seconds, and the remaining time during the response window (3.5 seconds – response time) would be added to the ITI.

Clinical symptom variation

Participants varied in their PTSD symptom severity (Fig 2A) assessed by the Clinician-Administered PTSD Scale (CAPS) for DSM-4 (Diagnostic and Statistical Manual of Mental Disorders, 4[th] Edition) (Blake et al. 1995). Veterans with PTSD had higher total CAPS score compared to controls (PTSD, N = 23: *Mean* = 72.13, *SD* = 15.04; control, N = 34: *Mean* = 6.21, *SD* = 9.68; $t(34) = 18.58$, $p < 0.001$). PTSD symptoms as captured by the 5-factor model of CAPS (Harpaz-Rotem et al. 2014) were highly correlated with symptoms of depression, anxiety and dissociative experiences (Fig 2B, see Table 1 for descriptive statistics of all measures). In order to account for the influence of clinical symptoms on the behavior and neural activity during the task, we conducted principal component analysis (PCA) on these clinical symptoms. Since the severity of psychopathology may be affected by the degree of stress exposure, we also included measures of combat exposure (CES) and childhood trauma (CTQ) in the PCA. The first three components accounted for ~80% of the variance in those data (Supplementary Fig 1A). The first component was affected by all clinical symptoms (PTSD, depression, anxiety, and dissociative experiences) and might reflect a general affective factor. This component was highly consistent with PTSD symptom severity (correlation with CAPS Spearman's $\rho = 0.94$, $n = 55$, $p < 0.001$), and could be used to clearly classify PTSD diagnosis (Supplementary Fig 1C). The second component was mostly affected by re-experiencing, avoidance and anxious arousal clusters of CAPS, as well as the degree of combat exposure, potentially representing a fear learning-updating deficit or general hyperarousal. The third component was affected by combat exposure and childhood trauma. Components 2 and 3 were not strongly correlated with PTSD symptom severity (n

= 55, Component 2: correlation with CAPS Spearman's $\rho = 0.11$, $p = 0.43$; Component 3: correlation with

CAPS Spearman's $\rho = 0.029$, $p = 0.84$).

Figure 2. Participants' symptom severity

A: Distribution of CAPS total score, colored by group (combat veterans with or without PTSD diagnoses). One PTSD participant included in the analysis did not have complete CAPS data. B: PTSD, depression and anxiety symptom severities were highly correlated. Numbers in the upper right panels indicate pair-wise Pearson correlation coefficients. Significance levels: ***, p < 0.001; **, p < 0.01; *, p < 0.05. Lower left panels show pairwise scatter plots and smoothed curves using locally weighted polynomial regression. Panels in the diagonal show distributions and density curves for each measure. Labels of measures: CAPS-ReExp: re-experiencing, CAPS-Avoid: avoidance, CAPS-Numb: numbing, CAPS-DysA: dysphoric arousal, CAPS-AnxA: anxious arousal, BDI: Beck Depression Inventory, STAI-1: State Anxiety, STAI-2: Trait Anxiety, DES: Dissociative Experiences Scale, CES: Combat Exposure Scale, CTQ: Childhood Trauma Questionnaire.

Table 1. Descriptive statistics of demographics and clinical measures of participants reported in the behavioral results

	PTSD	Control
Number of participants	24	34
Age	34.70 (6.44)	39.17 (10.03)
Kaufman Brief Intelligence Test (KBIT)	105.63 (10.52)	111.59 (13.22)
Clinician Administered PTSD Scale (CAPS)-Total Score	72.13 (15.04)	6.21 (9.68)
CAPS-Re-experiencing	19.83 (6.79), n = 23	1.00 (1.86)
CAPS-Avoidance	10.65 (3.41), n = 23	0.47 (1.38)
CAPS-Emotional Numbing	18.13 (6.61), n = 23	1.03 (3.03)
CAPS-Dysphoric Arousal	15.09 (3.15), n = 23	2.00 (3.65)
CAPS-Anxious Arousal	8.44 (2.56), n = 23	1.71 (2.87)
PTSD Checklist for DSM-5 (PCL-5)	41.54 (15.79)	10.88 (15.95)
Beck Depression Inventory (BDI)	26.11 (13.80)	5.50 (7.88)
State Anxiety (STAI-1)	47.92 (12.72)	32.62 (9.43)
Trait Anxiety (STAI-2)	47.32 (16.23)	31.33 (10.94)
Dissociative Experience Scale (DES)	44.04 (34.53)	17.31 (20.75), n = 33
Combat Exposure Scale (CES)	19.96 (9.06), n = 23	13.00 (8.57)
Childhood Trauma Questionnaire (CTQ)	39.04 (14.03), n = 23	34.43 (8.76)

Mean (standard deviation)

Fewer number of participants for several measures was due to incomplete data.

PTSD symptom severity is associated with increased ambiguity aversion in the loss domain, and increased risk aversion in the gain domain

For each participant, we estimated risk and ambiguity attitudes for gains and losses, using the combined data from both scanning sessions (see equations 1 and 2 in Methods: Model-based risk and ambiguity attitudes estimation; see Supplementary Fig 2A for an example from one participant). We then investigated the associations between these attitudes and PTSD diagnosis status, as well as PTSD symptom severity. All attitudes were transformed such that negative numbers indicate aversion to risk or ambiguity, and positive

numbers indicate seeking. Based on the previous behavioral finding that PTSD symptom severity was associated with higher aversion to ambiguity in losses (Ruderman et al. 2016), we first investigated ambiguity attitudes. At the group level, participants were not significantly averse to ambiguity in the domain of losses (Fig 3A; PTSD: *Mean* = -0.25, $t(23)$ = -1.81, p = 0.11; Control: *Mean* = 0.003, $t(33)$ = 0.040, p = 0.97), and were significantly averse to ambiguity in the domain of gains (Fig 3A; PTSD: *Mean* = -0.35, $t(23)$ = -3.45, $p < 0.01$; Control: *Mean* = -0.42, $t(33)$ = -7.27, $p < 0.001$). However, a two-way ANOVA of ambiguity attitude with domain as the within-subject factor and group as the between-subject factor showed a significant interaction between domain and group ($F(1,56)$ = 4.34, $p < 0.05$, η^2 = 0.0279). Post-hoc comparisons showed that veterans with PTSD were marginally more averse to ambiguity under losses (p = 0.081), but not under gains (p = 0.53). A dimensional analysis (Fig 3B) of this symptom–behavior relationship, regardless of PTSD diagnosis, revealed a negative correlation between ambiguity attitudes in the loss domain and CAPS total score (Spearman's ρ with CAPS total score = -0.30, $p < 0.05$), indicating that higher symptom severity was related to higher aversion to ambiguity under losses. Since many control participants had a CAPS score of zero, we also repeated the analysis using PCL-5 scores instead of CAPS and overserved a similar effect (Supplementary Fig 2B, Pearson's r with PCL-5 = -0.31, $p < 0.05$).

Next, we examined risk attitudes. Both the PTSD and control groups exhibited risk aversion in the domain of gains (PTSD: *Mean* = -0.54, $t(23)$ = -13.34, $p < 0.001$; Control: *Mean* = -0.28, $t(33)$ = -4.80, $p < 0.001$). In the domain of losses, veterans with PTSD exhibited risk seeking (Fig 3C; PTSD: *Mean* = 0.34, $t(23)$ = 5.43, $p < 0.001$), while combat controls exhibited marginal risk seeking (Control: *Mean* = 0.20, $t(33)$ = 1.82, p = 0.078, FDR corrected for four comparisons). A two-way ANOVA of risk attitude with domain (gain or loss) as the within-subject factor and group as the between-subject factor revealed a significant interaction between domain and group ($F(1,56)$ = 6.29, $p < 0.05$, η^2 = 0.0521). Post-hoc comparisons showed that veterans with PTSD were more averse to risk under gains ($p < 0.01$), but not under losses (p = 0.34), compared with combat controls. Examining this relationship further with a dimensional approach (Fig 3D and Supplementary Fig 2C), we observed a similar effect: PTSD symptom severity was negatively

correlated with risk attitudes in the gain domain (Spearman's ρ with CAPS total = -0.39, $p < 0.01$; Pearson's r with PCL5 = -0.36, $p < 0.01$).

Figure 3. Uncertainty attitudes and PTSD symptom severity

A: Group comparison of ambiguity attitudes in gains and losses between veterans with PTSD and combat controls. B: PTSD symptom severity was negatively correlated with ambiguity attitude in losses. One participant was not included in the analysis due to missing CAPS. C: Group comparison of risk attitudes in gains and losses between veterans with PTSD and combat controls. D: PTSD symptom severity was negatively correlated with risk attitude in gains. One participant was not included in the analysis due to missing CAPS. In A and C, comparisons of each group's attitudes with zero were FDR-corrected across all four comparisons in each uncertainty type. Post-hoc comparisons between groups in A and C are FDR-corrected. Significance level: *, p<0.05; **, p<0.01; ***, p<0.001.

To control for differences in age, income, education and intelligence, we also looked at the relationship between PTSD symptoms and uncertainty attitudes accounting for these factors. We included PTSD severity (CAPS total) together with age, income, education, and intelligence in a linear regression model to explain uncertainty attitudes. For risk attitude in the gain domain, only the effect of CAPS total score was significant (see Methods: Model-based risk and ambiguity attitudes estimation; multi-factor ANOVA by Generalized Linear Model: $F(1, 42) = 8.97$, $p < 0.01$), but not the four demographic factors. Similarly for ambiguity attitude in losses, CAPS total score (multi-factor ANOVA, $F(1, 42) = 5.54$, $p < 0.05$) was the only significant factor.

Because seven of the combat-control veterans in this study sample also participated in the previous behavioral study, we also repeated the analysis excluding these returning participants, to yield a completely independent dataset. The negative relationships between PTSD symptom severity and ambiguity attitude in losses (Spearman's ρ with CAPS total= -0.31, $p < 0.05$, $n = 50$), and between PTSD symptom severity and risk attitude in gains (Spearman's ρ with CAPS total = -0.42, $p < 0.01$, $n = 50$) still held in this independent sample (Supplementary Fig 3B, D).

We also assessed participants' risk-taking attitudes through the Domain-Specific Risk-Taking (DOSPERT) Scale self-report questionnaire, but none of the domains (Ethical, Financial, Health/Safety, Recreational, and Social) was correlated with PTSD symptoms severity measured by CAPS total. Among the other self-report measures, CAPS total was correlated with total score of Behavioral Inhibition Scale (BIS, Spearman's $\rho = 0.37$, $p < 0.01$, $n = 57$), and with total score of Barratt Impulsiveness Scale (BIS11, Spearman's $\rho = 0.47$, p < 0.001, $n = 57$).

PTSD symptom severity is associated with diminished neural response to decision making under uncertainty

To investigate the neural mechanisms of the stronger aversion to uncertainty observed in veterans with PTSD, we first examined the general neural activity during decision making. Because the key process of our task was evaluating the subjective values of the uncertain and sure options, we looked at the neural activity during the 6-second period of options presentation on each trial (see descriptive statistics of participants included in the neural analyses in Table 2). In a whole-brain analysis, we explored the relationship between PTSD symptom severity and the general neural activity during this valuation process (compared to baseline). We found that activity in a vmPFC area was negatively correlated with CAPS total score ($p < 0.001$, cluster-based corrected, Fig 4A), during the second session of the task. This negative relationship was not specific to a particular condition – rather, it was consistent across all four decision contexts (Fig 4B; Pearson's r(risky gains) = -0.50, r(ambiguous gains) = -0.51, r(risky losses) = -0.51, r(ambiguous losses) = -0.40). Veterans with higher overall PTSD symptom severity showed more vmPFC deactivation during valuation of uncertain options generally.

Table 2. Descriptive statistics of demographics and clinical measures of participants reported in the neural results

	PTSD	Control
Number of participants	19	28
Age	35.59 (6.86)	38.62 (10.42)
Kaufman Brief Intelligence Test (KBIT)	104.16 (11.24)	112.97 (13.42)
Clinician Administered PTSD Scale (CAPS)-Total Score	71.42 (15.52)	5.55 (9.40)
CAPS-Re-experiencing	19.84 (7.41)	1.00 (1.95)
CAPS-Avoidance	10.21 (3.52)	0.38 (1.21)
CAPS-Emotional Numbing	17.95 (6.91)	0.72 (2.14)
CAPS-Dysphoric Arousal	14.89 (2.81)	1.79 (3.40)
CAPS-Anxious Arousal	8.53 (2.41)	1.66 (2.93)
PTSD Checklist for DSM-5 (PCL-5)	42.37 (15.46)	9.93 (15.99)
Beck Depression Inventory (BDI)	24.95 (13.40)	5.07 (8.08)
State Anxiety (STAI-1)	46.96 (11.41)	31.69 (8.69)
Trait Anxiety (STAI-2)	45.62 (16.07)	30.12 (10.32)
Dissociative Experience Scale (DES)	47.53 (36.50)	16.17 (20.61)
Combat Exposure Scale (CES)	21.06 (9.70), n = 18	13.14 (8.19)
Childhood Trauma Questionnaire (CTQ)	39.05 (15.53), n = 18	34.64 (9.25)

Mean (standard deviation)

For imaging analysis, 10 participants were excluded from those reported in the behavioral results because of fMRI data quality. Fewer number of participants for several measures was due to incomplete data.

Figure 4. Reduced vmPFC activity during valuation is related to PTSD symptom severity

A: A whole-brain analysis revealed that activity in vmPFC during valuation was negatively correlated with CAPS total score, regardless of decision condition. B: Visualization of this negative correlation between general activity in the vmPFC and CAPS total score in each decision condition.

The subjective value of uncertain gains and losses is encoded in different brain regions

The association between vmPFC activity and PTSD symptom severity in our economic decision-making task is consistent with our hypothesis regarding the involvement of the valuation system in PTSD. We next turned to directly examine the neural correlates of valuation in the task. For each participant, we calculated the subjective value of the lottery presented on each trial, based on the behavioral model (see equation 1 in Methods), using the participant-specific risk and ambiguity attitudes under gains and losses. We then included the subjective values (positive for gain lotteries, negative for loss lotteries) in the GLM, separately for each of the four decision conditions. Replicating previous results (Levy et al. 2010), a whole-brain analysis revealed a strong representation of subjective value of gains in medial brain regions, including vmPFC, ACC, PCC, and bilateral caudates (Fig 5A). Under losses, however, the picture was very different. Neural representations of subjective value were mostly found in lateral regions of the brain, including posterior parietal cortex (PPC), middle frontal cortex, and cerebellum (Fig 5B; for a complete list of regions for all four decision conditions, see Supplementary Table 1). These results suggest differences in basic mechanisms of neural encoding of the subjective value of rewards and punishments, and provide a basis for further investigating the effect of PTSD symptoms on the neural encoding of value.

Figure 5. Neural representation of subjective value of gains and losses among all participants

Whole-brain analyses revealed neural subjective-value signals of all veteran participants (n = 48, PTSD and control veterans combined), under domains of: A. gains, and B. losses. All maps were corrected using cluster-based method controlling family-wise error at 0.05, when thresholded at p < 0.001 at the voxel level.

PTSD symptom severity is associated with altered neural encoding of subjective value of uncertain options

Since choices in the ambiguous-loss condition were related to PTSD symptoms both in a previous behavioral study (Ruderman et al. 2016) and in the current imaging study, we focused our analysis on this decision condition. First, we examined the encoding of subjective value in whole-brain analyses, separately in the PTSD and combat control groups. In controls (Fig 6A, right), activation patterns were similar to those observed in the combined group (Fig 5B), exhibiting positive correlation with subjective value (decreased activity for increased losses) in bilateral frontal, parietal, and cerebellar regions. Conversely, in veterans with PTSD, subjective value of ambiguous losses was represented in a negative manner (increased activity for increased losses) in bilateral temporal regions (Fig 6A left). Our behavioral data demonstrated an additional association between PTSD symptom severity and risk attitude in the gain domain. We therefore also examined the effect of PTSD on subjective-value encoding in this condition. Interestingly, subjective value of risky gains was encoded positively in the brains of veterans with PTSD (increased activity for increased gains) in medial prefrontal cortex (mPFC), ACC, PCC, caudate and temporal regions, but was not strongly encoded in the brains of controls (Fig 6B). For completion, we also investigated the other two conditions: subjective values of ambiguous gains were represented in medial brain regions in both groups (Supplementary Fig 4A); there was very little representation of subjective value of risky losses in either group (Supplementary Fig 4B; for a complete list of subjective-value-encoding regions from the whole-brain analysis, see Supplementary Table 2). Overall, these results demonstrate potential differences in the direction of loss-encoding, and in the magnitude of gain-encoding, between the two groups.

Figure 6. Neural representation of subjective value in PTSD and control

Neural representation of subjective value in veterans with PTSD (left) and combat controls (right), under A: ambiguous losses, and B: risky gains. All maps were corrected using cluster-based method controlling family-wise error at 0.05, when thresholded at p < 0.001 at the voxel level.

To confirm these group differences, we directly contrasted the neural representation of subjective value between veterans with PTSD and combat controls in a whole-brain analysis. Veterans with PTSD showed more negative subjective-value signals for ambiguous losses in left inferior frontal regions (IFG) and bilateral occipital regions (Fig 7A; for statistics of all regions, see Supplementary Table 3). We then used a leave-one-subject-out (LOSO) procedure to define regions around the inferior frontal gyrus (IFG) region and sample activation in an unbiased manner (see Methods: Leave-one-subject-out (LOSO) procedure). The subjective-value signal of ambiguous losses in IFG was negatively correlated with PTSD symptom severity (Fig 7B; Spearman's ρ = -0.35, $p < 0.05$, $n = 48$), such that higher symptom severity was associated with more negative subjective-value signal. Veterans with PTSD showed more positive subjective-value signals for risky gains in right orbitofrontal cortex (OFC) in a whole-brain analysis (Fig 7C), and PTSD symptoms severity was positively correlated with subjective-value signal of risky gains in this OFC region

(Fig 7D; Spearman's $\rho = 0.52$, $p < 0.001$, $n = 48$). For completion, we also looked at the other two conditions. Veterans with PTSD showed more positive encoding of subjective value of ambiguous gains in the thalamus and right cerebellum (Supplementary Fig 4C), and there was no group difference in the subjective-value encoding of risky losses.

Figure 7. Neural representation of subjective value directly contrasting PTSD and control

Whole-brain comparisons of neural subjective-value signals between veterans with PTSD and combat controls, under A: ambiguous losses, and C: risky gains. All maps were corrected using cluster-based method controlling family-wise error at 0.05, when thresholded at p < 0.001 at the voxel level. B: Neural subjective-value representation of ambiguous losses in the left IFG was negatively correlated with PTSD symptom severity. D: Neural subjective-value representation of risky gains in the right OFC was positively correlated with PTSD symptom severity. ROIs in B and D were defined by a leave-one-subject-out (LOSO) approach.

To further probe group and individual differences in value encoding, we examined the subjective-value signals of each group in the classical value areas – the vmPFC and the ventral striatum – as defined in a meta-analysis by Bartra and colleagues (Bartra, McGuire, and Kable 2013). We again focused on the conditions of ambiguous losses and risky gains (Fig 8). In vmPFC, the subjective-value signal of risky gain lotteries was positively correlated with PTSD symptom severity (Fig 8A, Spearman's ρ with CAPS = 0.31, $p < 0.05$). In ventral striatum, subjective-value signal of ambiguous loss lotteries was negatively correlated with PTSD symptom severity (Fig 8B, Spearman's ρ with CAPS = -0.35, $p < 0.05$). PTSD symptom severity was not significantly associated with the subjective-value signal of ambiguous losses in vmPFC (Fig 8A, Spearman's ρ with CAPS = -0.18, $p = 0.22$), or with the subjective-value signal of risky gains in ventral striatum (Fig 8B, Spearman's ρ with CAPS = 0.22, $p = 0.14$; see Supplementary Fig 5A and B for correlations with PCL5). These relationships could also be revealed in the group comparison (Fig 8 C and D). The subjective-value signal of ambiguous losses was more negatively encoded in ventral striatum in veterans with PTSD compared with combat controls (Fig 8D, $t = -2.77$, $p < 0.01$). Conversely, the subjective-value signal of risky gains was marginally more positively encoded in vmPFC in veterans with PTSD than in combat controls (Fig 8C, $t = 1.97$, $p = 0.054$).

The relationships between subjective-value signals and PTSD symptom severity held after controlling for age, income, education and intelligence. The subjective-value signal of ambiguous losses in ventral striatum was affected by CAPS (see Methods: fMRI GLM second-level analysis; multi-factor ANOVA by Generalized Linear Model, $F(1, 33) = 6.01$, $p < 0.05$), and not by the four demographic factors. The subjective-value signal of risky gain lotteries in vmPFC was marginally affected by CAPS (multi-factor ANOVA by Generalized Linear Model: $F(1, 33) = 3.53$, $p = 0.069$), and not by the four demographic factors.

Figure 8. Neural subjective-value signals in external ROIs of vmPFC and ventral striatum were related to PTSD symptom severity

A: In vmPFC, correlations between subjective-value signals of ambiguous losses and risky gains and PTSD symptom severity (CAPS total). B: In ventral striatum, correlations between subjective-value signals of ambiguous losses and risky gains and PTSD symptom severity (CAPS total). C: In vmPFC, group comparison

of neural subjective-value signals between veterans with PTSD and combat controls. D: In ventral striatum, group comparison of neural subjective-value signals between veterans with PTSD and combat controls. E: In ventral striatum, value-encoding of subjective values was observed in combat controls but not in veterans with PTSD; saliency-encoding of subjective values was observed in veterans with PTSD but not in combat controls. In panels C and D, comparisons are post-hoc FDR-corrected after ANOVA within each figure. In panel E, comparisons with zero for each PTSD and Control group were FDR-corrected across four comparisons in two figures. Significance level: *, p<0.05; **, p<0.01; ***, p<0.001.

A shift from value-encoding to saliency-encoding of ambiguous losses in PTSD

Our results so far point to differences in the mechanisms of subjective-value encoding between veterans with PTSD and combat controls. This difference is most notable for ambiguous losses: in combat controls, ambiguous losses were encoded in a positive manner (decreased activity for increased losses) consistent with a monotonic representation of value. Conversely, in the brains of veterans with PTSD, losses were encoded negatively (increased activity for increased losses), consistent with a U-shaped saliency-encoding mechanism. This difference in representation was particularly striking in the ventral striatum (Fig 8D). To directly confirm this group difference, however, we need to examine gains and losses on the same scale. To this end, we constructed a GLM with one predictor for the value of ambiguous gains and losses, and another predictor for the saliency of the same gains and losses. Subjective values of the lotteries were used for the value predictor, and saliency was computed as the absolute value of these subjective values (Fig 8E; see Methods: fMRI GLM first-level analysis). While the ventral striatum in controls significantly encoded value (one-sample t test GLM beta compared with 0, $t(28) = 3.4$, $p < 0.01$), but not saliency ($t(28) = -0.62$, $p = 0.54$), the opposite pattern was observed in veterans with PTSD: activity in the same brain area in the PTSD group encoded saliency ($t(18) = 2.7$, $p < 0.05$), but not value (($t(18) = 0.99$, $p = 0.45$; all p values were FDR corrected for four comparisons). Furthermore, the saliency-encoding patterns were significantly different between veterans with PTSD and combat controls (two-sample t test: $t(39.3) = -2.5$, $p < 0.05$).

Neural activity explains symptoms variation better than behavioral uncertainty attitudes

Having revealed both behavioral and neural alteration related to PTSD, we explored whether PTSD symptom variation could be better explained by neural activation patterns, behavioral uncertainty attitudes, or a combination of both. We constructed three linear models to predict PTSD symptom severity indicated by CAPS total score, using (1) magnitudes of neural responses to the four decision conditions in vmPFC (Fig 4); (2) behavioral risk and ambiguity attitudes under gains and losses; and (3) both neural responses and behavioral attitudes. All models controlled for age and intelligence (KBIT) (see details in Methods: Using behavioral and neural variation to predict PTSD symptom variation). The model including only neural measures best explained the variation of PTSD symptom severity (BIC(neural model) = 132.8, BIC(behavioral model) = 154.7, BIC(full model) = 143.8, Supplementary Fig 6).

Emotional numbing plays the key role in diminished vmPFC general neural activity

So far, we investigated the relationship between PTSD overall symptom severity and valuation under uncertainty. We further looked at whether a specific symptom cluster was the main source of influence, considering the multi-dimensional nature of PTSD. From the vmPFC ROI in which neural activity was negatively correlated with CAPS total score in the whole-brain analysis (Fig 4A), we sampled general neural activity during the valuation phase of the task (GLM beta) from each participant. We then constructed a linear regression model to explain this region's activity with all five clusters of CAPS, including re-experiencing, avoidance, emotional numbing, dysphoric arousal, and anxious arousal, accounting for age and intelligence. Only the emotional numbing cluster significantly contributed to the negative correlation between vmPFC activity and PTSD symptom severity (standardized regression coefficient, *Beta* = -0.72, t = -2.32, $p < 0.05$, Supplementary Fig 7A). Age and intelligence (KBIT) did not significantly influence vmPFC neural activity (standardized regression coefficient, Age: *Beta* = -0.14, t = -1.13, $p = 0.26$; intelligence: *Beta* = -0.044, t = 0.33, $p = 0.75$). Variable selection using exhaustive search also indicated that including only the emotional numbing cluster out of all PTSD symptom clusters best

explained the relationship between vmPFC neural activity and PTSD symptom severities (Supplementary Fig 7B, BIC = 112.8; see details in Methods: fMRI GLM second-level analysis).

Influence of clinical symptoms beyond PTSD

Because our participants showed high levels of comorbidity with other clinical symptoms, especially depression and anxiety (Fig 2B), we also investigated how the behavioral and neural mechanisms of valuation were influenced by symptoms beyond PTSD. We examined the correlation between the first three principal components of all clinical measures (Supplementary Fig1) and the behavioral uncertainty attitudes. Principal component 1 (general affective symptom) was negatively correlated with risk attitude under gains (Pearson's r = -0.35, p < 0.01) and ambiguity attitude under losses (Pearson's r = -0.29, p < 0.05), consistent with the effect of the overall PTSD severity indicated by CAPS total score (Fig 3). We did not find any relationship between uncertainty attitudes and the second (fear learning-updating) or the third (trauma severity) principal components.

We also examined potential relationships between subjective-value signals and the three principal components. In vmPFC, the first component (general affective symptom) was positively correlated with encoding of subjective value of risky gains (Pearson's r = 0.30, n = 47, p < 0.05), consistent with the effect of PTSD symptom severity (CAPS total score). The third component (trauma severity) was negatively correlated with encoding of subjective value of ambiguous losses (Pearson's r = -0.29, n = 47, p < 0.05), in the same direction as the correlation between subjective value of ambiguous losses and PTSD symptom severity. In ventral striatum, no correlation survived our statistical thresholds (see all correlations in Supplementary Fig 5C).

Discussion

In this study, we explored the neural basis of valuing uncertain monetary rewards and punishments, in veterans exposed to combat trauma with a wide range of PTSD symptoms. Behaviorally, symptom severity was associated with increased aversion to ambiguous losses, and increased aversion to risky gains. These two conditions were also the ones in which PTSD symptom severity influenced the neural representations of subjective value. Two main effects were observed: first, while ambiguous losses were encoded in a positive manner in the brains of controls, individuals with PTSD showed, on average, negative encoding of ambiguous losses in several brain areas (Fig 6A, Fig 7A, Fig 8D), consistent with a saliency representation. Second, increased PTSD symptoms were associated with stronger representation of subjective value of risky gains in value-related areas, including the vmPFC, PCC and OFC. In fact, the average activation pattern for subjective value of risky gains in the PTSD group (Fig. 6B left) was very similar to activation patterns observed in the general population (Levy et al. 2010; Bartra, McGuire, and Kable 2013), whereas controls did not show strong value-encoding signals (Fig 6B right). An intriguing possibility is that the altered processing of ambiguous losses is a marker for vulnerability to PTSD, whereas the altered processing of risky gains is a marker of resiliency to PTSD. Future research, and in particular longitudinal studies that compare individuals exposed to trauma to those who never experienced trauma, are needed to explore this possibility.

Using behavioral economics to identify markers of psychopathology

Our results add to a growing body of research, demonstrating the utility of behavioral economics in studying psychopathology (Paulus and Yu 2012; Pushkarskaya et al. 2015; Buckholtz et al. 2017; Konova et al. 2019; Mukherjee et al. 2020). Replicating the previous behavioral study (Ruderman et al. 2016), we confirmed the association between higher PTSD symptom severity and greater ambiguity aversion in losses, in an independent combat veteran sample. We also identified greater aversion to risk in gains in veterans with PTSD, likely due to a task design with increased range and variance of monetary outcomes, that

provided higher sensitivity for capturing true uncertainty attitudes. Our neural measure allowed us to also quantify individual and group differences in neural sensitivity to rewards and punishments. Previous studies have shown alterations in the neural processing of aversive outcomes in individuals with PTSD, in various brain areas, including several medial and lateral prefrontal regions. Many of these studies, however, used fear and trauma-related stimuli (Hayes, Hayes, and Mikedis 2012). Here we show that activation in the same brain areas is affected by PTSD symptoms even in an economic decision task, completely unrelated to the trauma. This raises the possibility of developing diagnostic methods in the domain of decision making under uncertainty, which do not require patients to recall the traumatic experience. Several previous studies have also reported altered reward processing in PTSD (Nawijn et al. 2015), including reduced expectation of uncertain monetary outcomes (Hopper et al. 2008; May and Wisco 2019) and decreased differentiation between monetary gains and losses in the striatum (Elman et al. 2009). Our experimental approach allowed us to estimate individual uncertainty attitudes during active decision making under four unique contexts. We applied a well-established computational model to infer these behavioral individual differences from the observed choice behavior (rather than estimating them through self-reports) and used the individual differences in the analysis of the neural data. Interestingly, participants' self-reported risk-taking on the DOSPERT questionnaire was not strongly correlated with their PTSD symptom severity, suggesting that our method for estimating uncertainty attitudes through a behavioral task may be more sensitive for capturing subtle differences associated with clinical symptoms. An intriguing question remains to be answered is what contributes to the context-specific differences associated with PTSD symptom severity. We found higher behavioral aversion only to ambiguous losses and risky gains in PTSD, and the most striking neural difference of subjective-value encoding was revealed in the context of ambiguous losses. Compared with risky outcomes, of which both outcome magnitude and probability are known, ambiguous outcomes lack the exact information of outcome probability. This additional level of uncertainty may be more relevant to the nature of battlefield, making negative ambiguous outcomes more relatable to combat exposure. Our task design separating decision contexts enabled us to pinpoint specific cognitive processes affected by PTSD in combat veterans, but replication in larger samples is needed to confirm the results.

Neural processing of rewards and punishments is associated with PTSD symptoms

By including both monetary gains and losses in the task design, our results illuminate the basic mechanisms of neural processing of rewards and punishments in two ways. First, our data suggest that neural processing of positive and negative outcomes may be realized through two distinct circuits in our sample of combat veterans, regardless of clinical symptoms. While rewards are processed primarily in medial parts of the brain, punishments are processed in lateral regions. This medial-lateral segregation and, more generally, the separate neural representation of rewards and punishments have been reported in previous fMRI studies (O'Doherty et al. 2001; Yacubian et al. 2006; Kim et al. 2014). Single unit recordings from non-human primates also indicate different populations of vmPFC neurons (Monosov and Hikosaka 2012) and dopaminergic neurons (Fiorillo 2013) responding to primary appetitive and aversive outcomes. Other studies, however, reported overlap between representations of gains and losses, supporting the common currency hypothesis (Tom et al. 2007; Fujiwara et al. 2009; Kahnt et al. 2014). Some of the discrepancies between the findings may stem from differences in task design. In particular, simultaneous processing of gains and losses, for example when considering mixed lotteries (Tom et al 2007), may lead to overlapping representations of gains and losses, whereas presentation of gains and losses in separate blocks, like in our design, may unravel differences in their neural representation. The separate involvement of medial and lateral brain regions in the valuation of gains and losses also points to an intriguing link to the default mode network which includes the mPFC and PCC, and the executive function network which includes the dlPFC and PPC. Also referred to as the task-negative and task-positive networks, these two networks are thought to be involved in the self-referential thinking and executive control processes respectively, as well as in reward processing (Lesage and Stein 2016). Future studies could investigate whether valuation of rewards and punishments is also reflected in the dynamics of connectivity in these two networks, and if the difference between processing rewards and punishments is related to shifting attention inward and outward. In individuals with PTSD, studies have found reduced overall DMN intrinsic connectivity during resting state, as well as finer abnormalities within the network including prefrontal dysconnectivity and increased

cohesiveness in posterior DMN (Akiki et al. 2018). Our study identified reduced vmPFC activity during valuation associated with symptom of emotional numbing. Further research into DMN dynamics during value-based decision-making task would be helpful to better understand the role of vmPFC in PTSD symptom development.

Second, we identified a shift from value-encoding to saliency-encoding in the brains of individuals who developed PTSD following trauma exposure (Fig 9). This shift could potentially imply an attention or arousal signal, that leads to avoidance of aversive outcomes like uncertain monetary gains or losses. Several previous studies examined the neural processing of value and saliency and revealed both distinct and overlapping regions for each type of encoding. Value signals were found in ventral striatum, parietal cortex, OFC, rostral ACC, and saliency signals were found in ventral striatum, rostral ACC, dorsal ACC, anterior insula by both univariate and multivariate analyses (O'Doherty et al. 2001; Yacubian et al. 2006; Tom et al. 2007; Fujiwara et al. 2009; Litt et al. 2011; Kim et al. 2014; Kahnt et al. 2014; Zhang et al. 2017). To our knowledge, our results are the first to recognize the influence of psychiatric symptoms in humans on the value/saliency-encoding pattern. Interestingly, recent research in mice shows a similar flip in representation, where acute stress transforms reward responses in the lateral habenula into punishment responses (Shabel et al. 2019). Neurons in the nucleus accumbens of rats can also flexibly shift their preferences between rewards and punishments, based on the emotional environment (Reynolds and Berridge 2008), suggesting that what we observe here may reflect a stress coping mechanism.

Figure 9. A shift from value- to saliency- encoding of subjective value in veterans with PTSD, compared with controls

In lateral brain areas, neural activity for increasing potential losses led to increased activation in the brains of veterans with PTSD (consistent with saliency encoding). Conversely, in the brains of control, increased losses were associated with reduced activity.

PTSD is highly comorbid with symptoms of depression and anxiety. Through PCA, we were able to disentangle three main symptom components, and showed that the component of general affective symptoms was likely the main source of influence (Supplementary Fig 5). In addition, we also found that the component of trauma symptoms was related to the neural representation of subjective value of ambiguous losses in vmPFC, raising the possibility that trauma exposure additionally influences sensitivity to aversive monetary outcomes, independent from general affective symptoms. Trauma symptoms were assessed through combat exposure and childhood trauma in our sample of veterans. Future research could investigate more systematically how other types of trauma exposure could additionally influence the neural processing of valuation of uncertain outcomes.

One concern in our investigation of neural representation of value is that the range of subjective values is lower in the group of veterans of PTSD because of their higher aversion to uncertainty, which could influence the sensitivity of the neural response to value differences. It should be noted, however, that our main conclusion is based on a difference in the direction of correlation (negative vs. positive), rather than a difference in the magnitude of slope of the correlation (Fig 6A, 7A, and 8D). This represents a substantial difference in the shape of subjective-value encoding and would not be affected by group difference in the range of subjective values.

Neural markers of vulnerability and resilience to PTSD

Previous studies of PTSD often focused on the neural processing of fear and trauma, and identified both functional and structural abnormalities in amygdala, hippocampus, and vmPFC (Hayes, Hayes, and Mikedis 2012; Admon, Milad, and Hendler 2013; Wolf and Herringa 2015; Rabellino et al. 2016; Homan et al. 2019). Other studies have looked into more general cognitive processes and found blunted neural activation to monetary rewards (Sailer et al. 2008; Elman et al. 2009). In our study, using a more nuanced computational approach, PTSD symptoms were associated with increased neural sensitivity to rewards and opposite direction of sensitivity to punishments. While the sensitivity to rewards may seem at odds with the previous studies, it should be noted that in those studies reward signals were defined as the difference in activation to gains and losses. A weaker contrast in individuals with PTSD could stem from a weaker reward signal, but also from a stronger punishment signal, consistent with a U-shaped saliency representation, as we report here, in which both highly salient positive and highly salient negative outcomes elicit similar magnitude of neural activation. With this being said, in the reward domain, we do find evidence of surprisingly stronger value sensitivity in PTSD (Fig 6 and 7C). Note that reduced activation to rewards in individuals with PTSD was previously observed in comparison to controls who *were not exposed to trauma* (Sailer et al. 2008). In our study, veterans with PTSD exhibited neural patterns for potential rewards which were similar to what has been observed in the general population (Bartra, McGuire, and

Kable 2013; Levy et al. 2010). Combat controls, who were exposed to trauma, but did not develop PTSD, were the ones who differed from the general population, suggesting that they have exhibited a resiliency marker. Other important methodological details may contribute to the difference in results between the studies, including a focus on outcome delivery, rather than decision value, female participants, and mixed types of trauma. These differences suggest interesting directions for future research. Of particular interest is the interaction between uncertainty and combat trauma, as uncertainty is a central component of the battlefield experience. Comparing behavior and neural mechanisms in individuals who experienced combat trauma and those who did not, will help to shed light on the potential vulnerability and resiliency markers proposed here.

In line with the NIMH Research Domain Criteria (RDoC), we did not exclude veterans with history of substance abuse, to allow for a diverse representative sample of trauma exposed symptomatology. We controlled for substance abuse by conducting urine test and breathalyzer for anyone with substance abuse history or if we suspected any intoxication, and excluded those with positive results. The severity for substance abuse history in our sample was low and did not vary too much as measured by the Addiction Severity Index (ASI-alcohol: median = 0.089, range = [0, 1.47]; ASI-drug: median = 0, range = [0, 0.092]). Future research could better control for substance abuse history and medication, and potentially look into the pharmacological effect involving the dopamine and serotonin systems, which are crucial for value-based decision making (Castrellon et al. 2019; Macoveanu 2014).

It should also be noted that our study could not establish causal relationship between decision making under uncertainty and the development of PTSD symptoms. Heightened aversion to uncertainty could possibly predispose individuals to developing PTSD symptoms, and on the other hand, acquiring PTSD symptoms could result in altered uncertainty attitudes. There is some evidence, however, that risk attitude is correlated with relatively stable biomarkers including structural volume of right posterior parietal cortex (Gilaie-Dotan et al. 2014), structural and functional connectivity of the amygdala (Jung et al. 2018) and genetic variations (Zyphur et al. 2009). These pieces of evidence might indicate that risk attitude is a personal trait, raising

the possibility of its predisposing effect on the development of PTSD symptoms. Less evidence exists for biomarkers of ambiguity attitude, although there is some evidence for a genetic association among females (Chew, Ebstein, and Zhong 2012). Further longitudinal studies comparing veterans pre- and post- military service may disentangle the role of pre-existing uncertainty attitudes on the development of PTSD from the subsequent impact of PTSD symptomatology on uncertainty attitudes.

Variations in decision making under uncertainty, and especially under ambiguity, have also been reported in other psychiatric disorders, including higher ambiguity aversion and choice inconsistency in individuals with Obsessive Compulsive Disorder (Pushkarskaya et al. 2015), and decreased ambiguity aversion in individuals with antisocial personality disorder (Buckholtz et al. 2017). Interestingly, a recent longitudinal study demonstrated transient increases in tolerance to ambiguity before relapses in opioid users undergoing treatment (Konova et al. 2019). Overall, these efforts to study psychiatric disorders using behavioral economics approaches could collectively lead to both early identification of behavioral and biological risk factors for symptom development, and more effective treatment.

Methods

Participants

68 male veterans (ages: 23.6-74.6; mean ± standard deviation: 39.4 ± 11.5), who had been deployed and exposed to combat, were recruited. Participants either had current diagnoses of PTSD at the time of the study or were never diagnosed with PTSD (controls). Due to the small proportion of female combat veterans (15% of female in Army 2019, statistics from Department of Defense), we only included male participants. PTSD diagnosis was based on the Clinician Administered PTSD Scale for DSM-4 (CAPS) (Blake et al. 1995). Data from 10 participants were excluded due to a large number of missing responses or low correct response rate in catch trials in the main task (see Task design and Manipulation check), resulting in 58 participants (ages: 23.6-67.0; mean ± standard deviation: 37.3 ± 8.9) whose behavioral data were reported. Among these 58 participants, CAPS was missing for one PTSD participant, so we only included him in the

group comparison but not the dimensional analysis. Imaging data from 10 additional participants were excluded due to excessive movement in the scanner, or because their data were collected using different scanning parameters (5 participants), resulting in 48 participants (ages: 23.6-67.0; mean ± standard deviation: 37.4 ± 9.2), whose neural results were reported. Full characteristics of the sample included in the analysis were reported in Table 1 (behavioral) and Table 2 (neural). Seven participants also took part in a previous behavioral study (Ruderman et al. 2016) using a similar paradigm, and these participants were included in both behavioral and imaging analyses. Two of these participants were diagnosed as PTSD in the previous behavioral study but were grouped into combat controls who never developed PTSD in this imaging study. This discrepancy could be due to the inaccuracy of subjective report at the time of the behavioral study, or of the imaging study, and could not be resolved. Behavioral results excluding these seven recurring participants are also reported in Supplementary Fig 3.

The study was approved by the Yale University Human Investigating Committee and the Human Subjects Subcommittee of the VA Connecticut Healthcare System, and compliance with all relevant ethical regulations was ensured throughout the study. All participants gave informed consent and were compensated with $100 for their participation, plus a variable bonus ($0-$240) based on choices they made in the task (see Experimental design).

Experimental design

The study was composed of three separate visits on three different days (Fig 1A). On the first day (Screening Day), recruited participants went through clinical interviews for screening. Eligible participants continued to two fMRI sessions, on two separate days. In these sessions, participants performed a task of decision making under uncertainty. They made a series of decisions between a sure monetary outcome and an uncertain monetary outcome with either known (risky) or unknown (ambiguous) outcome probability, in scenarios of both gaining and losing money (Fig 1B, see details in Task design below). On each trial, participants viewed the two options side-by-side for a fixed duration of 6 seconds, and then made a choice

(Fig 1D). To prevent learning, the outcome of the chosen option was not presented during the scan. At the end of the experiment, one randomly selected trial was realized. The scans were conducted over two days in order to limit the scanning time in each visit; task designs were identical for Day1 and Day2. Participants were introduced to the task at the beginning on the first scanning day (Day1) and were reminded of the study on the second scanning day (Day2). Additional questionnaires and intelligence tests were administered at the end of Day2.

Clinical assessment and exclusion criteria

Participants were recruited through flyers and were screened by clinicians at West Haven Veterans Affairs hospital. PTSD sympotms and diagnoses were determined by the Structured Clinical Interview for DSM-4 (SCID) (First et al. 1997) and the Clinician Administered PTSD Scale (CAPS) (Blake et al. 1995). We also collected the following measurements: PTSD Checklist for DSM-5 (PCL-5) (Weathers et al. 2013), Beck's Depression Inventory (BDI) (Beck et al. 1961), State-Trait Anxiety Inventory (STAI) (Spielberger and Gorsuch 1983), Dissociative Experiences Scale (DES) (E. M. Bernstein and Putnam 1986), Combat Exposure Scale (CES) (Keane et al. 1989), and Childhood Trauma Questionnaire (CTQ) (D. P. Bernstein et al. 2003). Participants with the following conditions were excluded after screening: psychosis, bipolar disorder, traumatic brain injury, neurologic disorder, learning disability, and ADHD. We did not exclude veterans with history of substance abuse, to allow for a diverse representative sample of trauma-exposed symptomatology. However, we conducted urine test and breathalyzer for anyone with substance abuse history, or if we suspected any intoxication, and excluded those with positive results. Participants also completed other questionnaires including demographic information, Behavioral Avoidance/Inhibition (BIS/BAS) Scales (Carver and White 1994), the Barratt Impulsiveness Scale (BIS-11) (Patton, Stanford, and Barratt 1995), and Doman-Specific Risk-Taking (DOSPERT) Scale (Blais and Weber 2006). Kaufman Brief Intelligence Test (KBIT) (Kaufman 1990) was administered after scanning as a measure of non-verbal intelligence.

Decision making under risk and ambiguity
Task design

The experimental design was based on a previous neuroimaging study (Grubb et al. 2016) and similar to the design of a previous behavioral study in combat veterans (Ruderman et al. 2016).

The experiment consisted of choices about risky and ambiguous gains and losses. On gain trials, participants made choices between a fixed monetary gain ($5) and a lottery with chance of a monetary gain but also chance of no gain ($0) (Fig 1B, left). Lottery outcome probability was represented by an image of a rectangle with blue and red areas. On each trial, one color was associated with a monetary gain, and the other color was associated with the null outcome ($0). The size of each colored area represented the probability of getting the outcome associated with it. In half of the trials, outcome probability was fully known (25%, 50% or 75%; 'Risk', Fig 1C left). In the other half of the trials, probability was only partially known. This was achieved by covering the middle part of the colored image with a grey bar ('Ambiguity', Fig 1C right). On different trials the bar covered 24%, 50% or 74% of the image, creating three levels of ambiguity. For example, the lower left example in Fig 1B represents a lottery with a chance between 25% and 75% of winning $27. Each of the risk and ambiguity levels corresponded to an actual physical bag with a total number of 100 red and blue chips inside. For risky bags (corresponding to the risk level images), the exact numbers of red and blue chips were equal to the numbers shown on the images. For ambiguous bags (corresponding to the ambiguity level images), the exact numbers of red and blue chips were within the range shown on the image. Participants were shown these bags during study introduction and were informed that the lottery images they saw during the task corresponded to these bags. The bags would be used to actualize participants' choice, and they were free to inspect the contents of the bags after finishing the study. The potential monetary gain of the lottery varied within a wide range ($5, 6, 7, 8, 10, 12, 14, 16, 19, 23, 27, 31, 37, 44, 52, 61, 73, 86, 101, and 120). The non-zero monetary outcome was randomly related to red or blue, so color was independent from outcome preference.

Loss trials were similar to gain trials, except that participants chose between losing $5 for sure, and playing a lottery with chance of losing money, but also chance of not losing any money ($0) (Fig 1B, right). Same as gain trials, half of the loss lotteries were risky and half of them were ambiguous, and the outcome probability was presented in the same way. The potential monetary loss of the lottery varied within the same range but of negative monetary outcomes (–$5, 6, 7, 8, 10, 12, 14, 16, 19, 23, 27, 31, 37, 44, 52, 61, 73, 86, 101, and 120).

Fig 1D illustrates the timing of each trial. Participants had 6 seconds to look at the options and consider their decision. A green circle then appeared in the middle of the screen, cueing participants to make a choice within 3.5 seconds. If they did not register a response, the response would be counted as missing. A feedback image of two side-by-side squares was shown for 0.5 second immediately following the button press to confirm the choice, with the yellow square indicating the side participants chose. The feedback was followed by a jittered inter-trial-interval (ITI) of a white circle, the duration of which was among 4, 6, 8 seconds plus the remainder of the response time limit (3.5s – reaction time (RT)). To avoid learning, chosen lotteries were not played during the scan. In both the gain and the loss domains, all risk and ambiguity levels were paired with all amount levels, resulting in 120 unique gain trials ((3 + 3) × 20) and 120 unique loss trials. Each trial type was presented once. All trials were grouped into 8 blocks of 30 trials each. Each block contained either only gain trials or loss trials, but risky and ambiguous trials were mixed in a random order. This resulted in 4 gain blocks and 4 loss blocks. Due to the length of the experiment, these blocks were divided into two scanning sessions on two separate days. The interval between the two scanning sessions was on average 12.6 days.

Manipulation check

To verify that participants understood the task, and that they aimed to maximize earnings and minimize losses, we included 12 trials in which the potential lottery outcome was identical to the certain amount (±$5). In these trials, one option is clearly better than the other (e.g. a certain gain of $5 should be preferred

over a 50% chance of gaining $5). Data from participants who chose the inferior option on more than 50% of these trials were excluded.

Task administration

On Day1, participants were introduced to the task, and were required to correctly respond to several questions to make sure they understand the task and the lottery presentation. They also practiced 16 trials before the actual task. After the introduction, each participant was endowed with $120 (the maximal possible loss), and then went through two gain and two loss blocks, whose order was counterbalanced across participants (either Gain-Gain-Loss-Loss or Loss-Loss-Gain-Gain). Each block contained 30 trials together with an additional trial in the beginning (a choice between +/-$5 and a lottery offering +/-$4) to capture the initial burst of activity. The added one trial was excluded from analysis.

On Day2, participants were first reminded of the task, and then went through another four blocks of choices (two gain and two loss), in an opposite order to what they had on Day1. Following the scanning, one trial out of the 240 trials (including both gains and losses) was randomly selected and realized. If the participant chose the sure option, $5 were added to or subtracted from the $120 endowment. If the participant chose the lottery, he\she would play the lottery by pulling a chip out of the physical bag corresponding to the lottery image, and the outcome related to the color of the pulled-out chip would be added to or subtracted from the $120. At the end of the second session, KBIT, demographic questionnaires, BIS/BAS, and BIS-11 were collected.

MRI scans

MRI data were collected with two scanners (due to scanner upgrade) at the Yale Magnetic Resonance Research Center: Siemens 3T Trio (37 participants, 29 reported in imaging results) and 3T Prisma (31 participants, 19 reported in imaging results), using a 32-channel receiver array head coil. High resolution structural images were acquired by Magnetization-Prepared Rapid Gradient-Echo (MPRAGE) imaging (TR

= 2.5 s, TE = 2.77 ms, TI = 1100 ms, flip angle = 7°, 176 sagittal slices, voxel size = 1 ×1 × 1 mm, 256 ×

256 matrix in a 256 mm field-of- view, or FOV). Functional MRI scans were acquired while the participants

were performing the choice task, using a multi-band Echo-planar Imaging (EPI) sequence (TR= 1000 ms,

TE= 30ms, flip angle=60°, voxel size = 2 × 2× 2 mm, 60 2 mm-thick slices, in-plane resolution = 2 × 2

mm, FOV= 220mm).

Analysis

Analysis of clinical symptoms and trauma-related measures

We used the 5-factor model of PTSD (Harpaz-Rotem et al. 2014) to assess multidimensional PTSD

symptoms, including re-experiencing, avoidance, emotional numbing, dysphoric arousal, and anxious

arousal. We calculated both the overall symptom severity, and the 5 factors' symptom severities by breaking

down the questionnaire items. To account for comorbidities, we conducted principal component analysis

(PCA) on all clinical and trauma-exposure measurements, including the 5 factors of CAPS, Beck's

Depression Inventory (BDI), state and trait anxiety indexes separately from State-Trait Anxiety Inventory

(STAI), Dissociative Experiences Scale (DES), Combat Exposure Scale (CES), and Childhood Trauma

Questionnaire (CTQ).

Model-based risk and ambiguity attitudes estimation

We fitted each participant's choice data into a behavioral economics model that was used in previous studies

(Levy et al. 2010; Ruderman et al. 2016). The model fitting was conducted separately for gain and loss

trials. The model separates the decision process into two steps: valuation and choice. In the valuation step,

the subjective value (SV) of each option is modelled by equation (1),

$$SV = \left[P - \beta\left(\frac{A}{2}\right)\right] \times V^{\alpha} \tag{1}$$

where P is the outcome probability (0.25, 0.50, or 0.75 for risky lotteries, 0.5 for ambiguous lotteries, and 1 for the certain option); A is the ambiguity level (0.24, 0.5, or 0.74 for ambiguous lotteries; 0 for risky lotteries and the certain amount); V is the non-zero outcome magnitude of the lottery or the amount of money of the certain option. For choices in the loss domain, amounts are entered with a positive sign. Risk attitude was modeled by discounting the objective outcome magnitude by a participant-specific parameter, α. In the gain domain, a participant is risk averse when $\alpha < 1$, and is risk seeking when $\alpha > 1$. Because we fitted the choice data in the loss domain using positive outcome magnitudes, the participant is risk averse when $\alpha > 1$, and is risk seeking when $\alpha < 1$. Ambiguity attitude was modeled by discounting the lottery probability linearly by the ambiguity level, weighted by a second participant-specific parameter, β. A participant is averse to ambiguity when $\beta > 0$, and is ambiguity seeking when $\beta < 0$ in the gain domain. In the loss domain, participant is averse to ambiguity when $\beta < 0$, and ambiguity seeking when $\beta > 0$.

The choice process is modeled by a standard soft-max function (equation 2),

$$P_V = \frac{1}{1+e^{\gamma(SV_L - SV_C)}} \tag{2}$$

where P_V is the probability of choosing the lottery option, SV_C and SV_L are the subjective values of the certain option and the lottery respectively, calculated by equation (1); γ is a participant-specific noise parameter.

The model-fitting was conducted in MATLAB (Version R2018b) through maximum likelihood. We primarily used MATLAB function fminunc to minimize the negative log-likelihood function, and switched to using fminsearch if it failed to converge. Under our task design, we could detect risk parameters (α) in the range of [0.0905, 7.6036], and ambiguity parameters β in the range of [-4.0303, 4.1667]. These constraints were calculated based on the range of uncertainty levels and monetary magnitudes in our task

design, same as the procedure used in previous studies (Ruderman et al. 2016). The lower boundary of α was determined by equating the subjective value of the best lottery (75% chance of $120) with the subjective value of the certain option ($5), using equation (1). Similarly, to determine the upper boundary of α, we equated the subjective values of the worst lottery (25% chance of $6) and the certain option ($5). Because the choices of ambiguous lotteries also depend on the risk attitude α, the boundary values of β also depend on α. So we simply calculated all possible β's by equating the subjective values of each ambiguous lottery and the certain option ($5), using the boundary values of α. The boundary values of β were then determined by the minimum and maximum of all calculated β's. Even without constraining α and β during the modeling fitting procedure, the fitted results of all participants included in behavioral and imaging analyses were within this range. We fitted each participant's choices combining data from two sessions and obtained four attitudes: risk attitudes for gains and losses, ambiguity attitudes for gains and losses. For consistency, we transformed all attitudes in the following way such that negative values indicate aversion and positive values indicate seeking: risky gains: $\alpha - 1$, risky losses: $1 - \alpha$, ambiguous gains: $-\beta$, ambiguous losses: β. Since participants performed the task on two separate sessions, we also fitted each session's choice data separately. These fitted parameters from separate sessions were used to calculate trial-wise subjective values of the lotteries for GLM neural analysis, because they could capture the subjective values more accurately for searching neural activity change induced by variations of subjective values.

We then analyzed the fitted risk and ambiguity attitudes both in group comparisons between veterans with PTSD and combat controls, and through a dimensional approach by looking at their correlation with clinical symptoms and other continuous measurements. We also accounted for the potential influence of demographic factors (age, income, education, and intelligence) on uncertainty attitudes, by conducting multi-factor ANOVAs through a Generalized Linear Model:

Uncertainty attitude in one decision condition (e.g. Risk attitude in gains) ~ CAPS total + age + income (categorical) + education (categorical) + intelligence

In the data we collected, age and intelligence were continuous variables, income was a categorical variable with 10 possible levels, and education was a categorical variable with 6 possible levels. All continuous variables were standardized before fitting the linear model.

MRI data preprocessing

Preprocessing of MRI data were conducted in BrainVoyager (Version 20.2.0.3065). Anatomical images were normalized to the standard brain template in Talairach space for each participant. Preprocessing of functional data included motion correction, slice scan time correction (cubic spline interpolation), temporal filtering (high-pass frequency-space filter with cut-off cycle of 3), spatial smoothing (Gaussian filter with 8mm full-width at half-maximum), co-registration with high-resolution standardized anatomical data, and normalization to Talairach space. Scan data with movement of over 2 mm in any direction were excluded from analysis.

fMRI GLM first-level analysis

Analysis of fMRI data were conducted in the Neuroelf toolbox (Version 1.0, https://neuroelf.net/) through MATLAB (Version R2018b) for the purpose of fitting generalized linear models (GLM), extracting regions-of-interest (ROI) data, and visualizing brain statistical maps. Further statistical analyses and visualization were conducted in R (Version 3.5.1) (R Core Team 2018) with packages ez (Lawrence 2013), psych (Revelle 2018), nlme (Pinheiro et al. 2013), emmeans (Russell 2018), ggplot2 (Wickham 2016), and PerformanceAnalytics (Peterson and Carl 2019). We investigated the neural response under different decision conditions by fitting pre-processed functional signals with generalized linear models (GLM). The pre-processed fMRI signal time course was first converted to percent signal change within each scanning block, and activity of each voxel was modeled by GLM predictors convolved with a standard double-gamma hemodynamic response function.

In the first GLM, we looked at the general activity during decision making, by including four binary predictors for all four decision conditions: ambiguous gains, risky gains, ambiguous losses, and risky losses. Each binary predictor was modeled as a box-car function, with the duration of choice display (6TR). We modeled choice response of all trials by another binary predictor with the duration of 1TR at the time of button press, and missing responses were not modeled. We also included nuisance predictors of 6 motion correction parameters (translation and rotation in the x, y, and z directions) in the GLM to account for influence of head motions on the neural activity.

In a second GLM, we modeled the neural response to the variation of trial-wise subjective value of the lottery by including the subjective value as a parametric modulator for each of the four decision-condition binary predictors. Subjective value of the lottery in each trial was calculated uniquely for each participant by equation (1) (See section Model-based risk and ambiguity attitudes estimation), by taking the fitted α and β for each participant under each domain of either gains or losses. Because we fitted the choice data in the loss domain by inputting the positive outcome value, we flipped the sign of the calculated subjective value back in the loss domain. We calculated the subjective values taking α's and β's fitted from the two sessions separately, because it would make the estimate of neural response to subjective value variation more accurate. Subjective values were normalized within each scanning block before GLM fitting, so that the estimated effect reflected each participant's neural response to the variation of subjective value, rather than to its absolute magnitude. Predictor of choice response and nuisance predictors of motion correction were included as in the first GLM.

In the third and fourth GLMs, we aimed to further investigate the shape of the neural representation of subjective values. In both GLMs, we compared gains and losses on the same scale, and only separated trials by uncertainty types. Thus, we included two binary predictors, ambiguous trials and risky trials, in both GLMs, and modeled them as box-car functions with a duration of choice display (6TR). In the third GLM, we included the subjective value itself as a parametric modulator to accompany each binary predictor, to look at the monotonic value-encoding of subjective values. In the third GLM, we included the absolute

value of subjective value as a parametric modulator to accompany each binary predictor, to look at the U-shaped saliency-encoding of subjective values. The predictor of choice response and nuisance predictors of motion correction were included as in the first GLM.

fMRI GLM second-level analysis

After individual GLM fitting, random-effect group analysis was conducted to test whether the mean effect of interest was significantly different from zero across participants, or significantly different between groups, by contrasting veterans with PTSD and combat controls. We also took a dimensional approach to test whether the predictor effects were related to the severity of PTSD and other clinical symptoms.. The tests were conducted both in a whole-brain search and in ROIs.

In whole-brain analyses, all statistical maps were thresholded at $p < 0.001$ per voxel, and corrected for multiple comparisons using cluster-extent correction methods through Alphasim by AFNI (Cox 1996) to control family-wise error (FWE) rate at 0.05.

In ROI analyses, investigating the general neural activity during decision making based on the first GLM, we chose the region in vmPFC whose activity showed negative correlation with CAPS total in the whole-brain analysis. We further investigated which factor of the PTSD symptoms drove this negative relationship, by fitting a linear model including all five symptoms based on the CAPS measurements together with age and intelligence:

Averaged GLM beta over all four decision conditions ~ re-experiencing + avoidance + emotional numbing + dysphoric arousal + anxious arousal + age + intelligence

We then conducted variables subset selection to identify which symptom cluster(s) best influenced vmPFC neural activity, using exhaustive search through the package "leaps" (Lumley and Miller 2009) in R. We compared regression models including all possible combinations of variables for each given number of predictors of this linear model (ranging from including only one predictor to including all seven predictors),

and selected the best model with the lowest Bayesian Information Criterion (BIC). The best linear model identified by this exhaustive approach could identify both the best number of symptom clusters to include, as well as which symptom cluster(s).

In ROI analyses, investigating subjective value representations based on the second, third and fourth GLMs, we chose two brain regions, vmPFC and ventral striatum, based on the meta-analysis by Bartra et al. (Bartra, McGuire, and Kable 2013), which identified these areas as encoding the value of different reward categories. To control for the potential influence of demographic factors on neural activity, we conducted multi-factor ANOVAs through a Generalized Linear Model that included the PTSD total symptom severity and demographic factors, to explain the neural subjective value signal, similar to the behavioral analysis of uncertainty attitudes:

Neural subjective value signal of a type of lottery (e.g. risk gain lottery) ~ CAPS total + age + income

(categorical) + education (categorical) + intelligence

All continuous variables were standardized before fitting the linear models.

Leave-one-subject-out (LOSO) procedure

After identifying regions from the whole-brain analysis, in which the neural representation of subjective values was influenced by PTSD symptom severity, we took a leave-one-subject-out (LOSO) approach to define these ROIs in an un-biased way for each participant. For each left-out participant, we defined an ROI from a whole-brain analysis using data from all other participants, so this ROI definition was not influenced by the left-out participant. We then sampled neural signals of the left-out participant's data from this ROI. We repeated the process for all participants.

Using behavioral and neural measures to predict PTSD symptom variation

To investigate whether symptom variation could be better explained by behavioral uncertainty attitudes or by neural activity during decision making, we constructed an additional model. This model included the betas of each decision condition (ambiguous losses, risky losses, ambiguous gains, and risky gains) in the first GLM, sampled from the vmPFC area identified in the whole-brain analysis:

Neural model: CAPS total ~ neural activity under risky gains + neural activity under ambiguous gains + neural activity under risky losses+ neural activity under ambiguous losses + age + intelligence

Similarly, we constructed a linear model using behavioral uncertainty attitudes in the four decision conditions to explain PTSD symptom severity:

Behavioral model: CAPS total ~ risk attitude in gains + ambiguity attitude in gains + risk attitude in losses + ambiguity attitude in losses + age + intelligence

We also constructed a full model including both neural and behavioral measures:

Full model: CAPS total ~ risk attitude in gains + ambiguity attitude in gains + risk attitude in losses + ambiguity attitude in losses + neural activity under risky gains + neural activity under ambiguous gains + neural activity under risky losses+ neural activity under ambiguous losses + age + intelligence

All variables were standardized before fitting the linear model. All models included age and intelligence to control for these two demographic factors. We compared these three models by BIC.

Acknowledgement

This research project is supported by National Center for PTSD, and NIMH Grant R21MH102634. We also thank graduate student training grants from the Interdepartmental Neuroscience Program at Yale and the China Scholarship Council. We thank Alicia Roy and Dr. Erin O'Brien for the recruitment and clinical assessment of veterans; Dr. Michael A. Grubb for development of the task paradigm; Simon Podhajsky,

Yumiko Nakamura, and Pooja Salhotra for help with running experiments and data pre-processing. We also thank Dr. Jutta Joormann and Dr. Hyojung Seo for suggestions and comments on data analyses.

Reference

Admon, Roee, Mohammed R. Milad, and Talma Hendler. 2013. "A Causal Model of Post-Traumatic Stress Disorder: Disentangling Predisposed from Acquired Neural Abnormalities." *Trends in Cognitive Sciences* 17 (7): 337–47. https://doi.org/10.1016/j.tics.2013.05.005.

Akiki, Teddy J., Christopher L. Averill, Kristen M. Wrocklage, J. Cobb Scott, Lynnette A. Averill, Brian Schweinsburg, Aaron Alexander-Bloch, et al. 2018. "Default Mode Network Abnormalities in Posttraumatic Stress Disorder: A Novel Network-Restricted Topology Approach." *NeuroImage* 176 (April): 489–98. https://doi.org/10.1016/j.neuroimage.2018.05.005.

Bartra, Oscar, Joseph T. McGuire, and Joseph W. Kable. 2013. "The Valuation System: A Coordinate-Based Meta-Analysis of BOLD FMRI Experiments Examining Neural Correlates of Subjective Value." *NeuroImage* 76: 412–27. https://doi.org/10.1016/j.neuroimage.2013.02.063.

Beck, A T, C H Ward, M Mendelson, J Mock, and J Erbaugh. 1961. "An Inventory for Measuring Depression." *Arch Gen Psychiatry* 4: 561–71. https://www.ncbi.nlm.nih.gov/pubmed/13688369.

Bernstein, David P, Judith A Stein, Michael D Newcomb, Edward Walker, David Pogge, Taruna Ahluvalia, John Stokes, Leonard Handelsman, Martha Medrano, and David Desmond. 2003. "Development and Validation of a Brief Screening Version of the Childhood Trauma Questionnaire." *Child Abuse & Neglect* 27 (2): 169–90.

Bernstein, Eve M, and Frank W Putnam. 1986. "Development, Reliability, and Validity of a Dissociation Scale."

Blais, A R, and E U Weber. 2006. "A Domain-Specific Risk-Taking (DOSPERT) Scale for Adult

Populations." *Judgment and Decision Making Journal* 1 (1): 33–47.

Blake, D D, F W Weathers, L M Nagy, D G Kaloupek, F D Gusman, D S Charney, and T M Keane. 1995. "The Development of a Clinician-Administered PTSD Scale." *J Trauma Stress* 8 (1): 75–90. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_ui ds=7712061.

Blankenstein, Neeltje E, and Anna C K van Duijvenvoorde. 2019. "Neural Tracking of Subjective Value under Riskand Ambiguity in Adolescence." *Cognitive, Affective, & Behavioral Neuroscience* 19 (6): 1364–78. https://doi.org/10.3758/s13415-019-00749-5.

Brown, Vanessa M, Lusha Zhu, John M Wang, B Christopher Frueh, Brooks King-Casas, and Pearl H Chiu. 2018. "Associability-Modulated Loss Learning Is Increased in Posttraumatic Stress Disorder." Edited by Michael J Frank. *ELife* 7 (January): e30150. https://doi.org/10.7554/eLife.30150.

Buckholtz, Joshua W., Uma Karmarkar, Shengxuan Ye, Grace M. Brennan, and Arielle Baskin-Sommers. 2017. "Blunted Ambiguity Aversion during Cost-Benefit Decisions in Antisocial Individuals." *Scientific Reports* 7 (1): 1–9. https://doi.org/10.1038/s41598-017-02149-6.

Carver, C S, and T L White. 1994. "Behavioral-Inhibition, Behavioral Activation, and Affective Responses to Impending Reward and Punishment - the BIS/BAS Scales." *Journal of Personality and Social Psychology* 67 (2): 319–33.

Castrellon, Jaime J., Jacob S. Young, Linh C. Dang, Ronald L. Cowan, David H. Zald, and Gregory R. Samanez-Larkin. 2019. "Mesolimbic Dopamine D2 Receptors and Neural Representations of Subjective Value." *BioRXiv Preprint*.

Chew, Soo Hong, Richard P. Ebstein, and Songfa Zhong. 2012. "Ambiguity Aversion and Familiarity Bias: Evidence from Behavioral and Gene Association Studies." *Journal of Risk and Uncertainty* 44 (1): 1–18. https://doi.org/10.1007/s11166-011-9134-0.

Cox, R W. 1996. "AFNI: Software for Analysis and Visualization of Functional Magnetic Resonance Neuroimages." *Comput Biomed Res* 29 (3): 162–73.

Elman, Igor, Steven Lowen, Blaise B Frederick, Won Chi, Lino Becerra, and Roger K Pitman. 2009. "Functional Neuroimaging of Reward Circuitry Responsivity to Monetary Gains and Losses in Posttraumatic Stress Disorder." *Biol Psychiatry* 66 (12): 1083–90. https://doi.org/10.1016/j.biopsych.2009.06.006 S0006-3223(09)00732-X [pii].

Fiorillo, Christopher D. 2013. "Two Dimensions of Value: Dopamine Neurons Represent Reward but Not Aversiveness." *Science* 341 (6145): 546–49. https://doi.org/10.1126/science.1238699.

First, Michael B, Miriam Gibbon, Robert L Spitzer, Lorna Smith Benjamin, and Janet B W Williams. 1997. *Structured Clinical Interview for DSM-IV® Axis II Personality Disorders SCID-II*. American Psychiatric Pub.

Foa, Edna B, Terence M Keane, and Matthew J Friedman. 2000. "Guidelines for Treatment of PTSD." *Journal of Traumatic Stress* 13 (4): 539–88. https://doi.org/10.1023/A:1007802031411.

Fujiwara, Juri, Philippe N. Tobler, Masato Taira, Toshio Iijima, and Ken Ichiro Tsutsui. 2009. "Segregated and Integrated Coding of Reward and Punishment in the Cingulate Cortex." *Journal of Neurophysiology* 101 (6): 3284–93. https://doi.org/10.1152/jn.90909.2008.

Gilaie-Dotan, Sharon, Agnieszka Tymula, Nicole Cooper, Joseph W. Kable, Paul W. Glimcher, and Ifat Levy. 2014. "Neuroanatomy Predicts Individual Risk Attitudes." *Journal of Neuroscience* 34 (37): 12394–401. https://doi.org/10.1523/JNEUROSCI.1600-14.2014.

Gleichgerrcht, Ezequiel, Agustín Ibáñez, María Roca, Teresa Torralva, and Facundo Manes. 2010. "Decision-Making Cognition in Neurodegenerative Diseases." *Nature Reviews Neurology* 6 (11): 611–23. https://doi.org/10.1038/nrneurol.2010.148.

Grubb, Michael A., Agnieszka Tymula, Sharon Gilaie-Dotan, Paul W. Glimcher, and Ifat Levy. 2016.

"Neuroanatomy Accounts for Age-Related Changes in Risk Preferences." *Nature Communications* 7 (May): 1–5. https://doi.org/10.1038/ncomms13822.

Harpaz-Rotem, Ilan, Jack Tsai, Robert H Pietrzak, and Rani Hoff. 2014. "The Dimensional Structure of Posttraumatic Stress Symptomatology in 323,903 U.S. Veterans." *J Psychiatr Res* 49: 31–36. https://doi.org/10.1016/j.jpsychires.2013.10.020.

Hayes, Jasmeet P, Scott M Hayes, and Amanda M Mikedis. 2012. "Quantitative Meta-Analysis of Neural Activity in Posttraumatic Stress Disorder." *Biology of Mood & Anxiety Disorders* 2 (1): 1. https://doi.org/10.1186/2045-5380-2-9.

Homan, Philipp, Ifat Levy, Eric Feltham, Charles Gordon, Jingchu Hu, Jian Li, Robert H. Pietrzak, et al. 2019. "Neural Computations of Threat in the Aftermath of Combat Trauma." *Nature Neuroscience* 22 (3): 470–76. https://doi.org/10.1038/s41593-018-0315-x.

Hopper, James W., Roger K. Pitman, Zhaohui Su, Gene M. Heyman, Natasha B. Lasko, Michael L. Macklin, Scott P. Orr, Scott E. Lukas, and Igor Elman. 2008. "Probing Reward Function in Posttraumatic Stress Disorder: Expectancy and Satisfaction with Monetary Gains and Losses." *Journal of Psychiatric Research* 42 (10): 802–7. https://doi.org/10.1016/j.jpsychires.2007.10.008.

Jung, Wi Hoon, Sangil Lee, Caryn Lerman, and Joseph W. Kable. 2018. "Amygdala Functional and Structural Connectivity Predicts Individual Risk Tolerance." *Neuron* 98 (2): 394-404.e4. https://doi.org/10.1016/j.neuron.2018.03.019.

Kable, Joseph W, and Paul W Glimcher. 2009. "The Neurobiology of Decision: Consensus and Controversy." *Neuron* 63 (6): 733–45. https://doi.org/S0896-6273(09)00681-3 [pii] 10.1016/j.neuron.2009.09.003.

Kahnt, Thorsten, Soyoung Q. Park, John-Dylan Dylan Haynes, and Philippe N. Tobler. 2014. "Disentangling Neural Representations of Value and Salience in the Human Brain." *Proceedings of*

the National Academy of Sciences of the United States of America 111 (13): 5000–5005.
https://doi.org/10.1073/pnas.1320189111.

Kaufman, Alan S. 1990. *Kaufman Brief Intelligence Test: KBIT*. AGS, American Guidance Service Circle
Pines, MN.

Keane, Terence M, John A Fairbank, Juesta M Caddell, Rose T Zimering, Kathryn L Taylor, and
Catherine A Mora. 1989. "Clinical Evaluation of a Measure to Assess Combat Exposure."
Psychological Assessment: A Journal of Consulting and Clinical Psychology 1 (1): 53.

Kim, Sang Hee, Heung Sik Yoon, Hackjin Kim, and Stephan Hamann. 2014. "Individual Differences in
Sensitivity to Reward and Punishment and Neural Activity during Reward and Avoidance
Learning." *Social Cognitive and Affective Neuroscience* 10 (9): 1219–27.
https://doi.org/10.1093/scan/nsv007.

Kolk, Bessel van der, Mark Greenberg, Helene Boyd, and John Krystal. 1985. "Inescapable Shock,
Neurotransmitters, and Addiction to Trauma: Toward a Psychobiology of Post Traumatic Stress."
Biological Psychiatry 20 (3): 314–25. https://doi.org/10.1016/0006-3223(85)90061-7.

Konova, Anna B., Silvia Lopez-Guzman, Adelya Urmanche, Stephen Ross, Kenway Louie, John
Rotrosen, and Paul W. Glimcher. 2019. "Computational Markers of Risky Decision-Making for
Identification of Temporal Windows of Vulnerability to Opioid Use in a Real-World Clinical
Setting." *JAMA Psychiatry* 08855. https://doi.org/10.1001/jamapsychiatry.2019.4013.

Lawrence, Michael A. 2013. "Ez: Easy Analysis and Visualization of Factorial Experiments." *R Package
Version* 3 (2).

Lesage, Elise, and Elliot A Stein. 2016. "Networks Associated with Reward." In *Neuroscience in the 21st
Century*, 1–27. https://doi.org/10.1007/978-1-4614-6434-1.

Levy, Ifat, Jason Snell, Amy J. Nelson, Aldo Rustichini, and Paul W. Glimcher. 2010. "Neural

Representation of Subjective Value Under Risk and Ambiguity." *Journal of Neurophysiology* 103 (2): 1036–47. https://doi.org/10.1152/jn.00853.2009.

Litt, Ab, Hilke Plassmann, Baba Shiv, and Antonio Rangel. 2011. "Dissociating Valuation and Saliency Signals during Decision-Making." *Cereb Cortex* 21 (1): 95–102. https://doi.org/10.1093/cercor/bhq065.

Lumley, Thomas, and Alan Miller. 2009. "Leaps: Regression Subset Selection." *R Package Version* 2: 2366.

Macoveanu, Julian. 2014. "Serotonergic Modulation of Reward and Punishment: Evidence from Pharmacological FMRI Studies." *Brain Research*. https://doi.org/10.1016/j.brainres.2014.02.003.

May, Casey L., and Blair E. Wisco. 2019. "Reward Processing and Decision-Making in Posttraumatic Stress Disorder." *Behavior Therapy*, November. https://doi.org/10.1016/J.BETH.2019.11.005.

Monosov, Ilya E., and Okihide Hikosaka. 2012. "Regionally Distinct Processing of Rewards and Punishments by the Primate Ventromedial Prefrontal Cortex." *Journal of Neuroscience* 32 (30): 10318–30. https://doi.org/10.1523/JNEUROSCI.1801-12.2012.

Mukherjee, Dahlia, Sangil Lee, Rebecca Kazinka, Theodore D. Satterthwaite, and Joseph W Kable. 2020. "Multiple Facets of Value-Based Decision Making in Major Depressive Disorder." *Scientific Reports* 10 (1): 3415. https://doi.org/10.1038/s41598-020-60230-z.

Nawijn, Laura, Mirjam van Zuiden, Jessie L. Frijling, Saskia B.J. Koch, Dick J. Veltman, and Miranda Olff. 2015. "Reward Functioning in PTSD: A Systematic Review Exploring the Mechanisms Underlying Anhedonia." *Neuroscience and Biobehavioral Reviews*. https://doi.org/10.1016/j.neubiorev.2015.01.019.

O'Doherty, J., M.L. L Kringelbach, E.T. T Rolls, J. Hornak, and C. Andrews. 2001. "Abstract Reward and Punishment Representations in the Human Orbitofrontal Cortex." *Nat Neurosci* 4 (1): 95–102.

https://doi.org/10.1038/82959.

Patton, J H, M S Stanford, and E S Barratt. 1995. "Factor Structure of the Barratt Impulsiveness Scale." *J Clin Psychol* 51 (6): 768–74. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_ui ds=8778124.

Paulus, Martin P., and Angela J. Yu. 2012. "Emotion and Decision-Making: Affect-Driven Belief Systems in Anxiety and Depression." *Trends in Cognitive Sciences*. https://doi.org/10.1016/j.tics.2012.07.009.

Peterson, Brian G, and Peter Carl. 2019. "PerformanceAnalytics: Econometric Tools for Performance and Risk Analysis." https://cran.r-project.org/package=PerformanceAnalytics.

Pinheiro, J, D Bates, S DebRoy, and D Sarkar. 2013. "R Development Core Team (2013)'Nlme: Linear and Nonlinear Mixed Effects Models.'R Package Version 3.1-107." *R Foundation for Statistical Computing, Vienna. Available: Cran. r-Project. Org/Web/Packages/Nlme./(February 2013)*.

Pushkarskaya, Helen, David Tolin, Lital Ruderman, Ariel Kirshenbaum, J. MacLaren Kelly, Christopher Pittenger, and Ifat Levy. 2015. "Decision-Making under Uncertainty in Obsessive–Compulsive Disorder." *Journal of Psychiatric Research* 69 (October): 166–73. https://doi.org/10.1016/J.JPSYCHIRES.2015.08.011.

R Core Team. 2018. "R: A Language and Environment for Statistical Computing."

Rabellino, Daniela, Maria Densmore, Paul A Frewen, Jean Théberge, C Mckinnon, and Ruth A Lanius. 2016. "Aberrant Functional Connectivity of the Amygdala Complexes in PTSD during Conscious and Subconscious Processing of Trauma-Related Stimuli." *PLoS One*, 1–12. https://doi.org/10.1371/journal.pone.0163097.

Revelle, William. 2018. "Psych: Procedures for Personality and Psychological Research. Northwestern

University, Evanston."

Reynolds, Sheila M., and Kent C. Berridge. 2008. "Emotional Environments Retune the Valence of Appetitive versus Fearful Functions in Nucleus Accumbens." *Nature Neuroscience* 11 (4): 423–25. https://doi.org/10.1038/nn2061.

Ruderman, Lital, Daniel B. Ehrlich, Alicia Roy, Robert H. Pietrzak, Ilan Harpaz-Rotem, and Ifat Levy. 2016. "Posttraumatic Stress Symptoms and Aversion To Ambiguous Losses in Combat Veterans." *Depress Anxiety* 33 (7): 606–13. https://doi.org/10.1002/da.22494.

Russell, L. 2018. "Emmeans: Estimated Marginal Means, Aka Least-Squares Means." *R Package Version* 1 (2).

Sailer, Uta, Simon Robinson, Florian Ph S. Fischmeister, Dorothea König, Claudia Oppenauer, Brigitte Lueger-Schuster, Ewald Moser, Ilse Kryspin-Exner, and Herbert Bauer. 2008. "Altered Reward Processing in the Nucleus Accumbens and Mesial Prefrontal Cortex of Patients with Posttraumatic Stress Disorder." *Neuropsychologia* 46 (11): 2836–44. https://doi.org/10.1016/j.neuropsychologia.2008.05.022.

Shabel, Steven J., Chenyu Wang, Bradley Monk, Sage Aronson, and Roberto Malinow. 2019. "Stress Transforms Lateral Habenula Reward Responses into Punishment Signals." *Proceedings of the National Academy of Sciences of the United States of America* 116 (25): 12488–93. https://doi.org/10.1073/pnas.1903334116.

Spielberger, Charles Donald, and Richard L Gorsuch. 1983. *State-Trait Anxiety Inventory for Adults: Manual and Sample: Manual, Instrument and Scoring Guide*. Consulting Psychologists Press.

Tom, Sabrina M., Craig R. Fox, Christopher Trepel, and Russell A. Poldrack. 2007. "The Neural Basis of Loss Aversion in Decision-Making under Risk." *Science* 315 (5811): 515–18. https://doi.org/10.1126/science.1134239.

Weathers, Frank W, Brett T Litz, Terence M Keane, Patrick A Palmieri, Brian P Marx, and Paula P
Schnurr. 2013. "The PTSD Checklist for DSM-5 (PCL-5)." *Scale Available from the National
Center for PTSD at Www. Ptsd. va. Gov* 10.

Wickham, Hadley. 2016. *Ggplot2: Elegant Graphics for Data Analysis*. Springer.

Wolf, Richard C, and Ryan J Herringa. 2015. "Prefrontal – Amygdala Dysregulation to Threat in
Pediatric Posttraumatic Stress Disorder." *Neuropsychopharmacology* 41 (3): 822–31.
https://doi.org/10.1038/npp.2015.209.

Yacubian, Juliana, Jan Glascher, Katrin Schroeder, Tobias Sommer, Dieter F Braus, and Christian
Buchel. 2006. "Dissociable Systems for Gain- and Loss-Related Value Predictions and Errors of
Prediction in the Human Brain." *J Neurosci* 26 (37): 9530–37.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_ui
ds=16971537.

Zhang, Zhihao, Jennifer Fanning, Daniel B. Ehrlich, Wenting Chen, Daeyeol Lee, and Ifat Levy. 2017.
"Distributed Neural Representation of Saliency Controlled Value and Category during Anticipation
of Rewards and Punishments." *Nature Communications* 8 (1): 1907. https://doi.org/10.1038/s41467-
017-02080-4.

Zyphur, Michael J., Jayanth Narayanan, Richard D. Arvey, and Gordon J. Alexander. 2009. "The
Genetics of Economic Risk Preferences." *Journal of Behavioral Decision Making* 22 (4): 367–77.
https://doi.org/10.1002/bdm.