Short Title: Genetic Architecture of Maize Rind Strength

Author for Correspondence: Rajandeep S. Sekhon

Genetic Architecture of Maize Rind Strength Revealed by the Analysis of Divergently Selected
Populations

Rohit Kumar[1], Abiskar Gyawali[2], Ginnie D. Morrison[2], Christopher A. Saski[3], Daniel J.
Robertson[4,] Nishanth Tharayil[3], Robert J. Schaefer[5], Timothy M. Beissinger[6,7], Rajandeep S.
Sekhon[1]
8
1 Department of Genetics and Biochemistry, Clemson University, Clemson, SC 29634, US
2 University of Missouri, Division of Biological Sciences, 105 Tucker Hall, Columbia, MO,
65211, USA
3 Department of Plant and Environmental Sciences, Clemson University, Clemson, SC 29634,
USA
4 Department of Mechanical Engineering, University of Idaho, Moscow, ID, USA
5 Linkage Analytics, LLC, Denver, CO 80216, USA
6 University of Göttingen, Department of Plant Breeding Methodology, Göttingen, 37075,
Germany
7 University of Göttingen, Center for Integrated Breeding Research, Göttingen, 37075, Germany
19

20   One sentence summary: Selection mapping and transcriptomic analysis of divergently selected
21   populations identifies key candidate genes for improvement of rind penetrometer resistance and
22   stalk strength in maize.
23

24   Author Contributions: R.S.S. and T.M.B designed the research; R.K., A.G., N.T., T.M.B., and
25   R.S. performed the research; R.K., A.G., G.D.M., C.A.S., D.J.R., R.J.S., and T.M.B. analyzed
26   the data; R.K., T.M.B., and R.S.S. wrote the paper.
27

Finding: Funding for this study was provided by National Science Foundation grant #1826715.

Author for contact email: Rajandeep S. Sekhon; sekhon@clemson.edu

ABSTRACT

Stalk lodging, breakage of the stalk at or below the ear, causes substantial yield losses in maize.

The strength of the stalk rind, commonly measured as rind penetrometer resistance (RPR), is an

important contributor to stalk lodging resistance. To enhance RPR genetic architecture, we

conducted selection mapping on populations developed by 15 cycles of divergent selection for

high (C15-H) and low (C15-L) RPR. We also performed time-course transcriptome and

metabolic analyses on developing stalks of high (Hrpr1) and low (Lrpr1) RPR inbred lines

derived from the C15-H and C15-L populations, respectively. Divergent selection significantly

altered allele frequencies at 3,656 and 3,412 single nucleotide polymorphisms (SNP) in the C15-

H and C15-L populations, respectively. While the majority of the SNPs under selection were

unique, 110 SNPs were common in both populations indicating the fixation of alleles with

alternative effects. Remarkably, preferential selection on the genomic regions associated with

lignin and polysaccharide biosynthesis genes was observed in C15-H and C15-L populations,

respectively. This observation was supported by higher lignification and lower extractability of

cell wall-bound sugars in Hrpr1 compared to Lrpr1. Tricin, a monolignol important for

incorporation of lignin in grass cell walls, emerged as a key determinant of the different cell wall

properties of Hrpr1 and Lrpr1. Integration of selection mapping with transcriptomics and

previous genetic studies on RPR identified 40 novel candidate genes including Zm*MYB31*,

ZmNAC25, ZmMADS1, two *PAL* paralogues, two lichenases, *ZmEXPA2, ZmIAA41,* and

Caleosin. Enhanced mechanistic and genetic understanding of RPR provides a foundation for

improved stalk lodging resistance.

INTRODUCTION

Maize (*Zea mays* L.) is one of the most important cereal crops in the world. Stalk lodging, which refers to the breakage of the stalk at or below the ear, is estimated to cause 5% - 25% annual yield losses in maize (Zuber and Kang, 1978; Flint-Garcia et al., 2003). Furthermore, stalk lodging often results in decreased grain quality and an increased presence of pests and disease due to the decay of fallen (i.e., lodged) grains. Increased use of nitrogen and higher planting density associated with the cultivation of higher-yielding maize hybrids are poised to further increase stalk lodging incidence (Yang et al., 2019). Improving the extractability of sugars for forage or biofuels also accompanies decreased stalk strength and increased lodging (Pedersen et al., 2005; Feltus and Vandenbrink, 2012). Finally, ever-worsening climate and associated extreme weather events are expected to enhance yield losses associated with lodging. A comprehensive understanding of the genetic architecture of stalk strength is key to the successful breeding and biotechnological intervention targeted to improve grain yield and biomass quality.

Phenotypic assessment of stalk lodging resistance has been challenging primarily because of difficulties in determining the aspects of stalk strength that translate into higher lodging resistance in field conditions. Indirect methods for prediction of stalk lodging resistance include counting lodged plants in field evaluations (Robertson et al., 2016), analyzing the chemical composition of stalks (Davidson and Phillips, 1930; Appenzeller et al., 2004; Ching et al., 2006), measuring stalk bending strength (Robertson et al., 2014; Robertson et al., 2016), stalk crushing strength (Zuber and Grogan, 1961; Undersander et al., 1977; Zuber et al., 1980), and rind penetrometer resistance (Zuber et al., 1980; Sibale et al., 1992). Rind strength is proposed to contribute to 50–80% of stalk strength (Zuber et al., 1980). Rind strength can be assessed by forcing a small probe through a plant stalk and measuring the maximum strength required to puncture the rind. This method, which measures rind penetrometer resistance (RPR), also known as rind puncture resistance, has been used throughout most of the 20th century to investigate stalk strength (Khanna, 1935). RPR has been shown to be significantly and negatively correlated with naturally occurring stalk lodging in the field (Dudley, 1994; Kang et al., 1999; Jampatong et al., 2000; Hu et al., 2012; Sekhon et al., 2019). In a recent and comprehensive study, RPR data collected on 47 maize hybrids in three environments was shown to be a reliable predictor of stalk lodging incidence of these hybrids observed in 98 temporally and spatially distinct environments

(Sekhon et al., 2019). To summarize, RPR offers a high-throughput phenotyping method for

artificial selection in breeding programs and for genetic studies aimed to understand the genetic

architecture of lodging resistance.

Several genetics studies have reported RPR as a quantitative trait and identified

quantitative trait loci (QTL) or single nucleotide polymorphisms (SNP) putatively associated

with this trait (Heredia et al., 1996; Flint-Garcia et al., 2003; Hu et al., 2012; Peiffer et al., 2013).

To exploit RPR for improvement of stalk lodging resistance, a divergent (i.e., bidirectional)

selection experiment in Missouri Second Cycle Stiff Stalk Synthetic (MoSCSSS), a yellow dent

synthetic population formed from intermating of 14 inbred lines, was initiated using recurrent S_0

selection. While the original MoSCSSS population, designated as cycle 0 (C0), had RPR value

of 3.55 kgf, twelve cycles of selection for high and low RPR from this population resulted in the

development of high and low RPR populations with RPR value of 8.52 kgf and 2.0 kgf,

respectively (Martin et al., 2004). Multiple rounds of such divergent selection have the potential

to fix alternative alleles with contrasting effects on RPR or to substantially alter the allele

frequencies of alternate alleles in these populations. Therefore, the populations developed by

divergent selection were for the discovery of QTL underlying RPR (Flint-Garcia et al., 2003). To

this end, $F_{2:3}$ families obtained from the high RPR population derived from 10 cycles of selection

were crossed with: 1) $F_{2:3}$ families obtained from the low RPR population derived from 11 cycles

of selection, 2) an S_1 plant with low RPR selected from an unrelated population (MoSQB-Low),

and 3) with an inbred line Mo47. The resulting mapping populations were used to identify 26

QTL for RPR (Flint-Garcia et al., 2003). Evaluation of recombinant inbred lines (RILs) from a

maize nested association mapping population and a large collection of diverse inbred lines

identified several significant single nucleotide polymorphisms (SNPs) and QTL associated with

RPR (Peiffer et al., 2013). Evaluation of RILs derived from B73 and a high kernel oil inbred

resulted in the identification of nine QTL for RPR (Hu et al., 2012), and evaluation of two sets of

RILs reported seven QTL (Li et al., 2014). These studies provide rich information about genomic

regions that control RPR, and identification of the underlying genes will help enhance stalk

strength. Recently, cloning of a major QTL underlying RPR resulted in the identification of a

novel gene *stiff1*, and knockdown of *stiff1,* either naturally through a transposon insertion in the

gene promoter or experimentally through gene editing, enhanced stalk strength (Zhang et al.,

2020). However, progress in cloning and characterizing the underlying genes has been slow and

requires novel genomic approaches to improve the resolution of existing genetic information on

the genomic regions governing RPR.

Divergent selection experiments enable a direct assessment of the response of a genome

to selection for a trait. This in turn leads to better understanding and a greater ability to

characterize the genetic architecture of the given trait. Populations derived from divergent

selection are an example of experimental evolution (Turner et al., 2011), and hence exhibit

differences in allele frequency between pre- and post-selection populations. Modern genotyping

approaches allow the assessment of allele frequency changes in experimentally-evolved

populations to map putatively causal loci (Hirsch et al., 2014; Lorenz et al., 2015; Kelly and

Hughes, 2019). The field of gene-mapping has been dominated by phenotype-to-genotype

correlation analyses including genome-wide association (GWA) and QTL mapping for several

decades (Pascual et al., 2016; Tam et al., 2019). Scanning for selective sweeps is an additional

tactic that can be effectively utilized to identify trait-relevant genomic regions and genes in

selected populations (Nielsen et al., 2005; Kimura et al., 2007; Qanbari et al., 2012; Beissinger et

al., 2014; Chen et al., 2018; Grainger et al., 2018; You et al., 2018). For agricultural species,

previous breeding and long-term selection studies have already generated many such populations

(Olsen and Wendel, 2013). In contrast to GWA and QTL mapping studies, selection scans do not

depend on phenotypes from genotyped individuals. Instead, with the given knowledge that

selection for a specific trait or traits has occurred, a scan for selected loci may proceed without

any additional phenotyping (Gerke et al., 2015). In addition to eliminating time-intensive

phenotyping, selection mapping can serve as an independent mapping technique to corroborate

region-trait associations identified via other methods.

The current study presents phenotypic, genomic, and transcriptomic analyses of

divergently selected maize populations to gain mechanistic insights into RPR in maize.

Specifically, we 1) examined the phenotypic changes associated with divergent selection in the

population obtained from additional cycles of selections than those reported earlier, and inbred

lines developed from these populations, 2) examined the selection signatures to identify the

divergently selected genomic segments in the populations, 3) performed transcriptomic and

metabolic analyses on the inbred lines with high and low RPR derived from the aforementioned

divergently selected populations, and 4) combined the data from the three experiments and

previously published studies to begin constructing a comprehensive picture of the genetic

architecture of RPR in maize. Results from this study will boost efforts towards the identification

of genes and genetic elements underlying RPR and alleviation of lodging-related losses in

cereals.

RESULTS

Divergent selection resulted in significant changes in rind penetrometer resistance

Phenotypic changes after twelve cycles of divergent selection for RPR on MoSCSSS (C0)

population have already been reported (Martin et al., 2004). To test the effect of three additional

cycles of selection in each direction, we recorded RPR in C15-H and C15-L populations and

found significant differences among the three populations (Figure 1). Mean RPR of C0, C15-L,

and C15-H was 4.55 Kgf, 2.1 Kgf, and 11.9 kgf, respectively.

Divergent selection significantly modified the frequency of several SNPs genome-wide

We computed F_{ST} and changes in allele frequency between C0 and C15-L, and between C0 and

C15-H, as well as F_{ST} values for these comparisons (Figure 2). Highly significant correlations

between changes in allele frequency and F_{ST} were observed both for high RPR selection ($r =$

0.9644; $p <$ 1e-324) and low RPR selection ($r = $ 0.9641; $p <$ 1e-324). Using allele frequency

change from C0 to C15-H to establish significance thresholds, we identified 3,656 significant

SNPs that were putatively selected for high RPR (Figure 2, Table S1). Likewise, based on allele

frequency change from C0 to C15-L, we identified 3,412 SNPs that showed significant evidence

of selection for low RPR (Figure 2, Table S1). In both cases, observed allele frequency changes

cannot be explained by drift alone, although we cannot exclude the possibility that other non-

neutral force(s) drove changes in allele frequency. Those forces could include unintentional

selection on additional traits besides high RPR and low RPR, such as general fitness. Of these

significant SNPs, 110 SNPs were selected in both directions (Figure 2, Table S1) suggesting

fixation of alternative alleles of certain genes. Bootstrap resampling indicated that the extent of

such overlap is significant ($p \leq$ 1e-4). For selection in both directions, there were clusters of

many SNPs showing significant selection signals. Such clusters included the distal-arm of

chromosome 2 for C15-H and the distal-arm of chromosome 7 for C15-L, respectively (Figure

2). Regions represented by these clusters likely harbor large-effect genes that experienced strong

selection during early generations of the experiment.

Divergence in RPR phenotype accompanied with distinct chemical composition of internodes

C15-H and C15-L populations are maintained in a heterozygous state (Flint-Garcia et al., 2003;

Martin et al., 2004). To fix and study a subset of alleles that underlie the diverged phenotype and

likely regulate RPR, we developed Hrpr1 and Lrpr1 inbred lines from C15-H and C15-L

populations, respectively. The RPR phenotype and transcriptome associated with RPR was

measured on the 12^{th} internode which, on average, represented the internode below the ear

bearing node in the inbred lines. The internodes were immature and completely lacked RPR at 0

DAS stage as indicated by a lack of detected RPR value by the rind penetrometer (Figure 3A).

The two inbred lines started to significantly diverge at 6 DAS, and 9 DAS appeared to be a

distinctive stage with a sharp increase in RPR of Hrpr1 compared to Lrpr1. While RPR

continued to increase in both inbreds after 12 DAS, albeit at a lower rate, Hrpr1 maintained

significantly higher RPR compared to Lrpr1.

The constituents of cell wall, including polysaccharides (cellulose and hemicellulose) and

secondary metabolites (primarily lignin), are proposed to be important determinants of stalk

strength (Bosch et al., 2011; Peiffer et al., 2013). To assess their role in the observed differences

in RPR, we examined the accumulation of these metabolites in the internodes of Hrpr1 and

Lrpr1. Remarkably, RPR levels were negatively correlated with the amount of these

monosaccharides released from the internode cell wall, and significantly higher glucose and

pentose was obtained from Lrpr1 compared to Hrpr1 (Figure 3B). Histological staining of the

internode sections revealed higher lignification of sclerenchyma and rind-region parenchyma in

Hrpr1 compared to Lrpr1 (Figure 3C). The differences in histological staining, and the release of

sugars from the lignocellulosic biomass (i.e., digestibility), in grasses are attributed to increased

lignification (Akin, 1989). Lignin is primarily composed of syringyl, guaiacyl, and, to a much

lesser extent, p-hydroxyphenyl monomers (Halpin, 2019; Ralph et al., 2019). Lignin extraction

in our study, performed in oxidizing conditions, measured syringic acid (S), vanillic acid (V),

and p-hydroxybenzoate (H) (Wang et al., 2015). The total amount of these three monolignols

was comparable up to 9 DAS stage in both inbred lines, but the Hrpr1 internodes accumulated

higher amounts at the later stages (Figure 3D). Accumulation of S and V lignin followed the

same trend (Figure 3E-F, Figure S1A-F). Accumulation of H lignin was quite low in both

inbreds albeit Lrpr1 internode accumulated slightly but significantly higher amounts of this

monomer compared to Hrpr1 at early stages, but this trend was reversed later in development

(Figure 3G, Figure S1G-I).

Incorporation of lignin monomers in the cell wall is of paramount importance for

imparting mechanical strength (Tobimatsu and Schuetz, 2019). Polymerization of lignin

monomers in grass cell walls is achieved by the activity of ferulic acid (FA) and *p*-coumaric acid

(pCA) (Hatfield et al., 2017). Interestingly, both pCA and FA were slightly higher in Lrpr1 at 3

DAS but eventually increased to significantly higher amounts in Hrpr1 (Figure 3H-I). Tricin is

an important monolignol involved in lignification by acting as a nucleation site for

polymerization of lignin monomers in grass cell walls (Lan et al., 2015). The cell wall-bound

fraction of tricin, which signifies the tricin incorporated in the lignin polymer, was slightly

higher in Hrpr1 at 3 DAS and, while the differences diminished at 9 DAS, Hrpr1 internodes had

higher amounts in the later stages of development (Figure 3J). Conversely, the unbound fraction

of tricin showed a reverse trend with a higher accumulation of Lrpr1 (Figure 3K). In summary,

while the differences in lignification were minimal at the early stages of development, higher

lignification of internode cell walls of Hrpr1 as compared to Lrpr1 was evident at the later stages

of development.

Divergent selection for high and low RPR resulted in enrichment of distinct pathways of cell

wall synthesis

Since lignin and polysaccharides emerged as two key classes of metabolites associated with

distinct RPR phenotypes in the inbred lines, we examined the effect of divergent selection on

genes related to the synthesis of these metabolites. We first identified the genes and gene

families associated with the synthesis of cellulose, hemicellulose, and lignin in the maize

genome. This analysis identified 504 genes associated with cellulose and hemicellulose

synthesis, and 341 genes associated with lignin synthesis (Table 1, Table S2). We then assessed

if the significant SNPs associated with these genes were enriched in SNPs identified through

selection mapping (Table S1), during divergent selection for high and low RPR. In the genome

space covered by the lignin pathway genes, 45 SNPs showed a significant signal of selection in

C15-H population compared to 26 SNPs the C15-L population. Bootstrap analysis further

showed that C15-H population was significantly enriched for SNPs linked to the lignin pathway
genes compared to whole-genome background levels of selection ($p \leq 0.004$) while C15-L
population did not show enrichment ($p \leq 0.491$) (Figure 4). In contrast, the SNPs in the genome
space covering the polysaccharide synthesis genes showed an opposite pattern. Eighty SNPs
showed a significant signal of selection in the C15-L population compared to 60 enriched SNPs
in the C15-H population. Bootstrap analysis indicated that the C15-L population was enriched
for significant SNPs in polysaccharide genome space ($p \leq 0.001$) while the C15-H population
was not enriched ($p \leq 0.34$) (Figure 4).

Characterization of transcriptome underlying RPR
To understand the transcriptome dynamics underlying the divergently selected phenotypes for
low and high RPR, we performed RNA-seq on the developing internodes of Lrpr1 and Hrpr1 at
key developmental stages (Figure 5). A snapshot of transcriptome dynamics obtained by
principal component analysis (Figure 5A) was consistent with developmental divergence in the
RPR phenotype in Lrpr1 and Hrpr1 (Figure 3A). Both inbred lines showed close clustering at 0
and 3 DAS but dispersed considerably in the later stages indicating distinct transcriptomes of the
internodes.
Comparative analysis of transcriptome changes between Lrpr1 and Hrpr1 at each of the
different stages (0 DAS to 27 DAS) showed that 8327 (21%) genes were differentially expressed
(DE) during one or more stages (Figure 5B, Table S3). Consistent with phenotypic changes
(Figure 3A), a larger number of DE genes were observed after 3 DAS. Furthermore, while the
early stages at 3, 6, and 9 DAS were characterized by more DE genes showing upregulation in
Hrpr1 relative to Lrpr1 and suggesting higher RPR to be a biologically more active process, this
trend was reversed in the later stages. Gene ontology (GO) enrichment analysis of upregulated
and downregulated genes identified several important enriched biological processes (GOBP)
(Figure 5C). Notable upregulated GOBP in Hrpr1 consisted of phenylpropanoid biosynthesis,
amino acid transport, dicarboxylic acid biosynthesis, pyridine-containing compound synthesis,
and quinolinate metabolism. Major upregulated GOBP in Lrpr1 consisted of xylan biosynthesis,
cellular glucan metabolic process, amino sugar metabolic process, cell wall organization, defense
response, gibberellic acid- and jasmonic acid-mediated signaling pathways, lignin catabolic
process, secondary cell wall biogenesis, and response to oxidative stress.

To further understand the distinct transcriptome dynamics during the development of the

high and the low RPR phenotype in the internodes, we compared the transcriptome at 0 DAS

with subsequent developmental stages for each of the inbred lines (Figure S2). The total number

of DE genes at various stages of RPR development relative to 0 DAS was similar in Hrpr1

(10758, 27.1%) and Lrpr1 (11701, 29.5%) (Figure S2A-B). However, more DE genes were

detected in Hrpr1 relative to Lrpr1 at the early stages of 3, 6, 9 DAS. Furthermore, consistent

with the combined analysis (Figure 5B), more DE genes were upregulated that downregulated in

Hrpr1 at compared to Lrpr1 at these early stages (Figure S2A-B), further supporting enhanced

biological activity underlying high RPR phenotype. Finally, among the DE genes, 2060 (15%)

and 3005 (21.8%) were unique for Hrpr1 and Lrpr1, respectively, while 8696 (63.2%) were

common for both inbreds (Figure 5D) suggesting that relatively fewer genes are responsible for

distinct RPR phenotype. The GO enrichment analysis of unique and common genes showed that

oxidation-reduction and phosphatidylcholine metabolic process predominantly enriched in Hrpr1

while nucleic acid phosphodiester bond hydrolysis and RNA modification were enriched in

Lrpr1 (Figure S2C).

Integrating selection mapping with gene expression identified high priority genes associated

with RPR

Given the large number of candidate genes obtained from selection mapping and from

differential expression in Lrpr1 and Hrpr1, we applied an integrative approach to identify the

high priority RPR associated genes. From the SNPs that underwent significant change in

frequency during divergent selection, we chose 5% SNPs with the highest F_{ST} resulting in 183

and 171 high priority SNPs for C15-H and C15-L populations, respectively (Table S4). We

mapped these SNPs to nearby two genes within a 100 kb (100 kb on each side) window and

further filtered them based on their intersection with differentially expressed genes (fold change

>2) between Lrpr1 and Hrpr1 at each of the developmental stages. Remarkably, 98 and 76 genes

were found to be associated with low and high RPR, respectively (Table S5). The GO analysis of

these genes showed the notable enrichment of pigments and glucuronoxylan synthesis in low

RPR phenotype while enrichment of lignin biosynthesis, drought recovery, stress response, and

salicylic acid catabolic process in the high RPR phenotype (Figure 6, Table S6).

Meta-analysis of the genetic architecture of RPR

To generate a comprehensive overview of the genetic architecture of RPR and for the
identification of high confidence candidate genes, we combined the findings of our study with
other diverse lines of evidence. The criteria used for meta-analysis were 1) location of each of
the 354 high priority SNPs (Table S4); 2) genomic regions identified from previous genetic
studies including QTL (Flint-Garcia et al., 2003; Hu et al., 2012; Li et al., 2014) and GWAS
(Peiffer et al., 2013) analyses; 3) differential expression of the genes identified from overlapping
SNPs (Table S3); 4) overlap with genes associated with SNPs enriched in polysaccharide and
lignin genome space. Since flanking markers were not available for the QTL derived from
MoSCSSS-derived populations (Flint-Garcia et al., 2003), we added 10 Mb windows on both
sides of the marker designated as the QTL position in the study. For the NAM GWAS SNPs,
genes present within ±100 kb region of associated SNPs were included in this analysis. This
analysis identified several key regions, particularly at chromosomes 2, 3, 7, and 9 that
overlapped among multiple studies and, therefore, potentially harbor genes and/or regulatory
elements associated with RPR (Figure 7). Using at least three of the four criteria defined for
meta-analysis, forty candidate genes were identified in these regions (Table 2). For instance, a
genomic region on chromosome 2, supported by two additional studies, identified a Myb
transcription factor *ZmMYB31* as a candidate gene involved in specifying the high RPR
phenotype. Another region on chromosome 3, supported by at least two additional studies,
identified a glucosidase encoded by *geb1*. Interestingly, two gene *lic1* and *lic2* identified on
chromosome 6 region were supported by an additional study. Another important region on
chromosome 9, supported by three additional studies, harbors two candidate genes including a
MADSs box gene *ZmMADS1* and an expansin encoded by *Expansin-like A2 (ZmEXPA2)*. Other
key candidate genes in the genomic regions identified by at least one more genetic study besides
the current study for high RPR include *phenylalanine ammonia lyases* (*ZmPAL7, ZmPAL9*),
Caleosin protein and Xyloglucan galactosyltransferase MUR3. The important genes identified to
be associated with low RPR phenotype and supported by additional genetics studies include
ZmNAC25, *receptor-like protein kinase HSL1*, *ZmWRKY55*, *DELLA protein RGA* (*gras46*),
xyloglucan 6-xylosyltransferase 2 (*ZmGT5*), *histidine kinase5* (*hk5*), and *auxin-responsive*
protein IAA41 (*ZmIAA41*). The genes identified from the meta-analysis are promising candidates
for RPR manipulation in future studies.

DISCUSSION

Rind strength is an important predictive phenotype of stalk strength and stalk lodging resistance. Rind penetrance resistance (a.k.a. rind puncture resistance) is an effective measure of rind strength that has been shown to be negatively correlated with the stalk lodging incidence in several field studies (Dudley, 1994; Hu et al., 2012; Sekhon et al., 2019). However, lack of knowledge about the genetic mechanisms underlying RPR has been a major obstacle to improving stalk strength and stalk lodging resistance. We employed selection mapping in diverse populations, transcriptomics of inbred lines with extreme RPR phenotypes, metabolite analyses, and combined these findings with the previous genetic studies to provides a comprehensive overview of the genetic architecture of the RPR trait in maize.

Divergent selection significantly altered allele frequencies at putative RPR-associated regions

Divergently selected populations provide a unique and valuable resource for mining genes and alleles underlying RPR and, therefore, stalk strength. A frequent and valid critique of selection mapping experiments carried out on wild or domesticated populations (Hufford et al., 2012; Beissinger et al., 2016; Plassais et al., 2019) is that these approaches rely on "outlier tests" for identification of divergently selected SNPs/regions. Consequently, these tests are not equivalent to significance tests (Narum and Hess, 2011; Lorenz et al., 2015). To avoid this criticism, we established significance by comparing observed allele frequency differentiation between selected and unselected populations. This approach allowed us to account for differentiation based on genetic drift alone. The rationale for applying this approach is that, like in a subset of other breeding studies carried out within over a short time span of a few to tens of generations on an evolutionary scale (Lorenz et al., 2015), accurate demographic parameters were recorded during the RPR selection experiment (Martin et al., 2004). This approach makes our analysis more comparable to experimental evolution studies in model species (Kofler and Schlötterer, 2014) than to most selection mapping studies in crops (e.g., Zhou et al., 2015; Gage et al., 2018). In contrast to wild populations (Plassais et al., 2019) or domesticated populations studied over an evolutionary timeframe (Hufford et al., 2012; Beissinger et al., 2016), wherein the accurate demographic parameters are unknown, knowledge of such parameters in our study make it straight forward to establish statistical significance via no-selection simulations.

Significant evidence of selection for a large number of SNPs, 3656 for C15-H and 3412

for C15-L, indicates a major change in the genome landscape of the selected population

compared to C0 population. Our observations implicating a large number of significant SNPs are

consistent with those observed in experimentally evolved populations of model species including

drosophila (Turner et al., 2011; Turner and Miller, 2012) and yeast (Araya et al., 2010).

Discovery of such a large number of SNPs results, in part, from the fact that the resolution and

power of such studies depend heavily on levels of linkage disequilibrium before selection

commenced (Kofler and Schlötterer, 2014). Moreover, a SNP with significantly altered allele

frequency due to selection is not necessarily causal for the trait of interest, and may instead be

linked to a causal site through a phenomenon known as hitchhiking (Smith and Haigh, 1974).

Although our analysis showed significant enrichment for SNPs selected in the high and

low RPR directions, the number of overlapping SNPs (110) was a small fraction of the total

number of significant SNPs. There are several possible reasons, both practical and biological, for

this outcome. First, selection mapping evaluated by quantification of the change in allele

frequency, as implemented here, is most powerful to identify SNPs that have moved from a low

frequency to a high frequency due to selection, or vice versa (Vlachos and Kofler, 2019).

Therefore, even when selection operated on the same SNP in both populations, our study may

only have the power to detect the SNP the population where the frequency of the SNP was

significantly altered from C0. Second, our drift simulation approach for declaring significance

included only single SNP tests and, therefore, it is possible that different SNPs associated with

high and low RPR in the same genomic region. Indeed, manual inspection of significantly

selected SNPs in either direction identifies several hotspots of SNPs in the high and low RPR

that may correspond to selection on the same underlying gene (Figure 7). Third, the relatively

small amount of overlap may indicate that distinct biological processes underlie high and low

RPR. This argument is supported by the enrichment of significant SNPs in the lignin genome

space in the high RPR direction and enrichment in the polysaccharide genome space in the low

RPR direction.

Transcriptomic analysis reveals both expected and novel process related to RPR

Consistent with large changes in the genome landscape indicated by selection mapping, high

levels of biological activity associated with RPR is evident from differential expression of 21%

of the annotated maize genes. Global transcriptome analysis showed that the high and low RPR

phenotypes start to diverge very early during the development of an internode. While

differentiation of secondary cell wall has been considered as the major contributor to strength,

such differentiation happens after the internode elongation is complete (Cosgrove and Jarvis,

2012; Zhang et al., 2019). Therefore, high transcriptome activity at early stages indicates that

differences in RPR are not limited to secondary cell wall biosynthesis. Furthermore, over 2000

and 3000 unique DE genes associated with high and low RPR, respectively, outnumber 845

genes associated with primary and secondary cell wall synthesis. While this gap will shrink with

the discovery of additional genes related to cell wall synthesis, and cell wall is one of the most

important determinants of RPR, identification of novel biological activities and processes will

further elucidate the determination of rind strength.

GO enrichment based on DE genes provides a global view of the key processes and

activities involved in the determination of rind strength. In the internodes of high RPR inbred,

Hrpr1, synthesis of secondary metabolites is a prominent process as indicated by enrichment of

phenylpropanoid biosynthetic processes. Phenylpropanoid pathway is responsible for a large

number of natural plant products including lignin, coumarins, flavonoids, anthocyanins, phenolic

acids, and stilbenes (Vogt, 2010). Besides providing resistance to various biotic and abiotic

stresses, these compounds are needed for reinforcing the cell wall and, therefore, providing

mechanical strength to plants (Dixon et al., 2002; Vogt, 2010; Fraser and Chapple, 2011; Gray et

al., 2012). Enrichment of quinolinate metabolism indicates increased synthesis of NAD, which is

known to facilitate lignification and oxidative cross linking of lignin and polysaccharides in the

cell wall (Pétriacq et al., 2012; Higuchi et al., 2014). Enrichment of amino acid transport

suggests an important albeit less understood role of this process in the determination of RPR.

Amino acids are involved in the synthesis of cell wall protein and enzymes, and act as the

precursors for monolignol synthesis (Maeda and Dudareva, 2012).

In the transcriptome of low RPR inbred, Lrpr1, enrichment of xylan and glucan

metabolism supports higher flux of photoassimilates towards polysaccharides which, together

with lower lignification, explains higher recovery of cell wall polysaccharides. Enriched

metabolism of amino sugars indicates a higher accumulation of glycoproteins that may be

important for providing mechanical strength in the absence of optimal lignification. Amino

sugars are important components of chitin and recent studies suggest their role in the

determination of plant architecture (Vanholme et al., 2014). Enrichment of auxin signaling, and

response to ethylene, brassinosteroids (BR), jasmonic acid, and gibberellic acid in Lrpr1

internodes indicates important, yet sparsely understood, role of hormones in the determination of

RPR. In Arabidopsis, BR are known to regulate cellulose synthesis, and defects in BR synthesis

and signaling result in impaired cellulose accumulation in the cell walls (Xie et al., 2011;

Sánchez-Rodríguez et al., 2017). Auxin concentration is linked to cell expansion during cell wall

synthesis and thereby affects the mechano-chemical aspects of plant development (Braybrook

and Peaucelle, 2013; Paque et al., 2014; Lehman et al., 2017; Majda and Robert, 2018).

Gibberellic acid has also been found to regulate cell wall extension and, interestingly, enhance

the lignification of xylem (Bai et al., 2012; Falcioni et al., 2018).

Selection on high and low RPR operated on distinct gene sets

Selection mapping, transcriptomic, and metabolic analyses indicate that plants with high and low

RPR employ distinct approaches to partitioning photoassimilates towards cell wall biosynthesis.

The annotation of the cell wall genes and examination of their enrichment during divergent

selection revealed that selection preferentially acted on polysaccharide synthesis genes during

selection for low RPR and lignin biosynthetic genes during selection for high RPR. This altered

genome landscape is well reflected in the transcriptional activity and chemical composition of

the internode cell walls. The enrichment of the lignin pathway during selection for high RPR

phenotype is consistent with elevated expression of lignin biosynthetic genes and the higher

lignin content in Hrpr1. The enrichment of polysaccharides pathway during selection for low

RPR is consistent with lower lignification and an increased amount of extractable sugars from

the cell wall in Lrpr1.

Chemical composition of rind cell walls, which generally refers to lignin and

polysaccharides, is an important determinant of stalk strength (Zuber et al., 1980; Appenzeller et

al., 2004). However, whether these constituents impart a positive or negative impact on stalk

strength is debated (Sekhon et al., 2019). Cellulose has been proposed to either positively

associate with stalk strength or to be no consequence (Appenzeller et al., 2004; Ye et al., 2016).

Reduction in lignin content, observed in the maize brown midrib (bmr) mutants, has been

associated with decreased stalk strength (Miller et al., 1983). Study of a monocot-specific

microRNA (ZmmiR528) also showed that increased lignification is positively associated with

improved RPR (Sun et al., 2018). The current study clearly indicates the opposing effect of the

accumulation of lignin and polysaccharides on RPR. Attempt to relate the gross amount of

polysaccharides and lignin with RPR or stalk strength, however, is an over-simplification of a

complex structural phenomenon. The microstructure of these components and the effect of

interactions between these constituents must be considered to understand their effect on stalk

strength.

While considering the lignification of secondary cell walls, the emphasis is often placed

on the content of the three major monolignols, and the processes involved in the polymerization

of these lignin monomers is overlooked (Tobimatsu and Schuetz, 2019). Detailed metabolic

analysis of Hrpr1 and Lrpr1 internodes highlights the role of additional phenolics in cell wall

lignification especially during the early stages of internode development. Tricin, a monolignol

that provides nucleation sites for incorporation of the three major monolignols and initiates

lignification of grass cell walls (Lan et al., 2015), emerged as an important determinant of

lignification associated with RPR. High tricin content would result in more lignin chains, and

interaction of these chains with each other and with other molecules (e.g. hemicellulose) would

result in higher rind strength. Consistent with this notion, higher FA and pCA content associated

with high RPR further indicates that higher cross-linking, and polymerization of lignin and other

constituents is important for high rind strength.

A roadmap to enhanced mechanistic understanding and exploitation of RPR for improving

lodging resistance

While 'stand-alone' omics investigations, including those looking at genome or transcriptome

landscapes underlying a complex trait, provide useful information about the underlying biology

and genetics, results from such studies are also often riddled with false positives and false

negatives. Integrating selection mapping and transcriptome analyses with previously reported

QTL and GWAS studies provides a more compressive view of the genetic architecture of RPR

and allows the identification of a core set of candidate genes (Figure 7). Several genomic

hotspots detected by examining selection signatures in the divergently selected RPR populations

that co-localize with previously reported regions likely harbor large effect genes. Conversely,

regions exclusively detected by selection mapping capture the allelic variation present in the

inbred lines used to develop the base (C0) population.

Many of the promising candidate genes highlighted by the integrated analysis are
associated with cell wall biology. For instance, both selection mapping and transcriptome
analyses identified a region on chromosome harboring two genes (Zm*PAL7*, Zm*PAL9*) encoding
for phenylalanine ammonia lyase which catalyzes the first reaction of the phenylpropanoid
pathway and regulates the flux towards the synthesis of all secondary metabolites (Vanholme et
al., 2010). Impairment of phenylpropanoid biosynthesis in *pal1* and *pal2* mutants, leads to a
reduced amount of lignin and decreased cell wall strength in Arabidopsis (Rohde et al., 2004).
Low RPR phenotype and low lignin content are associated with reduced expression of *ZmPAL7*
and *ZmPAL8* (Sun et al., 2018). Likewise, a candidate transcription factor, ZmMYB31, on
chromosome 2 has been established as a regulator of lignin biosynthesis in maize (Fornalé et al.,
2006). Overexpression of *ZmMYB31* in Arabidopsis reduced the lignin biosynthesis but did not
alter the composition of the lignin polymer (Fornalé et al., 2010). Of multiple candidate genes
present on chromosome 9, a MADS-box transcription factor encoded by *ZmMADS1* has been
implicated in auxin transport and signaling (Khanday et al., 2013) further supports the role of
auxin in the determination of RPR. Putative orthologs of two other candidate genes, *ZmNAC25*
and *ZmEXPA2*, have been shown to be involved in cell expansion in Arabidopsis through
GA/DELLA-NAC25/NAC1L-EXPA2 regulatory network (Sánchez-Montesino et al., 2019).
Given that GA signaling has multiple regulatory effects on cellulose synthesis (Huang et al.,
2015; Felipo-Benavent et al., 2018; Sánchez-Montesino et al., 2019), the role of *ZmNAC25* in
RPR warrants detailed investigations. Finally, while some of the genes identified in this study
can be directly or indirectly linked with some aspects of stalk strength, many lacks such
association due to absence of experimental evidence. These genes have the potential to provide a
wealth of information related to the biology of stalk strength in maize and related grasses.

Conclusions

Despite technological advances, progress in deciphering the genetic architecture of
complex traits through omics approaches has been slow. In the case of stalk lodging resistance, a
trait impacted by a number of internal and external factors, lack of reliable and scalable methods
for phenotypic evaluation of the germplasm adds additional challenges. To this end, systematic
analysis of populations divergently selected for RPR, a reliable predictor of lodging resistance,
offers a unique approach to generate novel insights into the molecular genetic mechanisms

governing stalk strength. Combining the results from selection mapping with those from the

current state-of-the-art approaches provide higher quality candidate mechanism and genes that

will guide functional studies to understand the molecular architecture of RPR and stalk strength.

Furthermore, the valuable information generated by this study will boost efforts for genetic

improvement of stalk strength in maize, sorghum, and other grasses.

MATERIALS AND METHODS

Genetics stocks

The MoSCSSS C0 population, hereafter called C0, was derived from 14 inbred lines (A657,

A632Ht, B14AHt, B37Ht, B68, B73, B76, B84, CM105, H84, N28Ht, N104, Oh514, and

Pa864P) that were either direct derivatives or closely related to the Iowa Sty Stalk Synthetic

heterotic group (Martin et al., 2004). Populations derived from 15 cycles selection for high and

low RPR by Dr. Larry Darrah and colleagues at the University of Missouri (Figure 1A) along

with the C0 population were kindly provided by Dr. Sherry Flint-Garcia. The high and low

populations, originally named as MoSCSSS H24 (High Rind Penetrometer [HRP]) C15 and

MoSCSSS H25 (Low Rind Penetrometer [LRP]) C15 were renamed as C15-H and C15-L,

respectively (Figure 1). Each of these three populations was multiplied separately by making 250

unique crosses within the population such that each of the 500 parent plants was either used as a

male or female only once, and by bulking 100 seeds from each of the resulting crossed ears.

For the development of inbred lines, one randomly chosen plant each from C15-H and

C15-L populations was self-pollinated. A single row of 15-20 plants was grown from each

resulting ear, and one randomly chosen plant from each row was again self-pollinated. This

process was repeated six times to attain near-homozygosity. Seed from each ear was then used to

multiply seed for each of the inbred lines. The inbred lines derived from C15-H and C15-L

populations were designated as Hrpr1 and Lrpr1, respectively.

Measurement of rind penetrometer resistance

For phenotypic analysis, the Hrpr1 and Lrpr1 inbred lines were grown in a randomized complete

block design with three replications at Clemson University Calhoun Field Laboratory, Clemson,

SC in summer 2017. In each block, a four-row plot of each inbred line was grown with row

length and row-to-row distance being 4.57 meters and 0.762 meters, respectively, resulting in a

total pot size of 13.93 m^2. Data was recorded on 12th internode as, on an average, this internode

lies below the primary ear-bearing node in these inbred lines. As each maize leaf is attached to a

node, leaves were used to identity the 12th internode. Since the juvenile leaves senesce and

become undetectable at later developmental stages, at the vegetative 4 (V4) stage marked by the

presence of four fully extended leaves, a hole was punched in the fifth leaf and this leaf was

subsequently used to identify the 12th internode. The development of the 12th internode was

visually followed by dissecting plants every 3 days. The phenotypic data and sample collection

were started at the V8 stage attained 45 days after sowing (DAS) when the 12th internode was

approximately 0.5 cm long. The samples were collected at a three-day interval. RPR was

measured with a 2 mm diameter probe with flat tip (49FL81, Grainger Inc., Lake Forest, IL)

installed on an Imada® Digitial Force Gauge (ZTA-DPU). A steel plate covering the base of the

probe was installed on the force gauge to prevent secondary contact after the initial puncture. For

the C0, C15-H, and C15-L populations, RPR was recorded on the internode below the primary

ear-bearing node after careful removal of the leaf sheath. The puncture test was performed in the

center of the internode, perpendicular to the minor axis of the stalk cross-section (i.e., in the

direction of the major axis of the stalk cross-section).

Genotyping-by-sequencing and SNP calling

Individual leaf samples from 96 plants of each of the three populations (C0, C15-L, and C15-H)

were collected from field-grown plants at the V8 stage, flash-frozen in liquid nitrogen, and stored

at -80°C. DNA of each plant was extracted using the cetyl(trimethyl)ammonium bromide

(CTAB) method (Saghai-Maroof et al., 1984). DNA was cleaned using a modification of the

Qiagen DNeasy 96 Plant kit (Qiagen, Germantown, MD), diluted to 20ng/µl, and approximately

200ng DNA was used for modified *Ape*KI genotyping-by-sequencing (GBS) (Elshire et al.,

2011). The modification included separating each 96 well plate into 4 pools of 24 adapter-ligated

samples. Barcodes were designed on DeenaBIO and synthesized by IDT (Coralville, Iowa, USA.

Pre-polymerase chain reaction of the pooled sample and PCR amplification was done using

ThermoFisher Phusion II master mix. Quantification of each enriched library pool was done by

Qubit (Invitrogen, Carlsbad, CA) and distribution was analyzed on Agilent Bioanalyzer (Agilent

Technologies, Santa Clara, CA) high sensitivity DNA chip. All 288 individually barcoded

samples were pooled. NextSeq High Output single-end 75-bp sequencing was performed at

University of Missouri, Columbia DNA Core (https://dnacore.missouri.edu/).

SNP calling, filtration, and estimating allele frequency

Sequenced reads were aligned to the maize B73 reference genome version 4 (AGPv4) (Jiao et

al., 2017) using the Tassel 5 GBS v2 pipeline (Glaubitz et al., 2014). This alignment resulted in a

total of 219,758 initial SNP markers with the mean read depth of ~2.6x. These SNPs underwent

very stringent filtering due to the fact that GBS often yields a high proportion of missing data

(Beissinger et al., 2013). First, only SNPs with two alleles at a particular locus were included.

Individuals were then filtered to remove SNPs with greater than 40% of markers missing,

resulting in the removal of 37 individuals from C0, 15 individuals from C15-H, and 31

individuals from C15-L population. Furthermore, SNPs missing at greater than 50% of

observations were also filtered from each population. The resulting SNPs were further filtered to

remove low confidence SNPs with read count less than 10. After filtering, 159,849 SNPs were

retained from 205 individuals for the downstream analysis. Reference allele frequency was

computed in each population using VCFtools (Danecek et al., 2011).

Detection of selected loci

Two techniques were used to identify putatively selected regions of the genome. We used a

pairwise estimate of F_{ST} calculated in R version 3.4.1 (R Core Team, 2013) based on a published

method (Weir and Cockerham, 1984) and the published analysis pipeline (Beissinger et al.,

2014), such that

$F_{ST} = \dfrac{\sigma^2}{\bar{p}(1-\bar{p})+\frac{\sigma^2}{n}}$,

where σ^2 is the sample variance between the populations, $\bar{p}$ is the average allele frequency

between two populations, and n is the number of populations. For every locus, a single F_{ST} value

was computed.

F_{ST} depends on initial allele frequency (Jakobsson, Edge, & Rosenberg, 2013) and cannot

alone be used to generate significance thresholds (Akey, 2009). Therefore we implemented a

simulation-based test that incorporated the demographic history of the C15-H and C15-L

populations to allow for significance testing. Our strategy involved comparing the observed

allele frequency divergence at each site to the expected magnitude of genetic drift over the

course of the experiment. We simulated a drift-only breeding program based on selection for

high and low RPR and evaluated the distribution of allele frequencies after breeding. In our

simulations, breeding proceeded for 15 generations with 60 males and 120 females mated each

generation in agreement with the scheme used for the development of these populations (Martin

et al., 2004). Furthermore, our simulation strategy incorporated sampling of 96 individuals from

generation 0 (G0) and G15 for genotyping. To implement these simulations, first we randomly

sampled 96 individuals from a pool with allele frequency set between 0.01 and 0.99 with

increments of 0.01. This sampling was conducted 1 million times for each initial frequency.

Next, the frequency of each random sample was used as an initial frequency for 15 generations

of drift simulations, with 60 contributing males and 120 contributing females each generation.

The frequencies at the end of G15 were then used for a final sampling of 96 individuals. We used

the final simulated frequencies to calculate the 1e-7 and 1-1e-7 quantiles of allele frequency

change. Then, we compared the observed pre-selection allele frequencies in C0 to the post-

selection allele frequencies in C15-L and C15-H to identify SNPs that changed in frequency

more than that expected under drift and sampling alone in each population. SNPs with minor

allele frequency below 0.01 in C0 were not included in this analysis resulting in 158,196 SNPs.

The SNPs with allele frequency change greater than the 1e-7 and 1-1e-17 quantiles are putative

candidates for selection, with a 1e-7 false positive rate. Since we evaluated 158,196 SNPs, the

experiment-wide false positive rate was fewer than one SNP. Further details, including our

simulation scripts, are provided as Supplemental Scripts.

Annotation of cell wall genes

The cell wall related protein sequences involved in the biosynthesis of cellulose, hemicellulose,

and lignin reported in an earlier study (Guillaumie et al., 2007) were used as query in BlastP (e-

value 10^{-5}) against the latest maize ensemble protein database (version B73_RefGen_v4) to

identify complete gene families. The top hit sequences were confirmed and annotated by

searching against the Conserved Domains Database

(www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi) and NCBI non-redundant database using e-

value of $10^{-5.}$ The confirmed cell wall genes were analyzed for their associated SNPs enrichment

during divergent selection for low and high RPR.

Testing for the enrichment of selected SNPs within lignin and polysaccharide synthesis genes

A resampling analysis was conducted to test the hypothesis that subsets of genes related to the

lignin and the polysaccharide biosynthetic pathways experienced elevated levels of selection as

compared to the rest of the genome. For this test, the number of significant SNPs within the total

genome space of annotated genes in lignin and polysaccharide biosynthetic pathways were

counted. The genome space was defined as positions from 5 kb upstream of the AGPv4 predicted

transcription start position to 5 kb downstream of the predicted transcription stop position of the

lignin and polysaccharide biosynthetic pathways. All SNPs within these spaces were designated

as potentially related to lignin or polysaccharide synthesis, respectively, and subjected to

enrichment analysis. Separate counts of the number of lignin and polysaccharide synthesis

associated SNPs were calculated for selection in the high and low RPR directions. Then, 1,000

permutations were conducted, where for each permutation, SNP p-values (calculated based on

allele frequency change) were shuffled over all SNPs positions to generate a set of *permuted-*

significant SNPs. For each permutation, the number of *permuted-significant* SNPs within the

genome space corresponding to the lignin and the polysaccharide pathway was counted. Based

on these permutations, we computed 95% and 99% confidence intervals for the number of

significant SNPs that should fall within each genome space if it was not enriched for significant

SNPs. We compared the permutation distribution to the observed number of significant SNPs in

the lignin and polysaccharide pathway genome space to identify one-sided p-values for

enrichment.

Histochemical staining

Mature maize stems were hand sectioned with a razor blade and the cross-sections were stained

in phloglucinol-HCl satin following a published protocol (Ruzin, 1999). Samples were imaged

with a Zeiss Axioplan2 imaging microscope.

Tissue collection for metabolic analysis and RNA sequencing

The 12[th] internode used for measurement of RPR was used for metabolic analysis. All samplings

were performed between 8:00 AM and 12:00 PM. At each stage and for each inbred line, three

biological replicates of each tissue were collected. A biological replicate consisted of pooled

tissue from two different, competitive (non-border and well-spaced from neighbors) plants

harvested from the same block and dissected to separate the 12[th] internode. Samples were

chopped into small pieces, flash-frozen in liquid nitrogen, and stored at -80°C.

Quantification of polysaccharides

For metabolic analyses, the samples were ground to very fine powder under liquid nitrogen with

a cryogenic grinder (6875 Freezer/Mill, Spex SamplePrep, Metuchen, NJ) using a 45 second

pulse followed by 30 seconds resting period and another 45 second pulse. The resulting fine

ground powder was transferred to 15 ml tubes, freeze-dried, and used for various metabolic

analyses. The glucose and pentose monomer in cell wall were measured according to a published

protocol (Santoro et al., 2010) with modifications described previously (Sekhon et al., 2016).

Phenolics/lignin extraction

Sequential extraction was used for the separation of phenolic compounds according to published

protocol (Li et al., 2015) with modifications. The tricin was extracted with methanol, the

ester/ether-bound phenolics were recovered following a mild-base hydrolysis at high

temperature, and monolignols were extracted after oxidative degradation of lignins at a higher

temperature and pressure in the presence of CuO (Hedges and Ertel, 1982). For quantifying the

extractable phenols, 500 mg of freeze-dried tissues was extracted with 2.5 ml of methanol for 3 h

with shaking at room temperature. After centrifugation at 1500 g for 15 min, 1.5 ml of

supernatant was transferred to a glass vial and stored at -20°C for tricin quantification. The

remaining pellet was washed twice with 2.5 ml of methanol, dried overnight and, hydrolyzed

with 6 ml of 1M NaOH in a glass tube, sparged with Ar, and incubated at 90°C for 3 h in the

dark. After cooling on ice, the supernatant was used for the analysis of ester/ether-bound

phenolics. The pH of the supernatant was reduced to < 2 using 50% HCl, and the resulting

precipitate was discarded after centrifugation. The solution in the tubes was extracted with 2 ml

of ethyl acetate and ethyl acetate layer was transferred into GC vials and stored at −20°C for

further analysis.The residual pellet was washed twice with 5 ml of deionized water, dried, and

stored at -20°C for lignin analysis.

The lignins were depolymerized in 23 ml Acid Digestion Vessels (model 4749 Parr,

Instrument Co., Moline, IL, USA) (Kaiser and Benner, 2012). 500 mg of CuO, 75 mg

$Fe(NH_4)_2(SO4)_2 \cdot 6H_2O$ and 5 ml of freshly prepared 2 M NaOH (pre-sparged with Ar for 30 min)

was combined with the base hydrolyzed pellet in Teflon cups. The vessel was sealed and

incubated at 155°C for 160 min and then immediately cooled in an ice bath to room temperature.

The digestion mixture was transferred to fresh tubes and 50 µl of ethyl vanillin (400 mg l^{-1}) was

added as an internal standard. The pH of the solution was reduced to < 2 by adding of H_2SO_4 and

mixed gently. After centrifugation, 8 ml of the supernatant was transferred to a clean glass tube,

and the lignin-derived phenols were extracted by liquid–liquid partitioning with 2 ml of ethyl

acetate.

Tricin analysis using LC-MS/MS

The methanol extracts were profiled using a quadrupole-orbitrap-iontrap mass spectrometer

(Orbitrap Fusion™ Tribrid™ Mass Spectrometer; Thermo Fisher Scientific, Waltham, MA,

USA). Compounds were separated on a C18 reversed-phase column (ACQUITY HSS T3, 150 x

2.1 mm, 2.6 μm) using an UHPLC (Dionex Ultimate 3000), with gradient elution employing

binary solvent system involving 0.1% formic acid and acetonitrile. The gradient of acetonitrile

mix was increased from the initial proportion of 5% to 95% over 25 min and held for 5 minutes

before re-equilibrating the column to the initial solvent condition. The solvent flow rate was

maintained at 0.15 mL/min, and the column was maintained at 35°C. The samples were

introduced into the mass spectrometer through a heated electrospray ionization (H-ESI) interface

operated in a negative ionization mode. The Orbitrap Fusion mass spectrometer was operated in

high-resolution (120,000 FWHM at 200 m/z) full-scan mode (120-1000 m/z). The optimized

parameters were set as follows: sheath gas 60, auxiliary gas 15, sweep gas 1, spray voltage 3.5

kV, probe temperature 350°C, and transfer capillary temperature 300°C. The fragmentation was

achieved via collision-induced dissociation with collision energy set at 35%. From the full

spectral scans, the extracted ion chromatograms of tricin were derived using theoretical exact

masses of [M-H]- ion with 4 significant figures and a scan width of ± 1.0 ppm mass error.

LC-MS Quantitation of Lignin Phenols

The phenolic monomers resulting from the base hydrolysis of bound phenolics and CuO

oxidation of lignins were measured using liquid chromatography coupled to a triple quadrupole

mass spectrometer (LC-MS/MS Shimadzu LC-20AT + Shimadzu 8040 mass analyzer; Kyoto,

Japan). Ethyl acetate partitioned samples were dried under nitrogen gas stream and re-dissolved

in methanol (300 μl volume maintained) for LC-MS/MS analysis. Compounds were separated on

a C18 reversed-phase column (Kinetex XB-C18, 150 x 3 mm, 2.6 μm) using a binary solvent

gradient. Solvent A was water containing 0.1% formic acid and solvent B was acetonitrile. The

solvent flow rate was maintained at 0.3 mL/min and the gradient of acetonitrile was increased

709 from the initial proportion of 25% to 95% over eight minutes on a linear gradient and held for

710 one minute before re-equilibrating the column to initial solvent condition. The samples were

711 introduced into the mass spectrometer through an electrospray ionization (ESI) interface

712 operated in a negative ionization mode. The ESI interface parameters were set as follows:

713 nitrogen nebulizing gas 3 L/min, nitrogen drying gas 12 L/min, capillary voltage -3.5 kV, heat

714 block 400°C, and desolvation line temperature 250°C. The triple quadrupole mass spectrometer

715 was operated in MS/MS mode based on the following multiple reaction monitoring parameters

716 that were optimized using authentic standards: p-coumaric acid (162.8 > 119.1, 92.95, 117.1),

717 dihydroxybenzoic acid (152.9 > 109.05, 65, 67.05), ferulic acid (192.9 > 134.1, 178.1, 149.2), p-

718 hydroxybenzoic acid (136.9 > 93.1, 64.95, 41.1), p-hydroxybenzaldehyde (121 > 92.05, 93,

719 41.05), p-hydroxyacetophenone (134.95 > 92.05, 93, 120.05), syringic acid (196.85 > 182.1,

720 123.1 95), syringaldehyde (180.85 > 166.1, 151.1, 123), acetosyringone (194.9 > 180.2, 165.1,

721 137), vanillic acid (166.85 > 152.05, 108.05, 123.1), vanillin (150.95 > 136.1, 92.05, 108.05),

722 and acetovanillone (164.95 > 150.05, 122.05, 79). Quantitation was performed using a series of

723 13-point external calibration curve ranging from 12 ppb to 12.5 ppm containing a mixture of all

724 12 compounds.

725 *RNA Sequencing*

726 RNA was extracted from the frozen chopped internode samples using TRIzol reagent

727 (Invitrogen, Carlsbad, CA, USA) following the manufacturer's protocol. The mRNA isolation,

728 cDNA synthesis, and library preparation were performed by Novogene Corporation (Chula

729 Vista, CA) following the standard Illumina (Illumina Inc., San Diego, CA) protocol. The 150 bp

730 paired-end reads generated after sequencing by Illumina HiSeq platform were trimmed of low-

731 quality bases and adapter sequences with the trimmomatic (Bolger et al., 2014). Reads were

732 aligned to reference maize genome (Zea_mays.AGPv4.40.gtf) (Jiao et al., 2017) with the

733 Tophat2 v. 2.1.1 (Trapnell et al., 2012) using Bowtie2 (v. 2.3.4.1) (Langmead and Salzberg,

734 2012). Cufflinks v.2.2.1 was used to determine Fragments per kilobase per million (FPKM)

735 (Trapnell et al., 2012). The read count was determined with the HTSeq-count v.0.10.0 (Anders et

736 al., 2015). The differential expression between different pairs of stages and genotypes (Hrpr1 and

737 Lrpr1) were calculated using DESeq2 (Love et al., 2014). Genes with $p \leq 0.05$ were corrected by

738 the false discovery rate (FDR) < 0.05 (Benjamini and Hochberg, 1995). Genes with $\log_2$ fold

change > 2 and FDR corrected $p < 0.05$ were selected for subsequent analyses. Gene ontology

(GO) enrichment was carried out with the GOSeq software tool (Young et al., 2010) and

enriched categories were filtered by FDR corrected $p < 0.05$.

Integrating of selection mapping high priority SNPs and gene expression

The top 5% F_{ST} SNPs for high and low RPR were mapped to nearby genes using the SNP-to-

gene mapping tool implemented in Camoco (v0.6.3) (Schaefer et al., 2018), using the parameters

of selecting up to 2 flanking genes within 100 kb windows both upstream and downstream of

SNPs. Candidate genes were filtered to only genes that were statistically significant and at least

2-fold differentially expressed between Hrpr1 and Lrpr1 internodes. Gene ontology (GO)

enrichment was performed using the hypergeometric statistic from utilities within Camoco to

identify the putative biological function of DE genes near extreme F_{ST} SNPs.

Analysis scripts

All scripts used for selection signature analysis are available as supplemental files. Drift

simulations were performed using scripts Supplemental Script 1 and Supplemental Script 2 that

were extracted from beissingerlab.github.io and originally published elsewhere (Lorenz et al.,

2015). Significance thresholds from drift simulations were compiled using Supplemental Script

3. Supplemental Script 4 was used for SNP calling and allele frequency calculations. F_{ST} was

calculated using Supplemental Script 5 that was extracted from beissingerlab.github.io/Software/

and originally published elsewhere (Beissinger et al., 2014). Supplemental Script 6 was used to

compare changes in allele frequency to drift simulations to determine significance, to conduct

bootstrap analysis of the number of overlapping SNPs selected by chance, to perform enrichment

analysis testing for selection on polysaccharide and lignin genes, and to visualize the meta-

analysis (Figure 7).

Accession Numbers

All sequence reads are available at Sequence data from this article can be found at the National

Center for Biotechnology Information Sequence Read Archive

(https://www.ncbi.nlm.nih.gov/sra) under accession number PRJNA623937

(https://dataview.ncbi.nlm.nih.gov/object/PRJNA623937?reviewer=su7vgercui58lfpvquk278b13

t)

ACKNOWLEDGMENTS

Authors thank Dr. Sherry Flint Garcia for providing the divergently selected RPR populations,

Kate Guill for genotyping, Jackyung Yi, Arlyn Ackerman, and a number of talented

undergraduate students for help in the collection of data and samples. Funding for this study was

provided by the National Science Foundation grant #1826715 to R.S.S. Support for SNP

genotyping included resources from the USDA-ARS, Plant Genetics Research Unit.

Table 1. Identification of maize cell wall synthesis gene families.

Cell wall component/process	Gene family	Number of genes
Polysaccharides biosynthesis (cellulose and hemicellulose)	*Cellulose synthase* (*CesA*/*Csl*) superfamily	83
	GT8 superfamily (galacturonosyl transferases)	57
	GT47 superfamily (galactosyl, arabinosyl, xylosyl, and glucuronosyl transferases)	54
	GT31 superfamily (galactosyl transferases)	32
Cell wall assembly and rearrangement	*Expansin* gene family	95
	Xyloglucan endotransglucosylase/hydrolase (*XTH*) superfamily	35
Cell wall modification and disassembly	*Polygalacturonase/ Pectin lyase* (*PGases*) gene family	53
	Pectinesterase (*PME*) gene family	48
Nucleotide-sugar synthesis and interconversion	*UDP-Glc dehydrogenase* (*UGD*), *UDP-GlcA decarboxylase*, *UDP-Glc 4-epimerases* (*UGE*), *UDP-Xyl 4-epimerases* (*UXE*), *4,6-dehydratase* (*GMD*)	47
Lignin biosynthesis	*Phenyl ammonia lyase* (*PAL*) gene family	18
	Caffeic acid o-methyltransferase (*COMT*) gene family	32
	Caffeoyl-CoA O-methyltransferase (*CCoAOMT*) gene family	5
	4-Coumaric acid CoA-ligase (*4CL*) gene family	17
	Cinnamoyl-CoA reductases (*CCR*) gene family	36
	p-Coumaroyl-shikimate/quinate 3'-hydroxylase (*C3'H*), Cinnamate 4-hydroxylase (*C4H*), *Ferulate 5-hydroxylase* (*F5H*) gene family	47
	Cinnamyl alcohol dehydrogenase (*CAD*) gene family	8
	Hydroxycinnamoyl-CoA transferase (*HCT*) gene family	42
	Peroxidase gene family	114
	Laccase gene family	22
Total genes		845

786 **Table 2.** Key genes related to rind penetrance resistance identified from the metanalysis. Criteria

787 refer to the four criteria defined in the text that were used to identify candidate genes.

RPR	SNP position	F_{ST}	Gene ID	Chr	Gene annotation	Fold change	Criteria
High	202,580,532	0.63	Zm00001d006236	2	*ZmMYB31*	2.6	1, 2, 3
High	197,606,711	0.81	Zm00001d006082	2	*PIN-formed protein11 (pin11)*	4.9	1, 2, 3
High	210,185,759	0.58	Zm00001d006511	2	*Glucose-6-phosphate/phosphate translocator 2 (gpm334)*	3.0	1, 2, 3
High	211,751,720	0.91	Zm00001d006591	2	*F-box domain containing protein*	-3.9	1, 2, 3
High	212,878,293	0.59	Zm00001d006627	2	*L-type lectin-domain containing receptor kinase IX.1*	5.6	1, 2, 3
High	213,548,099	0.64	Zm00001d006653	2	*AP2-EREBP transcription factor (ereb62)*	4.5	1, 2, 3
High	218,576,041	0.67	Zm00001d006883	2	*Calmodulin binding protein*	-2.6	1, 2, 3
Low	185,589,850	0.52	Zm00001d005741	2	*NAD(P)H-quinone oxidoreductase subunit L*	6.9	1, 2, 3
Low	186,080,069	0.80	Zm00001d005749	2	*ZmWRKY55*	-4.6	1, 2, 3
Low	189,506,978	0.92	Zm00001d005813	2	*Protein TIFY 10B (zim13)*	4.5	1, 2, 3
High	153,354,929	0.88	Zm00001d042143	3	*Glucan endo-1,3-beta-glucosidase homolog1 (geb1)*	-4.9	1, 2, 3
High	212,853,636	0.82	Zm00001d043881	3	*Digalactosyldiacylglycerol synthase 1*	24.2	1, 2, 3
High	213,845,025	0.87	Zm00001d043921	3	*ZmNAC82*	-3.8	1, 2, 3
High	214,419,702	0.76	Zm00001d043947	3	*Lipid phosphate phosphatase 2*	7.8	1, 2, 3
Low	193,474,769	0.86	Zm00001d043257	3	*G-type lectin S-receptor-protein kinase CES101*	2.5	1, 2, 3
Low	218,247,550	0.83	Zm00001d044065	3	*DELLA protein RGA (gras46)*	-2.6	1, 2, 3
High	39,730,201	0.77	Zm00001d049678	4	*ZmNAC115*	-5.5	1, 2, 3
High	155,986,150	0.81	Zm00001d051361	4	*VAN3-binding protein*	-4.9	1, 2, 3
High	169,990,594	0.52	Zm00001d051800	4	*Caleosin protein*	-2.5	1, 2, 3
High	191,466,721	0.60	Zm00001d017275	5	*Phenylalanine ammonia lyases (ZmPAL7/ZmPAL9)*	-2.5	1, 2, 3, 4
High	146,400,206	0.56	Zm00001d038049	6	*Lichenase (lic1/lic2)*	-6.4	1, 2, 3
Low	151,942,963	0.67	Zm00001d038221	6	*ZmNAC20*	4.9	1, 2, 3
Low	164,388,708	0.90	Zm00001d038791	6	*Receptor-like protein kinase HSL1*	2.4	1, 2, 3
Low	165,372,185	0.88	Zm00001d038831	6	*Type IV inositol polyphosphate 5-phosphatase*	3.4	1, 2, 3
Low	166,774,503	0.89	Zm00001d038905	6	*putative beta-14-xylosyltransferase IRX10L*	2.1	1, 2, 3
Low	163,561,967	0.85	Zm00001d021818	7	*ZmNAC25*	-6.9	1, 2, 3
High	103,637,970	0.81	Zm00001d046725	9	*D-mannose binding lectin receptor-like protein kinase*	-3.8	1, 2, 3
High	104,342,643	0.83	Zm00001d046748	9	*Putative polyphenol oxidase family protein*	-6.0	1, 2, 3
High	155,402,586	0.66	Zm00001d048389	9	*Xyloglucan galactosyltransferase MUR3*	3.6	1, 2, 3, 4
Low	111,179,135	0.57	Zm00001d046928	9	*Histidine kinase5 (hk5)*	-2.4	1, 2, 3
Low	154,806,326	0.83	Zm00001d048344	9	*Expansin-like A2 (ZmEXPA2)*	2.7	1, 2, 3, 4
Low	156,836,924	0.54	Zm00001d048469	9	*Hydroxylase5 (hyd5)*	-2.9	1, 2, 3
Low	15,993,004	0.88	Zm00001d045203	9	*Aux/IAA-transcription factor 41 (ZmIAA41)*	25.8	1, 2, 3
Low	156,985,230	0.81	Zm00001d048474	9	*ZmMADS1*	5.7	1, 2, 3
High	139,066,037	0.92	Zm00001d026127	10	*Patched family protein (mmp71)*	2.1	1, 2, 3
Low	129,135,133	0.82	Zm00001d025774	10	*Peroxidase superfamily protein*	-2.4	1, 2, 3
Low	130,349,712	0.60	Zm00001d025814	10	*Beta-16-galactosyltransferase (ZmGALT29A)*	-4.0	1, 2, 3
Low	131,515,767	0.89	Zm00001d025851	10	*Xyloglucan 6-xylosyltransferase 2 (ZmGT5)*	-5.6	1, 2, 3
Low	133,094,359	0.59	Zm00001d025886	10	*Cytokinin-O-glucosyltransferase 3*	8.7	1, 2, 3
Low	138,350,968	0.81	Zm00001d026095	10	*Sugar isomerase (SIS) family protein*	4.9	1, 2, 3

 FIGURE LEGENDS

Figure 1. Divergent selection for rind penetrometer resistance (RPR) and the associated

phenotypic changes. A. Schema for 15 cycles of divergent selection performed by Dr. Larry

Darrah and colleagues. B. Box plots showing phenotypic divergence of the two populations from

the C0 population. Populations are shown on the *x*-axis and RPR in kilograms of force (Kgf) is

shown on the *y*-axis. Two asterisks indicate significance at $p < 0.01$.

Figure 2. Identification of SNPs associated with divergent selection. Shown here are plots of

significantly selected SNPs obtained from comparison between C15-H and C0 (A), and between

C15-L and C0 (B). Each dot represents a SNP, vertical position of the dot represents F_{ST} value

shown on *y*-axis, and the horizontal location of the dot represents physical location of the SNP

on a chromosome shown on *x*-axis. Grey and black color differentiates the chromosomes. Red

and blue colored dots represent SNPs that were significant according to our simulation strategy.

Blue colored dots represent significant SNPs that are common in both high (C15-H) and low

(C15-L) populations.

Figure 3. Changes in phenotype and chemical composition of internodes associated with

divergent selection for rind penetrometer resistance. (A) Time course analysis showing

divergence of RPR in the Hrpr1 and Lrpr1 lines. The *x*-axis shows days after sampling and the *y*-

axis denotes RPR in kilograms of force (Kgf) measured by the rind penetrometer. (B) Glucose

and pentose accumulation in internode of Lrpr1 and Hrpr1 inbreds. (C) Histochemical staining

with Phloroglucinol-HCl of stem section of Lrpr1 (left) and Hrpr1 (right). (D-G) Accumulation

of lignin monomers syringic acid (S), vanillic acid (V), and p-hydroxy benzoate (H) in

internodes of Lrpr1 and Hrpr1 inbreds. Shown here is time course accumulation of SVH (D), S

(E), V (F) and, H (G). (H-K) Accumulation of phenolics involved in polymerization of lignin

monomers in the cell wall. Shown here is time course accumulation of p-Coumaric acid (H) and

ferulic acid (I), lignin bound tricin (J), and unbound tricin (K) in the internodes of Lrpr1 and

Hrpr1 inbreds.

Figure 4. Selection of cell wall associated SNPs in low RPR (C15-L) and high RPR (C15-H)

populations. Red diamonds indicate the observed value of enriched SNPs for lignin (left) and

polysaccharide (right) associated genes.

Figure 5. Transcriptomic dynamics during RPR development. (A) Clustering based on PCA

performed for different developmental stages of Lrpr1 (blue triangles) and Hrpr1 (red ovals)

internodes. The developmental stages (days after sampling; D) are mentioned with corresponding

triangles and ovals. (B) Differentially expressed genes during internode development. The *x*-axis

represents number of upregulated and downregulated genes in Hrpr1 compared to Lrpr1 and *y*-

axis represents developmental stages. (C) GO enrichment analysis showing significantly

enriched biological processes with FDR corrected $p < 0.05$, based on downregulated (D) and

upregulated (U) genes in Hrpr1 compare to Lrpr1. (D) Unique and common differentially

expressed genes in Hrpr1 and Lrpr1.

Figure 6. GO terms enriched in Lrpr1 and Hrpr1 based on the genes identified by combining

selection mapping and transcriptome analyses. GO terms represented here are significant at $p <$

0.01. Asterisks next to GO term description denote significance after Bonferroni correction.

Figure 7. Genetic architecture of RPR deduced from integration of different datasets. Orange

dots in the upper and lower panel indicate significantly selected SNPs in high (C15-H) and low

(C15-L) RPR, respectively. Middle panel indicates the QTL (blue, brown and black lines) and

GWAS SNPs (red dots) associated with RPR reported in earlier genetic studies (Flint-Garcia et

al., 2003; Hu et al., 2012; Peiffer et al., 2013; Li et al., 2014). Black circles highlight the SNPs

lying in the vicinity of genes with at least 2-fold differential expression in Hrpr1 and Lrpr1

internodes. Crosses, and triangles represent the SNPs associated with polysaccharide and lignin

pathways genes, respectively. Key genes associated with RPR identified from different

approaches are indicated with black squares.

SUPPLEMENTARY DATA

Table S1. Significant SNPs divergently selected for high RPR (C15-H) and low RPR (C15-L)
relative to C0

Table S2. Annotation of maize cell wall synthesis genes

Table S3. Differentially expressed genes (fold change >2; p(adj) < 0.05) in different
development stages (DAS) of Lrpr1 compared to Hrpr1

Table S4. High priority SNPs selected based on 5% F_{ST} from divergent selection for high RPR
(C15-H) and low RPR (C15-L)

Table S5. Genes associated with RPR identified from integration of selection mapping and
transcriptome data

Table S6. Gene ontology (GO) enrichment of genes identified by integration of selection
mapping and differential gene expression

Figure S1. Individual monolignols constituents in internodes of Lrpr1 and Hrpr1 inbreds. Shown
here are syringic acid, syringaldehyde and acetosyringone of S lignin (A-C); and vanillic acid,
vanillin and acetovanillone of V lignin (D-F); p-hydroxy benzoic acid, p-hydroxy benzaldehyde,
and p-hydroxy acetophenone of H lignin (G-I).

Figure S2. Transcriptomic changes during Lrpr1 and Hrpr1 development. Differentially
expressed (DE) genes at each developmental stage relative to 0 DAS in Hrpr1 (A) and Lrpr1 (B).
The x-axis represents number of upregulated and downregulated genes relative to 0 DAS and y-
axis represents different stages. (C) GO enrichment analysis showing significantly enriched
biological processes (FDR corrected $p < 0.05$) based on unique DE genes in Hrpr1 and Lrpr1,
and common DE genes in both inbred lines.

REFERENCES

Akey JM (2009) Constructing genomic maps of positive selection in humans: Where do we go from here? Genome Research **19:** 711-722

Akin DE (1989) Histological and physical factors affecting digestibility of forages. Agronomy Journal **81:** 17-25

Anders S, Pyl PT, Huber W (2015) HTSeq-a Python framework to work with high-throughput sequencing data. Bioinformatics **31:** 166-169

Appenzeller L, Doblin M, Barreiro R, Wang H, Niu X, Kollipara K, Carrigan L, Tomes D, Chapman M, Dhugga KS (2004) Cellulose synthesis in maize: isolation and expression analysis of the cellulose synthase (CesA) gene family. Cellulose **11:** 287-299

Araya CL, Payen C, Dunham MJ, Fields S (2010) Whole-genome sequencing of a laboratory-evolved yeast strain. BMC Genomics **11:** 88

Bai M-Y, Shang J-X, Oh E, Fan M, Bai Y, Zentella R, Sun T-p, Wang Z-Y (2012) Brassinosteroid, gibberellin and phytochrome impinge on a common transcription module in Arabidopsis. Nature Cell Biology **14:** 810

Beissinger TM, Hirsch CN, Sekhon RS, Foerster JM, Johnson JM, Muttoni G, Vaillancourt B, Buell CR, Kaeppler SM, de Leon N (2013) Marker density and read depth for genotyping populations using genotyping-by-sequencing. Genetics **193:** 1073

Beissinger TM, Hirsch CN, Vaillancourt B, Deshpande S, Barry K, Buell CR, Kaeppler SM, Gianola D, de Leon N (2014) A genome-wide scan for evidence of selection in a maize population under long-term artificial selection for ear number. Genetics **196:** 829

Beissinger TM, Wang L, Crosby K, Durvasula A, Hufford MB, Ross-Ibarra J (2016) Recent demography drives changes in linked selection across the maize genome. Nature Plants **2:** 16084

Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate - a practical and powerful approach to multiple testing. Journal of the Royal Statistical Society Series B-Methodological **57:** 289-300

Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics **30:** 2114-2120

Bosch M, Mayer C-D, Cookson A, Donnison IS (2011) Identification of genes involved in cell wall biogenesis in grasses by differential gene expression profiling of elongating and non-elongating maize internodes. Journal of Experimental Botany **62:** 3545-3561

Braybrook SA, Peaucelle A (2013) Mechano-chemical aspects of organ formation in *Arabidopsis thaliana*: the relationship between auxin and pectin. PloS One **8:** e57813-e57813

Chen M, Wang J, Wang Y, Wu Y, Fu J, Liu J-F (2018) Genome-wide detection of selection signatures in chinese indigenous laiwu pigs revealed candidate genes regulating fat deposition in muscle. BMC genetics **19:** 31-31

Ching A, Dhugga KS, Appenzeller L, Meeley R, Bourett TM, Howard RJ, Rafalski A (2006) *Brittle stalk 2* encodes a putative glycosylphosphatidylinositol-anchored protein that affects mechanical strength of maize tissues by altering the composition and structure of secondary cell walls. Planta **224:** 1174-1184

Cosgrove D, Jarvis M (2012) Comparative structure and biomechanics of plant primary and secondary cell walls. Frontiers in Plant Science **3**

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE,
Lunter G, Marth GT, Sherry ST, McVean G, Durbin R, Genomes Project Analysis
G (2011) The variant call format and VCFtools. Bioinformatics (Oxford, England) **27:**
2156-2158

Davidson J, Phillips M (1930) Lignin as a possible factor in lodging of cereals. Science **72:** 401

Dixon RA, Achnine L, Kota P, Liu C-J, Reddy MSS, Wang L (2002) The phenylpropanoid
pathway and plant defence—a genomics perspective. Molecular Plant Pathology **3:** 371-
390

Dudley JW (1994) Selection for rind puncture resistance in two maize populations. Crop
Science **34:** 1458-1460

Elshire RJ, Glaubitz JC, Sun Q, Poland JA, Kawamoto K, Buckler ES, Mitchell SE (2011)
A robust, simple genotyping-by-sequencing (GBS) approach for high diversity species.
PLoS ONE **6:** e19379

Falcioni R, Moriwaki T, de Oliveira DM, Andreotti GC, de Souza LA, Dos Santos WD,
Bonato CM, Antunes WC (2018) Increased gibberellins and light levels promotes cell
wall thickness and enhance lignin deposition in xylem fibers. Frontiers in Plant Science
9: 1391-1391

Felipo-Benavent A, Úrbez C, Blanco-Touriñán N, Serrano-Mislata A, Baumberger N,
Achard P, Agustí J, Blázquez MA, Alabadí D (2018) Regulation of xylem fiber
differentiation by gibberellins through DELLA-KNAT1 interaction. Development **145:**
dev164962

Feltus FA, Vandenbrink JP (2012) Bioenergy grass feedstock: current options and prospects
for trait improvement using emerging genetic, genomic, and systems biology toolkits.
Biotechnology for Biofuels **5:** 80

Flint-Garcia SA, Jampatong C, Darrah LL, McMullen MD (2003) Quantitative trait locus
analysis of stalk strength in four maize populations. Crop Science **43:** 13-22

Fornalé S, Shi X, Chai C, Encina A, Irar S, Capellades M, Fuguet E, Torres J-L, Rovira P,
Puigdomènech P, Rigau J, Grotewold E, Gray J, Caparrós-Ruiz D (2010)
ZmMYB31 directly represses maize lignin genes and redirects the phenylpropanoid
metabolic flux. The Plant Journal **64:** 633-644

Fornalé S, Sonbol F-M, Maes T, Capellades M, Puigdomènech P, Rigau J, Caparrós-Ruiz
D (2006) Down-regulation of the maize and arabidopsis thaliana caffeic acid O-methyl-
transferase genes by two new maize R2R3-MYB transcription factors. Plant Molecular
Biology **62:** 809-823

Fraser CM, Chapple C (2011) The phenylpropanoid pathway in Arabidopsis. The Arabidopsis
Book **2011**

Gage JL, White MR, Edwards JW, Kaeppler S, de Leon N (2018) Selection signatures
underlying dramatic male inflorescence transformation during modern hybrid maize
breeding. Genetics **210:** 1125

Gerke JP, Edwards JW, Guill KE, Ross-Ibarra J, McMullen MD (2015) The genomic
impacts of drift and selection for hybrid performance in maize. Genetics **201:** 1201

Glaubitz JC, Casstevens TM, Lu F, Harriman J, Elshire RJ, Sun Q, Buckler ES (2014)
TASSEL-GBS: A high capacity genotyping by sequencing analysis pipeline. PLOS ONE
9: e90346

Grainger CM, Letarte J, Rajcan I (2018) Using soybean pedigrees to identify genomic selection signatures associated with long-term breeding for cultivar improvement. Canadian Journal of Plant Science **98:** 1176-1187

Gray J, Caparrós-Ruiz D, Grotewold E (2012) Grass phenylpropanoids: Regulate before using! Plant Science **184:** 112-120

Guillaumie S, San-Clemente H, Deswarte C, Martinez Y, Lapierre C, Murigneux A, Barrière Y, Pichon M, Goffner D (2007) Maizewall. Database and developmental gene expression profiling of cell wall biosynthesis and assembly in maize. Plant Physiology **143:** 339-363

Halpin C (2019) Lignin engineering to improve saccharification and digestibility in grasses. Current Opinion in Biotechnology **56:** 223-229

Hatfield RD, Rancour DM, Marita JM (2017) Grass cell walls: A story of cross-linking. Frontiers in Plant Science **7:** 2056-2056

Hedges JI, Ertel JR (1982) Characterization of lignin by gas capillary chromatography of cupric oxide oxidation products. Analytical Chemistry **54:** 174-178

Heredia DO, Alsirt A, L. DL, Coe EH (1996) Allelic frequency changes in the MoSCSSS maize synthetic in response to bi-directional recurrent selection for rind penetrometer resistance. Maydica **41:** 65–76

Higuchi K, Iwase J, Tsukiori Y, Nakura D, Kobayashi N, Ohashi H, Saito A, Miwa E (2014) Early senescence of the oldest leaves of Fe-deficient barley plants may contribute to phytosiderophore release from the roots. Physiologia Plantarum **151:** 313-322

Hirsch CN, Flint-Garcia SA, Beissinger TM, Eichten SR, Deshpande S, Barry K, McMullen MD, Holland JB, Buckler ES, Springer N, Buell CR, de Leon N, Kaeppler SM (2014) Insights into the effects of long-term artificial selection on seed size in maize. Genetics **198:** 409

Hu H, Meng Y, Wang H, Liu H, Chen S (2012) Identifying quantitative trait loci and determining closely related stalk traits for rind penetrometer resistance in a high-oil maize population. Theoretical and Applied Genetics **124:** 1439-1447

Huang D, Wang S, Zhang B, Shang-Guan K, Shi Y, Zhang D, Liu X, Wu K, Xu Z, Fu X, Zhou Y (2015) A gibberellin-mediated DELLA-NAC signaling cascade regulates cellulose synthesis in rice. The Plant cell **27:** 1681-1696

Hufford MB, Xu X, van Heerwaarden J, Pyhäjärvi T, Chia J-M, Cartwright RA, Elshire RJ, Glaubitz JC, Guill KE, Kaeppler SM, Lai J, Morrell PL, Shannon LM, Song C, Springer NM, Swanson-Wagner RA, Tiffin P, Wang J, Zhang G, Doebley J, McMullen MD, Ware D, Buckler ES, Yang S, Ross-Ibarra J (2012) Comparative population genomics of maize domestication and improvement. Nature Genetics **44:** 808

Jampatong S, Darrah LL, Krause GF, Barry BD (2000) Effect of one- and two-eared selection on stalk strength and other characters in maize. Crop Science **40:** 605-611

Jiao Y, Peluso P, Shi J, Liang T, Stitzer MC, Wang B, Campbell MS, Stein JC, Wei X, Chin C-S, Guill K, Regulski M, Kumari S, Olson A, Gent J, Schneider KL, Wolfgruber TK, May MR, Springer NM, Antoniou E, McCombie WR, Presting GG, McMullen M, Ross-Ibarra J, Dawe RK, Hastie A, Rank DR, Ware D (2017) Improved maize reference genome with single-molecule technologies. Nature **546:** 524-527

Kaiser K, Benner R (2012) Characterization of lignin by gas chromatography and mass
spectrometry using a simplified CuO oxidation method. Analytical Chemistry **84:** 459-
464
Kang MS, Din AK, Zhang Y, Magari R (1999) Combining ability for rind puncture resistance
in maize. Crop Science **39:** 368-371
Kelly JK, Hughes KA (2019) Pervasive linked selection and intermediate-frequency alleles are
implicated in an evolve-and-resequencing experiment of *Drosophila simulans*. Genetics
211: 943
Khanday I, Yadav SR, Vijayraghavan U (2013) Rice *LHS1/OsMADS1* controls floret
meristem specification by coordinated regulation of transcription factors and hormone
signaling pathways. Plant physiology **161:** 1970-1983
Khanna KL (1935) An improved instrument for testing rind hardness in sugarcane. Agriculture
and live-stock in India **5:** 156-158
Kimura R, Fujimoto A, Tokunaga K, Ohashi J (2007) A practical genome scan for
population-specific strong selective sweeps that have reached fixation. PLOS ONE **2:**
e286
Kofler R, Schlötterer C (2014) A guide for the design of evolve and resequencing studies.
Molecular Biology and Evolution **31:** 474-483
Lan W, Lu F, Regner M, Zhu Y, Rencoret J, Ralph SA, Zakai UI, Morreel K, Boerjan W,
Ralph J (2015) Tricin, a flavonoid monomer in monocot lignification. Plant Physiology
167: 1284-1295
Langmead B, Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. Nature methods
9: 357-359
Lehman TA, Smertenko A, Sanguinet KA (2017) Auxin, microtubules, and vesicle trafficking:
Conspirators behind the cell wall. Journal of Experimental Botany **68:** 3321-3329
Li F, Zhang M, Guo K, Hu Z, Zhang R, Feng Y, Yi X, Zou W, Wang L, Wu C, Tian J, Lu
T, Xie G, Peng L (2015) High-level hemicellulosic arabinose predominately affects
lignocellulose crystallinity for genetically enhancing both plant lodging resistance and
biomass enzymatic digestibility in rice mutants. Plant Biotechnology Journal **13:** 514-525
Li K, Yan J, Li J, Yang X (2014) Genetic architecture of rind penetrometer resistance in two
maize recombinant inbred line populations. BMC Plant Biology **14:** 152-152
Lorenz AJ, Beissinger TM, Silva RR, de Leon N (2015) Selection for silage yield and
composition did not affect genomic diversity within the Wisconsin quality synthetic
maize population. G3: Genes|Genomes|Genetics **5:** 541
Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biology **15:** 550
Maeda H, Dudareva N (2012) The shikimate pathway and aromatic amino acid biosynthesis in
plants. Annual Review of Plant Biology **63:** 73-105
Majda M, Robert S (2018) The role of auxin in cell wall expansion. International journal of
molecular sciences **19:** 951
Martin SA, Darrah LL, Hibbard BE (2004) Divergent selection for rind penetrometer
resistance and its effects on european corn borer damage and stalk traits in corn. Crop
Science **44:** 711-717
Miller JE, Geadelmann JL, Marten GC (1983) Effect of the brown midrib-allele on maize
silage quality and yield1. **23:** 493-496

Narum SR, Hess JE (2011) Comparison of F_{ST} outlier tests for SNP loci under selection.
Molecular Ecology Resources **11:** 184-194

Nielsen R, Williamson S, Kim Y, Hubisz MJ, Clark AG, Bustamante C (2005) Genomic
scans for selective sweeps using SNP data. Genome Research **15:** 1566-1575

Olsen KM, Wendel JF (2013) A bountiful harvest: Genomic insights into crop domestication
phenotypes. Annual Review of Plant Biology **64:** 47-70

Paque S, Mouille G, Grandont L, Alabadí D, Gaertner C, Goyallon A, Muller P, Primard-
Brisset C, Sormani R, Blázquez MA, Perrot-Rechenmann C (2014) AUXIN
BINDING PROTEIN1 links cell wall remodeling, auxin signaling, and cell expansion in
Arabidopsis. The Plant Cell **26:** 280

Pascual L, Albert E, Sauvage C, Duangjit J, Bouchet J-P, Bitton F, Desplat N, Brunel D, Le
Paslier M-C, Ranc N, Bruguier L, Chauchard B, Verschave P, Causse M (2016)
Dissecting quantitative trait variation in the resequencing era: complementarity of bi-
parental, multi-parental and association panels. Plant Science **242:** 120-130

Pedersen JF, Vogel KP, Funnell DL (2005) Impact of reduced lignin on plant fitness. Crop
Science **45:** 812-819

Peiffer JA, Flint-Garcia SA, De Leon N, McMullen MD, Kaeppler SM, Buckler ES (2013)
The genetic architecture of maize stalk strength. PLoS ONE **8:** e67066

Pétriacq P, de Bont L, Hager J, Didierlaurent L, Mauve C, Guérard F, Noctor G, Pelletier
S, Renou J-P, Tcherkez G, Gakière B (2012) Inducible NAD overproduction in
Arabidopsis alters metabolic pools and gene expression correlated with increased
salicylate content and resistance to *Pst-AvrRpm1*. The Plant Journal **70:** 650-665

Plassais J, Kim J, Davis BW, Karyadi DM, Hogan AN, Harris AC, Decker B, Parker HG,
Ostrander EA (2019) Whole genome sequencing of canids reveals genomic regions
under selection and variants influencing morphology. Nature Communications **10:** 1489

Qanbari S, Strom TM, Haberer G, Weigend S, Gheyas AA, Turner F, Burt DW, Preisinger
R, Gianola D, Simianer H (2012) A high resolution genome-wide scan for significant
selective sweeps: An application to pooled sequence data in laying chickens. PLOS ONE
7: e49525

R Core Team (2013) R: A language and environment for statistical computing. *In*. R Foundation
for Statistical Computing, Vienna, Austria

Ralph J, Lapierre C, Boerjan W (2019) Lignin structure and its engineering. Current Opinion
in Biotechnology **56:** 240-249

Robertson D, Smith S, Gardunia B, Cook D (2014) An improved method for accurate
phenotyping of corn stalk strength. Crop Science **54:** 2038-2044

Robertson DJ, Lee SY, Julias M, Cook DD (2016) Maize stalk lodging: Flexural stiffness
predicts strength. Crop Science **56:** 1711-1718

Rohde A, Morreel K, Ralph J, Goeminne G, Hostyn V, De Rycke R, Kushnir S, Van
Doorsselaere J, Joseleau J-P, Vuylsteke M, Van Driessche G, Van Beeumen J,
Messens E, Boerjan W (2004) Molecular phenotyping of the *pal1* and *pal2* mutants of
Arabidopsis thaliana reveals far-reaching consequences on phenylpropanoid, amino acid,
and carbohydrate metabolism. The Plant cell **16:** 2749-2771

Ruzin SE (1999) Plant microtechnique and microscopy. Oxford University Press, New York

Saghai-Maroof MA, Soliman KM, Jorgensen RA, Allard RW (1984) Ribosomal DNA
spacer-length polymorphisms in barley: mendelian inheritance, chromosomal location,

and population dynamics. Proceedings of the National Academy of Sciences **81:** 8014-
8018

Sánchez-Montesino R, Bouza-Morcillo L, Marquez J, Ghita M, Duran-Nebreda S, Gómez
L, Holdsworth MJ, Bassel G, Oñate-Sánchez L (2019) A regulatory module
controlling GA-mediated endosperm cell expansion is critical for seed germination in
Arabidopsis. Molecular Plant **12:** 71-85

Sánchez-Rodríguez C, Ketelaar K, Schneider R, Villalobos JA, Somerville CR, Persson S,
Wallace IS (2017) BRASSINOSTEROID INSENSITIVE2 negatively regulates cellulose
synthesis in *Arabidopsis* by phosphorylating cellulose synthase 1. Proceedings of the
National Academy of Sciences **114:** 3533-3538

Santoro N, Cantu S, Tornqvist C-E, Falbel T, Bolivar J, Patterson S, Pauly M, Walton J
(2010) A high-throughput platform for screening milligram quantities of plant biomass
for lignocellulose digestibility. BioEnergy Research **3:** 93-102

Schaefer RJ, Michno J-M, Jeffers J, Hoekenga O, Dilkes B, Baxter I, Myers CL (2018)
Integrating coexpression networks with gwas to prioritize causal genes in maize. The
Plant Cell **30:** 2922

Sekhon RS, Breitzman MW, Silva RR, Santoro N, Rooney WL, de Leon N, Kaeppler SM
(2016) Stover composition in maize and sorghum reveals remarkable genetic variation
and plasticity for carbohydrate accumulation. Frontiers in Plant Science **7**

Sekhon RS, Joyner CN, Ackerman AJ, McMahan CS, Cook DD, Robertson DJ (2019) Stalk
bending strength is strongly associated with maize stalk lodging incidence across
multiple environments. bioRxiv**:** 675793

Sibale EM, Darrah LL, Zuber MS (1992) Comparison of two rind penetrometers for
measurement of stalk strength in maize. Maydica **37:** 111-114

Smith JM, Haigh J (1974) The hitch-hiking effect of a favourable gene. Genetical Research **23:**
23-35

Sun Q, Liu X, Yang J, Liu W, Du Q, Wang H, Fu C, Li W-X (2018) Microrna528 affects
lodging resistance of maize by regulating lignin biosynthesis under nitrogen-luxury
conditions. Molecular Plant **11:** 806-814

Tam V, Patel N, Turcotte M, Bossé Y, Paré G, Meyre D (2019) Benefits and limitations of
genome-wide association studies. Nature Reviews Genetics **20:** 467-484

Tobimatsu Y, Schuetz M (2019) Lignin polymerization: how do plants manage the chemistry
so well? Current Opinion in Biotechnology **56:** 75-81

Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, Pimentel H, Salzberg SL,
Rinn JL, Pachter L (2012) Differential gene and transcript expression analysis of RNA-
seq experiments with TopHat and Cufflinks. Nature protocols **7:** 562-578

Turner TL, Miller PM (2012) Investigating natural variation in Drosophila courtship song by
the evolve and resequence approach. Genetics **191:** 633

Turner TL, Stewart AD, Fields AT, Rice WR, Tarone AM (2011) Population-based
resequencing of experimentally evolved populations reveals the genetic basis of body
size variation in *Drosophila melanogaster*. PLOS Genetics **7:** e1001336

Undersander DJ, Bauman LF, Lechtenberg VL, Zuber MS (1977) Effect of cyclic selection
for high and low crushing strength on rind, pith, and whole stalk composition in corn.
Crop Science **17:** 732-734

Vanholme B, Vanholme R, Turumtay H, Goeminne G, Cesarino I, Goubet F, Morreel K,
Rencoret J, Bulone V, Hooijmaijers C, De Rycke R, Gheysen G, Ralph J, De Block

M, Meulewaeter F, Boerjan W (2014) Accumulation of N-acetylglucosamine oligomers in the plant cell wall affects plant architecture in a dose-dependent and conditional manner. Plant Physiology **165**: 290-308

Vanholme R, Demedts B, Morreel K, Ralph J, Boerjan W (2010) Lignin biosynthesis and structure. Plant Physiology **153**: 895

Vlachos C, Kofler R (2019) Optimizing the power to identify the genetic basis of complex traits with Evolve and Resequence studies. bioRxiv**:** 583682

Vogt T (2010) Phenylpropanoid biosynthesis. Molecular Plant **3:** 2-20

Wang J-J, Tharayil N, Chow AT, Suseela V, Zeng H (2015) Phenolic profile within the fine-root branching orders of an evergreen species highlights a disconnect in root tissue quality predicted by elemental- and molecular-level carbon composition. New Phytologist **206:** 1261-1273

Weir BS, Cockerham CC (1984) Estimating F-statistics for the analysis of population structure. Evolution **38:** 1358-1370

Xie L, Yang C, Wang X (2011) Brassinosteroids can regulate cellulose biosynthesis by controlling the expression of *CESA* genes in *Arabidopsis*. Journal of Experimental Botany **62:** 4495-4506

Yang D, Cai T, Luo Y, Wang Z (2019) Optimizing plant density and nitrogen application to manipulate tiller growth and increase grain yield and nitrogen-use efficiency in winter wheat. PeerJ **7:** e6484

Ye DL, Zhang YS, Al-Kaisi MM, Duan LS, Zhang MC, Li ZH (2016) Ethephon improved stalk strength associated with summer maize adaptations to environments differing in nitrogen availability in the north China plain. The Journal of Agricultural Science **154:** 960-977

You FM, Xiao J, Li P, Yao Z, Jia G, He L, Kumar S, Soto-Cerda B, Duguid SD, Booker HM, Rashid KY, Cloutier S (2018) Genome-wide association study and selection signatures detect genomic regions associated with seed yield and oil quality in flax. International Journal of Molecular Sciences **19:** 2303

Young MD, Wakefield MJ, Smyth GK, Oshlack A (2010) Gene ontology analysis for RNA-seq: accounting for selection bias. Genome biology **11:** R14

Zhang Y, Legland D, El Hage F, Devaux M-F, Guillon F, Reymond M, Méchin V (2019) Changes in cell walls lignification, feruloylation and p-coumaroylation throughout maize internode development. PLOS One **14:** e0219923

Zhang Z, Zhang X, Lin Z, Wang J, Liu H, Zhou L, Zhong S, Li Y, Zhu C, Lai J, Li X, Yu J, Lin Z (2020) A large transposon insertion in the *stiff1* promoter increases stalk strength in maize. Plant Cell **32:** 152

Zhou Z, Jiang Y, Wang Z, Gou Z, Lyu J, Li W, Yu Y, Shu L, Zhao Y, Ma Y, Fang C, Shen Y, Liu T, Li C, Li Q, Wu M, Wang M, Wu Y, Dong Y, Wan W, Wang X, Ding Z, Gao Y, Xiang H, Zhu B, Lee S-H, Wang W, Tian Z (2015) Resequencing 302 wild and cultivated accessions identifies genes related to domestication and improvement in soybean. Nature Biotechnology **33:** 408

Zuber M, Grogan C (1961) A new technique for measuring stalk strength in corn. Crop Science **1:** 378-380

Zuber MS, Colbert TR, Darrah LL (1980) Effect of recurrent selection for crushing strength on several stalk components in maize. Crop Science **20:** 711-717

Zuber MS, Kang MS (1978) Corn lodging slowed by sturdier stalks. Crops Soils **30:** 13-15

Figure 1

Figure 1. Divergent selection for rind penetrometer resistance (RPR) and the associated phenotypic changes. A. Schema for 15 cycles of divergent selection performed by Dr. Larry Darrah and colleagues. B. Box plots showing phenotypic divergence of the two populations from the C0 population. Populations are shown on the *x*-axis and RPR in kilograms of force (Kgf) is shown on the *y*-axis. Two asterisks indicate significance at *p* < 0.01.

Figure 2

Figure 2. Identification of SNPs associated with divergent selection. Shown here are plots of significantly selected SNPs obtained from comparison between C15-H and C0 (A), and between C15-L and C0 (B). Each dot represents a SNP, vertical position of the dot represents F_{ST} value shown on *y*-axis, and the horizontal location of the dot represents physical location of the SNP on a chromosome shown on *x*-axis. Grey and black color differentiates the chromosomes. Red and blue colored dots represent SNPs that were significant according to our simulation strategy. Blue colored dots represent significant SNPs that are common in both high (C15-H) and low (C15-L) populations.

Figure 3

Figure 3. Changes in phenotype and chemical composition of internodes associated with divergent selection for rind penetrometer resistance. (A) Time course analysis showing divergence of RPR in the Hrpr1 and Lrpr1 lines. The *x*-axis shows days after sampling and the *y*-axis denotes RPR in kilograms of force (Kgf) measured by the rind penetrometer. (B) Glucose and pentose accumulation in internode of Lrpr1 and Hrpr1 inbreds. (C) Histochemical staining with Phloroglucinol-HCl of stem section of Lrpr1 (left) and Hrpr1 (right). (D-G) Accumulation of lignin monomers syringic acid (S), vanillic acid (V), and p-hydroxy benzoate (H) in internodes of Lrpr1 and Hrpr1 inbreds. Shown here is time course accumulation of SVH (D), S (E), V (F) and, H (G). (H-K) Accumulation of phenolics involved in polymerization of lignin monomers in the cell wall. Shown here is time course accumulation of p-Coumaric acid (H) and ferulic acid (I), lignin bound tricin (J), and unbound tricin (K) in the internodes of Lrpr1 and Hrpr1 inbreds.

Figure 4

Figure 4. Selection of cell wall associated SNPs in low RPR (C15-L) and high RPR (C15-H) populations. Red diamonds indicate the observed value of enriched SNPs for lignin (left) and polysaccharide (right) associated genes.

Figure 5

Figure 5. Transcriptomic dynamics during RPR development. (A) Clustering based on PCA performed for different developmental stages of Lrpr1 (blue triangles) and Hrpr1 (red ovals) internodes. The developmental stages (days after sampling; D) are mentioned with corresponding triangles and ovals. (B) Differentially expressed genes during internode development. The *x*-axis represents number of upregulated and downregulated genes in Hrpr1 compared to Lrpr1 and *y*-axis represents developmental stages. (C) GO enrichment analysis showing significantly enriched biological processes with FDR corrected $p < 0.05$, based on downregulated (D) and upregulated (U) genes in Hrpr1 compare to Lrpr1. (D) Unique and common differentially expressed genes in Hrpr1 and Lrpr1.

Figure 6

Figure 6. GO terms enriched in Lrpr1 and Hrpr1 based on the genes identified by combining selection mapping and transcriptome analyses. GO terms represented here are significant at *p* < 0.01. Asterisks next to GO term description denote significance after Bonferroni correction.

Figure 7

Figure 7. Genetic architecture of RPR deduced from integration of different datasets. Orange dots in the upper and lower panel indicate significantly selected SNPs in high (C15-H) and low (C15-L) RPR, respectively. Middle panel indicates the QTL (blue, brown and black lines) and GWAS SNPs (red dots) associated with RPR reported in earlier genetic studies (Flint-Garcia et al., 2003; Hu et al., 2012; Peiffer et al., 2013; Li et al., 2014). Black circles highlight the SNPs lying in the vicinity of genes with at least 2-fold differential expression in Hrpr1 and Lrpr1 internodes. Crosses, and triangles represent the SNPs associated with polysaccharide and lignin pathways genes, respectively. Key genes associated with RPR identified from different approaches are indicated with black squares.

Parsed Citations

Akey JM (2009) Constructing genomic maps of positive selection in humans: Where do we go from here? Genome Research 19: 711-722
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Akin DE (1989) Histological and physical factors affecting digestibility of forages. Agronomy Journal 81: 17-25
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Anders S, Pyl PT, Huber W (2015) HTSeq-a Python framework to work with high-throughput sequencing data. Bioinformatics 31: 166-169
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Appenzeller L, Doblin M, Barreiro R, Wang H, Niu X, Kollipara K, Carrigan L, Tomes D, Chapman M, Dhugga KS (2004) Cellulose synthesis in maize: isolation and expression analysis of the cellulose synthase (CesA) gene family. Cellulose 11: 287-299
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Araya CL, Payen C, Dunham MJ, Fields S (2010) Whole-genome sequencing of a laboratory-evolved yeast strain. BMC Genomics 11: 88
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Bai M-Y, Shang J-X, Oh E, Fan M, Bai Y, Zentella R, Sun T-p, Wang Z-Y (2012) Brassinosteroid, gibberellin and phytochrome impinge on a common transcription module in Arabidopsis. Nature Cell Biology 14: 810
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Beissinger TM, Hirsch CN, Sekhon RS, Foerster JM, Johnson JM, Muttoni G, Vaillancourt B, Buell CR, Kaeppler SM, de Leon N (2013) Marker density and read depth for genotyping populations using genotyping-by-sequencing. Genetics 193: 1073
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Beissinger TM, Hirsch CN, Vaillancourt B, Deshpande S, Barry K, Buell CR, Kaeppler SM, Gianola D, de Leon N (2014) A genome-wide scan for evidence of selection in a maize population under long-term artificial selection for ear number. Genetics 196: 829
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Beissinger TM, Wang L, Crosby K, Durvasula A, Hufford MB, Ross-Ibarra J (2016) Recent demography drives changes in linked selection across the maize genome. Nature Plants 2: 16084
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate - a practical and powerful approach to multiple testing. Journal of the Royal Statistical Society Series B-Methodological 57: 289-300
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics 30: 2114-2120
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Bosch M, Mayer C-D, Cookson A, Donnison IS (2011) Identification of genes involved in cell wall biogenesis in grasses by differential gene expression profiling of elongating and non-elongating maize internodes. Journal of Experimental Botany 62: 3545-3561
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Braybrook SA, Peaucelle A (2013) Mechano-chemical aspects of organ formation in Arabidopsis thaliana: the relationship between auxin and pectin. PloS One 8: e57813-e57813
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Chen M, Wang J, Wang Y, Wu Y, Fu J, Liu J-F (2018) Genome-wide detection of selection signatures in chinese indigenous laiwu pigs revealed candidate genes regulating fat deposition in muscle. BMC genetics 19: 31-31
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ching A, Dhugga KS, Appenzeller L, Meeley R, Bourett TM, Howard RJ, Rafalski A (2006) Brittle stalk 2 encodes a putative glycosylphosphatidylinositol-anchored protein that affects mechanical strength of maize tissues by altering the composition and structure of secondary cell walls. Planta 224: 1174-1184
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Cosgrove D, Jarvis M (2012) Comparative structure and biomechanics of plant primary and secondary cell walls. Frontiers in Plant Science 3

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth GT, Sherry ST, McVean G, Durbin R, Genomes Project Analysis G (2011) The variant call format and VCFtools. Bioinformatics (Oxford, England) 27: 2156-2158

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Davidson J, Phillips M (1930) Lignin as a possible factor in lodging of cereals. Science 72: 401

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Dixon RA, Achnine L, Kota P, Liu C-J, Reddy MSS, Wang L (2002) The phenylpropanoid pathway and plant defence-a genomics perspective. Molecular Plant Pathology 3: 371-390

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Dudley JW (1994) Selection for rind puncture resistance in two maize populations. Crop Science 34: 1458-1460

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Elshire RJ, Glaubitz JC, Sun Q, Poland JA, Kawamoto K, Buckler ES, Mitchell SE (2011) A robust, simple genotyping-by-sequencing (GBS) approach for high diversity species. PLoS ONE 6: e19379

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Falcioni R, Moriwaki T, de Oliveira DM, Andreotti GC, de Souza LA, Dos Santos WD, Bonato CM, Antunes WC (2018) Increased gibberellins and light levels promotes cell wall thickness and enhance lignin deposition in xylem fibers. Frontiers in Plant Science 9: 1391-1391

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Felipo-Benavent A, Úrbez C, Blanco-Touriñán N, Serrano-Mislata A, Baumberger N, Achard P, Agustí J, Blázquez MA, Alabadí D (2018) Regulation of xylem fiber differentiation by gibberellins through DELLA-KNAT1 interaction. Development 145: dev164962

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Feltus FA, Vandenbrink JP (2012) Bioenergy grass feedstock: current options and prospects for trait improvement using emerging genetic, genomic, and systems biology toolkits. Biotechnology for Biofuels 5: 80

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Flint-Garcia SA, Jampatong C, Darrah LL, McMullen MD (2003) Quantitative trait locus analysis of stalk strength in four maize populations. Crop Science 43: 13-22

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Fornalé S, Shi X, Chai C, Encina A, Irar S, Capellades M, Fuguet E, Torres J-L, Rovira P, Puigdomènech P, Rigau J, Grotewold E, Gray J, Caparrós-Ruiz D (2010) ZmMYB31 directly represses maize lignin genes and redirects the phenylpropanoid metabolic flux. The Plant Journal 64: 633-644

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Fornalé S, Sonbol F-M, Maes T, Capellades M, Puigdomènech P, Rigau J, Caparrós-Ruiz D (2006) Down-regulation of the maize and arabidopsis thaliana caffeic acid O-methyl-transferase genes by two new maize R2R3-MYB transcription factors. Plant Molecular Biology 62: 809-823

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Fraser CM, Chapple C (2011) The phenylpropanoid pathway in Arabidopsis. The Arabidopsis Book 2011

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Gage JL, White MR, Edwards JW, Kaeppler S, de Leon N (2018) Selection signatures underlying dramatic male inflorescence transformation during modern hybrid maize breeding. Genetics 210: 1125

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Gerke JP, Edwards JW, Guill KE, Ross-Ibarra J, McMullen MD (2015) The genomic impacts of drift and selection for hybrid performance in maize. Genetics 201: 1201

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Glaubitz JC, Casstevens TM, Lu F, Harriman J, Elshire RJ, Sun Q, Buckler ES (2014) TASSEL-GBS: A high capacity genotyping by sequencing analysis pipeline. PLOS ONE 9: e90346
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Grainger CM, Letarte J, Rajcan I (2018) Using soybean pedigrees to identify genomic selection signatures associated with long-term breeding for cultivar improvement. Canadian Journal of Plant Science 98: 1176-1187
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Gray J, Caparrós-Ruiz D, Grotewold E (2012) Grass phenylpropanoids: Regulate before using! Plant Science 184: 112-120
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Guillaumie S, San-Clemente H, Deswarte C, Martinez Y, Lapierre C, Murigneux A, Barrière Y, Pichon M, Goffner D (2007) Maizewall. Database and developmental gene expression profiling of cell wall biosynthesis and assembly in maize. Plant Physiology 143: 339-363
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Halpin C (2019) Lignin engineering to improve saccharification and digestibility in grasses. Current Opinion in Biotechnology 56: 223-229
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Hatfield RD, Rancour DM, Marita JM (2017) Grass cell walls: A story of cross-linking. Frontiers in Plant Science 7: 2056-2056
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Hedges JI, Ertel JR (1982) Characterization of lignin by gas capillary chromatography of cupric oxide oxidation products. Analytical Chemistry 54: 174-178
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Heredia DO, Alsirt A, L. DL, Coe EH (1996) Allelic frequency changes in the MoSCSSS maize synthetic in response to bi-directional recurrent selection for rind penetrometer resistance. Maydica 41: 65–76
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Higuchi K, Iwase J, Tsukiori Y, Nakura D, Kobayashi N, Ohashi H, Saito A, Miwa E (2014) Early senescence of the oldest leaves of Fe-deficient barley plants may contribute to phytosiderophore release from the roots. Physiologia Plantarum 151: 313-322
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Hirsch CN, Flint-Garcia SA, Beissinger TM, Eichten SR, Deshpande S, Barry K, McMullen MD, Holland JB, Buckler ES, Springer N, Buell CR, de Leon N, Kaeppler SM (2014) Insights into the effects of long-term artificial selection on seed size in maize. Genetics 198: 409
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Hu H, Meng Y, Wang H, Liu H, Chen S (2012) Identifying quantitative trait loci and determining closely related stalk traits for rind penetrometer resistance in a high-oil maize population. Theoretical and Applied Genetics 124: 1439-1447
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Huang D, Wang S, Zhang B, Shang-Guan K, Shi Y, Zhang D, Liu X, Wu K, Xu Z, Fu X, Zhou Y (2015) A gibberellin-mediated DELLA-NAC signaling cascade regulates cellulose synthesis in rice. The Plant cell 27: 1681-1696
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Hufford MB, Xu X, van Heerwaarden J, Pyhäjärvi T, Chia J-M, Cartwright RA, Elshire RJ, Glaubitz JC, Guill KE, Kaeppler SM, Lai J, Morrell PL, Shannon LM, Song C, Springer NM, Swanson-Wagner RA, Tiffin P, Wang J, Zhang G, Doebley J, McMullen MD, Ware D, Buckler ES, Yang S, Ross-Ibarra J (2012) Comparative population genomics of maize domestication and improvement. Nature Genetics 44: 808
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Jampatong S, Darrah LL, Krause GF, Barry BD (2000) Effect of one- and two-eared selection on stalk strength and other characters in maize. Crop Science 40: 605-611
  Pubmed: Author and Title
  Google Scholar: Author Only Title Only Author and Title

Jiao Y, Peluso P, Shi J, Liang T, Stitzer MC, Wang B, Campbell MS, Stein JC, Wei X, Chin C-S, Guill K, Regulski M, Kumari S, Olson A, Gent J, Schneider KL, Wolfgruber TK, May MR, Springer NM, Antoniou E, McCombie WR, Presting GG, McMullen M, Ross-Ibarra J, Dawe RK, Hastie A, Rank DR, Ware D (2017) Improved maize reference genome with single-molecule technologies. Nature 546: 524-527

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Kaiser K, Benner R (2012) Characterization of lignin by gas chromatography and mass spectrometry using a simplified CuO oxidation method. Analytical Chemistry 84: 459-464

Kang MS, Din AK, Zhang Y, Magari R (1999) Combining ability for rind puncture resistance in maize. Crop Science 39: 368-371

Kelly JK, Hughes KA (2019) Pervasive linked selection and intermediate-frequency alleles are implicated in an evolve-and-resequencing experiment of Drosophila simulans. Genetics 211: 943

Khanday I, Yadav SR, Vijayraghavan U (2013) Rice LHS1/OsMADS1 controls floret meristem specification by coordinated regulation of transcription factors and hormone signaling pathways. Plant physiology 161: 1970-1983

Khanna KL (1935) An improved instrument for testing rind hardness in sugarcane. Agriculture and live-stock in India 5: 156-158

Kimura R, Fujimoto A, Tokunaga K, Ohashi J (2007) A practical genome scan for population-specific strong selective sweeps that have reached fixation. PLOS ONE 2: e286

Kofler R, Schlötterer C (2014) A guide for the design of evolve and resequencing studies. Molecular Biology and Evolution 31: 474-483

Lan W, Lu F, Regner M, Zhu Y, Rencoret J, Ralph SA, Zakai UI, Morreel K, Boerjan W, Ralph J (2015) Tricin, a flavonoid monomer in monocot lignification. Plant Physiology 167: 1284-1295

Langmead B, Salzberg SL (2012) Fast gapped-read alignment with Bowtie 2. Nature methods 9: 357-359

Lehman TA, Smertenko A, Sanguinet KA (2017) Auxin, microtubules, and vesicle trafficking: Conspirators behind the cell wall. Journal of Experimental Botany 68: 3321-3329

Li F, Zhang M, Guo K, Hu Z, Zhang R, Feng Y, Yi X, Zou W, Wang L, Wu C, Tian J, Lu T, Xie G, Peng L (2015) High-level hemicellulosic arabinose predominately affects lignocellulose crystallinity for genetically enhancing both plant lodging resistance and biomass enzymatic digestibility in rice mutants. Plant Biotechnology Journal 13: 514-525

Li K, Yan J, Li J, Yang X (2014) Genetic architecture of rind penetrometer resistance in two maize recombinant inbred line populations. BMC Plant Biology 14: 152-152

Lorenz AJ, Beissinger TM, Silva RR, de Leon N (2015) Selection for silage yield and composition did not affect genomic diversity within the Wisconsin quality synthetic maize population. G3: Genes|Genomes|Genetics 5: 541

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biology 15: 550

Maeda H, Dudareva N (2012) The shikimate pathway and aromatic amino acid biosynthesis in plants. Annual Review of Plant Biology 63: 73-105

Majda M, Robert S (2018) The role of auxin in cell wall expansion. International journal of molecular sciences 19: 951
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Martin SA, Darrah LL, Hibbard BE (2004) Divergent selection for rind penetrometer resistance and its effects on european corn borer damage and stalk traits in corn. Crop Science 44: 711-717
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Miller JE, Geadelmann JL, Marten GC (1983) Effect of the brown midrib-allele on maize silage quality and yield1. 23: 493-496
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Narum SR, Hess JE (2011) Comparison of FST outlier tests for SNP loci under selection. Molecular Ecology Resources 11: 184-194
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Nielsen R, Williamson S, Kim Y, Hubisz MJ, Clark AG, Bustamante C (2005) Genomic scans for selective sweeps using SNP data. Genome Research 15: 1566-1575
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Olsen KM, Wendel JF (2013) A bountiful harvest: Genomic insights into crop domestication phenotypes. Annual Review of Plant Biology 64: 47-70
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Paque S, Mouille G, Grandont L, Alabadí D, Gaertner C, Goyallon A, Muller P, Primard-Brisset C, Sormani R, Blázquez MA, Perrot-Rechenmann C (2014) AUXIN BINDING PROTEIN1 links cell wall remodeling, auxin signaling, and cell expansion in Arabidopsis. The Plant Cell 26: 280
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Pascual L, Albert E, Sauvage C, Duangjit J, Bouchet J-P, Bitton F, Desplat N, Brunel D, Le Paslier M-C, Ranc N, Bruguier L, Chauchard B, Verschave P, Causse M (2016) Dissecting quantitative trait variation in the resequencing era: complementarity of bi-parental, multi-parental and association panels. Plant Science 242: 120-130
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Pedersen JF, Vogel KP, Funnell DL (2005) Impact of reduced lignin on plant fitness. Crop Science 45: 812-819
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Peiffer JA, Flint-Garcia SA, De Leon N, McMullen MD, Kaeppler SM, Buckler ES (2013) The genetic architecture of maize stalk strength. PLoS ONE 8: e67066
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Pétriacq P, de Bont L, Hager J, Didierlaurent L, Mauve C, Guérard F, Noctor G, Pelletier S, Renou J-P, Tcherkez G, Gakière B (2012) Inducible NAD overproduction in Arabidopsis alters metabolic pools and gene expression correlated with increased salicylate content and resistance to Pst-AvrRpm1. The Plant Journal 70: 650-665
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Plassais J, Kim J, Davis BW, Karyadi DM, Hogan AN, Harris AC, Decker B, Parker HG, Ostrander EA (2019) Whole genome sequencing of canids reveals genomic regions under selection and variants influencing morphology. Nature Communications 10: 1489
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Qanbari S, Strom TM, Haberer G, Weigend S, Gheyas AA, Turner F, Burt DW, Preisinger R, Gianola D, Simianer H (2012) A high resolution genome-wide scan for significant selective sweeps: An application to pooled sequence data in laying chickens. PLOS ONE 7: e49525
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

R Core Team (2013) R: A language and environment for statistical computing. In. R Foundation for Statistical Computing, Vienna, Austria
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ralph J, Lapierre C, Boerjan W (2019) Lignin structure and its engineering. Current Opinion in Biotechnology 56: 240-249
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Robertson D, Smith S, Gardunia B, Cook D (2014) An improved method for accurate phenotyping of corn stalk strength. Crop Science

54: 2038-2044
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Robertson DJ, Lee SY, Julias M, Cook DD (2016) Maize stalk lodging: Flexural stiffness predicts strength. Crop Science 56: 1711-1718
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Rohde A, Morreel K, Ralph J, Goeminne G, Hostyn V, De Rycke R, Kushnir S, Van Doorsselaere J, Joseleau J-P, Vuylsteke M, Van Driessche G, Van Beeumen J, Messens E, Boerjan W (2004) Molecular phenotyping of the pal1 and pal2 mutants of Arabidopsis thaliana reveals far-reaching consequences on phenylpropanoid, amino acid, and carbohydrate metabolism. The Plant cell 16: 2749-2771
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ruzin SE (1999) Plant microtechnique and microscopy. Oxford University Press, New York
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Saghai-Maroof MA, Soliman KM, Jorgensen RA, Allard RW (1984) Ribosomal DNA spacer-length polymorphisms in barley: mendelian inheritance, chromosomal location, and population dynamics. Proceedings of the National Academy of Sciences 81: 8014-8018
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Sánchez-Montesino R, Bouza-Morcillo L, Marquez J, Ghita M, Duran-Nebreda S, Gómez L, Holdsworth MJ, Bassel G, Oñate-Sánchez L (2019) A regulatory module controlling GA-mediated endosperm cell expansion is critical for seed germination in Arabidopsis. Molecular Plant 12: 71-85
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Sánchez-Rodríguez C, Ketelaar K, Schneider R, Villalobos JA, Somerville CR, Persson S, Wallace IS (2017) BRASSINOSTEROID INSENSITIVE2 negatively regulates cellulose synthesis in Arabidopsis by phosphorylating cellulose synthase 1. Proceedings of the National Academy of Sciences 114: 3533-3538
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Santoro N, Cantu S, Tornqvist C-E, Falbel T, Bolivar J, Patterson S, Pauly M, Walton J (2010) A high-throughput platform for screening milligram quantities of plant biomass for lignocellulose digestibility. BioEnergy Research 3: 93-102
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Schaefer RJ, Michno J-M, Jeffers J, Hoekenga O, Dilkes B, Baxter I, Myers CL (2018) Integrating coexpression networks with gwas to prioritize causal genes in maize. The Plant Cell 30: 2922
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Sekhon RS, Breitzman MW, Silva RR, Santoro N, Rooney WL, de Leon N, Kaeppler SM (2016) Stover composition in maize and sorghum reveals remarkable genetic variation and plasticity for carbohydrate accumulation. Frontiers in Plant Science 7
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Sekhon RS, Joyner CN, Ackerman AJ, McMahan CS, Cook DD, Robertson DJ (2019) Stalk bending strength is strongly associated with maize stalk lodging incidence across multiple environments. bioRxiv: 675793
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Sibale EM, Darrah LL, Zuber MS (1992) Comparison of two rind penetrometers for measurement of stalk strength in maize. Maydica 37: 111-114
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Smith JM, Haigh J (1974) The hitch-hiking effect of a favourable gene. Genetical Research 23: 23-35
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Sun Q, Liu X, Yang J, Liu W, Du Q, Wang H, Fu C, Li W-X (2018) Microrna528 affects lodging resistance of maize by regulating lignin biosynthesis under nitrogen-luxury conditions. Molecular Plant 11: 806-814
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Tam V, Patel N, Turcotte M, Bossé Y, Paré G, Meyre D (2019) Benefits and limitations of genome-wide association studies. Nature Reviews Genetics 20: 467-484
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Tobimatsu Y, Schuetz M (2019) Lignin polymerization: how do plants manage the chemistry so well? Current Opinion in Biotechnology 56: 75-81

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, Pimentel H, Salzberg SL, Rinn JL, Pachter L (2012) Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. Nature protocols 7: 562-578

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Turner TL, Miller PM (2012) Investigating natural variation in Drosophila courtship song by the evolve and resequence approach. Genetics 191: 633

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Turner TL, Stewart AD, Fields AT, Rice WR, Tarone AM (2011) Population-based resequencing of experimentally evolved populations reveals the genetic basis of body size variation in Drosophila melanogaster. PLOS Genetics 7: e1001336

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Undersander DJ, Bauman LF, Lechtenberg VL, Zuber MS (1977) Effect of cyclic selection for high and low crushing strength on rind, pith, and whole stalk composition in corn. Crop Science 17: 732-734

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Vanholme B, Vanholme R, Turumtay H, Goeminne G, Cesarino I, Goubet F, Morreel K, Rencoret J, Bulone V, Hooijmaijers C, De Rycke R, Gheysen G, Ralph J, De Block M, Meulewaeter F, Boerjan W (2014) Accumulation of N-acetylglucosamine oligomers in the plant cell wall affects plant architecture in a dose-dependent and conditional manner. Plant Physiology 165: 290-308

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Vanholme R, Demedts B, Morreel K, Ralph J, Boerjan W (2010) Lignin biosynthesis and structure. Plant Physiology 153: 895

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Vlachos C, Kofler R (2019) Optimizing the power to identify the genetic basis of complex traits with Evolve and Resequence studies. bioRxiv: 583682

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Vogt T (2010) Phenylpropanoid biosynthesis. Molecular Plant 3: 2-20

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Wang J-J, Tharayil N, Chow AT, Suseela V, Zeng H (2015) Phenolic profile within the fine-root branching orders of an evergreen species highlights a disconnect in root tissue quality predicted by elemental- and molecular-level carbon composition. New Phytologist 206: 1261-1273

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Weir BS, Cockerham CC (1984) Estimating F-statistics for the analysis of population structure. Evolution 38: 1358-1370

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Xie L, Yang C, Wang X (2011) Brassinosteroids can regulate cellulose biosynthesis by controlling the expression of CESA genes in Arabidopsis. Journal of Experimental Botany 62: 4495-4506

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Yang D, Cai T, Luo Y, Wang Z (2019) Optimizing plant density and nitrogen application to manipulate tiller growth and increase grain yield and nitrogen-use efficiency in winter wheat. PeerJ 7: e6484

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ye DL, Zhang YS, Al-Kaisi MM, Duan LS, Zhang MC, Li ZH (2016) Ethephon improved stalk strength associated with summer maize adaptations to environments differing in nitrogen availability in the north China plain. The Journal of Agricultural Science 154: 960-977

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

You FM, Xiao J, Li P, Yao Z, Jia G, He L, Kumar S, Soto-Cerda B, Duguid SD, Booker HM, Rashid KY, Cloutier S (2018) Genome-wide association study and selection signatures detect genomic regions associated with seed yield and oil quality in flax. International Journal of Molecular Sciences 19: 2303

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Young MD, Wakefield MJ, Smyth GK, Oshlack A (2010) Gene ontology analysis for RNA-seq: accounting for selection bias. Genome biology 11: R14

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Zhang Y, Legland D, El Hage F, Devaux M-F, Guillon F, Reymond M, Méchin V (2019) Changes in cell walls lignification, feruloylation and p-coumaroylation throughout maize internode development. PLOS One 14: e0219923

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Zhang Z, Zhang X, Lin Z, Wang J, Liu H, Zhou L, Zhong S, Li Y, Zhu C, Lai J, Li X, Yu J, Lin Z (2020) A large transposon insertion in the stiff1 promoter increases stalk strength in maize. Plant Cell 32: 152

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Zhou Z, Jiang Y, Wang Z, Gou Z, Lyu J, Li W, Yu Y, Shu L, Zhao Y, Ma Y, Fang C, Shen Y, Liu T, Li C, Li Q, Wu M, Wang M, Wu Y, Dong Y, Wan W, Wang X, Ding Z, Gao Y, Xiang H, Zhu B, Lee S-H, Wang W, Tian Z (2015) Resequencing 302 wild and cultivated accessions identifies genes related to domestication and improvement in soybean. Nature Biotechnology 33: 408

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Zuber M, Grogan C (1961) A new technique for measuring stalk strength in corn. Crop Science 1: 378-380

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Zuber MS, Colbert TR, Darrah LL (1980) Effect of recurrent selection for crushing strength on several stalk components in maize. Crop Science 20: 711-717

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Zuber MS, Kang MS (1978) Corn lodging slowed by sturdier stalks. Crops Soils 30: 13-15

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title