Single-nucleus transcriptomics reveals functional compartmentalization in syncytial

skeletal muscle cells

Minchul Kim[1,4], Vedran Franke[2,4], Bettina Brandt[1], Simone Spuler[3], Altuna Akalin[2,*] and

Carmen Birchmeier[1,*]

[1] Developmental Biology/Signal Transduction, Max Delbrueck Center for Molecular Medicine,

Berlin, Germany

[2] Bioinformatics, Berlin Institute for Medical Science Biology, Max Delbrueck Center for

Molecular Medicine, Berlin, Germany

[3] Muscle Research Unit, Experimental and Clinical Research Center, Charité Medical Faculty

and Max Delbrueck Center for Molecular Medicine, Berlin, Germany

[4] These authors contributed equally to this work

[*] Co-correspondending authors: cbirch@mdc-berlin.de; altuna.akalin@mdc-berlin.de

Abstract

All cells must compartmentalize their intracellular space in order to properly function[1]. Syncytial

cells face an additional challenge to this fundamental problem, which is to coordinate gene

expression programs among many nuclei. Using the skeletal muscle fiber as a model, we

systematically investigated the functional heterogeneity among nuclei inside a syncytium. We

performed single nucleus RNA sequencing (snRNAseq) of isolated myonuclei from uninjured

and regenerating muscle, which revealed a remarkable and dynamic heterogeneity. We identified

distinct nuclear subtypes unrelated to fiber type diversity, completely novel subtypes as well as

the expected neuromuscular[2-5] and myotendinous[6-8] junction subtypes. snRNAseq of fibers from

the *Mdx* dystrophy mouse model uncovered additional nuclear populations. Specifically, we

identified the molecular signature of degenerating fibers and found a nuclear population that

expressed genes implicated in myofiber repair. We confirmed the existence of this population in

patients with muscular dystrophy. Finally, modifications of our approach revealed an astonishing

compartmentalization inside the rare and specialized muscle spindle fibers. In summary, our

work shows how regional transcription shapes the architecture of multinucleated syncytial

muscle cells, and provides an unprecedented roadmap to molecularly dissect distinct

compartments of the muscle.

Multinucleated skeletal muscle cells possess functionally distinct nuclear compartments to meet the needs for specialized gene expression in different parts of the large syncytium. The best documented nuclear subtype locates at the neuromuscular junction (NMJ) where the nerve instructs myonuclei to express genes involved in synaptic transmission[2-5]. Another specialized compartment is found at the myotendinous junction (MTJ) where the myofibers attach to the tendon. It has been shown that specialized cell adhesion and cytoskeletal proteins are enriched at the MTJ to allow force transmission[6-8]. However, whether the corresponding transcripts are specifically produced from MTJ myonuclei is often unclear, especially in mammals. More generally, a systematic analysis of the heterogeneity among myonuclei is lacking, and we do not know what kinds of nuclear subtypes exist. Such knowledge will provide important insights into how myofibers can orchestrate their many functions.

Previous studies on gene expression in the muscle relied on the analysis of selected candidates by *in situ* hybridization or on profiling the entire muscle tissue. The former is difficult to scale up, whereas the latter averages the transcriptomes of all nuclei. More recently, several studies have used single cell approaches to reveal the cellular composition of the entire muscle tissue[9-11]. However, these approaches did not sample nuclei in the myofiber and thus did not investigate myonuclear transcriptomes. Single-nucleus RNA-Seq (snRNAseq) using cultured human myoblasts failed to detect transcriptional heterogeneity among myotube nuclei[12], underscoring the importance of studying the heterogeneity in an *in vivo* context where myofibers interact with surrounding cell types.

Single-nucleus RNA-Seq analysis of uninjured and regenerating muscles

We genetically labeled mouse myonuclei by crossing a myofiber-specific Cre driver (*HSA-Cre*) with a Cre-dependent H2B-GFP reporter. H2B-GFP is deposited at the chromatin, which allows us to isolate single myonuclei using flow cytometry. Nuclei of regenerating fibers were also efficiently labeled 7 days after cardiotoxin-induced injury (7 days post injury; 7 d.p.i.) (Extended Data Fig. 1a). We confirmed the efficiency and specificity of the labeling (Extended Data Fig. 1a and 1b). H2B-GFP was absent in endothelia (Cd31+), Schwann cells (Egr2+), tissue resident macrophages (F4/80+) and muscle stem cells (Pax7+) (Extended Data Fig. 1c).

We next established a protocol for rapid isolation of intact myonuclei. Conventional methods
involve enzymatic dissociation of muscle fibers at 37°C, which causes secondary changes in
gene expression[13,14]. In contrast, our procedure uses mechanical disruption on ice and takes only
20 minutes from dissection to flow cytometry. Indeed, we did not detect nuclear populations
expressing stress-induced genes in the subsequent analysis (data not shown).

For snRNAseq profiling, we used the CEL-Seq2 technology[15], which has superior gene detection
sensitivity over other existing methods[16]. We used a stringent filter and considered only those
genes that were detected in at least 5 nuclei for further analysis. As a result, we detected 1000-
2000 genes per nucleus and ~5000 genes per cluster. Throughout our study, we extensively
tested multiple algorithms and parameters, the choice of which did not affect any of the
biological findings. We analyzed nuclei from uninjured (476 nuclei) and regenerating *tibialis*
anterior (TA) muscle (7 and 14 d.p.i., 958 and 1,675 nuclei, respectively). T-distributed
stochastic neighbor embedding (t-SNE) projection of these datasets revealed heterogeneity
among myonuclei (Fig. 1a). All nuclei expressed high levels of *Ttn*, a pan-muscle marker (Fig.
1b). The TA muscle contains three different fiber types (IIA - intermediate, IIB - very fast and
IIX - fast) that express distinct myosin genes. The major population (cluster 1) could be sub-
divided into nuclei from distinct fiber types (Fig. 1b); *Myh1*- (IIX) or *Myh4* (IIB)-positive nuclei
were most abundant and present roughly in a ratio of 1:1. *Myh2* (IIA)-expressing nuclei
represented a minor population, consistent with the reported proportion of fiber types[17].

To understand the molecular processes associated with regeneration, we selected top ~50 genes
that distinguish each time point and performed gene ontology (GO) term analysis. Genes
expressed specifically in uninjured muscle were generally associated with metabolism (e.g.
Gdap1, Gpx3 and *Smox*) and negative regulation of signal transduction (e.g. *Igfbp5*). Genes
expressed at 7 d.p.i. were associated with processes related to Pol II transcription (e.g. Med16),
suggesting that fiber growth is occurring. Indeed, genes known to induce fiber growth such as
Igf1 and Mettl21c were up-regulated at 7 d.p.i[18,19]. Lastly, genes expressed at 14 d.p.i. were
enriched for terms like muscle contraction and sarcomere organization (e.g. *Actn2, Lmod2* and
Dmd), suggesting that normal muscle function is being recovered at late regeneration stage.

In addition to the major population, we detected small nuclear populations with very distinct transcriptomes (clusters 2-8, Fig. 1c). Different myosin genes were evenly distributed in these, indicating that the heterogeneity is not caused by fiber type differences (Fig. 1b). We first searched for and identified a cluster specifically expressing known NMJ marker genes such as *Prkar1a, Chrne, Chrna*1, *Ache, Colq, Utrn* and *Lrp4* (cluster 4; Fig. 1d and 1e)[20]. Further, our data identified other genes not previously known to be specifically expressed at the NMJ such as *Ufsp1*, *Vav3* and *Ablim2.* (Fig. 1e). Double fluorescence *in situ* hybridization (FISH) of newly identified markers and *Prkar1a* confirmed their specific expression at the NMJ (Fig. 1f).

Two distinct nuclear populations at the myotendinous junction

We found two nuclear populations that expressed MTJ-related genes, and designated them MTJ-A and MTJ-B (Fig. 2a and 2b). MTJ-A expressed genes whose protein products are known to be enriched at the MTJ (e.g. *Itgb1*)[21] as well as specific collagens (e.g. *Col22a1* and *Col24a*). *Col22a1* has been functionally characterized in zebrafish using morpholino knockdown causing a disruption of MTJ development[22]. MTJ-B nuclei expressed an alternative set of collagens that are known to be deposited at the MTJ such as *Col6a1*, *Col6a3* and *Col1a2*[23]. *Col6a1* expression was particularly notable because its mutation causes Bethlem myopathy, which is characterized by deficits at the MTJ[24].

We validated the two top marker genes of MTJ-A (*Tigd4* and *Col22a1*) using single molecule FISH (smFISH), and observed their expression in nuclei at the fiber endings (Fig. 2c and Extended Data Fig. 2). These transcripts were exclusively expressed from H2B-GFP positive myonuclei and present only at the MTJ. Their expression became much more pronounced at 14 d.p.i. compared to uninjured muscle, indicating that at this stage of regeneration the MTJ is re-forming (Fig. 2c). To visualize heterogeneity within the syncytium, we isolated single fibers and performed double FISH (Fig. 2d). As expected, *Tigd4* FISH signals were detected at the fiber ends, whereas *Ufsp1* transcripts appeared at the middle of the fiber where the NMJ is located.

We detected markers of MTJ-B (*Pdgfrb* and *Col6a3* transcripts) expressed from H2B-GFP nuclei at the MTJ in both uninjured muscle and at 14 d.p.i. (Extended Data Fig. 3). Unlike the genes of MTJ-A, these transcripts were also found in H2B-GFP negative nuclei located outside

128   the fiber and distally to the MTJ (Extended Data Fig. 3). These signals likely originate from

129   interstitial fibroblasts or muscle stem cells[9,10,25]. Unlike MTJ-A, MTJ-B nuclei were not found in

130   every fiber. In addition, we confirmed that Ebf1 protein, another marker of MTJ-B, is present in

131   H2B-GFP positive myonuclei (Fig. 2c).

Identification of unprecedented nuclear populations

Other clusters represent novel myonuclear compartments. The top markers of cluster 5 were

maternally imprinted lncRNAs that are located at one genomic locus (also known as Dlk1-Dio3

locus) (Fig. 3a). This locus encodes a very large number of microRNAs expressed from the

maternal allele, among them microRNAs known to target transcripts of mitochondrial proteins

encoded by the nucleus[26]. smFISH against one of the lncRNAs (*Rian*) showed a clear and strong

expression in a subset of myonuclei (Fig. 3b), which was observed regardless of the animal's sex

(data not shown). smFISH on isolated fibers showed dispersed localization of *Rian* expressing

nuclei without clear positional preference (Fig. 3c). A previous study reported that the Dlk1-

Dio3 locus becomes inactive during myogenic differentiation[27]. However, our results show that

some myonuclei retain the expression, which might be important to shape the metabolisms of the

fiber.

Intriguingly, cluster 2 contained many genes that function in endoplasmic reticulum (ER)-

associated protein translation and trafficking (Fig. 3a). Tmem170a induces formation of ER

sheets, which is associated with active protein translation[28], and Rab40b is known to localize to

the Golgi/endosome and regulate trafficking[29]. Furthermore, srpRNA (signal recognition particle

RNA), an integral component of ER-bound ribosomes[30], was markedly enriched in this

population (Fig. 3a). The enrichment of srpRNA was also observed when expression of repeat

elements was quantified (Extended Data Fig. 4). smFISH against *Gssos2*, one of the marker

transcripts, showed a heterogeneous and strong expression in a subset of myonuclei (Fig. 3b).

Re-clustering of the major population identified in Figure 1a revealed additional nuclear

subpopulations with distinct transcriptomes (clusters 1a and 1b; Fig. 3d, 3e and Extended Data 5).

We verified cluster 1a using smFISH against the top marker gene *Muc13*. Myonuclei expressing

Muc13 were always located at the very outer part of the muscle tissue near the perimysium (Fig.

3f and 3g). One ultrastructural study using bovine muscle suggested that myofibers and

perimysium establish specialized adhesion structures[31], and we suggest that we have detected the

myonuclear compartment participating in this process.

snRNAseq of fibers in *Mdx* dystrophy model

To begin to understand whether and how myonuclei heterogeneity is altered in muscle disease,

we conducted snRNAseq on *Mdx* fibers (1,939 nuclei), a mouse model of Duchenne muscular

dystrophy (DMD) (Fig. 4a). DMD is a degenerative muscle disease caused by mutations in the

dystrophin gene, which is also mutated in *Mdx* mice. To examine how the *Mdx* transcriptome is

related to those of uninjured and regenerating muscle, we calculated gene signature scores of

each nucleus based on the top 50 genes that distinguish different time points after injury. This

showed that the overall gene signature of *Mdx* resembles nuclei of the regenerating muscle (14

d.p.i.) (Fig. 4b).

We next tried to identify NMJ and MTJ populations in *Mdx* fibers. However, nuclei with strong

NMJ signature could not be identified. Consistently, double smFISH against two NMJ markers

(*Ufsp1* and *Prkar1a1*) showed that the strict co-localization seen in control muscle was lost, but

these genes were instead expressed in a dispersed manner in *Mdx* muscles (Extended Data Fig. 6).

Notably, the histological structure of the NMJ is known to be fragmented in *Mdx* mice[32,33], and

our data suggest that also postsynaptic nuclei are incompletely specified. The histology of the

MTJ was reported to be disrupted in patients and mice with *Dystrophin* mutations[34-36].

Interestingly, we did not find nuclei with an MTJ-B signature in *Mdx* mice, whereas MTJ-A

nuclei were present but did not cluster in the t-SNE map (Extended Data Fig. 6). Nevertheless,

marker genes of MTJ-A (*Tigd4* and *Col22a1*) robustly labeled the MTJ of *Mdx* muscle. We

interpret that fiber-level heterogeneity (e.g. dying fibers, newly formed fibers, fibers in various

stages of maturity, repairing fibers) strongly drives the shape of the t-SNE map in *Mdx*,

interfering with the clustering of MTJ-A nuclei.

Emergence of new nuclear populations in *Mdx* dystrophy model

Several populations newly emerged in the *Mdx* muscle (especially clusters 1, 3 and 5), and we
further characterized two of these. Cluster 3 was highly enriched in various non-coding
transcripts (Fig. 4c and Extended Data 7a) and further experiments demonstrated that these
transcripts were located in dying fibers. In particular, staining of mouse IgG that identifies
degenerating fibers with leaky membranes demonstrated the expression of these ncRNAs in
IgG+ fibers (Fig. 4c and Extended Data Fig. 7b). Further, fibers strongly expressing these
ncRNAs were highly infiltrated with H2B-GFP negative cells likely corresponding to
macrophages (Extended Data Fig. 7c). Whether the ncRNAs are the consequence or active
contributors to fiber death needs further investigation.

Marker genes of cluster 5 showed high enrichment of ontology terms related to human muscle
disease (Fig. 4d). Indeed, many top marker genes were previously reported to be mutated in
human myopathies (*Klhl40*, *Flnc* and *Fhl1*)[37-39] or to directly interact with proteins whose
mutation causes disease (Ahnak interacts with Dysferlin, and Hsp7b or Xirp1 with Flnc)[40-42].
Combinatorial smFISH in tissue sections confirmed their co-expression in a subset of nuclei of
Mdx muscle, but such co-expressing nuclei were not present in control muscle (Fig. 4e, Extended
Data Fig. 8a and 8b). Further, nuclei expressing these genes were frequently closely spaced. We
also verified that *FLNC* and *XIRP1* were co-expressed in muscle tissues of Becker's patients, a
muscular dystrophy caused by *DYSTROPHIN* mutation, but not in healthy human muscle (Fig.
4f and Extended Data Fig. 8a).

Previous studies have established that Flnc and Xirp1 proteins localize to sites of myofibrillar
damage to repair such insults[41,43,44], whereas Dysferlin, an interaction partner of Ahnak,
functions during repair of muscle membrane damage[45]. Our analysis shows that these genes are
transcriptionally co-regulated, and suggests that this occurs in response to micro-damage. To
substantiate that this signature is not specific to muscular dystrophy caused by *Dystrophin*
mutation, we investigated whether these nuclei are also present in *Dysferlin* deficient muscle
where the continuous micro-damage to the membrane is no longer efficiently repaired[45]. Again,
we observed nuclei co-expressing *Flnc* and *Xirp1* in this mouse disease model and in biopsies
from human patients (Extended Data Fig. 8c and 8d). We propose that this cluster represents a
'repair signature'.

Nuclear heterogeneity in muscle spindle fibers

In principle, our approach can be used to explore nuclear heterogeneity in specific fiber types.
We thus aimed to investigate heterogeneity in muscle spindles that detect muscle stretch and
function in motor coordination[46]. Muscle spindles are extremely rare and ~10 spindles exist in a
TA muscle of the mouse[47]. They contain bag and chain fibers, and their histology forecasts
further compartmentalization (Fig. 5a). *HSA-Cre* labels myonuclei of the spindle (Fig. 5b), but
the overwhelming number of nuclei derive from extrafusal fibers. To overcome this, we used
Calb1-Cre to specifically isolate spindle myofiber nuclei (Fig. 5b) and discovered a pronounced
transcriptional heterogeneity (Fig. 5c).

Bag fibers are slow fibers and express *Myh7b*[48,49], whereas chain fibers are fast. A cluster
expressing *Myh7b* and *Tnnt1*, a slow type troponin isoform, was assigned to mark Bag fibers. In
contrast, two clusters expressed *Myh13*, a fast type Myosin, or *Tnnt3*, a fast type troponin, which
we named Chain1 and Chain2. Strikingly, we identified a cluster that expressed overlapping but
not identical set of genes as those identified in NMJ nuclei of extrafusal fibers, e.g. *Chrne*, *Ufsp1*
and *Ache*, which we assign as the NMJ of the spindle (spdNMJ) (Fig. 5d and 5e). Furthermore,
the spindle myotendinous nuclei (spdMTJ) expressed an overlapping but not identical set of
genes as those identified in MTJ-B nuclei of extrafusal fibers (Fig. 5d and 5e). MTJ-A markers
were not detected. Notably, the clusters Bag and spdNMJ expressed the mechanosensory channel
Piezo2[50]. To verify the assignment and to define the identity of an additional large compartment
(labeled as Sens), we validated the expression of different marker genes in H2B-GFP positive
fibers of *Calb1-Cre* muscle in tissue sections (Extended Data Fig. 9) and after manual isolation
of fibers under a fluorescent dissecting microscope (Fig. 5f). smFISH of *Calcrl*, a marker of the
Sens cluster, showed specific localization in the central part of spindle fibers containing densely
packed nuclei, the site where sensory neurons innervate (Fig. 5f). In the same fiber, transcripts of
the spdNMJ marker gene *Ufsp1* located laterally as distinct foci. In contrast, *Piezo2* was
expressed throughout the lateral contractile part of the fiber, but was excluded from the central
portion. Thus, the central non-contractile part of the muscle spindle that is contacted by sensory
neurons represents a distinct fiber compartment with specialized myonuclei clearly
distinguishable from the spdNMJ.

Profiling transcriptional regulators across distinct compartments

To gain insights into the transcriptional control of the distinct nuclear compartments, we investigated the expression profile of transcription factors and epigenetic regulators (Extended Fig. 10a). Notably, Etv5 (also known as Erm), a transcription factor known to induce the NMJ transcriptome[51], was identified as one of the top factor in NMJ nuclei. To begin to assess the functionality of new candidates, we chose Ebf1, the most strongly enriched transcription activator in MTJ-B. ChIP-Seq data of Ebf1 (ENCODE project) showed that Ebf1 directly binds to about ~70% of the genes we identified as MTJ-B markers (Extended Fig. 10b). We generated a C2C12 cell line in which Ebf1 expression is induced by doxycycline (Extended Fig. 10c). RT-qPCR analysis of selected markers showed that many of them were induced in a dose-dependent manner upon Ebf1 over-expression (Extended Fig. 10d). We also verified that ColVI protein was induced by Ebf1 over-expression (Extended Fig. 10c). To validate these findings *in vivo*, we analyzed *Ebf1* mutant muscle. Using smFISH for *Ttn* to identify myonuclei, we observed a strong reduction of *Col6a3* and *Fst1l* transcripts in the *Ebf1* mutant compared to control MTJ myonuclei (Extended Fig. 10e). Their expression from non-myonuclei that also express Ebf1 was also diminished. Thus, our dataset provides a template for the identification of regulatory factors that establish or maintain these compartments.

Discussion

Here, we used single-nucleus sequencing to define the transcriptome of myofiber nuclei, which identified nuclear populations at anatomically distinct locations such as NMJ, MTJ and an extremely rare nuclear population at the perimysium. Intriguingly, we found two distinct populations at the MTJ in the uninjured muscle, but only one of these were present in *Mdx* and spindle fibers. Moreover, we identified two new nuclear populations that are present throughout the fiber. Thus, myonuclear populations are not always associated with distinctive anatomical features. How these nuclear subtypes emerge and are regulated, e.g. whether they are associated with other cell types in the muscle tissue, induced by certain local signals or arise stochastically, needs further study. By investigating the *Mdx* dystrophy model, we identified the molecular

signature of degenerating fibers and a transcriptional program that appears to be associated with
fiber repair. These newly identified gene signatures might be useful for a quantitative and rapid
assessment of muscle damage in the clinic. Lastly, by restricting recombination to muscle
spindle fibers, we demonstrated compartmentalization inside these rare fibers, especially the
presence of a distinct compartment at the site of innervation by proprioceptive sensory neurons.
Taken together, our results show how regional transcription shapes the architecture of
multinucleated skeletal muscle cells, reveal the complexity of gene expression regulation in the
syncytium, and provide a comprehensive resource for studying distinct muscle compartments.

Given the large size and complexity of the muscle tissue, the full diversity of myonuclei likely
needs further exploration. In this aspect, our strategy of genetic labeling to enrich nuclei of
interest appears advantageous as demonstrated by our success with the identification of
perimysium and muscle spindle myonuclei, which are very rare and therefore hard to detect
when the entire tissue is used. Although we concentrated here on the analysis of uninjured and
regenerating muscle in health and disease, our strategy should be promising in identifying new
nuclear subtypes in other contexts. In the accompanying manuscript, snRNAseq successfully
discovered novel nuclear subtypes during early development and in aged muscle (Petrany MJ,
Swoboda CO, Sun C, Chetal K, Chen X, Weirauch MT, Salomonis N, Millay DP. Single-nucleus
RNA-seq identifies transcriptional heterogeneity in multinucleated skeletal myofibers. Submitted
to *BioRxiv*). More generally, our approach should be instrumental in investigating other syncytial
cell types such as the placental trophoblasts or osteoclasts. In addition to charting and dissecting
new nuclear populations, another important question will be to understand the next layers
towards establishing specialized compartments such as RNA/protein trafficking.

Methods

Mouse lines and muscle injury

All experiments were conducted according to regulations established by the Max Delbrück Centre for Molecular Medicine and LAGeSo (Landesamt für Gesundheit und Soziales), Berlin. *HSA-Cre*, *Calb1-IRES-Cre* and *Mdx* mice were obtained from the Jackson laboratory. *Rosa26-Lsl-H2B-GFP* reporter line was a kind gift from Martin Goulding (Salk institute) and was described[52]. For experiments regarding uninjured and regenerating muscles, homozygous *Rosa26-LSL-H2B-GFP* mice with heterozygous *HSA-Cre* were used. For *Calb1-IRES-Cre* and *Mdx* experiments, the *H2B-GFP* allele was heterozygous. Nuclei were isolated from muscle of 2.5 months old mice. Genotyping was performed as instructed by the Jackson laboratory. For genotyping of the *Rosa26-Lsl-H2B-GFP* reporter, the following primers were used. Rosa4; 5'-TCA ATGGGCGGGGGTCGTT-3', Rosa10; 5'-CTCTGCTGCCTCCTGGCTTCT-3', Rosa11; 5'-CGAGGCGGATCACAAGCAATA-3'. Ebf1 mutants[53] and Dysferlin mis-sense mutants[54] were described. To induce muscle injury, 30µl of cardiotoxin (10µM, Latoxan, Porte les Vaence, France) was injected into the tibialis anterior (TA) muscle.

Isolation of nuclei from TA muscle

For each sorting, we pooled two TA muscles from two individual mice. Dissected muscle was cut into small pieces, suspended in 1ml hypotonic buffer (250mM sucrose, 10mM KCl, 5mM $MgCl_2$, 10mM Tris-HCl pH 8.0, 25mM HEPES pH 8.0, 0.2mM PMSF and 0.1mM DTT supplemented with protease inhibitor tablet from Roche), and transferred to a 'Tissue homogenizing CKMix' tube (Bertin instruments) containing ceramic beads of mixed size. After incubating on ice for 15 minutes, samples were homogenized with Precellys 24 tissue homogenizer (Bertin instruments) for 20 seconds at 5,000 rpm. Homogenized samples were passed once through 70µm filter (Sysmex), twice through 20µm filter (Sysmex), and once though 5ml filter cap FACS tube (Corning 352235). DAPI (Sigma) was added to final concentration of 300nM to label DNA. GFP and DAPI double positive nuclei were sorted using ARIA Sorter III (BD).

Isolation of nuclei from muscle spindle

We pooled all TA muscles from 3-4 mice for each experiment. Two different protocols were
employed. The first procedure (protocol 1 in Extended Data Fig. 9a) employed the same
procedure as the one used to isolate TA nuclei, pooling up to 4 TA muscles/homogenizer tube,
and yielded 96 spindle nuclei.

The second procedure (protocol 2 in Extended Data Fig. 9a), we shorten isolation time. Briefly,
0.1% Triton X-100 was added to the hypotonic buffer to solubilize the tissue debris, which are
otherwise detected as independent particles during FACS. After homogenization and filtration,
nuclei were pelleted by centrifuging at 200g for 10 minutes at 4°C. After discarding the
supernatants, nuclei were resuspended in 300ul hypotonic buffer and passed through the FACS
tube. This yielded 192 spindle nuclei.

Library generation and sequencing

96-well plates for sorting were prepared using an automated pipetting system (Integra Viaflo).
Each well contained 1.2µl of master mix (13.2µl 10% Triton X-100, 25mM dNTP 22µl, ERCC
spike-in 5.5µl and ultrapure water up to 550µl total) and 25ng/µl barcode primers. Plates were
stored at -80°C until use.

After sorting single nuclei into the wells, plates were centrifuged at 4,000g for 1 minute,
incubated on incubated on 65°C for 5 minutes and immediately cooled on ice. Subsequent library
generation was performed using the CEL-Seq2 protocol as described by Hashimshony et al[15].
After reverse transcription and second strand synthesis, products of one plate were pooled into
one tube and cleaned up using AMPure XP beads (Beckman Coulter). After in vitro transcription
and fragmentation, aRNA was cleaned up using RNAClean XP beads (Beckman Coulter) and
eluted in 7µl ultrapure water. 1ul of aRNA was analyzed on Bioanalyzer RNA pico chip for a
quality check. To construct sequencing library, 5µg aRNA was used for reverse transcription
(Superscript II, Thermofisher) and library PCR (Phusion DNA polymerase, Thermofisher). After
clean up using AMPure XP beads, 1µl sample was ran on Bioanalyzer using a high sensitivity

DNA chip to measure size distribution, which we demonstrated the presence of a peak of around

400bp length. Sequencing was performed using Illumina HiSeq2500 (for uninjured and

regenerating muscle) or HiSeq4000 (for *Mdx* and spindle).

Bioinformatics analysis

Single nucleus RNA-Sequencing data was processed using PiGx-scRNAseq pipeline - a

derivative of a CellRanger pipeline, but enabling deterministic analysis reproducibility (version

0.1.3)[55]. In short, polyA sequences were removed from the reads. The reads were mapped to the

genome using STAR[56]. Number of nuclei, for each sample, was determined using dropbead[57].

Finally, a combined digital expression matrix was constructed, containing all sequenced

experiments.

Digital expression matrix post processing was performed using Seurat[58]. The raw data was

normalized using the NormalizeData function. The expression of each nucleus was then

normalized by multiplying each gene with the following scaling factor: 10000/(total number of

raw counts), log(2) transformed, and subsequently scaled. Number of detected genes per nucleus

was regressed out during the scaling procedure.

Variable genes were defined using the FindVariableGenes function with the default parameters.

Samples were processed in three groups with differing parameters. Samples originating from

uninjured, 7 d.p.i. and 14 d.p.i. were processed as one group, samples from the *Mdx* mouse as the

second group and muscle spindle nuclei as the third group.

For samples of the first group, nuclei with less than 500 detected genes were filtered out.

Subsequently, genes which were detected at least in 5 nuclei were kept for further analysis. To

remove the putative confounding effect between time of sample preparation and biological

variable (injury), the processed expression matrices were integrated using the

FindIntegrationAnchors function with reciprocal PCA, from the Seurat package. The function

uses within batch covariance structure to align multiple datasets.

The integration was based on 2000 top variable features, and first 30 principal components. T-
SNE was based on the first 15 principal components. Outlier cluster detection was done with
dbscan ([10.18637/jss.v091.i01](10.18637/jss.v091.i01).), with the following parameters eps = 3.4, minPts = 20.

Mdx samples were processed by filtering out all nuclei with less than 100 detected genes. Top
100 most variable genes were used for the principal component analysis. t-SNE and Leuvain
clustering were based on the first 10 principal components. Resolution parameter of 1 was used
for the Leuvain clustering.

Spindle cell samples were processed by filtering out all cells with less than 300 detected genes.
Top 200 most variable genes were used for the principal component analysis. t-SNE and Leuvain
clustering were based on the first 15 principal components. Resolution parameter of 1 was used
for the Leuvain clustering.

For all datasets, multiple parameter sets were tested during the analysis, and the choice of
parameters did not have a strong influence on the results and the derivative biological
conclusions. Genes with cluster specific expression were defined using Wilcox test, as
implemented in the FindAllMarkers function from the Seurat package. Genes that were detected
in at least 25% of the cells in each cluster were selected for differential gene expression analysis.

NMJ Nuclei Definition: NMJ nuclei were identified based on the expression of three previously
known markers (*Prkar1a*, *Chrne* and *Ache*), using the AddModuleScore function from the Seurat
package. All cells with a score greater than 1 were selected as NMJ positive cells. NMJ marker
set was expanded by comparing the fold change of gene expression in averaged NMJ positive to
NMJ negative cells.

MTJ A and B Nuclei Definition: The original gene sets were extracted from cluster specific
genes detected in the uninjured, 7 d.p.i. and 14 d.p.i. experiment. Cells were scored as MTJ A/B
using the aforementioned gene set, with the AddModuleScore function. All cells which had a
respective score greater than 1 were labeled as MTJ A/B positive cells.

Mdx nuclei scoring by uninjured and regenerating signatures: First, gene signatures specific for
each time point were selected using the FindAllMarkers function from the Seurat library, using

the default parameters. *Mdx* samples were scored using the top 50 genes per time point with the

AUCell method (10.1038/nmeth.4463).

Repetitive element annotation was downloaded from the UCSC Browser database

(10.1093/nar/gky1095) on 21.01.2020. Pseudo-bulk bigWig tracks were constructed for each

defined cluster in uninjured, 7.d.p.i, and 14.d.p.i. The tracks were normalized to the total number

of reads. Repetitive element expression was quantified using the ScoreMatrixBin function from

the genomation (10.1093/bioinformatics/btu775) package, which calculates the average per-base

expression value per repetitive element. The expression was finally summarized to the repetitive

element family (class) level by calculating the average expression of all repeats belonging to the

corresponding family (class).

Transcription factor compendium, used in all analyses was downloaded from AnimalTFDB

(https://doi.org/10.1093/nar/gky822). The expression map (Fig. 5d) for spindle fibers was created

using the DotPlot from the Seurat package on a selected set of cluster specific genes. Gene

ontology analysis was performed using the Enrichr program (10.1093/nar/gkw377).

(http://amp.pharm.mssm.edu/Enrichr/).

Preparation of tissue sections

421   Freshly isolated TA muscles were embedded in OCT compound and processed as previously

422   described[59]. Frozen tissue blocks were sectioned to 12-16µm thickness, which were stored at -

423   80°C until future use.

Single-molecule FISH (RNAscope)

RNAscope_V2 kit was used according to manufacturer's instructions (ACD/bio-techne). We

used Proteinase IV. When combined with antibody staining, after the last washing step of RNA

Scope, the slides were blocked with 1% horse serum and 0.25% BSA in PBX followed by

primary antibody incubation overnight on 4°C. The subsequent procedures were the same as

regular immunohistochemistry. Slides were mounted with Prolonged Antifade mounting solution

(Thermofisher). The following probes were used in this study; Ttn (483031), Rian (510531),

Vav3 (437431), Col6a3 (552541), Egr1 (423371), Gm10800 (479861) and Calcrl (452281). The

following probes were newly designed; Prkar1a (c1), Tigd4 (c1), Muc13 (c1), Gssos2 (c1), Flnc

(c1), Klhl40 (c1), Col22a1 (c2), Ufsp1 (c2), Gm10801 (c2), Xirp1 (c2), Fstl1 (c2), human Flnc

(c2), Ablim2 (c3) and human Xirp1 (c3).

Preparation of fluorescent in situ hybridization (FISH) probes

Probes of 500-700bp length spanning exon-exon junction parts were designed using software in

NCBI website. Forward and reverse primers included Xho1 restriction site and T3 promoter

sequence, respectively. cDNA samples prepared from E13.5-E14.5 whole embryos were used to

amplify the target probes using GO Taq DNA polymerase (Promega). PCR products were cloned

into pGEM-T Easy vector (Promega) according to manufacturer's guideline, and the identity of

the inserts was confirmed by sequencing. 2µg of cloned plasmid DNA was linearized, 500ng

DNA was subjected to *in vitro* transcription with T3 polymerase and DIG- or FITC- labeled

ribonucleotides (All Roche) for 2 hours at 37°C. Synthesized RNA probes were purified using

RNeasy kit (Qiagen). Probes were eluted in 50µl ultrapure water (Sigma), and 50µl formamide

was added. We checked the RNA quality and quantity by loading 5µl RNA to 2% agarose gel.

Until future use, probes were stored in -80°C. The followings are the annealing sequences of the

FISH probes used in this study. Prkar1a For; 5'-CTGCACCACTGAATTACCGTA-3', Prkar1a

Rev; 5'-ACAAGATAAGTGTGCCTCGTT-3', Ufsp1 For; 5'-CACTGCCCGTAGCTAGTCC-

3', Ufsp1 Rev; 5'-CAGGCCCAAACTTGATCGC-3', Phldb2 For; 5'-

GATGAGGAGTCTGTGTTCGAGG-3', Phldb2 Rev; 5'-GCCTCTCTAGCTTTTCCCTCTC-3',

Col6a3 For; 5'-GCAGGCTAGTGTGTTCTCGT-3', Col6a3 Rev; 5'-

CGTTGTCGGAGCCATCCAAA, Pdgfrb For; 5'-AGTGATACCAGCTTTAGTCCT-3', Pdgrfb

Rev; 5'-GTTCACACTCACTGACACGTT-3', Tigd4 For; 5'-

CAATGGTTGGCTGGATCGTTTT-3' and Tigd4 Rev; 5'-CTTTGGAGACCTGAATCCTGCT-

3'.

Fluorescence in situ hybridization (FISH) and immunohistochemistry

Basic procedure for FISH was described before[59] with minor modifications to use fluorescence

for final detection. After hybridizing the tissue sections with DIG- labeled probes, washing,

RNase digestion and anti-DIG antibody incubation, amplification reaction was carried out using

TSA-Rhodamine (1:75 and 0.001% H_2O_2). After washing, slides were mounted with Immu-

Mount (Thermo Scientific). When applicable, GFP antibody was added together with anti-DIG

antibody.

When conducting double FISH, the tissue was hybridized with DIG- and FITC-labeled probes,

after detection of the DIG signal, slides were treated with 3% H_2O_2 for 15 minutes and then with

4% PFA for one hour at room temperature to eliminate residual peroxidase activity. The second

amplification reaction was performed using anti-FITC antibody and TSA-biotin (1:50), which

was visualized using Cy5-conjugated anti-streptavidin.

Antibodies used for this study were: GFP (Aves labs, 1:500), Col3 (Novus, 1:500), ColIV

(Millipore, 1:500), CD31-PE (Biolegend, 1:200), F4/80 (Abcam ab6640, 1:500) and Laminin

(Sigma L9393, 1:500). Pax7 (1:50) and Ebf1 (1:20) anti-serum was described[59,60]. Egr2 (1:10000)

anti-serum was a kind gift from Michael Wegner (Friedrich-Alexander University, Erlangen,

Germany). For Pax7, we used an antigen retrieval step. For this, after fixation and PBS washing,

slides were incubated in antigen retrieval buffer (diluted 1:100 in water; Vector) pre-heated to

80°C for 15 minutes. Slides were washed in PBS and continued at permeabilization step. Cy2-,

Cy3- and Cy5-conjugated secondary antibodies were purchased from Dianova.

FISH experiments using isolated single fibers

481   We isolated single extensor digitorum longus (EDL) muscle fibers as described before[61]. Isolated

482   EDL fibers were immediately fixed with 4% PFA, and were subjected hybridization in 1.5ml

483   tubes. After DAPI staining, fibers were transferred on slide glasses and mounted. Spindle fibers

484   were vulnerable to collagenase treatment. Thus, we pre-fixed the EDL tissue and peeled off

485   spindles under the fluorescent dissecting microscope (Leica).

Acquisition of fluorescence images

Fluorescence was visualized by laser-scanning microscopy (LSM700, Carl-Zeiss) using Zen

2009 software. Images were processed using ImageJ and Adobe Photoshop, and assembled using

Adobe Illustrator.

Cell culture

C2C12 cell line was purchased from ATCC, and cultured in high glucose DMEM (Gibco)

supplemented with 10% FBS (Sigma) and Penicilllin-Streptomycin (Sigma). To engineer C2C12

cells with doxycycline (Sigma) inducible Ebf1, mouse Ebf1 cDNA (Addgene) was cloned into

pLVX Tet-One Puro plasmid (Clontech), packaged in 293T cells (from ATCC) using psPAX2

and VsvG (Addgene), followed by viral transduction to C2C12 cells with 5µg/µl polybrene

(Millipore). Transduced cells were selected using 3µg/µl puromycin (Sigma).

Western blotting

Cell pellets were resuspended in NP-40 lysis buffer (1% NP-40, 150mM NaCl, 50mM Tris-Cl

pH 7.5, 1mM MgCl2 supplemented with protease (Roche) and phosphatase (Sigma) inhibitors),

and incubated on ice for 20 minutes. Lysates were cleared by centrifuging in 16,000g for 20 min

at 4°C. Protein concentration was measured by Bradford assay (Biorad), and lysates were boiled

in Laemilli buffer with beta-mercaptoethanol for 10 minutes. Denatured lysates were fractionated

by SDS-PAGE, transferred into nitrocellulose membrane, blocked with 5% milk and 0.1%

Tween-20 in PBS, and incubated overnight in 4°C with primary antibodies diluted in 5% BSA

and 0.1% Tween-20 in PBS. After three times washing with PBST, membranes were incubated

with secondary antibodies diluted in blocking solution for one hour at room temperature. After

PBST washing, membranes were developed with prime ECL (Amarsham). The antibodies used

for this study were β-actin (Cell Signaling, 1:1000), ColVI (Abcam, 1:2000) and Ebf1 (1:1000).

RT-qPCR

Cell pellets were resuspended in 1ml Trizol (Thermofisher). RNA was isolated according to

manufacturer's guideline. 1µg of isolated RNA and random hexamer primer (Thermofisher)

were used for reverse transcription using ProtoSciprt II RT (NEB). Synthesized cDNA was

diluted five times in water, and 1µl was used per one qPCR reaction. qPCR was performed using

2X Sybergreen mix (Thermofisher) and CFX96 machine (Biorad). We used β-actin for

normalization. Primers were designed using the 'Primer bank' website. Following primers were

used for this study. β-actin For; 5'-GGCTGTATTCCCCTCCATCG-3', β-actin Rev; 5'-

CCAGTTGGTAACAATGCCATGT-3', Pdgfrb For; 5'-TTCCAGGAGTGATACCAGCTT-3',

Pdgfrb Rev; 5'-AGGGGGCGTGATGACTAGG-3', Col1a1 For; 5'-

GCTCCTCTTAGGGGCCACT-3', Col1a1 Rev; 5'-CCACGTCTCACCATTGGGG-3', Col5a3

For; 5'-CGGGGTACTCCTGGTCCTAC-3', Col5a3 Rev; 5'-GCATCCCTACTTCCCCCTTG-

3', Col6a1 For; 5'-CTGCTGCTACAAGCCTGCT-3', Col6a1 Rev; 5'-

CCCCATAAGGTTTCAGCCTCA-3', Col6a3 For; 5'-GCTGCGGAATCACTTTGTGC-3',

Col6a3 Rev; 5'-CACCTTGACACCTTTCTGGGT-3', Lrp1 For; 5'-

ACTATGGATGCCCCTAAAACTTG-3', Lrp1 Rev; 5'-GCAATCTCTTTCACCGTCACA-3',

Mgp For; 5'-GGCAACCCTGTGCTACGAAT-3', Mgp Rev; 5'-

CCTGGACTCTCTTTTGGGCTTTA-3', Fstl1 For; 5'-CACGGCGAGGAGGAACCTA-3',

Fstl1 Rev; 5'-TCTTGCCATTACTGCCACACA-3', Cilp For; 5'-

ATGGCAGCAATCAAGACTTGG-3' and Cilp Rev; 5'-AGGCTGGACTCTTCTCACTGA-3'.

Human biopsies

Human muscle biopsy specimens were obtained from M. vastus lateralis. The tissue was snap

frozen under cryoprotection. Research use of the material was approved by the regulatory

agencies (EA1/203/08, EA2/051/10, EA2/175/17, Charité Universitätsmedizin Berlin, Germany).

Informed consent was obtained from the donors.

References

1   Stoeger, T., Battich, N. & Pelkmans, L. Passive Noise Filtering by Cellular
Compartmentalization. *Cell* **164**, 1151-1161, doi:10.1016/j.cell.2016.02.005 (2016).

2   Sanes, J. R. & Lichtman, J. W. Development of the vertebrate neuromuscular junction.
Annual review of neuroscience **22**, 389-442, doi:10.1146/annurev.neuro.22.1.389 (1999).
3   Hall, Z. W. & Sanes, J. R. Synaptic structure and development: the neuromuscular
junction. *Cell* **72 Suppl**, 99-121, doi:10.1016/s0092-8674(05)80031-5 (1993).
4   Jasmin, B. J., Lee, R. K. & Rotundo, R. L. Compartmentalization of acetylcholinesterase
mRNA and enzyme at the vertebrate neuromuscular junction. *Neuron* **11**, 467-477,
doi:10.1016/0896-6273(93)90151-g (1993).
5   Schaeffer, L., de Kerchove d'Exaerde, A. & Changeux, J. P. Targeting transcription to the
neuromuscular synapse. *Neuron* **31**, 15-22, doi:10.1016/s0896-6273(01)00353-1 (2001).
6   Charvet, B., Ruggiero, F. & Le Guellec, D. The development of the myotendinous
junction. A review. *Muscles, ligaments and tendons journal* **2**, 53-63 (2012).
7   Maartens, A. P. & Brown, N. H. The many faces of cell adhesion during Drosophila
muscle development. *Developmental biology* **401**, 62-74,
doi:10.1016/j.ydbio.2014.12.038 (2015).
8   Schweitzer, R., Zelzer, E. & Volk, T. Connecting muscles to tendons: tendons and
musculoskeletal development in flies and vertebrates. *Development* **137**, 2807-2817,
doi:10.1242/dev.047498 (2010).
9   Giordani, L. *et al.* High-Dimensional Single-Cell Cartography Reveals Novel Skeletal
Muscle-Resident Cell Populations. *Molecular cell* **74**, 609-621 e606,
doi:10.1016/j.molcel.2019.02.026 (2019).
10   Dell'Orso, S. *et al.* Single cell analysis of adult mouse skeletal muscle stem cells in
homeostatic and regenerative conditions. *Development* **146**, doi:10.1242/dev.174177
(2019).
11   Rubenstein, A. B. *et al.* Single-cell transcriptional profiles in human skeletal muscle.
Scientific reports **10**, 229, doi:10.1038/s41598-019-57110-6 (2020).
12   Zeng, W. *et al.* Single-nucleus RNA-seq of differentiating human myoblasts reveals the
extent of fate heterogeneity. *Nucleic acids research* **44**, e158, doi:10.1093/nar/gkw739
(2016).
13   van den Brink, S. C. *et al.* Single-cell sequencing reveals dissociation-induced gene
expression in tissue subpopulations. *Nature methods* **14**, 935-936,
doi:10.1038/nmeth.4437 (2017).
14   Machado, L. *et al.* In Situ Fixation Redefines Quiescence and Early Activation of
Skeletal Muscle Stem Cells. *Cell reports* **21**, 1982-1993,
doi:10.1016/j.celrep.2017.10.080 (2017).
15   Hashimshony, T. *et al.* CEL-Seq2: sensitive highly-multiplexed single-cell RNA-Seq.
Genome biology **17**, 77, doi:10.1186/s13059-016-0938-8 (2016).
16   Ziegenhain, C. *et al.* Comparative Analysis of Single-Cell RNA Sequencing Methods.
Molecular cell **65**, 631-643 e634, doi:10.1016/j.molcel.2017.01.023 (2017).
17   Augusto, V. *et al.* Skeletal muscle fiber types in C57BL6J mice. *Braz. J. morphol. Sci.* **21**,
2, 89-94 (2004).
18   Schiaffino, S. & Mammucari, C. Regulation of skeletal muscle growth by the IGF1-
Akt/PKB pathway: insights from genetic models. *Skeletal muscle* **1**, 4, doi:10.1186/2044-
5040-1-4 (2011).
19   Wiederstein, J. L. *et al.* Skeletal Muscle-Specific Methyltransferase METTL21C
Trimethylates p97 and Regulates Autophagy-Associated Protein Breakdown. *Cell reports*
23, 1342-1356, doi:10.1016/j.celrep.2018.03.136 (2018).

20   Tintignac, L. A., Brenner, H. R. & Ruegg, M. A. Mechanisms Regulating Neuromuscular
Junction Development and Function and Causes of Muscle Wasting. *Physiological*
reviews **95**, 809-852, doi:10.1152/physrev.00033.2014 (2015).

21   Bao, Z. Z., Lakonishok, M., Kaufman, S. & Horwitz, A. F. Alpha 7 beta 1 integrin is a
component of the myotendinous junction on skeletal muscle. *Journal of cell science* **106**
(Pt 2), 579-589 (1993).

22   Charvet, B. *et al.* Knockdown of col22a1 gene in zebrafish induces a muscular dystrophy
by disruption of the myotendinous junction. *Development* **140**, 4602-4613,
doi:10.1242/dev.096024 (2013).

23   Can, T. *et al.* Proteomic analysis of laser capture microscopy purified myotendinous
junction regions from muscle sections. *Proteome science* **12**, 25, doi:10.1186/1477-5956-
12-25 (2014).

24   Jobsis, G. J. *et al.* Type VI collagen mutations in Bethlem myopathy, an autosomal
dominant myopathy with contractures. *Nature genetics* **14**, 113-115, doi:10.1038/ng0996-
113 (1996).

25   Baghdadi, M. B. *et al.* Reciprocal signalling by Notch-Collagen V-CALCR retains
muscle stem cells in their niche. *Nature* **557**, 714-718, doi:10.1038/s41586-018-0144-9
(2018).

26   Labialle, S. *et al.* The miR-379/miR-410 cluster at the imprinted Dlk1-Dio3 domain
controls neonatal metabolic adaptation. *The EMBO journal* **33**, 2216-2230,
doi:10.15252/embj.201387038 (2014).

27   Wust, S. *et al.* Metabolic Maturation during Muscle Stem Cell Differentiation Is
Achieved by miR-1/133a-Mediated Inhibition of the Dlk1-Dio3 Mega Gene Cluster. *Cell*
metabolism **27**, 1026-1039 e1026, doi:10.1016/j.cmet.2018.02.022 (2018).

28   Christodoulou, A., Santarella-Mellwig, R., Santama, N. & Mattaj, I. W. Transmembrane
protein TMEM170A is a newly discovered regulator of ER and nuclear envelope
morphogenesis in human cells. *Journal of cell science* **129**, 1552-1565,
doi:10.1242/jcs.175273 (2016).

29   Jacob, A. *et al.* Rab40b regulates trafficking of MMP2 and MMP9 during invadopodia
formation and invasion of breast cancer cells. *Journal of cell science* **126**, 4647-4658,
doi:10.1242/jcs.126573 (2013).

30   Hainzl, T., Huang, S. & Sauer-Eriksson, A. E. Structural insights into SRP RNA: an
induced fit mechanism for SRP assembly. *Rna* **11**, 1043-1050, doi:10.1261/rna.2080205
(2005).

31   Passerieux, E. *et al.* Structural organization of the perimysium in bovine skeletal muscle:
Junctional plates and associated intracellular subdomains. *Journal of structural biology*
154, 206-216, doi:10.1016/j.jsb.2006.01.002 (2006).

32   Haddix, S. G., Lee, Y. I., Kornegay, J. N. & Thompson, W. J. Cycles of myofiber
degeneration and regeneration lead to remodeling of the neuromuscular junction in two
mammalian models of Duchenne muscular dystrophy. *PloS one* **13**, e0205926,
doi:10.1371/journal.pone.0205926 (2018).

33   Pratt, S. J. P., Valencia, A. P., Le, G. K., Shah, S. B. & Lovering, R. M. Pre- and
postsynaptic changes in the neuromuscular junction in dystrophic mice. *Frontiers in*
physiology **6**, 252, doi:10.3389/fphys.2015.00252 (2015).

34   Law, D. J. & Tidball, J. G. Dystrophin Deficiency Is Associated with Myotendinous
Junction Defects in Prenecrotic and Fully Regenerated Skeletal-Muscle. *American*
Journal of Pathology **142**, 1513-1523 (1993).
35   Banks, G. B., Judge, L. M., Allen, J. M. & Chamberlain, J. S. The Polyproline Site in
Hinge 2 Influences the Functional Capacity of Truncated Dystrophins. *PLoS genetics* **6**,
doi:ARTN e1000958

10.1371/journal.pgen.1000958 (2010).
36   Bell, C. D. & Conen, P. E. Histopathological Changes in Duchenne Muscular Dystrophy.
J Neurol Sci **7**, 529-&, doi:Doi 10.1016/0022-510x(68)90058-0 (1968).
37   Ravenscroft, G. *et al.* Mutations in KLHL40 are a frequent cause of severe autosomal-
recessive nemaline myopathy. *American journal of human genetics* **93**, 6-18,
doi:10.1016/j.ajhg.2013.05.004 (2013).
38   Vorgerd, M. *et al.* A mutation in the dimerization domain of filamin c causes a novel type
of autosomal dominant myofibrillar myopathy. *American journal of human genetics* **77**,
297-304, doi:10.1086/431959 (2005).
39   Windpassinger, C. *et al.* An X-linked myopathy with postural muscle atrophy and
generalized hypertrophy, termed XMPMA, is caused by mutations in FHL1. *American*
journal of human genetics **82**, 88-99, doi:10.1016/j.ajhg.2007.09.004 (2008).
40   Huang, Y. *et al.* AHNAK, a novel component of the dysferlin protein complex,
redistributes to the cytoplasm with dysferlin during skeletal muscle regeneration. *FASEB*
journal : official publication of the Federation of American Societies for Experimental
Biology **21**, 732-742, doi:10.1096/fj.06-6628com (2007).
41   Molt, S. *et al.* Aciculin interacts with filamin C and Xin and is essential for myofibril
assembly, remodeling and maintenance. *Journal of cell science* **127**, 3578-3592,
doi:10.1242/jcs.152157 (2014).
42   Juo, L. Y. *et al.* HSPB7 interacts with dimerized FLNC and its absence results in
progressive myopathy in skeletal muscles. *Journal of cell science* **129**, 1661-1670,
doi:10.1242/jcs.179887 (2016).
43   Leber, Y. *et al.* Filamin C is a highly dynamic protein associated with fast repair of
myofibrillar microdamage. *Human molecular genetics* **25**, 2776-2788,
doi:10.1093/hmg/ddw135 (2016).
44   Otten, C. *et al.* Xirp proteins mark injured skeletal muscle in zebrafish. *PloS one* **7**,
e31041, doi:10.1371/journal.pone.0031041 (2012).
45   Bansal, D. *et al.* Defective membrane repair in dysferlin-deficient muscular dystrophy.
Nature **423**, 168-172, doi:10.1038/nature01573 (2003).
46   Albert, Y. *et al.* Transcriptional regulation of myotube fate specification and intrafusal
muscle fiber morphogenesis. *The Journal of cell biology* **169**, 257-268,
doi:10.1083/jcb.200501156 (2005).
47   Cheret, C. *et al.* Bace1 and Neuregulin-1 cooperate to control formation and maintenance
of muscle spindles. *The EMBO journal* **32**, 2015-2028, doi:10.1038/emboj.2013.146
(2013).
48   Schiaffino, S. & Reggiani, C. Fiber types in mammalian skeletal muscles. *Physiological*
reviews **91**, 1447-1531, doi:10.1152/physrev.00031.2010 (2011).
49   Rossi, A. C., Mammucari, C., Argentini, C., Reggiani, C. & Schiaffino, S. Two
novel/ancient myosins in mammalian skeletal muscles: MYH14/7b and MYH15 are

expressed in extraocular muscles and muscle spindles. *The Journal of physiology* **588**,
353-364, doi:10.1113/jphysiol.2009.181008 (2010).

50   Coste, B. *et al.* Piezo1 and Piezo2 are essential components of distinct mechanically
activated cation channels. *Science* **330**, 55-60, doi:10.1126/science.1193270 (2010).

51   Hippenmeyer, S., Huber, R. M., Ladle, D. R., Murphy, K. & Arber, S. ETS transcription
factor Erm controls subsynaptic gene expression in skeletal muscles. *Neuron* **55**, 726-740,
doi:10.1016/j.neuron.2007.07.028 (2007).

52   Li, Y. *et al.* Molecular layer perforant path-associated cells contribute to feed-forward
inhibition in the adult dentate gyrus. *Proceedings of the National Academy of Sciences of*
the United States of America **110**, 9106-9111, doi:10.1073/pnas.1306912110 (2013).

53   Lin, H. & Grosschedl, R. Failure of B-cell differentiation in mice lacking the
transcription factor EBF. *Nature* **376**, 263-267, doi:10.1038/376263a0 (1995).

54   Malcher, J. *et al.* Exon Skipping in a Dysf-Missense Mutant Mouse Model. *Molecular*
therapy. Nucleic acids **13**, 198-207, doi:10.1016/j.omtn.2018.08.013 (2018).

55   Wurmus, R. *et al.* PiGx: reproducible genomics analysis pipelines with GNU Guix.
GigaScience **7**, doi:10.1093/gigascience/giy123 (2018).

56   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21,
doi:10.1093/bioinformatics/bts635 (2013).

57   Alles, J. *et al.* Cell fixation and preservation for droplet-based single-cell transcriptomics.
BMC biology **15**, 44, doi:10.1186/s12915-017-0383-5 (2017).

58   Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell
transcriptomic data across different conditions, technologies, and species. *Nature*
biotechnology **36**, 411-420, doi:10.1038/nbt.4096 (2018).

59   Muller, T. *et al.* The homeodomain factor lbx1 distinguishes two major programs of
neuronal differentiation in the dorsal spinal cord. *Neuron* **34**, 551-562,
doi:10.1016/s0896-6273(02)00689-x (2002).

60   Roessler, S. *et al.* Distinct promoters mediate the regulation of Ebf1 gene expression by
interleukin-7 and Pax5. *Molecular and cellular biology* **27**, 579-594,
doi:10.1128/MCB.01192-06 (2007).

61   Vogler, T. O., Gadek, K. E., Cadwallader, A. B., Elston, T. L. & Olwin, B. B. Isolation,
Culture, Functional Assays, and Immunofluorescence of Myofiber-Associated Satellite
Cells. *Methods in molecular biology* **1460**, 141-162, doi:10.1007/978-1-4939-3810-0_11
(2016).

Acknowledgements

We thank Dr. K. Song for advice and protocols on snRNAseq, E. Lowenstein and Dr. T. Muller

for critical reading of the manuscript, C. Paeseler and P. Stallerow for help with animal care (all

MDC), as well as the MDC core facilities for flow cytometry (led by Dr. Hans-Peter Rahn) and

next generation sequencing (led by Dr. Sascha Sauer). We are grateful to Dr. Martyn Goulding

(Salk institute) for providing the H2B-GFP reporter mouse line, and Drs. M. Derecka and R.

Grosschedl (MPI for Immunobiology and Epigenetics) for providing Ebf1 tissues and reagents.

This work was supported by the Helmholtz association (C.B) and AVH postdoctoral fellowship
(M.K).

Author contributions

M.K and C.B conceived the work and designed the project. M.K. led the project, performed the
experiments and analyzed the data. V.F and A.A performed the bioinformatics analysis. B.B
contributed to the experiments, especially generated the sequencing library. S.S collected and
prepared human patient biopsies and *Dysferlin* mutant mouse samples. M.K and C.B wrote the
manuscript with comments by all authors.

Competing interests: We have no conflicting interests.

Materials & correspondence: Requests for reagents should be addressed to cbirch@mdc-
berlin.de

Data availability: All the raw sequencing data have been deposited in Array Express (Accession
number: E-MTAB-8623). The data will be made public after publication.

Figure 1. Nuclear heterogeneity in uninjured and regenerating muscles. a, t-SNE map of transcripts detected in nuclei of uninjured and regenerating muscles; the colors identify different nuclear populations (left) or nuclei in uninjured or regenerating muscle (right). **b**, Expression of *Ttn* and *Myosin* genes identifies myonuclei. **c**, Heat-map of specific genes enriched in clusters 2-8. **d**, Detection of NMJ nuclei. **e**, Volcano plot of NMJ nuclei enriched genes; Exp., expression. **f**, Upper row - double FISH against a known NMJ marker (*Prkar1a1*) and newly identified NMJ genes (green). Bottom row – double single molecule FISH (smFISH; RNAscope) against *Ufsp1* (red) and other newly identified NMJ genes (green). Expression patterns were validated in 2 or more individuals. Scale bar, 10μm.

Figure 2. Two distinct nuclear populations at the myotendinous junction. a, Detection of MTJ-A and MTJ-B clusters. **b**, Enriched genes expressed in MTJ-A and MTJ-B nuclei, and comparison of genes enriched in MTJ-A and -B clusters. **c**, Upper two rows – smFISH of two MTJ-A markers (*Tigd4* and *Col22a1*) in uninjured or 14 d.p.i TA muscle expressing H2B-GFP in myonuclei. Bottom row – Ebf1 (MTJ-B marker) immunohistochemistry in uninjured TA muscle expressing H2B-GFP in myonuclei. Shown are MTJ regions; T, tendon and M, myofiber. Arrowheads indicate co-localization of MTJ marker genes and GFP. Scale bar, 30µm. **d**, Double FISH experiment in isolated single EDL fiber. Insets show magnification of MTJ (a) and NMJ (b) regions. Scale bar, 100µm. Expression patterns were validated in 2 or more individuals.

Figure 3. Identification of novel nuclear subtypes. **a**, Illustrations of potential functions of clusters 5 and 2. Cluster 5 might regulate local mitochondrial metabolism through microRNAs embedded in Dlk1-Dio3 locus, whereas cluster 2 potentially regulates local protein synthesis and entry into the secretory pathway. **b**, Validation of the top markers of clusters 5 and 2 by smFISH. Note their strong expression in a subset of myonuclei. Scale bar, 30μm. **c**, Expression of *Rian* in isolated EDL fibers. Insets show magnifications of indicated areas. Scale bar, 100 μm. **d**, t-SNE map of re-clustering of the major population identified in Figure 1a. Note the emergence of two more distinct sub-populations, 1a and 1b. **e**, Heat-map showing differentially expressed genes in clusters 1a and 1b. **f**, Location of the perimysial sub-population (1a) in the t-SNE map. **g**, Validation of Muc13 in myonuclei adjacent to the perimysium by smFISH. Scale bar, 30μm. Expression patterns were validated in 2 or more individuals.

Figure 4. snRNAseq analysis of *Mdx* muscle. a, t-SNE map of Mdx myonuclei (1,939 nuclei). **b**, Each Mdx myonucleus was assigned a gene signature score, i.e. a gene expression score indicating similarity with uninjured (uninj.), 7 or 14 d.p.i. myonuclei. Each column represents an individual nucleus. **c**, Location of the cluster representing degenerating (necrotic) fibers in the t-SNE map (left). An ncRNA identified in this cluster (*Gm10801*) is expressed in IgG-positive fibers, indicating that it defines nuclei of necrotic fibers. **d**, Nuclear population expressing high levels of various genes implicated in fiber repair that are mutated in myopathies. **e**, Indicated marker genes of the cluster in d are co-expressed in longitudinal sections of muscle of *Mdx* mice. **f**, Co-expression of *FLNC* and *XIRP1* in the muscle from Becker's syndrome patients. Control images for e and f are shown in Extended Data Fig. 8a. All scale bars, 50μm.

Figure 5. Functional compartments inside muscle spindle fibers. a, Cartoon showing the structure of the muscle spindle. **b**, Specific labeling of spindle myonuclei using *Calb1-Ires-Cre.* Arrows indicate muscle spindles. **c,** t-SNE map of muscle spindle myonuclei (260 nuclei). **d,** Expression map of nuclear populations identified in c. **e,** Venn diagram comparing 'spdNMJ vs extrafusal fiber NMJ' and 'spdMTJ vs extrafusal fiber MTJ-B'. Genes enriched in each population (average logFC > 0.7) were used to generate the diagrams. Statistical analysis was performed using hypergeometric test using all genes detected in uninjured/regenerating and spindle datasets as background. **f,** Double smFISH experiments of isolated muscle spindle fibers. Arrows indicate spdNMJ, and asterisks the central non-contractile parts of spindle fibers. All scale bars, 50μm.