1

# Molecular Assays of Pollen Use Consistently Reflect Pollinator Visitation Patterns in a System of Flowering Plants

Running head: Molecular pollen networks reflect visitation

AUBRIE R. M. JAMES[1,2]*†, MONICA A. GEBER[1], and DAVID P. L. TOEWS[1,3]*

[1] Department of Ecology and Evolutionary Biology, Cornell University, Ithaca, New York, USA

[3] Department of Biology, Pennsylvania State University, University Park, PA, USA

† Corresponding author aubrie.james@uq.edu.au

*A.R.M. James and D.P.L. Toews contributed equally to this work

[2] Current Address: School of Biological Sciences, University of Queensland, St. Lucia, Queensland, Australia

**ABSTRACT**

Determining how pollinators visit plants versus how they carry and transfer pollen is an ongoing project in pollination ecology. The differences between how pollinators visit flowers versus how they carry pollen can be appreciable, but the current tools for identifying the pollens that bees carry have different strengths and weaknesses when used for ecological inference. In this study we use three methods to better understand a system of congeneric, co-flowering plants in the genus *Clarkia* and their bee pollinators: observations of plant-pollinator contact in the field, and two different molecular methods to estimate the relative abundance of each *Clarkia* pollen in samples collected from pollinators. We use these methods to investigate if observations of plant-pollinator contact in the field correspond to the pollen bees carry; if individual bees carry *Clarkia* pollens in predictable ways, based on previous knowledge of their foraging behaviors; and how the three approaches differ for understanding plant-pollinator interactions. We find that observations of plant-pollinator contact are generally predictive of the pollens that bees carry while foraging, and network topologies using the three different methods are statistically indistinguishable from each other. Results from molecular pollen analysis also show that while bees can carry multiple species of *Clarkia* at the same time, they often carry one species of pollen. Our work contributes to the growing body of literature aimed at resolving how pollinators use floral resources. We suggest our novel relative amplicon quantification method as another tool in the developing molecular ecology and pollination biology toolbox.

*Keywords:* pollination biology, bee foraging, DNA metabarcoding, relative abundance, species interactions, pollen networks

**INTRODUCTION**

Discovering how plant-pollinator interactions play out in natural systems is critical for understanding how such mutualisms can persist. This is because understanding resource use among mutualists is fundamental to determining the extent to which they rely on each other for population persistence (Roulston and Goodell 2011), support ecosystem functioning (Lucas et al. 2018), and mediate interspecific competition and coexistence (Johnson 2019). Two standard approaches for understanding pollination ecology in natural systems are building networks of species interactions using observations of contact between plants and pollinators, and making inferences about plant-pollinator interactions using observations of pollinator foraging behavior (Godoy et al. 2018, Manincor et al. 2020). Building networks and analyzing their topologies has been a reliable toolkit for illuminating how the community assembly of mutualistic actors operates across space and time (e.g. Alarcón et al. 2008; Bosch et al. 2009; Alarcón 2010; Tur et al. 2016; Godoy et al. 2018; Valdovinos 2019). On the other hand, studying mutualisms through the lens of individual pollinator foraging decisions can illustrate the role of pollinators in plant-plant interactions, which are the building blocks of coexistence theory and community ecology. For example, a well-documented and globally consistent trend in outcrossing flowering plants is incompatible pollen transfer between plant species, a competitive interaction that occurs when pollinators carry multiple species of pollen at once (Mitchell et al. 2009, Carvalheiro et al. 2014, Arceo-Gómez et al. 2019a,b).

Both network structure and pollinator foraging behaviors are understood using observations of pollinator visitation, and each of the following methodological approaches for documenting visitation reveals different information. First, networks of plants and pollinators can be built using observations of contact between pollinators and flowers in the field. Such observations require little work aside from identifying the pollinator and plant species at the time of observation to build a network and develop a basic understanding of how plant-pollinator communities are

organized. Despite widespread application, however, evidence suggests that observations of

contact between pollinators and flowers are potentially poor stand-ins for understanding how

pollinators transfer pollen between plants (Mayfield 2001, Popic et al. 2013, Ballantyne et al.

2015, 2017, Barrios et al. 2016). Because of this, some studies identify and quantify pollen

found on flower stigmas or pollinators using microscopy (Martin and Harvey 2017), which

guarantees that plants and pollinators have interacted in a way that is relevant for both plants

(pollen transfer) and pollinators (pollen resource acquisition). In some cases, however, pollens

are too morphologically similar to distinguish different plant species from one another; in these

situations, a third option is to molecularly interrogate pollen samples using metabarcoding

(Mitchell et al. 2009, Galliot et al. 2017). Metabarcoding identifies pollen species in a sample by

using single-locus PCR amplicons and high-throughput DNA sequencing to compare sample

reads to a database of putative DNA sources. As a tool, metabarcoding has been applied quite

successfully to identify which species of pollen pollinators carry (Wilson et al. 2010, Galimberti

et al. 2014, Sickel et al. 2015, Bell et al. 2016, Bell et al. 2017, Lucas et al. 2018a,b).

Using multiple approaches to study plant-pollinator interactions has the potential to address a

two-headed problem in contemporary pollination ecology: first, the desire to make sense of how

plants and pollinators interact with each other, and second, the desire to compare the

information we can gain using different methodological approaches in this endeavor. Field

observations of plant-pollinator contact are an effective but low-resolution method to reveal

species-scale interactions of plants and pollinators, and identifying pollens on individual

pollinators is a higher-resolution means of understanding how plants and pollinators interact

with each other with regard to pollen use and transfer. Importantly, both kinds of data can be

employed to compare how field observations of contact are similar or different from

observations of pollen found on pollinators; such comparisons are ultimately relevant for

questions of how individuals, populations, and species mutually rely on each other as

resources.

In this study, we use observations of plant-pollinator contact in the field and two different

molecular methods to quantify the relative abundance of pollen sources on bee pollinators

visiting a group of sympatric winter annual plants in the genus *Clarkia* (Onagraceae). This group

of plants – *C. cylindrica* ssp. *clavicarpa* (Jeps.) Lewis & Lewis*, C. speciosa* ssp. *polyantha* Lewis

& Lewis*, C. unguiculata* Lindl.*, C. xantiana* ssp. *xantiana* A. Gray — are sympatric in the

woodland-chaparral areas of the southern foothills of the Sierra Nevada mountain range (from

here, we do not use subspecies epithets). These four species of *Clarkia* rely on a handful of bee

pollinators specialized on the genus *Clarkia* rather than any single species (MacSwain et al.

1973, Moeller 2005). The specialization of these bees on *Clarkia* is either incidental - *Clarkia*

flower much later in the growing season than the vast majority of co-occurring flowering annual

plants, and as such are the only appreciable pollen resource for bees when and where they

occur – or evolutionary: some solitary bees have morphological and/or behavioral adaptations to

accommodate *Clarkia*'s large pollen grains (MacSwain et al. 1973). As in many instances of

closely-related groups of flowering plants, though these *Clarkia* have distinct adult phenotypes,

their pollen grains are morphologically indistinguishable. Furthermore, the *Clarkia* also co-occur

with each other more often than they occur alone in plant communities in their range of

sympatry, and assemblages can contain one to four species of *Clarkia* (Eisen and Geber 2018).

Studies of the four *Clarkia* in their range of sympatry have alternately found signatures of

*Clarkia* reproductive interference (Arceo-Gomez et al. 2016, James 2020), facilitation of

pollination success (Moeller 2004), and character displacement of floral traits (Eisen and Geber

2018) in response to pollinator sharing, thereby implicating pollen transfer as an important

ecological force in the system. However, these do not demonstrate how pollinators actually

carry *Clarkia* pollens. Previous research has shown that the most common pollinator

*Hesperapis regularis* (Melittidae) preferentially visits *C. xantiana* (James 2020). The second

most common pollinator in the system, bees in the *Lasioglossum* genus (Halictidae), have been

shown to either visit all *Clarkia* species at relatively the same rates (Singh 2014), or to

preferentially visit *C. xantiana* and *C. cylindrica* (James 2020). Both *Hesperapis regularis* and

*Lasioglossum spp.* visit *C. cylindrica, C. unguiculata,* and *C. xantiana* regularly despite their

preferences, and are inconstant when foraging in mixed species experimental floral arrays,

sequentially visiting multiple species while foraging (James 2020). Therefore, these pollinators

may transfer incompatible pollen between plants. The third most common bee pollinator in the

system, *Diadasia angusticeps* (Apidae), is more specialized on one *Clarkia* species, *C.*

*speciosa*, and rarely visits the other species of *Clarkia* (Singh 2013, James 2020). The next step

in understanding if these pollinator behaviors translate to trends in *Clarkia* reproductive

interactions is to determine if pollinators collect and carry *Clarkia* pollens in the same ways that

they visit flowers when foraging.

The variation in pollinator behaviors, the wealth of published knowledge of plant-pollinator

interactions and natural history, the limited number of species, and the morphological similarity

of pollens make this particular *Clarkia* system ideal for testing the use of multiple approaches to

understand its pollination ecology. The three different approaches we use to compare *Clarkia*

network topology and individual pollinator foraging decisions are: (1) field observations of plant-

pollinator contacts, (2) relative read abundance (RRA) of *Clarkia* amplicons from PCR

amplification, and, due to established skepticism of RRA in accurately quantifying relative

abundance of species in samples, (3) a novel method we call "quantitative amplicon

sequencing" or "qAMPseq".

Quantitative amplicon sequencing is a method that quantifies the relative abundance of *Clarkia*

pollen in mixed-species samples using the amplification curve of PCR as a backbone for

quantification. This method identifies single nucleotide polymorphisms private to each species,

and then uses PCR to amplify these regions with the goal of post-amplification sequencing, as

in metabarcoding. The PCR amplification is then stopped at four different cycles in order to

estimate the point at which each species' amplification curve crosses a critical threshold (as in

quantitative PCR or qPCR, Figure 1; c.f. Baksay et al. 2020). This point is used to estimate the

relative abundance of each species in each sample. The goal of using qAMPseq analysis of

nuclear DNA markers is to mitigate the risk of two common problems encountered when

quantifying amplicons with metabarcoding: copy number bias and amplification bias. Copy

number bias presents a problem for quantification because it is unclear how the abundance of a

marker is related to overall pollen abundance (e.g. plastid DNA in metabarcoding using cpDNA;

Oldenburg & Bendich, 2004; Golczyk et al., 2014; Bell et al. 2019). If the ratio of pollen grains to

copy number is not one-to-one and/or varies among species, this could skew estimates of

relative abundance in a pollen sample (Richardson et al. 2015, Bell 2019). Thus, our analysis of

nuclear DNA markers means we can more reliably assume a one-to-one ratio. Amplification

bias refers to the fact that the final concentration of amplicon DNA after a full PCR protocol is

not necessarily directly correlated to input DNA concentration (Suzuki and Giovanni 1996; Bell

et al. 2017). Instead, amplicon concentration may be a function of exhausted reaction reagents

rather than the original concentrations of input DNA, especially in samples with lower initial

concentrations (Bell et al. 2017, Bell et al. 2019).

We aim to answer the following questions in this study: (1) how do network topologies of the

*Clarkia* pollination system differ depending on which method we use to understand plant-

pollinator interactions? (2) Do observations of plant-pollinator contact in the field correspond to

the pollens that bees carry (as identified by molecular techniques)? And finally, (3) do individual

bees carry *Clarkia* pollens in predictable ways, based on previous knowledge of their visitation

behaviors? If each of the methods we use presents a different picture of plant-pollinator

interactions, the implication would be that the choice of method is critically important in studying

plant-pollinator systems. If the methods present the same or similar pictures of the system, then

we will have demonstrated that results are robust to different approaches. We predict that

network topologies using the three methods (observation of contact/visitation, RRA, and

qAMPseq) will differ according to method: networks built using observations of contact will have

fewer connections than the networks built using pollen identification and quantification from

RRA or qAMPseq methods, and as a result will appear more specialized than the pollen

networks. In terms of the three most common pollinator species, we hypothesize that if

previously-observed trends in pollinator behavior match what they carry in their pollen samples,

then *Hesperapis regularis* and *Lasioglossum spp.* will carry multiple species of pollen at once.

Moreover, because these particular bee taxa have an established behavioral preference for *C.*

*xantiana*, we expect that they will carry more *C. xantiana* than other *Clarkia* species. We also

predict that *Diadasia angusticeps* will carry only *C. speciosa* pollen.

**MATERIALS AND METHODS**

*Study Species and Field Sampling*

The four species of *Clarkia* in this study are sympatric in the Kern River Canyon in Kern County,

California. We sampled bees in 14 *Clarkia* communities throughout the four species' range of

sympatry from May-June of 2014 (Figure 2). Plant communities varied in *Clarkia* species

richness (one, two, or four species of *Clarkia*; Table 1), and were far enough apart that we did

not expect pollinators to meaningfully or readily move between them (Macswain et al. 1973,

Zurbuchen et al. 2010). In each community, we placed four 20m long transects through patches

of *Clarkia*. Each community was sampled twice on different days: once in the morning (between

8AM and 12PM) and once in the afternoon (between 1PM and 3:30PM). To sample, we walked

along each transect for 20 minutes and caught bees using a sweep net when they landed on

*Clarkia* flowers within one meter of the transect line (that is, the transects were 1/2 meter wide

on either side of the transect line). Nets were turned inside out and quickly wiped after each bee

was captured. Bees were sacrificed using ammonium carbonate. We stored, pinned, and

identified bee samples to species (or in the absence of species-level resolution, to subgenus or

genus) using Michener et al. (1994). We scraped the pollen contents off of all collected bees

and stored each pollen sample in 90% ethanol in centrifuge tubes at -20°C. Most of the

collected samples were not corbiculate bees (such as *Apis mellifera* or *Bombus sp.*), but rather

bees with hairy scopae on their legs or abdomens. *Clarkia* pollen is large (100-150 µm in length)

and connected by viscin threads which allows for pollen grains to be collected and deposited in

sticky pollen masses made up of many pollen grains (MacSwain et al. 1973). Because of this,

the vast majority of pollen samples we harvested were very loosely aggregated pollen masses

that were likely more subject to bee-to-flower transfer than pollen in the tight and neatly-formed

pollen balls of corbiculate bees.

We surveyed *Clarkia* floral abundance to estimate the relative abundance of each species in the

community at the end of each sampling period. To do so, we placed $\frac{1}{2}m^2$ quadrats every four

meters on either side of transect lines, counting all open flowers inside the quadrats. Relative

abundance of each *Clarkia* species was calculated as the number of open flowers of that

species divided by the total number of flowers we counted at the survey time.

In the summers of 2015 and 2016, we also collected pollen from each species of *Clarkia* for use

in testing the RRA and qAMPseq method. To do so, we collected mature anthers from all four

species of *Clarkia* in various communities throughout their range of overlap. We removed pollen

from the anthers and stored them in the same manner that we stored pollen samples taken from

bees.

*Transcriptome sequencing from greenhouse plants*

To identify nuclear genetic markers that could distinguish the four *Clarkia* species, we

performed transcriptome sequencing on tissue obtained from *Clarkia* plants we grew in the

greenhouse (five individuals per species; *n* = 20). We chose to use transcriptome sequencing

because it was a reduced-representation genomic approach that produced long, contiguous

sequences—as compared to ddRAD sequencing—that was necessary for us to develop

subsequent amplicon probes. To grow plants in the greenhouse, we first cold-stratified and

germinated seeds of each species of *Clarkia* in February 2016 in Ithaca, NY. Germinated

seedlings were transferred into D40L conetainers (Stuewe & Sons, Tangent, 208 Oregon, USA)

with a mix of 50% potting soil and 50% perlite. Once in the conetainers, seedlings were bottom-

watered and grown in common conditions in a greenhouse for two months. We harvested

seedlings for RNA extraction when seedlings had more than four true leaves, but before they

had started flowering (March-April 2016).

Leaf tissue of seedlings was harvested and flash frozen in liquid nitrogen. To extract total RNA,

we first mechanically homogenized approximately 100mg of leaf tissue with a nitrogen-chilled

mortar and pestle. We then mixed the homogenized tissue with 1mL of TRIzol and 200uL

chloroform, following the manufacturers guidelines for RNA isolation. We used 200uL of the

isolation to a RNeasy Mini Elute silica column (Qiagen). We added 5 ul of DNAase (NEB) to the

final 45 uL of the final elution, and aliquoted 20 uL of NEBNext Oligo d(t)$_{25}$ beads to isolate

mRNA from the total RNA pool. We then followed the protocol for NEBNext Ultra Directional

RNA Library prep kit (NEB #E7429L). Due to low mRNA yield, we modified the protocol such

that PCR enrichment included 30 cycles. We individually indexed the 20 samples and ran these

on a single lane of Illumina HighSeq, using single end 100 bp sequencing chemistry.

We combined data from the five *C. speciosa* individuals to generate a reference draft

transcriptome assembly using Trinity (Haas et al. 2013). We note here that many of our

sequence reads derived from likely chloroplast DNA (cpDNA). Our goal was to obtain DNA

markers within the nuclear genome as the presence and amount of choloroplasts within each

pollen grain is unknown. We note that future studies might instead sample non-leafy tissue

where possible, which will reduce the number of choloroplast reads. For the present study, we

removed these reads by initially aligning the total read pools from all individuals to the

chloroplast genome of a related species, *Oenothera picensis* (NCBI accession number

KX118607). From the pool of reads that *did not* align to the *O. picensis* cpDNA genome, we

aligned these reads from each individual to the draft *C. speciosa* transcriptome from above. We

then called SNPs using the GATK pipeline, using the same set of presets as in Toews et al.

(2016). We allowed for filtered out SNPs with more than 50% missing data and a minor allele

frequency of less than 5%.

Ideally, our goal was to identify a single genomic region that (1) we could PCR amplify and (2)

included derived, fixed SNPs for each species. To do this, we estimated Weir and Cockerham

per-SNP $F_{ST}$ estimates from VCFTools (Danecek et al. 2011). We generated $F_{ST}$ estimates for

one species compared to the other 15 individuals, and replicated this across all four species.

We then determined which transcriptome contig contained multiple SNPs that had $F_{ST} = 1$ for

each of the four species. We used BLAST to compare our top contig in our assembly to the

nucleotide database at NCBI Genbank. The top hit was a *Clarkia unguiculata* sequence (NCBI

accession number EF017402), and our contig aligned to a region that spans the *5.8S* rRNA

gene, the *internal transcribed spacer 2* gene, and the *26S* rRNA gene. To amplify this region

across additional samples, we used the forward primer sequence

[TCGTCGGCAGCGTC]GTGCCTCGGAGATCATCTGT and reverse primer sequence

[GTCTCGTGGGCTCG]GCCGTGAACCATCGAGTCTTT, with the brackets indicating the

portion of the sequence (P5 and P7, respectively) that would align to our dual-indexed (i5 and

i7) adaptors.

*Quantitative amplicon sequencing—An Overview*

We used two methods to quantify relative input DNA from the four *Clarkia*. First, we used a

common approach to quantify relative abundance of input DNA, which simply uses the relative

read abundance following the full PCR. We refer to the traditional sequencing approach—using

the relative read abundance of amplicons at-or-near the PCR plateau phase—as "RRA-plateau"

or "RRA". We contrast the RRA method with our method that utilizes a PCR cycle treatment,

which we refer to as quantitative amplicon sequencing (qAMPseq).

The premise of qAMPseq applies the theory of quantitative PCR (qPCR, A.K.A. real-time PCR)

with the ability to individually index, multiplex, and sequence hundreds of metabarcoded

samples (Figure 1). Quantitative PCR analysis uses a pre-determined threshold when the PCR

reaction is in an exponential phase of amplification, because the PCR cycle where a reaction

product moves into the exponential phase is directly related to the starting DNA concentration,

unlike the plateau stage (Kubista 2005). Quantitative PCR uses fluorescence (e.g. TaqMan

chemistry) quantified throughout thermocycling to determine the 'cycle number' where the

product fluorescence is higher than a background level, as the product is in the exponential

amplification phase. The estimated number of PCR cycles when the product hits this threshold

is known as threshold cycle (Ct). This Ct value can be compared across samples to compare

starting DNA concentrations.

In qAMPseq, we generate the same PCR amplicon in quadruplicate, with the same starting

conditions, but across different PCR cycling numbers (e.g. 20, 25, 30, and 35 cycles; Figure

2B). Subsequent cleanup and indexing steps preserve the relative DNA amounts in each of

these reactions, which are then individually indexed (i.e. each original sample has four unique

indexes, which correspond to the different cycle 'treatments') and pooled and sequenced with all

other samples (Figure 2C). Samples can then be de-multiplexed (Figure 2D) and, within each

sample and treatment, reads are assigned to predicted taxonomic units ("OTUs"; in this case,

the four *Clarkia* species). The read abundance across each sample and OTU can be used to

calculate Ct (Figure 2E), and a robust value of the relative contribution of input DNA (Figure 2F).

*Pollen DNA extraction and amplicon library preparation*

Pollen DNA was extracted from sample pollen samples (2015) and anther pollen (2015 and

2016) using a CTAB-Chloroform DNA preparation protocol (as in Agrawal et al. 2013), and

stored at -20°C until amplification and quantification. We first quantified the DNA concentration

in each sample using a Qubit fluorometer, and diluted each DNA sample to ~2 ng/uL. We also

created standard dilutions from 1:10 to 1:10000 in triplicate from a single sample of known

origin. We assayed 152 unknown origin pollen sample DNA samples split between two sets.

Each set included pollen DNA from 76 unknown samples, as well as the same 8 DNA samples

of known origin (two from each species), and 12 samples from the standard dilution in triplicate.

Each set of 96 was then transferred to four identical 96-well plates, where we ran a PCR

amplification. We conducted 10uL reaction volumes, including: 6.4 uL of ddH$_2$0, 1 uL of MgCl2,

1 uL of dNTPs, 0.2 uL of each forward and reverse primers (above), 0.1 uL (0.25 units) of

JumpStart Taq (Sigma-Aldrich), and 1 uL of template (at 2 ng/uL). For each set of four plates,

we then used four identical thermal cyclers to run the following protocol simultaneously: 94°C for

3 minutes, and then for plates 1, 2, 3 and 4 we had 20, 25, 30, or 35 cycles of 94°C for 30

seconds, 55°C for 30 seconds, and 72°C for 1 minute, respectively. We then used a final

extension time of 5 minutes.

This resulted in eight 96-well plates—four for each set—representing the different cycle

treatments. We cleaned up each reaction with 1.8X volume SeraPure beads: 10uL of sample

with 18uL of beads, and performed two 70% etOH washes. We eluted in 20 uL of resuspension

buffer (Illumina).

We then ran an individual indexing reaction for each sample within each set (i.e. 384 randomly

chosen, unique indexes for each set). The 20 uL indexing reaction included 4 uL of $ddH_2O$, 10

uL of HiFi Master Mix (KAPA Biosystems), 1 uL of each the forward and reverse i5 or i7

indexes, and 4 uL of template DNA from the amplification step. This was run with the following

thermal cycling conditions: $95^oC$ for 3 minutes, $98^oC$ for 30 seconds, followed by 8 cycles of

$98^oC$ for 30 seconds, $63^oC$ for 30 seconds, and $72^oC$ for 30 seconds. We had a final extension

time of 3 minutes.

Within each sample set, we pooled 5uL of each indexed sample from across the four-cycle

treatments, resulting in one plate for each of the two sample sets. As before, we used a 1.8X

SeraPure bead cleanup for the 20uL pooled samples, and completed two 70% etOH washes.

We eluted samples into 20uL of resuspension buffer. An equal volume of each sample was

pooled—within each set—into the final library. We sequenced each of the two final libraries

separately across two lanes of an Illumina MiSeq, with 2x150 paired end sequencing chemistry.

*Bioinformatics*

Demultiplexing resulted in 1,536 individual fastq files (192 samples across four cycle treatments

with forward and reverse reads). We used zgrep in bash to identify sequence motifs unique to

each of the four species, combining forward and reverse read counts (Supplemental

information).

We generated a standard curve by combining results from across the two sets (Figure S1). As

discussed, the critical number to determine relative abundance in qPCR is the Ct value.

Because qAMPseq data do not directly yield amplicon counts at the end of every cycle, we did

not have direct knowledge of the exact shape of the PCR curve – the important step in

determining relative abundance. To determine the cycle when samples crossed a Ct value

required using a different approach: first, we log-transformed read counts associated with each

of the four-cycle points for which we quantified amplicons. Log-transformation of a PCR curve

theoretically results in a linear relationship of cycle and amplicon number. We took advantage of

this by determining the slope of the amplification line, i.e. $\frac{\Delta \log (\text{amplicon count})}{\Delta \text{ cycle number}}$. We set our Ct

number as log(10,000 reads), and used it in a simple equation to determine the cycle that

corresponded to Ct for each amplification curve of each species in each sample. This is similar

to qPCR, where there is an arbitrary fluorescence value set above the 'background'

fluorescence which is consistently applied across samples. In this case, the Ct number was set

by identifying a region above the read 'noise' and below the maximal values. We henceforth call

this number the cycle count, and combine this information with the standard curve to estimate

relative abundance of each species in the sample.

We also set a second threshold unique to our method. If after 35 cycles a species in a sample

had fewer than 10,000 reads, it was assigned a cycle count value of zero. This is because, from

our known samples, we found non-specific reads can accumulate even if a species' DNA is not

present, which is to be expected in PCR to some extent. The number of non-specific reads

ranged from 64 to 16,227 reads in our known samples after 35 cycles (median read count of

724). By contrast, the on-target read counts after 35 cycles from our known samples ranged

from 19,770 to 744,218 reads (median read count of 135,921). Thus, the intent of this

conservative threshold is to avoid falsely identifying species presence in a sample due to noise

inherent in the method. We calculated cycle numbers and performed all of the following

analyses in R version 3.5.2 (R Core Team, 2018).

*Comparing Network Topologies*

We constructed three networks of plants and pollinators: one network with observations of the

*Clarkia* species bees were caught on, and two using the *Clarkia* pollen that we identified in

bees' pollen samples (one qAMPseq network and one RRA network; we used a 5% cutoff to be

considered 'present' in the sample for the RRA network, addressed in the following section). In

the case of the plant-pollinator contact ("visitation") network, the dataset consists of the number

of times each pollinator was caught on each of the *Clarkia* species. Both pollen use datasets

consist of the estimated proportions of *Clarkia* pollens in each sample, rather than a single

plant-pollinator connection or the presence/absence of *Clarkia* species in a pollen sample. To

build the datasets for the pollen use networks, we multiplied the proportion of each *Clarkia*

species in each pollen sample by 100, and rounded to the nearest whole number.

We used two different metrics to compare the visitation, qAMPseq, and relative read abundance

networks. First, we used a Procrustes analysis to compare network topology for each pair of

networks (a total of three analyses) using tools in the R package vegan (version 2.5.6; Oksanen

et al. 2019). In Procrustes analysis, two matrices are scaled to fit each other for maximum

similarity (minimizing sum-of-squares distances), and then a statistic, $m^2$, is produced to

measure the goodness-of-fit between the two matrices. This statistic is bounded between 0 and

1, where $m^2 = 0$ means the matrices are identical. The significance of network similarity is

calculated using a permutation test (Alarcón et al. 2008). Second, we measured and compared

network-level specialization, H2', in each network. We use H2' because, like Procrustes

analysis, it is robust to differences in the number of interactions (Blüthgen, Menzel & Blüthgen

2006). Values of H2' are between 0 and 1, where higher H2' values indicate that a network

contains more specialized relationships between plants and pollinators, and lower values

indicate the network has more generalized relationships. Network specialization should be the

same in the visitation versus molecular networks if pollinators carry *Clarkia* pollen at the same

rates that they visit *Clarkia*. Networks were built and H2' was calculated using the package

bipartite (version 2.14; Dorrman et al. 2009).

*Individual foraging and pollen loads*

We used both the qAMPseq method as well as the RRA method to determine the extent to

which bees carry multiple species of *Clarkia* pollen at once. In qAMPseq, we determined

presence/absence of *Clarkia* in our samples by asking simply if the cycle values were nonzero

(present) or zero (absent) for each species of *Clarkia*. In contrast, RRA yields relative reads

after full amplification (35 cycles). To determine presence/absence of *Clarkia* with RRA output,

we used three different sample proportion cutoffs to determine the presence of a species in

samples: 0%, where presence was defined as any nonzero read count; and 5% and 10 %

cutoffs, where presence was defined as anything above 5% or 10%, respectively. We used

these three different cutoffs due to the fact that the exact percentage is arbitrary, and we wanted

to account for potential differences in our RRA results due more or less conservative

presence/absence read count thresholds. The 5% threshold was used as the midpoint because

there is some support for using this threshold in the literature (Trevelline et al. 2018).

We compared the number of species in bee pollen samples to the amount of each flowering

*Clarkia* species present when bees were collected. When we sampled bees in *Clarkia*

communities, the communities contained one to four species of flowering *Clarkia*. If bees forage

at random, then we expect their pollen samples to contain the same number of species as the

communities in which they were captured. To test this, we tallied the number of bees caught in

communities with one to four flowering *Clarkia* species, as well as the presence/absence of

*Clarkia* species in each pollen sample (using our four different metrics: relative read abundance

with 0%, 5% and 10% sample proportion cutoffs, (RRA, RRA5 and RRA10), and qAMPseq). We

ran a Pearson's Chi-squared test to determine if the proportion of samples containing one to

four species of *Clarkia* pollen matched the proportion of bees caught in communities with one to

four species of flowering *Clarkia.* If bees are inconstant when pollen foraging, these proportions

would be the same, and the test would return a non-significant result.

Preference for different *Clarkia* species was estimated as the difference between the relative

amount of a species' pollen in a sample and the relative amount of that species' floral

abundance in a surveyed *Clarkia* community where the bee was captured (as in James 2020).

This measure of preference can only be calculated for communities with more than one *Clarkia*

species, because there is not an available 'choice' to make between plants in single-species

communities; as such, we only calculate preference using the samples from communities with

more than one *Clarkia* species. The calculation yields a value between -1 and 1 for each *Clarkia*

species in each pollen sample. Positive values indicate that pollinators carry a *Clarkia* species'

pollen more frequently than it is represented in the community, values of zero indicate that bees

do not preferentially forage for any species, and negative values indicate that pollinators carry a

*Clarkia* species' pollen less frequently than it is represented in the community. We calculated

preference using values generated by RRA with a 5% sample proportion cutoff and qAMPseq.

We used a paired t-test to determine if there was a significant difference in estimates of

preference using qAMPseq versus RRA. We then ran an ANOVA and Tukey's Honest

Significant Difference test to determine if pollinator preference for *Clarkia* species were

significantly different from each other, and t-tests to determine if pollinator preferences were

significantly different from zero*.*

Finally, for each *Clarkia* species, we also summed the number of bees carrying its pollen,

weighted by the proportion the *Clarkia* species was represented each pollen sample. This

weighted value reveals the relative (but not absolute) amount that each *Clarkia* species was

used by each pollinator.

**RESULTS**

In total, we analyzed 192 pollen samples, 40 of which were samples of known pollen contents

from field-collected *Clarkia* anthers, and 152 of which were pollen samples of unknown

composition harvested from bees in 2014. Sequencing resulted in 45,847,334 reads, 94% of

which aligned to one of the four *Clarkia* species reference sequences. All reads in known

samples post-amplification were consistent with the known composition of *Clarkia* in the sample

(Figure S2), barring one sample with a small number of reads. We attempted to analyze the

contents of all 152 pollen samples, but two contained pollen in such low amounts they were

excluded.

The final measurement of RRA with no cutoff returned all four species of *Clarkia* in 100% of the

samples, indicating the unlikely result that all pollinators are not only completely random

foragers, but visited all four *Clarkia* species - even when collected in communities containing

fewer than four *Clarkia* species. Not only is this result unlikely based on the biology of the

system, but it is exceedingly rare that any quantitative analysis with relative read abundance

would use raw read count in the analysis. As such, the rest of our reported results include only

information from RRA abundance reads with 5% (RRA5) and/or 10% (RRA10) cutoffs.

*Pollen use and network comparison*

All three networks were topologically similar (Figure 3) and statistically indistinguishable from

each other (qAMPseq vs. visitation network $m^2$=0.03, p=0.001; RRA5 vs. visitation network

$m^2$=0.03, p=0.001; qAMPseq vs. RRA5 network $m^2$=0.0006, p=0.001). Overall network

specialization, H2', was highest (more specialization) in the *Clarkia* visitation network

(H2'=0.38), lowest (less specialization) in the RRA5 pollen use network (H2'=0.28), and

intermediate in the qAMPseq pollen use network (H2'=0.32). Differences between networks

were most apparent in the less-abundant pollinators: *Apis mellifera* (Apidae)*, Bombus sp.*

(Apidae), and *Megachile sp.* (Megachilidae). Each of these species was only captured on a

subset of *Clarkia*, but carried multiple species of *Clarkia*. The European honeybee, *Apis*

*mellifera*, was only caught on *C. xantiana*, but qAMPseq analysis suggest it carried both *C.*

*cylindrica* and *C. xantiana* pollens (on the other hand, RRA5 only identified *C. xantiana* pollen in

*Apis mellifera* samples). *Bombus sp.* was only caught on *C. unguiculata* but carried both *C.*

*unguiculata* and *C. xantiana* pollens according to qAMPseq, and carried *C. unguiculata, C.*

*xantiana,* and *C. cylindrica* according to RRA5. Finally, *Megachile sp.* was caught on *C.*

*cylindrica* and *C. speciosa*, but carried all four *Clarkia* pollens according to both molecular

methods (Figure 3).

*Pollen sample species composition*

Relative abundance measurements of *Clarkia* in pollen samples were largely the same between

qAMPseq, RRA5, and RRA10, and most similar between the qAMPseq and RRA10

measurement methods (Table 2 and Figure 5; Panel B). Of the 150 pollen samples we analyzed

from bees, 21 out of 150 bees (14%) were caught on *Clarkia* flowers that were different from the

majority of the *Clarkia* pollen found in their pollen samples, indicating some amount of

inconstant foraging (Figure 6). The contents of the pollen samples, however, frequently show

that pollinators carried one species of pollen at a time. Estimates of single-species pollen

samples varied among methods, with 66% (RRA5), 74% (RRA10), or 76% (qAMPseq) of

samples containing only one *Clarkia* species. This indicates a striking level of pollinator fidelity,

emphasized by the fact that 70% of bees we sampled were captured in multi-species *Clarkia*

communities. Furthermore, the Pearson's Chi-squared test comparing the number of bees from

communities with one to four flowering *Clarkia* species versus the number of pollen samples

with one to four flowering *Clarkia* species was significant ($X^2$ (3)=77.05, p<0.001), confirming

that even in diverse *Clarkia* communities, bees often carried only one kind of pollen (Figure 5,

Panel B; Table 2).

A quarter (qAMPseq, 24%; RRA10, 26%) to a third (RRA5, 33%) of the bees carried mixed-

species pollen loads, and mixtures were most often two species of *Clarkia* pollen. The most

common multi-species combination in pollen samples was *C. cylindrica* and *C. unguiculata*

(Figure 6). No pollen samples contained all four species of *Clarkia*. Bees carrying *C. speciosa*

pollen tended to only carry *C. speciosa* pollen: when present, *C. speciosa* was the only species

of pollen in the sample in 12/14 cases. Bees carrying the other three species of *Clarkia* often

carried mixtures of the three (Figure 6).

Pollinator species also carried markedly different proportions of each *Clarkia* pollen. We were

able to distinguish two different *Lasioglossum* taxa in our study, and found that the two

identifiable taxa of *Lasioglossum* exhibited different rates of carrying each species of *Clarkia*. A

putative *Clarkia* specialist, *Lasioglossum pullilabre*, carried all four species but was most

associated with *C. cylindrica*, whereas *L. (Dialictus) sp.,* a likely generalist, carried *C. xantiana*

at higher rates (Figure 4). Furthermore, the *Clarkia* specialist *Hesperapis regularis* carried *C.*

*cylindrica, C. unguiculata*, and *C. xantiana* at almost equivalent rates, while *Diadasia*

*angusticeps* used *C. speciosa* almost to the exclusion of all other *Clarkia* (Figure 4).

*Preference*

The two methods we used to determine pollinator preference, RRA5 and qAMPseq, did not

differ in their estimates of preference (t(607)=0.828, p=0.41). Preferences for *Clarkia* species

were significantly different from each other ($F_{(4, 604)}=4.899$, $p=0.002$). On the whole, bees

carried *C. cylindrica* pollen significantly less than it was represented in *Clarkia* communities

(preference = -0.07±0.0 95% CI). Bees carried the other three species at roughly the same

frequency they occurred in sampled communities (*C. speciosa* $t(604)=0.077$, $p=0.93$; *C.*

*unguiculata* $t(604)=1.086$, $p=0.27$; *C. xantiana* $t(604)=1.59$, $p=0.11$; Figure 5, Panel A).

**DISCUSSION**

In community ecology, mapping the relationships between plants and pollinators is critical to

understanding how pollinators contribute to plant community functioning and vice-versa. Though

observations of pollinator visitation can be used to understand plant-pollinator community

assembly, visitation does not necessarily correspond to pollen transfer or pollen use (Mayfield

2001, Popic et al. 2013, Ballantyne et al. 2015, 2017, Barrios et al. 2016). Metabarcoding has

been critical for understanding if and when bees use certain pollen resources in plant

communities (Galliot et al. 2017, Lucas et al. 2018*b*), but some studies suggest that relative

read abundance from PCR could be unreliable for use in measuring abundance in mixed-

species pollen samples (Bell et al. 2016, Bell et al. 2019).

In this study, we have used visitation observations and molecular methods to uncover how

plants and pollinators interact. We asked if different methods reveal differences in how bee

pollinators visit plants versus how they carry and transport pollen, as well as if pollinators carried

pollen in predictable ways given what is known about their foraging behaviors. We discovered

that the way pollinators visit plants in the field and the pollen they carry in their pollen loads are

almost identical. This result is particularly striking because whereas our visitation network

documents contacts between single pollinators with single flowers, our pollen networks

document potentially tens or hundreds of contacts between a single pollinator and many flowers

in a community. That the visitation and pollen networks correspond so closely not only in

presence/absence but also in frequency of contact indicates high pollinator fidelity. This is

confirmed with further interpretation of the data- our analyses show that pollinators

predominantly carry a single species of pollen at a time. By quantifying relative abundance of

*Clarkia* in pollen loads, we shed light on how pollinator species actually collect pollen from

different *Clarkia* species: when pollinators do carry more than one species of pollen at a time, it

is not at equal frequencies, as presence/absence results alone (as in standard metabarcoding

analysis) might suggest.

*Pollen and Pollinators*

With this study, we have contributed to the growing body of literature that uses molecular

methods to understand pollen collection and transport by bees (Mitchell et al 2009; Galimberti

2014; Bell et al. 2017, Galliot et al 2017, Lucas et al. 2018, Bell et al. 2019), and in the process

provided some evidence for the potential of heterospecific pollen transfer by pollinators in

flowering plant communities (Mitchell et al. 2009, Arceo-Gomez et al. 2019). That said, pollen

samples in our study typically comprised one species of *Clarkia*, which indicates that

heterospecific pollen transfer may not be as common in the *Clarkia* system as in other flowering

plant systems (i.e. Arceo-Gómez et al. 2019). The bees that did carry heterospecific pollen

carried combinations of *C. cylindrica, C. unguiculata* and *C. xantiana*, suggesting that these

three species are more likely to receive heterospecific pollen than their counterpart, *C.*

*speciosa.* To decisively conclude the potential for heterospecific pollen transfer, however, more

research is needed to determine how pollen is transferred between bees and flowers after

pollen has been collected, as pollen contents found on pollinators does not necessarily

correspond to pollen transfer.

There were two important results revealed by our pollen analysis that would not have been

available using presence/absence metabarcoding or visitation observations alone. First, *C.*

*cylindrica* was carried with higher total representation in pollen samples than other species of

*Clarkia*, but it was still carried at lower frequencies than it was represented in *Clarkia*

communities (Figures 4 and 5). This is probably because *C. cylindrica* has the highest average

floral abundance of all the species throughout the communities we sampled (James

*unpublished data*). Consequently, it is possible that pollinators might limit the competitive

dominance of *C. cylindrica* via introducing pollen limitation to seed production. Second, we were

able to resolve differences in foraging between *Lasioglossum* taxa. Because we collected and

sacrificed the bee specimens in this study, we could identify *Lasioglossum* with higher resolution

than observing visitation without sampling, and show that the taxa exhibit differences in their

relationships to *Clarkia*: the putative specialist on the *Clarkia* genus, *Lasioglossum pullilabre*

(Moeller 2004; Eckhart et al. 2006), carries *C. cylindrica* with higher frequency, whereas the

likely generalist, *L. (Dialictus) sp.*, carries *C. xantiana* with higher frequency (Figure 4).

With this study, we were also able to better delineate *Clarkia* use by rare pollinators in our

dataset. The rarest pollinators, *Apis mellifera*, *Bombus sp.*, and *Megachile sp.* (rare to the

dataset, but not rare in the ecosystem; Moeller 2005, Eckhart 2006, and Singh 2013) all carried

more species of *Clarkia* pollen than they had been observed visiting. This explains the

differences in specialization of the pollinator *Clarkia* visitation network versus that of the pollen-

use networks: pollinators carried more diverse pollens (and the network was therefore less

specialized) than the plants we observed pollinators had visited. Given that sampling effort is a

perennial issue in network analyses, pollen networks like this one and others (for example:

Alarcón 2010, Galliot et al. 2017, Lucas et al. 2018) are a means to understand plant-pollinator

relationships when sampling effort is constrained. We say this with caution, however: pollen

analysis data should complement, not supplant, well-designed sampling methods. For example,

it is highly likely that *Apis mellifera* and *Bombus sp.* use all four *Clarkia* pollens (Singh 2013),

but our sample size of a few bees per species makes that impossible to say with certainty.

Other patterns in pollen use were noticeably similar to what we predicted based on standing

knowledge of the pollination ecology of the system. The two *Clarkia* species most often found in

multi-species pollen samples, *C. cylindrica* and *C. unguiculata*, have been shown to occur

together with higher frequency than any other *Clarkia* species pair in this system, overlap in

flowering time more than they do with other *Clarkia*, and exhibit pollinator-mediated character

displacement in floral traits (Eisen and Geber 2018). Given the frequency with which they occur

together in pollen samples, it is possible that character displacement in the floral traits of these

two species could be driven by the competitive effects of heterospecific pollen transfer.

Furthermore, *Diadasia angusticeps* bees carried pollen samples of single-species composition

(*C. speciosa*), which corresponds to previous observations that the species is specialized on *C.*

*speciosa* (Singh 2013, James 2020).

*Molecular methods for quantifying relative abundance in pollen samples*

This study presents a novel method of meta-DNA relative abundance analysis, qAMPseq, with

the motivation of avoiding some of the biases and problems that can arise when using plastid

makers and RRA. However, we unexpectedly found that the two analytical approaches to

quantifying pollen relative abundances in the molecular data yielded nearly identical results. The

only appreciable difference between the RRA network and the qAMPseq network that we

observed is that the network built using qAMPseq exhibited fewer connections overall, and thus

qAMPseq was slightly more conservative than the RRA approach (Figure 3). These slight

differences do not change our community-scale interpretation of the networks. Further, whereas

we expected that the RRA method would yield more spurious associations between plants and

pollinators, we found in almost all analyses that the methods yielded the same results. It is likely

that this similarity was due in part to the fact that all of our samples were diluted to the same

starting concentration of DNA (2 ng/uL). Therefore, in studies wishing to use RRA, it should be

noted that RRA may not yield accurate estimates of relative abundance if DNA concentrations

among samples are highly variable (Bell et al. 2017, Bell et al. 2019).

Furthermore, an important difference between our study and studies that use universal primers

to amplify ITS2 and rcbL is that we had specific target species in our samples. We designed

primers to amplify regions where we knew there were SNPs that differentiated our target

species from each other, rather than relying on variation in ITS2 and rcbL to distinguish *Clarkia*

species from each other. In the end, however, the region that ended up being the most

informative among our four species included ITS2, suggesting it is an ideal starting point for

future studies. It is also important to note that, because we were using closely related species

within the *Clarkia* genus, our application of the metabarcoding approach may not have been

subject to many of quantitative biases identified by Bell et al. (2019). These biases include copy

number variation of the amplified gene, differences in DNA isolation efficacy among samples,

and variation in primer amplification efficiency. If others are to use this method with primers that

target a broader range of possibly more divergent species, these additional biases need to be

carefully considered in experimental design.

Given the dramatic decline in sequencing costs, the sensitivity of current sequencing methods

to detecting SNPs, and the potential biases that may arise when quantifying PCR products from

metabarcoding in mixed-species samples (Bell et al. 2019), the qAMPseq method represents a

lower-cost alternative to florescence-based qPCR (as well as being more viable; see

supplement S3), and offers an alternative to relative read abundance analyses. That said, the

qAMPseq protocol is time-intensive, requires that researchers identify private SNPs for

constituent species in pollen samples, and could be streamlined. One example of streamlining is

that it may be acceptable to perform fewer bead cleaning steps to reduce cost and time at the

bench. Our protocol also necessitated four concurrently running thermocyclers, but we can

envision a modification of thermocycler heating blocks that might allow for variation in the

number of reaction cycles, which would allow for reactions to be run on the same machine (e.g.

Schicke and Hofmann, 2007). There are also standing issues inherent with both molecular

methods that we could not address: for example, our results using qAMPseq versus RRA

depended on the tolerance with which we filtered raw RRA values and the critical threshold

value in qAMPseq, both of which were arbitrarily selected. Furthermore, our analysis does not

incorporate information about the size of the sampled pollen sample, so we cannot draw

conclusions about the amount of pollen that different pollinators transport in their pollen loads.

Irrespective of approach, we highlight that care must be taken in determining the concentration

of pre-amplification samples with any particular primer/target combination, as well the

interpretation of the resulting data.

*Conclusions*

As the theories of plant species interactions continue to incorporate mutualistic, complex

community dynamics (Morales-Castilla et al. 2015, Vázquez et al. 2015, Valdovinos 2019),

molecular approaches can facilitate a more robust understanding of species interactions

between mutualists. Here we have presented evidence that field observations of plant-pollinator

contact correspond to molecular observations of how pollinators carry pollen, indicating a high

level of pollinator fidelity. The networks we built using molecular approaches also show that

pollinators sometimes carry mixed-species pollen balls that might lead to incompatible pollen

transfer between plants. Finally, the two different molecular approaches we used yielded similar

results indicating that both are valid ways of molecularly interrogating pollen samples of mixed

species composition.

**Acknowledgements**

The authors acknowledge Steve Bogdanowicz, the Cornell Evolutionary Genetics Core Facility, the Cornell Lab of Ornithology, Bronwyn Butcher, the Lovette Lab, and the Agrawal Lab for help in method development and for providing lab space and time. Dave Moeller and Indrani Singh helped with bee identification and provided helpful natural history knowledge, and Amy Hastings helped in DNA extraction methods. We thank Julia Brokaw, who was instrumental in the field collections of bees and pollen, as well as Alyssa Anderson, who aided in data preparation. This work was supported by the National Science Foundation Doctoral Dissertation Improvement Grant program grant number 1701675 awarded to ARMJ and MAG, and NSF DEB-1754299 awarded to MAG. Finally, the authors acknowledge that the samples in this research were collected on traditionally Tübatulabal land. The Tübatulabal tribe are currently seeking legal recognition from the federal government.

**REFERENCES**

Agrawal, A. A., Johnson, M. T. J., Hastings, A. P.and Maron, J. L. 2013. A Field Experiment
Demonstrating Plant Life-History Evolution And Its Eco-Evolutionary Feedback To Seed
Predator Populations. American Naturalist 181: S35-S45.

Arceo-Gómez, G., Raguso, R.A., and Geber, M.A. 2016. Can Plants Evolve Tolerance
Mechanisms to Heterospecific Pollen Effects? An Experimental Test of the Adaptive Potential in
Clarkia Species. Oikos 125 (5): 718–25. https://doi.org/10.1111/oik.02594.

Arceo-Gómez, G., Kaczorowski, R. L., Patel, C., and Ashman, T. L. 2019*a*. Interactive Effects
Between Donor And Recipient Species Mediate Fitness Costs Of Heterospecific Pollen Receipt
In A Co-Flowering Community. Oecologia 189: 1041-1047.

Arceo-Gómez, G., Schroeder, A., Albor, C., Ashman, T. L., Knight, T. M., Bennett, J. M., and
Suarez, B. et al. 2019*b*. Global Geographic Patterns Of Heterospecific Pollen Receipt Help
Uncover Potential Ecological And Evolutionary Impacts Across Plant Communities Worldwide.
Scientific Reports 9.

Alarcón, R, Waser, N. M., Ollerton, J. 2008. Year-to-Year Variation in the Topology of a Plant-
Pollinator Interaction Network 117 (12): 1796–1807.

Alarcón, R. 2010. Congruence between Visitation and Pollen-Transport Networks in a California
Plant-Pollinator Community. Oikos 119: 35–44.

Baksay, Sandra, André Pornon, Monique Burrus, Jérôme Mariette, Christophe Andalo, and
Nathalie Escaravage. 2020. Experimental Quantification of Pollen with DNA Metabarcoding
Using ITS1 and TrnL. Scientific Reports 10 (1): 1–9.

Ballantyne, G., K. C. R. Baldock, L. Rendell, and P. G. Willmer. 2017. Pollinator Importance
Networks Illustrate The Crucial Value Of Bees In A Highly Speciose Plant Community. Scientific
Reports.

Ballantyne, G., Baldock K. C. R., and Willmer, P. G. 2015. Constructing More Informative Plant-
Pollinator Networks: Visitation And Pollen Deposition Networks In A Heathland Plant
Community. Proceedings of the Royal Society B: Biological Sciences.

Barrios, B., Pena, S. R., Salas, A. and Koptur, S. 2016. Butterflies Visit More Frequently, But
Bees Are Better Pollinators: The Importance Of Mouthpart Dimensions In Effective Pollen
Removal And Deposition. AoB PLANTS.

Bell, K. L., De Vere, N., Keller, A., Richardson, R.T., Annemarie Gous, Kevin S. Burgess, and
Berry J. Brosi. 2016. Pollen DNA Barcoding: Current Applications and Future Prospects.
*Genome*, 59:629–40.

Bell, K. L., Fowler, J., Burgess, K.S., Dobbs, E.K., Gruenewald, D., Lawley, B., Morozumi, C.
and Brosi, B.J. 2017. Applying Pollen DNA Metabarcoding to the Study of Plant–Pollinator
Interactions. Applications in Plant Sciences 5.

745    Bell, K. L., Burgess, K.S., Botsch, J.C., Dobbs, E.K, Read, T.D., and Brosi, B.J. 2019.
746    Quantitative and Qualitative Assessment of Pollen DNA Metabarcoding Using Constructed
747    Species Mixtures. Molecular Ecology 28: 431–55.

748    Blüthgen, N., Menzel, F., & Blüthgen, N. 2006. Measuring Specialization In Species Interaction
749    Networks. BMC Ecology, 6:9-12.
750
751    Cane, J. H., and Sipes, S.S. 2006. Characterizing Floral Specialization By Bees: Analytical
752    Methods And A Revised Lexicon For Oligolecty. Plant–pollinator interactions: From
753    specialization to generalization.
754
755    Carvalheiro, L. G., Biesmeijer, J. C., Benadi, G., Fründ, J., Stang, M., Bartomeus, I., Kaiser-
756    Bunbury, C. N., et al. 2014. The Potential For Indirect Effects Between Co-Flowering Plants Via
757    Shared Pollinators Depends On Resource Abundance, Accessibility And Relatedness. Ecology
758    Letters 17:1389–1399.
759
760    Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker, R.E.,
761    Lunter, G., Marth, G.T., Sherry, S.T. and McVean, G., 2011. The Variant Call Format And
762    Vcftools. Bioinformatics, 27: 2156-2158.
763
Danforth, B. N., Minckley, R. L., and Neff, J. L. 2019. The Solitary Bees. Princeton University
Press, Princeton, New Jersey.

Deiner, K., Bik, H.M., Mächler, E., Seymour, M., Lacoursière‐Roussel, A., Altermatt, F., Creer,
S., et al. 2017. Environmental DNA Metabarcoding: Transforming How We Survey Animal and
Plant Communities. Molecular Ecology 26 (21): 5872–95.

Eckhart, V.M., Rushing, N.S., Hart, G.M., and Hansen, J.D. 2006. Frequency-Dependent
Pollinator Foraging in Polymorphic Clarkia xantiana ssp. xantiana Populations: Implications for
Flower Colour Evolution and Pollinator Interactions. Oikos 112: 412–21.

Eisen, K. E., and Geber, M.A. 2018. Ecological Sorting And Character Displacement Contribute
To The Structure Of Communities Of Clarkia Species. Journal of Evolutionary Biology.
774
Manincor, Natasha de, Nina Hautekèete, Clément Mazoyer, Paul Moreau, Yves Piquot,
Bertrand Schatz, Eric Schmitt, Marie Zélazny, and François Massol. 2020. How Biased Is Our
Perception of Plant-Pollinator Networks? A Comparison of Visit- and Pollen-Based
Representations of the Same Networks. Acta Oecologica 105: 103551.

Mitchell, R. J., Irwin, R. E., Flanagan, R. J., and Karron, J. D. 2009. Ecology and Evolution of
Plant-Pollinator Interactions. M Annals of Botany 103:1355-1363.

Moeller, D. A. 2004. Facilitative Interactions Among Plants via Shared Pollinators. Ecology 85
(12): 3289–330.

Galliot, J. N., Brunel, D., Bérard, A., Chauveau, A., Blanchetête, A., Lanore, L., and Farruggia,
A. 2017. Investigating A Flower-Insect Forager Network In A Mountain Grassland Community
Using Pollen DNA Barcoding. Journal of Insect Conservation 21:827–837.

Galimberti, A., De Mattia, F. Bruni, I., Scaccabarozzi, D., Sandionigi, A., Barbuto, M., Casiraghi,
M., and Labra, M. 2014. A DNA Barcoding Approach to Characterize Pollen Collected by
Honeybees. PLoS ONE.

Golczyk, H., Greiner, S., Wanner, G., Weihe, A., Bock, R., Börner, T., & Herrmann, R. G.
(2014). Chloroplast DNA In Mature And Senescing Leaves: A Reappraisal. The Plant cell, 26:
847–854.

Haas B.J., Papanicolaou A., Yassour M., Grabherr M., Blood P.D., Bowden J., Couger M.B.,
Eccles D., Li B., Lieber M., Macmanes M.D., Ott M., Orvis J., Pochet N., Strozzi F., Weeks N.,
Westerman R., William T., Dewey C.N., Henschel R., Leduc R.D., Friedman N., Regev A. De
novo Transcript Sequence Reconstruction From RNA-seq Using The Trinity Platform For
Reference Generation And Analysis. 2013. Nature Protocols. 8:1494-512.
797
798  James, A.R.M. 2020. Pollinator Sharing and Spatial Partitioning Affect Flowering Plant
799  Coexistence. Cornell University.
800
James, A., Geber, M.A., Toews, D.P.L.  2021. Molecular Assays of Pollen Use Consistently
Reflect Pollinator Visitation Patterns in a System of Flowering Plants, Dryad,
Dataset, https://doi.org/10.5061/dryad.0p2ngf1x0

Johnson, C. A., and Bronstein, J.L. 2019. Coexistence And Competitive Exclusion In Mutualism.
Ecology 100:e02708.

Kubista, M., Andrade, J.M., Bengtsson, M., Forootan, A., Jonák, J., Lind, K., Sindelka, R.,
Sjöback, R., Sjögreen, B., Strömbom, L., Ståhlberg, A., Zoric, N. 20016. The Real-Time
Polymerase Chain Reaction. Molecular Aspects of Medicine 27: 95-125.

Lucas, A., Bodger, O., Brosi, B. J., Ford, C. R., Forman, D. W., Greig, C., Hegarty, M., et al.
2018. Generalisation and Specialisation In Hoverfly (Syrphidae) Grassland Pollen Transport
Networks Revealed By DNA Metabarcoding. Journal of Animal Ecology 87:1008-1021.
812
813  MacSwain, J. W., Raven, P. H., and Thorp, R. W. 1973. Comparative Behavior Of Bees And
814  Onagraceae. IV. Clarkia bees of the Western United States. University of California Publications
815  in Entomology 70:1–80.
816
817  Martin, A. C., and Harvey, W. J. 2017. The Global Pollen Project: A New Tool For Pollen
818  Identification And The Dissemination Of Physical Reference Collections. Methods in Ecology
819  and Evolution 8:892–897.
820
821  Mayfield, M. M., Waser, N. M., and Price, M. V. 2001. Exploring The "Most Effective Pollinator
822  Principle" With Complex Flowers: Bumblebees And Ipomopsis Aggregata. Annals of Botany 88:
823  591-596.
824
Mitchell, R. J., Irwin, R. E., Flanagan, R. J., and Karron, J. D. 2009. Ecology And Evolution Of
Plant-Pollinator Interactions. Annals of Botany 103: 1355-1363.

Michener, C. D., McGinley, R. J., and Danforth, B. N. 1994. The Bee Genera Of North And
Central America. Edited by Rosemary Sheffield. Washington: Smithsonian Institution Press.

Moeller, D. A. 2005. Pollinator Community Structure And Sources Of Spatial Variation In Plant-
Pollinator Interactions In Clarkia xantiana ssp. xantiana. Oecologia 142: 28-37.
831
832     Morales-Castilla, I., Matias, M. G., Gravel, D. and Araújo, M. B. 2015. Inferring Biotic
833     Interactions From Proxies. Trends in Ecology and Evolution 30: 347–356.
834
835     Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R.,
836     O'Hara, R. B., Simpson, G.L., Solymos, P., Henry, M., Stevens, H., Szoecs, E. and Wagner, H.
837     2019. vegan: Community Ecology Package. R package version 2.5-6. https://CRAN.R-
838     project.org/package=vegan
839
840     Oldenburg, D. J., & Bendich, A. J. 2004. Changes In The Structure Of DNA Molecules And The
841     Amount Of DNA Per Plastid During Chloroplast Development In Maize. Journal of Molecular
842     Biology, 344: 1311– 1330.
843
Popic, T. J., Wardle, G. M., and Davila, Y. C. 2013. Flower-Visitor Networks Only Partially
Predict The Function Of Pollen Transport By Bees. Austral Ecology 38: 76-86.

Richardson, D. M., Allsopp, N., D'antonio, C. M., Milton, S. J., and Rejmánek, M. 2000. Plant
Invasions — the Role of Mutualisms. Biological Reviews 75: 65–93.

Roslin, T., Traugott, M., Jonsson, M., Stone, G. N., Creer, S., and Symondson, W.O.C. 2019.
Introduction: Special Issue on Species Interactions, Ecological Networks and Community
Dynamics – Untangling the Entangled Bank Using Molecular Techniques. *Molecular Ecology*
28: 157–64.

Roulston, T. H., and Goodell, K. 2011. The Role of Resources and Risks in Regulating Wild Bee
Populations. Annual Review of Entomology 56 (1): 293–312. https://doi.org/10.1146/annurev-
ento-120709-144802.

Sickel, W., Ankenbrand, M. J., Grimmer, G., Holzschuh, A., Härtel, S., Lanzen, J., Keller, A. et
al. 2015. Increased Efficiency In Identifying Mixed Pollen Samples By Meta-Barcoding With A
Dual-Indexing Approach. BMC Ecology 15: 20.
858
859     Singh, I. 2013. Master's Thesis. Cornell University.
860
Schicke, K. and Hofmann, C. 2007. Thermocycler With A Temperature Control Block Driven In
Cycle. US Patent number US 2007/0212774 A1.

Sousa, L.L., Silva, S. M., and Xavier, R. 2019. DNA Metabarcoding in Diet Studies: Unveiling
Ecological Aspects in Aquatic and Terrestrial Ecosystems. Environmental DNA 1: 199–214.

Suzuki, Marcelino T., and Stephen J. Giovannoni. 1996. Bias Caused by Template Annealing in
the Amplification of Mixtures of 16S RRNA Genes by PCR. Applied and Environmental
Microbiology 62: 625–30.

Tur, C., Sáez, A. Traveset, A. and Aizen, M.A. 2016. Evaluating the Effects of Pollinator-
Mediated Interactions Using Pollen Transfer Networks: Evidence of Widespread Facilitation in
South Andean Plant Communities. Ecology Letters. https://doi.org/10.1111/ele.12594.

Trevelline, B. K., Nuttle, T., Hoenig, B.D., Brouwer, N.L., Porter, B.A., and Latta, S.C. 2018.
DNA Metabarcoding of Nestling Feces Reveals Provisioning of Aquatic Prey and Resource
Partitioning among Neotropical Migratory Songbirds in a Riparian Habitat. Oecologia 187 (1):
85–98. https://doi.org/10.1007/s00442-018-4136-0.

Valdovinos, Fernanda S. 2019. Mutualistic Networks: Moving Closer to a Predictive Theory.
Ecology Letters 22 (9): 1517–34. https://doi.org/10.1111/ele.13279.

Vázquez, D. P., Ramos-Jiliberto, R., Urbani, P., and Valdovinos, F. S. 2015. A conceptual
framework for studying the strength of plant -- animal mutualistic interactions. Ecology Letters
18: 385-400.

Wilson, Erin E., C. Sheena Sidhu, Katherine E. Levan, and David A. Holway. 2010. Pollen
Foraging Behaviour of Solitary Hawaiian Bees Revealed through Molecular Pollen Analysis.
Molecular Ecology 19 (21): 4823–29.

Zurbuchen, Antonia, Lisa Landert, Jeannine Klaiber, Andreas Müller, Silke Hein, and Silvia
Dorn. 2010. Maximum Foraging Ranges in Solitary Bees: Only Few Individuals Have the
Capability to Cover Long Foraging Distances. Biological Conservation 143: 669–76.

886
## 887 Data Accessibility
888 Pollen sample composition data will be archived in the Dryad digital repository.

## 890 Author Contributions

ARMJ, MAG, and DPLT designed the study. ARMJ and DPLT performed research, analyzed

data, and wrote the paper. All authors contributed to editing and revising the manuscript.

Figure 1. Quantitative amplicon sequencing schematic. **A** A mixed-composition sample. Each circle represents a pollen grain and
species identity is indicated by color. The numbers of each pollen grain in the mixed sample are shown below (i.e. mixed sample #1
had 11 grains from red, 7 grains from yellow, etc.). **B** This mixed sample is amplified via PCR across four different thermal cyclers,
each with different cycle number "treatments". **C** Each of these samples are then uniquely indexed to keep track of sample identify
and PCR cycle treatment. All samples are pooled, post-PCR, and run on a single Illumina MiSeq lane, where they are sequenced. **D**
After sequencing, samples are de-multiplexed. **E** Read abundance for each species in each sample is used to calculate the Ct value
from a simplified read abundance 'curve'. **F** These Ct values are used to calculate the proportion of each species in each sample (i.e.
mixed sample #1 had more red pollen grains, a lower Ct, and subsequently a higher relative proportion in the sample).

902
903
904 Figure 2. Plant and pollinator sampling locations in the Kern River Canyon in Kern County, California, USA. Pie charts show the
905 *Clarkia* species community composition in the first round of sampling. Colors: red, *C. xantiana*; blue, *C. cylindrica*; orange, *C.*
906 *unguiculata*; and yellow, *C. speciosa* (see Table 1 for details).
907

908
Figure 3. Bipartite networks of plants and pollinators. The network on the left shows the relationships of pollinator taxa (bottom row)
and the plant species they were captured on (top row; C, *Clarkia cylindrica;* S, *C. speciosa;* U, *C. unguiculata;* X, *C. xantiana*). The
network in the middle and on the right show the relationships of pollinator taxa and the plant species in their pollen loads using the
qAMPseq method and relative read abundance (RRA5) method. The size of the bars in the top and bottom rows are proportional to
the number of bees/plants in the data, and the size of the blue connecting bars are proportional to the frequency with which a
plant/pollinator combination occurred.
915

917 Figure 4. Relative abundances of *Clarkia* pollen on pollinators. *Clarkia* present in a pollen sample were weighted by their relative
918 abundance in the sample and then summed to yield a value for that species' relative pollen contents for each pollinator. Larger bars
919 indicate higher pollinator counts, where *Lasioglossum pullilabre* was the most frequently caught pollinator, followed by *Hesperapis*
920 *regularis, Lasioglossum (Dialictus) sp.,* and *Diadasia angusticeps.*
921

922
Figure 5. **A** Preference for different *Clarkia* pollen based on samples taken from foraging bees. Preference was calculated as the
difference between frequency of a *Clarkia* species blooming in the community and frequency it was represented in bees' pollen
samples. Mean preference is plotted with a 95% confidence interval, and was estimated using quantitative amplicon sequencing.
Estimates of preference did not significantly differ between methods (not shown). **B** Bar plots showing the number of *Clarkia* species
in pollen samples. The far-left bar shows what we would expect if bees are random foragers: for example, the 34 bees caught in
communities with four co-flowering *Clarkia* species (black section of the far-left bar) should have four *Clarkia* species in their pollen
samples. We estimated the number of *Clarkia* species found on bees in four ways: quantitative amplicon sequencing (qAMPseq),
and relative read abundance (RRA) at different sample proportion cutoffs. Bees carried fewer species in their pollen samples than
were flowering in their corresponding *Clarkia* communities. The far-right bar is relative read abundance with no sample proportion
cutoff; with this method, all samples contained all four species.

Figure 6. Proportion plot for each sample of pollen harvested by bees, represented by bars. Small circles above each bar indicate the species of *Clarkia* the bee was caught on, while larger pie charts indicate the *Clarkia* species composition of the community each bee came from. White bars indicate that the amount of pollen found on the bee was too low for compositional analysis.

938    Table 1. Sites sampled in the study and the sampling schedule.

| Species composition | Site name | Dates sampled (2014) | Degree Decimal Latitude / Longitude |
|---|---|---|---|
| *C. cylindrica* *C. unguiculata* | Little Tree (LT) | 15 May 19 May | 35.527197 -118.66324 |
| | Summer Camp (SC) | 17 May 21 May | 35.531156 -118.64774 |
| | Site 31.5 (S31.5) | 24 May | 35.566167 -118.56695 |
| *C. speciosa* *C. xantiana* | Green Rock East (GRE) | 23 May 29 May | 35.597436 -118.51016 |
| | Site 8 (S8) | 24 May 30 May | 35.593325 -118.51219 |
| | Black Gulch (BG) | 29 May 30 May | 35.592425 -118.52681 |
| *C. cylindrica* *C. unguiculata* *C. speciosa* *C. xantiana* | Lower China Gardens (LCG) | 16 May 26 May | 35.538564 -118.64998 |
| | Kingsnake (KS) | 16 May 26 May | 35.529239 -118.6601 |
| | Mill Creek (MC) | 17 May 23 May | 35.537633 -118.61411 |
| *C. xantiana* | Borel (BR) | 28 May 30 May | 35.590131 -118.5188 |
| | Upper China Gardens (CG) | 18 May 31 May | 35.578939 -118.52383 |
| *C. cylindrica* | Coyote Gulch (UCG) | 24 May 25 May | 35.536933 -118.64963 |
| *C. speciosa* | Tip Top (TT) | 29 May 31 May | 35.596622 -118.50561 |

941 Table 2. The number of pollen samples collected in communities, and the number of flowering
942 species in the pollen samples estimated with different methods – quantitative amplicon
943 sequencing (qAMPseq) and relative read abundance (RRA) at different sample proportion
944 cutoffs.

| | Number of flowering species in community | | | 946 |
|---|---|---|---|---|
| | **1** | **2** | **3** | **4** |
| **Number of pollen samples collected** | 44 | 63 | 9 | 34 947 |
| **qAMPseq** | 114 | 31 | 5 | 0 948 |
| **RRA** | 0 | 0 | 0 | 150 |
| **RRA, 5%** | 100 | 42 | 8 | 0 |
| **RRA, 10%** | 111 | 35 | 4 | 0 |