2

Habitat Selection by the Ringed Kingfisher (*Megaceryle torquata stictipennis*) in Basse-Terre, Guadeloupe: Possible Negative Association with Chlordecone Pollution

8   Pascal Villard[1], Alain Ferchal[2], Philippe Feldmann[1], Claudie Pavis[1] and

Christophe Bonenfant[3]

12   [1]AEVA (the society for the study and protection of vertebrates and plants in the Lesser Antilles),

Hauteurs Lézarde, 97170 Petit-Bourg, Guadeloupe; P.V. e-mail: pascalvillard@yahoo.fr, P.F. e-mail:

14   feldmann@cirad.fr, C.P. e-mail: claudie.pavis@gmail.com; [2]Parc National de la Guadeloupe,

Montéran 97120 St Claude, Guadeloupe; e-mail: alain.ferchal@guadeloupe-parcnational.fr; [3]UMR

16   CNRS 5558, Laboratoire Biométrie et Biologie Évolutive, Université Claude Bernard - Lyon 1, 43

Boulevard du 11 Novembre 1918, 69622 Villeurbanne Cedex, France; e-mail:

christophe.bonenfant@univ-lyon1.fr

Header: The Ringed Kingfisher in Guadeloupe

Abstract: In the Lesser Antilles the Ringed Kingfisher *Megaceryle torquata stictipennis* is found on Martinique, Dominica and Guadeloupe. On Martinique and Guadeloupe, farmers

growing bananas have made massive use of chlordecone from 1973 until 1993. Chlordecone is a remnant organochlorine insecticide which bioaccumulates in organisms easily. We carried

out a survey of the Ringed Kingfisher in 2009 on Basse-Terre, Guadeloupe, traveling 270 km on 44 rivers to assess the effects of chlordecone on its habitat selection behaviour. We

recorded 47 encounters of Ringed Kingfishers over the survey. A habitat selection analysis revealed that the Ringed Kingfishers were located mainly in sections of rivers flowing

through the ombrophilous forest, but absent on the ocean shoreline or in the lower parts of rivers. The Ringed Kingfisher's distribution on Guadeloupe could be negatively associated

with areas heavily polluted by the chlordecone. We propose that the widespread use of chlordecone in banana plantations on Guadeloupe between 1973 and 1993 has adverse

ecological consequences, and may be responsible for the absence of Ringed Kingfishers in lowland habitat on the island.

Keywords: Antilles, bioindicator, habitat selection, Kepone, soil pollution

40   *Résumé*: Dans les Petites Antilles, le Martin-pêcheur à ventre roux *Megaceryle torquata*

stictipennis ne vit que sur trois îles: Martinique, Dominique et Guadeloupe. En 2009, nous

42   avons effectué un recensement en Guadeloupe, sur 270 km parcourus dans 44 rivières. Douze

territoires de Martin-pêcheur à ventre roux ont été trouvés, et la taille totale de la population

44   est estimée à 54-64 individus. Une analyse de la sélection de l'habitat montre que le Martin-

pêcheur à ventre roux se trouvait principalement sur les rivières dans la forêt ombrophile et

46   non pas sur le bord de l'océan ni dans la partie basse des rivières. En Guadeloupe, le Martin -

pêcheur à ventre roux est négativement associé avec les surfaces hautement polluées par un

48   insecticide organochloré, la chlordécone. Nous suggérons que l'épandage massif de la

chlordécone de 1973 à 1993, dans les bananeraies de Guadeloupe, doit être grandement

50   responsable de l'absence du Martin-pêcheur à ventre roux dans les habitats de basses altitudes

de l'île.

Mots-clés: bioindicateur, chlordécone, Martin-pêcheur à ventre roux, pollution du sol,

54   sélection d'habitat

Resumen: En las Antillas Menores, el Martín Gijante Neotropical *Megaceryle torquata*

stictipennis solo vive en tres islas: Martinica, Santo Domingo y Guadalupe. En 2001,

realizamos un censo en Guadalupe sobre 270 km recorridos en 44 ríos. Doce territorios de

Martín Gijante Neotropical han sido registrados, y el volumen total de la población se estima

a 54-64 individuales. Un análisis de la selección del hábitat muestra que el Martín Gijante

Neotropical se encontraba principalmente en los ríos en el bosque lluvioso y no a la orilla del

mar tampoco en la parte baja de los ríos. En Guadalupe, el Martín Gijante Neotropical esta

negátivamente asociado con las superficies muy poluadas por un insecticida organicloro, la

chlordecona. Sugeremos que la difusión masiva de la chlordecona entre 1973 y 1993, en las

plantaciones de plátanos de Guadalupe, debe de ser altamente responsable de la aucencia del

Martín Gijante Neotropical en los hábitats de baja altitud en la isla.

Palabras claves: bioindicador, polución del suelo, selección de hábitat;

The Ringed Kingfisher *Megaceryle torquata* is a quite common and widely distributed species

72   across Central and South America (Woodall 2001). In the Antilles, however, the presence of

the species is heterogeneous among islands. For instance, the Ringed Kingfisher does not

74   occur in the Greater Antilles, and in the Lesser Antilles *M. t. stictipennis* is found on

Dominica, Martinique and Guadeloupe (Remsen 1990), suggesting that favorable ecological

76   conditions are met on a few islands only, or that anthropogenic activities could be limiting the

Ringed Kingfisher distribution. Of particular interest for the distribution of many bird species

78   are the agricultural practices known to affect many aspects of their biology and ecology (see

Geiger *et al.* 2010 for a review). Banana production on Guadeloupe, an important and

80   intensive agricultural activity of many people living on tropical islands, have among other

things led to marked modifications of the landscape like the widespread deforestation.

82   Tightly associated to banana farming is chlordecone, an organochlorine insecticide, that

has been used heavily and extensively by farmers to control regular damage by a root borer,

84   the banana weevil *Cosmopolites sordidus,* from 1973 until its ban on Guadeloupe in

September 1993 by French law (Cabidoche and Lesueur 2011). The widespread use of

86   chlordecone resulted in extended pollution of soils, waters and riverbed sediments (Crabit *et*

al. 2016), which ecological consequences remains poorly documented and studied on

88   Guadeloupe. Since spreading of chemicals in the landscape for agriculture can potentially

alter the distribution of birds (Douthwaite 1982; Parsons *et al.* 2010; Mineau and Whiteside

90   2013), one factor that may have influenced Ringed Kingfisher populations in the Guadeloupe

could be the historical use of chlordecone.

92   Chlordecone is a long lasting organic pollutant with carcinogenic, mutagen and/or

reprotoxic consequences for exposed organisms. On wildlife, negative effects of chlordecone

94   were reported with strong effects on the reproductive biology of many species (see

Eroschenko 1981 for a review). In birds, bioaccumulation of chlordecone manifested itself in the eggs of osprey *Pandion haliaetus* living along a chlordecone-polluted river in the USA (Wiemeyer *et al*. 1988). Besides, organochlorides can damage the nervous system altering the breeding behavior of seabirds (Schreiber and Burger 2002). Organochlorides such as chlordecone can affect negatively bird biology to a large extent, ranging from organism functioning, to reproductive success, scaling up to ecological processes such population abundance and viability (Saaristo *et al*. 2018). For instance the reproductive failure induce by DDT because of fragile egg shells was accounted for the decreased in population abundance of the brown pelican *Pelicanus occidentalis* California, USA (Risebrough *et al*. 1971).

Lying at the top of the aquatic food chain, bioaccumulation of chlordecone in body tissues should be particularly acute for Ringed Kingfishers (Clarkson 1995). Two types of ecological consequences of pesticides are usually recognized in the literature (Saaristo *et al*. 2018). The direct effects act on the physiology of exposed organisms to pollutant in turn altering the metabolism, the cognitive ability or the behavior of individuals, and ultimately leads to reduced survival and reproduction. Indirect effects are generated by cascading effects of pollution on the abundance of preys or by changes in the habitat quality, which indirectly impacts on reproduction and survival of individuals of higher levels in the food chain (Campbell *et al*. 1997). Indirect effects are expected to modify the behavior of individuals, such as lowered predator avoidance or predation efficiency, or decreased movement ability, which all contributes to magnify the direct effects of contaminant on populations (Saaristo *et al*. 2018). Given reported adverse effects on the life histories of birds of many organochlorine compounds, especially for fish-eating birds in general (Moore 1965; USEPA 1975; Schäfer *et al*. 2011), the historical use of the chlordecone could explain the distribution and habitat selection of Ringed Kingfisher on Guadeloupe. Accordingly, many kingfisher species are used

as biomonitors of waterways polluted by various contaminants including mercury (White and

Cristol 2014, Zamani *et al*. 2009) and organochlorides such as DDT (Tanabe *et al*. 1998,

Evans and Bouwman 2000).

Here we investigated whether chlordecone pollution could be related to Ringed

Kingfisher habitat selection. We conducted presence-absence surveys for this species on 44

rivers located on Basse-Terre island, Guadeloupe, and calculated selection ratios to determine

whether Ringed Kingfishers avoid polluted areas. If chlordecone negatively affects the

abundance of Ringed Kingfishers, they should avoid areas with chlordecone contamination,

i.e. we should find proportionally less individuals in the polluted areas than its availability in

the landscape (Manly *et al.* 2007). To lend further support to that the potential negative

association between chlordecone pollution and Ringed Kingfisher habitat selection, we

accounted for those ecological variables that we believe would capture the main spatial

structure of Basse-Terre in terms of available habitats and river characteristics, and could

potentially confound the effect of chlordecone..

Methods

We carried out our study in Guadeloupe, an archipelago of the Lesser Antilles in the

Caribbean sea. On Basse-Terre island (848 km^2, 16°10' N 61°40' W), a mountain range runs

from north to south mostly between 500-800 m above sea level. The highest point is the 1,467

m summit of the southern volcano La Soufrière. Because the mountain range blocks the

prevailing easterly winds from the Atlantic ocean, the east coast of Basse-Terre is wetter with

a drier leeward west coast. Two main forest types have been described along the elevation

gradient of the volcano (Van Laere *et al.* 2016). At low elevation lies the lower rainforest

dominated by *Amanoa caribaea, Tapura latifolia* and *Dacryodes excelsa* trees. At higher

elevation, the rainforest is mostly dominated by *Richeria grandis.* Many rivers run from the

mountain range to the ocean, especially along the wetter east coast (Fig. 1b). We sampled 21

rivers on the windward east coast and 23 rivers on the leeward west coast; with 14 rivers (< 5

km), 15 (5 <10 km), 13 (10 < 25 km) and the 2 longest rivers (> 25 km). For the shorter 42

rivers, looking on a map of Basse-Terre, we selected the rivers to be equally spread around the

island and with different levels of chlordecone pollution (Fig. 1a).

We conducted fieldwork in search of Ringed Kingfishers from 20 March to 20 August

2009. To obtain a baseline for the bird habitat use, we first studied the behavior of one pair,

observing the birds for 33.5 hours over four days (on March 20th-23rd-24th and 26th 2009). We

then surveyed Ringed Kingfisher using two methods: from kayak where possible in the lower

sections of rivers or by walking along rivers. We assessed the Ringed Kingfisher presence on

a river section by moving upstream slowly mostly in the river bed at a speed of 0.8 km per

hour (1 km/h with a kayak), and starting at sunrise. We sampled the rivers only once and

usually walked down away from rivers back to the coast afterwards. During the survey, we

searched for the birds in the vegetation of riverbeds (with binoculars *Swarovski* 10x25) to spot

any perched kingfisher. We recorded the GPS location of any Ringed Kingfisher heard or

seen. Our experience showed that when a Ringed Kingfisher was present, by being careful

enough (watching and listening) during the survey, we were quite confident not to miss the

bird. Because some birds flew ahead of us as we surveyed rivers, we sometimes made

repeated observations of a single individual. We always recorded the locations where Ringed

Kingfishers were initially flushed. If the bird flew upstream, we also recorded the subsequent

locations where it perched. We visually kept track of the flying bird until it reached its

territory limit and turned back. We used this behavior to differentiate Ringed kingfishers

individually, and any new bird later flushed upstream was considered as a different one. We

acknowledge here that birds may have different habitat use between early spring and summer

168   that could affect our results, even if it could be easier to see adult birds flying when feeding

nestlings. Besides, because birdwatchers have been recording new, opportunistic Ringed

170   Kingfishers observations between 2013 and December 2017 since our study was conducted

(Karunati data base, accessed on May 10[th] 2019). We retrieved the observation location

172   reported by the observers from the Internet to reinforce our own observations and increase our

sample size for the statistical analyses. We hence assumed observers spotted different Ringed

174   Kingfishers, a likely met assumption given the rather large distance between the different

observations we retained (mean: 7.1 km; range: 2.2 and 12.2 km). During the survey, we also

176   made some anecdotal observations on adults Ringed Kingfisher bringing preys to their chicks

and identified what species they feed on.

178   We described the local environmental conditions faced by Ringed Kingfisher along the

sampled rivers and its surroundings with four ecological variables. With a GIS, we first

180   created 250 m long river segments, starting from the river mouth toward its spring. For each

segments we used a digital elevation model to calculate (*i*) the height difference in degree

182   (*pente_moy*) between the start and the end of the river segment, which is associated with the

speed, depth and turbidity of the waters and which we thus believe is associated with the

184   fishing conditions for the birds; (*ii*) the average elevation in m of the 250 m long river

segment (*alt_moy*). We believe this accounts for the elevation gradient in plant composition

186   and land use by farmers and other human activities, because at high elevation the rainforest

dominates, while in the lowland, forests were often cleared for banana and crops plantation,

188   particularly on the east coast; (*iii*) the bearing in degrees (*angle*). We believe this accounts for

the contrasting ecosystems between the windward and leeward coasts, because ombrophilous

190   forest is generally found on both coasts, while seasonal evergreen and semi-deciduous forest,

in contrast, are found only on the west coast (Van Laere *et al.* 2016). Finally, we added (*iv*)

192 the chlordecone pollution level according to the classification and map of land pollution risks

by Rochette *et al*. (2017). Rochette et al. (2017) assessed soil pollution using a multiresidue

194 analysis of pesticides of 227 samples on randomly chosen rivers plus 120 samples collected at

the estuary of 59 different watersheds, and interpolated this data with a Kriging procedure to

196 produce a map of chlordecone pollution risks (Rochette *et al*. 2017). Class 5 soils have "very

high risk" of chlodercone pollution (>90%) covering 4,761 ha (5.6% of Basse-Terre land); a

198 class 4 "high risk" pollution (~80%) making 1,629 ha (1.9%); class 3 soil has "low risk" of

pollution (~30%) covering 181 ha (0.2%); Class 2 soils have "low to negligible risk" of

200 chlordecone pollution risks (<10%), covering 27,929 ha of the island (32.9%); Finally the last

class 1 should show "no pollution by chlordecone", representing the remaining 59.3% of the

202 Basse-Terre area. We assessed the pollution risk of each river segment (*val_tc_chlor*) by

assigning it the highest pollution risk class of all contiguous areas.

204 To explore whether Ringed Kingfishers avoided polluted areas, we calculated selection

ratios (noted SR), a classical measure of preference and avoidance in habitat selection studies

206 (Manly *et al.* 2007). An SR is calculated by dividing the proportional use of a given habitat by

the proportional availability of that habitat. An SR value of one means that birds use the focal

208 habitat randomly (i.e., use is equal to availability). When individuals avoid a particular

habitat, it will be used to a lower extent than its availability and the SR will be <1.

210 Conversely, a preferred habitat will have a SR >1. Note that habitat preference and avoidance

are the standard terms used in habitat selection studies and describe an observed pattern that

212 can be the result of many different behaviors and factors—e.g., the active avoidance of a

threat, or the absence of a species in a certain habitat due to a lack of suitable resources. For

214 all sampled river segments, we calculated the proportions falling into each of the five

chlordecone pollution risk classes. We repeated this calculation for the subset of river segments where we observed Ringed Kingfishers. We then calculated selection ratios for each pollution class by dividing, for a given pollution class, the proportion of river segments with Ringed Kingfisher observations by the proportion of all river segments in that class. We computed the variance for each SR according to Aho and Bowyer (2015) for the variance of the two ratios.

We complemented the selection ratio analysis of Ringed Kingfisher with a multivariate approach called the MADIFA (Mahalanobis Distance Factorial Analysis, Calenge *et al.* 2008). A principal component analysis is first performed on the environmental descriptors to uncover the most influential variables structuring the landscape. This multivariate description of the environment corresponds to the ecological space available to the birds. Then the method compares, in the ecological space defined above, where birds are present and absent. If birds select the habitat along some variables, the locations where birds were detected should be associated with the one or several axes. An advantage of the method over the selection ratios (Manly *et al.* 2007) is that spatial covariation among the different environmental descriptors we included is visualized and accounted for by the MADIFA. As a side note, none of the methods could account for potential pseudo-replication in the data arising from the repeated observations of the same individuals. We hence treated each observation as independent record, and assumed little or no effect of pseudo-replication on our results. We ran all analyses using R 3.5 (R core team 2019) and the associated package *asbio* (Aho 2019) and *adehabitatHS* (Calenge 2006).

Results

We surveyed 270 km of river habitat over a period of 376.6 hours (93.8% of the habitat by foot and 6.2% by kayak). Thus we surveyed 69% of the total length (395 km) of the 44 rivers on Basse-Terre that we sampled representing 27% of the total length of permanent rivers on Basse-Terre that were available to the Ringed Kingfisher. We recorded a total of $n =$ 47 Ringed Kingfisher locations. From these observations, we tentatively assessed the possible number of different birds to 16 among which 12 were seen as singletons, and 2 couples.. On Basse-Terre the total length of rivers with permanent water was 1,012 km. During our survey we noticed that rivers shorter than 2 km long and representing 113 km out of the 1,012 km were not occupied by Ringed Kingfishers. Thus, river habitat available to Ringed Kingfishers that we later used for our habitat selection analyses reduced to 899 km long (including 395 km for the 44 rivers sampled). Since our study was conducted, 13 new Ringed Kingfisher observations have been recorded between 2013 and 2017 by birdwatchers (Fig. 1a). Note that these new locations where birds were seen fit in our Ringed Kingfisher habitat selection patterns of an avoidance of polluted river segments whether these observations were made in areas we previously visited during our sampling (9 observations) or not (4 observations). .

We overlaid the data on Ringed Kingfisher distribution with the map of level of chlordecone contamination risk (Fig. 1a) and forest habitat (Fig. 1b). Our results suggest that Ringed Kingfishers (*i*) are found outside of high chlordecone contamination areas (> 80% polluted, classes 4 and 5 of chlordecone contamination level), and (*ii*) mostly in the ombrophilous forest. The statistical analysis of habitat selection confirmed the visual inspection of Ringed Kingfisher locations on the map. River segments with no chlordecone were used slightly more relative to availability ($SR_1 = 1.18 \pm 0.01$; Fig. 2). Conversely, Ringed Kingfisher used river segments with any risk of chlordecone pollution significantly less relative to its availability (for all river segments with pollution risks of 2 to 5: $SR_{2-5} =$

0.46 ± 0.30, p = 0.07). Biologically speaking it means that the odds to find a Ringed Kingfisher are 2.56 higher in unpolluted than in polluted river segments on Basse-Terre, on average. When considering each pollution level separately, all estimates of selection ratios of polluted river segments were < 1, although not statistically different from one (see Fig 2; SR_2 = 0.32 ± 0.98; SR_3 = 0.00 ± 2.15; SR_4 = 0 .00 ± 2.08; SR_5 = 0.90 ± 0.50).

The multivariate habitat selection analyses lend further supports to the selection ratio results. The correlation circle (Fig. 3A) showed that elevation and bearing correlated positively because high elevation areas are found mostly in the southwest of the island. Importantly, the level of chlordecone pollution was orthogonal to the other environmental variables, suggesting that the use of the pollutant in the landscape has been (statistically) independent of the other landscape variables used to described the Ringed Kingfisher environment. It also means that any effect of chlordecone we would find on the Ringed Kingfisher habitat selection would not be the outcome of potential confounding effect of the described landscape structure. When projecting the points on the first two axes of the ecological space, the MADIFA shows that Ringed Kingfisher (black dots, Fig. 3B) selected high elevation river segments rather than low elevation segments. Similarly, river segments with too high a denivelation were used less intensively than flatter ones. Furthermore, none of the river segments where birds were seen are on the lower-left quarter of the plot , which represents the ecological space with chlordecone pollution (black dots, Fig. 3B). . Nonetheless, coupled with the selection ratios, this multivariate analysis is congruent with the idea that on Guadeloupe, the Ringed Kingfisher might use river segments polluted by chlordecone less than their availability. Accordingly, none of the river segments where birds were seen were found on the upper-right quarter of the plot (black dots, Fig. 3B), which represents the ecological space with chlordecone pollution.

Discussion

288   Given the particular agricultural history of some islands in the Antilles, we put forward the

hypothesis that long-lasting effects of chloredone pollution, which persists in the environment

290   for centuries (Crabit *et al*. 2016), could have had adverse ecological conditions on freshwater

ecosystems of Guadeloupe, including the Ringed Kingfisher. On Basse-Terre, we document a

292   lower relative use of the lower sections of rivers by the Ringed Kingfisher, which run though

the heavily chlordecone contaminated estuary on the east coast (Fig. 1). A study carried out on

294   the European Dipper (*Cinclus cinclus*) found a similar absence of this river-dwelling species

in 93.7% of polluted or strongly polluted streams in Italy (Sorace *et al*. 2002). Chlordecone

296   pollution could therefore at least partly explain the past and current limited distribution of

Ringed Kingfishers on Basse-Terre that we describe here (Fig. 1a). In the 1970's, Ringed

298   Kingfishers were regularly recorded in Sofaïa, Capesterre and Trois-Rivières on this island

(Pinchon 1976). Forty years later, we detected the species in the first two locations still, but

300   only in upstream areas; one possible explanation is that chlordecone has not been spread by

banana farmers in upstream areas. Overall, the Ringed Kingfisher is not found in the lower

302   part of rivers or the ocean front on Basse-Terre (Fig. 1). This could be related to the 84.1 % of

watersheds analyzed having pesticides such as chlordecone (Rochette *et al*. 2017). Finally

304   during the 10 years following our survey, birdwatchers recorded several sightings of Ringed

Kingfishers on Guadeloupe, but these were all made outside of areas of high chlordecone

306   pollution risks. Regarding the four Ringed Kingfishers detected in polluted area (class 4 and

5), they were found where land pollution was patchily distributed, with access to some

308   unpolluted areas as well.

The spatial avoidance of polluted river segments by Ringed Kingfisher could result

310   from direct and indirect effects of chlordecone on the birds. Direct effects usually lead to the

death of individuals or regular reproduction failures because of organism dysfunction when

312   exposed to chemical pollutants (Saaristo *et al.* 2018). Although anecdotal, we could observe

the feeding of Ringed Kingfisher chicks with basket shrimp *Atya innocous,* stream shrimp

314   *Macrobrachium crenulatum,* and the river goby *Awaous banana*. An ecolotoxicology study

carried out on southern Basse-Terre river evidenced substantial amount of chlordecone in

316   those three prey species (Coat *et al.* 2011). Although we currently lack of published material

on the Ringed Kingfisher diet in the tropics to strengthen this observation, we can expect a

318   direct effect of chlordecone *via* bioaccumulation in its tissue from polluted food resource in

lowlands of Guadeloupe. Clearly, an epidemiology study on the Ringed Kingfisher is most

320   needed to assert the physiological effects of chlordecone on the bird organism and life

histories. Moreover, indirect effects of chlordecone may proceed from cascading effects of

322   contamination on prey species community or size structure of the populations and, in turn, can

lower the suitability of habitat for predators in terms of food resources (Saaristo *et al.* 2018).

324   With no bird banding we cannot assess whether kingfishers actually move from one river to

another to actively avoid chlordecone pollution, but over time both direct and indirect effects

326   could have led to the absence of individuals from polluted areas as we suggest is the case on

Guadeloupe.

328   Our results indicate that other ecological factors might also be related to the distribution

of the Ringed Kingfisher on Guadeloupe. For example, river segments located on the eastern

330   coast are used more than expected from its availability by Ringed Kingfishers (Fig. 3), and we

observed the Ringed Kingfisher most on the windward east coast (Fig. 1b), where

332   ombrophilous forests grow. Rivers running though this type of forest are long with gentle

slopes, and river banks offer many clay cliffs to the birds for nesting. In this ecosystem, food

334 resources are likely available to Ring Kingfishers given the many school fishes we saw while

walking the rivers. The vegetation structure of the forest also provides numerous perches ideal

336 for birds to fish. This ombrophilous forest hence appears a rather suitable habitat for the

Ringed Kingfisher both from a food resource viewpoint, and for nest excavation too (Woodall

338 2001). Conversely, on the dryer leeward west coast fish abundance is lower and rocks make

most of the river banks. On average rivers are also shorter (1.8 km *vs*. 3.0 km) and steeper

340 (16.6° *vs.* 11.0°) hence presenting more rapid water streams. We noted previously that no

kingfisher were using streams of < 2km long, maybe because it is too shallow for diving.

342 Generally speaking the west coast habitat seems less favorable to Ringed kingfishers because

of river characteristics, low food resource and the lack of soft banks forcing birds to fly far

344 from waters to dig its burrow nest (Woodall 2001). Accordingly, we detected only two

individuals on the west coast. One exception was an individual living in the northwest (Fig.

346 2), that we could easily associate with the presence of a fish farm where it is seen fishing

regularly.

348   Elevation is another influential ecological variable of Ringed Kingfisher habitat

selection. Here, elevation likely serves as a proxy of river physical characteristics, as water

350 speed, depth, temperature or turbidity all vary gradually from the high elevation river source

to the estuary (Allan 1995). It is known that the species needs fairly deep water to dive and

352 should hence be most abundant in the lowlands (Skutch 1972). We found the opposite pattern

and attributed this difference to a combination of chlordecone pollution and unsuitable

354 habitats for Ringed Kingfishers. On Basse-Terre, although chlordecone pollution of soils and

waters decreases with altitude (Fig. 1), our river sampling resulted in a weak spatial

356 correlation between the two variables at the landscape level (Fig. 3a). Statistically speaking,

we could decipher the effect of chlordecone pollution risks and of elevation, and found that

Ringer Kingfisher was present relatively less in polluted river segments or in those located at

low elevation (Fig. 3b). That the birds avoid unpolluted lowland river segments could be

explained by the marked human footprints on the beds, banks and surroundings of rivers in

Guadeloupe. For instance, the riverbed of the river flowing through the city of Basse-Terre is

made of concrete mostly, which probably reduces fish abundance drastically. Similarly in

banana plantations the forest is sometimes clear cut down to the river edge, except from a

narrow strip of trees along river banks, here again modifying the vegetation structure and

density (note that here a lack of perches could not explain the low relative use of Ringed

Kingfishers in banana plantation). Lowlands of Guadeloupe islands are at the heart of human

activities, which generates all kinds of perturbations for wildlife and possibly for the Ringed

Kingfisher. Light pollution, chlordecone upstream pollution, active disturbance, presence of

predators or competition for food with fishermen could all contribute to deter Ringed

Kingfishers from lowland rivers beside the polluted river segments.

Our study suggests that the Ringed Kingfisher could be a useful bioindicator of areas

with organochlorine pollution, and most specifically chlordecone. We contend that our habitat

selection analysis lacks statistical power but given the potentially dramatic consequences for

the Ringed Kingfisher, the low level of statistical significance should not hamper the

biological signal we document here (see Yoccoz 1991 for a discussion of statistical

significance). Although we cannot prove the generalized ecological effects of chlordecone in

the Antilles, we advance that historical chlordecone use by banana farmers could have had

and be still having an ongoing influence on Ringed Kingfisher populations and distribution on

certain islands, including Guadeloupe. To further explore this potential issue, researchers

should collect data on the ecology of the Ringed Kingfisher with a detailed descriptions of

diet through the year, habitat, territory size, and behavior, and also to investigate the

382 ecological consequences of chlordecone pollution could have on individuals, populations and

avian communities.

Acknowledgements

386 This work was supported by the national park of Guadeloupe, DEAL Guadeloupe (the

Regional Department of Environment) and AEVA. We thank Stéphane Di Mauro for preys

388 determination, Gavin Hunt for editing the English and an anonymous reviewer for his

constructive comments. We thank Shirley Dolo for the Spanish version of the summary. We

390 acknowledge the insightful and extensive help of Alice McBride who greatly improved our

manuscript.

Literature Cited

394 Aho, K. 2019. Asbio: A collection of statistical tools for biologists. R package version 1.5-5.

https://CRAN.R-project.org/package=asbio.

396 Aho, K., and R.T. Bowyer. 2015. Confidence intervals for ratios of proportions: implications

for selection ratios. Methods in Ecology and Evolution 6:121-132.

398 Allan, J.D. (1995). Stream ecology: structure and function of running waters. Springer

Science and Business Media.

400 Cabidoche, Y.M., and M. Lesueur Jannoyer. 2011. Pollution durable des sols par la

chlordécone aux Antilles : comment la gérer ? Innovations Agronomiques 16:117-133.

402 Calenge, C., G. Darmon, M. Basille, A. Loison, and J.M. Jullien. 2008. The factorial

decomposition of the Mahalanobis distances in habitat selection studies. Ecology,

404 89:555-566.

Calenge, C. 2006. The package adehabitat for the R software: tool for the

406   analysis of space and habitat use by animals. Ecological Modelling, 197:1035.

Campbell L.H., M.I. Avery, P. Donald, A.D. Evan, R.E. Green and J.D. Wilson 1997. A review

408   of the indirect effects of pesticides on birds. Joint Nature Conservation Committee

report 227.

410   Clarkson, T.W. 1995. Environmental contaminants in the food chain. The American Journal of

Clinical Nutrition, 61:682-686.

412   Coat, C., D. Monti, P. Legendre, C. Bouchon, F. Massat, and G. Lepoint. 2011.

Organochlorine pollution in tropical rivers (Guadeloupe): Role of ecological factors in

414   food web bioaccumulation. Environmental Pollution 159:1692-1701.

Crabit, A., P. Cattan, F. Colin, and M. Voltz. 2016. Soil and river contamination patterns of

416   chlordecone in a tropical volcanic catchment in the French West Indies (Guadeloupe).

Environmental Pollution 212:615-626.

418   Douthwaite, R.J. 1982. Changes in pied kingfisher (*Ceryle rudis*) feeding related to

endosulfan pollution from tsetse fly control operations in the Okavango Delta,

420   Botswana. Journal of Applied Ecology 19:133–141.

Eroschenko, V.P. 1981. Estrogenic activity of the insecticide chlordecone in the reproductive

422   tract of birds and mammals. Journal of Toxicology and Environmental Health 8:731-

742.

424   Evans, S.W., and H. Bouwman. 2000. The geographic variation and potential risk of DDT in

the blood of Pied Kingfishers from northern KwaZulu-Natal, South Africa. Ostrich

426   71:351-354.

Geiger, F., J. Bengtsson, F. Berendse, W.W. Weisser, M. Emmerson, M.B. Morales, and S.

428   Eggers 2010. Persistent negative effects of pesticides on biodiversity and biological

control potential on European farmland. Basic and Applied Ecology, 11:97-105.

430 Karunati. The regional platform of the Nature and Landscape Information System (SINP) of

Guadeloupe. http://karunati.fr/.

432 Manly, B.F.L., L. McDonald, D.L. Thomas, T.L. McDonald, and W.P. Erickson. 2007.

Resource selection by animals: statistical design and analysis for field studies. Springer

434   Science and Business Media.

Mineau P. and M. Whiteside 2013. Pesticide acute toxicity is a better correlate of U.S.

436   grassland bird declines than agricultural intensification. PloS One 8:e57457.

Moore, N.W. 1965. Pesticides and birds—A review of the situation in Great Britain in 1965.

438   Bird Study 12:222-252

Parsons, K. C., P. Mineau and R.B. Renfrew 2010. Effects of pesticide use in rice fields on

440   birds. Waterbirds 33:193-218.

Pinchon, R. 1976. Faune des Antilles Françaises : les oiseaux. Fort-de-France.

442 Remsen, J.V.J. 1990. Community ecology of neotropical kingfishers. PhD, University

California Press, Berkeley and Los Angeles.

444 Rochette, R., P. Andrieux, V. Bonnal, and P. Cattan. 2017. Contamination des bassins versants

de la Guadeloupe continentale par la chlordécone. Actualisation des connaissances et

446   cartographie des zones à risque de contamination.

http://daaf.guadeloupe.agriculture.gouv.fr/IMG/pdf/Projet_ChlEauTerre_-

448   _Rapport_Final_2017_cle8e6961.pdhf

Risebrough R.W., F.C. Sibley and M.N Kirven (1971). Reproductive failure of the Brown

450   Pelican on Anacapa Island in 1969. American Birds 25:8-9.

452 Saaristo M., T. Brodin, S. Balshine, M.G. Bertram, B.W. Brooks, S.M. Ehlman, E.S. McCallum, A. Sih, J. Sundin, B.B.M. Wong and K.E. Arnold 2018. Direct and indirect effects of chemical contaminants on the behaviour, ecology and evolution of wildlife.

454 Proceedings of the Royal Society of London, Series B 285:20181297.

Schreiber, E.A., and J. Burger. 2002. Effects of chemicals and pollution on seabirds. Pp. 485-

456 525 in Biology of marine birds (E.A. Schreiber and J. Burger, eds.). CRC Press, Boca Raton.

458 Schäfer, R.B., P.J. van den Brink and M. Liess 2011. Impacts of pesticides on freshwater ecosystems. In *Ecological Impacts of Toxic Chemicals*, eds Sánchez-Bayo F., van den

460 Brink P.J., Mann R.M. (Bentham Science Publishers Ltd., Bussum, The Netherlands).

Skutch, A.F. 1972. Ringed Kingfisher C*eryle torquata*. Pp. 88-101 in Studies of tropical

462 American birds. Publications of the Nuttall Ornithological Club, Cambridge, Massachusetts.

464 Sorace A., P. Formichetti, A. Boano, P. Andreani, C. Gramegna, and L. Mancini. 2002. The presence of a river bird, the dipper, in relation to water quality and biotic indices in

466 central Italy. Environmental Pollution 118: 89-96.

Tanabe, S., K. Senthilkumar, K. Kannan, and A.N. Subramanian. 1998. Accumulation features

468 of polychlorinated biphenyls and organochlorine pesticides in resident and migratory birds from South India. Archives of Environmental Contamination and Toxicology 34:

470 387-397.

USEPA (U.S. Environmental Protection Agency) 1975. DDT - A review of scientific and

472 economic aspects of the decision to ban its use as a pesticide. Washington D.C.

Van Laere, G., Y. Gall and A. Rousteau 2016. The forest ecosystems observatory in

474 Guadeloupe (FWI). Caribbean Naturalist 1:108-115.

White, A.E., and D.A. Cristol. 2014. Plumage coloration in belted kingfishers (*Megaceryle*

476  *alcyon*) at a mercury-contaminated river. Waterbirds 37:144-152.

Wiemeyer, S.N., C.M. Bunck, and A.J. Krynitsky. 1988. Organochlorine pesticides,

478  polychlorinated biphenyls, and mercury in osprey eggs—1970-79—and their

relationships to eggshell thinning and productivity. Archives of Environmental

480  Contamination and Toxicology 17: 767-787.

Woodall, P.F. 2001. Family Alecedinidae. Pp 130-149 *in* Handbook of the birds of the world

482  (J. del Hoyo, A. Elliott and J. Sargatal, eds.). Vol. 6. Mousebirds to hornbills. Lynx

Edicions, Barcelona.

484 Yoccoz, N.G. 1991. Use, overuse, and misuse of significance tests in evolutionary biology

and ecology. Bulletin of the Ecological Society of America 72:106-111.

486 Zamani-Ahmadmahmoodi, R., A. Esmaili-Sari, S.M. Ghasempouri, and M. Savabieasfahani.

2009. Mercury levels in selected tissues of three kingfisher species; *Ceryle rudis,*

488  *Alcedo atthis,* and *Halcyon smyrnensi,* from Shadegan Marshes of Iran. Ecotoxicology

18:319-324.

Contributions

492 P.V. designed the study, carried out fieldwork and wrote the paper; A.F. made all maps and

extracted with a SIG the data for analysis; P.F. and C.P. helped shape the report and C.B.

494 performed the statistical analysis, wrote their interpretation and contributed to the writing.

Figure legend

496 **Fig. 1:** Location of Ringed Kingfisher (*Megaceryle torquata stictipennis*) encounters (2009 to

2017) on Basse-Terre, Guadeloupe (source for maps: BDTOPO – Institut Géographique

498 National Paris; SRTM – DAAF Guadeloupe). In total, we recorded 47 observations of Ringed

Kingfishers but the number of different birds remains uncertain, because some indivdiuals

500 might have been seen several times if disturbed by the observer. A gross estimate of the

number of different birds we have detected is 16. We overlappedthe observation locations on

502 the map of (a) chlordecone exposure risk and (b) the map of forest habitats.

504 **Fig. 2:** Selection ratios (with associated 95% confidence limits) based on Ringed Kingfisher

observations made along river segments with five different levels of chlordecone pollution on

506 Basse-Terre, Guadeloupe between 2009 and 2017 ($n = 47$ bird locations on approximately 16

different individuals). Chlordecone levels from 1 to 5 are increasing risks to find polluted

508 river segments, while the last category pooled all polluted segments whatever its risks.

Although selection ratios are not statistically different from 1, Ringed Kingfishers seem to use

510 non-polluted segments of rivers (chlordecone level 1) more relative to availability, contrasting

with river segments with low to high levels of pollution (chlordecone level 2 to 5) which are

512 used less relative to availability.

514 **Fig. 3**: Graphical output of the MADIFA (Mahalanobis Distance Factorial Analysis, Calenge

et al. 2008) displaying the ecological conditions prevailing in river segments where Ringed

516 Kingfishers (*Megaceryle torquata stictipennis*) were observed (black dots) compared to

available river segments on the island (grey dots), on Basse-Terre, Guadeloupe, between 2009

518 and 2017 ($n = 47$ bird locations on approximately 16 different individuals). We used 4

variables to describe the environmental condition for each 250 long river segments: the

520   average height difference in degree (Elevation diff.), the average elevation in m (Elevation),

the bearing in degrees (angle), and the chlordecone pollution level (chlordecone). (a)

522   Correlation circle of the principal component analysis of the river segments characteristics

showing that chlordecone pollution is only weakly associated with other environmental

524   descriptors we used; (b) the MADIFA shows that none of the river segments where birds were

seen were found on the upper-right quarter of the plot (black dots), which represents the

526   ecological space with low to high chlordecone pollution.

Fig. 1

a)

Source : BDTOPO - IGN Paris - SRTM - DAAF Guadeloupe - BD ObsOcc PNG

554   b)

Fig. 2

Fig. 3

(A)

(B)