1   **Deciphering cellular transcriptional alterations in Alzheimer's disease brains**

3   Xue Wang[1*], Mariet Allen[2], Shaoyu Li[3], Zachary S. Quicksall[1], Tulsi A. Patel[2], Troy P.

4   Carnwath[2], Joseph S. Reddy[1], Minerva M. Carrasquillo[2], Sarah J. Lincoln[2], Thuy T. Nguyen[2],

5   Kimberly G. Malphrus[2], Dennis W. Dickson[2], Julia E. Crook[1], Yan W. Asmann[1], Nilüfer

6   Ertekin-Taner[2,4*]

[1] Department of Health Sciences Research, Mayo Clinic Florida, Jacksonville, FL, USA.

[2] Department of Neuroscience, Mayo Clinic Florida, Jacksonville, FL, USA.

[3] Department of Mathematics and Statistics, University of North Carolina at Charlotte, Charlotte,

NC, USA.

[4] Department of Neurology, Mayo Clinic Florida, Jacksonville, FL, USA.

* Corresponding Authors

E-mails: Wang.Xue@mayo.edu (XW) and Taner.Nilufer@mayo.edu (NET)

ABSTRACT

Large-scale brain bulk-RNAseq studies identified molecular pathways implicated in Alzheimer's disease (AD), however these findings can be confounded by cellular composition changes in bulk-tissue. To identify cell intrinsic gene expression alterations of individual cell types, we designed a bioinformatics pipeline and analyzed three AD and control bulk-RNAseq datasets of temporal and dorsolateral prefrontal cortex from 685 brain samples. We detected cell-proportion changes in AD brains that are robustly replicable across the three independently assessed cohorts. We applied three different algorithms including our in-house algorithm to identify cell intrinsic differentially expressed genes in individual cell types (CI-DEGs). We assessed the performance of all algorithms by comparison to single nucleus RNAseq data. We identified consensus CI-DEGs that are common to multiple brain regions. Despite significant overlap between consensus CI-DEGs and bulk-DEGs, many CI-DEGs were absent from bulk-DEGs. Consensus CI-DEGs and their enriched GO terms include genes and pathways previously implicated in AD or neurodegeneration, as well as novel ones. We demonstrated that the detection of CI-DEGs through computational deconvolution methods is promising and highlight remaining challenges. These findings provide novel insights into cell-intrinsic transcriptional changes of individual cell types in AD and may refine discovery and modeling of molecular targets that drive this complex disease.

Introduction

Alzheimer's disease (AD) is a neurodegenerative disease that affects ~5.7 million patients with annual cost of more than \$230 billion in the US[1]. Effective disease-modifying drugs are still elusive despite the urgent need and increasing global burden[2,3]. Pathologically, AD is marked by amyloid-beta plaques and neurofibrillary tangles, along with neuronal loss and gliosis in the affected brain regions. Transcriptome-wide expression levels have been analyzed from bulk brain tissue of hundreds of AD patients and neuropathologically normal controls[4-8] to discover genes and biological pathways that are perturbed in and/or lead to AD. Systems biology and bioinformatics analysis of these data have implicated altered pathways in AD including immune response [6] and myelin metabolism [4,5]. However, a fundamental knowledge gap remains concerning whether disease-associated changes in brain gene expression are due to changes in cellular composition of the AD samples secondary to disease neuropathology, or due to changes in the intrinsic regulation/activity of genes in the central nervous system (CNS) cells. From a clinical perspective, it is difficult to target changes in cellular composition secondary to neuropathology, whereas intrinsic changes in gene expression that may drive AD progression are potentially "druggable".

We expect that identifying cell-intrinsic differentially expressed genes (CI-DEGs) of individual CNS cell types will reveal novel insights into the genes and pathways that could potentially identify drug targets and lead to AD therapeutics. This approach circumvents the influence of cell-composition changes that can impact disease associated DEGs obtained from bulk tissue transcriptome analysis. RNA sequencing (RNAseq) studies from single cell, single nucleus or purified adult human CNS cells[9-11] can be used to identify CI-DEGs. Even though such single cell-type RNAseq data can effectively serve as a reference to annotate bulk-tissue

transcriptome data[4], such approaches remain costly, cumbersome and limited in sample sizes. On

the other hand, there exist large-scale bulk brain RNAseq datasets[5,8,12], which can be leveraged to

identify CI-DEGs through analytic deconvolution of bulk tissue expression into signals of

individual cell types by using cell proportions or their proxies[13,14].

In this study, we describe the design and application of a bioinformatics pipeline that uses

cell type marker genes to estimate cell proportion[15,16] to deconvolute bulk tissue transcriptome

data using three computational approaches[13,14,17] and to subsequently identify CI-DEGs. We

applied our pipeline to the analysis of three post-mortem brain datasets, one from dorsolateral

prefrontal cortex (DLPFC)[8] and two from temporal cortex (TCX)[4,12,18] regions, comprised of 685

unique samples. Consensus CI-DEGs common to both TCX and DLPFC regions were analyzed

for enrichment of gene ontology (GO) terms. We compared the results of consensus CI-DEGs to

consensus bulk-DEGs. In addition, for the DLPFC[8] dataset that had both bulk and single nucleus

RNAseq[19] (snRNAseq) data, we compared the CI-DEGs from the computational deconvolution

to CI-DEGs obtained from snRNAseq[19].

To our knowledge, this is the first study to detect consensus CI-DEGs and their enriched

gene ontology (GO) terms from multiple brain regions using multiple computational

deconvolution algorithms for AD and control RNAseq samples. Our study illustrates the cell

proportion landscape of AD and control brain regions assessed in three independent RNAseq

studies[4,7,8,12]. We identify consensus CI-DEGs many of which are not observed in bulk-DEG

analysis and characterize their cell-type specificity. GO terms that are enriched for CI-DEGs

implicate cell intrinsic transcriptional alterations that may influence AD, rather than be a result

of cell-proportion changes in this disease. These CI-DEGs and their biological pathways may

serve as refined molecular targets for therapeutic discoveries and disease modeling in AD. Our

study also demonstrates that detection of CI-DEGs through computational deconvolution

methods is promising while some challenges remain.

Results

Cellular composition in three brain cohorts from two brain regions

We analyzed three cohorts each consisting of post-mortem brains from AD and control

subjects (**Table S1**), namely the Rush Religious Orders Study and Memory and Aging Project

dorsolateral prefrontal cortex (DLPFC)[7,8], Mayo Clinic temporal cortex (TCX-Mayo)[4,12], and

Mount Sinai VA Medical Center Brain Bank temporal cortex (TCX-MSBB)[18]. We generated the

TCX-Mayo RNAseq dataset, and downloaded DLPFC and TCX-MSBB RNAseq datasets from

the AMP-AD knowledge portal on Synapse (www.synapse.org).

Cell proportions (**Table S2**) were estimated for DLPFC, TCX-Mayo and TCX-MSBB

datasets independently using the digital sorting algorithm (DSA) method[16] and the top 100

marker genes (**Table S3**) obtained from R package BREGITEA[15] for each of the following cell

types – neuron, oligodendrocyte, microglia, oligodendrocyte progenitor cell (OPC), astrocyte and

endothelial cell.

An inspection about the pairwise correlation between marker genes (**Fig 1a**) revealed that

markers of OPC have poor median pairwise Pearson correlation values of 0.12 in DLPFC, 0.11

in TCX-Mayo and 0.06 in TCX-MSBB respectively, whereas among the other five cell types

neuronal markers have the highest median correlation (0.68 in DLPFC, 0.78 in TCX-Mayo and

0.67 in TCX-MSBB), and microglia markers have the lowest correlation (0.37 in DLPFC, 0.42

in TCX-Mayo and 0.44 in TCX-MSBB). In addition, a computer simulation study (**Fig S1**)

demonstrated that the estimated proportions of OPC were not robust upon using different

selection of marker genes. Therefore, we did not include OPC in downstream analyses in this

study.

In all three datasets, neuronal cell proportion estimates were significantly lower in AD

compared to controls (**Fig 1b**). The magnitude of this decrease was the greatest for TCX-Mayo

(AD mean proportion = 28.0%, Control = 35.7%; ratio of AD:control cell proportions=0.78),

followed by TCX-MSSM (AD = 42.3%, control = 49.3%; ratio=0.87) and DLPFC (AD = 42.4%,

control = 47.4%; ratio=0.89). The estimated proportions of microglia were significantly higher in

AD vs. controls for all datasets, with higher magnitude in TCX-Mayo (AD:control ratio=1.19)

and TCX-MSBB (AD:control ratio=1.19) than for DLPFC (AD:control ratio=1.06). The

estimated proportions of astrocytes and endothelial cells were significantly higher in AD vs.

controls for DLPFC and TCX-Mayo datasets, although the magnitude was greater in TCX-Mayo

(1.40 and 1.30 respectively) than in DLPFC (1.07 and 1.14 respectively) for both cell types.

Oligodendrocyte proportion is significantly higher in AD in DLPFC with AD:control ratio 1.14

and TCX-MSBB with AD:control ratio 1.27, although remains unchanged in TCX-Mayo with

the ratio 0.94. Collectively, these findings demonstrate that the proportions of CNS cell types are

different in post-mortem AD vs. control brains for most cell types. Although these proportional

changes with AD are mostly consistent across the different studies, their extent varies across

brain regions, with TCX tending towards higher magnitude of neuronal loss and microglia

proliferation than DLPFC.

Fig.1: **a)** Pearson correlation between marker gene expressions in six cell types. Marker genes are from literature. **b)** Estimated cell proportions in DLFPC, TCX-Mayo and TCX-MSBB datasets in five cell types. Red asterisk indicates differences between cell proportions in AD and control groups at nominal p value 0.05 from Wilcoxon rank sum test.

Differential Expression Analyses

In this study, three computational approaches were applied to identify cell intrinsic differential expression in individual cell types (CI-DEGs, **Table S4-S6**), namely CellCODE[14], PSEA[13] and our method WLC. Differentially expressed genes from bulk brain tissue (bulk-DEGs) were identified through linear regression without adjusting for cellular composition (**Table S7**). For the DLPFC, TCX-Mayo and TCX-MSBB datasets, we obtained bulk-DEGs and

CI-DEGs from the three computer algorithms for neuronal, oligodendrocytic, microglial,

astrocytic and endothelial cell types respectively.

We compared bulk-DEGs across the three datasets (**Fig 2a**, top panel). Similarly, CI-

DEGs from CellCODE, PSEA and WLC are compared across datasets (**Fig 2a**, lower panels),

such that CI-DEGs shared between datasets are required to be consistent in the designated cell

type. All DEGs are identified at nominal p-value cutoff 0.05 and shared CI-DEGs have the same

direction of change in the compared datasets. The ratio of overlap between any two datasets over

all DEGs, i.e. the number in overlapping areas of the Venn diagram over the total number (**Fig**

2a, top panel), is 30.0% or 1711/5697 in up-regulated bulk-DEGs, and 34.8% or 2214/6371 in

down-regulated bulk-DEGs. This ratio of overlap in bulk-DEGs is much higher than that in CI-

DEGs (2.7%, 4.7% and 10.0% in up-regulated genes from CellCODE, PSEA and WLC

respectively; 3.1%, 6.8% and 9.3% in down-regulated genes from CellCODE, PSEA and WLC

respectively).

Fig.2: **a)** Overlap across three independent RNAseq datasets of bulk-DEGs (upper panel) and CI-DEGs (lower panels) from three computational approaches. **b)** Consensus CI-DEGs between DLPFC and TCX brain regions, which consist of consensus CI-DEGs between DLPFC and TCX-Mayo, or between DLPFC and TCX-MSBB. c) Overlap between consensus CI-DEGs and consensus bulk-DEGs, per cell type. The p-values of overlap are from hypergeometric tests.

Consensus CI-DEGs between DLPFC and TCX

165   To obtain the consensus CI-DEGs that are shared between DLPFC and TCX brain

166 regions, we selected those CI-DEGs that are detected in "DLPFC and TCX-Mayo" or in

167 "DLPFC and TCX-MSBB" under any of the three algorithms (**Fig S2**). We combined all such

168 genes, which collectively comprised the consensus CI-DEGs for each cell type (**Fig 2b**).

169 Similarly, consensus bulk-DEGs were the combined set of bulk-DEGs shared between "DLPFC

170 and TCX-Mayo" or "DLPFC and TCX-MSBB".

171   Most consensus CI-DEGs are from neuronal cells (N=554), followed by

172 oligodendrocytes (N=249), whereas microglia contributed the least number (N=100). The

173 majority (65.7% or 364/554) of neuronal CI-DEGs is down-regulated in AD, and the majority

174 (65.9% or 139/211) of endothelial CI-DEGs is up-regulated in AD, with other cell types lying in

175 between. Some of these CI-DEGs are also among the 1000 marker genes of the corresponding

176 cell type from BRETIGEA[15]; 14.8% or 82/554 of neuronal CI-DEGs are also neuronal markers,

177 26.5% or 66/249 of oligodendrocyte CI-DEGs are also oligodendrocyte markers, and other cell

178 types lie in between.

179   With regards to consensus bulk-DEGs (**Fig S3**), 28.2% of them (885/3135) are cell type

180 markers; 10.4% neuronal markers, 5.6% oligodendrocyte, 3.4% microglia, 4.8% astrocyte and

181 4.0% endothelial markers. The above observations indicate that computational deconvolution

182 algorithms could identify CI-DEGs for both marker genes and non-marker genes. Importantly,

183 the proportion of non-marker CI-DEGs is greater than that in bulk-DEGs. This suggests that

184 compared to bulk-DEGs, CI-DEGs may be capturing a greater proportion of expression changes

185 that are not due to mere cell population changes.

186   We also compared the consensus bulk-DEGs with consensus CI-DEGs (**Fig S4**). We

187 determined that only a small fraction (14.7% or 28/190) of the up-regulated neuronal CI-DEGs

was also present in up bulk-DEGs although the overlap is still significant (**Fig 2c**). In

comparison, most of the up-regulated CI-DEGs of the other four cell types were included in up

bulk-DEGs. On the other hand, most (85.0% or 308/365) of the down-regulated neuronal CI-

DEGs were also present in down bulk-DEGs, whereas most of the down-regulated CI-DEGs of

the other four cell types were absent from this group. Since bulk-DEGs did not adjust for

neuronal loss and gliosis in AD (**Fig 1b**), its ability to identify up-regulated neuronal genes and

down-regulated glial genes is likely to be compromised. For the same reason, bulk-DEGs may

have a false inflation of detecting down-regulated neuronal and up-regulated glial genes.

Enriched GO terms of consensus CI-DEGs between DLPFC and TCX

198   To identify pathways implicated by CI-DEGs that are robust across brain regions, we

199   performed Gene Ontology (GO) enrichment analysis[20,21] for the consensus CI-DEGs, assessing

200   separately those that are up vs. down in AD subjects (**Table S8-S17**). **Fig 3** illustrates the top

201   two enriched GO terms by enrichment p-values, after filtering out terms that encompass less than

202   four CI-DEGs or are cellular compartments.

203   Consensus CI-DEGs revealed biological pathways that are perturbed in AD in specific

brain cell types. Some of these pathways have previously been implicated in AD and others are

novel. Down-regulated neuronal CI-DEGs were enriched in neuropeptide hormone activity

(GO:0005184) and hormone activity (GO:0005179) pathways, which include *VGF* (a.k.a.

neuroendocrine regulatory peptide 1)[22] and corticotropin releasing hormone (CRH)[23] (**Table**

S13). Consensus up-regulated neuronal CI-DEGs were significantly enriched in potassium

channel activity (GO:0005267) and regulation of ion transport (GO:0043269) pathways (**Table**

S8). The latter GO term encompasses most of the genes from the former, and also includes other

genes involved in neuronal functions such as the glutamate ionotropic receptor NMDA type

subunit 1, *GRIN1*[24] and *SYT1*, which encodes the synaptic vesicle protein, synaptotagmin[25].

Many of the most significant GO terms are related to key functions of the respective cell

types for the glial CI-DEGs, as well. The top enriched pathway of down-regulated CI-DEGs in

oligodendrocytes is myelination (GO:0042552), including myelin basic protein (*MBP*)[4],

plasmolipin (*PLLP*)[4,5], myelin and lymphocyte protein (*MAL*), and myelin-associated

glycoprotein (*MAG*)[26] (**Table S14**). Up-regulated CI-DEGs of oligodendrocytes are enriched in

ceramide biosynthetic process (GO:0046513) including ceramide synthase 4 (*CERS4*) and UDP

glycosyltransferase 8 (*UGT8*)[5] (**Table S9**). Ceramide is a constituent of sphingomyelin, a

sphingolipid which is particularly found in the myelin sheath; and also a multi-functional

signaling molecule[27,28]. Hence, both the down-regulated and the up-regulated oligodendroglial

consensus CI-DEGs highlight different components of the myelin biology that are perturbed in

AD.

Similarly, microglial, astrocytic and endothelial CI-DEGs also highlight processes

pertinent to the functions of these cell types. Microglial up-regulated CI-DEGs are enriched in

inflammatory response (GO:000695) and leukocyte activation (GO:0002696), which includes

complement C3a receptor 1 (*C3AR1*)[29], interleukin 18 (*IL18*)[30,31] and CCAAT enhancer binding

protein alpha (*CEBPA*)[32] genes (**Table S10**).

Astrocytes, a cell type that plays a critical role in maintaining brain energy dynamics[33]

and metabolism[34], show enrichment of oxidoreductase activity (GO:0016491) and drug

metabolic process (GO:0017144) in down-regulated CI-DEGs which includes genes glutathione

S-transferase mu 2 (*GSTM2*)[35] and thioredoxin2 (*TXN2*)[36] (**Table S16**). Astrocytic up-regulated

consensus CI-DEGs are enriched for cell-cell junction assembly (GO:0007043) process (**Table**

S11), including the astrocytic gap junction protein connexin43 (*GJA1*)[37], which was identified as

a key regulator associated with AD related outcomes. The other top GO process for astrocytic

up-regulated consensus CI-DEGs is adenylate cyclase-inhibiting G protein-coupled receptor

signaling pathway (GO:0007193), which harbors adenylate cyclase 8 (*ADCY8*), involved in

memory functions[38].

Finally, endothelial cells, which are crucial in maintaining blood-brain barrier

integrity[39,40], show enrichment of up-regulated DEGs in cytoskeleton organization

(GO:0007010) and actin filament-based process (GO:0030029) (**Table S12**).

Importantly, some CI-DEGs highlight protein translation as a top perturbed biological

pathway. Down-regulated microglial consensus CI-DEGs show enrichment in processes

involved in protein translation (GO:0006614 and GO:0006613), which include ribosomal protein

encoding genes[41-43] (**Table S15**). Similarly, down-regulated endothelial consensus CI-DEGs also

harbor ribosomal protein encoding genes, with enrichment in protein translation related GO

processes (GO:0006413 and GO:0006613) (**Table S17**).

Fig 3: Top two enriched GO terms in up (red) or down-regulated (blue) consensus CI-DEGs

between DLPFC and TCX regions, per cell type.

Comparison of CI-DEGs from computational deconvolution vs. snRNAseq

We determined the extent to which each of the three computational deconvolution algorithms could detect CI-DEGs from bulk tissue by comparison of their results with those obtained in a published snRNAseq study[19]. The ROSMAP dataset utilized in our study has both bulk RNAseq from DLPFC (bulk-DLPFC) as well as snRNAseq (snDLPFC) in a subset of its participants[19]. We compared the bulk-DLPFC data deconvoluted with three different algorithms with the published snDLPFC[19] data. Endothelial CI-DEGs were not available from the snRNAseq study, therefore overlap of results could be assessed only for four cell types.

We tested the overlap between the top CI-DEGs for each cell type obtained from deconvoluted bulk-DLPFC and those from snDLPFC ranked by their p values (**Fig 4a**). We evaluated the overlap for a range of top CI-DEGs up to top 1,000 genes. Overlap for CI-DEGs that are either up (**Fig 4a, upper panel**) or down (**Fig 4a, lower panel**) in AD were assessed separately. Hence, overlapping genes had both similar ranks and direction of effect in both deconvoluted bulk-DLPFC and snDLPFC analyses. We established the significance of overlap using simulations for a range of top ranked genes (N=200, 600 and 1,000) (**Table S18**).

Neuronal CI-DEGs retained their significance of overlap across all comparisons and for all algorithms, except for the top 1,000 up-regulated neuronal CI-DEGs deconvoluted with PSEA. Microglial CI-DEGs had the least numbers of significant overlap for their top ranked genes. Astrocytic and oligodendrocytic top ranked CI-DEGs had significance of overlap between the neuronal and microglial results (**Table S18**). These findings are reflective of the abundance of these cell types, with the most abundant neurons having the most overlap for the top ranked CI-DEGs between deconvoluted bulk-DLPFC and snDLPFC.

Amongst these comparisons, we determined that the significance for overlap was best for

all algorithms for the top ranked 600 genes. Using WLC deconvoluted results, the overlap for the

top 600 CI-DEGs from bulk-DLPFC and snDNPFC are statistically significant for all eight

comparisons (**Fig 4a**). For the top 600 genes, overlap with CellCODE results is significant for all

except down-regulated oligodendrocyte and up-regulated astrocyte CI-DEGs. For PSEA, none of

the microglia CI-DEGs had significant overlap. PSEA results for the top 600 genes were

otherwise significant for all but up-regulated oligodendrocyte and down-regulated astrocyte

genes.

We also performed a comparison of CI-DEGs identified at nominal significance (p-value

<0.05) with each algorithm from bulk-DLPFC to nominally significant snDLPFC results (**Fig**

4b, Table S19). As with the above comparison, genes that are either up or down in both

deconvoluted bulk-DLPFC and snDLPFC data were analyzed separately for each cell type.

Not surprisingly, down-regulated neuronal CI-DEGs have the greatest overlap (537/3732

or 14.4% for WLC, 292/3213 or 9.1% for CellCODE, 415/3516 or 11.8% for PSEA). These

overlaps are significant for all three algorithms (**Table S19**). Down-regulated CI-DEGs in

microglia show the least proportion of overlap (9/723 or 1.2% for WLC, 4/609 or 0.66% for

CellCODE, 16/820 or 2.0% for PSEA) (empirical p-value > 0.05). Significant overlap detected

with WLC (all but down-regulated microglia) and PSEA (all but microglial results and up-

regulated oligodendrocytes) were similar, whereas CellCODE results had significant overlaps

only for the neuronal CI-DEGs (**Table S19**).

Fig 4: Comparison of CI-DEGs from computational deconvolution with CI-DEGs from snRNAseq on DLPFC dataset. **a.** Upper panel: number of overlapping genes (y-axis) between the top N (x-axis) up-regulated genes in snDLPFC and top N up-regulated genes from bulk-DLPFC deconvoluted with PSEA, WLC and CellCODE, respectively. Lower panel: number of overlapping genes (y-axis) between the top N (x-axis) down-regulated genes in snDLPFC and top N down-regulated genes from deconvoluted bulk-DLPFC. **b.** Venn diagram of CI-DEGs from computational deconvolution methods and those from snRNAseq. Overlap is evaluated for both bulk-DLPFC and snDLPFC CI-DEGs detected at nominal p value ≤0.05.

Discussion

There is an increasing number of large scale RNAseq-based transcriptome datasets generated in bulk tissue for many diseases, including brain tissue from patients with AD, other neurodegenerative diseases and controls[5-8,12]. Comparison of such transcriptome data from patient and control individuals has been instrumental in the identification of genes and co-expression networks that are altered in and may therefore be risk factors for these diseases[4-7,44]. The discovery that many of these transcriptional networks harbor genes with disease risk variants provides support for the utility of this bulk-transcriptome approach in deciphering molecules and pathways that are risk factors for these conditions. Nevertheless, there is clear evidence for abundant transcriptome alterations in bulk tissue from affected organs of patients with disease, which appears to be due to changes in cellular composition of that tissue as a consequence of the disease processes[4,45]. Given this, there is a strong need to detect cell-intrinsic transcriptional alterations to be able to distinguish gene expression changes that are upstream of and may therefore be causative factors for disease from those that are merely a result of cell proportion changes that occur due to disease pathology. The discovery of molecules and pathways that are upstream of and risk factors for disease pathology is of paramount importance for development of targeted therapies. This information can also aid in the identification of refined disease biomarkers reflective of disease-causative expression alterations in these conditions. Detection of cell-specific transcriptional changes can also help develop more accurate disease models harboring these cellular alterations. Further, discovery of cell-specific transcriptional alterations in disease may uncover expression changes, particularly in less abundant cell types, which may be missed by the analysis of bulk transcriptome. Thus, there is a growing effort to identify cell-specific expression alterations in human diseases[9-11,14,15,17,46-48].

There are two general approaches to decipher cell-specific transcriptional changes in AD.

One approach is to conduct single nucleus (snRNAseq), single cell (scRNAseq) or purified cell

RNAseq experimentally, followed by data analyses. The alternative approach is to design

relatively complex bioinformatics pipelines to decipher cell-intrinsic information of individual

cell types from bulk tissue microarray or RNAseq data. The first approach is limited due to

significantly higher cost and experimental challenges. Additionally, these approaches may have

the drawback that the procedures of dissociating cells break cell-to-cell communication and thus

may not reflect the true expression signal *in vivo*. The alternative bioinformatics approach to

decipher cell-specific transcriptome alterations from bulk tissue has the potential to avoid the

above weaknesses of the experimental approach. Furthermore, a bioinformatics approach can

make use of the large amount of existing RNAseq data[4-8,12] from fresh or frozen bulk brain

tissues with minimum added cost, and may better reflect the true situation in brain where

different cells are in communication.

In this study, applying three different algorithms[13,14] including our own novel approach,

we estimated cell-intrinsic gene expression for deconvoluted cell types from three large bulk

RNAseq datasets[4,8,12,18] from two brain regions. We identified consensus cell-intrinsic

transcriptional alterations (CI-DEGs) in AD, which are conserved across cohorts and brain

regions. We also performed an in-depth comparison of these CI-DEGs with bulk brain RNAseq

data obtained from the same datasets, collectively comprised of 685 unique brain samples. To

our knowledge, this is the first study to detect CI-DEGs and their enriched gene ontology (GO)

terms in computationally deconvoluted large-scale RNAseq data from AD and control brain

samples. Additionally, we conducted a detailed comparison of CI-DEGs deconvoluted from

bulk-DLPFC data with the three algorithms and those obtained from snRNAseq[19] (snDLPFC) of

a subset of the samples from the same cohort[7,8].

354   The main findings from our study can be summarized as follows: **1)** The direction of

change in cellular proportions in AD is consistent across two brain regions and three datasets for

most cell types, although the magnitude of change seems to vary. Our findings revealed greater

neuronal loss and microgliosis in TCX compared to DLPFC. **2)** We identified CI-DEGs and bulk

tissue DEGs (bulk-DEGs) independently in two TCX and one DLPFC datasets. The overlap in

bulk-DEGs across datasets is greater than that for CI-DEGs. **3)** We performed an in-depth

comparison of the consensus CI-DEGs, common to both TCX and DLPFC against the consensus

bulk-DEGs detected in these same datasets. We identified significant overlap between consensus

CI-DEGs and consensus bulk-DEGs. The extent of overlap between consensus bulk-DEGs and

consensus CI-DEGs was greatest for *down-regulated neuronal* genes (p=5.6E-207). This was

followed by *up-regulated non-neuronal* genes (p ranging from 1.4E-77 for endothelia to 4.3E-14

for microglia). **4)** Despite the statistically significant overlap between consensus bulk vs. CI-

DEGs, the majority of the consensus CI-DEGs for *up-regulated neuronal* and *down-regulated*

non-neuronal genes were not detected in bulk tissue. This finding highlights the potential ability

of computational deconvolution approach to identify CI-DEGs that may be missed in bulk-DEGs

especially for genes that are not moving in the direction of cell proportion changes. **5)** We

identified GO-terms enriched for consensus CI-DEGs, and detected processes that have

previously been implicated in AD as well as novel ones. **6**) Using an snRNAseq[19] dataset as

comparison, we assessed the performance of our CI-DEG detection algorithm (WLC), and the

published CellCODE[14] and PSEA[13] approaches. We determined that WLC had comparable or

superior performance in the detection CI-DEGs that had significant overlap with snDLPFC

results.

Our findings highlight the consistency and reproducibility of our findings across two

different brain regions from three different studies conducted separately. We identified similar

directions of change in AD:Control cell proportions in TCX and DLPFC. As expected from

known AD neuropathology, neuronal populations are significantly lower, and microglial

populations are significantly higher in AD vs. control brains in all datasets. Consistent with this

pattern of reproducibility, we also found significant overlap of consensus CI-DEGs detected in

TCX and DLPFC for all cell types and for both directions of change, i.e. up or down in AD, with

consensus bulk-DEGs.

Using consensus CI-DEGs, we identified GO terms, which include processes and genes

that have previously been implicated in AD, thereby providing further validation of our

approach. Detailed discussion of all of the pathways identified with the consensus CI-DEGs is

beyond the scope of this study. Instead, we herein highlight some of the CI-DEG enriched

pathways.

Down-regulated neuronal CI-DEGs were enriched in neuropeptide hormone activity

(GO:0005184) pathway. These terms include *VGF* (a.k.a. neuroendocrine regulatory peptide 1),

which is selectively expressed in some neurons and shown to be reduced in AD and Parkinson's

disease[22]. Corticotropin releasing hormone (CRH), which is a neuronally expressed peptide that

mediates stress in the hypothalamic-pituitary-adrenal axis, is also a member of the same GO

term. CRH has been implicated in both adverse outcomes related to AD pathology in model

systems and epidemiology studies, as well as having an important role in learning and memory[23].

Neuronal reduction of *CRH* may either be a potentially neuro-protective response in AD brains

or lead to further negative impact on memory. Although the biological implications of our

finding remain to be uncovered, our results are aligned with the potential importance of the

neuroendocrine system in AD.

Interestingly, CI-DEGs also implicate biological processes that are enriched for neuronal-

DEGs that are *up* in AD, despite reductions in neuronal cell populations in this condition for both

TCX and DLPFC. This suggests that our cell type specific transcriptome deconvolution

successfully captures transcript changes that occur in the direction opposite to that of cell-

population changes, and that may therefore be missed in bulk-DEG approaches.

Indeed, the significant GO terms potassium channel activity (GO:0005267) and

regulation of ion transport (GO:0043269) harbor many potassium channels, which are up in AD

for neuronal CI-DEGs in both TCX and DLPFC, but do not have consistent results in bulk-DEGs

from the same cohorts. These findings highlight the potential utility of cell-specific transcript

deconvolution approaches in reducing noise from cell population changes, thereby resulting in

consistent detection of true signal. Notably, potassium channels have been reported to be up in

AD brains and mouse models of AD[49-51], leading to their suggestion as a potential therapeutic

avenue in this condition.

Some consensus CI-DEGs point to AD-related perturbations of key cellular functions for

the specific cell type. An example of this is consensus oligodendrocyte CI-DEGs. The *down-*

regulated oligodendrocyte CI-DEGs are enriched for the myelination GO term (GO:0042552)

and those that are *up* in this cell type are enriched in ceramide biosynthetic process

(GO:0046513).

Down-regulated oligodendrocyte CI-DEGs include myelin basic protein (*MBP*)[4],

plasmolipin (*PLLP*)[4,5], myelin and lymphocyte protein (*MAL*), and myelin-associated

glycoprotein (*MAG*)[26], even though bulk-DEGs for these genes did not show consistent changes.

We[4] and others[5] demonstrated lower levels in AD of co-expression networks of genes implicated

in myelination, which is consistent with the present findings from oligodendrocyte CI-DEGs.

Ceramide dysregulation has been implicated in AD[52,53]. Increased ceramide species were

observed in AD and other neuropathological disorders compared to controls[53], and the activation

of the neutral sphingomyelinase–ceramide pathway induces oligodendrocyte death[54]. Our present

observation from oligodendrocyte CI-DEGs are consistent with these findings.

Despite the biological insights gained from computationally deconvoluted CI-DEGs, they

also have some shortcomings. Compared to bulk-DEGs, CI-DEGs between different datasets

show less degree of overlap, regardless of the deconvolution algorithm utilized. This highlights

the challenge in the field for the ultimate goal of minimizing detection of transcriptional

perturbations due to cell proportion changes while maximizing discovery of those that lead to

disease. Put differently, CI-DEGs may enhance discovery of true cell-specific transcriptional

changes but this may come at the expense of increased false negative findings. In contrast, bulk-

DEGs may capture a larger number of perturbed transcripts, but some may be merely due to cell

population differences between diseased and healthy tissue. Ultimately, findings from both

approaches may provide the greatest utility in detecting true positives.

Comparison of CI-DEGs deconvoluted from bulk-DLPFC and those detected using an

orthogonal approach of snRNAseq[19] from the same cohort (ROSMAP) demonstrates the ability

of these computational approaches to identify true cell-specific expression changes in AD. Using

our in-house WLC algorithm, there was significant overlap with the results from snRNAseq and

CI-DEGs of most cell types (**Fig 4, Tables S18-S19**). CellCODE and PSEA also identified

significant overlaps, but to a lesser extent, especially for rarer cell types such as microglia.

Hence, our WLC algorithm demonstrates at least comparable performance to these two

algorithms[13,14]. This is also supported by the higher degree of overlap among different cohorts

for CI-DEGs detected by WLC than the other two algorithms (**Fig 2a**). Due to the challenges in

deconvoluting noisy data from human series, different computational approaches may be utilized

and combined, and that calls for a more devoted effort in developing such algorithms.

In summary, using three distinct computational approaches, we deconvoluted brain bulk-

RNAseq data from three large and independent cohorts[8,12,18]. We detected cell population

changes that are observed consistently across cohorts, and congruent with the known disease

pathology. Although there is significant overlap between consensus CI-DEGs and consensus

bulk-DEGs, there are more unique CI-DEGs that change in the direction opposite to that of cell

population changes. This suggests that CI-DEGs may have utility in detecting disease-related

transcriptional changes above and beyond those due to cell proportion changes. Consensus CI-

DEGs identify GO terms, including those for hormone activity, myelin biology and channel

activity. The enriched CI-DEGs include genes previously implicated in AD or

neurodegeneration, such as *VGF, CRH, MOBP* and *MBP*, and other novel genes.

This study demonstrates the utility of our analytic approach in deciphering cell-specific

transcriptional alterations using bulk tissue in a complex disease, provides a comprehensive

comparison of our pipeline to existing ones, identifies patterns of cell proportions in AD and

control samples across brain regions, discovers novel CI-DEGs with replication across

independent cohorts and highlights biological processes with cell-specific expression changes in

AD. These findings are expected to refine discovery of molecular therapeutic targets, biomarkers

that reflect cellular transcriptional alterations in AD and accelerate generation of more accurate

disease models.

Methods

Analysis datasets

We generated the TCX-Mayo data, which consists of temporal cortex RNAseq

measurement of 80 AD patients and 28 controls diagnosed according to neuropathologic

criteria[4,12]. RNAseq data were processed and quality control (QC) was conducted as

described[4,12]. ROSMAP DLPFC[7,8] and TCX-MSBB[18] datasets were downloaded from the AMP-

AD Knowledge Portal on Synapse (syn8691134 and syn8691099). We further filtered out non-

Caucasian samples and those that had incongruent sex based on provided covariate vs.

transcriptome data. All samples were classified as AD or control based on neuropathological

data. All TCX-Mayo AD samples had Braak neurofibrillary tangle (NFT) score ≥ 4. TCX-Mayo

controls had Braak score ≤ 3 and were without any neurodegenerative disease diagnoses. TCX-

MSBB AD samples had Braak ≥ 4 and CERAD ≥ 2; and controls had Braak ≤ 3 and CERAD ≤ 1.

DLPFC AD samples form ROSMAP had Braak score ≥ 4 and CERAD neuritic plaque score ≤ 2.

ROSMAP control samples had Braak ≤ 3 and CERAD neuritic plaque score ≥ 3.

It should be noted that the CERAD[55] neuritic plaque score as applied by the ROSMAP

study is defined such that high CERAD indicates lower neuritic plaque burden and decreased

probability of AD. In the MSBB study, higher CERAD indicates higher plaque burden.

Raw RNA read counts were normalized using conditional quantile normalization (CQN)

method[56] implemented in R cqn package, as previously described[4]. This normalization takes into

consideration library size, gene length and GC content. It also performs a log2 transformation so

that the resulting distribution for each gene is Gaussian-like. We also determined covariates that

contributed significantly to the variation of gene expression in these RNAseq cohorts (**Fig S5**)

for adjustment in the analyses.

Cell proportion estimation

Digital sorting algorithm (DSA)[16] was applied to estimate cell proportions through R

DSA package, function DSA. For each cell type, DSA first computes the average marker gene

expression in the analysis dataset, the purpose of which is to construct a variable that better

reflects cell proportion variation among subjects. To reduce the effect of outlier expression that

is occasionally seen in RNAseq data, we modified the original DSA so that the median instead of

mean expression was computed.

CI-DEG analysis for individual cell types

In this study we identified CI-DEGs from deconvoluted bulk RNAseq data using three

different algorithms, namely PSEA[13], CellCODE[14] and our in-house algorithm WLC. All

analyses adjusted for the following variables: Sex, RIN, age at death and batch for DLPFC and

TCX-MSBB datasets, and sex, RIN and age at death for TCX-Mayo dataset (**Fig S5**).

PSEA[13] applies model selection procedures to select cell type(s) that should be included

in baseline (control) or AD condition, and then estimate differential expression in specific cell

types (CI-DEGs). We used the R package PSEA to obtain CI-DEG results of PSEA approach,

through functions em_quantvg (to generate candidate models) and lmfitst (to fit all candidate

models and pick the best one). Expression values used in PSEA are in linear scale (non log-

transformed).

CellCODE[14] assesses overall gene expression difference between AD and control groups

with adjustment of relative cell proportion, followed by assigning the cell type that correlates

best with the difference (CI-DEGs). R package CellCODE was used to obtain DEG results of

CellCODE approach, through functions getAllSPVs (to construct surrogate variable using

marker genes through singular value decomposition) and lm.coef (to estimate difference between

AD and control groups). Strictly speaking, CellCODE measures the overall differential

expression rather than CI-DEGs but identifies the cell type that is most correlated with the

observed difference between AD and control using cellPopT function. However, for simplicity,

we refer to the DEGs from CellCODE as CI-DEGs in this study. Expression values used in

CellCODE are in log scale.

Our in-house method WLC, described in the method section, applies <u>w</u>eighted <u>l</u>inear

regression with <u>c</u>onstraints and model selection procedures, which also estimates differential

expression in specific cell types (CI-DEGs). It guarantees the fitted relative gene expression is

non-negative. By weighing the expression, it smooths out the extreme data points. The

procedures of this algorithm is illustrated by the following high level pseudo code.

____

Assume cell type 1,2,3,4,5 are neu, oli, mic, ast and end respectively

Fit Eq.(1) to identify a set of cell type T in which the gene is significantly expressed

If the set T is not empty:

Fit Eq.(2) to identify a set of cell type $\Phi \subseteq T$ in which the gene is differentiallly expressed

Let $\Omega = \{each\ cell\ type\ in\ T, \Phi, T\ \}$

For each element $\Theta \in \Omega$

533   *Fit Eq.(3) with adjustment for other covariates C_k ($1 \le k \le m$)*

534   *Keep Akaike information criterion (AIC) of this model fitting*

535   *Identify Θ_{best} that gives the best AIC.*

536   *Use the estimated values from the best model*

$$y_i \sim \alpha_0 + \sum_{t \in \{1,2,3,4,5\}} \alpha_t x_{i,t} + \epsilon, \qquad \text{s.t. } \alpha_t \geq 0 \ (1 \leq t \leq 5) \qquad\qquad (1)$$

$$y_i \sim \beta_0 + \sum_{t \in T} \beta_t x_{i,t} + d_i \sum_{t \in T} \beta_t^\Delta x_{i,t} + \epsilon, \quad \text{s.t. } \beta_t \geq 0, \beta_t + \beta_t^\Delta \geq 0 \ (t \in T) \qquad (2)$$

$$y_i \sim \gamma_0 + \sum_{t \in T} \gamma_t x_{i,t} + d_i \sum_{t \in \Theta} \gamma_t^\Delta x_{i,t} + \sum_{k=1}^m \lambda_k C_{i,k} + \epsilon, \text{s.t. } \gamma_t \geq 0 (t \in T), \ \gamma_t + \gamma_t^\Delta \geq 0 \ (t \in \Theta) \ (3)$$

____

In Eqs.1-3, y_i is the observed expression of a gene in subject i; $x_{i,t}$ is the median marker gene

expression of cell type t in subject i; $C_{i,k}$ is covarite k in subject i. In Eq.1, α_t is the overall

relative expression in cell type t. In Eq.2, β_t is relative expression at the baseline condition in cell

type t; $d_i = 0$ if subject i is in control group, and $d_i = 1$ if subject i is in AD group; therefore,

β_t^Δ is the difference of relative gene expression between baseline condition and AD condition in

cell type t. Of note, due to the biological meaning these coefficents, they must satisfy constraints

such that α_t, β_t, $\beta_t + \beta_t^\Delta$ be non-negative. In addition, y_i is in linear scale rather than in log

scale[57], and these non-log-transformed expression values tend to have extreme data points that

need to be weighted down. Based on the above considerations, Eq. 1 was fitted by weighted least

square linear regression with contraints, which is implemented in lsei function in R package

limSolve. The weight of each observation (w_i) is determined by formulae Eq.3 and Eq.4 below.

Intuitively, if the expression of a gene in a sample is extremely distant from the median

expression of all samples in the same diagnosis group, the weight of that sample is smaller than 1

for that gene.

$$w_i = 1/(1 + \frac{|y_i - \text{median}(y_{AD})|}{1 + \text{median}(y_{AD})^2}) \qquad \text{sample } i \text{ is AD} \qquad (4)$$

$$w_i = 1/(1 + \frac{|y_i - \text{median}(y_{control})|}{1 + \text{median}(y_{control})^2}) \qquad \text{sample } i \text{ is control} \qquad (5)$$

GO enrichment analysis

Using genes included in the CI-DEG analysis as background genes, p-value for GO term enrichment with consensus CI-DEGs was calculated by "enrichmentAnalysis" function from WGCNA R package[35].

Comparison of CI-DEGs from computational deconvolution vs. snRNAseq:

To determine if computational bulk-tissue RNAseq could reveal true CI-DEGs, we downloaded and utilized a published snRNAseq study[19] from frozen DLPFC tissues (snDLPFC) which compared gene expression from 24 individuals with AD-pathology to that from 24 individuals without AD-pathology in six cell types - excitatory neurons, inhibitory neurons, oligodendrocyte, microglia, oligodendrocyte precursor cells and astrocytes. These snDLPFC samples are from the ROSMAP project, of which we analyzed 474 bulk-DLPFC RNAseq data in our current study. Twenty-four (9 AD cases, 15 controls) of the 48 individuals in snDLPFC study are also included in the bulk-DLPFC dataset. Hence both the snDLPFC and bulk-DLPFC are from the same cohort with significant overlap. Three deconvolution methods were included in this comparison – CellCODE[14], PSEA[13] and WLC, our in-house method.

Two types of comparisons were made between the deconvoluted bulk-DLPFC and snDLPFC results. In the first comparison, we ranked the genes by their p-values of differential expression between AD and control subjects, per cell type. We compared the top *N* up- or down-genes from the snDLPFC study to those identified by each deconvolution algorithm, per cell

type. Genes common to both the deconvoluted bulk-DLPFC and snDLPFC were counted and

plotted.

In the second set of comparisons, CI-DEGs identified from deconvolution methods at

nominal significance (p-value <0.05) were compared to those identified in the snRNAseq data

(p-value <0.05).

To assess if the observed overlap from each set of analyses is significant with regards to

overlap between random selections, we obtained empirical p values from computer simulations

described below.

Empirical p-value for the number of overlapping genes

The empirical p-values for the number of overlapping genes between snDLPFC and bulk-

DLPFC was obtained using a computer simulation as follows. (**A**) Let S_{sn} stand for all genes in

snDLFC and S_{bulk} for all genes in bulk-DLFC; (**B**) randomly assign up-regulation on each S_{sn}

gene at probability 0.5, and on each S_{bulk} gene at probability 0.5; (**C**) randomly pick N genes

from up-genes of S_{sn} , randomly pick N genes from up-genes of S_{bulk}, and count the number of

overlapping genes; (**D**) Steps B-C were repeated 10000 times, and the numbers of overlaps

$(M_1, M_2, ..., M_{10000})$ were obtained; (**E**) Let M be the number of observed overlapping genes, and

the empirical p-value is (1+number of occurrences that $M_i >= M$)/10001.

Declarations:

Ethics, consent and permission:

All data were generated from deceased individual's autopsied brain tissue. This study

was approved by Mayo Clinic Institutional Review Board.

Consent for publication:

All authors reviewed and approved the final manuscript.

Availability of Data and Materials:

The data used in this manuscript is available to the research community through the

AMP-AD knowledge portal on Sage Synapse as follows:

Mayo Clinic RNAseq: https://www.synapse.org/#!Synapse:syn8690799

ROSMAP RNAseq: https://www.synapse.org/#!Synapse:syn8691134

MSBB RNAseq: https://www.synapse.org/#!Synapse:syn8691099

Scripts to perform the analysis and results reported here will be made available upon

acceptance of the manuscript for publication.

Competing interests:

The authors declare that they have no competing interests.

Funding and Acknowledgements:

We thank the patients and their families for their participation, without them these studies

would not have been possible. The Mayo RNAseq study data was led by Dr. Nilüfer Ertekin-

Taner, Mayo Clinic, Jacksonville, FL as part of the multi-PI U01 AG046139 (MPIs Golde,

Ertekin-Taner, Younkin, Price) using samples from The Mayo Clinic Brain Bank. Data

collection was supported through funding by NIA grants P50 AG016574, R01 AG032990, U01

AG046139, R01 AG018023, U01 AG006576, U01 AG006786, R01 AG025711, R01

AG017216, R01 AG003949, NINDS grant R01 NS080820, CurePSP Foundation, and support

from Mayo Foundation. This study was in part funded by NIH RF1 AG051504 and R01

AG061796 (NET). We thank the Mayo Clinic Advanced Genomic Technology Center staff and

bioinformatics core facility for all gene expression measurements. XW thanks Dr. E. Aubrey

Thompson for text editing and Jeremy Burgess for provide insightful inputs. XW was funded in

part by a pilot grant from the Mayo Clinic ADRC (P50 AG016574).

The results published here are in whole or in part based on data obtained from the AMP-

AD Knowledge Portal (https://adknowledgeportal.synapse.org). Study data were provided by the

Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago. Data collection

was supported through funding by NIA grants P30AG10161 (ROS), R01AG15819 (ROSMAP;

genomics and RNAseq), R01AG17917 (MAP), R01AG30146, R01AG36042 (5hC methylation,

ATACseq), RC2AG036547 (H3K9Ac), R01AG36836 (RNAseq), R01AG48015 (monocyte

RNAseq) RF1AG57473 (single nucleus RNAseq), U01AG32984 (genomic and whole exome

sequencing), U01AG46152 (ROSMAP AMP-AD, targeted proteomics), U01AG46161(TMT

proteomics), U01AG61356 (whole genome sequencing, targeted proteomics, ROSMAP AMP-

AD), the Illinois Department of Public Health (ROSMAP), and the Translational Genomics

Research Institute (genomic).

637   The results published here are in whole or in part based on data obtained from the AMP-

AD Knowledge Portal (https://adknowledgeportal.synapse.org/). These data were generated from

postmortem brain tissue collected through the Mount Sinai VA Medical Center Brain Bank and

were provided by Dr. Eric Schadt from Mount Sinai School of Medicine.

Author contributions:

X.W., S.L., M.A. and N.E-T conceived the idea. X.W., M.A. and N.E-T developed and designed

the study, and wrote the manuscript. X.W. and S.L. developed the mathematical theory and X.W.

performed the computations. D.D. provided tissue from the Mayo Clinic Brain Bank and

neuropathologically characterized TCX-Mayo brain samples. T.N., K.M., S.L., M.A. and M.C.

performed sample selection and library preparation of the TCX-Mayo dataset. M.A., Z.Q., T.C.,

T.P. and J.R. performed quality control of DLPFC and TCX-MSBB datasets. J.C. and Y.A.

supervised the analytical aspects of the project. N.E-T. provided funding, supervision and

direction for the whole study. All authors provided critical feedback and approved the final

manuscript.

References:

1. 2018 Alzheimer's disease facts and figures. *Alzheimer's & Dementia* **14**, 367-429 (2018).
2. Mehta, D., Jackson, R., Paul, G., Shi, J. & Sabbagh, M. Why do trials for Alzheimer's disease drugs keep failing? A discontinued drug perspective for 2010-2015. *Expert opinion on investigational drugs* **26**, 735-739 (2017).
3. Anderson, R.M., Hadjichrysanthou, C., Evans, S. & Wong, M.M. Why do so many clinical trials of therapies for Alzheimer's disease fail? *The Lancet* **390**, 2327-2329 (2017).
4. Allen, M. *et al.* Conserved brain myelination networks are altered in Alzheimer's and other neurodegenerative diseases. *Alzheimer's & Dementia* (2017).
5. McKenzie, A.T. *et al.* Multiscale network modeling of oligodendrocytes reveals molecular components of myelin dysregulation in Alzheimer's disease. *Molecular Neurodegeneration* **12**, 82 (2017).
6. Zhang, B. *et al.* Integrated systems approach identifies genetic nodes and networks in late-onset Alzheimer's disease. *Cell* **153**, 707-720 (2013).
7. Mostafavi, S. *et al.* A molecular network of the aging human brain provides insights into the pathology and cognitive decline of Alzheimer's disease. *Nature Neuroscience* **21**, 811-819 (2018).
8. De Jager, P.L. *et al.* A multi-omic atlas of the human frontal cortex for aging and Alzheimer's disease research. *Scientific Data* **5**, 180142 (2018).
9. Zhang, Y. *et al.* Purification and Characterization of Progenitor and Mature Human Astrocytes Reveals Transcriptional and Functional Differences with Mouse. *Neuron* **89**, 37-53 (2016).
10. Darmanis, S. *et al.* A survey of human brain transcriptome diversity at the single cell level. *Proceedings of the National Academy of Sciences* **112**, 7285-7290 (2015).
11. Lake, B.B. *et al.* Integrative single-cell analysis of transcriptional and epigenetic states in the human adult brain. *Nature biotechnology* **36**, 70-80 (2018).
12. Allen, M. *et al.* Human whole genome genotype and transcriptome data for Alzheimer's and other neurodegenerative diseases. *Scientific Data* **3**, 160089 (2016).
13. Kuhn, A., Thu, D., Waldvogel, H.J., Faull, R.L.M. & Luthi-Carter, R. Population-specific expression analysis (PSEA) reveals molecular changes in diseased brain. *Nat Meth* **8**, 945-947 (2011).
14. Chikina, M., Zaslavsky, E. & Sealfon, S.C. CellCODE: a robust latent variable approach to differential expression analysis for heterogeneous cell populations. *Bioinformatics* **31**, 1584-1591 (2015).
15. McKenzie, A.T. *et al.* Brain Cell Type Specific Gene Expression and Co-expression Network Architectures. *Scientific Reports* **8**, 8868 (2018).
16. Zhong, Y., Wan, Y.-W., Pang, K., Chow, L.M. & Liu, Z. Digital sorting of complex tissues for cell type-specific gene expression profiles. *BMC Bioinformatics* **14**, 89 (2013).
17. Kuhn, A. *et al.* Cell population-specific expression analysis of human cerebellum. *BMC genomics* **13**, 610 (2012).
18. Wang, M. *et al.* The Mount Sinai cohort of large-scale genomic, transcriptomic and proteomic data in Alzheimer's disease. *Sci Data* **5**, 180185 (2018).
19. Mathys, H. *et al.* Single-cell transcriptomic analysis of Alzheimer's disease. *Nature* (2019).
20. Ashburner, M. *et al.* Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nature genetics* **25**, 25-29 (2000).
21. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics* **9**, 559 (2008).

22.   Cocco, C. *et al.* Distribution of VGF peptides in the human cortex and their selective changes in
Parkinson's and Alzheimer's diseases. *Journal of anatomy* **217**, 683-93 (2010).
23.   Futch, H.S., Croft, C.L., Truong, V.Q., Krause, E.G. & Golde, T.E. Targeting psychologic stress
signaling pathways in Alzheimer's disease. *Molecular neurodegeneration* **12**, 49 (2017).
24.   Wang, R. & Reddy, P.H. Role of Glutamate and NMDA Receptors in Alzheimer's Disease. *J*
Alzheimers Dis **57**, 1041-1048 (2017).
25.   Berchtold, N.C. *et al.* Brain gene expression patterns differentiate mild cognitive impairment
from normal aged and Alzheimer's disease. *Neurobiol Aging* **35**, 1961-72 (2014).
26.   McAleese, K.E. *et al.* Parietal white matter lesions in Alzheimer's disease are associated with
cortical neurodegenerative pathology, but not with small vessel disease. *Acta neuropathologica*
134, 459-473 (2017).
27.   Crivelli, S.M. *et al.* Sphingolipids in Alzheimer's disease, how can we target them? *Adv Drug Deliv*
Rev (2020).
28.   Olsen, A.S.B. & Faergeman, N.J. Sphingolipids: membrane microdomains in brain development,
function and neurological diseases. *Open Biol* **7**(2017).
29.   El Gaamouch, F. *et al.* VGF-derived peptide TLQP-21 modulates microglial function through
C3aR1 signaling pathways and reduces neuropathology in 5xFAD mice. *Molecular*
neurodegeneration **15**, 4-4 (2020).
30.   Tzeng, T.-C. *et al.* Inflammasome-derived cytokine IL18 suppresses amyloid-induced seizures in
Alzheimer-prone mice. *Proceedings of the National Academy of Sciences of the United States of*
America **115**, 9002-9007 (2018).
31.   White, C.S., Lawrence, C.B., Brough, D. & Rivers-Auty, J. Inflammasomes as therapeutic targets
for Alzheimer's disease. *Brain Pathol* **27**, 223-234 (2017).
32.   Gao, T. *et al.* Transcriptional regulation of homeostatic and disease-associated-microglial genes
by IRF1, LXRβ, and CEBPα. *Glia* **67**, 1958-1975 (2019).
33.   Deitmer, J.W., Theparambil, S.M., Ruminot, I., Noor, S.I. & Becker, H.M. Energy Dynamics in the
Brain: Contributions of Astrocytes to Metabolism and pH Homeostasis. *Frontiers in Neuroscience*
13(2019).
34.   Bélanger, M., Allaman, I. & Magistretti, Pierre J. Brain Energy Metabolism: Focus on Astrocyte-
Neuron Metabolic Cooperation. *Cell Metabolism* **14**, 724-738 (2011).
35.   Zhang, Q. *et al.* Integrated proteomics and network analysis identifies protein hubs and network
alterations in Alzheimer's disease. *Acta neuropathologica communications* **6**, 19-19 (2018).
36.   Lovell, M.A., Xie, C., Gabbita, S.P. & Markesbery, W.R. Decreased thioredoxin and increased
thioredoxin reductase levels in alzheimer's disease brain. *Free Radical Biology and Medicine* **28**,
418-427 (2000).
37.   Kajiwara, Y. *et al.* GJA1 (connexin43) is a key regulator of Alzheimer's disease pathogenesis. *Acta*
Neuropathol Commun **6**, 144 (2018).
38.   de Quervain, D.J. & Papassotiropoulos, A. Identification of a genetic cluster influencing memory
performance and hippocampal activity in humans. *Proc Natl Acad Sci U S A* **103**, 4270-4 (2006).
39.   Shi, Y. *et al.* Rapid endothelial cytoskeletal reorganization enables early blood–brain barrier
disruption and long-term ischaemic reperfusion brain injury. *Nature Communications* **7**, 10523
(2016).
40.   Stamatovic, S.M., Keep, R.F. & Andjelkovic, A.V. Brain endothelial cell-cell junctions: how to
"open" the blood brain barrier. *Current neuropharmacology* **6**, 179-192 (2008).
41.   Evans, H.T., Benetatos, J., van Roijen, M., Bodea, L.G. & Gotz, J. Decreased synthesis of
ribosomal proteins in tauopathy revealed by non-canonical amino acid labelling. *EMBO J* **38**,
e101174 (2019).

42.   Garcia-Esparcia, P. *et al.* Altered machinery of protein synthesis is region- and stage-dependent
and is associated with alpha-synuclein oligomers in Parkinson's disease. *Acta Neuropathol*
Commun **3**, 76 (2015).
43.   Koren, S.A. *et al.* Tau drives translational selectivity by interacting with ribosomal proteins. *Acta*
Neuropathol **137**, 571-583 (2019).
44.   Conway, O.J. *et al.* ABI3 and PLCG2 missense variants as risk factors for neurodegenerative
diseases in Caucasians and African Americans. *Molecular neurodegeneration* **13**, 53 (2018).
45.   Srinivasan, K. *et al.* Untangling the brain's neuroinflammatory and neurodegenerative
transcriptional responses. *Nature communications* **7**, 11295 (2016).
46.   Lin, M.-Y. *et al.* Releasing Syntaphilin Removes Stressed Mitochondria from Axons Independent
of Mitophagy under Pathophysiological Conditions. *Neuron* **94**, 595-610.e6 (2017).
47.   Newberg, L.A., Chen, X., Kodira, C.D. & Zavodszky, M.I. Computational de novo discovery of
distinguishing genes for biological processes and cell types in complex tissues. *PloS one* **13**,
e0193067 (2018).
48.   Li, Z. *et al.* Genetic variants associated with Alzheimer's disease confer different cerebral cortex
cell-type population structure. *Genome medicine* **10**, 43 (2018).
49.   Angulo, E. *et al.* Up-regulation of the Kv3.4 potassium channel subunit in early stages of
Alzheimer's disease. *J Neurochem* **91**, 547-57 (2004).
50.   Maezawa, I. *et al.* Kv1.3 inhibition as a potential microglia-targeted therapy for Alzheimer's
disease: preclinical proof of concept. *Brain* **141**, 596-612 (2018).
51.   Yi, M. *et al.* KCa3.1 constitutes a pharmacological target for astrogliosis associated with
Alzheimer's disease. *Mol Cell Neurosci* **76**, 21-32 (2016).
52.   Czubowicz, K., Jęśko, H., Wencel, P., Lukiw, W.J. & Strosznajder, R.P. The Role of Ceramide and
Sphingosine-1-Phosphate in Alzheimer's Disease and Other Neurodegenerative Disorders.
Molecular neurobiology **56**, 5436-5455 (2019).
53.   Filippov, V. *et al.* Increased ceramide in brains with Alzheimer's and other neurodegenerative
diseases. *Journal of Alzheimer's disease : JAD* **29**, 537-547 (2012).
54.   Lee, J.-T. *et al.* Amyloid-beta peptide induces oligodendrocyte death by activating the neutral
sphingomyelinase-ceramide pathway. *The Journal of cell biology* **164**, 123-131 (2004).
55.   Mirra, S.S. *et al.* The Consortium to Establish a Registry for Alzheimer's Disease (CERAD). Part II.
Standardization of the neuropathologic assessment of Alzheimer's disease. *Neurology* **41**, 479-
86 (1991).
56.   Hansen, K.D., Irizarry, R.A. & Wu, Z. Removing technical variability in RNA-seq data using
conditional quantile normalization. *Biostatistics* **13**, 204-216 (2012).
57.   Zhong, Y. & Liu, Z. Gene expression deconvolution in linear space. *Nat Meth* **9**, 8-9 (2012).

Figure S1: Pearson correlation of estimated cell proportions using different sets of marker genes randomly selected. The steps of this simulation study are as follows. **(1)** Obtain candidate markers from R package BRETIGEA[1] for neurons (top 500 out of 1000 listed), oligodendrocyte (top 500 out of 1000 listed), microglia (top 500 out of 1000 listed), OPC (top 250 out of 500 listed), astrocyte (top 500 out of 1000 listed) and endothelial (top 500 out of 1000 listed). **(2)** From candidate markers, randomly select 1/10 genes for each cell type, that is 50 selected genes for all cell types except OPC which has 25 selected genes. **(3)** Estimate cell proportion using DSA algorithm[2]. **(4)** Repeat (2)-(3) 100 times, and compute Pearson correlation of estimated cell proportion between different runs.

Figure S2: Illustration of obtaining consensus CI-DEGs between DLPFC and TCX regions, using up-regulated CI-DEGs in neuronal cells as an example.

Figure S3: Left panel: the number of up-regulated and down-regulated consensus bulk-DEGs between DLPFC and TCX-Mayo, or between DLPFC and TCX-MSSM. Right panel: percent of consensus CI-DEGs that are also cell type marker genes. Cell type markers are from BRETIGEA[1], containing 1000 markers for each of the five cell types.

Figure S4: Venn diagram for all consensus bulk-DEGs and consensus CI-DEGs.

Figure S5: Source of variance analysis of in the DLPFC RNAseq dataset (left panel), TCX-Mayo (middle panel) and TCX-MSBB (right panel). For each gene, a full model was fitted in which cqn normalized gene expression with gene expression as dependent variable and RIN, age at death, sex, batch, and diagnosis group as independent variables (for DLPFC); RIN, age at death, sex, flowcell, and diagnosis group as independent variables (for TCX-Mayo); RIN, age at death, sex, batch, and diagnosis group as independent variables (for TCX-MSBB). Partial models were fitted using the same dependent variable and all but one independent variable. F statistics were obtained by comparing the full model and partial model for each independent variable. Y-axis is the mean of values of F statistics over all genes. In DLPFC and TCX-MSBB, diagnosis, age at death, sex, RIN and batch contributed more than random errors to the variation of gene expression, whereas in TCX-Mayo diagnosis, age at death, sex, and RIN contributed more than random errors to the variation of gene expression.

References:

64   1.   McKenzie, A.T. *et al.* Brain Cell Type Specific Gene Expression and Co-expression Network
65   Architectures. *Scientific Reports* **8**, 8868 (2018).
66   2.   Zhong, Y., Wan, Y.-W., Pang, K., Chow, L.M. & Liu, Z. Digital sorting of complex tissues for
67   cell type-specific gene expression profiles. *BMC Bioinformatics* **14**, 89 (2013).