INSECT HERBIVORY ON URBAN TREES: COMPLEMENTARY EFFECTS OF TREE NEIGHBOURS AND PREDATION

Alex Stemmelen[1], Alain Paquette[2] *, Marie-Lise Benot[1], Yasmine Kadiri[1], Hervé Jactel[1] and Bastien Castagneyrol[1,*]

1 - BIOGECO, INRAE, Univ. Bordeaux, 33610 Cestas, France

2 – Département des sciences biologiques, Centre d'étude de la forêt (CEF), Université du Québec à Montréal, Centre-Ville Station, P.O. Box 8888, Montréal, Qc H3C 3P8, Canada

* Corresponding authors:

- Bastien Castagneyrol, INRAE UMR BIOGECO, 69 route d'Arcachon, FR-33612 Cestas Cedex France, bastien.castagneyrol@inrae.fr
- Alain Paquette, Département des sciences biologiques, Centre d'étude de la forêt (CEF), Université du Québec à Montréal, Centre-Ville Station, P.O. Box 8888, Montréal, Qc H3C 3P8, Canada, paquette.alain@uqam.ca.

Abstract

1. Insect herbivory is an important component of forest ecosystems functioning and can affect tree growth and survival. Tree diversity is known to influence insect herbivory in natural forest, with most studies reporting a decrease in herbivory with increasing tree diversity. Urban ecosystems, on the other hand, differ in many ways from the forest ecosystem and the drivers of insect herbivory in cities are still debated.

2. We monitored 48 urban trees from five species – three native and two exotic – in three parks of Montreal (Canada) for leaf insect herbivory and predator activity on artificial larvae, and linked herbivory with both predation and tree diversity in the vicinity of focal trees.

3. Leaf insect herbivory decreased with increasing tree diversity and with increasing predator attack rate.

4. Our findings indicate that tree diversity is a key determinant of multitrophic interactions between trees, herbivores and predators in urban environments and that managing tree diversity could contribute to pest control in cities.

Keywords : Artificial prey, Insect herbivory, Tree diversity, Top-down control, Urban biodiversity

Introduction

Insect herbivores have a major impact on tree growth and survival, henceforth on the functioning of forest ecosystems (Metcalfe et al., 2014; Visakorpi et al., 2018; Zvereva, Zverev, & Kozlov, 2012). Tree diversity significantly influences insect herbivory in forest ecosystems (Castagneyrol, Jactel, Vacher, Brockerhoff, & Koricheva, 2014; Jactel et al., 2017). Most studies reported negative relationship between tree diversity and herbivory (*i.e.*, associational resistance, Barbosa et al., 2009), although the opposite was also found (Haase et al., 2015; Schuldt et al., 2011). Recently, the interest about tree diversity effects on insect herbivory has expanded to include urban forests (Clem & Held, 2018; Dale & Frank, 2018; Frank, 2014), where pest damage can compromise the ecological and aesthetic values of urban trees (Nuckols & Connor, 1995; Tooker & Hanks, 2000; Tubby & Webber, 2010). Urban forests differ from natural forests in many ways. For example, most of the trees in cities are planted, found in lower density and/or mixed with native and exotic ornamental species, which are rarely encountered in natural forests. Henceforth, given those numerous peculiarities, it is still unclear how and why tree diversity might influence insect herbivory in urban trees.

By determining the amount and the quality of food and habitat resources available to herbivores and their enemies, both the density and diversity of trees can have strong impact on the bottom-up and top-down forces acting upon insect herbivores (Haase et al., 2015; Muiruri, Rainio, & Koricheva, 2016; Setiawan, Vanhellemont, Baeten, Dillen, & Verheyen, 2014). For example, some insect herbivores, in particular generalist species, could take advantage of tree diversity to acquire more abundant, complementary food resources or benefit from a more balanced food mix, thus causing more damage in mixed forests (Lefcheck, Whalen, Davenport, Stone, & Duffy, 2013). On the contrary, insect herbivores generally find it easier to identify and orientate towards the signals emitted by their host trees when the latter are more concentrated (*the resource concentration hypothesis*, Hambäck & Englund, 2005; Root, 1973) while non-host trees can emit volatile compounds that interfere with the ability of herbivores to detect their preferred host (Jactel, Birgersson, Andersson, & Schlyter, 2011). Finally, the abundance and diversity of predatory birds and arthropods generally increases with plant density and diversity, which would result in a better top-down regulation of insect herbivores (*the enemies hypothesis*, Risch, Andow, & Altieri, 1983; Root, 1973). However, the evidence available to support this hypothesis in forest is controversial (Muiruri et al., 2016; Riihimäki, Kaitaniemi, Koricheva, & Vehviläinen, 2005) and the contribution of natural enemies to the control of herbivores in urban area remains poorly explored.

Tree diversity and density vary widely between and within cities (Ortega-Álvarez, Rodríguez-Correa, & MacGregor-Fors, 2011; Sjöman, Östberg, & Bühler, 2012). A consequence of this variability is that even within a common urban environment, herbivory may be reduced in some tree species and increased in others (Clem & Held, 2018; Frank, 2014), and the relative importance of bottom-up and top-down forces responsible for these effects may also differ. In addition, non-native trees have been widely planted in urban habitats (Cowett & Bassuk, 2014; Moro, Westerkamp, & de Araújo, 2014). While they often escape from herbivory by native

insects ('*the enemy escape hypothesis*', Adams et al., 2009; Keane & Crawley, 2002), cases of
native herbivores spilling-over onto exotic trees have been recorded (e.g. Branco, Brockerhoff,
Castagneyrol, Orazio, & Jactel, 2015). Non-native tree species can also provide habitats to
predatory birds or arthropods (Gray & van Heezik, 2016). It is thus difficult to predict the effect
of mixing native and exotic trees on insect herbivory in urban habitats (Clem & Held, 2018;
Frank, 2014).

In this study, we investigated the effect of tree density, tree diversity, presence of conspecific
trees, tree origin and predator activity on insect herbivory in urban trees of the city of Montreal
(Quebec, Canada). We measured leaf area removed or damaged by insect herbivores on 48
trees of five species – three native and two exotic – in three urban parks. We concomitantly
assessed predator activity by using artificial caterpillars exposed on tree branches. We tested
the following hypotheses: (1) insect herbivory decreases with tree density, number of non-
conspecific trees (host dilution) and diversity (associational resistance) around focal trees, (2)
predator activity increases with increasing tree density and diversity and (3) predation and
herbivory have different responses to tree diversity on native and exotic trees. By doing so, our
study builds toward a better understanding of the drivers of pest insect damage on urban trees.

Materials and methods

Study site

96 The study was conducted in the city of Montreal (Canada, 45°50'N, -73°55'W), where the
97 climate is temperate cold, with 6.8°C average temperature and 1000.3 mm annual rainfall
98 during the 1981-2010 period (Pierre Elliott Trudeau airport weather station, www.canada.ca).
99 The study took place in three parks of the southwest part of the city: Angrignon, Marguerite
100 Bourgeoys and Ignace-Bourget (Table 1).

Tree selection

Every tree in Angrignon, Ignace-Bourget and Marguerite-Bourgeoys parks had been
previously geolocalized and identified to the species level. This information was accessible
through the city database for urban trees (http://donnees.ville.montreal.qc.ca/dataset/arbres).
We selected a total of 48 trees of five deciduous species (Table 1). Three species are native to
the study area (*Acer saccharinum* L.*, Tilia americana* L.*, Quercus rubra* L.) while two are
exotics, from Europe (*Acer platanoides* L.*, Tilia cordata* Mill.). These species are amongst the
most abundant tree species in the city of Montreal where together they represent 37% of all the
trees of the public domain. In agreement with the city of Montreal administration, we only
selected trees with a diameter at breast height (DBH) greater than 8 cm (mean ± SD: 18.38 ±
9.36) (to withstand the sampling of leaves required for the experiment) and with low branches
that could be easily accessed using a stepladder (for safety).

Table 1. Mean (± SD) diameter at breast height (in cm) and number of trees selected for each park and species.

Species	Angrignon (AN) (45°26'N, -73°36')	Marguerite-Bourgeoys (MB) (45°47'N, -73°36'W)	Ignace-Bourguet (IB) (45°45'N, -73°60'W)
Acer saccharinum	37.55 (n = 2)	37.55 (n = 2)	15.1 (n = 2)
Acer platanoides	21.60 (n = 1)	23.68 ± 1.97 (n = 6)	26.25 (n = 2)
Tilia cordata	22.40 (n = 1)	30.60 ± 3.37 (n = 5)	9.67 ± 0.51 (n = 4)
Tilia americana	10.52 ± 0.55 (n = 4)	22.06 ± 1.87 (n = 3)	27.60 ± 1.20 (n = 3)
Quercus rubra	8.96 ± 0.37 (n = 5)	*NA*	12.30 ± 1.45 (n = 5)

Predation rate assessment

We used artificial caterpillars made with modelling clay to estimate predation rate on sampled trees (Ferrante, Lo Cacciato, & Lovei, 2014; Howe, Lövei, & Nachman, 2009). We installed 15 artificial caterpillars per tree. We haphazardly selected three low (2.5-3.5 m from ground) branches facing opposite directions and installed five artificial caterpillars per branch (total: 720 caterpillars). Caterpillars were 3 cm long, and modelled to match the approximate form and size of real caterpillars. They were modelled using a 1-cm ball of non-toxic and odourless green modelling clay *(Sculpey III String Bean colour)* and secured on thin branches using a 12-cm long, 0.5 mm diameter, non-shiny metallic wire.

We exposed artificial caterpillars for 11 days in late spring (from May 29[th] to June 9[th], 2018) and for 6 days in early summer (from July 18[th] to July 24[th], 2018). These seasons were chosen to cover the main activity period of both predators and herbivores. Artificial caterpillars were left untouched for the full duration of each survey. We estimated total predator attack rate as the number of artificial larvae with any predation mark, divided by the total length of the observation period in days. There were uncertainties regarding predator identity responsible for predation marks. Most of the marks were attributable to birds or arthropods, while very few were attributable to small mammals, therefore, we chose to combine predation marks primarily attributed to birds or arthropods into a single category, which we refer to as total predation.

Branches of three trees were accidentally pruned by city workers in late spring so that the predation rate could not be estimated on these trees for the first survey. Three new trees of the same species were selected for the second survey, in early summer.

Leaf insect herbivory

We estimated background leaf insect herbivory (Kozlov et al., 2017) as the percentage of leaf area removed or impacted by insect herbivores. At the end of the second predation survey, we collected 10 leaves per branch on the same branches on which we had exposed artificial caterpillars, starting with the most apical, fully-developed, leaf to the 10th leaf down to branch basis (Total: 30 leaves per tree). We estimated total herbivory (i.e., total leaf area consumed or impacted by herbivores, regardless of their identity) as well as damage made by chewing, mining and sap-feeding herbivores at the level of individual leaves by using an ordinal scale of eight percentage classes of defoliation: 0%; 0-1%, 1-5%; 6-10%; 11-25%; 26-50%; 51-75% and 76-100%. We counted the number of galls per leaf. Most damage was made by leaf chewers, while other damage had a much skewed distribution, preventing detailed analyses for each type of damage separately. We therefore analysed total herbivory by averaging herbivory at the level of individual trees and using the median of each class of defoliation. Herbivory was scored by a single observer (BC), who was blind to tree identity.

Tree neighbourhood

We used three variables to describe tree neighbourhood in a 20-m radius around each focal tree: tree density (defined as the number of neighbouring trees in that radius), tree species diversity (Shannon diversity index) and the number of conspecific trees around each focal tree. Those variables were obtained using QGIS Geographic Information System software (QGIS Development Team, 2018). Excluding focal tree species, the most common tree species in the vicinity of focal trees were the smooth serviceberry (*Amelanchier leavis* Wiegand), the white spruce (*Picea glauca* Voss), the green ash (*Fraxinus pennsylvanica* Marshall) and the eastern cottonwood (*Populus deltoides* Marshall), all of them native to the region. We should note that, as focal trees were not necessarily 20m or more apart, we could not avoid that some "neighbour" trees were used in more than one neighbourhood, and some focal trees were also within the neighbourhood of another focal tree.

Statistical analyses

We used the information theory framework to identify the best model fitting our data and applied model averaging whenever necessary to estimate model coefficient parameters (Grueber, Nakagawa, Laws, & Jamieson, 2011). We first build a beyond optimal model including tree density (*Density*), tree diversity (*Diversity*), number of conspecifics (*Conspecific*), origin of the focal tree (*Origin*, native of exotic), park (*Park*), predation rate (*Predation*) as fixed factor and tree species identity (*Species*) as a random factor:

$$Y_{ij} \sim \beta_0 + \beta_1 \times Density_{ij} + \beta_2 \times Diversity_{ij} +$$
$$\beta_3 \times Conspecific_{ij} + \beta_4 \times Origin_{exotic,\ ij} + \beta_5 \times Park_{IB,\ ij} +$$
$$\beta_6 \times Park_{MB,\ ij} + \beta_6 \times Predation_{ij} +$$
$$\gamma_j + \varepsilon_{ij} \qquad\qquad (1)$$
$$\gamma \sim N(0;\ \sigma_{species}^2) \qquad \varepsilon \sim N(0;\ \sigma_e^2)$$

Where Y_{ij} is the herbivory on tree individual i in tree species j, β are model coefficient
parameters for fixed effects, γ_j is the random effect of tree species identity and ε the residuals.

To ease the interpretation of parameter estimates after model averaging, we standardized the
input variables using Gelman's approach (Gelman, 2008). We then applied a procedure of
model selection based on the Akaike's criterion corrected for small sample size (AICc) by
running every models nested within the beyond optimal model. As tree density and tree
diversity were correlated (Pearson's correlation: $r = 0.71$), we excluded all sub-models that
included these predictors together. We ranked all models based on difference in AICc between
each model and the top ranked model with the lowest AICc (ΔAICc). Models with a ΔAICc <
2 are generally considered equally supported by the data or not differentiable from the top
ranked model. Finally, we estimated model fit by calculating marginal (R²m) and conditional
(R²c) R² values, corresponding to variance explained by fixed effects only (R²m) and by fixed
and random effects (R²c) (Nakagawa & Schielzeth, 2013). When multiple models had a ΔAICc
< 2, we used a model averaging approach to build a consensus model including all variables
found in the set of best models. We considered that a given predictor had a significant effect if
its 95% confidence interval did not overlap zero. When only one model had a ΔAICc < 2, we
used it as the best model. We used a square-root transformation of insect herbivory to satisfy
model assumptions of normality and homogeneity of residuals.

We used the same approach to test the effect of tree neighbourhood on predation rate. We log-
transformed predation rate to satisfy model assumptions. Model equation (2) included the fixed
effect of sampling season (*Season*) and the random effect of tree identity (τ_k), nested within
tree species identity as an additional random factor accounting for repeated measurements of
the same individuals:

$\texttt{Y}_{ijk} \sim \beta_0 + \beta_1 \times \texttt{Density}_{ijk} + \beta_2 \times \texttt{Diversity}_{ijk} +$
$\qquad \beta_3 \times \texttt{Conspecific}_{ijk} + \beta_4 \times \texttt{Origin}_{exotic,\ ijk} + \beta_5 \times \texttt{Park}_{IB,\ ijk} +$
$\qquad \beta_6 \times \texttt{Park}_{MB,\ ijk} + \beta_6 \times \texttt{Season}_{summer,\ ijk} +$
$\qquad \gamma_j + \tau_{k|j} + \varepsilon_{ijk}$ (2)
$\qquad \gamma \sim N(0;\ \sigma_{species}^2) \qquad \tau \sim N(0;\ \sigma_{individual}^2) \qquad \varepsilon \sim N(0;\ \sigma_e^2)$

200
201 Statistical analyses were performed using the R software version 3.4.4 (R Core Team 2019)
202 with packages *lme4* (Bates, Mächler, Bolker, & Walker, 2015) and *MuMIn* (Barton 2019).

Results

204 ***Insect herbivory*** – Damaged leaf area was on average (± SE) 7.19 ± 0.70 % ($n = 48$). Leaf
205 damage was lower in *Acer platanoides* (3.53 ± 0.54) and *A. saccharinum* (3.86 ± 0.47) than in
206 *Quercus rubra* (8.77 ± 1.65), *Tilia americana* (10.3 ± 1.37) and *T. cordata* (8.75 ± 1.75) (Fig.
207 1A).
208

There were six models competing with the top ranked model in a range of 2 units of ΔAICc
(Table 2). These models included tree Shannon diversity, predation rate and tree origin as
predictors. Insect herbivory decreased significantly with increasing tree diversity (average
model coefficient parameter estimate ± CI: –0.482 ± [-0.91;-0.05], Fig. 2A, Table 3) and with
increasing predation rate (–0.473 ± [-0.91; -0.003]) (Fig. 2B, Table 3). Others predictors had
no significant effect on insect herbivory. Among the set of best models, fixed effects explained
between 7 and 12% of variability in insect herbivory. Fixed and random effects together
explained between 47 and 65% of variability in insect herbivory.

Figure 2. Effect of tree species identity on insect herbivory (A) and predation rate (B). Black dots and solid lines represents mean ± SE calculated on raw data. Herbivory is the percentage of leaf area removed or impacted by herbivores in early summer. Predation rate is the number of caterpillars attacked per day in late spring

Predation – Of the 1,315 artificial caterpillars that we installed, 198 displayed marks
unambiguously attributable to predators (*i.e.*, 15%). Predation rate varied between 0 and 0.87
caterpillars.day^{-1} (Fig. 1B).
Only one model had a ΔAICc < 2 and was then selected as the best model. Only Season was
included in the best model and had a significant effect on predation rate, which was two times
higher in late spring (mean ± CI: 0.44 ± [0.31, 0.58] caterpillars·day^{-1}) than in early summer
(0.20 ± [0.16, 0.24] caterpillars·day^{-1}). Others predictors had no significant effects on predation
rate. Season explained 56 % of variability in predation rate and, collectively, fixed and random
effects explained 59 % of variability in predation rate.

Table 2. Summary results of model selection of tree neighbourhood effect on herbivory rate: set of models with ΔAICc < 2. Only predictors that were present at least once in the set of best models are represented. R²m and R²c represent fixed and fixed *plus* random factor, respectively.

	Model covariates				Model selection			
Model	Intercept	Predation	Origin	Diversity	K	Log L	$\Delta AICc$	R²m (R²c)
1	2.53			-0.52	1	-46.44	0.00	0.09 (0.46)
2	2.52	-0.52		-0.44	2	-45.18	0.04	0.12 (0.58)
3	2.51	-0.51			1	-46.79	0.70	0.07 (0.56)
4	2.53	-0.44	0.171	-0.43	3	-44.64	1.67	0.12 (0.65)
5	2.53		0.078	-0.52	2	-46.07	1.82	0.08 (0.53)
6	2.53	-0.53	0.357		2	-46.12	1.92	0.08 (0.62)

Table 3. Summary results after model averaging: effects of each parameter presents on the set of best models on herbivory rate. Bold parameter are significant. Relative importance is a measure of the prevalence of each parameter in each model used in model averaging.

Parameter	Estimate	Adjusted SE	Confidence interval	Relative importance
(Intercept)	2.53	0.31	(1.91, 3.14)	
Diversity	-0.48	0.21	(-0.91, -0.05)	0.72
Predation	-0.47	0.22	(-0.91, -0.003)	0.64
Origin	0.19	0.71	(-1.20, 1.60)	0.31

Figure 2. Effects of tree diversity (A) and predation rate (B) on insect herbivory. Solid and dashed lines represent prediction and adjusted standard error of the average model respectively (Table 3). Herbivory is the percentage of leaf area removed or impacted by herbivores in early summer. Tree diversity is represented by Shannon's diversity index. Predation rate is the number of caterpillars attacked per day in late spring.

Discussion

We confirmed that tree diversity can influence insect herbivory on urban trees. Specifically, we found that insect herbivory decreased with increasing tree diversity (i.e. associational resistance). We also found a negative correlation between predator attack rate and insect herbivory. Although further analyses are needed to confirm this trend, our findings provide support to the view that increasing tree diversity can enhance regulation of insect herbivores by natural enemies in urban forests.

Our results are in line with several studies having reported reduced herbivory in trees surrounded by heterospecific neighbours (reviewed by Castagneyrol et al., 2014; Jactel et al., 2017). It also adds to the growing number of studies documenting diversity-resistance relationships in urban environments (Clem & Held 2018; Doherty, Meagher, & Dale 2019; Frank 2014). However, it conflicts with other results suggesting an increase in herbivore

abundance with increasing plant diversity and vegetation volume in urban environments (Mata
et al., 2017), although the relationship between herbivore abundance and actual herbivory is
not always positively correlated (Barbosa et al., 2009; Schueller, Paul, Payer, Schultze, &
Vikas, 2019). Tree diversity may have influenced the probability of focal trees being found and
colonized by herbivores. Theory predicts that specialist herbivores have greater difficulties
finding their host trees when they are surrounded by heterospecific neighbours (Castagneyrol
et al., 2014; H. Jactel, Brockerhoff, & Duelli, 2009). It is possible that non-host neighbours
disrupted the physical and chemical cues used by insect herbivores to locate their hosts
(Damien et al., 2016; H. Jactel et al., 2011; Zhang & Schlyter, 2004). However, and contrary
to our expectations, we did not find any significant effect of conspecific tree density on insect
herbivory, thus ruling out the resource concentration hypothesis in this particular case. Because
our study was by essence observational, we could not separate the effect of conspecific
neighbour density vs. heterospecific neighbour density. In the absence of data on the identity
of herbivores responsible for herbivory, further speculation would be hazardous.

Insect herbivory varied across tree species but did not differ between native and non-native
species, which contradicts the predictions of the enemy release hypothesis (Cincotta, Adams,
& Holzapfel, 2009; Meijer, Schilthuizen, Beukeboom, & Smit, 2016). One possible
explanation for this result could be that native herbivores spilled over exotic tree species from
neighbouring native tree species, as it was recorded in previous studies (Branco et al., 2015).
This would have been facilitated by the fact that exotic tree species (from Europe) had
congeneric species in Canada. It is also important to note that a large part of the variability in
leaf insect damage was attributable to the species on which leaf samples were collected. In
particular, both *Acer platanoides* and *A. saccharinum* were far less damaged than *Tilia cordata*,
T. americana and *Quercus rubra*. In a recent study in Michigan, Schueller et al., (2019) also
reported greater insect herbivory (and herbivore diversity) on *Quercus* species as compared to
Acer species, which is consistent with the view that plant species identity can drive arthropods
community and abundance on forest host trees (Burghardt, Tallamy, & Gregory Shriver, 2009;
Pearse & Hipp, 2009).

We found a significant negative correlation between predator attack rate and insect herbivory
measured later in the season. This finding suggests a potential relationship between herbivory
and predation in urban environments (Faeth, Warren, Shochat, & Marussich, 2005; Kozlov et
al., 2017 but see Long & Frank, 2020). However, we refrain from concluding that, in our
system, predation was the main driver of insect herbivory for several reasons. First, the effect
size of the herbivory-predation relationship was small, as was model R^2 (Table 3). Second,
concerns remain about how much predation on artificial preys is representative of actual
predation (Lövei & Ferrante, 2017; Rößler, Pröhl, & Lötters, 2018). In particular, artificial
caterpillars used to assess predation rate modelled lepidopteran-like leaf chewing caterpillars
and thus, caution is needed when it comes to extrapolate predator attack rates to other herbivore
feeding guilds. Third, we had no information on actual natural prey density in focal and
neighbouring trees. Yet, prey availability may have influenced the functional response of bird
insectivores (e.g. optimal foraging) such that we cannot exclude that herbivory actually drove
predation rate instead of the other way around. Finally, the putative effect of predation on

herbivory may be weak in respect to other factors acting directly upon herbivores in urban
environments such as drought (Huberty & Denno, 2004; Mattson, 1980; Meineke & Frank,
2018), extreme heat (Dale & Frank, 2014; Meineke, Dunn, Sexton, & Frank, 2013) and
pollution leading to altered foliage quality (Kozlov et al., 2017; Mattson, 1980; Moreira et al.,
2019).

Contrary to tree species identity effect on insect herbivory, tree species had no clear influence
on predation rate on dummy caterpillars, which contradicts the view that tree species identity
can modulate attack rates of caterpillars by birds (Mooney & Singer, 2012; Nell et al., 2018).
Variation in predator density between plants is often related to an indirect effect of the plant on
the density (Bailey et al. 2006) or quality (Brower et al. 1967, Clancy and Price, 1987) of their
preys (herbivores). However, such effect of plant identity is not relevant when using dummy
caterpillars, as neither their abundance nor their quality can be affected by plant species
identity, which could explain the contradiction between past results and our study.

Predation was greater during the first survey, in late spring, than during the second survey, in
early summer. This result could be explained either by a lower foliage density in trees in spring,
making it easier for predators to detect artificial caterpillars, or by greater predator activity
matching the phenology of wild caterpillars and feeding period of chicks (Coley, 1980; Raupp
& Denno, 1983). We cannot either exclude that birds have learnt how to avoid artificial
caterpillars, thus resulting in much lower predation pressure during the second survey.
However, unless bird ability to avoid artificial caterpillar varied between tree species and
neighbourhood, we do not see this possibility as a major threat to our inferences.

Conclusion

Our study highlights several ecological factors driving leaf insect herbivory in the urban trees
of the Montreal city. In particular, we found that insect herbivory decreased with increasing
tree diversity and was probably affected by predator activity. While biological invasions and
global warming are increasing the risk upon urban trees, more and more cities choose to ban
or reduce the use of pesticides in urban parks and green areas (Sustainable Use of Pesticides
Directive 2009), such as in Montreal. In this context, diversifying urban tree cover in urban
parks might help to reduce insect damage, which could result in a better provision of services
provided by trees in cities (Beyer et al., 2014; Bowler, Buyung-Ali, Knight, & Pullin, 2010;
Nowak, Hirabayashi, Bodine, & Greenfield, 2014).

Authors' contributions

312 BC and AP conceived the study. YK selected trees and coordinated field work. YK and BC
313 acquired data. AS, MLB and BC analysed the data. AS, MLB and BC wrote the first draft.
314 All authors contributed critically to the final manuscript.

Acknowledgements

This study has been carried out with financial support from (i) the French National Research Agency (ANR) in the frame of the Investments for the future Programme to BC, within the Cluster of Excellence COTE (ANR-10-LABX-45), (ii) the Conseil Franco-Québécois de Coopération Universitaire (FRQNT – Samuel-de-Champlain fund) to BC and AP, and (iii) a Discovery grant to AP from the Natural Sciences and Engineering Research Council of Canada. We thank Charles Desroches, Elyssa Cameron, Christian Messier, summer interns at UQAM, and the city of Montréal, Arrondissement du Sud-Ouest for their help with field work. We also thank Frédéric Barraquand and Benjamin Brachi for their constructive comments on the preliminary version of the paper. We thank Luc Barbaro and Steve Frank for their friendly reviews early drafts of this paper. Finally, we are grateful to Ian Pearse and Freerk Molleman for their reviews of prior manuscript versions.

Data accessibility

Data and script available from the Data INRAE repository: https://doi.org/10.15454/R4NESA

Conflict of interest disclosure

The authors of this article declare that they have no financial conflict of interest with the content of this article. Bastien Castagneyrol is one of the PCI Ecology recommenders.

References

Adams, J. M., Fang, W., Callaway, R. M., Cipollini, D., Newell, E., & Transatlantic Acer platanoides Invasion Network (TRAIN). (2009). A cross-continental test of the Enemy Release Hypothesis: leaf herbivory on Acer platanoides (L.) is three times lower in North America than in its native Europe. *Biological Invasions*, *11*(4), 1005–1016. doi: 10.1007/s10530-008-9312-4

Akaike, H. 1973. Information theory as an extension of the maximum likelihood principle. *In: Second International Symposium on Information Theory (B.N. Petrov & F. Csaki, eds), pp. 267–281. Akademiai Kiado, Budapest.*

Barbosa, P., Hines, J., Kaplan, I., Martinson, H., Szczepaniec, A., & Szendrei, Z. (2009). Associational Resistance and Associational Susceptibility: Having Right or Wrong Neighbors. *Annual Review of Ecology, Evolution, and Systematics*, *40*(1), 1–20. doi: 10.1146/annurev.ecolsys.110308.120242

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, *67*(1). doi: 10.18637/jss.v067.i01

Beyer, K. M. M., Kaltenbach, A., Szabo, A., Bogar, S., Nieto, F. J., & Malecki, K. M. (2014). Exposure to Neighborhood Green Space and Mental Health: Evidence from the Survey of the Health of Wisconsin. *International Journal of Environmental Research and Public Health*, *11*(3), 3453–3472. doi: 10.3390/ijerph110303453

Bowler, D. E., Buyung-Ali, L. M., Knight, T. M., & Pullin, A. S. (2010). A systematic review of evidence for the added benefits to health of exposure to natural environments. *BMC Public Health*, *10*, 456. doi: 10.1186/1471-2458-10-456

Castagneyrol, B., Jactel, H., Vacher, C., Brockerhoff, E. G., & Koricheva, J. (2014). Effects of plant phylogenetic diversity on herbivory depend on herbivore specialization. *Journal of Applied Ecology*, *51*(1), 134–141. doi: 10.1111/1365-2664.12175

Cincotta, C. L., Adams, J. M., & Holzapfel, C. (2009). Testing the enemy release hypothesis: a comparison of foliar insect herbivory of the exotic Norway maple (Acer platanoides L.) and the native sugar maple (A. saccharum L.). *Biological Invasions*, *11*(2), 379–388. doi: 10.1007/s10530-008-9255-9

Clem, C. S., & Held, D. W. (2018). Associational Interactions Between Urban Trees: Are Native Neighbors Better Than Non-Natives? *Environmental Entomology*, *47*(4), 881–889. doi: 10.1093/ee/nvy071

Coley, P. D. (1980). Effects of leaf age and plant life history patterns on herbivory. *Nature*, *284*, 545–546. doi: 10.1038/284545a0

Cowett, F. D., & Bassuk, N. L. (2014). Statewide assessment of street trees in New York State, USA. *Urban Forestry & Urban Greening*, *13*(2), 213–220. doi: 10.1016/j.ufug.2014.02.001

Dale, A. G., & Frank, S. D. (2018). Urban plants and climate drive unique arthropod interactions with unpredictable consequences. *Current Opinion in Insect Science*, *29*, 27–33. doi: 10.1016/j.cois.2018.06.001

Dale, A. G., & Frank, S. D. (2014). Urban warming trumps natural enemy regulation of herbivorous pests. *Ecological Applications: A Publication of the Ecological Society of America*, *24*(7), 1596–1607. doi: 10.1890/13-1961.1

Damien, M., Jactel, H., Meredieu, C., Régolini, M., van Halder, I., & Castagneyrol, B. (2016) Pest damage in mixed forests: Disentangling the effects of neighbor identity, host density and host apparency at different spatial scales. *Forest Ecology and Management*, *378*, 103–110. doi: 10.1016/j.foreco.2016.07.025

Doherty, E. M., Meagher, R. L., & Dale, A. G. (2019). Turfgrass Cultivar Diversity Provides

Associational Resistance in the Absence of Pest Resistant Cultivars. *Environmental Entomology*, *48*(3), 623–632. doi: 10.1093/ee/nvz026

Faeth, S. H., Warren, P. S., Shochat, E., & Marussich, W. A. (2005). Trophic Dynamics in Urban Communities. *BioScience*, *55*(5), 399–407. doi: 10.1641/0006-3568(2005)055[0399:TDIUC]2.0.CO;2

Ferrante, M., Lo Cacciato, A., & Lövei, G. (2014). Quantifying predation pressure along an urbanisation gradient in Denmark using artificial caterpillars. *European Journal of Entomology*, *111*(5), 649–654.

Frank, Steven D. (2014). Bad neighbors: urban habitats increase cankerworm damage to non-host understory plants. *Urban Ecosystems*, *17*(4), 1135–1145. doi: 10.1007/s11252-014-0368-x

Gelman, A. (2008). Scaling regression inputs by dividing by two standard deviations. *Statistics in Medicine*, *27*(15), 2865–2873. doi: 10.1002/sim.3107

Grueber, C. E., Nakagawa, S., Laws, R. J., & Jamieson, I. G. (2011). Multimodel inference in ecology and evolution: challenges and solutions. *Journal of Evolutionary Biology*, *24*(4), 699–711. doi: 10.1111/j.1420-9101.2010.02210.x

Hambäck, P. A., & Englund, G. (2005). Patch area, population density and the scaling of migration rates: the resource concentration hypothesis revisited. *Ecology Letters*, *8*(10), 1057–1065. doi: 10.1111/j.1461-0248.2005.00811.x

Howe, A., Lövei, G. L., & Nachman, G. (2009). Dummy caterpillars as a simple method to assess predation rates on invertebrates in a tropical agroecosystem. *Entomologia Experimentalis et Applicata*, *131*(3), 325–329. doi: 10.1111/j.1570-7458.2009.00860.x

Huberty, A. F., & Denno, R. F. (2004). Plant Water Stress and Its Consequences for Herbivorous Insects: A New Synthesis. *Ecology*, *85*(5), 1383–1398.

Jactel, H., Birgersson, G., Andersson, S., & Schlyter, F. (2011). Non-host volatiles mediate associational resistance to the pine processionary moth. *Oecologia*, *166*(3), 703–711. doi: 10.1007/s00442-011-1918-z

Jactel, H., Brockerhoff, E., & Duelli, P. (2009). A Test of the Biodiversity-Stability Theory: Meta-analysis of Tree Species Diversity Effects on Insect Pest Infestations, and Re-examination of Responsible Factors. In M. Scherer-Lorenzen, C. Körner, & E.-D. Schulze (Eds.), *Forest Diversity and Function: Temperate and Boreal Systems* (pp. 235–262). doi: 10.1007/3-540-26599-6_12

Jactel, H., Bauhus, J., Boberg, J., Bonal, D., Castagneyrol, B., Gardiner, B., … Brockerhoff, E. G. (2017). Tree Diversity Drives Forest Stand Resistance to Natural Disturbances. *Current Forestry Reports*, *3*(3), 223–243. doi: 10.1007/s40725-017-0064-1

Keane, R., & Crawley, M. J. (2002). Exotic plant invasions and the enemy release hypothesis. *Trends in Ecology & Evolution*, *17*(4), 164–170. doi: 10.1016/S0169-5347(02)02499-0

Kozlov, M. V., Lanta, V., Zverev, V., Rainio, K., Kunavin, M. A., & Zvereva, E. L. (2017). Decreased losses of woody plant foliage to insects in large urban areas are explained by bird predation. *Global Change Biology*, *23*(10), 4354–4364. doi: 10.1111/gcb.13692

Kozlov, M. V., Zverev, V., & Zvereva, E. L. (2017). Combined effects of environmental disturbance and climate warming on insect herbivory in mountain birch in subarctic forests: Results of 26-year monitoring. *Science of The Total Environment*, *601–602*, 802–811. doi: 10.1016/j.scitotenv.2017.05.230

Lefcheck, J. S., Whalen, M. A., Davenport, T. M., Stone, J. P., & Duffy, J. E. (2013). Physiological effects of diet mixing on consumer fitness: a meta-analysis. Ecology, 94(3), 565-572.

Long, L. C., & Frank, S. D. (2020). Risk of bird predation and defoliating insect abundance are greater in urban forest fragments than street trees. *Urban Ecosystems*. doi: 10.1007/s11252-020-00939-x

Lövei, G. L., & Ferrante, M. (2017). A review of the sentinel prey method as a way of quantifying invertebrate predation under field conditions. *Insect Science*, *24*(4), 528–542. doi: 10.1111/1744-7917.12405

Mata, L., Threlfall, C. G., Williams, N. S. G., Hahs, A. K., Malipatil, M., Stork, N. E., & Livesley, S. J. (2017). Conserving herbivorous and predatory insects in urban green spaces. *Scientific Reports*, *7*. doi: 10.1038/srep40970

Mattson, W. J. (1980). Herbivory in Relation to Plant Nitrogen Content. *Annual Review of Ecology and Systematics*, *11*(1), 119–161. doi: 10.1146/annurev.es.11.110180.001003

Meijer, K., Schilthuizen, M., Beukeboom, L., & Smit, C. (2016). A review and meta-analysis of the enemy release hypothesis in plant-herbivorous insect systems. *PeerJ*, *4*, e2778. doi: 10.7717/peerj.2778

Meineke, E. K., Dunn, R. R., Sexton, J. O., & Frank, S. D. (2013). Urban warming drives insect pest abundance on street trees. *PloS One*, *8*(3), e59687. doi: 10.1371/journal.pone.0059687

Meineke, E. K., & Frank, S. D. (2018). Water availability drives urban tree growth responses to herbivory and warming. *Journal of Applied Ecology*, *55*(4), 1701–1713. doi:10.1111/1365-2664.13130

Metcalfe, D. B., Asner, G. P., Martin, R. E., Espejo, J. E. S., Huasco, W. H., Amézquita, F. F. F., … Malhi, Y. (2014). Herbivory makes major contributions to ecosystem carbon and nutrient cycling in tropical forests. *Ecology Letters*, *17*(3), 324–332. doi: 10.1111/ele.12233

Mooney, K.A. & Singer, M.S. (2012). Plant effects on herbivore-enemyinteractions in natural systems. InTrait-Mediated Indirect Interactions:Ecological and Evolutionary Perspectives (eds Ohgushi, T., Schmitz, O.,Holt, R.D.). Cambridge University Press, Cambridge, pp. 107–130.

Moreira, X., Abdala-Roberts, L., Teran, J. C. B. M. y, Covelo, F., Mata, R. de la, Francisco, M., Tack, A. J. M. (2019). Impacts of urbanization on insect herbivory and plant defences in oak trees. *Oikos*, *128*(1), 113–123. doi: 10.1111/oik.05497

Moro, M. F., Westerkamp, C., & de Araújo, F. S. (2014). How much importance is given to native plants in cities' treescape? A case study in Fortaleza, Brazil. *Urban Forestry & Urban Greening*, *13*(2), 365–374. doi: 10.1016/j.ufug.2014.01.005

Muiruri, E. W., Rainio, K., & Koricheva, J. (2016). Do birds see the forest for the trees? Scale-dependent effects of tree diversity on avian predation of artificial larvae. *Oecologia*, *180*(3), 619–630. doi: 10.1007/s00442-015-3391-6

Nakagawa, S., & Schielzeth, H. (2013). A general and simple method for obtaining R2 from generalized linear mixed-effects models. *Methods in Ecology and Evolution*, *4*(2), 133–142. doi: 10.1111/j.2041-210x.2012.00261.x

Nell, C., Abdala-Roberts, L., Parra-Tabla, V. & Mooney, K. (2018).Tropical tree diversity mediates foraging and predatory effects ofinsectivorous birds.Proc. R Soc. B, 285, 20181842

Nowak, D. J., Hirabayashi, S., Bodine, A., & Greenfield, E. (2014). Tree and forest effects on air quality and human health in the United States. *Environmental Pollution (Barking, Essex:1987)*, *193*, 119–129. doi: 10.1016/j.envpol.2014.05.028

Nuckols, M. S., & Connor, E. F. (1995). Do trees in urban or ornamental plantings receive more damage by insects than trees in natural forests? *Ecological Entomology*, *20*(3), 253–260. doi: 10.1111/j.1365-2311.1995.tb00455.x

Ortega-Álvarez, R., Rodríguez-Correa, H. A., & MacGregor-Fors, I. (2011). Trees and the City: Diversity and Composition along a Neotropical Gradient of Urbanization. *International Journal of Ecology*, doi: https://doi.org/10.1155/2011/704084

Pearse, I. S., & Hipp, A. L. (2009). Phylogenetic and trait similarity to a native species

predict herbivory on non-native oaks. *Proceedings of the National Academy of Sciences*,
106(43), 18097–18102. doi: 10.1073/pnas.0904867106

QGIS Development Team (2019). QGIS Geographic Information System. Open Source
Geospatial Foundation Project.

R Core Team (2019). R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Raupp, & Denno. (1983). Leaf age as a predictor of herbivore distribution and abundance
[Herbivorous insects including the willow leaf beetle]. In *Academic Press. Variables*
Plants and Herbivores in Natural and Managed Systems.

Revision of World Urbanization Prospects | Multimedia Library - United Nations
Department of Economic and Social Affairs Retrieved May 30, 2019, from
https://www.un.org/development/desa/publications/2018-revision-of-world-urbanization-
prospects.html

Riihimäki, J., Kaitaniemi, P., Koricheva, J., & Vehviläinen, H. (2005). Testing the enemies
hypothesis in forest stands: the important role of tree species composition. *Oecologia*,
142(1), 90–97. doi: 10.1007/s00442-004-1696-y

Root, R. B. (1973). Organization of a Plant-Arthropod Association in Simple and Diverse
Habitats: The Fauna of Collards (Brassica Oleracea). *Ecological Monographs*, *43*(1),
95–124. doi: 10.2307/1942161

Sjöman, H., Östberg, J., & Bühler, O. (2012). Diversity and distribution of the urban tree
population in ten major Nordic cities. *Urban Forestry & Urban Greening*, *11*(1), 31–39.
doi: 10.1016/j.ufug.2011.09.004

Sustainable use of pesticides. Retrieved May 30, 2019, from
https://ec.europa.eu/food/plant/pesticides/sustainable_use_pesticides_en

Tubby, K. V., & Webber, J. F. (2010). Pests and diseases threatening urban trees under a
changing climate. *Forestry: An International Journal of Forest Research*, *83*(4), 451–
459. doi: 10.1093/forestry/cpq027

Visakorpi, K., Gripenberg, S., Malhi, Y., Bolas, C., Oliveras, I., Harris, N., … Riutta, T.
(2018). Small-scale indirect plant responses to insect herbivory could have major impacts
on canopy photosynthesis and isoprene emission. *The New Phytologist*, *220*(3), 799–810.
doi: 10.1111/nph.15338

Zhang, Q.-H., & Schlyter, F. (2004). Olfactory recognition and behavioural avoidance of
angiosperm nonhost volatiles by conifer-inhabiting bark beetles. *Agricultural and Forest*
Entomology, *6*(1), 1–20. doi: 10.1111/j.1461-9555.2004.00202.x

Zvereva, E. L., Zverev, V., & Kozlov, M. V. (2012). Little strokes fell great oaks: minor but
chronic herbivory substantially reduces birch growth. *Oikos*, *121*(12), 2036–2043. doi:
10.1111/j.1600-0706.2012.20688.x