Genome-wide identification, characterization, and expression analysis of the monovalent cation-proton antiporter superfamily in maize, and functional analysis of its role in salt tolerance

Mengsi Kong[1,2†], Meijie Luo[2†], Jingna Li[2], Zhen Feng[2,3], Yunxia Zhang[2], Wei Song[2], Ruyang Zhang[2], Ronghuan Wang[2], Yuandong Wang[2], Jiuran Zhao[2*], Yongsheng Tao[1*], Yanxin Zhao[2*]

[1] College of Agronomy, Hebei Agricultural University, Baoding 071001, Hebei, China.

[2] Beijing Key Laboratory of Maize DNA Fingerprinting and Molecular Breeding, Maize Research Center, Beijing Academy of Agriculture and Forestry Sciences (BAAFS), Beijing 100079, China.

[3] Plant Science and Technology College, Beijing University of Agriculture, Beijing 102206, China.

[†] Mengsi Kong and Meijie Luo contributed equally to this work.

[*] Correspondence: rentlang2003@163.com (Yanxin Zhao); maizezhao@126.com (Jiuran Zhao); yshtao@126.com (Yongsheng Tao)

ABSTRACT

Na$^+$, K$^+$ and pH homeostasis are important for plant life and they are controlled by the monovalent cation proton antiporter (CPA) superfamily. The roles of *ZmCPAs* in salt tolerance are not fully elucidated. In this study, we identified 35 *ZmCPAs* comprising 13 Na$^+$/H$^+$ exchangers (*ZmNHXs*), 16 cation/H$^+$ exchanger (*ZmCHXs*), and 6 K$^+$ efflux antiporters (*ZmKEAs*). All ZmCPAs have transmembrane domains and maize protoplasts transient expression showed that ZmNHX5 and ZmCHX17 localized to cell membrane system. *ZmCHXs* were specifically highly expressed in anthers, while *ZmNHXs* and *ZmKEAs* showed high expression in various tissues. *ZmNHX5* and *ZmKEA2* were up-regulated in maize seedlings under both NaCl and KCl stresses. Yeast complementation experiments revealed the roles of *ZmNHX5*, *ZmKEA2* in NaCl tolerance. Analyses of the maize mutants further validated the salt tolerance functions of *ZmNHX5* and *ZmKEA2*. Our study highlights comprehensive information of *ZmCPAs* and provides new gene targets for salt tolerance maize breeding.

Keywords: CPA, Maize, Salinity, *ZmNHX*, *ZmCHX*, *ZmKEA*

1. Introduction

High salinity is a significant abiotic stress factor restraining crop production worldwide. High concentration of sodium chloride (NaCl) or potassium chloride (KCl) affects the osmotic potential of the soil solution, thus inhibiting water absorption by plant. In addition, Na^+ and Cl^- ions are easily absorbed to toxic levels by plant roots, which cause ion toxicity to plant cells [1, 2]. Under the above effects, salt stress eventually leads to plant metabolism disorder, oxidation-reduction system damage, photosynthesis limitation, and finally plant growth inhibition [3].

Many monovalent cation channels/transporters have been reported to play key roles in regulating salt tolerance in plants. In *Arabidopsis*, *AtNHX1–6* and *AtNHX7/AtSOS1* were shown to be involved in maintaining Na^+, K^+ and pH homeostasis, and are critical for *Arabidopsis* salt tolerance [4, 5]. *AtKEA1*, *AtKEA3* and *AtKEA4* can enhance the tolerance of *Arabidopsis* to high concentration of K^+, and *AtKEA2* overexpression enhanced the resistance of yeast mutants to Na^+ and K^+ stress. Functional characterization of *AtCHX17*, *AtCHX21* and *AtCHX23* knockout *Arabidopsis* mutants suggested their roles in K^+ acquisition and in *Arabidopsis* salt tolerance. In rice, the expression of *OsNHX1–5* and *OsNHX7/OsSOS1* were increased by salt stress and they can suppress sensitivity of rice to high Na^+ and K^+ concentration [6, 7]. The K^+/Rb^+ competition assay of *OsCHX14* in *Xenopus laevis oocytes* verified its function in K^+ homeostasis. In maize, the expression of *ZmNHX2* was up-regulated under NaCl stress in both shoot and root of maize seedling and it was presumed to be responsible for Na^+ sequestration into vacuoles [8]; *ZmNHX7* putatively encodes a Na^+ efflux plasma membrane protein and its expression increased in maize plants exposed to salt stress [9]. All these Na^+/H^+ exchangers (NHXs), K^+ efflux antiporters (KEAs) and cation/H^+ exchangers (CHXs) carried a conserved Na^+/H^+ exchanger protein domain (Pfam00999), and are members of the monovalent cation/proton antiporter (CPA) superfamily. However, the CPA gene family in maize has not been comprehensively studied, and the role of its members in salt tolerance have not been fully elucidated.

The CAP superfamily is grouped into the CPA1 and CPA2 families based on the Transporter Classification database (http://www. tcdb.org/) [10, 11] and the CPA1 family

comprises the *NHXs* subfamily, while the CPA2 family is comprised of the *KEAs* and the

CHXs subfamilies [4, 12]. The *NHXs* is predicted to contain 10–12 transmembrane (TM)

domains. The *AtNHXs* consist of 8 members in *Arabidopsis*, and the *AtNHX1–4* and

AtNHX5–6 are localized inside cell, while *AtNHX7/AtSOS1* and *AtNHX8* are localized to the

plasma membrane (PM) [4, 5]. The different subcellular location and tissue expression

specificity between members are consistent with their different biological functions. For

instance, the *AtNHX1–2* on vacuolar membrane are responsible for cell expansion in rapidly

elongating organs; the *AtNHX5* and *AtNHX6* located at the trans-Golgi network are important

for vesicle trafficking to the vacuole in different organs throughout plant development, while

AtNHX7 on PM affect membrane trafficking and vacuolar biogenesis in root cells [4, 5].

Compared with *Arabidopsis* and rice, the function of *ZmNHXs* in maize have not been well

characterized.

The CPA2-type transporters have 8–14 putative TM domains [4]. The *Arabidopsis* had

6 members in the KEA subfamily [13]. The *AtKEA1* and *AtKEA2* are anchored in the inner

envelope membrane of chloroplasts and are involved in chloroplasts division and membrane

formation. The *AtKEA3* is located in the thylakoid membrane and it benefits photosynthesis

efficiency [14, 15]. The *AtKEA4-6* were found mainly at the Golgi and they have a dramatic

impact on K^+, pH homeostasis in endomembrane compartments and in plant growth [4]. The

function of *OsKEAs* in rice is rarely known. For the CHX subfamily, the *Arabidopsis* had 28

members. Among them, 18 *AtCHX* genes are pollen-specific or pollen-enhanced, and 6 are

highly expressed in vegetative tissues, revealing the crucial role of *AtCHXs* in pollen

development [4, 16]. In rice, *OsCHX14* was demonstrated to locate at Endoplasmic reticulum

and regulate K^+ homeostasis in flowering organs [17]. In maize, the members of *ZmKEAs* and

ZmCHXs have not been investigated and their roles in salt tolerance need to be determined.

In this study, we conducted genome-wide identification of *ZmCPA* genes in maize. Their

encoded proteins were classified into ZmNHX, ZmKEA and ZmCHX subfamilies, and their

protein properties comprising molecular weight (MW), isoelectric point (pI), evolutionary

relationships, number of TM domains, sub-cellular localization, and motifs were determined.

The splice variants, exon–intron structure, and intron phases of identified genes were

investigated. The transcriptional patterns of *ZmCPA* genes were analyzed during numerous

developmental stages and under salinity stress. Six *ZmCPA* genes were functionally

characterized by expression in a salt-sensitive yeast mutant. Analyses of two maize mutants,

zmnhx5 and *zmkea2*, further verified the functions of *ZmNHX5* and *ZmKEA2* in salt tolerance.

2. Results

2.1. *Genome-wide identification and characterization of maize CPA superfamily genes*

Extensive BLAST searches based on known *AtCPAs* identified a total of 35 ZmCPA

superfamily proteins encoded in the genome of maize (Table S1). These were further

confirmed through searching the Pfam database (http://pfam.xfam.org/search) for the

existence of the signature Na^+/H^+ exchanger (PF00999) domain. The 35 genes were classified

into three subfamilies: *ZmNHX* (13 genes), *ZmCHX* (16 genes) and *ZmKEA* (6 genes). The

identified *CPA* genes were named *ZmNHX1–ZmNHX13, ZmCHX1–ZmCHX12,*

ZmCHX14–ZmCHX17, and ZmKEA1–ZmKEA6 according to previous reports or based on

their degree of similarity with the family members in other species. The *ZmCPA*s were

distributed on all chromosomes of maize, with 5, 3, 3, 4, 1, 3, 5, 4, 3, and 4 genes on

chromosome 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10, respectively (Fig. S1).

As most of the *ZmCPA* genes had more than one transcript annotated in the MaizeGDB

database (Table S1 and Fig. S2). The transcripts with a conserved gene structure similar to

that of their orthologs (*Arabidopsis*, rice, *Sorghum*, *Brachypodium distachyon*, and *Setaria*

italica) were selected for further analyses. The transcript of each *ZmCPA* gene model was

further validated using reverse transcription polymerase chain reaction (RT-PCR) assays with

the gene-specific primers listed in Table S2. Notably, the transcripts of five *ZmCPA* genes

verified by Sanger sequencing of cDNA were different from their annotation in MaizeGDB

(Fig. S3).

The gene structures of the *ZmCPA* genes were determined by aligning their genomic

sequences with cDNA sequences obtained by RT-PCR. In general, the *ZmCPA* genes in the

same subfamily shared a conserved gene structure and the majority of introns in each family

were in 0 phase (Fig. 1). The number of introns ranged from 11 to 25, from 0 to 5, and from

16 to 19 in *ZmNHXs*, *ZmCHXs* and *ZmKEAs*, respectively (Table 1). The average length of

ZmNHX, ZmCHX, and ZmKEA proteins was 640, 824, and 815 amino acid (aa) residues,

respectively. The average molecular weight (MW) of ZmNHX, ZmCHX, and ZmKEA

proteins was 70.8, 88.4, and 87.4 kDa, respectively. The isoelectric point (pI) value of

ZmNHX, ZmCHX, and ZmKEA proteins ranged from 5.28 to 9.07, from 5.95 to 9.88, and

from 5.22 to 6.01, respectively (Table 1).

2.2. Phylogenetic analysis of maize CPA genes

To analyze the evolutionary relationships among the identified *ZmCPA* genes, we aligned

their protein sequences with 26 OsCPA and 42 AtCPA proteins to construct a neighbor-joining

(NJ) phylogenetic tree. Detailed information for *AtCPA* and *OsCPA* genes is shown in Table

S3. Based on the topology of the phylogenetic tree, the *CPA* gene superfamily in plants can be

subdivided into three subfamilies; NHX, CHX, and KEA. The NHX, CHX, KEA subfamilies

can be further divided into groups N1–3, C1–3, and K1–2, respectively (Fig. 2). All groups

contained genes from the three species of *Arabidopsis*, rice, and maize, indicating similar

functions of orthologous genes in each group. We also performed a phylogenetic analysis

using *ZmCPAs* only. As shown in Fig. 3, *ZmCPA* genes were grouped into *ZmNHX*, *ZmCHX*,

and *ZmKEA* three subfamilies, which was consistent with the classification shown in Fig. 1

(Fig. 3).

2.3. Motif and subcellular localization analysis of ZmCPA proteins

Known conserved domains of ZmCPA proteins were identified by screening the Pfam

database (http://pfam.xfam.org/). All proteins contained the Na^+/H^+ exchanger domain

(PF00999), and three ZmKEAs and five ZmNHXs also contained the Trka_N domain

(PF02254) and the TatD_DNase domain (PF01026), respectively (Fig. S4). Putative motifs of

the ZmCPA proteins were mined using tools at the MEME server

(http://meme-suite.org/tools/meme). The 15 motif sequences identified are listed in Table S4.

Generally, ZmNHX proteins contained five to eight conserved motifs, the ZmCHX proteins

had six to seven conserved motifs and the ZmKEA proteins had two to three conserved motifs

(Fig. S4). Most closely related members in the same subfamily had common motifs,

indicative of functional similarities among ZmCPA proteins.

The conserved domains of ZmCPA proteins corresponded to partial motifs and covered

the TM domain (Figure S4–5). The TM helices within ZmCPA proteins were predicted using

tools at the TMHMM Server v. 2.0 (http://www.cbs.dtu.dk/services/TMHMM/). All ZmCPA

proteins had TM domains, and each subfamily had its own characteristics (Fig. S5). A

maximum of 12 TM regions were predicted in ZmNHX5, -7, ZmCHX7, -16, -17, and

ZmKEA1. The average occurrence of TM regions was 8, 10, and 10 in ZmNHX, ZmCHX

and ZmKEA proteins, respectively (Table S1). The TM domain was mainly located at the

C-terminal in ZmCHX subfamily, while at the N-terminal in ZmKEAs.

We performed transient expression assays in maize protoplasts to determine the

subcellular localization of ZmCHX17 and ZmNHX5 proteins. The ZmNHX5-GFP and

ZmCHX17-GFP fusion proteins co-localized with OsSCMP1-RFP protein (rice

SECRETORY CARRIER MEMBRANE PROTEIN 1 fused with RFP as a membrane protein

control [18]) suggesting that ZmNHX5 and ZmCHX17 localize to the membrane system (Fig.

4).

2.4. Gene expression analysis in different tissues and developmental stages

NimbleGen maize microarray data [19] (ZM37) including 60 tissues representing 11

major organs and various developmental stages of the B73 maize inbred line were analyzed to

detect the expression patterns of *ZmCPA* genes. Gene expression data for 31 *ZmCPA* genes

including 10 *ZmNHXs,* 16 *ZmCHXs* and 5 *ZmKEAs* were used for the cluster analysis. As

revealed by the heatmap, all *ZmNHX* genes except *ZmNHX4, ZmNHX3* and *ZmNHX9* were

highly expressed in nearly all 60 tissues. All five *ZmKEA* genes were highly expressed in all

60 tissues. All *ZmCHX* genes were expressed at lower levels than *ZmNHX* and *ZmKEA* genes,

and *ZmCHX* genes were only expressed at high levels in anther (Fig. 5).

To confirm the tissue-specific expression of *ZmCPA* genes as revealed by the microarray

data, semi-quantitative reverse transcription polymerase chain reaction (semi qRT-PCR)

analyses of nine *ZmCPA* genes were performed with total RNA isolated from the roots, leaves,

ears, immature tassels, pollen, anthers, silk, and whole seeds (20 days after pollination) of the

B73 inbred line. The primers for semi qRT-PCR are listed in Table S5. Generally, the

expression patterns of nine *ZmCPA* genes were consistent with the results from the microarray

data. The *ZmNHX5*, *ZmNHX6* and *ZmNHX8* were highly expressed in root and leaf of maize,

while the *ZmKEA1*, *ZmKEA2* and *ZmKEA5* were highly expressed in most of 8 tissues. The

ZmCHX8 showed low expression in all tissues, and the *ZmCHX5* together with *ZmCHX17*

were specifically highly expressed in anther and pollen of maize (Fig. 5 and Fig. 6).

2.5. Expression analysis of ZmCPA genes under 100 mM NaCl and 100 mM KCl stress

To understand the expressional responses of *ZmCPA* genes to salt stress, genes in two

gene subfamilies, *ZmNHX* (four genes) and *ZmKEA* (four genes), were selected for analysis

by real-time quantitative reverse transcription polymerase chain reaction (qRT-PCR) analyses.

The primers are listed in Table S6. We detected the gene expression response at 0h, 1 h, 2 h, 4

h, and 24 h of salt stress. In the leaf of plants treated with 100 mM NaCl, *ZmNHX2*, *ZmNHX5*,

ZmNHX6, *ZmKEA1* and *ZmKEA2* were up-regulated, while other genes were down-regulated.

In the root, *ZmNHX5 and ZmKEA2* were up-regulated after 4 h, while other genes were

down-regulated or did not change much. Under 100 mM KCl stress treatment, all all eight

*ZmCPA*s (*ZmNHX2*, *ZmNHX5*, *ZmNHX6*, *ZmNHX8*, *ZmKEA1*, *ZmKEA2, ZmKEA5* and

ZmKEA6) had a tendency of up-regulation in leaf. In the root, only *ZmNHX5* was

up-regulated while other genes were down-regulated or showed fluctuation (Fig. 7). These

results implied that *ZmCPA*s might play a role in salinity stress tolerance through expression

regulation.

2.6. Functional analysis of ZmCPA genes in a yeast mutant under NaCl stress

To test the functions of *ZmCPA*s in salt tolerance, the coding sequences of *ZmNHX5*,

ZmCHX2, *ZmCHX3*, *ZmCHX5*, and *ZmCHX17* and *ZmKEA2* were each cloned into the yeast

expression vector pDR196 under the control of the PMA1 promoter. Each vector was

introduced into the *Saccharomyces cerevisiae* mutant strain AXT3K, which lacks the

functions of four plasma membrane Na^+-ATPases (ScENA1–4), the plasma membrane Na^+,

K^+/H^+ antiporter ScNHA1, as well as the vacuolar Na^+, K^+/H^+ antiporter ScNHX1 [20].

Therefore, this mutant is very sensitive to Na^+ stress. Each transformed yeast strain was

grown on Arg phosphate (AP) medium with or without 20 mM NaCl. Results showed that the

transformed AXT3K lines expressing these genes grew much better than the AXT3K mutant

as well as the AXT3K mutant transformed with the empty pDR196 vector under 20 mM NaCl

stress (Fig. 8). These results confirmed the functions of *ZmNHX5*, *ZmCHX2*, *ZmCHX3*,

ZmCHX5, and *ZmCHX17* and *ZmKEA2* in salt tolerance.

2.7. Functional analysis of ZmNHX5 and ZmKEA2 under high concentration salt stress

We obtained two maize mutants, *zmnhx5* (EMS4-0a18d8) *and zmkea2* (EMS4-02c2af),

which were produced by EMS mutagenesis of the B73 inbred line, from the Maize

EMS-induced Mutant Database (MEMD) [21]. The *zmnhx5* and *zmkea2* mutants have

pre-termination mutations in *ZmNHX5* (*Zm00001d022504*) and *ZmKEA2* (*Zm00001d026645*),

respectively, resulting in the production of truncated proteins (Fig. 9A, E). The phenotypes of

inbred B73 lines and the two maize mutants were analyzed after 4 days of growth under salt

stress. Under normal conditions, the growth status of the wild type and mutants was not

significantly different (Fig. 9). However, under 100 mM NaCl treatment, the shoot length of

the *zmkea2* mutant was significantly lower than that of the wild type ($P<0.05$). The shoot dry

weight of *zmkea2* was reduced greater than that of the wild type (Fig. 9B–D). Similarly, under

100 mM KCl condition, the shoot length of the *zmnhx5* mutant were significantly lower than

that of the wild type ($P<0.05$). The shoot dry weight of *zmnhx5* was reduced greater than that

of the wild type. ((Fig. 9F–H)). These results further verified that the *ZmNHX5* and *ZmKEA2*

are important salt tolerance-related genes.

3. Discussion

The cation/proton antiporter superfamily is responsible for monovalent cations exchange

in fungi, bacteria, animals, and plants [12, 16]. Functional analyses of *CPA* genes have been

conducted in *Arabidopsis* [22, 23], rice [17], wheat [24] and *Arachis hypogea* [25]. Most of

them acted on maintaining the pH, Na^+ and K^+ homeostasis in plants under high salinity [26].

In maize, the *ZmNHX7* is homologous to *AtNHX7/SOS1*, which is critical for excluding Na$^+$

from plant cell. The expression of *ZmNHX7* was up-regulated upon NaCl stress, and it is

associated with the major quantitative trait locus (QTL), *qST1*, which confers salt tolerance on

maize plants [27, 28]. Till now, there are few studies on maize *CPAs* gene family, and the

functional analysis of *ZmCPAs* are rarely performed.

In this study, 35 *ZmCPA*s were identified in maize, and they were named according to

previous reports [29] or based on their sequence similarity to known plant *CPA* genes.

Analysis of chromosomal distribution revealed that *ZmCPA*s are evenly distributed on the 10

chromosomes of maize. Similarly, *CPA* genes are located on all chromosomes in wheat [24],

and 15 out of 17 chromosomes in pear [30]. A phylogenetic tree using CPA proteins from

Arabidopsis, rice and maize revealed that the homologous proteins from maize, *Arabidopsis*

and rice were tightly clustered. For example, the *ZmNHX7*, *AtNHX7* and *OsNHX7* clustered

together; the *ZmKEA3*, *AtKEA3* and *OsKEA3* gathered together; the *ZmCHX1, AtCHX1* and

OsCHX1 were in one clade. In *Arabidopsis*, genes with similar functions are clustered nearby,

e.g., the *AtNHX1–4* localized on vacuolar membrane, *AtNHX5–6* for vesicle trafficking and

AtNHX7–8 localized on PM were distributed together, respectively. Thus it was speculated

that the clustered homologous genes from maize and *Arabidopsis* or clustered maize genes

had similar functions. Consistent with previous studies in *Arabidopsis* and pear, we also

classified the CPA superfamily proteins into NHX, KEA and CHX subfamilies and into the

N1–N3, K1–K2, and C1–C3 groups [12, 30].

The predicted sub-cellular localization of these proteins in different species is consistent

to some extent. The *AtNHXs* exhibit vacuole, endosome, and PM localization [4] and the

ZmNHXs have similar predicted localizations. Like *AtKEA1*, *AtKEA2* and *AtKEA3*, the

ZmKEA4 was predicted to localize to the chloroplast. All *ZmCHX* proteins were predicted to

localize to the PM, like *AtCHX13* and *AtCHX14* [31]. The *ZmNHX5* and *ZmCHX17* were

selected for transient expression in maize protoplast cells. As predicted, they were localized to

the membrane system.

We analyzed the expression patterns of *ZmCPA* genes according to the NimbleGen maize

microarray data. The *ZmNHX* and *ZmKEA* genes showed high expression at multiple

developmental stages, suggesting that they are important for maize growth and development. The *ZmCHXs* were specifically expressed in anthers, indicating that they participate in reproductive organ development. Similar expression trends of *CPA* genes have been found in other plant species. For example, in wheat, *TaNHX2*, *TaNHX5* and *TaNHX8* are highly expressed at multiple developmental stages, and most *TaCHX* family genes are relatively highly expressed in anthers [24]. The expression patterns of selected genes in maize were verified by semi-qRT-PCR in this study.

The differential expression of *ZmCPAs* between the control and salinity treatments were analyzed using qRT-PCR. Under 100 mM NaCl treatment, the *ZmNHX5* and *ZmKEA2* were up-regulated in both leaf and root, providing evidence of their roles in NaCl stress in maize. Under 100 mM KCl treatment, the *ZmNHX2*, *ZmNHX5*, *ZmNHX6*, *ZmNHX8*, *ZmKEA1*, *ZmKEA2*, *ZmKEA5* and *ZmKEA6* were up-regulated in the leave. The up-regulated expression of them to K^+ stress indicates that they might be involved in maintaining K^+ homeostasis in maize. The *ZmNHX5*, *ZmKEA2*, *ZmCHX2*, *ZmCHX3*, *ZmCHX5* and *ZmCHX17* and were respectively introduced into the yeast mutant strain AXT3K. All constructs restored Na^+-resistance of the AXT3K mutant. Furthermore, the growth reduction of maize EMS mutants under salt stress validated the functions of *zmnhx5* and *zmkea2* in salt tolerance. These analyses provided evidence for the role of CHX and KEA subfamilies in salt tolerance in maize. Similarly, *AtNHX5* and *AtNHX6* recovered salt tolerance in yeast [23], indicating that *ZmNHXs* have a common mode of action as other plant species. Unlike *ZmCHX* and *ZmKEA* from maize, neither *AtKEAs* nor *AtCHXs* from *Arabidopsis* are shown to improve yeast growth under high salt condition [4, 13].

It is worth mentioning that both *ZmNHX5* and *ZmKEA2* played an important role in salt tolerance of maize according to the results obtained in this study. The expression of *ZmNHX5* and *ZmKEA2* were up-regulated under salt stress (NaCl and KCl), and they can restore the Na^+-resistance of yeast mutant. Moreover, the salt tolerance function (NaCl) of *ZmNHX5* and *ZmKEA2* were further verified by maize mutant experiment. The functional analysis of these two genes needs further study, and they are likely to be new gene targets for breeding salt tolerance maize.

4. Conclusion

We identified and characterized the *ZmCPA* superfamily, consisting of *ZmNHX*, *ZmKEA* and *ZmCHX* subfamily genes, from the maize genome. Gene and protein sequences analyses demonstrated high conservation of evolutionarily related members in each subfamily, while members differed among different groups. The existence of many helices, coils and TM regions in protein structure are consistent with the hydrophobic and membrane-bound nature of these proteins. Expression analyses revealed diverse expression patterns in various tissues and under different abiotic stresses, indicating that these genes are important for maize growth, development, and salt stress responses. Characterization of *ZmNHX5*, *ZmCHX2*, *-3*, *-5*, *-17* and *ZmKEA2* in yeast confirmed their roles in maintaining monovalent cation homeostasis and in salt stress tolerance. The functions of *ZmNHX5* and *ZmKEA2* were verified by phenotypic analyses of EMS mutants. This study not only describes numerous features of *ZmCPA* genes, but also provides the foundation for functional validation of these genes in further studies.

5. Materials and methods

5.1. Identification and bioinformatic analysis of ZmCPA gene superfamily

The known *CPA* genes in *Arabidopsis* were used to query the maize AGPv4 gene set (https://download.maizegdb.org/Zm-B73-REFERENCE-GRAMENE-4.0/) using a local BLASTP program with an E-value <1e-10. The putatively identified sequences were further confirmed by an HMMER search for the presence of signature Na^+/H^+ exchanger (PF00999) domain.

The phylogenetic tree was constructed using full-length CPA protein sequences of maize, *Arabidopsis*, and rice. Sequences were aligned using MUSCLE v3.8.31 [32, 33], and a phylogenetic tree was built using the neighbor-joining (NJ) method with MEGA 6.0 [34] with a bootstrap value of 1000 and amino acid substitution according to the Poisson model. The TM domains were predicted using tools at the TMHMM Server v. 2.0 (http://www.cbs.dtu.dk/services/TMHMM/). Predotar [35], TargetP [36], and WoLF PSORT (https://www.genscript.com/wolf-psort.html), three *in silico* programs, were used to predict

the organellar localization of ZmCPA proteins. Putative conserved motifs in maize CPA proteins were identified using the MEME Suite 5.1.1 (http://meme-suite.org/tools/meme) with the following parameters: motif length, 10–50 aa; maximum number of motifs, 15.

5.2. Gene structure analysis

The DNA and transcript sequences of *ZmCPA* genes obtained from the maize sequence annotation database MaizeGDB were used to design gene-specific PCR primers with Primer3 (http://primer3.ut.ee/). The DNA and cDNA sequences were validated using PCR and RT-PCR with B73 genomic DNA and total RNA as templates. The gene-specific primers shown in Table S2. Alignment of validated DNA and cDNA sequences of each maize *CPA* gene revealed the structure of *ZmCPA* genes. Gene structure display server (GSDS 2.0) was used to display the exon–intron structure, and intron phases [37].

5.3. Plant materials and treatments

The maize B73 inbred line was used in this study. For qRT-PCR, the sterilized seeds were grown in a hydroponic system as described previously [38] with sterile water in a greenhouse at 27/23°C under a 12-h light/12-h dark photoperiod. Four days later, the plants were incubated in 1× Hoagland's solution (PhytoTechnologies Ltd., Shawnee Mission, KS, USA) until the first leaves were fully expanded. Maize plants were grown with 1 × Hoagland's solution as the control (CK) and with 1 × Hoagland's solution containing 100 mM NaCl for the NaCl stress treatment or 100 mM KCl for the KCl stress treatment. These concentrations were maintained until the end of the experiments.

For maize EMS mutants experiment, the sterilized seeds were cultured hydroponically in a greenhouse at 27/23°C under a 12-h light/12-h dark photoperiod as described above. The 1× Hoagland's solution was exchanged every 2 days. Ten days later, 100 mM KCl and 100 mM NaCl were added. After 4 days of salt stress, phenotypic analyses were performed.

5.4. Expression analysis of ZmCPA genes in different tissues

To investigate the spatiotemporal expression patterns of *ZmCPA* genes, the

log2-transformed and RMA-normalized data for *ZmCPA* genes were downloaded from PLEXdb (http://www.plexdb.org/) [39]. A heat map was produced using the Lianchuan Bio Cloud Platform (https://www.lc-bio.cn/overview).

5.5. RNA isolation and cDNA synthesis

Total RNA was isolated from different tissues of the B73 inbred line, including seedling roots, leaves, 5-cm ears, immature tassels, anthers, pollens, silks, and seeds at 20 days after pollination, using Trizol reagent (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's protocol. All RNA was purified using DNase I (Thermo Scientific, Waltham, MA, USA). First-strand cDNA was synthesized from 1 μg total RNA (20 μL reaction volume) using a PrimeScript™ 1st Strand cDNA Synthesis Kit (Takara, Dalian, China) according to the manufacturer's protocol.

5.6. Semi-quantitative reverse transcription PCR

All gene-specific primers were designed as shown in Table S5. Specific primers were used to amplify maize *Actin1* (*GRMZM2G126010*) as an internal control. Reactions were performed with $2 \times$ Taq Master Mix (Vazyme, Nanjing, China) on a Bio-Rad Thermal Cycler (Bio-Rad, Hercules, CA, USA) using the following procedure: 5 min at 94 °C to start; 33 cycles of 30 s at 94 °C, 30 s at 59 °C and 2 min at 72 °C; and a final extension step of 72 °C for 10 min to complete the reaction. *Actin1* was amplified with 29 PCR cycles. Each PCR was performed in triplicate, mixtures without a template were employed as negative controls, and the maize *Actin1* amplicon served as an internal control for each gene investigated.

5.7. Real-time PCR

Real-time PCR was performed using TB Green Premix Ex Taq™ II (Takara, Dalian, China). An ABI 7500 real-time PCR system (Applied Biosystems, Foster City, CA, USA) was used with the following thermal cycling conditions: 95 °C for 5 min, followed by 40 cycles of 95 °C for 15 s, 60 °C for 5 s, and 72 °C for 34 s. The maize *Actin1* gene (*GRMZM2G126010*) was used as an endogenous control to normalize the samples. Based on the cDNA sequences

of *ZmCPA* genes, real-time PCR primers (Table S6) were designed with Primer 3

(http://primer3.ut.ee/). The experiment was performed with three technical replicates for each

sample. The specificity of each PCR reaction was confirmed by melting curve analysis of the

amplicons. The comparative $2^{-\Delta\Delta CT}$ method was used to calculate the relative transcript levels

in the samples [40].

5.8. Subcellular localization

The full-length cDNAs of *ZmCHX17* and *ZmNHX5* were amplified using the primers

listed in Table S7 and then were introduced into the pM999-EGFP vector to construct the GFP

fusion proteins ZmCHX17-GFP and ZmNHX5-GFP, respectively, with the ClonExpress II

One Step Cloning Kit (Vazyme). Constitutive expression of the fused constructs,

ZmCHX17-GFP and ZmNHX5-GFP, was driven by the cauliflower mosaic virus 35S

promoter. Maize mesophyll protoplasts were isolated and prepared from etiolated leaves

according to established protocols [41]. The plasmids harboring the ZmCHX17-GFP and

ZmNHX5-GFP fusion constructs were co-transfected with the OsSCAMP1-RFP construct

into protoplast cells, respectively. OsSCAMP1 is a known rice secretory carrier membrane

protein and was used as a membrane protein control [18]. The transformed protoplast cells

were cultured at room temperature overnight and then observed under a Leica SP8 confocal

microscope (Leica, Wetzlar, Germany).

5.9. Functional expression of ZmCPAs in yeast

The coding sequences of *ZmNHX5*, *ZmCHX2*, *ZmCHX3*, *ZmCHX5*, *ZmCHX17*, and

ZmKEA2 were each cloned into the PDR196 vector, and then transformed into the yeast strain

AXT3K (*ena1-4::HIS3,nha1::LEU2, nhx1::KanMX*). The transformed yeast cells were

cultured overnight at 29 °C in YPDA medium containing 1 mM KCl. Cell cultures were

diluted in water to an A_{600} of 0.8. To test cation tolerance, 5-μL aliquots from yeast cultures or

10-fold serial dilutions were spotted onto AP [42] plates containing 1 mM KCl and

supplemented with or without 20 mM NaCl.

List of abbreviations

CPA: monovalent cation proton antiporter; NHX: Na^+/H^+ exchanger; KEA: K^+ efflux antiporter; CHX: cation/H^+ exchanger; Na^+: sodium; Cl^-: chloride; TM: transmembrane; K^+: potassium; H^+: proton; MW: molecular weight; pI: isoelectric point; RT-PCR: reverse transcription polymerase chain reaction; DNA: deoxyribonucleic acid; cDNA: complementary DNA; NJ: neighbor-joining; aa: amino acid; kDa: kilodalton; TF: transcription factor; semi qRT-PCR: semi-quantitative reverse transcription polymerase chain reaction; AP: Arg phosphate; EMS: ethyl methanesulfonate; MEMD: Maize EMS-induced Mutant Database; QTL: quantitative trait locus; GFP: green fluorescent protein; RFP: red fluorescent protein.

Authors' contributions

Y.T., Y.Z., (Yanxin Zhao) and J.Z. conceived the experiment. M.K, M.L, J.L, Y.Z. (Yunxia Zhang), Z.F, W.S, R.Z, R.W., and Y.W. performed bioinformatic analysis and data acquisition. M.K, M.L, and Y.Z (Yanxin Zhao) analyzed the data and wrote the manuscript. All authors revised and approved the final manuscript, and agreed to be accountable for this work.

Funding

This research was financially supported by the Beijing Municipal Natural Science Foundation (6204041), the Construction of Collaborative Innovation Center of Beijing Academy of Agricultural and Forestry Sciences (Collaborative Innovation Center of Crop Phenomics, KJCX201917), Youth Research Fund of Beijing Academy of Agriculture and Forestry Sciences (QNJJ202028), and the Beijing Scholars Program (BSP041). The funding body does not play roles in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Declaration of Competing Interest

The authors declare that they have no competing interests.

Acknowledgements

We thank Prof. Liwen Jiang (Chinese University of Hong Kong) for kindly providing pM999-EGFP and membrane protein marker (OsSCAMP1-RFP) plasmids, Prof. Xiaoduo Lu (Qilu Normal University) for identifying and providing maize EMS mutants *zmkea2* and *zmnhx5*, and Profs. Yi Wang and Zhenxian Zhang (China Agricultural University) for kindly sharing yeast mutant AXT3K and yeast expression vector pDR196.

References

[1] Zhu JK. Plant salt tolerance. Trends Plant Sci. 2001, 6:66–71.

[2] Gupta B, Huang B. Mechanism of salinity tolerance in plants: physiological, biochemical, and molecular characterization. Int J Genom. 2014, 1:701596.

[3] Farooq M, Hussain M, Wakeel A, Siddique KHM. Salt stress in maize: effects, resistance mechanisms, and management. A review. Agron Sustain Dev. 2015(35), 35:461–481.

[4] Chanroj S, Wang G, Venema K, Zhang MW, Delwiche CF, Sze H. Conserved and diversified gene families of monovalent Cation/H^+ Antiporters from algae to flowering plants. Front Plant Sci. 2012, 3:25.

[5] Tian FX, Chang E, Li Y, Sun P, Hu JJ, Zhang J. Expression and integrated network analyses revealed functional divergence of NHX-type Na^+/H^+ exchanger genes in poplar. Sci Rep. 2017, 7:2607.

[6] Fukuda A, Nakamura A, Hara N, Toki S, Tanaka Y. Molecular and functional analyses of rice NHX-type Na^+/H^+ antiporter genes. Planta. 2011. 233(1):175–188.

[7] Sellamuthu G, Jegadeeson V, Sajeevan RS, Rajakani R, Parthasarathy P, Raju K, Shabala L, Chen ZH, Zhou MX, Sowdhamini R, Shabala S, Venkataraman G. Distinct evolutionary origins of intron retention splicing events in *NHX1* antiporter transcripts relate to sequence specific distinctions in *Oryza* species. Front Plant Sci. 2020, 11:267.

[8] Huanca-Mamani W, Ortiz MV, Cárdenas-Ninasivincha S, Acosta-Garcia G, Bastías E. Gene expression analysis in response to combined salt and boron (B) stresses in a tolerant maize landrace. Plant Omics. 2018, 11(2):80–88.

[9] Bosnic P, Bosnic D, Jasnic J, Nikolic M. Silicon mediates sodium transport and partitioning in maize under moderate salt stress. Environ Exp Bot. 2018, 155:681–687.

[10] Pires IS, Negrão S, Pentony MM, Abreu IA, Oliveira MM, Purugganan MD. Different evolutionary histories of two cation/proton exchanger gene families in plants. BMC Plant Biol. 2013, 13(1): 97.

[11] Saier MH, Reddy VS, Tamang DG, Västermark A. The transporter classification database. Nucleic Acids Res. 2014, 42(D1):D251–D258.

[12] Ye CY, Yang X, Xia X, Yin W. Comparative analysis of cation/proton antiporter superfamily in plants. Gene. 2013, 521(2):245–251.

[13] Zheng S, Pan T, Fan L, Qiu Q. A novel *AtKEA* gene family, homolog of bacterial K^+/H^+ antiporters, plays potential roles in K+ homeostasis and osmotic adjustment in *Arabidopsis*. PLoS One. 2013, 8(11):e81463.

[14] Kunz HH, Gierth M, Herdean A, Satoh-Cruz M, Kramer DM, Spetea C, Schroeder JI.
Plastidial transporters KEA1, -2, and -3 are essential for chloroplast osmoregulation,
integrity, and pH regulation in *Arabidopsis*. Proc Natl Acad Sci USA. 2014,
111(20):7480–7485.
[15] Aranda-Sicilia MN, Aboukila A, Armbruster U, Cagnac O, Schumann T, Kunz HH,
Jahns P, Rodríguez-Rosales MP, Sze H, Venema K. Envelope K^+/H^+ antiporters AtKEA1
and AtKEA2 function in plastid development. Plant Physiol. 2016, 172(1):441–449.
[16] Isayenkov SV, Dabravolski SA, Pan T, Shabala S. Phylogenetic diversity and
physiological roles of plant monovalent cation/H^+ antiporters. Front Plant Sci. 2020,
11:573564.
[17] Chen Y, Ma J, Miller AJ, Luo B, Wang M, Zhu Z, Ouwerkerk PBF. OsCHX14 is
involved in the K+ homeostasis in rice (*Oryza sativa*) flowers. Plant Cell Physiol. 2016,
57(7):1530–1543.
[18] Lam SK, Siu CL, Hillmer S, Jang S, An G, Robinson DG, Jiang L. Rice SCAMP1
defines clathrin-coated, *trans*-golgi-located tubular-vesicular structures as an early
endosome in tobacco BY-2 cells. Plant Cell. 2007, 19(1):296–319.
[19] Sekhon RS, Lin H, Childs KL Hansey CN, Buell CR, de Leon N, Kaeppler SM.
Genome-wide atlas of transcription during maize development. Plant J. 2011,
66(4):553–563.
[20] Quintero FJ, Blatt MR, Pardo JM. Functional conservation between yeast and plant
endosomal Na^+/H^+ antiporters. FEBS Lett. 2000, 471(2-3):224–228.
[21] Lu X, Liu J, Ren W, Yang Q, Chai Z, Chen R, Wang L, Zhao J, Lang Z, Wang H, Fan Y,
Zhao J, Zhang C. Gene-indexed mutations in maize. Mol Plant. 2018, 11(3):496–504.
[22] Apse MP, Sottosanto JB, Blumwald E. Vacuolar cation/H^+ exchange, ion homeostasis,
and leaf development are altered in a T-DNA insertional mutant of *AtNHX1*, the
Arabidopsis vacuolar Na^+/H^+ antiporter. Plant J. 2003;36:229–239.
[23] Bassil E, Ohto M, Esumi T, Tajima H, Zhu Z, Cagnac O, Belmonte M, Peleg Z,
Yamaguchi T, Blumwald E. The *Arabidopsis* intracellular Na^+/H^+ antiporters NHX5 and
NHX6 are endosome associated and necessary for plant growth and development. Plant
Cell. 2011, 23(1):224–239.
[24] Sharma H, Taneja M, Upadhyay SK. Identification, characterization and expression
profiling of cation-proton antiporter superfamily in *Triticum aestivum* L. and functional
analysis of *TaNHX4-B*. Genomics. 2020, 112(1):356–370.
[25] Zhang WW, Meng JJ, Xing JY, Yang S, Guo F, Li XG, Wan SB. The K^+/H^+ antiporter
AhNHX1 improved tobacco tolerance to NaCl stress by enhancing K^+ retention. J Plant
Biol. 2017, 60(3):259–267.
[26] Jia Q, Zheng C, Sun S, Amjad H, Liang K, Lin W. The role of plant cation/proton
antiporter gene family in salt tolerance. Biol Plantarum. 2018, 62(4):617–629.
[27] Luo M, Zhao Y, Zhang R, Xing J, Duan M, Li J, Wang N, Wang W, Zhang S, Chen Z,
Zhang H, Shi Z, Song W, Zhao J. Mapping of a major QTL for salt tolerance of mature
field-grown maize plants based on SNP markers. BMC Plant Biol. 2017, 17(1):140.
[28] Luo M, Zhang Y, Chen K, Kong M, Song W, Lu B, Shi Y, Zhao Y, Zhao J. Mapping of
quantitative trait loci for seedling salt tolerance in maize. Mol Breeding. 2019, 39:64.
[29] Zörb C, Noll A, Karl S, Leib K, Yan F, Schubert S. Molecular characterization of Na^+/H^+

antiporters (ZmNHX) of maize (*Zea mays* L.) and their expression under salt stress. J
Plant Physiol. 2005, 162(1):55–66.

[30] Zhou H, Qi K, Liu X, Yin H, Wang P, Chen J, Wu J. Zhang S. Genome-wide
identification and comparative analysis of the cation proton antiporters family in pear
and four other Rosaceae species. Mol Genet Genomics. 2016, 291(4):1727–1742.

[31] Zhao J, Li P, Motes CM, Park S, Hirschi KD, CHX14 is a plasma membrane K-efflux
transporter that regulates K^+ redistribution in *Arabidopsis thaliana*. Plant Cell Environ.
2015, 38(11):2223–2238.

[32] Luo MJ, Shi YX, Yang Y, Zhao YX, Zhang YX, Shi YM, Kong MS, Li CH, Feng Z, Fan
YL, Xu L, Xi SL, Lu BS, Zhao JR. Sequence polymorphism of the *waxy* gene in waxy
maize accessions and characterization of a new *waxy* allele. Sci Rep. 2020, 10:15851.

[33] Edgar RC. MUSCLE: a multiple sequence alignment method with reduced time and
space complexity.BMC Bioinformatics. 2004, 5:113.

[34] Tamura K, Stecher G, Peterson D, Filipski A, Kumar S.MEGA6: molecular evolutionary
genetics analysis version 6.0. Mol Biol Evol. 2013, 30(12):2725–2729

[35] Small I, Peeters N, Legeai F, Lurin C. Predotar: A tool for rapidly screening proteomes
for N-terminal targeting sequences. Proteomics. 2004, 4(6):1581–1590.

[36] Emanuelsson O, Brunak S, von Heijne G, Nielsen H. Locating proteins in the cell using
TargetP, SignalP and related tools. Nat Protoc. 2007, 2(4):953–971.

[37] Hu B, Jin J, Guo AY, Zhang H, Luo J, Gao G. GSDS 2.0: an upgraded gene feature
visualization server. Bioinformatic. 2015, 31:1296–1297.

[38] Luo M, Zhao Y, Wang Y, Shi Z, Zhang P, Zhang Y, Song W, Zhao J. Comparative
proteomics of contrasting maize genotypes provides insights into salt-stress tolerance
mechanisms. J Proteome Res. 2018, 17(1):141–153.

[39] Dash S, Van Hemert J, Hong L, Wise RP, Dickerson JA. PLEXdb: gene expression
resources for plants and plant pathogens. Nucleic Acids Res. 2010, 40(Database
issue):D1194-1201.

[40] Thomas S, Kenneth D, Livak J. Analyzing real-time PCR data by comparative CT
method. Nat Protoc. 2008, 3(6):1101–1108.

[41] Yoo SD, Cho YH, Sheen J. *Arabidopsis* mesophyll protoplasts: a versatile cell system for
transient gene expression analysis. Nat Protoc.2007, 2(7):1565-1572.

[42] Rodríguez-Navarro A, Ramos J. Dual system for potassium transport in *Saccharomyces*
cerevisiae. J Bacteriol. 1984, 159(3):940–945.

Figure legends

Fig. 1. Gene structures of *ZmCPA* genes.
Neighbor-joining tree of maize *CPA* genes at left is based on Fig. 3. Exon–intron organization of *ZmCPA* genes (right) as determined by aligning DNA and cDNA sequences of *CPA* genes at GSDS 2.0 server (http://gsds.gao-lab.org/index.php). Introns and exons are shown as thin lines and yellow boxes, respectively. Numbers 0, 1, and 2 represent different intron phases. Blue boxes at 5′ and 3′ ends represent untranslated regions (UTRs).

Fig. 2. Phylogenetic tree of *CPA* genes in maize, *Arabidopsis*, and rice.
Neighbor-joining tree constructed using MEGA 6.0 based on alignment of 103 CPA proteins including 35 ZmCPAs, 42 AtCPAs, and 26 OsCPAs. NHX, KEA, and CHX subfamilies are divided into three (N1, N2 and N3), three (C1, C2, and C3) and two (K1 and K2) groups, respectively.

Fig. 3. Phylogenetic tree of *ZmCPA* genes.
All 35 ZmCPA proteins were aligned using MUSCLE v3.8.31 and alignment was used to construct neighbor-joining tree with MEGA 6.0. *ZmCPA* genes are divided into three subfamilies; *ZmNHXs, ZmKEAs,* and *ZmCHXs*.

Fig. 4. Subcellular localization of ZmCHX17 and ZmNHX5 proteins.
ZmCHX17-GFP (upper panel) and ZmNHX5-GFP (lower panel) fusion proteins transiently expressed in maize protoplast cells co-transformed with OsSCAMP1-RFP fusion construct (membrane protein marker). GFP and RFP signals were detected by confocal fluorescent microscopy (Leica SP8). Scale bar = 5 μm.

Fig. 5. Gene expression profiles of *ZmCPA* genes in different tissues.
Normalized microarray expression data of *ZmCPA* genes were downloaded from PLEXdb (http://www.plexdb.org/). Cluster analysis was performed and heatmap was generated using tools in the Lianchuan BioCloud Platform (https://www.lc-bio.cn/overview).

Fig. 6. Expression analysis of *ZmCPA* genes in eight tissues of B73 inbred line by semi qRT-PCR.
Total RNA was isolated from eight tissues: seedling roots, seedling leaves, 5-cm immature ears, tassels, anthers, pollen, silk and whole seeds (20 days after pollination, DAP), and then used for semi qRT-PCR analyses of *ZmCPA* genes. Maize *Actin1* gene was the internal control.

Fig. 7. Expression analysis of *ZmCPA* genes under salt stress conditions by qRT-PCR.
Expression analysis of *ZmCPA* genes under 100 mM NaCl or 100 mM KCl as determined by quantitative reverse transcription-polymerase chain reaction (qRT-PCR). Y-axis represents relative transcript levels of *ZmCPA* genes compared with that of *actin1*. X-axis represents different time points after salt treatment in each group. Error bars represent standard deviations for three replicates.

Fig. 8. *ZmCPA* genes facilitate growth of yeast mutant strain AXT3K under salt stress.
Full-length CDS sequences of *ZmNHX5*, *ZmCHX2*, *ZmCHX3*, *ZmCHX5*, *ZmCHX17*,
ZmKEA2 were each cloned into PDR196 vector, then transformed into yeast strain AXT3K
(*ena1-4::HIS3,nha1::LEU2, nhx1::KanMX*). Yeast transformant cells were cultured overnight,
then were diluted with water to A_{600} of 0.8. Aliquots (5 μL) of 10-fold serial dilutions were
spotted on AP plates supplemented with 20 mM NaCl at pH 7.5 (right panel) and normal AP
plates as the control (left panel). Strains were grown at 29°C for 5 days and photographed.
639
Fig. 9. Growth performance of *zmnhx5* and *zmkea2* mutant seedlings under salt stress.
Point mutation sites of EMS mutants *zmnhx5* and *zmkea2* were identified by PCR sequencing.
Stop codons induced by EMS resulted in truncated proteins (a, e). Growth status (b, f),
seedling length (c, g) and dry weight (d, h) of mutant seedlings shoot under 100 mM NaCl or
under 100 mM KCl treatment for 4 days (n = 6, mean ± SEM). *, Significant difference
(Student's t-test; $P < 0.05$). NS, non significant.

676 **Table 1. Information of maize *CPA* genes identified in this study.**

Gene Name	Gene Model	Chromosomal Location	CDS (bp)	Protein Length (aa)
ZmNHX1	*Zm00001d028330*	Chr1: 31041489-31049951	1599	532
ZmNHX7	*Zm00001d031232*	Chr1: 182925112-182945175	3411	1136
ZmNHX11	*Zm00001d003728*	Chr2: 56632216-56643296	2367	788
ZmNHX6	*Zm00001d048732*	Chr4: 4469601-4475393	1641	546
ZmNHX8	*Zm00001d019978*	Chr7: 81486383-81509317	1611	536
ZmNHX3	*Zm00001d020892*	Chr7: 135679272-135687679	1491	496
ZmNHX10	*Zm00001d021844*	Chr7: 164344371-164361391	2820	939
ZmNHX5	*Zm00001d022504*	Chr7: 179047417-179052457	1620	539
ZmNHX4	*Zm00001d045883*	Chr9: 45473833-45477106	1638	545
ZmNHX12	*Zm00001d048459*	Chr9: 156705048-156709982	1440	479
ZmNHX2	*Zm00001d024832*	Chr10: 90134129-90141743	1680	559
ZmNHX13	*Zm00001d025052*	Chr10: 101802712-101807654	1563	520
ZmNHX9	*Zm00001d026118*	Chr10: 138832710-138862665	2115	704
ZmCHX11	*Zm00001d031077*	Chr1: 176192733-176196049	2388	795
ZmCHX15	*Zm00001d031078*	Chr1: 176200721-176205899	2061	686
ZmCHX7	*Zm00001d005032*	Chr2: 155039785-155042337	2553	850
ZmCHX4	*Zm00001d041198*	Chr3: 105028259-105030854	2529	842
ZmCHX9	*Zm00001d044623*	Chr3: 233503244-233505658	2415	804
ZmCHX8	*Zm00001d049663*	Chr4: 39074035-39076527	2493	830
ZmCHX6	*Zm00001d050509*	Chr4: 94925858-94928987	2232	743
ZmCHX2	*Zm00001d053237*	Chr4: 221575337-221578465	2478	825
ZmCHX1	*Zm00001d017805*	Chr5:207302222-207308368	2484	827
ZmCHX12	*Zm00001d035631*	Chr6:37793506-37814621	2571	856
ZmCHX16	*Zm00001d038517*	Chr6:159098609-159101687	2598	865
ZmCHX3	*Zm00001d021461*	Chr7:152741608-152744549	2424	807
ZmCHX14	*Zm00001d009889*	Chr8:87464523-87467367	2391	796
ZmCHX5	*Zm00001d010601*	Chr8:121526743-121529490	2475	824
ZmCHX10	*Zm00001d010629*	Chr8:122278283-122281316	2874	957
ZmCHX17	*Zm00001d012521*	Chr8:176021482-176024817	2628	875
ZmKEA5	*Zm00001d027466*	Chr1:5893819-5906194	1791	596
ZmKEA4	*Zm00001d001788*	Chr2:935639-948444	3480	1159
ZmKEA3	*Zm00001d041308*	Chr3:111434917-111440097	2397	798
ZmKEA1	*Zm00001d036981*	Chr6:108202268-108211871	1848	615
ZmKEA6	*Zm00001d046231*	Chr9:73621310-73635421	1872	623
ZmKEA2	*Zm00001d026645*	Chr10:149326566-149340249	3315	1104

Legend: ▨ CDS ▬ upstream/ downstream — Intron 0 1 2: intron phase

10kb

A

CAG/TAG

ZmKEA2   TACTCTCTG**G**TGGAATTCG
zmkea2   TACTCTCT**A**TGGAATTCG

B

Hoagland 100 mM NaCl   Hoagland 100 mM NaCl
ZmKEA2   *zmkea2*

C

■ Hoagland
■ 100 mM KCl

Shoot length (cm)

ns   94.6%   *

67.0%

ZmKEA2   *zmkea2*

D

Shoot dry weight (g)

ns   95.1%   ns

66.5%

ZmKEA2   *zmkea2*

E

TGG/TGA

ZmNHX5   CCGGCCCAC**C**ATATTACA
zmnhx5   CCGGCCCA**T**CATATTACA

F

Hoagland 100 mM KCl   Hoagland 100 mM KCl
ZmNHX5   *zmnhx5*

G

Shoot length (cm)

ns   94.2%   **

71.6%

ZmNHX5   *zmnhx5*

H

Shoot dry weight (g)

ns   103.1%   *

72.6%

ZmNHX5   *zmnhx5*