Macroevolutionary imprints on the assembly of frugivory networks

Gustavo Burin[1,2], Paulo R. Guimarães Jr.[1], Tiago B. Quental[1,3]

[1]Departamento de Ecologia, Instituto de Biociências, Universidade de São Paulo

[2]gustavoburin@usp.br

[3]tbquental@usp.br

Biological interactions are a key aspect of ecological communities (Delmas et al., 2018). Current interaction network structures are snapshots of dynamic processes of community assembly (Thompson, 2005), and represent the product of the evolutionary history of interacting species over millions of years. Thus, assessing the deep time mechanisms affecting the assembly of ecological networks are key to better understand biodiversity changes in broader time scales (Jablonski, 2008). Here we integrate tools from macroevolutionary studies with network science to show that more central species in frugivory networks belong to lineages with higher macroevolutionary stability. This association is more pronounced in warmer, wetter, less seasonal environments, which highlights the role of environmental factors in shaping ecological networks. Furthermore, our results suggest that these environments possess a more diverse (either in species number or ecology) assemblage of species that can be sorted during network assembly. Lastly, we found evidence that the macro-evolutionary contribution to network stability changes in geographical space. This reinforces the idea that the macroevolutionary sorting mechanism acts at the regional pool of species, rather than selecting absolute paces of diversification. Hence our results suggest an interplay between ecological roles and diversification regimes in shaping the fate of lineages of plants and seed dispersing birds.

Seed dispersal is a fundamental ecosystem service (Fleming & Kress 2013). This

mutualism between angiosperms and, mainly, vertebrates started about 80 Mya

(Eriksson, 2016, 2014), and currently between 70-90% of woody species rely on

vertebrate species to seed dispersal (Jordano, 2000). Accordingly, many vertebrate

groups have fruits as at least part of their diet (56% of bird families, and up to 95% of

species for some mammal groups, in addition to squamates, chelonians, fishes, and even

frogs – Jordano, 2000, Donatti et al. 2011). The radiations of important groups of seed-

dispersers, such as birds and mammals, are hypothesized to be linked to the angiosperm

rise and dominance during the Cenozoic (Eriksson, 2016), although the causal links are

elusive. In particular, some bird groups are consistently recognized as specialized

frugivores (Snow, 1981; Fleming et al. 1987, Kissling et al. 2009).

In the past decades, studies on seed-dispersal networks have: 1- provided

important insights of how mutualisms are structured and maintained at the local level

(Bascompte & Jordano, 2007); 2- have supported the role of coevolution in shaping traits

(Eriksson, 2016); 3- have shown how fast selection imposed by frugivores lead to trait

evolution (Galetti et al. 2013). However, most studies have focused on describing and

understanding patterns and processes at ecological timescales. The few studies with

broader temporal scales are either those investigating community phylogenetics (see

Harmon et al. 2019 and references therein), the potential association between

diversification rates and diet, including those specialized in fruit (e.g. Burin et al 2016),

the patterns of phylogenetic conservatism of specialization/generalization of interactions

(Gómez et al. 2010), or more broadly the effect of mutualisms on diversification rates in

vertebrates (e.g. Gómez & Verdú, 2012) and plants (e.g. Weber & Agrawal 2014).

On the other hand, we are beginning to understand how diversification dynamics

may affect the assembly process, and consequently, the structure of interaction networks

(e.g. Nuismer & Harmon, 2015, Roopnarine et al. 2019). Although the whole interaction

networks might be plastic both in time and space (Ponisio et al. 2017), some evidence

suggests that the core of seed dispersal network is somewhat robust to short-term (yearly)

fluctuations of fruit availability and bird species presence (Plein et al. 2013). Moreover,

theory suggests that predictable and stable species in a particular place should interact

with more partners (Olesen et al. 2008, Yang et al. 2013), supporting the idea that at least

the core of the network might show some temporal stability. Lastly, the identity of

interacting partners typically shows phylogenetic signal, meaning that related species

tend to share similar interactions (Rezende et al. 2007, Gómez et al. 2010). Although the

existence of non-random structures in seed-dispersal networks suggest some long-term

mechanisms driving the assembly of networks (Fleming & Kress, 2013), it is still

unknown whether or how these non-random structures are maintained and detectable

over longer timescales, and what are potential mechanisms behind it.

Using diversification rates estimated from molecular phylogenies, and a dataset

composed of 29 frugivory networks we tested the hypothesis that macroevolutionary

predictable lineages are more likely to contribute with species to the core of ecological

networks, providing an explicit macroevolutionary mechanism for network assemblage

(see figure 1 for schematic version of general approach, and data used). To test this idea,

we use a hierarchical Bayesian approach to investigate the potential association between

the species' ecological roles and their macroevolutionary stability, therefore integrating

two areas of knowledge. We define as stable lineages those that either have low

extinction fraction (i.e. relative low extinction when compared to speciation), and/or

higher net diversification (, i.e. lineages that can accumulate species at a faster pace). We

estimated the species' ecological role by using tools derived from network science. We

then incorporated phylogenetic and ecological uncertainties, along with environmental

factors, in order to properly address how the interplay between biotic and abiotic factors

drives the assembly of local networks due to macroevolutionary dynamics. Lastly, we

tested whether the sorting of macroevolutionary regimes takes place at the regional or at a

global scale.

We find that central species tend to belong to macroevolutionary stable lineages.

The standardized extinction fraction is negatively correlated with species' ecological role

represented by their centrality values (figure 2). Similarly, net diversification rates are

positively correlated with species centralities (figure 2). In both cases the posterior

distributions are clearly off centered with respect to a slope value of zero. These results

hold after accounting for multiple sources of uncertainty. Median standardized effects for

extinction fraction and net diversification were -0.14 and 0.19, respectively, indicating

the two components of macroevolutionary stability contribute with similar intensity to the

structuring of the networks. Sensitivity analysis based on the medians of the posterior

distributions for most parameters for each individual tree (figure 2) suggests that this

signal holds irrespective of phylogenetic uncertainty.

Our results provide evidence that this macro-evolutionary sorting mechanism is

predominantly a regional-scale phenomenon, since we only observed evidence for the

sorting when analyzing the standardized rates (deviations from the mean rate per

network), but not for the raw absolute rates (Fig S3). This suggests that the macro-

evolutionary sorting mechanism acts at the relative rank of stability for the available

lineages, and not on absolute values of extinction fraction or diversification rates. We

also found that environmental conditions affect the relationship between ecological role

and macroevolutionary stability, with warmer, wetter, non-seasonal environments having

a stronger sorting of macroevolutionary stability regimes (figure 2). Interestingly, these

environments harbor the highest frugivorous species richness (Kissling et al. 2009), and it

is possible that such species rich environments allow for a finer sub-division of

ecological roles on which this macroevolutionary sorting mechanism could more

efficiently act. In fact, networks found on those tropical-forest-like environments are

more centralized (i.e. a higher difference in centrality between the more central and more

peripheral species - see figure 2 panel PC1) than networks commonly found on colder,

drier, and seasonal environments. An analysis using a different dataset, in this case

species dietary composition, suggests a tendency for many networks from warm/wet

climates to show broader total amplitude in dietary "types" (figure 3A), corroborating the

finding that warm and humid environments possess a broader range of species types (not

only more species) on which the macro-evolutionary sorting can more efficiently act.

Moreover, warmer, more humid and less seasonal climates allow for a more

homogeneous occupation of the dietary space without increasing the dietary overlap

between species, as indicated by the positive association between temperature/humidity

and the average pairwise dietary distance (figure 3A), and by the absence of association

between climatic conditions and the shortest dietary distances between species (figure

3B)

Although some clades are known to be important seed-dispersers across multiple

continents (some families of Passeriformes, Galliformes, Piciformes, etc. - Snow 1981,

Fleming et al. 1987, Kissling et al. 2009), a spatial analysis of those networks reinforces

the idea that such macroevolutionary sorting mechanism acts regionally on the available

pool of species, and that the role of available species is, as expected, geographically

structured. To evaluate if species within the same lineage play a similar role in different

networks, we calculated the average centrality value for each different lineage (either

family or genus) for all networks. The association between the mean centrality value of

each lineage in different networks decays as a function of the geographical distances

between those networks (figure 4). These results suggest that geographically close

networks tend to have similar lineages playing similar ecological roles, but those roles are

taken by different lineages in different parts of the planet.

In summary, our results provide evidence for a macroevolutionary sorting

mechanism (species selection in a broad sense - Jablonski 2008) on network assembly

where central species tend to belong to evolutionary lineages that are more stable over

deep time. Moreover, we found that those central species not only interact with a higher

number of species, but also that those partner species belong to a wider range of plant

families (figure 3C). Hence our results suggest potential bi-directional selective regimes

between ecological roles and diversification regimes, which might shape the fate of

groups of very distantly related lineages (birds and plants) linked through ecological

interactions.

Figure 1: Conceptual framework and data used - Panel A (top left): Conceptual representation of the study; Panel B (top right): Geographical and climatic distribution of the 29 networks used in the study. Panel C (bottom): Exemplar pruned phylogeny with branch colors indicating the bird orders, along with the extinction fraction (purple dashes and density), net diversification rates (green dashes and density) in the center, and the PCA centrality values (top row of panel C) for each of the species of all species used in the analyses. Some species were present in more than one ecological network, and this is information is displayed by points or bars of different colors in the PCA centrality graph.

Figure 2: Association between network centrality and macroevolutionary stability - A - exemplar association between PCA centrality values and extinction fraction and net diversification rates estimated from one phylogenetic tree. B - Posterior distributions of all parameters (except the intercept for a matter of scale) for the Bayesian generalized linear mixed model analysis. The points within the violin plot represent the median values taken from the analysis done for each different phylogenetic tree (that are highly correlated with the mean and the mode, since the distributions are highly symmetrical). Posterior distribution of slopes for Extinction fraction (green violin plot) and Net diversification rates (purple violin plot) are clearly off-centered from the slope value of zero in the expected direction (negative for extinction fraction; positive for net diversification), indicating that species belonging to macroevolutionary more stable lineages (low Epsilon and/or high Net Diversification) are more central in the networks.

143
144

Figure 3: Different axes of specialization of bird species analyzed in the study - A) Plot showing that the average pairwise distance of species along dietary PC1 increases towards warmer, wetter, less seasonal environments (climatic PC1, p-value = 0.001, R² = 0.4316). . B) Scatterplot between mean nearest-neighbor distance and the climatic PC1, highlighting the much smaller variation in comparison to panel A. We used the same scale from panel A on the y-axis to highlight the much lower, non-significant association (p-value = 0.1196, R² = 0.0535). C) Association between PCA centrality and the phylogenetic diversity of plant families with which each bird species interacts, indicating that more central species interact with more diverse plant families (p-value < $2.2*10^{-16}$, R² = 0.7033). On all panels continuous lines indicate statistically significant associations while dotted lines a lack of significant association.

145
146

Figure 4: Association between mean centrality per family (panels A and B) or genus (panels C and D) and geographical distance - Each point represents a pairwise correlation between ecological roles in two different networks, and the red lines represent a loess smoothing to show the general trend of the data. The grey areas represent the null models that were built by randomizing centrality values per species within networks (panels A and C), or by randomizing the identity of networks for each geographical distance (panels B and D). Regardless of the null model selected, the associations between mean centrality of families/genera is higher than expected by the null model for geographically close networks, and this similarity dissipates with distance, suggesting that there is no single group driving the association between macroevolutionary stability and ecological role.

Data and Code Availability

149 Bird phylogenies were obtained from birdtree.org (Jetz et al. 2012), and ecological
150 networks were obtained from http://dx.doi.org/10.5061/dryad.2br2b (Pigot et al. 2016).
151 All data and code used in this study is also available for download at
152 https://doi.org/10.5281/zenodo.3560680. Codes are also available at
153 https://www.github.com/gburin/macroevoNet.
154

Author Contributions

156 GB, PRG and TBQ designed the study; GB gathered and analyzed the data; GB and TBQ
157 wrote the first draft of the manuscript; GB, PRG and TBQ significantly contributed to the
158 last version of the manuscript.

Acknowledgements

We thank Eduardo Santos, Marco Melo, Marcus Aguiar, Patrícia Morelatto, Marilia
Gaiarsa, and Daniel Caetano for their insightful comments during several steps of this
work. All authors thank FAPESP (Fundação de Amparo à Pesquisa do Estado de São
Paulo) for funding (GB: grants #2014/03621-9 and #2018/04821-2; PRG: grant
#2018/14809-0, TBQ: grants #2012/04072-3 and #2018/05462-6)

Supplementary information is available for this paper

Correspondence and requests for materials should be addressed to gustavoburin@usp.br
or tbquental@usp.br

References

Bascompte, J., & Jordano, P. (2007). Plant-Animal Mutualistic Networks: The
Architecture of Biodiversity. Annual Review of Ecology, Evolution, and Systematics,
38(1), 567–593.
Burin, G., Kissling, W. D., Guimarães, P. R., Şekercioğlu, Ç. H., & Quental, T. B.
(2016). Omnivory in birds is a macroevolutionary sink. Nature Communications, 7,
11250. doi:10.1038/ncomms11250
Delmas, E., Besson, M., Brice, M.-H., Burkle, L. A., Dalla Riva, G. V., Fortin,
M.-J., Gravel, D., Guimarães Jr., P.R., Hembry, D.H., Newman, E.A., Olesen, J.M.,

Pires, M.M., Yeakel, J.D., Poisot, T. (2018). Analysing ecological networks of species
interactions. Biological Reviews of the Cambridge Philosophical Society.
Donatti, C.I., Guimarães, P.R., Galetti, M., Pizo, M.A., Marquitti, F.M. and Dirzo,
R., (2011). Analysis of a hyper☐diverse seed dispersal network: modularity and
underlying mechanisms. Ecology letters, 14(8), pp.773-781.
Eriksson, O. (2016) "Evolution of angiosperm seed disperser mutualisms: the
timing of origins and their consequences for coevolutionary interactions between
angiosperms and frugivores." Biol. Rev. Camb. Philos. Soc. 91, 168–186.
Fleming, T. H., Breitwisch, R. & Whitesides, G. H. (1987). Patterns of tropical
vertebrate frugivore diversity. Annual Review of Ecology and Systematics 18 , 91–109.
Fleming, T. H., Kress, W. J. (2013). "The Ornaments of Life: Coevolution and
Conservation in the Tropics. Interspecific Interactions". Chicago (Illinois): University of
Chicago Press. 588 p., ISBN: 978-0-226-25341-1.
Galetti, M., Guevara, R., Côrtes, M.C., Fadini, R., Von Matter, S., Leite, A.B.,
Labecca, F., Ribeiro, T., Carvalho, C.S., Collevatti, R.G., Pires, M.M., Guimarães, P.R.,
Brancalion, P.H., Ribeiro, M.C., Jordano, P., (2013). Functional extinction of birds drives
rapid evolutionary changes in seed size. Science 340, 1086–1090.
Gómez, J. M., & Verdú, M. (2012). Mutualism with plants drives primate
diversification. Systematic Biology, 61(4), 567–577.
Gómez, J. M., Verdú, M. & Perfectti, F. Ecological interactions are evolutionarily
conserved across the entire tree of life. Nature 465, 918–921 (2010).
Harmon, L. J., Andreazzi, C. S., Débarre, F., Drury, J., Goldberg, E. E., Martins,
A. B., Melián, C.J, Narwani, A., Nusimer, S.L., Pennell, M.W., Rudman, S.M, Seehause,
O., Silvestro, D., Weber, M., Matthews, B. (2019). Detecting the macroevolutionary
signal of species interactions. Journal of Evolutionary Biology, 32(8), 769–782.
Jablonski, D. (2008). Biotic interactions and macroevolution: extensions and
mismatches across scales and levels. Evolution, 62(4), 715–739.
Jordano, P. (2000). Fruits and frugivory. In The Ecology of Regeneration in Plant
Communities . Second Edition (ed. M. Fenner), pp. 125–166. CAB International,
Wallingford.
Kissling, W.D., Böhning-Gaese, K., & Jetz, W. (2009). The global distribution of
frugivory in birds. Global Ecology and Biogeography, 18(2), 150–162.
Nuismer, S. L. & Harmon, L. J. Predicting rates of interspecific interaction from
phylogenetic trees. Ecol. Lett. 18, 17–27 (2015).
Olesen, J. M., Bascompte, J., Elberling, H., & Jordano, P. (2008). Temporal
dynamics in a pollination network. Ecology, 89(6), 1573–1582.
Plein, M., Längsfeld, L., Neuschulz, E. L., Schultheiß , C., Ingmann, L., Töpfer,
T., Böhning-Gaese, K. & Schleuning, M. (2013). Constant properties of plant-frugivore
networks despite fluctuations in fruit and bird communities in space and time. Ecology 94
, 1296–1306.
Ponisio, L. C., Gaiarsa, M. P., & Kremen, C. (2017). Opportunistic attachment
assembles plant-pollinator networks. Ecology Letters, 20(10), 1261–1272.
Rezende, E. L., Lavabre, J. E., Guimarães, P. R., Jordano, P., & Bascompte, J.
(2007). Non-random coextinctions in phylogenetically structured mutualistic networks.
Nature, 448(7156), 925–928.

Roopnarine, P. D., Angielczyk, K. D., Weik, A. & Dineen, A. Ecological
persistence, incumbency and reorganization in the Karoo Basin during the Permian-
Triassic transition. Earth-Science Reviews 189, 244–263 (2018).
Snow, D. W. (1981). Tropical frugivorous birds and their food plants: a world
survey. Biotropica 13 , 1–14.
Thompson, J. N. The Geographic Mosaic of Coevolution (Univ. Chicago Press,
2005)
Weber, M. G., & Agrawal, A. A. (2014). Defense mutualisms enhance plant
diversification. Proceedings of the National Academy of Sciences of the United States of
America, 111(46), 16442–16447.
Yang, S., Albert, R., & Carlo, T. A. (2013). Transience and constancy of
interactions in a plant-frugivore network. Ecosphere, 4(12), art147.
240

241 Material and Methods

To investigate the potential effect of diversification rates on the assembly of

current ecological networks we first characterized each frugivore species with respect to

its potential ecological role in the network, and then estimated speciation and extinction

rates related to each bird lineage using a comprehensive molecular phylogeny containing

approximately 67% of all bird species (see below). We then tested for an association

between the network metric and the macroevolutionary rate (see figure 1 main text for a

cartoon version of the general approach) using phylogenetic generalized linear mixed

models (MCMCglmm - Hadfield 2010). Phylogenetic correction was used because the

network metrics showed significant phylogenetic signal (mean value for Pagel's lambda

= 0.537, although for each tree the corresponding lambda value was used).

252 Diversification Rates

To estimate speciation and extinction rates we used the most recent bird

phylogeny, which comprises the 99.3% of the extant species (9993 species in the

phylogeny from an estimated total of 10,064 species – (Jetz et al. 2012). To avoid any

potential bias introduced by adding species without DNA sequences (using a pure-birth

algorithm) on our estimates of speciation and extinction rates (Rabosky 2015), we used

the trees that contain only species for which molecular data were available from the

original phylogenetic paper (Jetz et al 2012). This tree had 6670 species, but species not

sampled were taken into account by the model. We analyzed the bird molecular

phylogeny using a Bayesian framework (BAMM v. 2.5; Rabosky 2014) to estimate the

rates of speciation and extinction while explicitly taking into account the sampling

fraction of species. The incompleteness (about 1/3 of species without DNA data) was

considered within BAMM by using the percentage of missing species within each clade

of the maximum clade credibility tree obtained by Jetz et al. (2012). Rabosky (2014) has

developed an algorithm that finds sub-trees within a phylogeny, which share speciation

and extinction rates through a Bayesian framework using reversible-jump Markov Chain

Monte Carlo sampling. Each diversification shift configuration has an associated

posterior probability, and each configuration can assign different diversification rates to a

particular species or clade. Since it is impossible to not know which is the true

configuration, we used the mean rates averaged by the marginal posterior probabilities of

each distinct shift configuration. Lastly, we calculated our rates of interest: the extinction

fraction ($\varepsilon = \mu/\lambda$) and net diversification rate ($r = \lambda - \mu$) for all species of interest. These

two rates represent two different metrics used to quantify the stability/predictability of a

given lineage. A higher value of extinction fraction suggests a more volatile dynamics

within a given lineage and hence a higher chance of the entire lineage going extinct. A

lower net diversification rate suggests that a lineage accumulates species as a smaller

pace, and therefore has a higher chance of going extinct than lineages with higher net

diversification rate. To our purposes, we assume that those two rates (each of them plus

their interaction) affects the likelihood that a given lineage might be evolutionary reliable

at long temporal scales.

Ecological Networks

To evaluate the ecological role of different species we used data from 34 different

frugivory networks compiled by previous authors (Pigot et al 2016). Those networks

comprise both temperate and tropical areas and a total of 546 unique bird species (838

total unique interactions). From the original dataset, we discarded 5 networks due to

biases in species composition (networks that focused on specific bird or plant groups, and

therefore do not represent the whole interacting assemblage), resulting in a final dataset

of 468 species (700 unique interactions). Although some networks (23) had information

on the strength of interactions, we decided to binarize all networks for two reasons. First,

it allows us to work with a larger dataset. Second, it is known that interaction strengths

vary across time (e.g. Ponisio et al. 2017), rendering quantitative estimates not

necessarily meaningful as a long-term description. We also removed pairs of species that

only interacted with each other, and were, by definition, disconnected from the rest of the

network. Those represent only 14 species from only 6 different networks from a total of

468 species from 29 networks. This was done because some of the network metrics are

not defined for networks that are not fully connected.

To characterize each species' ecological role within the compiled dataset we

investigated different node (species) properties of the networks by calculating, using the

original bipartite networks, three different centrality metrics: degree (number of

interactions), closeness (the reciprocal of the sum of shortest distances, in links,

connecting the focal node to all other nodes in the network) and Katz (a measure of the

distance, in terms of all possible pathways, between the focal node and all other nodes of

the network, using $\alpha = 0.05$). Such metrics provide us with information about the

importance of each species network structure when considering only the direct

interactions (degree), the shortest pathways (closeness) or all pathways (Katz). To be able

to combine the ecological roles estimated for different networks (that have different

properties, such as number of species and connectance) into a single analysis, we first

standardized the three metrics by calculating their z-scores for each network separately.

After compiling each of the three standardized centrality metrics for all species in all

networks in a single dataset, we performed a Principal Component Analysis (PCA) in

order to obtain a single score that comprised different aspects of the network position (as

a proxy for the ecological role). We used only the first Principal Component since it

explained 93.3% of the variance in centrality values.

Sources of uncertainties

We incorporated in our analyses two distinct sources of uncertainty. The first one

is related to the true evolutionary history of birds. To incorporate the phylogenetic

uncertainty, we used 112 different topologies from the bird phylogeny from Jetz et al.

2012 (56 of each backbone) to both estimate the rates of diversification and to take into

account the phylogenetic structure of the residuals in the linear model. The other source

of uncertainty came from the networks. From all the species in the final dataset (N = 468

species), 139 are part of more than one network (up to 6), making it impossible to

attribute a single centrality value to those species. We thus incorporated this variation as a

source of uncertainty in the regression analysis by using species identity as a random

factor as described in the modeling section below.

Co-Factors

Because the nature and relevance of ecological interactions might be different at
different parts of the planet (Mittelbach & Schemske 2015) we ran a global analysis (with
all networks combined) using climatic variables as co-factors. Those were: 1- annual
mean temperature; 2- temperature seasonality; 3- annual precipitation; 4- precipitation
seasonality. To summarize environmental conditions and avoid over-parameterization of
the models, we ran a PCA with the four climatic variables, and used the first and second
PCs that in combination explain 81.92% of total variance. We note that those two PCs
aligned well with the raw temperature and precipitation variables (Annual Mean
Temperature, Temperature Seasonality, and Annual Precipitation have scores of 0.619, -
0.592, and 0.492 on PC 1, respectively. Annual Precipitation and Precipitation
Seasonality have scores of 0.331 and -0.942 on PC 2, respectively - figure S5). The
climatic data were obtained from the WorldClim database version 2.0 (Fick & Hijmans,
2017).

To test if the macroevolutionary effect on network is a phenomenon that depends
on the absolute or the relative diversification rates (i.e. if rates of a given species are
higher or lower than other species within the same network), we ran each of the three
classes of models previously described using raw and standardized (z-scores computed to
within each network) macroevolutionary rates.

Modelling

We tested the association between ecological roles and macroevolutionary
dynamics using a Bayesian generalized linear mixed model (MCMCglmm), implemented
in the R package MCMCglmm (Hadfield, 2010). This framework allows us to naturally
incorporate both sources of uncertainty. By using species identity and the phylogenetic
structure as random factors, we can account for intraspecific variation in the ecological
roles on the estimation of the regression parameters while simultaneously controlling for
the phylogenetic structure. Moreover, by combining the posterior distributions of
parameter values using rates from different trees (topologies), we also can account for
phylogenetic uncertainty. We assessed the effect size of each variable used in the model
by analyzing the asymmetries in both the combined posterior distributions in relation to

zero (no association), and in the medians of individual posterior distributions for each

tree (Holland, 2019; Wasserstein et al. 2019). By doing this we embrace sources of

uncertainty incorporated in our analysis (ecological and phylogenetic) while looking at

the magnitude of the effect (Holland, 2019). All MCMC chains were run for 5000000

iterations, with half of the total length discarded as burn-in, and sampling every 2500

iterations. We used inversed Wishart distributions as priors for both the fixed and the

random effects.

Geographical patterns of centrality per family/genus

Lastly, to check if the species that play central/peripheral roles in different

networks show a spatial structure with respect to their taxonomic identity (e.g. belong to

the same family/genus), we calculated the association between the average centrality per

lineages (either family or genus) for each pair of networks using Spearman's correlations,

ρ. and then plotted ρ as a function of the distance between each pair of networks. We used

family or genus as our data points for two reasons. First, it is unlikely that the same

species would be present in geographically distant networks. Second, the

macroevolutionary effect we are measuring is a lineage effect (e.g. different species

represent the macroevolutionary regimes of different lineages). To build a null model, we

used two different randomization approaches and two different ways to deal with missing

data. Hence, we ended up with four different null models for each analysis, one set for the

family and another for the genus level.

In the first randomization approach (figure 4 panels A and C), we randomized the

species centrality values within each network prior to calculating the average centrality

for each lineage (either per family or per genus). The association was then estimated for

all pair of networks and plotted as a function of the geographical distance of each

network pair. In the second randomization approach (figure 4 panels B and D), instead of

randomizing the species centrality values within each network, we randomly draw two

networks irrespective of their geographical distance, repeating this procedure for all a set

of geographical distances equal to the one seen in the empirical comparison. Hence in our

first randomization approach the species identity is randomized but the geographical

distances between pairs of networks is kept as in the empirical dataset. In the second

randomization approach it is the geographical distance that is randomized while keeping

the original species identity.

In both randomization approaches, there were instances where a given lineage

was only present in one of the two comparing networks. Hence to calculate the

association between networks when using lineage (e.g. family or genus) average

centralities, we had to deal with the limitation that "valid" average centrality value for a

given lineage might not be estimated for some of the networks. To overcome this

limitation, we filled the missing values in two different ways. In the first, we simply

replaced the missing values in a given network by 0 (hereafter called "single 0 scenario").

By doing this we are strongly "breaking" any possible association, which means that only

very strong association would still remain significant after adding those zero values

(figure 4). The second approach consisted in replacing the average centrality of a lineage

(family or genus depending on the analysis) by 0 not only for the network where that

lineage was missing but in both networks ("full 0 scenario"). This was done in order to

"completely remove" the effect of that lineage from the analysis, while still allowing for

the association between the families/genera that are present in both networks to be

estimated (supp. figure 4).

The prediction of the null model is constructed by doing a LOESS smoothing to

each randomized dataset (1000 datasets in total). The empirical LOESS smoothing line is

then compared to the group of LOESS smoothing lines defining the null model and the

relationship is considered significant if the empirical line does not overlap with the null

model lines.

Ecospace Occupation

To analyze how ecospace is occupied in different climates, we used an

independent dataset of dietary information (EltonTraits v1.0, Wilman et al., 2014), and a

phylogenetic Principal Component Analysis (phyPCA, Garland et al., 1993, Revell 2012)

to obtain the axes of greater variation in the diet of all species present in our dataset. The

phyPCA was done using only the 468 species present in our network dataset. Using the

first Principal Component (PC1 - greatest loading of fruits), we calculated three different

disparity metrics for each network: amplitude (that indicates how broad the dietary niche

is for the whole network), average pairwise distance and mean nearest-neighbor distance

(that indicates, on average, how far each species is from the closest species in their

respective network). We tested the relationship between each of the three metrics with

climatic PC 1 (which explains about 56% of the variation) to assess any trends of

ecospace occupation along climatic gradients.

The degree of interaction generalization

For each bird species in each network, we calculated both the number of unique

families with which they interact, and also the plant phylogenetic diversity by summing

all branch lengths in the pruned phylogeny containing only the plant families with which

the species interact. In order to test if more central species are more generalist (either

defined as interacting with a higher number of plant families or by a higher value of plant

phylogenetic diversity of their plant partners) we fitted linear models to test for the

association between ecological role (PCA centrality) and both the number of families and

phylogenetic diversity.

References
Fick, S.E. and R.J. Hijmans, 2017. Worldclim 2: New 1-km spatial resolution
climate surfaces for global land areas. International Journal of Climatology.
Garland, T., Dickerman, A. W., Janis, C. M., & Jones, J. A. (1993). Phylogenetic
analysis of covariance by computer simulation. Systematic Biology, 42(3), 265.
Hadfield, J.D. (2010). MCMC Methods for Multi-Response Generalized Linear
Mixed Models: The MCMCglmm R Package. Journal of Statistical Software, 33(2), 1-22.
Holland, S. M. (2019). Estimation, not significance. Paleobiology, 1–6.
Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K., & Mooers, A. O. (2012). The
global diversity of birds in space and time. Nature, 491(7424), 444–448.
Mittelbach, G. G., & Schemske, D. W. (2015). Ecological and evolutionary
perspectives on community assembly. Trends in Ecology & Evolution, 30(5), 241–247.
Newman, M.E.J., (2010) "Networks: An Introduction", Oxford University Press,
Oxford, New York, 772p.
Pigot, A. L., Bregman, T., Sheard, C., Daly, B., Etienne, R. S., & Tobias, J. A.
(2016). Quantifying species contributions to ecosystem processes: a global assessment of
functional trait and phylogenetic metrics across avian seed-dispersal networks.
Proceedings. Biological Sciences / the Royal Society, 283(1844).
doi:10.1098/rspb.2016.1597
Ponisio, L. C., Gaiarsa, M. P., & Kremen, C. (2017). Opportunistic attachment
assembles plant-pollinator networks. Ecology Letters, 20(10), 1261–1272.
Rabosky, D. L. (2014). Automatic detection of key innovations, rate shifts, and
diversity-dependence on phylogenetic trees. Plos One, 9(2), e89543.
doi:10.1371/journal.pone.0089543
Rabosky, D. L. (2015). No substitute for real data: A cautionary note on the use of
phylogenies from birth-death polytomy resolvers for downstream comparative analyses.
Evolution, 69(12), 3207–3216.

Revell, L. J. (2012). phytools: an R package for phylogenetic comparative biology
(and other things). Methods in Ecology and Evolution, 3(2), 217–223.
Wasserstein, R. L., Schirm, A. L., & Lazar, N. A. (2019). Moving to a world
beyond "p < 0.05." The American Statistician, 73(sup1), 1–19.
Wilman, H., Belmaker, J., Simpson, J., de la Rosa, C., Rivadeneira, M. M., &
Jetz, W. (2014). EltonTraits 1.0: Species-level foraging attributes of the world's birds and
mammals. Ecology, 95(7), 2027–2027.
465