2   **The application of community ecology theory to co-infections in wildlife hosts**

4   Chloe Ramsay[1*] and Jason R. Rohr[1]

[1]University of Notre Dame, Department of Biological Sciences Notre Dame, IN 46556, USA

*- Corresponding author:
Chloe Ramsay
Department of Biological Sciences
100 Galvin Life Science Center
University of Notre Dame, IN 46556, USA
email: cramsay@nd.edu

Abstract

Priority effect theory, a foundational concept from community ecology, states that the

order and timing of species arrival during species assembly can affect species composition.

Although this theory has been applied to co-infecting parasite species, it has almost always

been with a single time lag between co-infecting parasites. Thus, how the timing of parasite

species arrival affects co-infections and disease remains poorly understood. To address this

gap in the literature, we exposed post-metamorphic Cuban tree frogs (*Osteopilus*

septentrionalis) to Ranavirus, the fungus *Batrachochytrium dendrobatidis* (Bd), a nematode

Aplectana hamatospicula, or pairs of these parasites either simultaneously or sequentially at a

range of time lags and quantified load of the secondary parasite and host growth, survival and

parasite tolerance. Prior exposure to Bd or *A. hamatospicula* significantly increased viral

loads relative to hosts singly infected with Ranavirus, whereas *A. hamatospicula* loads in

hosts were higher when co-exposed to Bd than when co-exposed to Ranavirus. There was a

significant positive relationship between time since Ranavirus infection and Bd load, and

prior exposure to *A. hamatospicula* decreased Bd loads compared to simultaneous co-

infection with these parasites. Infections with Bd and Ranavirus either singly or in co-

infections decreased host growth and survival. This research reveals that time lags between

co-infections can affect parasite loads, in line with priority effects theory. As co-infections in

the field are unlikely to be simultaneous, an understanding of when co-infections are

impacted by time lags between parasite exposures may play a major role in controlling

problematic co-infections.

Keywords: priority effects, amphibians, co-infection, *Batrachochytrium dendrobatidis,*

Ranavirus, *Aplectana hamatospicula*

Introduction

Simultaneous infection with multiple species of parasites, hereafter referred to as co-

infection, is a regular occurrence for host species (Greub et al. 2000, Kim et al. 2003,

Swanson 2006). Direct interactions among these parasites can occur when each occupies

similar habitats in hosts, competing for space or resources and, in some cases, they can even

depredate one another (Kuris and Lafferty 1994, de Roode et al. 2004). Indirect interactions

of parasites can also occur. These can be facilitative or inhibitory and are often mediated by

the host immune system. As an example, microparasites generally activate the T helper 1

(Th1) arm of the adaptive immune system (fungal infections can also activate Th17; Berger

2000, Blanco and Garcia 2008, Zelante et al. 2009), whereas macroparasites activate the Th2

arm of the immune system (Paul and Zhu 2010). Because an investment in one arm generally

comes at the expense of investment in the other (i.e. a tradeoff; Abbas et al. 1996, Morel and

Oriss 1998, Yazdanbakhsh et al. 2002), micro-marcroparasite co-infections are often

facilitative (Abbas et al. 1996, Berger 2000), whereas micro-microparasite and macro-

macroparasite co-infections can be inhibitory though cross-immunity (Johnson and Buller

2011).

Interactions among parasites within hosts can affect the ability of hosts to tolerate

infections and thus can have important consequences on disease and host fitness and

population growth rates (Corbett et al. 2003, Bromenshenk et al. 2010, Ezenwa and Jolles

2015). For example, co-infections with multiple virus strains and microsporidia were detected

in honeybee hives suffering from colony collapse disorder (Bromenshenk et al. 2010).

Deworming a population of African buffalo increased the transmission of the deadlier

Mycobacterium bovis (Ezenwa and Jolles 2015). Mice co-infected with influenza virus and

Legionella pneumophila were less tolerant of infection than singly infected mice, increasing

their mortality (Jamieson et al. 2013).

Understanding how co-infecting parasites interact with one another and their host

could be informed by community ecology theory which highlights interactions among species

and their environment (Pedersen and Fenton 2007, Johnson et al. 2015). Priority effects in

ecology describe how the order and timing of species arrival during species assembly affect

species composition (Sutherland 1974, Drake 1991, Leopold et al. 2017). In community

ecology, priority effects have been shown to affect predator-prey interactions (Blaustein and

Margalit 1996), competition (Wilbur and Alford 1985, Sousa 1992), and community

succession (Facelli and Facelli 1993, Devevey et al. 2015, Leopold et al. 2017). Priority

effects have also been shown to be important in parasite-parasite interactions (Sousa 1992, de

Roode et al. 2005, Hoverman et al. 2013, Devevey et al. 2015). For example, relative to

simultaneous co-infections, infection with one strain of *Plasmodium chabaudi*, a causative

agent of malaria, reduced infections from a second strain of the pathogen when challenged

three days after the first (de Roode et al. 2005).

The definition of priority effects encompasses both order and timing of species arrival

(Sutherland 1974, Drake 1991, Leopold et al. 2017). However, most studies applying priority

effect theory in co-infecting parasite species have focused exclusively on the order and thus

have had a single time lag between co-infecting parasites (Hoverman et al. 2013, Devevey et

al. 2015, Leopold et al. 2017). Thus, how the timing of parasite species arrival affecting co-

infections and disease remains poorly understood (but see de Roode et al. 2005, Wuerthner et

al. 2017 for three time points). This is an important gap in the literature because co-infections

are almost always staggered in nature, making it important to understand how time lags

between co-infections affect disease. In community ecology, the initial population size of

interacting species can affect the outcome of their competition, with larger initial populations

being more likely to outcompete and potentially competitively exclude an introduced second

species (Chesson and Kuang 2008). Therefore, for microparasites that internally replicate and

are not easily cleared, longer time lags should allow the first parasite to reach higher densities

in the host before a second parasite attempts to infect (Rojas et al. 2005, Rosenblum et al.

2012).

The outcomes of co-infections are particularly important for amphibians because

amphibians are suffering from worldwide declines and extinctions due in large part to the

spread of several diseases. As a result, amphibians have become the most threatened

vertebrate taxon on the planet, with 32.5% of species threatened globally (Stuart et al. 2004,

Wake and Vredenburg 2008, Scheele et al. 2019). The primary goals of this study are to test

how a co-infecting parasite and the time lag between co-infections affects the secondary

parasite and amphibian host fitness (growth and survival) and whether these interactions are

mediated by host immune responses (indirect effects). To address these goals, we exposed

Cuban treefrogs (*Osteopilus septentrionalis*) to all pairwise combinations of a fungus

(microparasite), virus (microparasite), and nematode (macroparasite). The fungus we studied

is *Batrachochytrium dendrobatidis* (Bd), a chytrid fungus associated with hundreds of

amphibian declines and extinctions globally (Berger et al. 1998, Skerratt et al. 2007,

Kilpatrick et al. 2010). This fungus causes the disease chytridiomycosis, which leads to

cardiac arrest by fusing keratinous portions of amphibian skin and thus, limiting electrolyte

transport (Voyles et al. 2009, Kilpatrick et al. 2010). The virus we studied is frog virus 3

(FV3), a Ranavirus from the family *Iridoviridae* that replicates in the internal organs of

ectothermic vertebrates (Chinchar 2002), causing hemorrhaging and mass mortality (Gantress

et al. 2003, Gray et al. 2009). FV3 is also associated with mass die-offs of amphibians across

the planet (Green et al. 2002, Gray et al. 2009). In addition to these deadly microparasites,

treefrogs were also exposed to a macroparasite, specifically the gastrointestinal nematode,

Aplectana hamatospicula. This nematode is a common parasite of amphibians in the

Southeastern US and Latin America (Baker 1987, Vhora and Bolek 2013, Ortega et al. 2015).

Importantly, parasitic worms, Bd, and Ranaviruses are known to co-occur in amphibian hosts

in the field (Hoverman et al. 2012, Stutz et al. 2018, Watters et al. 2018). In the Peruvian

Andes, an amphibian biodiversity hotspot, up to 50% of frogs were coinfected with Bd and

Ranavirus (Warne et al. 2016).

Our goals were motivated by several hypotheses. First, the three focal parasites occupy

different microhabitats within the host; Bd infects the skin, *A. hamatospicula* dwells in the

gastrointestinal tract, and Ranavirus primarily infects the kidneys and liver(Gray et al. 2009,

Voyles et al. 2009, Knutie et al. 2017). Thus, we hypothesized that these parasites would

primarily interact indirectly, mediated by host immunity. Second, because microparasites

activate the Th1 arm and macroparasites activate the Th2 arm of the host immune system

(Abbas et al. 1996) and investment in each arm is mutually exclusive, we hypothesized that

co-infections that involve any microparasite (Bd and Ranavirus) and the macroparasite (*A.*

hamatospicula) would be facilitative. In contrast, we hypothesized that Bd and Ranavirus co-

infections might be inhibitory because they activate the same arm of the immune system.

Third, we hypothesized that longer time lags between initial and secondary infection would

increase the strength of co-infection priority effects for micro- but not macroparasites. This is

because microparasites, but not macroparasites, can replicate in the host and thus greater time

lags would allow for greater time for microparasite densities to increase. In this study, a

priority effect is defined as any significant relationship between the load of the second

parasite and the time lag between exposures to the co-infecting parasites. Fourth, from a host

health perspective, we hypothesized that host growth and survival would be lower if the

initial parasite *i*) was particularly virulent relative to the other co-infecting parasite and *ii*) had

a long time to replicate within the host, reaching high parasite loads (assuming it was a

microparasite). Fifth, when co-infecting parasites inhibit one another (either directly or

through the host immune system), we expected co-infection to increase host tolerance (e.g. a

host's ability to maintain high health despite high infection loads), whereas when co-infecting

parasites were facilitative of one another, we hypothesized co-infection to decrease host

tolerance.

158   **Methods**

Animal Husbandry

Mature Cuban treefrogs (*Osteopilus septentrionalis*) were collected from PVC pipes

(7.62 x 91.44 cm) hammered into the ground upright around wetlands in Flatwoods

Wilderness Park, Tampa, Florida (28.11° N, 82.31° W) from February to March 2016.

Individuals were housed separately in 57.75 cm^3 sterilized plastic cups containing an

unbleached paper towel wet with artificial spring water (ASW). Tree frogs were fed 2-week-

old crickets *ad libitum* twice a week at 25 °C (12hr photoperiod). Each of these plastic cups

was washed by hand, bleached in a 10% solution, thoroughly rinsed with deionized (DI)

water, and dried weekly.

Pre-inoculation

To eliminate any potential helminth infections in wild-caught frogs, collected

individuals were orally treated with ivermectin (0.2 mg/kg of host) two months before the

experiment. Following ivermectin exposures, husbandry containers were changed daily for

one week to reduce the potential for reinfection with any larval *A. hamatospicula*. Once frogs

did not shed helminths for four weeks, they were considered uninfected. Amphibian species

from Flatwoods Wilderness Park have not been found to carry Ranavirus or Bd and none of

the collected individuals showed symptoms of infection before the start of the experiment.

Two days before initial parasite exposure, frogs were moved to an environmental chamber to

acclimate to experimental conditions. The environmental chamber was kept at 17 °C (12h

photoperiod) because Cuban treefrogs can clear Bd at higher temperatures (McMahon et al.

2014, Cohen et al. 2017).

Parasite culture

A. hamatospicula were obtained from the gastrointestinal tract of euthanized Cuban

treefrogs collected from Flatwoods Wilderness Park. Extracted worms were stored in a Petri

dish (15 x 150 mm; $n \leq 5$ per dish) filled with 2mL of DI water and held under dark

conditions. We collected larval worms (J1) from adults and held each for ≥ 14 days to ensure

maturation to the infectious stage (*i.e.*, J3). Gastrointestinal (GI) content was collected from

uninfected frogs and stored identically to act as a sham inoculant.

Ranavirus (FV3) was cultured in fathead minnow cells (*Pimephales promelas*) and

stored in minimal essential media (MEM) at -80 °C.

Bd (strain SRS-JEL 212) solution was grown on 1% tryptone-agar Petri dishes,

aliquoted from an active culture. Bd growth occurred over eight days at 23°C until zoospores

were observed. Petri dishes were then flooded with 1 mL of DI water for five minutes to

permit zoosporangia to release infectious zoospores. Released zoospores and DI water were

then poured into a beaker with 15 mL of DI water and stirred to produce a homogeneous

slurry. To assess Bd zoospore concentration, we sampled the homogeneous slurry and

counted zoospore density using a hemocytometer. We added DI water to the slurry to obtain a

final Bd concentration of 10^5 zoospores/ml. Sterile 1% tryptone plates without Bd were

cultured at 23°C for eight days and rinsed with DI water to produce sham Bd and control

treatments.

Experimental Design

To examine how the timing of initial parasite exposures affects host parasite

dynamics, we exposed Cuban treefrogs to one of ten parasite treatments. Parasite treatments

included exposure to *A. hamatospicula* alone*,* Ranavirus alone, or Bd alone, exposure to

pairwise combinations of two parasites in sequential order (six co-infection treatments), and a

sham exposure (control; see below for details). Each of the six co-infection treatments had

five temporal subtreatments of staggered exposures (Table 1). Five frogs were assigned to

each of the 34 treatment combinations for a total of 170 experimental individuals.

Timing of parasite exposure

For the six co-infection treatments, timing of the first parasite exposure relative to the

second parasite was based on the generation time of the first parasite (Table 1). *A.*

hamatospicula takes approximately 11 days to reach the GI tract after initial infection, about

19 days to acquire antibodies, and 24 days to produce offspring (Knutie et al. 2017).

Therefore, frogs were first exposed to *hamatospicula* at 24, 19, 11, 1, or 0 days prior to a

second parasite exposure. Bd has a generation time of ~4 days at room temperature, with

growth leveling off over time (Piotrowski et al. 2004). Based on our experience with Bd

infection in Cuban treefrogs, Bd-induced mortality generally begins about three weeks after

exposure. Therefore, to avoid substantial mortality, frogs were exposed to Bd at 16, 12, 8, 4,

or 0 days prior to a second parasite exposure. Finally, Ranavirus replicates rapidly, causing

mortality in as few as two days for highly susceptible frog species, but can also be cleared in

one to two weeks in other frog species (Green et al. 2002, Gantress et al. 2003).

Consequently, frogs were first exposed to Ranavirus 12, 8, 4, 2, or 0 days prior to a second

parasite exposure. Time points were chosen to simulate natural points in disease progression

for each parasite (Gray et al. 2009, Voyles et al. 2009, Knutie et al. 2017). Longer time lags

between parasites were chosen so that the second parasite exposure occurred when the host

was recovering from or in the established stages of infection. Shorter time lags for each

parasite were chosen so that the second parasite exposure occurred when the host was in the

beginning stages of combating the initial parasite.

Parasite exposures

Depending on treatment, frogs in Petri dishes (25 x 100 mm) were exposed to a 1-ml

aliquot of DI water containing either 30 J3 *A. hamatospicula* larvae, a 1-ml aliquot of DI

water containing 10^5 Bd zoospores, and/or a 77-µl aliquot of MEM containing 10^5 plaque

forming units (PFU) of Ranavirus. Sham exposures consisted of the same volume of liquid

without the infectious agent for all parasites (1-ml DI water for *A. hamatospicula* and Bd

and/or 77-µl of MEM for Ranavirus). All hosts received sham exposures for parasites they

were not exposed to. For example, a host singly infected with Bd received sham exposures

for Ranavirus and *A. hamatospicula*. Control hosts received sham treatments for all three

parasites on day zero.

Amphibian responses to parasites

To measure the growth of amphibian hosts over the duration of the experiment, 24 h

after secondary parasite exposure, treefrogs were weighed and snout-vent length (SVL) was

measured and these measurements were taken weekly for four weeks. Survival was assessed

twice daily. At the time of death, we weighed deceased individuals, measured SVL, and

assessed infection for the secondary parasite to which the individual was exposed. For Bd-

and Ranavirus-exposed individuals, we swabbed the skin and conducted qPCR (see below).

For *A. hamatospicula* exposed individuals, we conducted dissections of the gut. Twenty-eight

days after exposure to the secondary parasite, all surviving frogs were euthanized and

dissected. Host growth rate was calculated as the final mas or snout-vent length (SVL) minus

initial mass or SVL divided by weeks alive (g wk^{-1}).

Assessing parasite loads

To assess *A. hamatospicula* infection, treefrog feces were checked weekly for larval

worms and amphibian hosts were dissected at time of morality and at the end of the

experiment to quantify adult *A. hamatospicula* infection load in the gastrointestinal tract. To

assess Bd and/or viral load, amphibians were swabbed 2, 4, 8, 16, and 24 days after exposure

to the second parasite and at the time of mortality. Individuals were swabbed 5 times from

hip to toe on both rear legs for Bd and around the cloaca and mouth to test for Ranaviral

infection. Swabs were placed in 2-ml sterile microcentrifuge tubes and stored at -80 °C until

processing.

257   Given the generation time of Ranavirus and Bd, DNA was extracted (Qiagen DNEasy

258   Blood & Tissue Kit) from 2-, 4, and 8-day swabs where Ranavirus was the second infecting

259   parasite and from day 8-, 16-, and 24-day swabs where Bd was the second infecting parasite.

260   Bd and Ranavirus swabs were analyzed using quantitative polymerase chain reaction (qPCR)

261   to quantify parasite load (Boyle et al. 2004, Picco et al. 2007).

262   *Assessing amphibian immune response*

263   We conducted a separate experiment that included 48 additional Cuban tree frogs to

quantify amphibian antibody responses to single and simultaneous co-infections with the

parasites that we tested above. Four individuals were tested for each of the three single

infection treatments, for a total of twelve singly infected individuals. Eight individuals were

tested for each of the three, pairwise, co-infection treatments and half of the individuals in

each pairwise co-infection treatment were tested for one parasite and the other half were

tested for the other. Finally, twelve individuals were tested as controls, with four individuals

tested to compare with each parasite (Table S3). To measure the immune response for each

parasite, frogs were euthanized as described above and blood was extracted from frogs with

microcapillary tubes 11, 19, and 28 days after exposure to Ranavirus, *A. hamatospicula*, and

Bd, respectively. These times were chosen to reflect when antibody immune responses to

each parasite should be detectable (Gantress et al. 2003, Ramsey et al. 2010, Knutie et al.

2017).

276   Blood plasma was used in enzyme-linked immunosorbent assays (ELISA) to detect

IgY antibody presence (adapted from Knutie et al. 2017). Individual frog serum was first

diluted at 1:100 in carbonate coating buffer (0.05 M, pH 9.60) and then added to ninety-six

well plates with 100 μL/well. Each individual sample was run in triplicate. After serum was

added, plates were incubated at 4 °C overnight. Plates were washed five times with 300

μL/well of a Tris-buffered saline wash solution. The plates were then filled with 200 μL/well

of bovine serum albumin (BSA) blocking buffer and incubated for 2 h at room temperature

on an orbital table. Plates were washed again five times, loaded with 100 μL/well of primary

detection antibody (Goat-αAlligator-IgG, diluted 1:1000; Bethyl), placed on an orbital table

for 1 h, washed again (five times), loaded with 100 μL/well of secondary detection antibody

(Rabbit-αGoat-IgG, diluted 1:5000; Bethyl) for 1 h, and then washed for the final time (five

times). Plates were then loaded with 100 μL/well of tetramethylbenzidine (TMB: Bethyl

Laboratories) and incubated for 30 mins before the reaction was stopped with 100 μL/well of

stop solution (Bethyl Laboratories). Finally, optical density (OD) of the wells was measured

with a spectrophotometer (BioTek, PowerWave HT, 450-nanometer filter).

Statistical Analyses

All analyses were run with R version 3.4.2 (R Core Team 2017). Plots were created

using the *visreg* package and function (Breheny and Burchett 2019).

Goal 1: Do prior infections and the timing of those infections change the abundance

of the secondary parasite?

Importantly, there are two time variables used for Bd and Ranavirus in the statistical

analysis. First, loads of Bd and Ranavirus were measured over time after co-infection or

single infection. This measurement is not present for *A. hamatospicula* because loads of *A.*

hamatospicula were only quantified through destructive sampling at the end of the

experiment. Second, there is the time lag between the first and second infection in the co-

infection situations. This measurement occurred for all parasites tested.

To test how the interaction between initial parasite identity and time since co-

exposure (the amount of time after co-infection) changed parasite growth, we compared the

load of the secondary parasite to load of the same parasite species in a single infection

treatment through time. These analyses were only done for Bd and Ranavirus because *A.*

hamatospicula load data was only collected, at the end of the experiment. We conducted a

general linear mixed model (lmer) with the load of the secondary parasite as the dependent

variable, initial parasite identity (none if single infection) and swabbing time points (time

since co-exposure) as interacting independent variables, and individual as a random effect.

For Bd, the error distribution was negative binomial because it was counts with no upper

bound. Viral loads were also counts with no upper bound, but the range of viral loads were

too large for the negative binomial model to converge; so, we were forced to use a log-

normal model. Tukey's post-hoc tests were run to compare parasite treatments of co-infecting

parasites to one another and to single infections (eg. Bd only, Ranavirus x Bd, *A.*

hamatospicula x Bd). All Tukey's post-hoc tests were run using the *multcomp* package and

glht function (Hothorn 2010).

To test the effects of the identity of the initial parasite and time lags between parasite

exposures (*i.e.* the time gap between co-infecting parasites) on secondary parasite loads

(loads measured on day 8 for Bd, day 2 for Ranavirus, day 24 for *A. hamatospicula*), we

conducted a generalized linear model (glm) with secondary parasite loads (Bd or Ranavirus)

on the first day the secondary parasite load was measured as the negative binomially

distributed dependent variable and the identity of the initial parasite and time since initial

parasite exposure as interacting independent variables. Zero-inflated negative binomial and

Poisson error distributions were also compared, but the negative binomial error distribution

was found to have the lowest AIC. For *A. hamatospicula,* we used a logistic regression (*glm*

function) with the number of larvae (second infection) that successfully infected (penetrated

the host, reached the gut, and reached maturity), out of the total number of larvae to which

hosts were exposed, as the dependent variable. Time since initial parasite exposure and the

identity of the initial parasite were crossed independent variables. Time since prior exposure

in co-infection treatments was a proportion (values between 0 and 1). To calculate this

proportion time since exposure for a given parasite treatment was divided by the maximum

time since initial exposure possible for each of the parasites (*e.g.* 24, 16, and 12 for *A.*

hamatospicula, Bd, and Ranavirus respectively). This allows for comparisons among the

various parasites even though the timing of initial exposures were different. We also tested

how timing of prior exposure altered parasite growth rate (see supplemental methods and

results).

Goal 2: Do prior infections and the timing of those infections affect host growth, survival,

and tolerance?

To test how the timing of initial infections alters host growth rate, we conducted a

general linear model with amphibian growth rate (mass and SVL) over four weeks ([$mass_{final}$

- $mass_{initial}$] /4 weeks) as the dependent variable. Time since initial parasite exposure, the

identity of the initial parasite, and the identity of the second (co-infecting) parasite were

interacting independent variables and starting weight was a covariate to control for the

different host sizes at the start of the experiment. Tukey post-hoc tests were run to compare

initial parasite identities to one another and second parasite identities to one another.

Dunnett's tests, using the *multcomp* package and *glht* function (Hothorn 2010), were used to

examine if the time lags between the parasite exposures matter when comparing each co-

infection time lag to the single infection.

To investigate how the timing of initial infections alters host survival, we conducted a

survival analysis using the *survival* package and the *coxph* function (Therneau and Lumley

2019). Individuals that survived to the end of the experiment were right-censored. We

included the time since initial exposure, the identity of the initial parasite, and the identity of

the second (co-infecting) parasite as interacting independent variables.

To test how the timing of the initial infection and the identity of the co-infecting

parasite alters amphibian tolerance (maintaining survival or growth with high parasite loads),

we re-conducted the previously described growth and survival analyses with secondary (co-

infecting) parasite load (loads measured at the 1st time point; log transformed for Bd or

Ranavirus, not log transformed for *A. hamatospicula*) as an additional crossed independent

variable.

Goal 3: Do co-infections affect host immunity?

To test how co-infections vs. single infections affects host immunity, we conducted a

general linear model with antibody levels as the dependent variable, and single vs. co-

infection treatments as the independent variable. To test how treatment group (*e.g.* controls,

Bd, Bd and Ranavirus, etc.) affects host immunity, we conducted a general linear model with

antibody levels as the dependent variable and treatment group as the independent variable.

Given that Bd is known to be immunosuppressive (Fites et al. 2014), we further tested how

Bd impacts host immunity by testing how exposure to Bd alters antibody levels compared to

parasites that are not known to be immunosuppressive. We conducted a general linear model

(lm) with Bd exposure (*e.g.* bd exposed vs. exposed to other parasites vs. controls) as the

independent variable and antibody levels as the dependent variable. Mass of individuals and

timing of blood removal were initially used as covariates for all the above analyses, but were

excluded in final analyses, as they were not significant.

Results

Goal 1: Do prior infections and the timing of those infections change the abundance of the

secondary parasite?

Importantly, there are two time variables for Bd and Ranavirus that we cover in this

Results section. There is a measurement of time lag between the initial and secondary

infection and there is the change in loads of Bd and Ranavirus through time after exposures

for both singly and co-infected hosts.

Ranavirus as the secondary parasite and response variable

We did not find evidence for priority effects when Ranavirus was the secondary

parasite, as time lags between parasite exposures did not affect viral loads (Time lag between

co-infections*:* $F_{1,44}=1.738$, $p=0.194$). However, we did find effects of co-infections on the

change in viral loads through time. More specifically, we found that the identity of the first

infecting parasite ($\chi^2_2=17.2$, $p<0.001$), time since hosts were co-infected ($\chi^2_2= 77.65$,

$p<0.001$), and an interaction between these two factors (Treatment x day: $\chi^2_4 = 12.12$,

$p=0.017$) affected viral load. Previous infection with *A. hamatospicula* or Bd increased host

viral loads relative to loads in hosts exposed to Ranavirus only (Ranavirus vs. Bd and

Ranavirus*: p=0.003*; Ranavirus vs. *A. hamatospicula* and Ranavirus: $p=0.034$) and this

response was independent of the time since co-exposure for *A. hamatospicula* but increased

with time since co-exposure for Bd (Parasite identity: $\chi^2_2=17.2$, $p<0.001$; time: $\chi^2_2= 77.65$,

$p<0.001$; parasite identity x time: $\chi^2_4 = 12.12$, $p=0.017$; Fig. 1).

Aplectana hamatospicula *as the secondary parasite and response*

We did not find evidence for priority effects when *A. hamatospicula* was the

secondary parasite, as time lags between parasite exposures did not affect *A. hamatospicula*

loads (Time lag: $\chi^2_1:0.073$, $p=0.78$). However, we did find effects of co-infection on *A.*

hamatospicula loads. Initial exposure to Bd, regardless of the time lags between co-

exposures, was associated with a higher load of *A. hamatospicula* in hosts compared to hosts

initially exposed to Ranavirus (Parasite identity: χ^2_1=9.563, *p*=0.008; Bd and *A.*

hamatospicula vs. Ranavirus and *A. hamatospicula*: *p*=0.037; Fig. 2B). *A. hamatospicula*

loads in co-infected hosts were not different from loads of hosts infected with *A.*

hamatopicula only (Ranavirus and *A. hamatopicula* vs. *A. hamatopicula*: *p*=0.752; Bd and *A.*

hamatopicula vs. *A. hamatopicula*: *p*=0.191; Fig. 2B).

Batrachochytrium dendrobatidis *as the secondary parasite and response*

There was evidence for priority effects when Bd was the secondary parasite (Parasite

identity x time lag: χ^2_1:6.78, *p*=0.009). When *A. hamatospicula was* the first parasite, time

lags between parasite exposures and Bd loads at 8 d were correlated negatively (*Z*=4.09,

p<0.001). In contrast, when Ranavirus was the first parasite, time lags between parasite

exposures and Bd loads at 8 d were correlated positively (*Z*=-1.89 *p*=0.058; Fig. 2A). When

looking at loads of Bd through time since co-infection, loads of Bd were positively associated

with time (Swab day: χ^2_1=173.78, *p*<0.001), but these trends were not affected by co-

infection or the identity of the co-infecting parasite (Swab day x treatment*:* χ^2_2: 1.65,

p=0.439).

Goal 2: Do prior infections and the timing of those infections affect host growth, survival,

and tolerance?

Growth as the response

Time lags between parasite exposures were negatively associated with host growth

based on SVL (mm/week) when both Bd and Ranavirus were the initial parasites, regardless

of whether these hosts were infected singly or co-infected ($F_{1,49}$=5.00, *p*=0.03; $F_{1,45}$=6.617,

p=0.013, respectively; Fig. S1). The growth rates of co-infected (at all initial infection time

points) and singly infected hosts did not differ significantly at any time since exposure (Fig.

S1; Table S1). These effects were similar when growth rate was calculated based on weight

(see supplemental results).

Survival as the response

Time lags between parasite exposures did not affect the survival of hosts (χ^2_1=1.72,

p= 0.19). However, when hosts were infected initially with Bd or Ranavirus hosts had a

significantly lower survival rate than hosts initially infected with *A. hamatospicula* (Identity

of first parasite: χ^2_2=22.92, *p*<0.001; Bd vs. *A. hamatospicula*: *p*=0.003; *Ranavirus vs. A.*

hamatospicula: p=0.043; Figure 5). Bd, Ranavirus, and *A. hamatospicula* did not

significantly affect host survival if they were the second infecting parasite (Identity of the

second parasite: χ^2_2=2.85, *p*=0.24).

Tolerance as a response

Time lags between parasite exposures did not alter host tolerance (the slope between

load and growth rate [g/week] or survival) to secondary infections (Viral load x time lag:

Host Growth: $F_{1,35}$=0.19, *p*=0.665; Host survival: χ^2_1=1.18, *p*=0.28; see supplement for

tolerance to Bd and *A. hamatospicula*). However, the identity of the initial parasite did impact

host tolerance to the secondary parasite. Previous infection with *A. hamatospicula* reduced

host tolerance (lower growth rate [g/week] and lower survival at higher loads) of Ranavirus

infections more than did previous Bd infections (Viral load x identity of second infecting

parasite: Host growth: $F_{1,35}$= 6.033, *p*<0.02; Fig. 3; Host survival: χ^2_1=6.98, *p*=0.008). In fact,

hosts that were exposed to Bd and then Ranavirus were hyper-tolerant of Ranavirus as growth

rates were positively correlated with viral loads (Fig. 3). Despite this, hosts infected with Bd

before Ranavirus were not significantly more tolerant of Ranavirus infections than hosts

singly infected with Ranavirus (Co-infection with Bd x viral load: $F_{1,22}$=0.4, *p*=0.8). Co-

infection and timing of co-infections did not change host tolerance to Bd, *A. hamatospicula*

(host growth or survival), or Ranavirus (survival; see table S2).

Goal 3: Do co-infections affect host immunity?

The antibody response of hosts was not altered by co-infections when compared to

single infections or controls (Number of infections: $F_{2,39}$=1.98, *p*=0.152) and did not differ

significantly when comparing all treatment groups (Treatment group: $F_{6,35}$=1.43, *p*=0.232;

Fig. S2). Bd is well documented to be immunosuppressive (Fites et al. 2014), whereas the

other two parasites are not. Thus, we hypothesized that Ranavirus and *A. hamatospicula*

would elevate antibodies relative to controls, but that Bd might not. This hypothesis was

supported as exposure to Ranavirus and *A. hamatospicula* (grouping individuals co-infected

and singly infected with each parasite) increased antibody levels relative to control animals

(Parasite identity: $F_{2,39}$=4.15, *p*=0.024; Ranavirus and *A. hamatospicula* vs. control:

p=0.024), but Bd did not *(*Bd vs. control: *p*=0.709; Fig. 5).

465   **Discussion**

Organisms in the wild are regularly co-infected with several species of parasites.

Furthermore, these co-infections are likely to be staggered and thus it is important to

understand if the time lags between these parasite exposures affect disease progression and

host health, a primary goal of this work. Our study found that the identity of the co-infecting

parasite, and the time lag between co-infecting parasites can alter disease progression in

hosts. (Table 2).

Hosts co-infected with *A. hamatospicula* and Ranavirus, had increased viral loads and

decreased *A. hamatospicula* loads relative to single infections, regardless of the timing of

those infections (Fig. 1,2). Therefore, for this co-infection pairing, we found evidence of

parasite interactions, but not priority effects. One of the potential drivers for these

interactions could be in-direct interactions mediated by the host immune system. In many

vertebrate species, Th1 and Th2 immune responses suppress one another and thus hosts have

difficulty mounting simultaneous defenses against macro and microparasites (Abbas et al.

1996, Morel and Oriss 1998, Yazdanbakhsh et al. 2002). Indeed, helminth infections can

suppress the INF- γ expression that promotes the Th1 arm of the immune system, which

targets most intracellular microparasites (Graham 2008). Consistent with the notion that Th1

and Th2 are mutually exclusive and thus tradeoff was the lack of an additive IgY antibody

response in hosts co-infected with Ranavirus and *A. hamatospicula* when compared to single

infections with these parasites (Fig. S4). Therefore, higher Ranavirus infection loads

following *A. hamatospicula* exposure may be driven by the host having previously mounted

an immune response that not only limits available resources to combat a second parasite, but

also suppresses the immune response necessary to combat Ranavirus. Conversely, *A.*

hamatospicula loads were lower with Ranavirus infection. This result could be mediated by

direct competition between parasites. Ranavirus infections commonly induce hemorrhaging

of intestinal epithelial cells, potentially compromising the resources or living space of *A.*

hamatospicula (Miller et al. 2011). If this direct effect was stronger than the indirect effect of

a tradeoff between the Th1 and Th2 immunity, then the effect of Ranavirus on *A.*

hamatospicula would be inhibitory rather than facilitative. These results suggest that

macroparasite co-infections could increase Ranavirus loads, which could significantly impact

amphibian population health (Green et al. 2002).

In co-infections with Bd and *A. hamatospicula*, *A. hamatospicula* loads increased and

Bd loads decreased when they were the second parasite (Fig.2). *A. hamatospicula* enters

through the skin of the frog (Knutie et al. 2017), which is also the site of Bd infection (Berger

et al. 2005). Therefore, higher *A. hamatospicula* loads after Bd infection suggests that Bd

infection facilitates the process of juvenile worms entering the skin. Several mechanisms are

possible for this facilitation. First, Bd disrupts the normal functioning of the skin (Voyles et

al. 2009, Kilpatrick et al. 2010), and can be locally immunosuppressive (Fig. 5; Fites et al.

2013). Therefore, *A. hamatospicula* loads could be higher after Bd exposure due to direct

interactions between these parasites or indirect interactions from Bd immune suppression.

When the order of infections was reversed, Bd loads were lower the longer the host was

previously infected with *A. hamatospicula* (Fig. 2), a priority effect. Given that *A.*

hamatospicula establishes as adults in the gut (Knutie et al. 2017) but Bd infects the skin

(Berger et al. 2005), this priority effect is unlikely to be a product of direct interactions

between these parasites once *A. hamatospicula* establishes. We predicted that Bd loads would

be facilitated by previous *A. hamatospicula* infection because of the Th1/Th2 immune

tradeoff. Although fungal infections cause the host to upregulate the Th1 arm of the immune

system (Blanco and Garcia 2008), they also frequently, but not always (Zelante et al. 2009),

upregulate the Th17 arm of the immune system. Whether amphibians uses the Th1 arm, the

Th17 arm, or a mix of the two to combat Bd infections has been debated and is still unknown

because of a limited understanding of whether the Th17 arm of the immune system is present

in amphibians (Fites 2014). Consequently, amphibian immune responses, especially those

combatting Bd, will need to be better understood to elucidate any immune mechanisms that

might have inhibited Bd growth after *A. hamatospicula* infections. If the pattern seen here

with *A. hamatospicula* and Bd holds up with other macroparasite infections, macroparasite

co-infections (when macroparasites infect first) could have significant positive impacts on

lowering the total Bd load in an environment, positively impacting frog populations in the

field (Kilpatrick et al. 2010).

In co-infections with Ranavirus and Bd, co-infection with either parasite increased the

load of the other. Additionally, there was a positive relationship between Bd loads and the

time lag between Ranavirus and Bd infections; the timing of the prior Ranavirus infection

altered Bd loads (*i.e.,* a priority effect, Fig. 2). In contrast, viral loads increased regardless of

the timing of the Bd co-infection (Fig. 1). These two parasites are unlikely to interact directly,

as Bd infects the skin (Berger et al. 2005) and Ranavirus infects internal organs (Gantress et

al. 2003, Gray et al. 2009). However, they have the potential for indirect interactions.

Infection with Bd significantly suppresses the host immune response (Fig. 5; Fites et al.

2013) and this could facilitate subsequent Ranaviral infection. If Bd activates the immune

system through the Th17 arm of the immune system (Zelante et al. 2009) in addition to the

Th1 arm (Blanco and Garcia 2008), then tradeoffs and complexities of mounting these

immune responses could account for the facilitative pattern observed in this co-infection

pairing. Additionally, these parasites replicate internally and can be costly and deadly to

amphibian hosts (Berger et al. 1998, Gantress et al. 2003, Skerratt et al. 2007, Gray et al.

2009, Kilpatrick et al. 2010). Given that immune responses are costly and can be traded off

with other demanding processes, such as reproduction and growth (Lochmiller and

Deerenberg 2000), mounting an immune response sufficient to combat both of these parasites

could have been beyond the resource capacity of our hosts. Field studies have found that

Ranavirus and Bd are not associated with each other at the site (Olori et al. 2018, Stutz et al.

2018) or individual level, but there are positive associations between these parasites at the

species level for some of the sampled amphibian species (Stutz et al. 2018). Despite this, co-

infections with Bd and Ranavirus, are common in nature (Whitfield et al. 2013, Warne et al.

2016, Watters et al. 2018) and our findings suggest that these co-infections can be facilitative

and thus appear to put host populations at greater risk than single infections.

When we tested how the time lags between infections and the identity of the co-

infecting parasites altered host growth, survival, and tolerance, we found that infection with

Bd or Ranavirus as the first parasite caused higher mortality than *A. hamastopicula* infections

(Fig. 4). We also found that the longer hosts were infected with Bd and Ranavirus, the lower

the growth rate of hosts (Fig. S2). These results support our hypothesis that host growth and

survival should be lower if the infecting parasite is virulent, like Bd and Ranavirus (Gray et

al. 2009, Voyles et al. 2009), and has longer to establish. Additionally, hosts were less

tolerant of Ranavirus infections (lower growth rate [g/week] with higher viral loads) when

the previous infecting parasite was *A. hamatospicula* but were more tolerant when hosts were

exposed to Bd before Ranavirus (Fig.4). Hosts not only tolerated viral loads when co-

exposed to Bd, but also grew more rapidly at higher viral loads than at lower viral loads when

co-infected by Bd. Amphibians display increased growth and development when exposed to

other inhospitable conditions such as rapidly drying ponds (Bekhet et al. 2014), pesticides

(Rohr et al. 2004), or predators (Relyea 2007), which could mean that hosts are using their

resources to grow rapidly when exposed to two deadly amphibian parasites rather than

attempting to resist or clear the infections (*i.e.*, a life-history trade-off).

563   Studies broadly support that the order of co-infections can alter parasite communities

(de Roode et al. 2005, Devevey et al. 2015, Leopold et al. 2017), but our research revealed

that time lags between these co-infections can affect parasite loads, likely as a result of

indirect competition mediated by the host immune system. Additionally, host tolerance,

growth, and survival can be impacted by these co-infections. Our results suggest that co-

infections with Ranavirus and Bd could lead to negative effects on amphibian populations.

Also, if our results with *A. hamatospicula* are generalizable to other macroparasite co-

infections, macroparasite co-infections could increase Ranavirus loads and decrease Bd loads.

This suggests that co-infections with macroparasites and these microparasites could have a

positive or negative overall effect on amphibian population health depending on host

susceptibility to these two deadly microparasites. Given this, further research is needed to

understand the generality of time lag effects and the mechanisms driving time lag effects on

co-infections. A better understanding of when time lags between co-infections are important

will guide researchers and practitioners in understanding when co-infections might be

particularly problematic. This knowledge may, in turn, allow practitioners to implement more

efficient control strategies targeted at co-infections for amphibians and other important hosts

such as pollinators (Bromenshenk et al. 2010), livestock (Ezenwa and Jolles 2015), and

humans (Corbett et al. 2003).

Acknowledgements: We would like to thank J. Hoverman for supplying the Ranavirus, S.

Knutie for guidance with immune assays, J. Cohen for general guidance, and all the

undergraduates who made this experiment possible. Funds were provided by grants to J.R.R.

from the National Science Foundation (EF-1241889), the National Institutes of Health

(R01GM109499, R01TW010286-01).

References:

Abbas, A. K., K. M. Murphy, and A. Sher. 1996. Functional diversity of helper T

    lymphocytes. Nature **383**:787-793.

Baker, M. R. 1987. Synopsis of the Nematoda Parasitic in amphibians and reptiles.

    Occasional papers in biology **11**.

Bekhet, G. A., H. A. Abdou, S. A. Dekinesh, H. A. Hussein, and S. S. Sebiae. 2014.

    Biological factors controlling developmental duration, growth and metamorphosis of

    the larval green toad, Bufo viridis viridis. The Journal of Basic & Applied Zoology

    67:67-82.

Berger, A. 2000. Th1 and Th2 responses: what are they? British Medical Journal **321**:424.

Berger, L., A. D. Hyatt, R. Speare, and J. E. Longcore. 2005. Life cycle stages of the

    amphibian chytrid *Batrachochytrium dendrobatidis*. Dis Aquat Organ **68**:51-63.

Berger, L., R. Speare, P. Daszak, D. E. Green, A. A. Cunningham, C. L. Goggin, R.

    Slocombe, M. A. Ragan, A. D. Hyatt, K. R. McDonald, H. B. Hines, K. R. Lips, G.

    Marantelli, and H. Parkes. 1998. Chytridiomycosis causes amphibian mortality

    associated with population declines in the rain forests of Australia and Central

    America. Proceedings of the National Academy of Sciences **95**.

Blanco, J. L., and M. E. Garcia. 2008. Immune response to fungal infections. Vet Immunol

    Immunopathol **125**:47-70.

Blaustein, L., and J. Margalit. 1996. Priority effects in temporary pools: nature and outcome

    of mosquito larva-toad tadpole interactions depend on order of entrance. British

    Ecological Society **65**:77-84.

Bolker, B. 2017. bbmle: Tools for general maximum likelihood estimation, R package

    version 1.0.20.

630 Boyle, D. G., D. B. Boyle, V. Olsen, A. T. Morgan, and A. D. Hyatt. 2004. Rapid

631  quantitative detection of chytridiomycosis (*Batrachochytrium dendrobatidis*) in

632  amphibian samples using real-time Taqman PCR assay. Dis Aquat Organ **60**:141-148.

633 Breheny, P., and W. Burchett. 2019. visreg: Visualization of regression models, R version

634  2.5-1.

635 Bromenshenk, J. J., C. B. Henderson, C. H. Wick, M. F. Stanford, A. W. Zulich, R. E.

636  Jabbour, S. V. Deshpande, P. E. McCubbin, R. A. Seccomb, P. M. Welch, T.

637  Williams, D. R. Firth, E. Skowronski, M. M. Lehmann, S. L. Bilimoria, J. Gress, K.

638  W. Wanner, and R. A. Cramer, Jr. 2010. Iridovirus and microsporidian linked to

639  honey bee colony decline. PLoS One **5**:e13181.

640 Chesson, P., and J. J. Kuang. 2008. The interaction between predation and competition.

641  Nature **456**:235-238.

642 Chinchar, V. G. 2002. Ranavirus (family *Iridoviridae*): emerging cold-blooded killers.

643  Archives of Virology **147**:447-470.

644 Cohen, J. M., M. D. Venesky, E. L. Sauer, D. J. Civitello, T. A. McMahon, E. A. Roznik, and

645  J. R. Rohr. 2017. The thermal mismatch hypothesis explains host susceptibility to an

646  emerging infectious disease. Ecol Lett **20**:184-193.

647 Corbett, E. L., C. J. Watt, N. Walker, D. Maher, B. G. Williams, M. C. Raviglione, and D. C.

648  2003. The growing burden of Tuberculosis: global trends and interactions with the

649  HIV epidemic. Arch Intern Med **163**:1009-1021.

650 de Roode, J. C., R. Culleton, S. J. Cheesman, R. Carter, and A. F. Read. 2004. Host

651  heterogeneity is a determinant of competitive exclusion or coexistence in genetically

652  diverse malaria infections. Proc Biol Sci **271**:1073-1080.

653   de Roode, J. C., M. E. Helinski, M. A. Anwar, and A. F. Read. 2005. Dynamics of multiple

654   infection and within-host competition in genetically diverse malaria infections. Am

655   Nat **166**:531-542.

656   Devevey, G., T. Dang, C. J. Graves, S. Murray, and D. Brisson. 2015. First arrived takes all:

657   inhibitory priority effects dominate competition between co-infecting Borrelia

658   burgdorferi strains. BMC Microbiol **15**:61.

659   Drake, J. A. 1991. Community-assembly mechanics and the structure of an experimental

660   species ensemble. American Naturalist **137**:1-26.

661   Ezenwa, V. O., and A. E. Jolles. 2015. Epidemiology. Opposite effects of anthelmintic

662   treatment on microbial infection at individual versus population scales. Science

663   **347**:175-177.

664   Facelli, J. M., and E. Facelli. 1993. Interactions after death: plant litter controls priority

665   effects in a successional plant community. Oecologia **95**:277-282.

666   Fites, J. S. 2014. Evasion of adaptive immune defenses by the lethal chytrid fungus

667   *Batrachochytrium dendrobatidis*. Vanderbilt University.

668   Fites, J. S., J. P. Ramsey, W. M. Holden, S. P. Collier, D. M. Sutherland, L. K. Reinert, A. S.

669   Gayek, T. S. Dermody, T. M. Aune, K. Oswald-Richter, and L. A. Rollins-Smith.

670   2013. The invasive chytrid fungus of amphibians paralyzes lymphocyte responses.

671   Science **342**:366-369.

672   Fites, J. S., L. K. Reinert, T. M. Chappell, and L. A. Rollins-Smith. 2014. Inhibition of local

673   immune responses by the frog-killing fungus Batrachochytrium dendrobatidis. Infect

674   Immun **82**:4698-4706.

675   Gantress, J., G. D. Maniero, N. Cohen, and J. Robert. 2003. Development and

676   characterization of a model system to study amphibian immune responses to

677   iridoviruses. Virology **311**:254-262.

Graham, A. L. 2008. Ecological rules governing helminth-microparasite coinfection. Proc

Natl Acad Sci U S A **105**:566-570.

Gray, M. J., D. L. Miller, and J. T. Hoverman. 2009. Ecology and pathology of amphibian

ranaviruses. Dis Aquat Organ **87**:243-266.

Green, D. E., K. A. Converse, and A. K. Schrader. 2002. Epizootiology of sixty-four

amphibian morbidity and mortality events in the USA, 1996-2001. Annals New York

Academy of Sciences **969**.

Greub, G., B. Ledergerber, M. Battegay, P. Grob, L. Perrin, H. Furrer, P. Burgisser, P. Erb,

K. Boggian, J. C. Piffaretti, B. Hirschel, P. Janin, P. Francioli, M. Flepp, and A.

Telenti. 2000. Clinical progression, survival, and immune recovery during

antiretroviral therapy in patients with HIV-1 and hepatitis C virus coinfection: the

Swiss HIV Cohort Study. The Lancet **356**:1800-1805.

Hothorn, W. 2010. multcomp: simultaneous inference in general parametric models, R

package version 1.4-10.

Hoverman, J. T., B. J. Hoye, and P. T. Johnson. 2013. Does timing matter? How priority

effects influence the outcome of parasite interactions within hosts. Oecologia

173:1471-1480.

Hoverman, J. T., J. R. Mihaljevic, K. L. Richgels, J. L. Kerby, and P. T. Johnson. 2012.

Widespread co-occurrence of virulent pathogens within California amphibian

communities. EcoHealth **9**:288-292.

Jamieson, A. M., L. Pasman, S. Yu, P. Gamradt, R. J. Homer, T. Decker, and R. Medzhitov.

2013. Role of tissue protection in lethal respiratory viral-bacterial coinfection. Science

340:1230-1234.

Johnson, P. T., J. C. de Roode, and A. Fenton. 2015. Why infectious disease research needs

community ecology. Science **349**:1259504.

Johnson, P. T. J., and I. D. Buller. 2011. Parasite competition hidden by correlated

coinfection: using surveys and experiments to understand parasite interactions.

Ecology **92**:535-541.

Kilpatrick, A. M., C. J. Briggs, and P. Daszak. 2010. The ecology and impact of

chytridiomycosis: an emerging disease of amphibians. Trends Ecol Evol **25**:109-118.

Kim, J., H. K. Chung, and C. Chae. 2003. Association of porcine circovirus 2 with porcine

respiratory disease complex. The Veterinary Journal **166**:251-256.

Knutie, S. A., C. L. Wilkinson, Q. C. Wu, C. N. Ortega, and J. R. Rohr. 2017. Host resistance

and tolerance of parasitic gut worms depend on resource availability. Oecologia

183:1031-1040.

Kuris, A. M., and K. D. Lafferty. 1994. Community structure: larval trematodes in snail

hosts. Annual Review of Ecology and Systematics **25**:189-217.

Leopold, D. R., J. P. Wilkie, I. A. Dickie, R. B. Allen, P. K. Buchanan, and T. Fukami. 2017.

Priority effects are interactively regulated by top-down and bottom-up forces:

evidence from wood decomposer communities. Ecol Lett **20**:1054-1063.

Lochmiller, R. L., and C. Deerenberg. 2000. Trade-offs in evolutionary immunology: just

what is the cost of immunity? Oikos **88**:87-98.

McMahon, T. A., B. F. Sears, M. D. Venesky, S. M. Bessler, J. M. Brown, K. Deutsch, N. T.

Halstead, G. Lentz, N. Tenouri, S. Young, D. J. Civitello, N. Ortega, J. S. Fites, L. K.

Reinert, L. A. Rollins-Smith, T. R. Raffel, and J. R. Rohr. 2014. Amphibians acquire

resistance to live and dead fungus overcoming fungal immunosuppression. Nature

511:224-227.

Miller, D., M. Gray, and A. Storfer. 2011. Ecopathology of ranaviruses infecting amphibians.

Viruses **3**:2351-2373.

Morel, P. A., and T. B. Oriss. 1998. Crossregulation between Th1 and Th2 cells. Crit Rev

Immunol **18**:275-303.

Olori, J. C., R. Netzband, N. McKean, J. Lowery, K. Parsons, and S. T. Windstam. 2018.

Multi-year dynamics of ranavirus, chytridiomycosis, and co-infections in a temperate

host assemblage of amphibians. Dis Aquat Organ **130**:187-197.

Ortega, N., W. Price, T. Campbell, and J. Rohr. 2015. Acquired and introduced

macroparasites of the invasive Cuban treefrog, Osteopilus septentrionalis. Int J

Parasitol Parasites Wildl **4**:379-384.

Paul, W. E., and J. Zhu. 2010. How are T(H)2-type immune responses initiated and

amplified? Nat Rev Immunol **10**:225-235.

Pedersen, A. B., and A. Fenton. 2007. Emphasizing the ecology in parasite community

ecology. Trends Ecol Evol **22**:133-139.

Picco, A. M., J. L. Brunner, and J. P. Collins. 2007. Susceptibility of the endangered

California tiger salamander, Ambystoma californiense, to ranavirus infection. J Wildl

Dis **43**:286-290.

Piotrowski, J. S., S. L. Annis, and J. E. Longcore. 2004. Physiology of *Batrachochytrium*

dendrobatidis, a chytrid pathogen of amphibians   Mycologia **96**:9-15.

R Core Team. 2017. R: a language and environment for statistical computing. R Foundation

for Statisical Computing, Vienne Austria. R version 3.4.2.

Ramsey, J. P., L. K. Reinert, L. K. Harper, D. C. Woodhams, and L. A. Rollins-Smith. 2010.

Immune defenses against Batrachochytrium dendrobatidis, a fungus linked to global

amphibian declines, in the South African clawed frog, Xenopus laevis. Infect Immun

78:3981-3992.

Relyea, R. A. 2007. Getting out alive: how predators affect the decision to metamorphose.

Oecologia **152**:389-400.

Rohr, J. R., A. A. Elskus, B. S. Shepherd, P. H. Crowley, T. M. McCarthy, J. H.

Niedzwiecki, T. Sager, A. Sih, and B. D. Palmer. 2004. Multiple stressors and

salamanders: effects of an herbicide, food limitation, and hydroperiod. Ecological

Applications **14**:1028-1040.

Rojas, S., K. Richards, J. K. Jancovich, and E. W. Davidson. 2005. Influence of temperature

on Ranavirus infection in larval salamanders Ambystoma tigrinum. Dis Aquat Organ

63:95-100.

Rosenblum, E. B., T. J. Poorten, M. Settles, and G. K. Murdoch. 2012. Only skin deep:

shared genetic response to the deadly chytrid fungus in susceptible frog species. Mol

Ecol **21**:3110-3120.

Scheele, B. C., F. Pasmans, L. F. Skerratt, L. Berger, A. Martel, W. Beukema, A. A.

Acevedo, P. A. Burrowes, T. Carvalho, A. Catenazzi, I. De la Riva, M. C. Fisher, S.

V. Flechas, C. N. Foster, P. Frías-Álvarez, T. W. Garner, B. Gratwicke, J. M.

Guayasamin, M. Hirschfeld, J. E. Kolby, T. A. Kosch, E. La Marca, D. B.

Lindermayer, K. R. Lips, A. V. Longo, R. Maneyro, C. A. McDonald, J. Medelson

III, P. Palacios-Rodriguez, G. Parra-Olea, C. L. Richards-Zawacki, M.-O. Rödel, S.

M. Rovito, C. Soto-Azat, L. F. Toledo, J. Voyles, C. Weldon, S. M. Whitfield, M.

Wilkinson, K. R. Zamudio, and S. Canessa. 2019. Amphibian fungal panzootic causes

catastrophic and ongoing loss of biodiversity. Science **363**:1459-1463.

Skerratt, L. F., L. Berger, R. Speare, S. Cashins, K. R. McDonald, A. D. Phillott, H. B. Hines,

and N. Kenyon. 2007. Spread of Chytridiomycosis Has Caused the Rapid Global

Decline and Extinction of Frogs. EcoHealth **4**:125-134.

Sousa, W. P. 1992. Interspecific interactions among larval trematode parasites of freshwater

and marine snails. American Zoologist **32**:583-592.

Stuart, S. N., J. S. Chanson, N. A. Cox, B. E. Young, A. S. L. Rodrigues, D. L. Fischman,

and R. W. Waller. 2004. Status and trends of amphibian declines and extinctions

worldwide. Science **306**:1783-1786.

Stutz, W. E., A. R. Blaustein, C. J. Briggs, J. T. Hoverman, J. R. Rohr, and P. T. J. Johnson.

2018. Using multi-response models to investigate pathogen coinfections across scales:

insights from emerging diseases of amphibians. Methods Ecol Evol **9**:1109-1120.

Sutherland, J. P. 1974. Multiple stable points in natural communities. Am Nat **108**:859-873.

Swanson, S. J., D. Neitzel, K.D. Reed, and E.A. Belongia. 2006. Coinfections Acquired from

Ixodes Ticks. Clinical Microbiology Reviews **19**:708-727.

Therneau, T. M., and T. Lumley. 2019. survival: Survival analysis, R version 2.44-1.1.

Vhora, M. S., and M. G. Bolek. 2013. New host and distribution records for Aplectana

hamatospicula (Ascaridida: Cosmocercidae) in Gastrophryne olivacea (Anura:

Microhylidae) from the Great Plains U.S.A. J Parasitol **99**:417-420.

Voyles, J., S. Young, L. Berger, C. Campbell, W. F. Voyles, A. Dinudom, D. Cook, R.

Webb, R. A. Alford, L. F. Skerratt, and R. Speare. 2009. Pathogenesis of

Chytridiomycosis, a cause of catastrophic amphibian declines. Science **326**:582-585.

Wake, D. B., and V. T. Vredenburg. 2008. Colloquium paper: are we in the midst of the sixth

mass extinction? A view from the world of amphibians. Proc Natl Acad Sci U S A

105 Suppl 1:11466-11473.

Warne, R. W., B. LaBumbard, S. LaGrange, V. T. Vredenburg, and A. Catenazzi. 2016. Co-

Infection by Chytrid Fungus and Ranaviruses in Wild and Harvested Frogs in the

Tropical Andes. PLoS One **11**:e0145864.

Watters, J. L., D. R. Davis, T. Yuri, and C. D. Siler. 2018. Concurrent Infection of

Batrachochytrium dendrobatidis and Ranavirus among Native Amphibians from

Northeastern Oklahoma, USA. J Aquat Anim Health **30**:291-301.

Whitfield, S. M., E. Geerdes, I. Chacon, E. Ballestero Rodriguez, R. R. Jimenez, M. A.

Donnelly, and J. L. Kerby. 2013. Infection and co-infection by the amphibian chytrid

fungus and ranavirus in wild Costa Rican frogs. Dis Aquat Organ **104**:173-178.

Wilbur, H. M., and R. A. Alford. 1985. Priority effects in experimental pond communities:

responses of Hyla to Bufo and Rana. Ecology **66**:1106-1114.

Wuerthner, V. P., J. Hua, and J. T. Hoverman. 2017. The benefits of coinfection: trematodes

alter disease outcomes associated with virus infection. J Anim Ecol **86**:921-931.

Yazdanbakhsh, M., P. G. Kremsner, and R. van Ree. 2002. Allergy, Parasites, and the

Hygiene Hypothesis. Science **296**:490-494.

Zelante, T., A. De Luca, C. D'Angelo, S. Moretti, and L. Romani. 2009. IL-17/Th17 in anti-

fungal immunity: what's new? Eur J Immunol **39**:645-648.

829   **Table 1.** An example of the general design of the timing of the initial and secondary infection are

830   shown below for one pairwise combination of parasites, single infections, and controls. The same

831   general design was applied to all other pairwise combinations of parasites. There were 5 individuals

832   for each treatment group. Bd = *Bactrachochytrium dendrobatidis*, Rv = Ranavirus.

	Proportion of time lags (time lag : the maximum time lag)				
Exposures	-1 to -0.76	-0.75 to -.51	-0.5 to -0.26	-0.25 to -0.01	0
Bd					Bd
Rv then Bd	Rv				Bd
		Rv			Bd
			Rv		Bd
				Rv	Bd
					Rv & Bd
Rv					Rv
Bd then Rv	Bd				Rv
		Bd			Rv
			Bd		Rv
				Bd	Rv
					Bd & Rv
None					Control

Table 2. A summary of the effects of parasite identity and time lag between co-infecting parasites on the abundance of the second parasite and host tolerance of infection, survival, and growth. Ah = *Aplectana hamatospicula*, Bd = *Batrachochytrium dendrobatidis*, Rv = Ranavirus, + = significant positive effect, - = significant negative effect, 0 = non-significant effect, pe = significant priority effect (significant association between the time lag between co-infecting parasites and the load of the second parasite)

Response variable	Second parasite	First parasite		
		Ah	Bd	Rv
Parasite load of second parasite	Ah	NA	+*	-*
	Bd	-, pe	NA	+, pe
	Rv	-	-	NA
Host tolerance	Ah	NA	0	0
	Bd	0	NA	0
	Rv	-	+	NA
Host survival	Ah	NA	-	0
	Bd	0	NA	0
	Rv	0	-	NA
Host growth (mm/week)	Ah	NA	+	+
	Bd	0	NA	+
	Rv	0	+	NA

Aplectana hamatospicula loads were significantly higher when Bd was the first parasite than when Ranavirus was the first parasite, but *A. hamatospicula* loads in co-infections were not significantly different from single infections.

Fig. 1. Log-transformed viral load at 2, 4, and 8 days after Ranavirus exposure for *O. septentrionalis*

singly infected with Ranavirus (Rv), or initially exposed to *Aplectana hamatospicula* (AhRv) or

Bactrachochytrium dendrobatidis (BdRv) before Ranavirus exposure. Data are shown as a conditional

plot (*i.e.* controlling for everything else in the model), with the expected value (black hashed line) and

partial residuals (points) displayed. Co-infected hosts had higher viral loads than singly infected hosts

(Treatment: χ^2_4 =17.24, p=0.0002) and these differences increased through time (Treatment x day: χ^2_4

= 12.12, p<0.02).

Fig. 2. (A) *Batrachochytrium dendrobatidis* (Bd) loads on *Osteopilus septentrionalis* hosts (8 days

after exposure to Bd) as a function of the standardized time lag (-time lag/maximum time lag) since

first exposed to either *Aplectana hamatospicula* (Ah) or Ranavirus (Rv). A -1.0 corresponds to the

longest time lag between co-infections and 0 corresponds to simultaneous co-infection. The time lag

between co-infecting parasites was associated positively with Bd loads for Ah (p=4.30e-5), but

negatively for Rv (p=0.058). **(B)** Prior exposure to Bd allowed more Ah to successfully establish in

hosts than prior exposure to Ranavirus (Initial parasite identity*: χ^2_1=9.563, p=0.008), but singly

infected hosts did not have significantly different loads from either co-infected host treatments

(p>0.190). Data in both panels are show as conditional plots (*i.e.* controlling for everything else in the

model), with the expected value (black hashed line), a 95% confidence bands or intervals for the

expected value (gray band), and partial residuals (points) displayed.

Fig. 3. *Osteopilus septentrionalis* tolerance (growth rate as a function of viral load) of loads of

Ranavirus when first exposed to either *Aplectana hamatospicula* (Ah) or *Batrachochytrium*

dendrobatidis (Bd). Data are shown as a conditional plot (*i.e.* controlling for everything else in the

model), with the expected value (the black line and black dotted line represent *A. hamatospicula* and

Bd respectively), a 95% confidence band for the expected value (light and darker grey shading

represent *A. hamatospicula* and Bd respectively), and partial residuals (points) displayed. Initial

infection with Ah or Bd decreased and increased host tolerance, respectively (Initial parasite identity x

viral load: $F_{35,1}$=0.056, p<0.02).

Fig. 4. Survival of *Osteopilus septentrionalis* hosts exposed to no parasites (controls, C) or first

exposed to *Aplectana hamatospicula* (Ah), *Bactrachochytrium dendrobatidis* (Bd), or Ranavirus (Rv)

averaged across both secondary parasites. Data are shown as survival curves, with the median survival

(solid and dashed lines) and the 95% confidence interval for the expected value (grey band) displayed.

Initial exposure to Bd and Rv reduced days alive relative to initial exposure to Ah (Bd vs. Ah*:*

p=0.003; Rv vs. Ah: *p*=0.043).

Fig. 5. IgY antibody levels for hosts exposed to *Bactrachochytrium dendrobatidis* (Bd), *Aplectana*

hamatospicula (Ah), Ranavirus (Rv) averaged across single infections and pairwise co-infections and

control individuals with no infections (C). Data are shown as a conditional plot (*i.e*. controlling for

everything else in the model), with the expected value (black hashed line), a 95% confidence interval

for the expected value (gray band), and partial residuals (points) displayed. Antibody levels were not

significantly different in hosts not exposed to parasites and those exposed to Bd (*p*=0.709), which is

known to be immunosuppressive (Fites et al. 2014). However, antibody levels were significantly

higher in hosts exposed to parasites other than Bd (*i.e*., those not known to be immunosuppressive)

relative to controls (Identity of parasite: $F_{2,39}$=4.15, *p*=0.024; non-Bd vs. control: *p*=0.024).

931 **Fig. 1.**

936
937 **Fig. 2.**

938
939

941   **Fig. 3.**

Fig. 4.

952
953 **Fig. 5.**