Varietas delectat: groups with diverse personalities mitigate physiological stress in a songbird

Csongor I. Vágási,[1,2,*,†] Attila Fülöp,[3,4,†] Gergely Osváth,[1,2,5] Péter L. Pap,[1,2] Janka Pénzes,[1] Zoltán Benkő,[1,6] Ádám Z. Lendvai[2] and Zoltán Barta[3,*]

[1]*Evolutionary Ecology Group, Hungarian Department of Biology and Ecology, Babeş-Bolyai University, Cluj-Napoca, Romania;* [2]*Department of Evolutionary Zoology, University of Debrecen, Debrecen, Hungary;* [3]*MTA-DE Behavioural Ecology Research Group, Department of Evolutionary Zoology, University of Debrecen, Debrecen, Hungary;* [4]*Juhász-Nagy Pál Doctoral School, University of Debrecen, Debrecen, Hungary;* [5]*Museum of Zoology, Babeş-Bolyai University, Cluj-Napoca, Romania;* [6]*Romanian Ornithological Society/BirdLife Romania, Cluj-Napoca, Romania*

Correspondence: E-mail: csvagasi@gmail.com; barta.zoltan@science.unideb.hu

[†]*These authors contributed equally to this work.*

https://orcid.org/0000-0002-8736-2391 (C.I.V.), https://orcid.org/0000-0002-7121-9865 (Z.Ba.)

AUTHORS' CONTRIBUTIONS

C.I.V., A.F., and Z.Ba. conceived and designed the study. C.I.V., A.F., P.L.P., O.G., J.P., and Z.Be. performed research. C.I.V., J.P., O.G., and Á.Z.L. contributed new reagents/analytic tools. C.I.V., A.F., and Z.Ba. analysed data. C.I.V., A.F., and Z.Ba. drafted the manuscript with major input from Á.Z.L. and P.L.P. All authors contributed revisions and approved the final version of the manuscript. The authors declare no conflict of interest.

ABSTRACT

Social groups often consist of diverse phenotypes, including personality types. Social selection

theory hypothesizes that group composition (i.e. social environment) influences the performance

of group members. However, this hypothesis remains experimentally unexplored, and it is still

contentious whether a homogeneous or a diversely composed group (i.e. social heterosis) is more

beneficial to the individual. We experimentally formed groups of house sparrows with high and

low diversity of personality, and found that their physiological state (body condition,

physiological stress, and oxidative stress) improved with increasing group-level diversity of

personality. These findings demonstrate that group composition affects the condition of group

members and individuals benefit from social heterosis (i.e. associating with a diverse set of

behavioural types). This aspect of the social life can play a key role in the evolutionary

coexistence of different personalities in nature and has implications for human teams and animal

welfare.

Keywords

Affiliation, exploratory behaviour, group composition, health, *Passer domesticus*, personality,

social environment, social evolution, social heterosis.

INTRODUCTION

Social groups were once considered aggregates of similar individuals, but more recently are viewed as a mixture of members with diverse phenotypes (Farine *et al.* 2015). Variation within group can occur in morphology (e.g. size), behavioural traits (e.g. reactive and proactive behavioural types (Sih *et al.* 2004)), or social roles (e.g. leaders/followers in human teams (King *et al.* 2009), or producers/scroungers in sparrow flocks (Fülöp *et al.* 2019)). Group composition has implications for emergent group-level processes such as decision-making, which ultimately drive group functioning (reviewed in (Aplin *et al.* 2014; Farine *et al.* 2015)). Ethnic diversity can, for instance, have a positive effect on research teams' scientific performance (AlShebli *et al.* 2018).

Personality, the consistent among-individual differences in behavioural phenotype (Réale *et al.* 2007), has strong relevance for social life (Webster & Ward 2011). Social groups can largely differ in their personality composition (Farine *et al.* 2015). Groups' personality composition can have effects at both the level of group as a whole ('upstream effects') and the level of individual group members ('downstream effects'). The group-level consequences of groups' personality composition have mostly been assessed in human teams (van Knippenberg & Schippers 2007; Dyer *et al.* 2009b), where personality composition may affect team performance, albeit in an inconsistent manner (Barrick *et al.* 1998; Halfhill *et al.* 2005; Stewart 2006). In non-human organisms, upstream effects of personality composition have been shown to influence within-group social network structure, collective behaviour, and group performance (Sih & Watters 2005; Dyer *et al.* 2009a; Aplin *et al.* 2014; Bengston & Jandt 2014; Brown & Irving 2014). Although individual-level aspects of sociality, e.g. rank in the dominance hierarchy, have been shown to influence the state of individual members (reviewed by (Sapolsky 2005; Creel *et*

al. 2013)), the downstream effects of group composition on individual state is surprisingly little

scrutinised (Farine *et al.* 2015). This happens despite that social selection theory postulates that

the fitness of an individual is contingent upon the phenotype of those with whom it affiliates (i.e.

social environment; (Wolf *et al.* 1999; Sih & Watters 2005)).

Health state can have a strong influence on the individual's performance (Kelly &

Vitousek 2017) and, at the same time, is moulded by changes of individual's social environment

(Creel *et al.* 2013). Therefore, it is reasonable to assume that physiological condition (e.g. body

condition, stress physiology, oxidative stress, and immune capacity) of group members is also

influenced downstream by the personality composition of their group. Correlative studies on

human work teams found that age and gender composition can be associated with subjective self-

reported health impairment (Wegge *et al.* 2008), but human studies with experimental

manipulation of group composition and actual health measurements are still lacking. Earlier

animal studies only assessed how individuals' position within the social structure (e.g. rank in

dominance hierarchy, an individual-level social attribute) affected their health or physiology

(reviewed by (Sapolsky 2005; Creel *et al.* 2013)). No experimental study on animals, however,

tested directly whether group phenotypic composition (a group-level social attribute) affects the

stress level and health state of group members.

We conducted an experimental study to explore whether manipulated personality

composition of groups *per se* or in interaction with individual personality type affects the

physiological condition of group members. Given that diverse groups might have advantages for

group functioning, we predicted that individuals in groups with more diverse personality

composition would improve their physiological condition as compared with individuals in groups

with more homogeneous personality composition. In this case, improved physiological condition

of group members is a legacy of being part of a diversely composed and better functioning group

and each group member shares this legacy. We do not know whether this potential benefit is

indeed uniform for each member. Alternatively, some members might harvest more the benefits

of better group-level functioning to the detriment of other group members. We assessed this

prediction by testing whether personality diversity of the groups interact with individual

personality type to influence physiological condition. A significant interaction might suggest that

individuals that either match or mismatch the group's personality composition benefit more than

other group members do.

METHODS

Study protocol

The study is based on a large sample of 240 house sparrows. We captured 40 sparrows (1:1 sex

ratio) per each of the six study replicates. These 40 birds were divided into four treatment groups

(see below) consisting of 10 birds each, which yielded 24 social groups for the entire study (four

treatment groups per study replicate × six study replicates). The study timeline was identical in

each study replicate as follows. Upon capture (day 0), birds were marked with an aluminium ring,

and their sex and body mass was recorded. Then they were housed in indoor aviaries for 18 days

at the Campus of Babeș-Bolyai University, Cluj-Napoca, Romania. On days 5–7, we recorded

exploratory behaviour as a well-established and ecologically relevant axis of personality (Wolf &

Weissing 2012) following the novel environment test of Dingemanse and co-authors

(Dingemanse *et al.* 2002). At day 9, we measured the body mass and tarsus length of the birds,

and took pre-treatment blood sample. Then the birds were allocated according to an *a priori*

defined protocol into one of four social treatment groups of 10 birds each: 'random' (random

subsample of birds of a given replicate), 'low exploratory' (only birds with low exploratory

scores), 'high exploratory' (only birds with high scores), and 'variable' (mixture of birds with

either low or high scores). To further characterise the birds' social environment, we calculated

the Shannon diversity index of exploratory behaviour for each group by dividing exploration

values into 10 ordered categories of roughly equal sizes. The social treatment period lasted nine

days until day 18, when we measured again the body mass and took a second blood sample to

measure the post-treatment physiological condition. Physiological state was characterised by

measuring the within-individual change in body condition (scaled body mass index, SMI),

heterophil-to-lymphocyte (H/L) ratio (an indicator of glucocorticoid-mediated stress response;

(Davis *et al.* 2008)), oxidative stress (damage to lipids – malondialdehyde concentration, MDA),

and innate immune capacity (natural antibodies – agglutination score; complement system – lysis

score). Within-individual change in physiological condition was computed as the difference

between the post-treatment and pre-treatment values. We provide a more detailed description of

study timeline, captivity conditions, measurement of exploratory behaviour, assignment to social

treatment groups, blood sampling and physiological measurement methods in the *SI Appendix*.

None of the birds died during the study and all of them were released at the site of capture

in good health on day 18. The study complies with the ethical guidelines of the Babeș-Bolyai

University (permit no. 30792) and the European laws regarding animal welfare, and adheres to

the ASAB guidelines for the use of animals in behavioural research.

Statistical procedures

Malondialdehyde level was log-transformed to normalize its distribution. Changes in each

physiological trait during the social treatment period were analysed in separate models as

dependent variables. These change variables were calculated by subtracting the pre-treatment

values from the post-treatment values. Exception from this are the scores of haemagglutination

and haemolysis that were highly zero-inflated. Therefore, we converted these variables into

binary variables (0 for absence and 1 for presence of agglutination or lysis), and scored the

change during treatment at a 4-point scale by considering all four possible transitions: 0 for $1 \rightarrow$

0 transition ("immune depression"); 1 for $0 \rightarrow 0$ transition ("stable weak immunity"); 2 for $1 \rightarrow$

1 transition ("stable good immunity"); 3 for $0 \rightarrow 1$ transition ("immunity improvement").

The explanatory variables were the same in all models as follows: sex and social

treatment were set as fixed factors with two and four levels, respectively, and individual

exploratory score as a continuous predictor with log(x+1)-transformation. In addition, all second-

order interactions between the three explanatory variables were also tested. Study replicate (six

levels) and group ID (24 levels) nested within study replicate were entered as random factors.

Within each model, the continuous response variables (i.e. body condition, heterophil-to-

lymphocyte ratio and malondialdehyde concentration) and exploration score were Z-transformed

to zero mean and unit standard deviation (Schielzeth 2010). We used linear mixed-effects models

with normal error distribution (LMMs; 'lmer' function of the R package 'lme4' (Bates *et al.*

2015)) for change in body condition, heterophil-to-lymphocyte ratio and malondialdehyde, while

we used cumulative link models (CLMMs; 'clmm' function of the R package 'ordinal'

(Christensen 2015)) for change in agglutination and lysis scores. The assumption of homogeneity

of variances among treatment groups were met for each response variable of the LMM models

(Levene test, all $P > 0.22$). We assessed the fulfilment of model assumptions by graphical

diagnosis; all assumptions were met for each model. Each model was simplified to obtain

minimal adequate models (MAMs) containing only significant main effects or their interactions

by sequentially dropping predictors with non-significant ($P > 0.05$) effects using the 'drop1' R

function. Table 1 presents the type II Anova results of MAMs, while Table S2 presents the

parameter estimates of both the full models and the MAMs.

To assess the relationships between groups' Shannon diversity of personality and changes

in physiological state, we computed the group-level mean of change in physiological variables

for each of the 24 groups by averaging the 10 group members' physiological change values.

These relationships were tested using LMMs with normal error distribution. In all models, the

group-level mean of change in physiological parameter was entered as dependent variable,

groups' Shannon diversity of exploration as a continuous predictor, and study replicate was

entered as a random factor. Both the dependent variables and the exploration diversity values

were Z-transformed. For all the statistical models the reported significance levels were calculated

using type II Wald Chi-square tests using the 'Anova' and 'Anova.clmm' functions of the R

packages 'car' (Fox & Weisberg 2011) and 'RVAideMemoire' (Hervé 2019), respectively.

RESULTS

The groups in the four experimental treatments differed in the mean, variance and Shannon

diversity index of their personality composition (Fig. 1A–C; statistics in Social treatment section

of Materials and Methods in *SI Appendix*). Treatment groups also clearly differed in within-

individual change in body condition (Fig. 1D), heterophil-to-lymphocyte ratio (Fig. 1E) and

oxidative stress (damage to lipids – malondialdehyde concentration; Fig. 1F), but not in change in

the activity of constitutive innate immunity (natural antibodies – agglutination score, and

complement system – lysis score; Fig. 1G and Fig. 1H) (Table 1). The differences in

physiological condition among treatment groups (Fig. 1D–F) are mostly congruent with the

group differences in personality diversity (cf. Fig. 1C) rather than with the mean or variance of

personality of the groups (cf. Fig. 1A–B). Individuals' personality was not associated with the

change in physiological condition by itself, but it was in interaction with social treatment in case

of agglutination and lysis: changes in agglutination scores increased with exploration in the low

exploratory groups, while changes in lysis scores decreased with exploration in the random

groups (Table 1; see also additional results in *SI Appendix* and Table S2).

Figure 1 Behavioural and physiological differences among the social treatment groups.

Treatment groups differ according to (A) mean, (B) variance and (C) Shannon diversity index of

personality (i.e. exploration score), change in (D) body condition (SMI), (E) heterophil-to-

lymphocyte ratio (H/L ratio, an indicator of glucocorticoid-mediated stress response), (F)

oxidative damage to lipids (i.e. malondialdehyde, MDA), and (G–H) constitutive immune

capacity, as expressed through agglutination score (G) and lysis score (H). Means ± SEM are

shown; raw data on panels A–C and model-predicted values on panels D–H. Different lowercase

letters denote significant differences ($P \leq 0.05$), while similar but italicized letters denote

marginal differences ($P < 0.1$) between social treatment groups based on pairwise comparisons

with Tukey-adjusted *P*-values. Male house sparrow drawing credit: Márton Zsoldos.

Table 1 Minimal adequate models containing predictors of changes in individual physiological state of house sparrows during the social treatment period. Statistically significant effects are marked in bold. SMI – Scaled Mass Index (body condition), H/L – heterophil-to-lymphocyte ratio (indicator of glucocorticoid-mediated stress response), MDA – malondialdehyde (oxidative damage to lipids), S – sex, T – social treatment, EB – exploratory behaviour.

Response	Predictor	χ^2	df	P
SMI	S	6.52	1	**0.011**
	T	17.15	3	**< 0.001**
H/L	T	7.94	3	**0.047**
MDA	S	0.13	1	0.719
	T	13.34	3	**0.004**
	S × T	8.35	3	**0.039**
Agglutination	T	0.32	3	0.955
	EB	0.05	1	0.831
	T × EB	9.53	3	**0.023**
Lysis	T	1.13	3	0.770
	EB	1.78	1	0.182
	T × EB	10.74	3	**0.013**

To assess the role of personality diversity of experimental social groups, we tested

whether the calculated Shannon diversity predicts the physiological responses to social treatment.

Because the Shannon diversity index is a group-level metric (i.e. one value for a 10 birds group,

hence 24 values in total), we computed the group-level mean of within-individual change in each

physiological trait by averaging the change values of the 10 birds per group and assessed their

association with groups' Shannon diversity of personality. We found that in groups with higher

personality diversity body condition increased more ($\beta = 0.515$, SE $= 0.152$, $t = 3.380$, $P < 0.001$;

Fig. 2A), while heterophil-to-lymphocyte ratio ($\beta = -0.279$, SE = 0.128, $t = 2.174$, $P = 0.030$; Fig.

2B) and oxidative damage to lipids (i.e. malondialdehyde; $\beta = -0.464$, SE = 0.189, $t = 2.457$, $P =$

0.014; Fig. 2C) became lower. Mean change in agglutination and lysis scores were not related to

group personality diversity (agglutination: $\beta = 0.058$, SE = 0.188, $t = 0.310$, $P = 0.757$; lysis: $\beta =$

-0.034, SE = 0.131, $t = 0.263$, $P = 0.792$).

Figure 2 Physiological condition improved in social groups with diverse personalities.

Relationships between personality diversity in social groups of house sparrows and mean change

in (A) body condition (SMI), (B) heterophil-to-lymphocyte (H/L) ratio, and (C) oxidative damage

to lipids (MDA) during the social treatment period. Dots are group-level means, regression lines

are model-predicted slopes with 95% confidence intervals (shaded area).

DISCUSSION

In the realm of human psychology, it is a long-standing debate whether uniformly or diversely

composed teams perform better (Barrick *et al.* 1998; Stewart 2006), but nothing has been known

about whether group composition affects the physiological condition of group members either in

humans or other animals. Here we showed that "variety is delighting," as condition improved,

and levels of glucocorticoid-mediated stress response and oxidative stress was lower for house

sparrows living in a social environment with diverse personalities. Therefore, living in social

groups with diverse composition provides benefits in terms of reduced physiological stress and

superior health state. The finding of no significant interactions between social treatment and

individual's exploration, at least in cases of improved physiological variables, suggests that all

individuals in diverse groups enjoy the benefits.

Diverse groups provide more opportunities for specialization (Bergmüller & Taborsky

2010; Farine *et al.* 2015) and are more likely to host keystone individuals, which are influential

individuals with disproportionately large effect on other group members and/or overall group

functioning (Modlmeier *et al.* 2014). Both role specialization and keystone individuals can lead

to superior group-level performance (upstream effect). Indeed, great tit affiliations consisting of

diverse personalities show the most effective coordinated action when exploring a habitat patch

(Aplin *et al.* 2014). Note that the group compositions simulated in this modelling study (Aplin *et*

al. 2014) are highly similar to our experimental setup. Similarly, a mixture of shy and bold fish

can be advantageous in reducing the trade-off between exploring novel foraging tasks and

antipredator vigilance (Dyer *et al.* 2009a). The minority of keystone individuals can also

substantially affect group-level behaviour and performance (Brown & Irving 2014). These group-

level advantages of diversely composed groups can bring about higher individual performance

(downstream effect) in terms of either fitness or condition (Bengston & Jandt 2014) by reducing

stress exposure and ultimately leaving group members in better physiological condition. Note

that we found a positive effect of social diversity on health in a set-up where food was unlimited.

This suggests that this benefit of diversity might arise because of the innate working of the group

(e.g. the type and intensity of interactions between members) rather than being the consequence

of more obvious benefits like improved foraging success, habitat exploration, defence against

predators, and decision-making. Therefore, preference for bonding with dissimilar individuals

(i.e. heterophily), and in turn, the better health state of individuals in diverse groups might

reinforce the improved group-level outcomes creating a positive feedback loop (Seebacher &

Krause 2017) between group-level and individual-level performances. Such a feedback loop can

then drive the evolutionary maintenance of heterophily (Pruitt & Riechert 2011).

Consistency of personality traits places a constraint on individuals because one is either

more reactive (shy, neophobic, less exploratory, less aggressive) or more proactive (bold, novelty

seeking, more exploratory, more aggressive) (Sih *et al.* 2004). However, if different personalities

affiliate, they can share mutual benefits; a concept termed social heterosis (Nonacs & Kapheim

2007). Social heterosis in associations of dissimilar personalities thus can explain why

behavioural (and genetic) diversity can evolutionarily persist (Nonacs & Kapheim 2007).

Negative frequency-dependent selection is another evolutionary explanation for the existence of

behavioural polymorphisms (producers–scroungers, hawks–doves, and leaders–followers), and

has been shown to maintain diversity in group personality composition (Pruitt & Riechert 2009).

Our results provide a physiological mechanism that could also be responsible for the evolutionary

maintenance of behavioural diversity in social groups. Behavioural diversity is a requisite for the

evolution of leadership (Johnstone & Manica 2011), cooperation (McNamara *et al.* 2004), and

social responsiveness, which in turn is necessary for the evolution of personality (Wolf *et al.*

2011). Our findings, thus, bring helpful insight into the study of social evolution, which is a

fundamental question in biology and has implications for human work teams in particular and for

human society in general. Although the role of group composition in human team performance is

still contentious (van Knippenberg & Schippers 2007), there is some evidence showing that

heterophily is advantageous in project groups that are less stable in time and are engaged in

creative tasks, but disadvantageous in production groups that are stable in time and are engaged

in routine tasks (Stewart 2006). If humans also may gain health benefits from belonging to a

diversely composed team as sparrows do, the fact that homophily is still the rule of thumb in

humans (McPherson *et al.* 2001) poses a challenge on the health benefits of heterophily, and thus

deserves more attention.

ACKNOWLEDGEMENTS

We thank A. Jakab for providing accommodation for A.F. during the experiment, M. Bán for

technical assistance, and V. Bókony and F. Bonier for highly useful comments. The study was

financed by the Hungarian National Research, Development and Innovation Office (#K112527 to

Z.Ba.) and a Postdoctoral Fellowship of the Hungarian Academy of Sciences (to C.I.V.). C.I.V.

was supported by the János Bolyai Fellowship of the Hungarian Academy of Sciences and by the

Hungarian National Research, Development and Innovation Office (#PD121166), A.F. by a PhD

scholarship and through the New National Excellence Program of the Ministry of Human

Capacities of Hungary (#ÚNKP-16-3-IV), Á.Z.L. by the Hungarian National Research,

Development and Innovation Office (#K113108), P.L.P. by a János Bolyai Fellowship of the

Hungarian Academy of Sciences, and Z.Ba. by the Higher Education Institutional Excellence

Program (NKFIH-1150-6/2019) of the Ministry of Innovation and Technology in Hungary,

within the framework of the FIK-DE Behavioural Ecology Research Group thematic program of

the University of Debrecen.

REFERENCES

AlShebli, B.K., Rahwan, T. & Woon, W.L. (2018). The preeminence of ethnic diversity in

scientific collaboration. *Nat. Commun.*, 9, 5163.

Aplin, L.M., Farine, D.R., Mann, R.P. & Sheldon, B.C. (2014). Individual-level personality

influences social foraging and collective behaviour in wild birds. *Proc. R. Soc. London B*,

281, 20141016.

Barrick, M.R., Stewart, G.L., Neubert, M.J. & Mount, M.K. (1998). Relating member ability and

personality to work-team processes and team effectiveness. *J. Appl. Psychol.*, 83, 377–391.

Bates, D., Mächler, M., Bolker, B. & Walker, S. (2015). Fitting linear mixed-effects models

using lme4. *J. Stat. Softw.*, 67, 1–48.

Bengston, S.E. & Jandt, J.M. (2014). The development of collective personality: the ontogenetic

drivers of behavioral variation across groups. *Front. Ecol. Evol.*, 2, 81.

Bergmüller, R. & Taborsky, M. (2010). Animal personality due to social niche specialisation.

Trends Ecol. Evol., 25, 504–511.

Brown, C. & Irving, E. (2014). Individual personality traits influence group exploration in a feral

guppy population. *Behav. Ecol.*, 25, 95–101.

Christensen, R.H.B. (2015). *ordinal - Regression models for ordinal data. R package version*

2015.6-28. http://www.cran.r-project.org/package=ordinal/.

Creel, S., Dantzer, B., Goymann, W. & Rubenstein, D.R. (2013). The ecology of stress: effects of

the social environment. *Funct. Ecol.*, 27, 66–80.

Davis, A.K., Maney, D.L. & Maerz, J.C. (2008). The use of leukocyte profiles to measure stress

in vertebrates: a review for ecologists. *Funct. Ecol.*, 22, 760–772.

Dingemanse, N.J., Both, C., Drent, P.J., van Oers, K. & van Noordwijk, A.J. (2002).

Repeatability and heritability of exploratory behaviour in great tits from the wild. *Anim.*

Behav., 64, 929–938.

Dyer, J.R.G., Croft, D.P., Morrell, L.J. & Krause, J. (2009a). Shoal composition determines

foraging success in the guppy. *Behav. Ecol.*, 20, 165–171.

Dyer, J.R.G., Johansson, A., Helbing, D., Couzin, I.D. & Krause, J. (2009b). Leadership,

consensus decision making and collective behaviour in humans. *Philos. Trans. R. Soc.*

London B, 364, 781–789.

Farine, D.R., Montiglio, P.-O. & Spiegel, O. (2015). From individuals to groups and back: the

evolutionary implications of group phenotypic composition. *Trends Ecol. Evol.*, 30, 609–

621.

Fox, J. & Weisberg, S. (2011). *An {R} Companion to Applied Regression*. 2nd edn. Sage,

Thousand Oaks, CA.

Fülöp, A., Németh, Z., Kocsis, B., Deák-Molnár, B., Bozsoky, T. & Barta, Z. (2019). Personality

and social foraging tactic use in free-living Eurasian tree sparrows (*Passer montanus*).

Behav. Ecol., 30, 894–903.

Halfhill, T., Sundstrom, E., Lahner, J., Calderone, W. & Nielsen, T.M. (2005). Group personality

composition and group effectiveness. *Small Gr. Res.*, 36, 83–105.

Hervé, M. (2019). *RVAideMemoire: testing and plotting procedures for biostatistics. R package*

version 0.9-73.

Johnstone, R.A. & Manica, A. (2011). Evolution of personality differences in leadership. *Proc.*

Natl. Acad. Sci. USA, 108, 8373–8378.

Kelly, A.M. & Vitousek, M.N. (2017). Dynamic modulation of sociality and aggression: an

examination of plasticity within endocrine and neuroendocrine systems. *Philos. Trans. R.*

Soc. London B, 372, 20160243.

King, A.J., Johnson, D.D.P. & Van Vugt, M. (2009). The origins and evolution of leadership.

Curr. Biol., 19, R911–R916.

van Knippenberg, D. & Schippers, M.C. (2007). Work group diversity. *Annu. Rev. Psychol.*, 58,

515–541.

McNamara, J.M., Barta, Z. & Houston, A.I. (2004). Variation in behaviour promotes cooperation

in the Prisoner's Dilemma game. *Nature*, 428, 745–748.

McPherson, M., Smith-Lovin, L. & Cook, J.M. (2001). Birds of a feather: homophily in social

networks. *Annu. Rev. Sociol.*, 27, 415–444.

Modlmeier, A.P., Keiser, C.N., Watters, J. V., Sih, A. & Pruitt, J.N. (2014). The keystone

individual concept: an ecological and evolutionary overview. *Anim. Behav.*, 89, 53–62.

Nonacs, P. & Kapheim, K.M. (2007). Social heterosis and the maintenance of genetic diversity.

J. Evol. Biol., 20, 2253–2265.

Pruitt, J.N. & Riechert, S.E. (2009). Frequency-dependent success of cheaters during foraging

bouts might limit their spread within colonies of a socially polymorphic spider. *Evolution*,

63, 2966–2973.

Pruitt, J.N. & Riechert, S.E. (2011). Within-group behavioral variation promotes biased task

performance and the emergence of a defensive caste in a social spider. *Behav. Ecol.*

Sociobiol., 65, 1055–1060.

Réale, D., Reader, S.M., Sol, D., McDougall, P.T. & Dingemanse, N.J. (2007). Integrating

animal temperament within ecology and evolution. *Biol. Rev.*, 82, 291–318.

Sapolsky, R.M. (2005). The influence of social hierarchy on primate health. *Science*, 308, 648–

652.

Schielzeth, H. (2010). Simple means to improve the interpretability of regression coefficients.

Methods Ecol. Evol., 1, 103–113.

Seebacher, F. & Krause, J. (2017). Physiological mechanisms underlying animal social

behaviour. *Philos. Trans. R. Soc. London B*, 372, 20160231.

Sih, A., Bell, A. & Johnson, J.C. (2004). Behavioral syndromes: an ecological and evolutionary

overview. *Trends Ecol. Evol.*, 19, 372–378.

Sih, A. & Watters, J. V. (2005). The mix matters: behavioural types and group dynamics in water

striders. *Behaviour*, 142, 1417–1431.

Stewart, G.L. (2006). A meta-analytic review of relationships between team design features and

team performance. *J. Manage.*, 32, 29–55.

Webster, M.M. & Ward, A.J.W. (2011). Personality and social context. *Biol. Rev.*, 86, 759–773.

Wegge, J., Roth, C., Neubach, B., Schmidt, K.-H. & Kanfer, R. (2008). Age and gender diversity

as determinants of performance and health in a public organization: the role of task

complexity and group size. *J. Appl. Psychol.*, 93, 1301–1313.

Wolf, J.B., Brodie III, E.D. & Moore, A.J. (1999). Interacting phenotypes and the evolutionary

process. II. Selection resulting from social interactions. *Am. Nat.*, 153, 254–266.

Wolf, M., Van Doorn, G.S. & Weissing, F.J. (2011). On the coevolution of social responsiveness

and behavioural consistency. *Proc. R. Soc. London B*, 278, 440–448.

Wolf, M. & Weissing, F.J. (2012). Animal personalities: consequences for ecology and

evolution. *Trends Ecol. Evol.*, 27, 452–461.

SI Appendix

METHODS

Study protocol

The study is based on a large sample of 240 house sparrows. We caught 40 sparrows during each

study replicate (1:1 sex ratio). Sparrows were caught with mist nets at a cattle farm near Bălcaciu

village, central Transylvania, Romania (46°11'N, 24°3'E) during six capture sessions (9

November 2014, 5 December 2014, 5 January 2015, 23 January 2015, 10 February 2015, and 28

February 2015). Upon capture (day 0), birds were marked with an aluminium ring, and their sex

and body mass (± 0.1 g) was recorded. The birds were transported to the Campus of the Babeș-

Bolyai University, Cluj-Napoca (46°46'N, 23°33'E) and housed in indoor aviaries for 18 days.

Study timeline

The study timeline was the same for all six replicates. We let the birds to habituate to captivity on

the day of capture (day 0) and the next two days (days 1–2). They were housed in four indoor

aviary rooms (3 m length × 2 m width × 2.5 m height) in which the birds were distributed

randomly in groups of equal sizes (10 birds in each room). The aviary rooms were visually

separated from each other. To assess the exploratory behaviour of birds in a novel environment,

we first transferred them into individual cages in the morning of day 3 and let them to habituate

for two days (days 3–4), then tested for exploration on days 5–7 (see below the details). Day 8

was a resting day. At day 9, we measured the body mass and tarsus length (± 0.01 mm) of the

birds, and took the pre-treatment blood sample (150–200 μL; see below the methods). Then the

birds were allocated according to an *a priori* defined protocol into one of four social treatment

groups of 10 birds each (see below). These four social groups of 10 birds in each of the six study

replicates were housed in the same four adjacent aviary rooms as mentioned above. Each social

group had an even or quasi-even sex ratio (Table S1). There was no significant difference in sex

ratio between the treatment groups in any of the study replicates (χ^2 test, all $P > 0.362$). The

social treatment period lasted nine days until day 18, when we measured again the body mass and

took a second blood sample to measure the post-treatment physiological condition. On the same

day, we released the birds at the site of capture.

Table S1. Sex ratio as shown by the sample sizes per each sex (F – female, M – male) per each

experimental group per each study replicate. Each group was formed by 10 birds during each

study replicate totalling 40 birds per study replicate and 240 birds for the entire study.

	Social treatment group							
	random		variable		low		high	
	F	M	F	M	F	M	F	M
replicate #1	4	6	5	5	6	4	5	5
replicate #2	6	4	6	4	4	6	4	6
replicate #3	5	5	3	7	7	3	5	5
replicate #4	5	5	3	7	7	3	5	5
replicate #5	7	3	4	6	5	5	4	6

replicate #6	7	3	4	6	5	5	4	6

Housing and ethical note

Birds were transported within max. 4 h from capture into aviaries. To increase the sparrow's comfort, aviaries were enriched with several perches and one nest box per bird for resting, hiding and roosting, and a water tank was full-time available for bathing. The artificial photoperiod was identical to the natural day–night cycle throughout. Birds were fed *ad libitum* with a seed mixture consisting of ground corn, barley, millet and sunflower, and this diet was supplemented with one grated boiled egg per aviary room every other day (Pap *et al.* 2008; Vágási *et al.* 2010). Fresh drinking water was provided on a daily basis. None of the birds died during the study and all of them were released at the site of capture in good health.

Exploratory behaviour

When transferred into individual cages (day 3), birds were randomly ordered from 1 to 40 and split into three clusters (first 13, next 14, and last 13 birds). Their exploration test was performed according to this 1–40 order on days 5–7 (one cluster was tested each day). We recorded exploratory behaviour as a well-established axis of personality following the novel environment test of Dingemanse and co-authors (Dingemanse *et al.* 2002). Sparrows were deprived of food and water for 1 h before the novel environment test started. Their cage was moved to the test room 10 min before the test run and was covered with a dark curtain, so birds were left to calm down in complete darkness and quietness before the test run. The birds entered the test room from their cage through a sliding door after being startled by knocking the wall of the cage, but

without being handled or seeing any person. They were tested alone by spending 10 min in the

test room (3 m length × 2 m width × 2 m height) that contained four artificial wooden trees with

four branches each and arranged symmetrically within the test room. Exploratory behaviour was

video recorded through a one-way window with a hand-held video camera (Panasonic HC-V510)

between 09:00 and 16:00 (schedule of the test runs: 09:00, 09:30, 10:00, 10:30, 11:00, 11:30,

12:00, 12:30, 13:00, 13:30, 14:00, 14:30, 15:00, and 15:30) by the same person (A.F.).

Exploratory score is the total number of hops (performed either on the trees or on the ground) and

flights during the 10-min test. The exploratory behaviour was scored by the same person (Z.Be.).

An additional set of 40 birds that were not involved in the social experiment were

assessed thrice for their exploratory behaviour in the same novel environment as the 240

experimental birds in order to verify whether this behavioural trait is consistent in time, a

prerequisite of personality traits. The timeline and housing condition for these 40 birds were

identical with those 240 birds that were involved in the six study replicates (i.e. they were housed

under the same conditions and spent the same number of days before the first test and between

the consecutive tests). Consistency of the exploratory behaviour was measured by calculating

individual repeatability (i.e. separating variation in exploratory score into a within-individual and

an among-individuals component) using a linear mixed-effects model (R package 'rptR' (Stoffel

et al. 2017)) as per Nakagawa and Schielzeth (Nakagawa & Schielzeth 2010). Exploration score

was first log-transformed and then Z-transformed (i.e. scaled to mean = 0 and standard deviation

= 1; (Schielzeth 2010)). We first built a full model in which exploration score was the dependent

variable with sex (male/female), exploration test repeat (first/second/third), aviary room (from

one to four) where the birds were kept between test repeats, and their second-order interaction

were entered as potential confounding fixed effects, and individual's ID, test day (three test days;

see above), and the novel environment test order (1–40) nested within test day were entered as

random factors. The minimal model was obtained by sequentially dropping all the non-significant

fixed predictors from the full model until only significant effects remained. Individuals were

significantly consistent in their exploratory behaviour across the three exploration test repeats in

both the full model and minimal model (full model: $R = 0.472$, SE $= 0.097$, 95% confidence

interval $= 0.316$–0.691, $P < 0.001$; minimal model: $R = 0.416$, SE $= 0.100$, 95% confidence

interval $= 0.199$–0.589, $P < 0.001$).

Social treatment

The social treatment consisted of creating four groups that differed in personality composition:

'random' (random subsample of birds of a given replicate), 'high variance' (mixture of birds with

either low or high exploration scores), 'low exploratory' (only birds with low scores), and 'high

exploratory' (only birds with high scores). For this, we first ranked the 40 birds of each study

replicate according to their exploration score in an increasing order (i.e. rank #1 is the least

exploratory bird). The 'random' group was set up by forming 10 quartets along this rank order

(i.e. first quartet consisting of birds ranking #1–4 and the last quartet of birds ranking #37–40),

randomizing the order within each quartet, and then choosing the first bird from each quartet. The

'high variance' group was set up by reordering the remaining 30 birds, forming 15 duos along

this rank order, randomizing the order of birds within each duo, and choosing the first bird from

the first five and the last five duos. The remaining 20 birds were reordered once again and the

first 10 birds along this exploration rank order formed the 'low exploratory' group, while the last

10 birds formed the 'high exploratory' group. The goodness of this protocol was *a priori* assessed

by generating 40 random exploration scores with uniform, normal or exponential distribution.

The group formation protocol worked for each of the three distribution types as the four groups

the protocol created differed both according to mean and to variance of exploration scores.

Social treatment protocol successfully created differences in the mean, variance and

Shannon diversity index of exploration score (personality) among groups (Fig. 1). The mean

exploratory score was smallest in the low exploratory group, intermediate in the random and

variable groups, and largest in the high exploratory group (ANOVA, $F = 18.87$, df $= 3,236$, $P <$

0.001) (Fig. 1A). Variance was lower in the low and high exploratory groups than in the random

and variable groups ($F = 16.21$, df $= 1,238$, $P < 0.001$) (Fig. 1B). Comparing the latter two

groups, variance was lower in the random group than the variable group ($F = 10.04$, df $= 1,118$, P

$= 0.002$) (Fig. 1B), which shows that the random group is an even sample of the entire range of

exploratory behaviour, while the variable group is in fact a mixture of low exploratory and high

exploratory groups in equal proportion. Indeed, opposite to the exploration score variance, the

Shannon diversity index computed for each social group of 10 sparrows (24 groups in total; four

treatment groups per study replicate × six study replicates) was highest in the random group, was

intermediate in the variable group, and lowest in the low and high exploratory groups (treatment

group effect: $\chi^2 = 68.118$, df $= 3$, $P < 0.001$; random vs. variable: $P = 0.008$) (Fig. 1C).

Blood sampling

Blood samples were collected on day 9 and day 18 to assess the physiological state of sparrows

before and after the social treatment period, respectively. Blood samples were collected into

heparinized capillaries by puncturing the brachial vein with insulin syringe. A drop of blood was

smeared onto a microscope slide for counting leucocytes. The capillaries with blood samples

were stored in dark cooling boxes at 4°C for max. 4 h until centrifuged (5 min at 6200 g) to

separate the plasma and erythrocyte fractions. Plasma was partitioned into aliquots for each

physiological parameter and all aliquots were stored at –50°C until the laboratory assay took

place.

Physiological parameters

We measured the following five parameters to describe the physiological state of the birds. First,

we computed a size-corrected body mass index to characterize the individuals' body condition

(i.e. the relative amount of energy stores in the form of muscle and fat). For this, we used the

Scaled Mass Index (Peig & Green 2009) (for details, see (Vágási *et al.* 2012)). Second, 50

leukocytes were counted from blood smears by G.O. (for details, see (Pap *et al.* 2011, 2015)).

Heterophil-to-lymphocyte ratio was used as an indicator of glucocorticoid-mediated stress

response (Davis *et al.* 2008). Because all the leukocytes were heterophils on some smears,

heterophil-to-lymphocyte ratio was calculated as heterophils / (heterophils + lymphocytes); thus,

a value close to 1 indicates higher physiological stress. Third, oxidative stress was assessed by

J.P. and C.I.V. by measuring the amount of oxidative damage to cell membrane phospholipids

via the plasma concentration of malondialdehyde, a toxic intermediate of oxidative lipid

decomposition (for details, see (Bókony *et al.* 2014)). Fourth, the level of natural antibodies

(agglutination score) and the activity of the complement system (lysis score) as two associated

measures of the constitutive innate immune system was assessed by J.P. and C.I.V. via a

haemagglutination–haemolysis assay (Matson *et al.* 2005) (for details, see (Pap *et al.* 2010)).

Higher scores mean that the immune system constituents of the plasma can agglutinate or lyse

foreign red blood cells at lower concentration (i.e. indicate better immune capacity).

ADDITIONAL RESULTS

Individual's exploration score had no clear association with the change in physiological condition either alone or in interaction with social treatment (Table 1; see Table S2 for full models). Exceptions are the cases of change in natural antibody level (i.e. agglutination score) that was positively associated with personality in the low exploratory group ($\beta = 0.893$, SE = 0.351, $z = 2.547$, $P = 0.011$), but not in other groups (all $P > 0.05$), and the change in complement activity (i.e. lysis score) that was negatively associated with personality in the random group ($\beta = -0.693$, SE = 0.217, $z = 3.188$, $P = 0.001$), but not in other groups (all $P > 0.05$).

Table S2. Parameter estimates of full models containing all the predictors of changes in individual physiological state of house sparrows during the social treatment period. Parameter estimates of minimal adequate models are also shown besides the corresponding full models. Statistically significant effects are marked in bold. (a) SMI – Scaled Mass Index (body condition), (b) H/L – heterophil-to-lymphocyte ratio (indicator of physiological stress), (c) MDA – malondialdehyde (oxidative damage to lipids), (e) Agglutination – level of natural antibodies, (f) Lysis – activity of the complement system. Predictors: HVG – high variance group (experimental group with high exploratory behaviour variance), HEG – high exploratory group (experimental group of birds with high exploratory behaviour), LEG – low exploratory group (experimental group of birds with low exploratory behaviour), S – sex (F – female is the reference level), EB – exploratory behaviour, REP – study replication, T – social treatment. For random effects, σ^2 is the residual variance, while τ_{00} is the variance explained by random factors.

549   (a) SMI

Fixed effects	Full model			Min. adequate model		
	β	SE	t	β	SE	t
Intercept	0.336	0.23	1.458	0.595	0.19	3.134
HVG	−0.257	0.258	0.996	−0.508	0.183	2.768
HEG	−0.524	0.293	1.792	−0.721	0.183	3.94
LEG	0.113	0.306	0.37	−0.54	0.182	2.958
S (F)	0.165	0.245	0.675	−0.305	0.12	2.553
EB	0.137	0.143	0.957			
HVG × S	−0.418	0.354	1.179			
HEG × S	−0.509	0.351	1.451			
LEG × S	−0.758	0.355	2.134			
HVG × EB	−0.142	0.159	0.894			
HEG × EB	−0.03	0.232	0.129			
LEG × EB	0.184	0.251	0.735			
S × EB	0.069	0.14	0.488			
Random effects						
σ^2	0.82			0.84		

	τ₀₀	0.02 REP:T; 0.09 REP	0.02 REP:T; 0.09 REP

τ_{00}	0.02 $_{REP:T}$; 0.09 $_{REP}$	0.02 $_{REP:T}$; 0.09 $_{REP}$
N	6 REP; 4 T	6 REP; 4 T
Observations	240	240
Marg. R^2; Cond. R^2	0.118; 0.231	0.086; 0.188

551   (b) H/L

	Full model			Min. adequate model		
Fixed effects	β	SE	t	β	SE	t
Intercept	–0.43	0.244	1.765	–0.271	0.199	1.362
HVG	0.299	0.248	1.208	0.333	0.169	1.972
HEG	0.454	0.285	1.59	0.459	0.169	2.72
LEG	0.362	0.3	1.209	0.293	0.169	1.736
S (F)	0.259	0.251	1.034			
EB	0.157	0.147	1.066			
HVG × S	0.145	0.363	0.4			
HEG × S	–0.033	0.36	0.092			
LEG × S	–0.177	0.364	0.486			
HVG × EB	–0.08	0.163	0.493			

	Full model			Min. adequate model		
HEG × EB	−0.016	0.237	0.069			
LEG × EB	−0.16	0.256	0.625			
S × EB	−0.104	0.144	0.722			
Random effects						
σ^2	0.87			0.85		
τ_{00}	0.00 REP:T; 0.15 REP			0.00 REP:T; 0.15 REP		
N	6 REP; 4 T			6 REP; 4 T		
Observations	240			240		
Marg. R^2; Cond. R^2	0.054; NA			0.032; NA		

553 (c) MDA

	Full model			Min. adequate model		
Fixed effects	β	SE	t	β	SE	t
Intercept	0.015	0.196	0.074	0.008	0.191	0.042
HVG	−0.403	0.262	1.538	−0.409	0.253	1.615
HEG	−0.028	0.301	0.092	−0.07	0.257	0.272
LEG	0.519	0.315	1.648	0.441	0.27	1.636
S (F)	−0.302	0.262	1.152	−0.278	0.254	1.094

	Full model			Min. adequate model		
	β	SE	t	β	SE	t
EB	−0.025	0.156	0.164			
HVG × S	0.478	0.382	1.252	0.517	0.362	1.428
HEG × S	0.904	0.385	2.349	0.864	0.367	2.355
LEG × S	−0.054	0.382	0.142	−0.035	0.359	0.098
HVG × EB	−0.002	0.174	0.013			
HEG × EB	−0.045	0.256	0.175			
LEG × EB	0.174	0.269	0.645			
S × EB	−0.089	0.154	0.578			
Random effects						
σ^2	0.95			0.94		
τ_{00}	0.00 REP:T; 0.00 REP			0.00 REP:T; 0.00 REP		
N	6 REP; 4 T			6 REP; 4 T		
Observations	231			231		
Marg. R^2; Cond. R^2	0.091; NA			0.088; NA		

555 (d) Agglutination

	Full model			Min. adequate model		
Fixed effects	β	SE	t	β	SE	t

HVG	0.884	0.56	0.22	0.844	0.426	0.399
HEG	0.433	0.644	1.299	0.565	0.504	1.133
LEG	1.859	0.656	0.945	1.758	0.523	1.079
S (F)	1.021	0.503	0.041			
EB	0.757	0.302	0.923	0.747	0.255	1.147
HVG × S	0.887	0.766	0.157			
HEG × S	1.641	0.748	0.662			
LEG × S	0.912	0.742	0.124			
HVG × EB	1.08	0.343	0.223	1.099	0.327	0.289
HEG × EB	2.443	0.519	1.722	2.241	0.505	1.597
LEG × EB	4.359	0.557	2.644	4.332	0.549	2.672
S × EB	0.975	0.311	0.081			

Random effects		
σ^2	3.29	3.29
τ_{00}	0.17 REP:T; 0.30 REP	0.17 REP:T; 0.29 REP
N	6 REP; 4 T	6 REP; 4 T
Observations	237	237
Marg. R^2; Cond. R^2	0.049; 0.167	0.045; 0.162

557   (e) Lysis

Fixed effects	Full model			Min. adequate model		
	β	SE	t	β	SE	t
HVG	1.416	0.575	0.605	1.217	0.402	0.488
HEG	0.678	0.684	0.568	0.991	0.489	0.019
LEG	3.358	0.69	1.756	3.053	0.523	2.136
S (F)	0.868	0.582	0.242			
EB	0.462	0.339	2.28	0.392	0.293	3.188
HVG × S	0.61	0.848	0.583			
HEG × S	2.188	0.859	0.912			
LEG × S	1.006	0.829	0.007			
HVG × EB	2.357	0.383	2.237	2.535	0.369	2.52
HEG × EB	2.455	0.569	1.578	2.318	0.542	1.551
LEG × EB	6.73	0.638	2.99	5.759	0.622	2.816
S × EB	0.677	0.34	1.151			
Random effects						
σ^2	3.29			3.29		

τ_{00}	0.00 REP:T; 0.89 REP	0.00 REP:T; 0.86 REP
N	6 REP; 4 T	6 REP; 4 T
Observations	237	237
Marg. R^2; Cond. R^2	0.108; NA	0.090; NA

REFERENCES

Bókony, V., Lendvai, Á.Z., Vágási, C.I., Pătraș, L., Pap, P.L., Németh, J., *et al.* (2014). Necessity
or capacity? Physiological state predicts problem-solving performance in house sparrows.
Behav. Ecol., 25, 124–135.

Davis, A.K., Maney, D.L. & Maerz, J.C. (2008). The use of leukocyte profiles to measure stress
in vertebrates: a review for ecologists. *Funct. Ecol.*, 22, 760–772.

Dingemanse, N.J., Both, C., Drent, P.J., van Oers, K. & van Noordwijk, A.J. (2002).
Repeatability and heritability of exploratory behaviour in great tits from the wild. *Anim.*
Behav., 64, 929–938.

Matson, K.D., Ricklefs, R.E. & Klasing, K.C. (2005). A hemolysis–hemagglutination assay for
characterizing constitutive innate humoral immunity in wild and domestic birds. *Dev. Comp.*
Immunol., 29, 275–286.

Nakagawa, S. & Schielzeth, H. (2010). Repeatability for Gaussian and non-Gaussian data: a
practical guide for biologists. *Biol. Rev.*, 85, 935–956.

Pap, P.L., Czirják, G.Á., Vágási, C.I., Barta, Z. & Hasselquist, D. (2010). Sexual dimorphism in

immune function changes during the annual cycle in house sparrows. *Naturwissenschaften*,

97, 891–901.

Pap, P.L., Vágási, C.I., Czirják, G.Á. & Barta, Z. (2008). Diet quality affects postnuptial molting

and feather quality of the house sparrow (*Passer domesticus*): interaction with humoral

immune function? *Can. J. Zool.*, 86, 834–842.

Pap, P.L., Vágási, C.I., Czirják, G.Á., Titilincu, A., Pintea, A., Osváth, G., *et al.* (2011). The

effect of coccidians on the condition and immune profile of molting house sparrows (*Passer*

domesticus). *Auk*, 128, 330–339.

Pap, P.L., Vágási, C.I., Vincze, O., Osváth, G., Veres-Szászka, J. & Czirják, G.Á. (2015).

Physiological pace of life: the link between constitutive immunity, developmental period,

and metabolic rate in European birds. *Oecologia*, 177, 147–158.

Peig, J. & Green, A.J. (2009). New perspectives for estimating body condition from mass/length

data: the scaled mass index as an alternative method. *Oikos*, 118, 1883–1891.

Schielzeth, H. (2010). Simple means to improve the interpretability of regression coefficients.

Methods Ecol. Evol., 1, 103–113.

Stoffel, M.A., Nakagawa, S. & Schielzeth, H. (2017). rptR: repeatability estimation and variance

decomposition by generalized linear mixed-effects models. *Methods Ecol. Evol.*, 8, 1639–

1644.

Vágási, C.I., Pap, P.L. & Barta, Z. (2010). Haste makes waste: accelerated molt adversely affects

the expression of melanin-based and depigmented plumage ornaments in house sparrows.

PLoS One, 5, e14215.

Vágási, C.I., Pap, P.L., Vincze, O., Benkő, Z., Marton, A. & Barta, Z. (2012). Haste makes waste

but condition matters: molt rate–feather quality trade-off in a sedentary songbird. *PLoS One*,

7, e40651.