Social groups with diverse personalities mitigate physiological stress in a songbird

Csongor I. Vágási[1,2,†], Attila Fülöp[3,4,†], Gergely Osváth[1,2,5], Péter L. Pap[1,2], Janka Pénzes[1], Zoltán Benkő[1,6], Ádám Z. Lendvai[2] and Zoltán Barta[3]

[1]Evolutionary Ecology Group, Hungarian Department of Biology and Ecology, Babeş-Bolyai University, Cluj-Napoca, Romania

[2]Department of Evolutionary Zoology, University of Debrecen, Debrecen, Hungary

[3]MTA-DE Behavioural Ecology Research Group, Department of Evolutionary Zoology, University of Debrecen, Debrecen, Hungary

[4]Juhász-Nagy Pál Doctoral School, University of Debrecen, Debrecen, Hungary

[5]Museum of Zoology, Babe□-Bolyai University, Cluj-Napoca, Romania

[6]Romanian Ornithological Society/BirdLife Romania, Cluj-Napoca, Romania

[†]These authors shared first authorship.

ORCID: CIV, 0000-0002-8736-2391; AF, 0000-0001-5337-336X; PLP: 0000-0002-3659-7684; ÁZL, 0000-0002-8953-920X; ZBa, 0000-0002-7121-9865

Author for correspondence:

Csongor I. Vágási and Zoltán Barta

e-mail: CIV, csvagasi@gmail.com; ZBa, barta.zoltan@science.unideb.hu

Social groups often consist of diverse phenotypes, including personality types, and this diversity

is known to affect the functioning of the group as a whole. Social selection theory proposes that

group composition (i.e. social environment) also influences the performance of individual group

members. However, the effect of group behavioural composition on group members remains

largely unexplored, and it is still contentious whether individuals benefit more in a social

environment with homogeneous or diverse behavioural composition. We experimentally formed

groups of house sparrows *Passer domesticus* with high and low diversity of personality

(exploratory behaviour), and found that their physiological state (body condition, physiological

stress and oxidative stress) improved with increasing group-level diversity of personality. These

findings demonstrate that group personality composition affects the condition of group members

and individuals benefit from social heterosis (i.e. associating with a diverse set of behavioural

types). This aspect of the social life can play a key role in affiliation rules of social animals and

might explain the evolutionary coexistence of different personalities in nature.

Keywords:

affiliation, group composition, personality, physiology, social environment, social heterosis

Subject Category:

evolution

Subject Areas:

ecology, evolution, physiology

1. Introduction

Social groups usually consist of a mixture of members with diverse phenotypes [1]. Variation within groups can occur in morphology (e.g. size), behavioural traits (e.g. reactive and proactive behavioural types [2], or social roles (e.g. leaders/followers in human *Homo sapiens* teams [3], or producers/scroungers in tree sparrow *Passer montanus* flocks [4]). Group composition has implications for emergent group-level processes such as decision-making, which ultimately drive group functioning (reviewed in [1,5]). Ethnic diversity can, for instance, have a positive effect on research teams' scientific performance [6].

Personality, the consistent among-individual differences in behavioural phenotype [7], has strong relevance for social life [8]. Social groups can largely differ in their personality composition some being more homogenous, while others more heterogeneous [1]. Groups' personality composition can have effects at both the level of group as a whole ('upstream effects') and the level of individual group members ('downstream effects'). The group-level consequences of groups' personality composition have mostly been assessed in human teams [9,10], where personality composition may affect team performance, albeit in an inconsistent manner [11–13]. In non-human organisms, group personality composition influences within-group social network structure, collective behaviour, and group performance [5,14–17]. Although individual-level aspects of sociality, e.g. rank in the dominance hierarchy, have been shown to influence the state of individual members (reviewed by [18,19]), the downstream effects of group behavioural composition on individual state is surprisingly little scrutinised [1]. This happens despite that social selection theory postulates that the fitness of an individual is contingent upon the phenotype of those with whom it affiliates (i.e. social environment; [15,20]).

Health state can have a strong influence on the individual's social behaviour [21] and, at

the same time, is moulded by changes of individual's social environment [19]. Therefore, it is

reasonable to assume that physiological condition (e.g. body condition, stress physiology,

oxidative stress and immune capacity) of group members is also influenced downstream by the

personality composition of their group. Correlative studies on human work teams found that age

and gender composition can be associated with subjective self-reported health impairment [22],

but human studies with experimental manipulation of group composition and actual health

measurements are still lacking. Earlier animal studies mostly assessed how individuals' position

within the social structure (e.g. rank in dominance hierarchy, an individual-level social attribute)

affected their health or physiology (reviewed by [18,19]). Experimental studies on animals that

tested directly whether group behavioural composition (a group-level social attribute) affects the

stress level and condition of group members are very scarce and each involve livestock species

[23,24]. No experimental study addressed this question in wild animals.

How could group composition affect the state of group members? Several non-exclusive

mechanisms can play a role. Diverse groups provide more opportunities for specialization [1,25]

and are more likely to host keystone individuals, which are influential individuals with

disproportionately large effect on other group members and/or overall group functioning [26].

Both role specialization and keystone individuals can lead to superior group-level performance

(upstream effect). Indeed, great tit *Parus major* affiliations consisting of diverse personalities

show the most effective coordinated action when exploring a habitat patch [5]. Similarly, a

mixture of shy and bold guppies *Poecilia reticulata* can be advantageous in reducing the trade-off

between exploring novel foraging tasks and antipredator vigilance [16]. The minority of keystone

individuals can also substantially affect group-level behaviour and performance [14].

Complementarity of different personalities might also be advantageous in groups of diverse

personalities. Stickleback *Gasterosteus aculeatus* shoals solve better a two-stage food acquisition

problem when the shoal contains fish that have experience with stage one and fish that have

experience with stage two (termed "experience-pooling"; [27]). Finally, groups with diverse

behavioural composition might experience less aggression in pigs *Sus scrofa domesticus* [23].

These group-level advantages of diversely composed groups can bring about higher individual

performance (downstream effect) in terms of either fitness or condition [17] by reducing stress

exposure and ultimately leaving group members in better physiological condition. Here we asked

whether group-level diversity of personalities might influence the physiological state of

individual group members.

We conducted an experimental study to explore whether manipulated personality

composition of groups *per se* or in interaction with individual personality type affects the

physiological condition of group members. Given that diverse groups might have advantages for

group functioning (see above) and potentially there is less aggression in behaviourally diverse

groups [23], we predicted that individuals in groups with more diverse personality composition

would improve their physiological condition as compared with individuals in groups with more

homogeneous personality composition. In this case, improved physiological condition of group

members is a legacy of being part of a diversely composed and better functioning group and each

group member shares this legacy. We do not know whether this potential benefit is indeed

uniform for each member. Alternatively, some members might harvest more the benefits of better

group-level functioning to the detriment of other group members. We assessed this prediction by

testing whether personality diversity of the groups interact with individual personality type to

influence physiological condition. A significant interaction might suggest that individuals that

either match or mismatch the group's personality composition benefit more than other group

members do.

2. Material and methods

(a) Study species

The house sparrow *Passer domesticus* is arguably one of the most popular model organisms in
animal ecology and evolution [28]. It is an ideal candidate to study sociality, since it exhibits a
wide spectrum of social behaviour including colonial breeding, social foraging or communal
roosting [29]. At our study site, house sparrows are year-round residents, breed in cavities of stall
buildings at the cattle farm and forage in flocks of various sizes, especially outside the breeding
season. Flock sizes vary from small to medium, containing from a few birds up to some dozen
individuals similarly to other populations (see [29]).

(b) Study protocol

The study is based on a large sample of 240 house sparrows. The same study protocol was used
in six study replicates. We captured 40 sparrows (1:1 sex ratio) per each study replicate
(Electronic Supplementary Material, ESM table S1). These 40 birds were divided into four
treatment groups (see below) consisting of 10 birds each, which yielded 24 social groups for the
entire study (four treatment groups per study replicate × six study replicates). There was no
significant difference in sex ratio between the treatment groups in any of the study replicates (χ^2
test, all $p > 0.362$).

The study timeline in each study replicate was as follows. Upon capture (day 0), birds

were marked with an aluminium ring, and their sex and body mass was recorded. Then they were

housed in indoor aviaries for 18 days at the campus of Babeș-Bolyai University, Cluj-Napoca,

Romania. On days 5–7, we recorded exploratory behaviour as a well-established and ecologically

relevant axis of personality [30] following the novel environment test of Dingemanse et al. [31].

At day 9, we measured the body mass and tarsus length of the birds, and took a pre-treatment

blood sample. Then the birds were allocated according to an *a priori* defined, stratification based

protocol (for details, see ESM) into one of four social treatment groups of 10 birds each:

'random' (random subsample of birds of a given replicate), 'low exploratory' (only birds with

low exploratory scores), 'high exploratory' (only birds with high scores), and 'variable' (equal

mixture of birds with either low or high scores) (see details of Social treatment methods in ESM).

Social treatment protocol successfully created differences in the mean and variance of

exploration score (personality) among groups as recommended by Farine et al. [1] (ESM figure

S1). The mean exploratory score was the smallest in the low exploratory group, intermediate in

the random and variable groups, and the largest in the high exploratory group (Linear mixed-

effects model with study replicate as random factor, treatment group effect: $\chi^2 = 632.92$, df = 3, *p*

< 0.001; figure S1a). The variance of exploratory score was the lowest in the low and high

exploratory groups, intermediate in the random group and the highest in the variable groups

(treatment group effect: $\chi^2 = 401.78$, df = 3, $p < 0.001$; low vs. high exploratory: $\chi^2 = 1.93$, df = 1,

$p = 0.165$; random vs. variable: $\chi^2 = 58.58$, df = 1, $p < 0.001$; figure S1b). The significant

difference in variance of exploratory score between the random and the variable groups is due to

the former being an even sample of the entire range of exploratory behaviour, while the latter

being a mixture of low exploratory and high exploratory individuals in equal proportion (see

details of Social treatment methods and figure S2 in ESM). Therefore, as an *a posteriori*

characterisation of the birds' social environment, we calculated the Shannon diversity index of

exploratory behaviour for each social group of 10 sparrows by dividing exploration values into

10 ordered categories of roughly equal sizes. As expected, the Shannon diversity index was the

lowest in the low and high exploratory groups, intermediate in the variable group, and the highest

in the random group (treatment group effect: $\chi^2 = 68.118$, df = 3, $p < 0.001$; low vs. high

exploratory: $\chi^2 = 1.30$, df = 1, $p = 0.254$; random vs. variable: $\chi^2 = 11.73$, df = 1, $p < 0.001$; figure

S1c).

The social treatment period lasted nine days until day 18, when we measured again the

body mass and took a second blood sample to measure the post-treatment physiological

condition. Physiological state was characterised by measuring body condition (scaled body mass

index, SMI), heterophil-to-lymphocyte (H/L) ratio, oxidative stress (damage to lipids –

malondialdehyde concentration, MDA), and innate immune capacity (natural antibodies –

agglutination score; complement system – lysis score) both during the pre- and post-treatment

sampling events. These traits were chosen for the following reasons. Changes in body condition

usually take more time than changes in other physiological traits. Among the factors that affect

body condition, exposure to stress stimuli is known to reduce body condition [32]. Ultimately,

impaired body condition can have widespread consequences for the organism. H/L ratio has been

shown to correlate with the glucocorticoid stress response governed by hypothalamic-pituitary-

adrenal axis [33,34], but is less sensitive to handling stress as compared with plasma levels of

corticosterone (the main glucocorticoid in birds). H/L ratio therefore can indicate the degree of

physiological stress. MDA is a versatile marker of oxidative stress for two reasons. MDA

indicates the damage to vital cell membrane lipids, which has substantial adverse consequences

from the cell level to the organism level. Besides being a direct measure of oxidative damage,

MDA is a pro-oxidant itself with a long half-life and hence can reach far from its site of origin

damaging other vital macromolecules [35]. Agglutination and lysis capacity of the plasma

describes the activity of the humoral innate immune system and hence is an indicator of the first

line of defence of vertebrate hosts against invading microorganisms [36]. There was no

significant difference among social treatment groups in the pre-treatment values of the five

physiological response variables (all $p > 0.188$). We provide a more detailed description of study

timeline, captivity conditions, measurement of exploratory behaviour, assignment to social

treatment groups, blood sampling and physiological measurement methods in the ESM.

(c) Statistical procedures

The statistical analyses were carried out in the R statistical environment (ver. 4.03; [37]). Since

we used a repeated-measures approach to analyse our data, each physiological variable contained

the values of both sampling events (i.e. pre-treatment and post-treatment values). Each

physiological trait was analysed in a separate statistical model. Heterophil-to-lymphocyte ratio

was arcsine square root transformed, malondialdehyde level was log-transformed and exploration

score was log(x+1)-transformed to normalize their distributions, while haemagglutination and

haemolysis scores were converted into binary variables (0 for absence and 1 for presence of

agglutination or lysis) because they were highly zero-inflated. To improve model convergence,

the continuous dependent variables (i.e. body condition, heterophil-to-lymphocyte ratio and

malondialdehyde concentration) and the continuous predictor variables (exploration score and

Shannon diversity) were Z-transformed to have zero mean and unit standard deviation [38].

In the first set of models, we assessed the effect of social treatment and the effect of

sampling event × social treatment interaction on the individual physiological responses of

sparrows. The explanatory variables were the same in all models as follows: sex (male or

female), social treatment (random, variable, high-exploratory and low-exploratory) and sampling

event (pre-treatment and post-treatment) were set as fixed factors, and individual exploratory

score as a continuous predictor. In addition, all second-order interactions between the four

explanatory variables were also tested. Note that, in this setup, a significant interaction with

sampling event indicates that the rate of change in the response variable is influenced by the other

explanatory variable in the interaction. Study replicate, group ID nested within study replicate,

and individual ID nested within study replicate and group ID were entered as random factors. We

used linear mixed-effects models with normal error distribution (LMMs; 'lmer' function of the R

package 'lme4'; [39]) for body condition, heterophil-to-lymphocyte ratio and malondialdehyde,

while we used generalized linear mixed-effects models with binomial error distribution

(GLMMs; 'glmer' function of the R package 'lme4') for agglutination and lysis scores. The

assumption of homogeneity of variances among treatment groups were met for each response

variable of the LMM models (Levene test, all $p > 0.195$). We assessed the fulfilment of model

assumptions by graphical diagnosis; all assumptions were met for each model. Each model was

simplified to obtain minimal adequate models (MAMs) containing only significant main effects

or their interactions by sequentially dropping predictors with non-significant ($p > 0.050$) effects

using the 'drop1' R function. The sampling event × social treatment interaction and its main

effects were kept in the model even if they were non-significant because our main interest is

related to the sampling event × social treatment interaction.

In the second set of models, we assessed the effect of Shannon diversity of group

personality and the effect of sampling event × Shannon diversity interaction on the individual

physiological responses of sparrows. For this, we used a similar approach as in the first model set

with the only difference that we entered groups' Shannon diversity of exploration as a continuous

predictor in the models instead of the fixed effect of social treatment.

The reported significance levels were calculated using type II Wald Chi-square tests using

the 'Anova' function of the R package 'car' [40]. The post-hoc comparisons of individual

changes in physiological variables between the two sampling events as a function of different

treatment groups (first set of models) or as a function of groups' Shannon diversity of exploration

(second set of models) were conducted using the R package 'emmeans' (functions 'emmeans'

and 'emtrends', respectively; [41]). Tables 1 and 2 present the type II ANOVA results of MAMs

for the first and second set of models, respectively. ESM tables S2 and S3 present the parameter

estimates of both the full models and the MAMs for the first and second set of models,

respectively.

3. Results

(a) Social treatment effects

In the first model set, we assessed the effect of social treatment on the individual's physiological

responses. The main effect of social treatment was non-significant for each response variable,

except for a marginal effect in case of the complement system of the innate immunity (i.e. lysis

score; table 1, table S2). Importantly, the sampling event $\times$ social treatment interaction was

significant for body condition and oxidative stress, and marginally significant for heterophil-to-

lymphocyte ratio, while it was non-significant for agglutination and lysis (table 1, table S2, figure

1).

Body condition increased only in the random group and decreased in the other three

groups (figure 1a). The improvement in body condition during the treatment period was

significantly higher in the random group as compared with the other three groups (see sampling

event $\times$ social treatment interaction in table S2a). Post-hoc tests revealed that body condition

improved significantly in the random group between the two sampling events ($\beta = -0.232$, s.e. =

0.074, df = 236, $t = 3.123$, $p = 0.008$), while body condition was not affected by the experimental

treatment in the other three treatment groups (all $p > 0.198$).

248   Heterophil-to-lymphocyte ratio (an indicator of physiological stress) decreased only in the

random group, while increased in the other three groups (figure 1b). The rate of increase in

heterophil-to-lymphocyte ratio was significantly higher in the high-exploratory group, and

marginally higher in the low-exploratory and variable groups as compared with the random group

(see sampling event $\times$ social treatment interaction in table S2b). Post-hoc tests showed a weak

decrease in heterophil-to-lymphocyte ratio in the random group between the two sampling events

($\beta = 0.378$, s.e. = 0.166, df = 235, $t = 2.270$, $p = 0.093$), while it remained unchanged in the other

three groups (all $p > 0.484$).

256   Malondialdehyde levels (i.e. oxidative damage to lipids) increased in high- and low-

exploratory groups, but decreased in the variable and random groups (figure 1c). The increase in

oxidative stress during the treatment period was significantly higher in the low-exploratory group

and marginally higher in the high-exploratory group as compared with the random group, while

the variable and random groups did not differ (see sampling event $\times$ social treatment interaction

in table S2c). The post-hoc tests revealed that individuals from the variable group showed a

marginally significant decrease in oxidative stress levels between the two sampling events ($\beta =$

0.439, s.e. = 0.179, df = 233, $t = 2.452$, $p = 0.058$), whereas birds from the other three treatment

groups were not affected by the experimental manipulation (all $p > 0.140$).

Individuals' personality was not associated with the individual responses in physiological

condition by itself or in interaction with social treatment, hence was dropped from the models

(table 1, table S2). Constitutive innate immunity improved during treatment as both agglutination

and lysis scores increased significantly between the two sampling events, and males had higher

body condition than females (table 1, table S2).

(b) Shannon diversity of group's personality

To assess the role of personality diversity of experimental social groups, we tested whether the

calculated Shannon diversity index predicted the individual's physiological responses to social

treatment. The main effect of Shannon diversity was non-significant for each response variable

(table 2, table S3). Importantly, the sampling event $\times$ Shannon diversity interaction was highly

significant for body condition and oxidative stress (figure 2), while it was non-significant for

heterophil-to-lymphocyte ratio and the activity of constitutive innate immunity as measured by

agglutination and lysis (table 2, table S3). Regarding the significant interactions, we found that

body condition increased ($\beta = 0.146$, s.e. $= 0.067$, df $= 24.9$, $t = 2.195$, $p = 0.038$), while oxidative

damage to lipids (i.e. malondialdehyde) decreased ($\beta = -0.170$, s.e. $= 0.067$, df $= 66.0$, $t = 2.532$,

$p = 0.014$) with increasing personality diversity in the post-treatment sample (i.e. after exposure

to social treatment), while these two response variables were unrelated to Shannon diversity in

the pre-treatment sample (i.e. prior to social treatment; all $p > 0.413$; table 2, table S3, figure 2).

The other three response variables (i.e. heterophil-to-lymphocyte ratio, and agglutination and

lysis) were unrelated to Shannon diversity in either the pre-treatment or the post-treatment

samples (all $p > 0.231$ and all $p > 0.578$, respectively).

4. Discussion

In the realm of human psychology, there is a long-standing debate on whether uniformly or diversely composed teams perform better [11,12], but almost nothing has been known about whether group behavioural composition affects the physiological condition of group members either in humans or other animals. Pigs housed in groups of uniform behavioural composition or in groups with a mixture of behavioural types did not differ in their glucocorticoid stress response or weight gain [23,24]. Here we showed that "variety is delighting," as condition significantly improved, heterophil-to-lymphocyte ratio (an indicator of physiological stress) was marginally reduced and oxidative stress level was significantly reduced for house sparrows living in a social environment with diverse personalities. Therefore, living in social groups with diverse composition can provide at least short-term benefits in terms of reduced physiological stress and superior condition. The finding of no significant interactions between social treatment and individual's exploration suggests that all individuals in diverse groups similarly enjoy the benefits.

It is important to note that we found a positive effect of social diversity on health in a set-up where ample amount of food was available. This suggests that the benefits of social diversity might arise because of the innate working of the group (e.g. the type and intensity of aggressive interactions between members) rather than being the consequence of more obvious benefits like improved foraging success, habitat exploration, defence against predators, and decision-making. Therefore, preference for bonding with dissimilar individuals (i.e. heterophily), and the resulting better health state of individuals in diverse groups might reinforce the improved group-level outcomes creating a positive feedback loop [42] between group-level and individual-level

performances. Indeed, affiliations of great tits and simulated social groups consisting of diverse

personalities show the most effective coordinated action when exploring a habitat patch [5],

which indicates an improved group-level performance. Note that the group compositions

simulated in this modelling study [5] are highly similar to our experimental setup. Therefore, our

study suggests that social groups composed of diverse personalities also improve individual-level

performance. This simultaneous group- and individual-level benefits of diversely composed

groups can generate a positive feedback loop, and such a feedback loop can then drive the

evolutionary maintenance of heterophily [43].

Consistency of personality traits places a constraint on individuals because one is either

more reactive (shy, neophobic, less exploratory, less aggressive) or more proactive (bold, novelty

seeking, more exploratory, more aggressive) [2]. However, if different personalities affiliate, they

can share mutual benefits; a concept termed social heterosis [44]. Social heterosis in associations

of dissimilar personalities thus can explain why behavioural (and genetic) diversity can

evolutionarily persist [44]. Negative frequency-dependent selection is another evolutionary

explanation for the existence of behavioural polymorphisms (producers–scroungers, hawks–

doves, and leaders–followers), and has been shown to maintain diversity in group personality

composition [45]. Our findings bring evidence to these potential explanations. Living in social

groups with diverse composition can provide benefits in terms of reduced physiological stress.

Because physiological state in one life-history stage can have carry-over effects on performance

in the subsequent life-history stage (e.g. from wintering to the breeding stage, see e.g. the case of

American redstart *Setophaga ruticilla* [46,47]), our results might provide a physiological

mechanism that could be responsible for the evolutionary maintenance of behavioural diversity in

social groups. These findings, thus, bring helpful insight into the study of social evolution, which

is a fundamental question in biology and has implications for human work teams. Although the

role of group composition in human team performance is still contentious [10], there is some

evidence showing that heterophily is advantageous in project groups that are less stable in time

and are engaged in creative tasks, but disadvantageous in production groups that are stable in

time and are engaged in routine tasks [12].

Ethics. None of the birds died during the study and all of them were released at the site of

capture in good health on day 18. The study complies with the ethical guidelines of the Babeș-

Bolyai University (permit no. 30792) and the European laws regarding animal welfare, and

adheres to the ASAB guidelines for the use of animals in behavioural research.

Data accessibility. Should the manuscript be accepted, all data supporting the results will be

provided as ESM or deposited in Dryad repository.

Authors' contributions. C.I.V., A.F., and Z.Ba. conceived and designed the study. C.I.V., A.F.,

P.L.P., O.G., J.P., and Z.Be. performed research. C.I.V., J.P., O.G., and Á.Z.L. contributed new

reagents/analytic tools. C.I.V., A.F., and Z.Ba. analysed data. C.I.V., A.F., and Z.Ba. drafted the

manuscript with major input from Á.Z.L. and P.L.P. All authors contributed revisions and

approved the final version of the manuscript. The authors declare no conflict of interest.

Competing interests. We declare we have no competing interests.

Funding. The study was financed by the Hungarian National Research, Development and

Innovation Office (#K112527 to Z.Ba.) and a Postdoctoral Fellowship of the Hungarian

Academy of Sciences (to C.I.V.). C.I.V. was supported by the János Bolyai Fellowship of the

Hungarian Academy of Sciences and by the Hungarian National Research, Development and

Innovation Office (#PD121166), A.F. by a PhD scholarship and through the New National

Excellence Program of the Ministry of Human Capacities of Hungary (#ÚNKP-16-3-IV), Á.Z.L.

by the Hungarian National Research, Development and Innovation Office (#K113108), P.L.P. by

a János Bolyai Fellowship of the Hungarian Academy of Sciences, and Z.Ba. by the Higher

Education Institutional Excellence Program (#NKFIH-1150-6/2019) of the Ministry of

Innovation and Technology in Hungary, within the framework of the FIK-DE Behavioural

Ecology Research Group thematic program of the University of Debrecen.

Acknowledgements. We thank A. Jakab for providing accommodation for A.F. during the

experiment, M. Bán for technical assistance, and V. Bókony and F. Bonier for highly useful

comments. The constructive comments raised by two anonymous reviewers considerably

improved our manuscript.

References

1.   Farine DR, Montiglio P-O, Spiegel O. 2015 From individuals to groups and back: the

evolutionary implications of group phenotypic composition. *Trends Ecol. Evol.* **30**, 609–

621. (doi:10.1016/j.tree.2015.07.005)

2.   Sih A, Bell A, Johnson JC. 2004 Behavioral syndromes: an ecological and evolutionary

overview. *Trends Ecol. Evol.* **19**, 372–378. (doi:10.1016/j.tree.2004.04.009)

3.   King AJ, Johnson DDP, Van Vugt M. 2009 The origins and evolution of leadership. *Curr.*

Biol. **19**, R911–R916. (doi:10.1016/j.cub.2009.07.027)

4.   Fülöp A, Németh Z, Kocsis B, Deák-Molnár B, Bozsoky T, Barta Z. 2019 Personality and

social foraging tactic use in free-living Eurasian tree sparrows (*Passer montanus*). *Behav.*

Ecol. **30**, 894–903. (doi:10.1093/beheco/arz026)

5. Aplin LM, Farine DR, Mann RP, Sheldon BC. 2014 Individual-level personality

influences social foraging and collective behaviour in wild birds. *Proc. R. Soc. London B*

281, 20141016. (doi:10.1098/rspb.2014.1016)

6. AlShebli BK, Rahwan T, Woon WL. 2018 The preeminence of ethnic diversity in

scientific collaboration. *Nat. Commun.* **9**, 5163. (doi:10.1038/s41467-018-07634-8)

7. Réale D, Reader SM, Sol D, McDougall PT, Dingemanse NJ. 2007 Integrating animal

temperament within ecology and evolution. *Biol. Rev.* **82**, 291–318. (doi:10.1111/j.1469-

185X.2007.00010.x)

8. Webster MM, Ward AJW. 2011 Personality and social context. *Biol. Rev.* **86**, 759–773.

(doi:10.1111/j.1469-185X.2010.00169.x)

9. Dyer JRG, Johansson A, Helbing D, Couzin ID, Krause J. 2009 Leadership, consensus

decision making and collective behaviour in humans. *Philos. Trans. R. Soc. London B* **364**,

781–789. (doi:10.1098/rstb.2008.0233)

10. van Knippenberg D, Schippers MC. 2007 Work group diversity. *Annu. Rev. Psychol.* **58**,

515–541. (doi:10.1146/annurev.psych.58.110405.085546)

11. Barrick MR, Stewart GL, Neubert MJ, Mount MK. 1998 Relating member ability and

personality to work-team processes and team effectiveness. *J. Appl. Psychol.* **83**, 377–391.

(doi:10.1037/0021-9010.83.3.377)

12. Stewart GL. 2006 A meta-analytic review of relationships between team design features

and team performance. *J. Manage.* **32**, 29–55. (doi:10.1177/0149206305277792)

13.   Halfhill T, Sundstrom E, Lahner J, Calderone W, Nielsen TM. 2005 Group personality

composition and group effectiveness. *Small Gr. Res.* **36**, 83–105.

(doi:10.1177/1046496404268538)

14.   Brown C, Irving E. 2014 Individual personality traits influence group exploration in a feral

guppy population. *Behav. Ecol.* **25**, 95–101. (doi:10.1093/beheco/art090)

15.   Sih A, Watters J V. 2005 The mix matters: behavioural types and group dynamics in water

striders. *Behaviour* **142**, 1417–1431. (doi:10.1163/156853905774539454)

16.   Dyer JRG, Croft DP, Morrell LJ, Krause J. 2009 Shoal composition determines foraging

success in the guppy. *Behav. Ecol.* **20**, 165–171. (doi:10.1093/beheco/arn129)

17.   Bengston SE, Jandt JM. 2014 The development of collective personality: the ontogenetic

drivers of behavioral variation across groups. *Front. Ecol. Evol.* **2**, 81.

(doi:10.3389/fevo.2014.00081)

18.   Sapolsky RM. 2005 The influence of social hierarchy on primate health. *Science* **308**, 648–

652. (doi:10.1126/science.1106477)

19.   Creel S, Dantzer B, Goymann W, Rubenstein DR. 2013 The ecology of stress: effects of

the social environment. *Funct. Ecol.* **27**, 66–80. (doi:10.1111/j.1365-2435.2012.02029.x)

20.   Wolf JB, Brodie III ED, Moore AJ. 1999 Interacting phenotypes and the evolutionary

process. II. Selection resulting from social interactions. *Am. Nat.* **153**, 254–266.

(doi:10.1086/303168)

21.   Kelly AM, Vitousek MN. 2017 Dynamic modulation of sociality and aggression: an

examination of plasticity within endocrine and neuroendocrine systems. *Philos. Trans. R.*

Soc. London B **372**, 20160243. (doi:10.1098/rstb.2016.0243)

22.   Wegge J, Roth C, Neubach B, Schmidt K-H, Kanfer R. 2008 Age and gender diversity as

determinants of performance and health in a public organization: the role of task

complexity and group size. *J. Appl. Psychol.* **93**, 1301–1313. (doi:10.1037/a0012680)

23.   Hayne SM, Gonyou HW. 2006 Behavioural uniformity or diversity? Effects on behaviour

and performance following regrouping in pigs. *Appl. Anim. Behav. Sci.* **98**, 28–44.

(doi:10.1016/j.applanim.2005.08.007)

24.   van Erp-van der Kooij E, Kuijpers AH, van Eerdenburg FJCM, Dieleman SJ, Blankenstein

DM, Tielen MJM. 2003 Individual behavioural characteristics in pigs—influences of

group composition but no differences in cortisol responses. *Physiol. Behav.* **78**, 479–488.

(doi:10.1016/S0031-9384(03)00002-7)

25.   Bergmüller R, Taborsky M. 2010 Animal personality due to social niche specialisation.

Trends Ecol. Evol. **25**, 504–511. (doi:10.1016/j.tree.2010.06.012)

26.   Modlmeier AP, Keiser CN, Watters J V., Sih A, Pruitt JN. 2014 The keystone individual

concept: an ecological and evolutionary overview. *Anim. Behav.* **89**, 53–62.

(doi:10.1016/j.anbehav.2013.12.020)

27.   Webster MM, Whalen A, Laland KN. 2017 Fish pool their experience to solve problems

collectively. *Nat. Ecol. Evol.* **1**, 1–5. (doi:10.1038/s41559-017-0135)

28.   Hanson HE, Mathews NS, Hauber ME, Martin LB. 2020 The house sparrow in the service

of basic and applied biology. *eLife* **9**, e52803. (doi:10.7554/eLife.52803)

29.   Anderson TR. 2006 *Biology of the Ubiquitous House Sparrow: From Genes to*

440   *Populations*. New York, NY: Oxford University Press.

441   30.   Wolf M, Weissing FJ. 2012 Animal personalities: consequences for ecology and evolution.

442   *Trends Ecol. Evol.* **27**, 452–461. (doi:10.1016/j.tree.2012.05.001)

443   31.   Dingemanse NJ, Both C, Drent PJ, van Oers K, van Noordwijk AJ. 2002 Repeatability and

444   heritability of exploratory behaviour in great tits from the wild. *Anim. Behav.* **64**, 929–938.

445   (doi:10.1006/anbe.2002.2006)

446   32.   Breuner CW, Delehanty B, Boonstra R. 2013 Evaluating stress in natural populations of

447   vertebrates: total CORT is not good enough. *Funct. Ecol.* **27**, 24–36. (doi:10.1111/1365-

448   2435.12016)

449   33.   Davis AK, Maney DL, Maerz JC. 2008 The use of leukocyte profiles to measure stress in

450   vertebrates: a review for ecologists. *Funct. Ecol.* **22**, 760–772. (doi:10.1111/j.1365-

451   2435.2008.01467.x)

452   34.   Goessling JM, Kennedy H, Mendonça MT, Wilson AE. 2015 A meta-analysis of plasma

453   corticosterone and heterophil□: lymphocyte ratios - is there conservation of physiological

454   stress responses over time? *Funct. Ecol.* **29**, 1189–1196. (doi:10.1111/1365-2435.12442)

455   35.   Del Rio D, Stewart AJ, Pellegrini N. 2005 A review of recent studies on malondialdehyde

456   as toxic molecule and biological marker of oxidative stress. *Nutr. Metab. Cardiovasc. Dis.*

457   **15**, 316–328. (doi:10.1016/j.numecd.2005.05.003)

458   36.   Matson KD, Ricklefs RE, Klasing KC. 2005 A hemolysis–hemagglutination assay for

459   characterizing constitutive innate humoral immunity in wild and domestic birds. *Dev.*

460   *Comp. Immunol.* **29**, 275–286. (doi:10.1016/j.dci.2004.07.006)

461   37.   R Core Team. 2020 *R: a language and environment for statistical computing*. R

462   Foundation for Statistical Computing, Vienna, Austria. http://www.R-project.org/.

463   38.   Schielzeth H. 2010 Simple means to improve the interpretability of regression coefficients.

464   *Methods Ecol. Evol.* **1**, 103–113. (doi:10.1111/j.2041-210X.2010.00012.x)

465   39.   Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting linear mixed-effects models using

466   lme4. *J. Stat. Softw.* **67**, 1–48. (doi:10.18637/jss.v067.i01)

467   40.   Fox J, Weisberg S. 2011 *An {R} Companion to Applied Regression*. 2nd edn. Thousand

468   Oaks, CA: Sage. See http://socserv.socsci.mcmaster.ca/jfox/Books/Companion.

469   41.   Lenth R. 2020 *emmeans: Estimated Marginal Means, aka Least-Squares Means. R*

470   *package version 1.4.4*. https://CRAN.R-project.org/package=emmeans.

471   42.   Seebacher F, Krause J. 2017 Physiological mechanisms underlying animal social

472   behaviour. *Philos. Trans. R. Soc. London B* **372**, 20160231. (doi:10.1098/rstb.2016.0231)

473   43.   Pruitt JN, Riechert SE. 2011 Within-group behavioral variation promotes biased task

474   performance and the emergence of a defensive caste in a social spider. *Behav. Ecol.*

475   *Sociobiol.* **65**, 1055–1060. (doi:10.1007/s00265-010-1112-z)

476   44.   Nonacs P, Kapheim KM. 2007 Social heterosis and the maintenance of genetic diversity.

477   *J. Evol. Biol.* **20**, 2253–2265. (doi:10.1111/j.1420-9101.2007.01418.x)

478   45.   Pruitt JN, Riechert SE. 2009 Frequency-dependent success of cheaters during foraging

479   bouts might limit their spread within colonies of a socially polymorphic spider. *Evolution*

480   **63**, 2966–2973. (doi:10.1111/j.1558-5646.2009.00771.x)

46.   Reudink MW, Marra PP, Kyser TK, Boag PT, Langin KM, Ratcliffe LM. 2009 Non-

breeding season events influence sexual selection in a long-distance migratory bird. *Proc.*

R. Soc. London B **276**, 1619–1626. (doi:10.1098/rspb.2008.1452)

47.   Angelier F, Vleck CM, Holberton RL, Marra PP. 2013 Telomere length, non☐breeding

habitat and return rate in male American redstarts. *Funct. Ecol.* **27**, 342–350.

(doi:10.1111/1365-2435.12041)

Tables

Table 1. Minimal adequate models containing predictors of individual responses in physiological

state of house sparrows to social treatment. Statistically significant effects ($p \leq 0.050$) are marked

in bold, while marginally significant effects ($0.050 < p \leq 0.100$) in italic. SMI – Scaled Mass

Index (body condition), H/L ratio – heterophil-to-lymphocyte ratio (physiological stress

response), MDA – malondialdehyde (oxidative damage to lipids).

response	fixed effects	χ^2	df	p
SMI	sampling event	0.00	1	1.000
	treatment	3.70	3	0.296
	sex	**13.66**	**1**	**< 0.001**
	sampling event × treatment	**14.32**	**3**	**0.003**
H/L ratio	sampling event	0.00	1	1.000
	treatment	1.38	3	0.709
	sex	0.02	1	0.879
	sampling event × treatment	*7.56*	*3*	*0.056*
	sampling event × sex	**3.85**	**1**	**0.050**
MDA	sampling event	0.00	1	0.979
	treatment	1.04	3	0.792
	sampling event × treatment	**13.83**	**3**	**0.003**

agglutination	**sampling event**	**12.54**	**1**	**< 0.001**
	treatment	1.51	3	0.680
	sampling event × treatment	0.81	3	0.848
lysis	**sampling event**	**16.17**	**1**	**< 0.001**
	treatment	*7.09*	*3*	*0.069*
	sampling event × treatment	0.38	3	0.945

494   **Table 2.** Minimal adequate models containing predictors of individual responses in physiological

495   state of house sparrows to social treatment in relation with the groups' personality diversity.

496   Statistically significant effects ($p < 0.050$) are marked in bold. SMI – Scaled Mass Index (body

497   condition), H/L ratio – heterophil-to-lymphocyte ratio (physiological stress response), MDA –

498   malondialdehyde (oxidative damage to lipids).

response	fixed effects	χ^2	df	p
SMI	sampling event	0.00	1	1.000
	sex	**12.68**	**1**	**< 0.001**
	Shannon diversity	1.57	1	0.210
	sampling event × Shannon diversity	**14.44**	**1**	**< 0.001**
H/L ratio	sampling event	0.00	1	1.000
	Shannon diversity	0.00	1	0.977
	sampling event × Shannon diversity	0.62	1	0.431
MDA	sampling event	0.00	1	0.983
	Shannon diversity	1.37	1	0.241
	sampling event × Shannon diversity	**6.10**	**1**	**0.014**
agglutination	**sampling event**	**12.36**	**1**	**< 0.001**
	Shannon diversity	0.18	1	0.675
	sampling event × Shannon diversity	1.49	1	0.222
lysis	**sampling event**	**16.10**	**1**	**< 0.001**

	Shannon diversity	0.02	1	0.882
	sampling event × Shannon diversity	0.10	1	0.756

Figure legends and Figures

Figure 1. Physiological responses to social treatment: (a) body condition (SMI), (b) heterophil-to-lymphocyte ratio (H/L ratio, an indicator of physiological stress response), (c) oxidative damage to lipids (i.e. malondialdehyde, MDA), and (d–e) constitutive immune capacity, as expressed through agglutination score (d) and lysis score (e). Model-predicted mean ± s.e. are shown for each treatment group both at pre-treatment (dark bars) and post-treatment (light bars) sampling events. Male house sparrow drawing credit: Márton Zsoldos.

Figure 2. Physiological condition improved in social groups with diverse personalities. Relationships between personality diversity in social groups of house sparrows and individual responses to social treatment in terms of (a) body condition (SMI) and (b) oxidative damage to lipids (MDA). Open symbols and continuous lines denote the pre-treatment sampling, while the filled symbols and dashed lines denote the post-treatment sampling. Regression lines are model-predicted slopes with 95% confidence intervals (shaded areas).

Figure 1

515   **Figure 2**

516