On identifying the eukaryotic *N*-glycosylation scramblase by activity correlation profiling

Alice Verchère[1], Andrew Cowton[2], Aurelio Jenni[2,3], Monika Rauch[2], Robert Häner[4], Johannes Graumann[5,6], Peter Bütikofer[2], Anant K. Menon[1]

[1] Department of Biochemistry, Weill Cornell Medical College, 1300 York Ave, New York, New York 10065, USA
[2] Institute of Biochemistry and Molecular Medicine, University of Bern, Bühlstr. 28, CH-3012 Bern, Switzerland
[3] Graduate School for Cellular and Biochemical Sciences, University of Bern, Mittelstr. 43, CH-3012 Bern, Switzerland
[4] Department of Chemistry and Biochemistry, University of Bern, Freiestr. 3, CH-3012, Bern, Switzerland
[5] Max Planck Institute for Heart and Lung Research, W.G. Kerckhoff Institute, Ludwigstr. 43, Bad Nauheim, Germany
[6] German Centre for Cardiovascular Research (DZHK), Rhine-Main site, Max Planck Institute for Heart and Lung Research, Bad Nauheim, Germany

Correspondence:
Anant K. Menon (akm2003@med.cornell.edu)
Peter Bütikofer (peter.buetikofer@ibmm.unibe.ch)

Abstract

The canonical pathway of *N*-linked protein glycosylation in yeast and humans involves transfer of the oligosaccharide moiety from the glycolipid $Glc_3Man_9GlcNAc_2$-PP-dolichol to select asparagine residues in proteins that have been translocated into the lumen of the endoplasmic reticulum (ER). Synthesis of $Glc_3Man_9GlcNAc_2$-PP-dolichol occurs in two stages, producing first the key intermediate $Man_5GlcNAc_2$-PP-dolichol (M5-DLO) on the cytoplasmic face of the ER, followed by translocation of M5-DLO across the ER membrane where the remaining glycosyltransfer reactions occur to complete the structure. The scramblase protein that mediates the translocation of M5-DLO across the ER membrane has not been identified, but activity assays provide compelling evidence that it is an ER membrane protein with exquisite substrate specificity. Here we report on our progress in identifying the M5-DLO/*N*-glycosylation scramblase via a mass spectrometry-based 'activity correlation profiling' approach.

Introduction
Asparagine *(N)*-linked protein glycosylation is found in all domains of life, and is highly conserved in eukaryotes [1-5]. *N*-glycosylation takes place in the lumen of the endoplasmic reticulum (ER), when oligosaccharyltransferase (OST) attaches a pre-synthesized oligosaccharide ($Glc_3Man_9GlcNAc_2$ in yeast and humans) to asparagine residues within glycosylation sequons in newly translocated proteins. The $Glc_3Man_9GlcNAc_2$ oligosaccharide is built by sequentially adding sugars to dolichyl phosphate, an isoprenoid glycosyl carrier lipid with a very long hydrocarbon chain (Fig. 1A). Sugar addition occurs in two stages and on different sides of the ER membrane. The first stage comprises seven reactions in which the sugar nucleotide donors UDP-GlcNAc and

GDP-Man are used to produce the key intermediate $Man_5GlcNAc_2$-PP-dolichol (M5-DLO) on the cytoplasmic face of the ER (Fig. 1A). M5-DLO is then flipped to the luminal side of the ER where seven, lipid-mediated glycosyltransfer steps occur to generate $Glc_3Man_9GlcNAc_2$-PP-dolichol (G3M9-DLO). The mannose and glucose residues needed for these latter reactions are donated by mannose phosphate dolichol (MPD) and glucose phosphate dolichol (GPD), both of which are synthesized on the cytoplasmic face of the ER and flipped to the luminal leaflet.

The assembly of G3M9-DLO thus requires the rapid translocation of three polar glycolipids - M5-DLO, MPD and GPD - from the cytoplasmic to the luminal leaflet of the ER. Biochemical assays provide compelling evidence that these transport events do not require metabolic energy (they are ATP-independent) and that they are mediated by ER membrane proteins with exquisite substrate specificity [6-9]. The molecular identity of these proteins (scramblases) is not known [10,11]. The M5-DLO scramblase is critically important as *N*-glycosylation cannot occur without it (we therefore also refer to it as the *N*-glycosylation scramblase). The polytopic ER membrane protein Rft1 was proposed as the M5-DLO scramblase almost two decades ago [12,13], but subsequent work showed that whereas Rft1 is clearly important for *N*-glycosylation, it appears to have no direct role in translocating M5-DLO across the ER membrane [6,14-17].

We developed methods based on the work by Snider and colleagues [18] to assay M5-DLO scramblase activity in large unilamellar vesicles reconstituted with a mixture of ER membrane proteins derived from yeast or rat liver. We used radiolabeled M5-DLO as the reporter lipid, and the α-mannosyl-binding lectin Concanavalin A as a topological probe [6,7,14]. The assay is described in detail in Fig. 1B. Using this assay, we could show that the M5-DLO scramblase activity is ATP-independent, and highly structure specific [10]. Thus, higher order structures such as M7-DLO and M9-DLO are not scrambled efficiently [6], nor is a structural isomer of M5-DLO in which the mannose residues correspond to the arrangement seen in 'processed glycans' [7]. We were also able to resolve the M5-DLO scramblase from other scramblases responsible for transporting phospholipids and MPD [6,9]. Despite these advances, a straightforward purification of the scramblase proved elusive indicating that a different approach may be useful.

Here we describe a mass spectrometry-based 'activity correlation profiling' approach [19,20] to attack this problem. Conceptually, this method entails using a chromatography technique of choice (we used velocity gradient sedimentation) to resolve a crude mixture of ER-solubilized membrane proteins into a series of fractions. Measurement of M5-DLO scramblase activity in the fractions provides an activity profile. Simultaneous identification of the relative abundance of individual proteins in the fractions by Tandem Mass Tag (TMT) multiplexing mass spectrometry generates protein profiles. Quantitative comparison of protein profiles with the activity profiles, coupled with suitable data curation, leads to the identification of M5-DLO scramblase candidates for activity testing. We report on our progress using this approach.

Results

Identifying M5-DLO scramblase candidates by activity correlation profiling. We developed an activity correlation profiling approach to identify scramblase candidates from a crude mixture of detergent-solubilized ER membrane proteins (Fig. 2). The underlying concept involves resolving the proteins in the crude mixture into a number of fractions using any separation technique of choice, measuring the scramblase activity of each fraction to generate an activity profile and, in

parallel, using mass spectrometry to determine the relative abundance of individual proteins across the fractions (Fig. 2A). Proteins whose profiles best match the activity profile (Fig. 2B) are subjected to a curation process to identify scramblase candidates.

Total membranes from a yeast cell homogenate were washed with high salt to remove peripheral proteins, and then treated with ice-cold Triton X-100 to extract ER membrane proteins as described previously [6]. The resulting 'Triton Extract' (TE) was fractionated by velocity gradient sedimentation using a continuous glycerol gradient. Sedimentation standards were analyzed in a parallel gradient. Gradient fractions were collected manually from the top and, after measurement of refractive index, the fractions were passed over a desalting spin column to remove glycerol. A portion of each fraction from the middle section of the gradient (fractions 5-11, shown in preliminary tests to contain the majority of M5-DLO scramblase activity) was reconstituted into large unilamellar vesicles and assayed for scramblase activity using [^{3}H]M5-DLO (Fig. S1), whereas the remainder was taken for mass spectrometry using 6-plex tandem mass tag (TMT) reagents [21].

The M5-DLO scramblase activity assay has been described in detail previously [6,7,14] and is explained in Fig. 1B. The assay exploits the organic-solvent resistant interaction between M5-DLO and the lectin Concanavalin A (Con A) to capture M5-DLO molecules located on the external surface of large unilamellar vesicles, and separate them from those located in the inner leaflet. In the absence of a reconstituted functional scramblase, M5-DLO molecules located in the inner leaflet of the vesicles are inaccessible to Con A, whereas if the vesicles contain a scramblase then these molecules are translocated from the inner leaflet to the external leaflet and captured by the lectin. A key point is that the rate of scrambling is greater than the rate at which M5-DLO is captured by Con A [6,7], even if the assay is conducted on ice to reduce the scrambling rate. Thus, the assay reports end-point data, ranging from 50-100% M5-DLO captured depending on the fraction of vesicles in the sample that is reconstituted with a functional M5-DLO scramblase. The reconstitution of M5-DLO scramblase molecules into an ensemble of lipid vesicles is governed by Poisson statistics. To know the amount of M5-DLO scramblase in a mixture of proteins, e.g. in a fraction from the velocity gradient, we would therefore have to carry out reconstitutions with a range of protein concentrations and use a Poisson model to calculate scramblase abundance. As this is an impractical undertaking when assaying fractions from the velocity gradient, we decided instead to estimate scramblase abundance by reconstituting just enough protein so that the fraction with the highest amount of M5-DLO scramblase would report approximately 70% M5-DLO capture. We determined empirically that this could be accomplished by reconstituting half of each fraction.

The quality of separation achieved on the velocity gradient is shown in Fig. 3A. The activity profile has a peak in fractions 7-8, corresponding to a nominal sedimentation coefficient of 6-7S, whereas the bulk of the protein peaks earlier in the gradient, in fraction 6. Approximately 3000 proteins were identified in the mass spectrometric analysis (Tables S1 and S2). To learn how the various proteins were resolved on the gradient we determined the fraction in which each protein had its maximum abundance. This allowed us to obtain the average molecular weight of proteins that had their maximum abundance in each fraction. The data show that this average molecular weight increases from ~40 kDa (fraction 5) to ~80 kDa (fraction 8), remaining approximately constant at ~80 kDa over fractions 8-10 (Fig. S2). This result is consistent with the general expectation that protein monomers and small complexes will predominate in the initial fractions

and higher order complexes will be found in fractions 8-10. In line with this conclusion, the multi-subunit OST complex and the glycosyl-phosphatidylinositol (GPI) transamidase complex sediment relatively rapidly in the gradient, with the majority of their constituent subunits being maximally detected in fraction 7 (Fig. S3).

We next quantitatively compared the protein abundance profiles (obtained from each of two technical replicates) with the activity profile, identifying 280 proteins whose profile matched that of the scramblase activity profile with a correlation score of R ≥ 0.9 (Fig. S4) (Tables S1 and S2). We subjected this list to several data curation steps. Hypothesizing that the *N*-glycosylation scramblase would be an essential, ER-localized membrane protein, we discarded proteins that did not meet these criteria. Thus, 62 of the 280 proteins are known to be essential in yeast, and 5 correspond to ER-localized polytopic membrane proteins with 3 or more transmembrane spans as predicted by the TMHMM web server (Fig. 3B). These latter proteins are listed in Fig. 3C, and their profiles in comparison to the activity profile are shown in Figs. 3D, 3E. We included the highly correlated but non-essential protein Gpi7 because it belongs to the same GPI phospho-ethanolamine-transferase (P-EtN) family as Mcd4 and Gpi13 (Fig. 3F, Fig. S4) that were identified in our analysis; all three are polytopic membrane proteins with a large lumenal loop (Fig. 3G). Remarkably, the five candidates identified in our analyses are enzymes of the ER-localized GPI biosynthetic pathway [22], and three of these (Gpi13, Gpi14, Gpi2) are ubiquitously found in eukaryotes. We noticed that the profile of Stt3, the catalytic subunit of OST (Fig. 1A), matched closely with the activity profile (correlation score of 0.89, close to our cut-off value of 0.9). Because of its connection to *N*-glycosylation we included Stt3 in our list of candidates.

Biochemical test of M5-DLO scramblase candidates. To determine whether the six candidates (Gpi13, Gpi7, Mcd4, Gpi14, Gpi2 and Stt3) play a role in M5-DLO scrambling, we determined whether the M5-DLO scramblase activity present in TE could be eliminated by quantitatively and specifically immunodepleting the candidate from the TE prior to reconstitution and assay. We previously used a similar approach to show that opsin accounts for all the phospholipid scramblase activity in detergent-solubilized retinal disc membranes [23] and that Rft1 does not contribute to M5-DLO scramblase activity in TE [14].

We obtained haploid yeast strains in which M5-DLO scramblase candidate of interest is functionally expressed from its chromosomal locus with a C-terminal Tandem Affinity Purification (TAP) tag. The TAP-tagged protein is thus the only version of that protein being expressed in the cell. TE was prepared from strains expressing individual TAP-tagged M5-DLO scramblase candidates and either mock-treated, or treated with IgG resin to eliminate the TAP-tagged protein. Western blotting indicated that the TAP-tagged candidates were quantitatively removed by IgG-resin treatment, whereas an irrelevant ER membrane protein - Dpm1 - was still present in the TE (Fig. 4). We reconstituted equivalent amounts of mock-treated and IgG-treated TE into vesicles for M5-DLO scramblase activity assay. As our assay provides an end-point readout that reports on the fraction of vesicles that is equipped with a functional scramblase (see above), we expected to see a complete loss of activity for one of the candidates, thus pointing to its role in scrambling M5-DLO. Surprisingly, scramblase activity was not significantly affected by the removal of any of the candidates (Fig. 4). Thus, none of the candidates contributes significantly to the scramblase activity of TE. However, as described below, we were nevertheless interested in further exploring the possible function of the GPI P-EtN transferases in M5-DLO scrambling.

Re-consideration of the GPI P-EtN transferase proteins as scramblase candidates. The three GPI P-EtN transferases have very similar membrane topology (Fig. 3G), carry out similar reactions (they each transfer P-EtN from phosphatidylethanolamine (PE) to mannose residues in the GPI core structure (Fig. 3F)), and are present at equivalent levels in the cell (Fig. S5). Previous reports indicate that they have common functions in the cell in addition to their recognized roles in the GPI biosynthetic pathway. Thus, over-expression of any of the three proteins in yeast unexpectedly results in enhanced secretion of ATP, consistent with the possibility that each of these proteins is involved in the transport of ATP into the lumen of secretory compartments [24]. Also, the Mcd4 ortholog in *C. elegans* and human was demonstrated to have a non-canonical role in ER protein quality control, distinct from its role in GPI anchoring [25]. Based on these indicators, we considered the possibility that the three GPI P-EtN transferases may each (redundantly) have M5-DLO scramblase activity. As the proteins are expressed at similar levels (Fig. S5), elimination of any one of them would result in a lowering of our scramblase assay read-out by one-third, whereas elimination of all three would result in total removal of scramblase activity from the TE. Although we did not see any significant difference in the assay readout when testing TE samples from which individual P-EtN transferases had been eliminated (Fig. 4), this could be a result of the potentially poor sensitivity of the assay to changes that are less than 2-fold. As described next, rather than continue with the reconstitution approach to explore a possible role of P-EtN transferases in M5-DLO scrambling, we took a genetic approach.

***In vivo* test of Gpi13 as an M5-DLO scramblase.** At the level of a single eukaryotic cell, *N*-glycosylation is essential for cell viability whereas GPI anchoring is generally not essential. Thus, mammalian cells in culture are non-viable when *N*-glycosylation is disrupted [26], but their growth is unaffected by a deficiency in GPI anchoring [27]. The same holds true for the early diverging eukaryote *Trypanosoma brucei* [28-31]. Interestingly, GPI anchors in *T. brucei* have only a single P-EtN, linked to the third mannose residue, providing the means to attach the anchor to proteins. Thus *T. brucei* has an ortholog of Gpi13 (TbGPI13), but lacks orthologs of Mcd4 and Gpi7. To test whether Gpi13 is essential for M5-DLO scrambling, we decided to generate TbGpi13 null mutants of insect stage (procyclic) *T. brucei*. We predicted that if TbGPI13 was solely involved in GPI anchoring the *T. brucei* GPI13Δ cells would be viable. However, the GPI13Δ cells would be inviable if TbGPI13 played an essential role in scrambling M5-DLO for protein *N*-glycosylation.

We sequentially disrupted the two alleles of TbGPI13 in procyclic-stage *T. brucei* using hygromycin and geneticin drug resistance cassettes (Fig. 5A) and recovered viable GPI13Δ cells that grew well, albeit approximately 2-fold more slowly than wild-type cells (Fig. 5B). Although our ability to generate viable GPI13Δ cells immediately indicates that Gpi13 does not play an essential role in *N*-glycosylation, we nevertheless characterized the cells to verify the expected disruption of GPI anchoring and to investigate the *N*-glycosylation status of specific glycoproteins. We performed metabolic radiolabeling with [^{3}H]ethanolamine and analyzed both polar lipids and proteins. Whereas both PE and the P-EtN-containing GPI anchor precursor PP1 were labeled with [^{3}H]ethanolamine in wild-type (WT) cells, PP1 was not labeled in GPI13Δ cells (Fig. 5C). Consistent with this observation, SDS-PAGE/fluorography of protein extracts showed radiolabeled GPI-anchored GPEET procyclin (arrow head) in WT but not GPI13Δ cells (Fig. 5D). Of note,

radiolabeling of eukaryotic elongation factor 1A, a protein that is modified by ethanolamine phosphoglycerol [32], was similar in both WT and GPI13Δ cells (Fig. 5D).

We tested the intactness of the *N*-glycosylation pathway in GPI13Δ cells by assessing the glycosylation status of the lysosomal *N*-glycoprotein p67 which has 14 *N*-glycosylation sites. An anti-p67 immunoblot of whole cell protein samples from WT and GPI13Δ cells revealed no change in the molecular mass of p67, running at ~100 kDa, consistent with normal *N*-glycosylation [33]. In contrast, the mass of the glycoprotein was lower in a control *N*-glycosylation defective cell line (ALG3Δ) that cannot mannosylate M5-DLO *en route* to synthesis of the optimal DLO for the OST reaction [34] (Fig. 5E).

Finally, we examined the status of non-GPI-anchored procyclins in the GPI13Δ cells. We found that the main consequence of the lack of GPI anchoring was that the proteins remain sequestered inside the cell, failing to exit the ER (Fig. 5F-H).

We conclude that GPI anchoring is disrupted as expected in GPI13Δ cells, whereas *N*-glycosylation is unperturbed. The inability to synthesize GPI-anchored surface coat proteins may explain the somewhat slower growth rate of the cells [29,31](Fig. 5B). Taken together with the biochemical data presented in Fig. 4, these results show that Gpi13 is not essential for M5-DLO scrambling as its deficiency does not impact *N*-glycosylation in a living cell.

Discussion

We hypothesized that M5-DLO scramblase activity is essential in eukaryotes, and that if it is due to a single protein, then that protein would necessarily be essential for cell viability. Using activity correlation profiling (Fig. 2) in conjunction with a data filtering strategy (summarized in Fig. 3B) to implement this idea, we generated a list of 6 M5-DLO scramblase candidates (Fig. 3C). All six proteins are ER residents with multiple transmembrane spans, and they have well-established functions in GPI anchoring and protein *N*-glycosylation. However, our biochemical data indicate that none of them contributes significantly to M5-DLO scramblase activity. Thus, quantitative removal (using TAP-tagging and immunodepletion) of any one of the candidates from a crude mixture of ER membrane proteins (TE) prior to membrane reconstitution did not decrease the potency of the TE to populate an ensemble of large unilamellar vesicles with M5-DLO scramblase (Fig. 4).

Despite the negative outcome of our biochemical tests, we were especially interested in the family of three GPI P-EtN transferases - Gpi13, Gpi7 and Mcd4 - that we had identified as scramblase candidates. These proteins catalyze essentially identical reactions (Fig. 3F), and have a similar membrane topology comprising multiple transmembrane spans (Fig. 3G). Of considerable interest, all three were shown to participate in a function distinct from their role in GPI anchoring, i.e. ATP secretion [24]. In addition, Mcd4 was shown to have a non-canonical role in protein quality control [25]. These intriguing moonlighting functions attributed to the GPI P-EtN transferases raised the possibility that these three proteins could potentially each, independently, contribute to M5-DLO scramblase activity. Our inability to detect these activities biochemically could be due to an inherent lack of sensitivity to the assay as performed. Thus, in order to identify the potential contributions of each of these equally expressed proteins (Fig. S5), we would have to (i) perform the assay over a range of protein-phospholipid ratios, and (ii) use a Poisson model to detect the change in abundance of M5-DLO scramblase after biochemical

elimination of any one of the three proteins. Alternatively, we could have purified each of the proteins and tested them for scramblase activity after reconstitution. Rather than carry out this latter sufficiency test which brings with it a unique set of issues, e.g. the protein may not be amenable to purification in functional form, we opted for a genetic approach. Knowing that *T. brucei* expresses only the Gpi13 ortholog of this 3-member family, we chose to knockout TbGPI13 and ask if we could recover viable cells. Our prediction was that if Gpi13 played an essential role in *N*-glycosylation, i.e. it is the sole M5-DLO scramblase in *T. brucei*, then we would not be able to generate GPI13Δ trypanosomes (as noted above, the GPI anchoring function of Gpi13 is not essential in *T. brucei* cells grown in culture)[28-31]. To our disappointment, we were able to generate viable GPI13Δ cells that had the expected deficiency in GPI anchoring (Fig. 5C, D), but no evident disruption of protein *N*-glycosylation (Fig. 5E). Thus, Gpi13 does not play an essential role in *N*-glycosylation.

It remains to discuss why we failed to identify the M5-DLO scramblase. There are several possibilities that we list here that may form the basis for future investigation.

(i) We detected the vast majority (255) out of 286 known yeast ER membrane proteins (M. Schuldiner, unpublished data) (the 31 proteins that we did not detect are listed in Table S3), but the M5-DLO scramblase may have eluded detection. This could be because it might have only a few lysine and/or arginine residues, and that the resulting long tryptic peptides are not optimal for detection in the mass spectrometer.

(ii) We filtered the mass spectrometric data by imposing the criterion that M5-DLO scramblase candidates must have $\geq$ 3 TM spans. Known phospholipid scramblases of the GPCR and TMEM16 protein families have at least 7 TM spans, consistent with their function as transporters. Indeed, if the M5-DLO scramblase would have fewer than 3 TM spans we would predict that it must homo- or hetero-oligomerize in order to form a functional transport entity.

(iii) We hypothesized that M5-DLO scramblase activity is essential, and extrapolated this to imply that the activity is due to a single protein and that this protein would therefore also be essential. What if the system is redundant, with several proteins individually possessing M5-DLO scramblase activity? In this case, the individual proteins would not necessarily be essential. Accordingly, we compiled a list of non-essential proteins whose profiles correlate well with the activity profile (Table S4). This list includes 18 ER-localized non-essential proteins with $\geq$ 3 predicted TM spans, plus 11 other proteins (also with $\geq$ 3 TM spans) whose subcellular location is not known. It would be interesting in the future to explore the possibility that these proteins may contribute to M5-DLO scramblase activity.

(iv) Perhaps the most significant reason for the failure of the activity profiling approach is that the M5-DLO scramblase may exist in two or more forms, a complication that could account for the difficulty (noted in the Introduction) in purifying it. There are several scenarios that we can consider. For example, Protein X is the scramblase, but when complexed with Protein Y it is inactive. The protein profile of Protein X would thus track the rising portion of the activity profile (corresponding to Protein X alone) and then continue more broadly (corresponding to a complex of Protein X and Protein Y) before returning to baseline. The resulting broader profile of Protein X would result in it being discarded in our data curation as it would correlate poorly with the activity profile. Examples of such profiles (Fig. S6) reveal potentially interesting proteins such as the GlcNAc phosphotransferase Alg7 (R=0.65 $\pm$ 0)(mean $\pm$ SD) and the mannosyltransferase Alg2

(R= 0.66 ± 0.05) (mean ± SD). Both proteins recognize dolichyl diphosphate and could potentially play a role in scrambling M5-DLO. The profiles of both proteins track the rising portion of the activity profile but then broaden out even as the activity decreases. Additional scenarios are possible, for example that scramblase activity is due to a complex of two proteins that are inactive as monomers – in this situation the protein profiles would diverge from the rising portion of the activity profile, and only track with the activity as it decreases in higher fractions.

In conclusion, we describe an innovative strategy to identify the M5-DLO scramblase. Although we did not identify the scramblase, we report an extensive data set that provides an opportunity to extend these studies in the future. Importantly, we identify potential pitfalls in our approach that will be useful in future work. Thus, our results eliminate the candidates studied here and raise the important possibility that the scramblase is either a heteromeric complex, or that its activity is regulated by complex formation as outlined above. This result will need to be factored in to future approaches to address this problem.

Materials and methods

Materials
Analytical-grade reagents were purchased from Sigma-Aldrich unless stated otherwise. Antibiotics were from Sigma-Aldrich, Invivogen or Invitrogen. Other reagents/materials were obtained as follows: D-2-[^{3}H]-mannose (20-30 Ci/mmol) (American Radiolabeled Chemicals Inc.), Micro BCA protein assay kit (Thermo Fisher Scientific), Triton X-100 (Roche), IgG-Agarose resin (GE Healthcare), anti-TAP antibody (Genescript), anti-Dpm1 antibody (Abcam), SM-2 Biobeads and P6-Biogel resin (Biorad) and egg phosphatidylcholine (Avanti Polar Lipids). Yeast strains used in this study are listed in Table S5.

Buffers
Buffer A: 10 mM HEPES-NaOH pH 7.4, 100 mM NaCl, 1% (w/v) Triton X-100; Buffer B: 10 mM HEPES-NaOH pH 7.4, 100 mM NaCl; Buffer C: Buffer B supplemented with 3 mM $MgCl_2$, 1 mM $CaCl_2$, 1 mM $MnCl_2$.

Preparation of [^{3}H]M5-DLO
Protocol 1. Procyclic form *T. brucei* lacking TbRft1 (TbRft1$^{-/-}$) were generated from *T. brucei* strain Lister 427 and maintained in culture as previously reported [17]. 10^9 cells were washed with glucose-free SDM-80, resuspended at a density of 1×10^8 cells/ml in the same medium, and incubated for 90 min at 27°C. D-[2-^{3}H]-mannose (200 μCi) was added and the cells were incubated for a further 2 hr at 27°C. The cells were then collected by centrifugation and lipids were sequentially extracted from the washed cell pellet with 3 x 5 ml chloroform/methanol (3:2, v/v), 3 x 5 ml chloroform/methanol/water (3:48:47, v/v/v), containing 1 M $MgCl_2$, and 1 x 5 ml chloroform/methanol/water (3:48:47, v/v/v) without $MgCl_2$. Finally, [^{3}H]M5-DLO was extracted with 2 x 5 ml chloroform/methanol/water (10:10:3, v/v/v). Extracts were pooled, dried under a stream of nitrogen, and partitioned between water and n-butanol. [^{3}H]M5-DLO was recovered in the n-butanol phase.

Protocol 2. Yeast cells (*alg3Δ*) were metabolically labeled with D-[2-^{3}H]mannose in low-glucose medium as previously described [6], and subjected to differential solvent extraction as for Protocol 1 to purify [^{3}H]M5-DLO.

Radiochemical purity of [^{3}H]M5-DLO preparations. The radiochemical purity of [^{3}H]M5-DLO preparations was evaluated by thin layer chromatography (TLC) of the intact lipid, and by HPLC analysis of the oligosaccharide portion. TLC was done on glass-backed Silica Gel 60 plates (Merck) with chloroform/methanol/water (10/10/3, v/v/v) as the solvent system. The chromatograms were visualized using a Raytest Rita* radioactivity TLC analyser (Berthold Technologies). HPLC analysis was carried out by Bobby Ng and Hudson Freeze (Sanford-Burnham-Prebys Medical Discovery Institute, La Jolla, CA) as described [35]. Briefly, the [^{3}H]M5-DLO preparation was dried and treated with a 1:1 (v/v) mixture of isopropanol and 0.2 N HCL for 30 min at 100°C. The released oligosaccharide was analyzed on a Dionex Ultimate 2000 HPLC system using an Agilent Microsorb NH2-column. Radioactivity was detected using an in-line Lablogic Radiodetector and the analysis was calibrated using a co-injected 2-aminobenzamide-labeled dextran ladder (ProZyme/Agilent) detected on a Dionex fluorescence detector.

Triton extract (TE) enriched in yeast ER membrane proteins

TE was prepared from yeast cells as described previously [36], except that an additional step was included to salt-wash the membranes prior to detergent solubilization. Briefly, 600 OD$_{600}$ units of cells (BY4741, or BY4741-derived strain expressing a TAP tagged version of the protein of interest) were harvested, washed and homogenized using glass beads. After low-speed centrifugation to clear unbroken cells, the homogenates were centrifuged at 200,000 x g_{av} for 30 min to pellet membranes. The membranes were salt-washed by resuspension in buffer containing 1 M sodium acetate, followed by incubation on ice for 30 min. After re-pelleting, the salt-washed membranes were resuspended in ice-cold Buffer B before being solubilized by gradually adding an equal volume of 2% (w/v) ice-cold Triton X-100 at a final concentration of 1 % (w/v). The sample was incubated on ice for 30 min before removing insoluble material by centrifugation at 200,000 x g_{av} for 1 hr to generate a clear supernatant (TE). We previously used SDS-PAGE immunoblotting to show that TE is enriched in ER membrane proteins such as Dpm1 and Sec61, and depleted of plasma membrane proteins (Pma1, Gas1), as well as proteins of the vacuole (Vph1) and Golgi/endosomes (Pep12) [6].

Fractionation of TE by velocity sedimentation on a glycerol gradient

1.5 mL TE was loaded onto a 10 mL continuous glycerol gradient (5% to 25% (w/v), prepared in Buffer A) and centrifuged for 30 h in an SW41 Ti rotor at 100,000 x g_{av}. A mixture of sedimentation standards (anhydrase, ovalbumin, bovine serum albumin, β-amylase) was loaded onto a parallel gradient. Fractions were collected manually or using an Auto Densi-Flow apparatus (Labconco). The refractive index of each fraction was measured before removing glycerol by buffer exchange on a desalting spin column packed with Biogel-P6 resin equilibrated in Buffer A. Migration of the sedimentation markers was determined by SDS-PAGE/Coomassie-staining of the fractions.

Immunodepletion of a candidate protein from TE

TE from yeast strains expressing a TAP-tagged protein of interest (Table S5) was divided into two aliquots. One aliquot was treated at 4°C for 2 h with IgG-agarose beads to remove specifically the

TAP-tagged protein, whereas the other was mock-incubated. Removal of the protein from the extract was verified by SDS-PAGE immunoblotting, using an anti-TAP antibody (1:1,000 dilution) and anti-Dpm1 (1:1,000). The Dpm1 immunoblot served as a specificity control as well as a loading control.

Quantitative mass spectrometry
Six samples (fractions 5 to 10 of the gradient) were analyzed by mass spectrometry as follows. Proteins in each sample were digested by trypsin after reduction and alkylation with dithiothreitol and iodoacetamide. The resulting peptides were labeled with 6-plex TMT reagents. The digests were combined and subjected to offline fractionation by high pH reversed phase chromatography to obtain 6 fractions. Each fraction was analyzed by LC-MS. Online chromatography was performed with a Thermo Easy nLC 1000 ultra-high-pressure HPLC system (Thermo Fisher Scientific) coupled online to an Orbitrap Fusion Lumos mass spectrometer with a NanoFlex source (Thermo Fisher Scientific). Analytical columns (~ 25 cm long and 75 µm inner diameter) were packed in-house with ReproSil-Pur C18 AQ 3 µM reversed phase resin (Dr. Maisch GmbH, Ammerbuch-Entringen, Germany). A peptide mixture was loaded onto the analytical column with buffer A (0.1% formic acid) and separated with a linear gradient of 3% to 32% buffer B (100% ACN and 0.1% formic acid) at a flow rate of 300 nL/min over 240 min. MS data were acquired using a data-dependent method, dynamically choosing the most abundant not-yet-sequenced precursor ions from the survey scans. Peptide fragmentation was performed via higher energy collisional dissociation. The MS data were processed by the MaxQuant software for protein identification and quantitation. The false discovery rate for protein identification was 1%. A Pearson correlation score (R) was obtained for each identified protein by comparing the measured profile profile and the activity profile.

Candidate selection
Proteins with R >0.9 were selected. Using Yeastmine [37] this list was intersected with the list of all membrane proteins [38], as well as all essential proteins in *Saccharomyces cerevisiae* (list generated using Yeastmine). This final list was manually completed with the number of transmembrane domains and cellular localization for each protein using information available in the Saccharomyces Genome Database.

Liposomes and proteoliposome preparation
Vesicles were reconstituted essentially as described [6,14]. Briefly, for each reconstitution sample, [^{3}H]M5-DLO (20,000 cpm) was mixed with 4.5 µmol egg PC, dried under a stream of N_2 and resuspended in 540 µl Buffer B. The lipids were sonicated before being supplemented with Triton X-100 (from a 10 % stock solution in water, final concentration 1 % (w/v)). For proteoliposomes, TE or a fraction from the velocity gradient was added; for control (protein-free) liposomes, an equivalent amount of Buffer A was added instead. The final volume of the sample was 1 ml. Vesicle formation was induced by removing detergent with SM2 biobeads (previously washed with methanol and water) added in two stages: 100 mg of the beads were added and the sample was incubated with end-over-end mixing at room temperature; following this incubation, a further 200 mg of the beads were added and vesicles were incubated with end-over-end mixing overnight at 4°C. The resulting vesicles were pelleted by centrifugation (200,000 x g_{av}, 1 hr, 4°C)

and resuspended in Buffer C. Vesicle protein was estimated using the Kaplan Pederson method [39] and phospholipid was measured as described previously [40].

M5-DLO scramblase assay
The assay was performed as previously described [6,7,14]. Briefly, 20 µL of vesicles (proteoliposomes or liposomes) were incubated in 40 µl of Buffer C ("buffer"), or Buffer C containing 3.6 mg/ml Concanavalin A ("ConA") or Buffer E containing 3.6 mg/ml Concanavalin A and Triton X-100 1.5% (w/v) ("ConA-TX") for 30 min on ice. Then 140 µl of blocking solution (BSA 20 mg/ml, Yeast Mannan 2 mg/ml) was added, followed immediately by 1.4 ml of chloroform/methanol (1:1, v/v). The sample was vortexed and centrifuged (20,000 x g_{av}, 5 min, 4°C). The supernatant was placed in a 20-ml scintillation vial for drying and the pellet was solubilized in 1 % (w/v) SDS overnight under vigorous shacking. Radioactivity in the pellet (P) and the supernatant (S) was measured by liquid scintillation counting. For each sample the percentage of M5-DLO capture was calculated as follows:

$100 \bullet ((P+S)_{ConA} - (P+S)_{buffer}) / ((P+S)_{ConA-TX} - (P+S)_{buffer})$.

Generation of TbGPI13 and TbGPI14 knock out strains in *T. brucei* procyclic forms
Gene disruptions were accomplished in *T. brucei* Cas9-expressing SmOx p9 procyclic forms by replacing both alleles of the target gene with antibiotic resistance genes using the CRISPR/Cas9 technique as described previously [41]. Geneticin and hygromycin replacement cassettes, flanked by 30 bp sequences homologous to the target gene UTRs, were PCR-amplified from pPOTv7 plasmids using gene-specific forward and reverse primers (Table S6). Short guide RNA (sgRNA) templates, consisting of a T7 RNA polymerase promoter sequence and a gene-specific sgRNA sequence to guide double-strand breaks to the UTRs of the target gene, were amplified by annealing gene-specific primers with the G00 sgRNA scaffold primer. All PCR products were purified using Promega Wizard SV PCR clean-up system before transfection. Gene-specific geneticin and hygromycin targeting fragments and 5' and 3' sgRNA templates were combined with 1 x TbBSF buffer (90 mM sodium phosphate, 5 mM potassium chloride, 0.15 mM calcium chloride, 50 mM HEPES, pH 7.3) in a total volume of 100 µl and transfected into 3 x 10^7 SmOx p9 *T. brucei* cells by electroporation using the Amaxa Nucleofector 4D (Lonza), program FI-115. Clonal populations resistant to both hygromycin and geneticin were obtained by limiting dilution, and complete loss of the target gene was verified by diagnostic PCR reactions.

SDS-PAGE and immunoblotting
T. brucei cells were harvested by centrifugation, washed twice with PBS and the pellet suspended directly in SDS sample buffer and heated at 55°C for 10 mins. Approximately 5x10^6 cell equivalents per sample were separated by SDS-PAGE under reducing conditions on a 12% polyacrylamide gel and transferred to nitrocellulose membrane. For all blots, blocking was carried out in 5% milk in TBS, and primary and secondary antibodies diluted in 1% milk in TBS + 0.1% Tween-20 (TBST), with the addition of 0.01% SDS for the secondary antibodies. The following primary antibodies were used: anti-p67 (J. Bangs, 1:5000), anti-ALBA1 (I. Roditi, 1:2500), anti-EP247 (Cedarlane, 1:1500), anti-GPEET K1 (I. Roditi, 1:2500), anti-GPEET 5H3 (I. Roditi, 1:5000) and anti-aldolase (P. Michels, 1:180,000). HRP-conjugated secondary anti-rabbit and anti-mouse antibodies (DAKO) were used at 1:1000 and 1:5000 respectively, and

chemiluminescent detection carried out using SuperSignal™ Pico West PLUS substrate (ThermoScientific).

Flow cytometric analysis

Trypanosomes (~$5x10^6$ cells) were centrifuged and the pellet was directly incubated with one of the following primary antibodies, anti-EP247, GPEET K1 or anti-GPEET 5H3, in SDM-79 media for 30 min. After two washes with SDM-79 samples were incubated with anti-rabbit or anti-mouse AlexaFluor 488 (Invitrogen) secondary antibody (both 1:1000) in SDM-79 for 30 min. Samples were washed two more times and analyzed with a Novo-Cyte flow cytometer (ACEA Biosciences). Samples and buffers were kept on ice throughout, and all centrifugations and antibody incubations were performed at 4°C.

Metabolic labeling and extraction of [^{3}H]-ethanolamine-labeled GPI precursors and GPI-anchored proteins

Procedures were as previously described [17]. Briefly, $5x10^8$ cells were cultured for 16 hours in the presence of 40 μCi of [^{3}H]ethanolamine. The cells were harvested by centrifugation and washed twice with ice-cold TBS (10 mM Tris-HCl, 144 mM NaCl, pH 7.4). Bulk phospholipids were extracted from the pellet with 2x 10 ml of chloroform/methanol 2:1 (v/v). GPI precursors and free GPIs were extracted with 3x 10 ml of chloroform/methanol/water 10:10:3 (v/v/v), and GPI-anchored proteins were extracted with 2x 10 ml of 9% butanol in water (v/v). GPI precursors and free GPIs were further separated by partitioning the extract between butanol and water. GPI precursors were analyzed by TLC while free GPIs and GPI-anchored proteins were visualized by SDS-PAGE followed by fluorography.

Acknowledgements

This work was supported by Swiss National Science Foundation Sinergia grant CRSII5_170923. We thank Jennifer Jelk for technical assistance, Guoan Zhang and his colleagues at the Proteomics and Metabolomics Core Facility (Weill Cornell Medicine) for mass spectrometry, Maya Schuldiner (Weizmann Institute) for sharing a list of yeast ER membrane proteins, Bobby Ng and Hudson Freeze (Sanford-Burnham-Prebys Medical Discovery Institute, La Jolla, CA) for oligosaccharide analysis of [^{3}H]M5-DLO preparations, Brenda Andrews (University of Toronto) and Markus Aebi (ETH Zürich) for yeast strains, and Jay Bangs (SUNY Buffalo), Isabel Roditi (Universität Bern) and Paul Michels (Research Unit for Tropical Diseases, International Institute of Cellular and Molecular Pathology) for antibodies.

Tables S1 and S2: Technical replicates of TMT mass spectrometric analysis of velocity gradient fractions (see Fig. 3A and 'Materials and methods' for details). The Tables are provided as .csv (comma separated value) files. A description of the column headings used in the Tables is given below.

Column Name	Description
Protein IDs	Identifier(s) of protein(s) contained in the protein group.
Protein names	Name(s) of protein(s) contained within the group.
Gene names	Name(s) of the gene(s) associated to the protein(s) contained within the group.
FASTA headers	FASTA headers(s) of protein(s) contained within the group. FASTA file is a text-based format containing protein sequences, in which amino acids are represented using single-letter codes.
Peptides	The total number of peptide sequences associated with the protein group (i.e. for all the proteins in the group).
Sequence coverage [%]	Percentage of the sequence that is covered by the identified peptides of the best protein sequence contained in the group.
Q-value	This is an indicator of the confidence for protein identification, with range of 0-0.01 (1%). A smaller Q-value represents higher confidence of protein identification.
Columns H-M	Summed up intensity of all the identified peptide sequences
cor_score	The correlation score is the Pearson correlation between the measured profile and the standard profile (see Fig. S4)

Table S3: Predicted ER membrane proteins that were not identified in the present study

Gene	Protein	Function	TM spans
YLL055W	YCT1	Yeast Cysteine Transporter	10
YLR004C	THI73	THI regulon	7
YLR023C	IZH3	Implicated in Zinc Homeostasis	7
YMR274C	RCE1	Ras and a-factor Converting Enzyme	7
YHR140W			6
YER140W	EMP65	ER Membrane Protein of 65 kDa	5
YKR053C	YSR3	Yeast Sphingolipid Resistance	5
YFR042W	KEG1	Kre6-binding ER protein responsible for Glucan synthesis	4
YLR246W	ERF2	Effect on Ras Function	4
YNL008C	ASI3	Amino acid Sensor-Independent	4
YNL194C			4
YLL014W	EMC6	ER Membrane protein Complex	3
YIL089W			2
YFR041C	ERJ5	Endoplasmic Reticulum located J-protein	2
YDR437W	GPI19	Glycosyl PhosphatidylInositol anchor biosynthesis	2
YGR105W	VMA21	Vacuolar Membrane Atpase	2
YLL052C	AQY2	AQuaporin from Yeast	2
YNL046W			2
YNL146W			2
YOR044W	IRC23	Increased Recombination Centers	2
YPL096C-A	ERI1	ER-associated Ras Inhibitor	2
YAL028W	FRT2	Functionally Related to TCP1	1
YBL100C			1
YBR162W-A	YSY6		1
YDL232W	OST4	OligoSaccharylTransferase	1
YLR238W	FAR10	Factor ARrest	1
YNL087W	TCB2	Three Calcium and lipid Binding domains (TriCalBins)	1
YOR324C	FRT1	Functionally Related to TCP1	1
YPL200W	CSM4	Chromosome Segregation in Meiosis	1
YER053C-A			1
YMR122W-A	NCW1	Novel Cell Wall protein	1

Table S4: Non-essential membrane proteins with R>0.9 and ≥3 TM spans

ORF	Protein		MW (Da)	TM spans
		ER localized		
YDR205W	MSC2	Meiotic Sister-Chromatid recombination	80578.2	13
YBR132C	AGP2	high-Affinity Glutamine Permease	67261.4	12
YJL212C	OPT1	OligoPeptide Transporter	91630.2	12
YOR067C	ALG8	Asparagine-Linked Glycosylation	67409.4	12
YGR227W	DIE2	Derepression of ITR1 Expression	61811	11
YEL031W	SPF1	Sensitivity to Pichia Farinosa killer toxin	135259.6	10
YOR002W	ALG6	Asparagine-Linked Glycosylation	62798.3	10
YFL025C	BST1	Bypass of Sec Thirteen	117747	8
YGL160W	AIM14	Altered Inheritance rate of Mitochondria	65849.6	7
YGL167C	PMR1	Plasma Membrane ATPase Related	104554.8	7
YLR130C	ZRT2	Zinc-Regulated Transporter	46342.2	7
YBR287W			47493.6	6
YJR015W			58116.5	6
YBL011W	SCT1	Suppressor of Choline-Transport mutants	85693.2	4
YKR067W	GPT2	Glycerol-3-Phosphate acylTransferase	83674.9	4
YOR085W	OST3	OligoSaccharylTransferase	39487.6	4
YER072W	VTC1	Vacuolar Transporter Chaperone	14381.3	3
YGR089W	NNF2		106476.7	3
		Unknown subcellular location		
YDL206W			85973.1	12
YOL103W	ITR2	myo-Inositol TRansporter	66715.8	12
YLR241W	CSC1	Calcium permeable Stress-gated cation Channel	89348.1	11
YOL158C	ENB1	ENteroBactin	66989.4	11
YER060W	FCY21	FluoroCYtosine resistance	58041.4	10
YHR078W			63331.2	8
YGR199W	PMT6	Protein O-MannosylTransferase	88018.7	8
YGR149W	GPC1	GlyceroPhosphoCholine acyltransferase	51652.9	7
YLR214W	FRE1	Ferric REductase	78879.7	7
YLR443W	ECM7	ExtraCellular Mutant	50314.2	4
YOR301W	RAX1	Revert to Axial	50160.1	3

Table S5: Yeast strains used in this study

Strain	Genotype	Source
GPI13-TAP	*MATα his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 GPI13-TAP::HIS3MX6*	Brenda Andrews
GPI7-TAP	*MATα his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 GPI7-TAP::HISMX6*	Dharmacon[1]
MCD4-TAP	*MATα his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 MCD4-TAP::HISMX6*	Dharmacon[1]
GPI2-TAP	*MATα his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 GPI12-TAP::HISMX6*	Dharmacon[1]
STT3-TAP	*MATα his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 STT3-TAP::HISMX6*	Dharmacon[1]
GPI14-TAP	*MATα his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 GPI14-TAP::HISMX6*	Brenda Andrews
BY4741	*MATα his3Δ1 leu2Δ0 met15Δ0 ura3Δ0*	Brenda Andrews
YG248	*MATα ade2-101 ura3-52 his3Δ200 lys2-801 Δalg3::HIS3*	Markus Aebi

[1]Dharmacon TAP-Fusion ORF collection (strains were confirmed by colony PCR)

Table S6: Primers used to create knockout *T. brucei* cell lines

Primer	Sequence
TbGPI13 -pPOT- Fwd	<u>AGAAGTGTGAGCACTAGACGCTGACTACCC</u>GTATAATGCAGACCTGCTGC
TbGPI13 -pPOT- Rev	<u>ACCATCAGCATCAACAACAACACACAAAGA</u>CCGGAACCACTACCAGAACC
TbGPI13 -sgRNA- 5'	GAAATTAATACGACTCACTATAGG<u>CTTTCAGCAAAGTTACACAC</u>GTTTTAGAGCTAGAAATAGC
TbGPI13 -sgRNA- 3'	GAAATTAATACGACTCACTATAGG<u>AAAAAAGGATGACGAGAACA</u>GTTTTAGAGCTAGAAATAGC
TbGPI14 -pPOT- Fwd	<u>GATAGAGGCTTATCAGCTAGAAAATTTCCG</u>GTATAATGCAGACCTGCTGC
TbGPI14 -pPOT- Rev	<u>ACCGCGAAGCAAACACCCACTTGCAAGCCA</u>CCGGAACCACTACCAGAACC
TbGPI14 -sgRNA- 5'	GAAATTAATACGACTCACTATAGG<u>AGACGTGGTGGAAAACTACA</u>GTTTTAGAGCTAGAAATAGC
TbGPI14 -sgRNA- 3'	GAAATTAATACGACTCACTATAGG<u>TGTTGTCGATGTCTTACAGT</u>GTTTTAGAGCTAGAAATAGC
G00	AAAAGCACCGACTCGGTGCCACTTTTTCAAGTTGATAACGGACTAGCCTTATTTTAACTTGCTA TTTCTAGCTCTAAAAC

<u>Gene-specific sequences are underlined</u>

Figures

Figure 1: Transbilayer movement of Man₅GlcNAc₂-PP-dolichol (M5-DLO).

A. Synthesis of $Glc_3Man_9GlcNAc_2$-PP-dolichol (G3M9-DLO), the oligosaccharide donor for protein *N*-glycosylation in yeast and humans, occurs in two stages. The first stage, comprising seven glycosyltransfer reactions, occurs on the cytoplasmic side of the ER to generate M5-DLO from dolichyl phosphate, UDP-GlcNAc and GDP-mannose. The remaining seven glycosyltransfer reactions occur in the ER lumen, adding four mannose residues (from dolichylphosphoryl mannose (dol-P-man)) and three glucose residues (from dolichylphosphoryl glucose (dol-P-Glc)) to M5-DLO to generate G3M9-DLO. The switch from the cytoplasmic to the lumenal side of the ER requires transbilayer movement of M5-DLO, which is facilitated by M5-DLO scramblase (red icon). The right side of the figure shows oligosaccharyl-transferase (OST)-mediated transfer of the $Glc_3Man_9GlcNAc_2$ oligosaccharide from G3M9-DLO to an asparagine residue within a glycosylation sequon in a nascent protein as it emerges from the protein translocon into the ER lumen. See text for details.

B. Recapitulation of M5-DLO scrambling in a reconstituted system. The assay exploits the organic-solvent-resistant interaction between M5-DLO and the lectin Concanavalin A (Con A) - thus whereas M5-DLO is soluble in organic solvent, M5-DLO:ConA complexes are precipitated. Proteoliposomes are reconstituted from egg phosphatidylcholine, [³H]M5-DLO and ER membrane proteins. Protein-free liposomes are prepared in parallel. Incubation of the liposomes with Con A results in the capture/precipitation of ~50% of [³H]M5-DLO (left), corresponding to the pool in the outer leaflet of the vesicles, whereas similar incubation of proteoliposomes containing M5-DLO scramblase results in the capture of ~100% of [³H]M5-DLO (right) as M5-DLO molecules in the inner leaflet are transported to the outer leaflet where they are

captured by Con A. Proteoliposomes reconstituted with proteins other than the M5-DLO scramblase, i.e. irrelevant proteins, are expected to behave similarly to protein-free liposomes (left).

Verchère et al.
Figure 2

Figure 2. Principle of the activity correlation profiling approach.
A. A Triton X-100 extract (TE) of salt-washed yeast membranes, i.e. an extract that is enriched in ER membrane proteins, is fractionated by velocity gradient sedimentation. Half of each fraction is reconstituted into large unilamellar vesicles and assayed for M5-DLO scramblase activity as outlined in Figure 1B to generate a profile of scramblase activity across the gradient. The remaining half of each fraction is subjected to mass spectrometric analysis using the tandem mass tag (TMT) system. Thus, each fraction is digested with trypsin, and the resulting peptides are labeled with TMT multiplex reagents (a unique TMT mass tag is used for each fraction). The digests are pooled and analyzed by tandem mass spectrometry to quantify the relative amount of specific proteins across fractions, thereby generating protein profiles. The protein profiles are quantitatively compared with the activity profile by obtaining a Pearson correlation score.
B. Schematic illustration of the readout from the activity correlation profiling experiment. The activity profile (solid blue line) can be compared to the profile of several different proteins (dashed lines) identified by tandem mass spectrometry. Only a protein whose profile is highly correlated with the activity profile is considered as a scramblase candidate. Other proteins that fractionate differently (brown and green dashed lines) are considered irrelevant to scramblase activity.

Figure 3: Identification of M5-DLO scramblase candidates by activity correlation profiling.

A. Yeast TE was fractionated by velocity sedimentation on a 5-25% glycerol gradient and fractions were collected from the top. Sedimentation standards (a-d, corresponding to carbonic anhydrase (2.8S), ovalbumin (3.6S), bovine serum albumin (4.2S) and β-amylase (8.9S)) were resolved in a parallel gradient and their migration positions are indicated. The refractive index of the fractions is indicated by the green line. The profiles of bulk protein (carmine) and M5-DLO scramblase activity (blue, duplicate measurements) are indicated for fractions 5-10.

B. Venn diagram describing the selection of M5-DLO scramblase candidates. The mass spectrometry analyses identified 280 proteins with a correlation coefficient R>0.9 (filled circle). Intersections between this group and the groups of all essential proteins in yeast (*Sc* essential proteins), and yeast membrane proteins with three or more predicted transmembrane spans (*Sc* membrane proteins (TM≥3, assessed by the TMHMM membrane protein topology prediction program)) yielded 5 proteins. One of these (Neo1) was discarded because it is not an ER resident.

C. List of M5-DLO scramblase candidates. The four ER membrane proteins obtained at the intersection of the Venn diagram are listed. The non-essential protein Gpi7 was added because its profile is highly correlated with that of M5-DLO scramblase activity (see panel D) and because it belongs to the family of GPI phosphoethanolamine transferases that includes the identified candidates Gpi13 and Mcd4 (see panels F and G). Stt3 was added to the list despite R<0.9 because of its relevance to *N*-glycosylation as the catalytic subunit of OST (Figure 1A).

D. Profile of candidates: profiles of Gpi13, Gpi7 and Mcd4 versus the activity profile (shaded and outlined in blue).

E. Profile of candidates: profiles of Gpi14, Gpi2 and Stt3 versus the activity profile (shaded and outlined in blue).

F. Role of Gpi13, Gpi7 and Mcd4 in P-EtN transfer reactions of GPI biosynthesis. The structure of the mature GPI anchor precursor in yeast and mammals is shown, with phospho-ethanolamine residues (P-EtN) decorating each of the mannose residues. Mcd4, Gpi7 and Gpi13 are the P-EtN transferases responsible for adding the first, second and third P-EtN residues, respectively. *T. brucei* lacks homologs of Mcd4 and Gpi7, and consequently has only the capping P-EtN residue transferred by Gpi13.

G. Membrane topology of Gpi13, Gpi7 and Mcd4 assessed using Protter (Protter: interactive protein feature visualization and integration with experimental proteomic data. Omasits U, Ahrens CH, Müller S, Wollscheid B. Bioinformatics. 2014 Mar 15;30(6):884-6. doi: 10.1093/bioinformatics/btt607.) The salmon-coloured strip is the ER membrane bilayer. The stretch of Gpi7 coloured red is a predicted signal sequence. The lumenal and cytoplasmic sides of the membrane are indicated. The number of predicted transmembrane spans differs slightly from the number predicted by the TMHMM server that was used to calculate the Venn diagram group '*Sc* membrane proteins (TM≥3)'.

Verchère et al.
Figure 4

Figure 4: Evaluating M5-DLO scramblase candidates by immunodepletion.

TE was prepared from yeast strains expressing a C-terminally TAP-tagged variant of the scramblase candidate in place of the endogenous protein. Tagging was done at the genomic locus. Six unique TE preparations were thus generated, corresponding to cells expressing Gpi13-TAP, Gpi7-TAP, Mcd4-TAP, Gpi14-TAP, Gpi2-TAP or Stt3-TAP. In every case the extract was mock-treated (green filled circles) or incubated with IgG-agarose beads to remove the TAP-tagged protein (blue open circles). Removal of the TAP-tagged protein from the extract was confirmed by SDS-PAGE/immunoblotting using anti-TAP antibodies, and specificity of immunodepletion and equivalence of sample loading was confirmed by immunoblotting to detect the ER resident protein Dpm1. Mock-treated and depleted extracts were reconstituted and assayed for M5-DLO scramblase activity. Reconstitutions were performed at different ratios of protein to phospholipid, in the approximate range 25 - 125 g/mol. The figure shows immunoblots

paired with the corresponding activity assays for each candidate. Error bars are obtained from duplicate measurements of activity, as well as duplicate measurements of the protein and phospholipid content of the reconstituted vesicles. The red dashed line, based on all data from the mock-treated samples, is intended to guide the eye.

Verchère et al.
Figure 5

Figure 5: Evaluating Gpi13 as the M5-DLO scramblase in procyclic form *Trypanosoma brucei*.
A. PCR confirmation of *TbGPI13* gene replacement using primers specific to the *TbGPI13* UTR and ORF, and HygR/G418R gene replacement cassettes. Lanes 1, 3, 5 and 7 correspond to wild-type cells, whereas lanes 2, 4, 6 and 8 correspond to TbGPI13 knockout cells.
B. Growth of TbGPI13 knockout cells (GPI13Δ) compared with wild-type (WT) cells. Data points represent mean values from 2 independent experiments.
C. Thin layer chromatographic analysis of polar lipid extracts from [³H]-ethanolamine-labeled WT (top trace) and GPI13Δ (bottom trace) cells showing loss of GPI synthesis precursor PP1 in GPI13Δ cells. The chromatograms were visualized using a radioactivity scanner. The WT trace is displaced upwards for clarity. PP1 and PE refer to the major GPI precursor in procyclic trypanosomes (structure shown in

schematic, corresponding to EtN-P-Man$_3$GlcN-(2-acyl-Inositol)-P-monoacylglycerol) and PE, respectively. The majority of PE is removed by extraction with a lower polarity solvent, before extracting PP1 in a more polar solvent.

D. SDS-PAGE/fluorography of protein extracts from [^{3}H]-ethanolamine-labeled WT and GPI13Δ cells showing loss of GPI-anchored GPEET procyclin (indicated by arrow head) in GPI13Δ cells. The band at ~50 kDa (indicated by filled circle) represents eukaryotic elongation factor 1A, a protein that is modified by ethanolamine phosphoglycerol and therefore metabolically labeled with [^{3}H]-ethanolamine (Signorell, A., Jelk, J., Rauch, M., Bütikofer, P. (2008) Phosphatidylethanolamine is the precursor of the ethanolamine phosphoglycerol moiety bound to eukaryotic elongation factor 1A. J. Biol. Chem. 283: 20320-20329)

E. Analysis of p67 *N*-glycosylation by anti-p67 immunoblot of whole cell protein samples from WT and GPI13Δ cells. A sample from a known *N*-glycosylation defective cell line (ALG3Δ) was used as a positive control. Alba1 was probed using anti-Alba1 antibody and used as a loading control.

F. Flow cytometry analysis of surface expression of GPI-anchored EP and GPEET procyclins in WT and GPI13Δ cells. The middle and right panels correspond to flow cytometry analysis of GPEET using 5H3 antibody (middle) and K1 antiserum (right). Control (Con) samples were generated by omitting the primary antibody.

G. Anti-EP and anti-GPEET (K1) procyclin immunoblots of whole cell samples from WT and GPI13Δ cells. Aldolase was probed using anti-aldolase antibody and used as a loading control.

H. WT and GPI13Δ cells were fixed and stained with antibodies against EP and BiP (as a marker for the endoplasmic reticulum) in combination with fluorophore-conjugated secondary antibodies and analyzed by fluorescence microscopy. DNA was stained with 4',6-diamidino-2-phenylindole (DAPI) in the merged panels. DIC, differential interference contrast. Scale bar = 5 μm.

Supplementary Figures

Figure S1. HPLC analysis of the oligosaccharide portion of [³H]M5-DLO. The pyrophosphate linkage in [³H]M5-DLO was hydrolyzed by mild acid treatment and the released [³H]oligosaccharide was analyzed by HPLC using an amino-column. Representative profiles (radioactivity (cpm), versus elution time) of samples of [³H]M5-DLO prepared via Protocol 1 (*Tb*) and Protocol 2 (*Sc*) are shown. The peak in each case co-elutes approximately with the glucose-9 oligomer in a dextran ladder that was co-injected with the samples. Data provided by Bobby Ng and Hudson Freeze.

Verchère et al.
Figure S2

Figure S2. Distribution of proteins in the velocity gradient. The mass spectrometry data were used to determine the peak fraction for each identified protein, i.e. the fraction in which the protein is maximally abundant. The molecular weights of the proteins were obtained from the YeastMine webserver. **A, B.** Box and violin plots showing the molecular weights of proteins peaking in each fraction for the two technical replicates of the mass spectrometric analysis. The median and quartiles of the data are indicated. **C.** Mean molecular weights of proteins maximally abundant in each fraction. The points show the mean (± standard deviation) of data obtained from two technical replicates. The line through the points is intended to guide the eye.

Figure S3. Profiles of OST and GPIT subunits in the velocity gradient. The distribution of subunits of the OST complex (panel A) and the GPI transamidase (GPIT) complex (panel B) in the velocity gradient were obtained from the mass spectrometry data. Yeast OST exists in two isoforms that contain either Ost3 or Ost6, in addition to seven other subunits: Stt3, Ost1, Wbp1, Swp1, Ost2, Ost4, Ost5. These complexes can be isolated from yeast membranes after solubilization in dodecylmaltoside detergent supplemented with cholesteryl hemisuccinate (Wild et al. *Science* **359**: 545-550 (2018)). In panel A, Triton X-100-solubilized OST complex remains largely intact, with the exception of the Ost5 subunit which sediments less rapidly than the rest of the subunits and appears to have dissociated from the complex.

Figure S4: Correlation between the activity profile and the profiles of Gpi13, Gpi7 and Mcd4.
A. Reproduced from Fig. 3D.
B. Quantification of correlation between the activity profile and profiles of Gpi13, Gpi7 and Mcd4 (taken from panel A). Values defining the profile of a specific protein, i.e. relative protein abundance in a particular fraction, are plotted against the corresponding values of activity and analyzed by simple linear regression. Linear regression yielded R squared values of >0.81, corresponding to R values of >0.90. The plot shows intuitively that activity values are high when relative protein abundance values are high, and correspondingly that activity values are low when relative protein abundance values are low.

A

 B

Figure S5: Gpi7, Gpi13 and Mcd4 are expressed at similar levels.

A. 6 OD of cells expressing chromosomally TAP-tagged Gpi7, Gpi13 or Mcd4 were precipitated with TCA, pellets were washed with acetone, resuspended in loading dye and analyzed by SDS-PAGE/immunoblotting using anti-TAP antibody. Immunoblotting of GAPDH served as a loading control.
B. The intensity of bands detected by anti-TAP and anti-GAPDH was quantified using Image Studio software. The ratio of intensities was taken as an indicator of abundance of the TAP-tagged protein. Data for three analyses are shown (mean ± standard deviation).

Figure S6: Profiles of Alg2 and Alg7.

Profiles of Alg2 and Alg7 illustrate the possibility that the activity correlation profile method may be complicated by complex formation (see Discussion for details).

REFERENCES

Lennarz, W. J. Protein glycosylation in the endoplasmic reticulum: current topological issues. *Biochemistry* **26**, 7205-7210, doi:10.1021/bi00397a001 (1987).

Hirschberg, C. B. & Snider, M. D. Topography of glycosylation in the rough endoplasmic reticulum and Golgi apparatus. *Annu Rev Biochem* **56**, 63-87, doi:10.1146/annurev.bi.56.070187.000431 (1987).

Schenk, B., Fernandez, F. & Waechter, C. J. The ins(ide) and out(side) of dolichyl phosphate biosynthesis and recycling in the endoplasmic reticulum. *Glycobiology* **11**, 61R-70R, doi:10.1093/glycob/11.5.61r (2001).

Larkin, A. & Imperiali, B. The expanding horizons of asparagine-linked glycosylation. *Biochemistry* **50**, 4411-4426, doi:10.1021/bi200346n (2011).

Breitling, J. & Aebi, M. N-linked protein glycosylation in the endoplasmic reticulum. *Cold Spring Harb Perspect Biol* **5**, a013359, doi:10.1101/cshperspect.a013359 (2013).

Sanyal, S., Frank, C. G. & Menon, A. K. Distinct flippases translocate glycerophospholipids and oligosaccharide diphosphate dolichols across the endoplasmic reticulum. *Biochemistry* **47**, 7937-7946, doi:10.1021/bi800723n (2008).

Sanyal, S. & Menon, A. K. Specific transbilayer translocation of dolichol-linked oligosaccharides by an endoplasmic reticulum flippase. *Proc Natl Acad Sci U S A* **106**, 767-772, doi:10.1073/pnas.0810225106 (2009).

Rush, J. S. & Waechter, C. J. Transmembrane movement of a water-soluble analogue of mannosylphosphoryldolichol is mediated by an endoplasmic reticulum protein. *J Cell Biol* **130**, 529-536, doi:10.1083/jcb.130.3.529 (1995).

Sanyal, S. & Menon, A. K. Stereoselective transbilayer translocation of mannosyl phosphoryl dolichol by an endoplasmic reticulum flippase. *Proc Natl Acad Sci U S A* **107**, 11289-11294, doi:10.1073/pnas.1002408107 (2010).

Sanyal, S. & Menon, A. K. Flipping lipids: why an' what's the reason for? *ACS Chem Biol* **4**, 895-909, doi:10.1021/cb900163d (2009).

Pomorski, T. G. & Menon, A. K. Lipid somersaults: Uncovering the mechanisms of protein-mediated lipid flipping. *Prog Lipid Res* **64**, 69-84, doi:10.1016/j.plipres.2016.08.003 (2016).

Ng, D. T., Spear, E. D. & Walter, P. The unfolded protein response regulates multiple aspects of secretory and membrane protein biogenesis and endoplasmic reticulum quality control. *J Cell Biol* **150**, 77-88, doi:10.1083/jcb.150.1.77 (2000).

Helenius, J. *et al.* Translocation of lipid-linked oligosaccharides across the ER membrane requires Rft1 protein. *Nature* **415**, 447-450, doi:10.1038/415447a (2002).

Frank, C. G., Sanyal, S., Rush, J. S., Waechter, C. J. & Menon, A. K. Does Rft1 flip an N-glycan lipid precursor? *Nature* **454**, E3-4; discussion E4-5, doi:10.1038/nature07165 (2008).

Rush, J. S., Gao, N., Lehrman, M. A., Matveev, S. & Waechter, C. J. Suppression of Rft1 expression does not impair the transbilayer movement of Man5GlcNAc2-P-P-dolichol in sealed microsomes from yeast. *J Biol Chem* **284**, 19835-19842, doi:10.1074/jbc.M109.000893 (2009).

Jelk, J. *et al.* Glycoprotein biosynthesis in a eukaryote lacking the membrane protein Rft1. *J Biol Chem* **288**, 20616-20623, doi:10.1074/jbc.M113.479642 (2013).

Gottier, P. *et al.* RFT1 Protein Affects Glycosylphosphatidylinositol (GPI) Anchor Glycosylation. *J Biol Chem* **292**, 1103-1111, doi:10.1074/jbc.M116.758367 (2017).

Snider, M. D. & Rogers, O. C. Transmembrane movement of oligosaccharide-lipids during glycoprotein synthesis. *Cell* **36**, 753-761, doi:10.1016/0092-8674(84)90355-6 (1984).

Sakurai, H., Kubota, K., Inaba, S., Takanaka, K. & Shinagawa, A. Identification of a metabolizing enzyme in human kidney by proteomic correlation profiling. *Mol Cell Proteomics* **12**, 2313-2323, doi:10.1074/mcp.M112.023853 (2013).

McAllister, F. E. & Gygi, S. P. Correlation profiling for determining kinase-substrate relationships. *Methods* **61**, 227-235, doi:10.1016/j.ymeth.2013.03.012 (2013).

Thompson, A. *et al.* Tandem mass tags: a novel quantification strategy for comparative analysis of complex protein mixtures by MS/MS. *Anal Chem* **75**, 1895-1904, doi:10.1021/ac0262560 (2003).

Orlean, P. & Menon, A. K. Thematic review series: lipid posttranslational modifications. GPI anchoring of protein in yeast and mammalian cells, or: how we learned to stop worrying and love glycophospholipids. *J Lipid Res* **48**, 993-1011, doi:10.1194/jlr.R700002-JLR200 (2007).

Menon, I. *et al.* Opsin is a phospholipid flippase. *Curr. Biol.* **21**, 149-153, doi:10.1016/j.cub.2010.12.031 (2011).

Zhong, X., Malhotra, R. & Guidotti, G. ATP uptake in the Golgi and extracellular release require Mcd4 protein and the vacuolar H+-ATPase. *J Biol Chem* **278**, 33436-33444, doi:10.1074/jbc.M305785200 (2003).

Ihara, S. *et al.* PIGN prevents protein aggregation in the endoplasmic reticulum independently of its function in the GPI synthesis. *J Cell Sci* **130**, 602-613, doi:10.1242/jcs.196717 (2017).

Kelleher, D. J. & Gilmore, R. An evolving view of the eukaryotic oligosaccharyltransferase. *Glycobiology* **16**, 47R-62R, doi:10.1093/glycob/cwj066 (2006).

Hyman, R. Somatic genetic analysis of the expression of cell surface molecules. *Trends Genet* **4**, 5-8, doi:10.1016/0168-9525(88)90120-5 (1988).

Guther, M. L., Lee, S., Tetley, L., Acosta-Serrano, A. & Ferguson, M. A. GPI-anchored proteins and free GPI glycolipids of procyclic form Trypanosoma brucei are nonessential for growth, are required for colonization of the tsetse fly, and are not the only components of the surface coat. *Mol Biol Cell* **17**, 5265-5274, doi:10.1091/mbc.e06-08-0702 (2006).

Hong, Y. *et al.* TbGPI16 is an essential component of GPI transamidase in Trypanosoma brucei. *FEBS Lett* **580**, 603-606, doi:10.1016/j.febslet.2005.12.075 (2006).

Lillico, S., Field, M. C., Blundell, P., Coombs, G. H. & Mottram, J. C. Essential roles for GPI-anchored proteins in African trypanosomes revealed using mutants deficient in GPI8. *Mol Biol Cell* **14**, 1182-1194, doi:10.1091/mbc.e02-03-0167 (2003).

Nagamune, K. *et al.* Critical roles of glycosylphosphatidylinositol for Trypanosoma brucei. *Proc Natl Acad Sci U S A* **97**, 10336-10341, doi:10.1073/pnas.180230697 (2000).

Signorell, A., Jelk, J., Rauch, M. & Butikofer, P. Phosphatidylethanolamine is the precursor of the ethanolamine phosphoglycerol moiety bound to eukaryotic elongation factor 1A. *J Biol Chem* **283**, 20320-20329, doi:10.1074/jbc.M802430200 (2008).

Alexander, D. L., Schwartz, K. J., Balber, A. E. & Bangs, J. D. Developmentally regulated trafficking of the lysosomal membrane protein p67 in Trypanosoma brucei. *J Cell Sci* **115**, 3253-3263 (2002).

Manthri, S., Guther, M. L., Izquierdo, L., Acosta-Serrano, A. & Ferguson, M. A. Deletion of the TbALG3 gene demonstrates site-specific N-glycosylation and N-glycan processing in Trypanosoma brucei. *Glycobiology* **18**, 367-383, doi:10.1093/glycob/cwn014 (2008).

Kim, S. *et al.* Dolichol phosphate mannose synthase (DPM1) mutations define congenital disorder of glycosylation Ie (CDG-Ie). *J Clin Invest* **105**, 191-198, doi:10.1172/JCI7302 (2000).

Chalat, M., Menon, I., Turan, Z. & Menon, A. K. Reconstitution of glucosylceramide flip-flop across endoplasmic reticulum: implications for mechanism of glycosphingolipid biosynthesis. *J Biol Chem* **287**, 15523-15532, doi:10.1074/jbc.M112.343038 (2012).

Balakrishnan, R. *et al.* YeastMine--an integrated data warehouse for Saccharomyces cerevisiae data as a multipurpose tool-kit. *Database (Oxford)* **2012**, bar062, doi:10.1093/database/bar062 (2012).

Vazquez, H. M. *et al.* Chemogenetic E-MAP in Saccharomyces cerevisiae for Identification of Membrane Transporters Operating Lipid Flip Flop. *PLoS Genet* **12**, e1006160, doi:10.1371/journal.pgen.1006160 (2016).

Kaplan, R. S. & Pedersen, P. L. Determination of microgram quantities of protein in the presence of milligram levels of lipid with amido black 10B. *Anal Biochem* **150**, 97-104, doi:10.1016/0003-2697(85)90445-2 (1985).

Rouser, G., Siakotos, A. N. & Fleischer, S. Quantitative analysis of phospholipids by thin-layer chromatography and phosphorus analysis of spots. *Lipids* **1**, 85-86, doi:10.1007/BF02668129 (1966).

Beneke, T. *et al.* A CRISPR Cas9 high-throughput genome editing toolkit for kinetoplastids. *R Soc Open Sci* **4**, 170095, doi:10.1098/rsos.170095 (2017).