Title: Temperatures that sterilise males better predict global species distributions than

lethal temperatures

Authors: Steven R Parratt[1*], Benjamin S Walsh[1], Soeren Metelmann[2], Nicola White[1], Andri

Manser[1†], Amanda J Bretman[3], Ary A Hoffmann[4], Rhonda R Snook[5], Tom AR Price[1*]

Affiliations:

[1]Department of Evolution, Ecology and Behavior, University of Liverpool, Liverpool, UK.

[2]Health Protection Research Unit in Emerging and Zoonotic Infections, University of

Liverpool, Liverpool, UK.

[3]Faculty of Biological Sciences, University of Leeds, Leeds, UK.

[4]School of BioSciences, Bio21 Institute, University of Melbourne, Australia.

[5]Department of Zoology, Stockholm University, Stockholm, Sweden

*Correspondence to: s.parratt@liverpool.ac.uk

†Current Address: Department of Evolutionary Biology and Environmental Studies,

University of Zürich, Zürich, Switzerland

Predicting how biodiversity will respond to increased temperatures caused by climate

change is vital. However, our understanding of the traits that determine species' response

to thermal stress remains incomplete. Laboratory measurements of lethal temperatures

have successfully been used to predict global species distributions and the vulnerability of

species to future climate change. However, although it has long been known that fertility is

sensitive to heat stress, temperatures that cause sterility have not been incorporated into

predictions about how climate change will affect biodiversity. Here we show that male

sterility temperatures predict the global distributions of 43 species of *Drosophila*

substantially better than their lethal temperatures. This strongly suggests that thermal

limits to reproduction can underpin how temperature affects species' distributions. High

temperatures impair male fertility across a broad range of animals and plants, so many

organisms may be more vulnerable to high temperatures than currently expected.

Main Text:

We urgently need to understand how rises in temperature will impact biodiversity[1]. To do this

the physiological, behavioral and evolutionary factors that underpin species' current thermal

distributions need to be considered[2,3]. Laboratory-derived estimates of the highest

temperatures at which an organism can function provide measures of species' thermal

tolerances[5,6]. These measures of upper thermal limits have improved the accuracy of

functional species distribution models[4] which can be extrapolated to climate change scenarios

to forecast future distributions[5]. Accurate predictions of species' distributions are invaluable

for prioritizing conservation efforts[6] and predicting the invasion of pests and disease vectors[7].

Upper thermal tolerance limits are usually based on the temperatures that cause loss of

coordinated movement, coma, respiratory failure, or death: the species' critical thermal limit.

Despite these being measured in artificial laboratory conditions, critical limits correlate

reasonably well with species' distributions[8,9] and have been used to estimate species' capacity

to tolerate temperature increases across their current range; their 'thermal safety margins'[3,8].

However, critical limits can be higher than the temperatures that cause seasonal population

declines in nature[10]. Therefore, population persistence may also be impacted by the effects of

sub-lethal temperatures on traits such as reproduction. Sub-lethal temperatures cause losses

in fertility in plants[11], insects[12-14], fish[15], aquatic invertebrates[16], birds[17] and mammals,

including humans[18]. Thermal stress disrupts fertility by impacting physiological

processes[12,17,19] and influencing behavior and phenology[20]. The temperature at which fertility

is lost, the thermal fertility limit (TFL)[20], may be an important part of species' true upper

thermal limits. If TFLs are lower than critical limits, then many organisms will be more

vulnerable to climate change than currently thought. If TFLs and critical limits correlate poorly

then we may misidentify which species are most at risk from rising temperatures. We need to

test whether TFLs measured in laboratory conditions can predict the distribution of natural

populations better than measures of lethal limits.

Comparing lethal and sterilizing temperatures

Fruit flies of the genus *Drosophila* have been a model taxon for thermal biology and

biogeography[8,21] due to their well understood biology, laboratory tractability, and the

diversity of thermal environments the species inhabit. Here, we recorded three measures of

upper thermal limits in adult males from 43 species of *Drosophila*. First, we compared fertility

and survival limits under identical heat-stress conditions. We exposed flies to a 4-hour static

heat stress at a range of temperatures from benign through to lethal. From these data we

estimated both the temperature that is lethal to 80% of individuals (LT80), and the

temperature at which 80% of surviving males are sterilized (TFL80). Measuring thermal traits

under static temperature stress rather than slowly increasing temperatures (i.e. ramping) has

received criticism[22]. However, ramping assays require an immediate observable response,

such as flies losing coordinated motor function. Unfortunately, sterilization is not immediately

observable, so we use static temperatures and assay fertility through subsequent matings. We

score fertility at two time points: (i) cumulatively over 1-6 days post-heat, to capture any

immediate sterilizing effect of heat, and (ii) 7-days after heat-stress to capture any recovery of

fertility or delayed sterility. To compare our estimates of TFL80 and LT80 with a measure of

lethal temperature under ramping thermal stress, we also assayed the CT_{MAX} of each species.

This is the temperature at which males lose coordinated motor function under gradually

increasing temperatures. CT_{MAX} is commonly used across animal systems to predict species'

sensitivity to thermal stress associated with climate change[3,8].

We found that 11 of 43 species experience an 80% loss in fertility at cooler-than-lethal

temperatures immediately following heat-stress (Extended Data Figure 1). Interestingly, rather

than seeing a recovery of fertility over time, the impact of high temperatures on fertility was

more pronounced 7-days post heat stress (Figure 1). Using this delayed measure of fertility,

44% of species (19/43) showed fertility loss at cooler-than-lethal temperatures. The difference

between lethal and fertility limits ranged from $0^{o}C$ to $4.3^{o}C$ (mean of all species = 1.15 ±

$0.22^{o}C$), and LT80 and TFL80 predict dramatically different relative ranking of species'

robustness to high temperature (Extended Data Figure 2). All three thermal limits significantly,

positively correlate with each other (Extended Data Table 1). Despite deriving from different

types of heat-stress, the correlation coefficient between CT_{MAX} and LT80 is larger than that

between TFL80 and either measure of lethal temperature. Relatively low correlations between

survival (measured as CT_{MAX} under dynamic conditions or LT80 under static stress) and fertility

(measured under static heat stress) suggests that they are distinct phenomena, and that

measuring both may be important for understanding how species respond to thermal stress.

Figure 1: Upper lethal temperature (LT80, black circles) and upper thermal fertility limits

(TFL80 measured 7-days after heat stress, pink points) of 43 species of *Drosophila*. 19 of 43

species show significantly lower thermal fertility limits than lethal limits. Species ranked by

LT80 with estimated TFL80 joined by pink bar. We see considerable variation in TFL80 across

the range of lethal values, with some species maintaining fertility at high temperatures. 95% CI

are shown as error bars for both measures, differences between a species' TFL80 and LT80

considered to be significant if these bars do not overlap. Our measure of TFL80 is capped at

the LT80 to reflect ecological realism (see Methods). Fertility loss was measured seven days

post heat stress (for fertility measured immediately following heat-stress see Extended Data

Figure 1).

Our data confirm that fertility loss at sub-lethal temperatures is common across *Drosophila.*

This suggests that studies based on lethal limits alone may overestimate the thermal tolerance

of many species, which could be problematic for efforts to preserve rare and endangered

species or control emerging pests and disease vectors. For example, we found that the crop

pest species *Zaprionus indianus* (African fig fly) has a relatively high upper lethal limit (LT80 =

37.9°C) but a much cooler upper fertility limit (TFL80 = 34.8°C). This switches the species'

thermal hardiness rank from 5th to 17th of species here (Extended Data Figure 2). *Z. indianus* is

currently expanding throughout the New World, and this disparity could have marked

consequences for forecasting its putative range and hence where to concentrate control

efforts[23].

Linking Thermal Limits to Distributions

Our work demonstrates that TFLs can be substantially lower than lethal temperatures, even

when measured under the same conditions in the laboratory. However, the key question is

whether TFLs are linked to organisms' distributions in nature. To test this, we integrated

existing distribution data of each sampled *Drosophila* species with global climate data. From

this we estimated the mean maximum air temperatures species are likely to encounter where

they occur in nature. Previous work using 95 *Drosophila* species found that the mean

maximum summer air temperature measured in this way significantly correlated with CT_{MAX} of

species from dry environments (<1000mm annual rainfall), but not for species from wetter

environments[8]. We first recapitulated this finding in our smaller dataset of 43 species; our

measurement of CT_{MAX} significantly predicts mean maximum environmental air temperature

(PGLS: T_{40} = 2.647, *P* = 0.012) albeit this relationship negatively interacts with annual rainfall

(PGLS: T_{40} = -2.077, P = 0.044) and has reasonably low explanatory power (adjR^2 = 0.186). We

then tested how well our measurements of LT80 and TFL80 predict the mean maximum air

temperature in species' distribution ranges (Extended data Table 1). LT80 significantly

predicted mean maximum environmental temperature (PGLS: T_{40} = 3.360, P = 0.002), but

again explained less than 20% of the variation (adjR^2 = 0.197). However, the relationship

between TFL80 and mean environmental maximum temperature was stronger, both when

TFL80 was measured immediately following heat-stress (PGLS: T_{40} = 4.225, P < 0.001, adjR^2

=0.286) and 7 days later (PGLS: T_{40} = 5.014, P < 0.001, adjR^2 = 0.365). We found no significant

interaction with annual rainfall for either LT80 or TFL80. Comparing across all best–fit models,

TFL80 measured 7-days after heat shock most strongly predicted mean maximum air

temperatures in species' environments, explaining 36.5% of the variation. This is an 85.3% and

95.8% improvement in accuracy (adjR^2) compared to models with LT80 or CT_{MAX} respectively.

Furthermore, we found qualitatively similar results when we used a more conservative 50%

threshold for LT and TFL estimates (Extended Data Table 4). These analyses suggest that TFLs

and species distributions are strongly linked in nature, and that fertility losses due to high

temperature are an important determinant of where species occur.

Our approach based on TFL80 provides a significant improvement over current measures of

upper thermal limits such as CT_{MAX}, which have been valuable in predicting species' responses

to climate change[3,6,8,9]. Nevertheless, all laboratory measures of thermal limits have issues in

reflecting ecological realism. They are optimized for high throughput of species and

individuals, rather than to precisely replicate natural conditions, and often measure heat

effects under artificial conditions. In our tests, we used some fly stocks that have been in

culture for many generations and so may have phenotypes compromised by adaptation to

benign laboratory temperatures or genetic drift. However, we find only weak evidence for lab

adaptation in LT80 and no significant signal for lab adaptation in either CT_{MAX} or TFL80

(Methods and Supplementary Information). We also have not considered variability of

physiological traits across different populations of a given species, and so cannot account for

local adaptation. Despite these limitations, our measure of TFL does appear to be a useful

measure, as it captures 37% of the variation in temperatures among species' distributions.

Figure 2: Distribution Thermal safety margins for 43 *Drosophila* species. Margins

calculated using the physiological limit minus mean maximum air temperature + 1SD

across all recorded locations for each species to capture populations at the boundary of

the thermal range[8]. Grey squares and line are margins calculated with CT_{MAX}, black

circles and lines with LT80, and pink points and lines with TFL80 measured 7-days after

heat stress. Points on or below the horizontal '0' line represent species with populations

that are currently exposed to sterilizing temperatures in the hottest part of the year.

Fitted lines from models summarized in Extended Data Table 3. For central safety

margins see extended Data Figure 3.

Predicting vulnerability to temperature

Predicting species' vulnerability to current and future temperature extremes is critical to protect biodiversity[1]. "Thermal safety margins", often measured as the difference between an organism's physiological thermal limit and the maximum air temperature it is likely to experience may indicate the species' vulnerability to climate change. Although safety margins can use complex measures of environmental temperature to improve accuracy (e.g. microhabitat availability, acclimation[24], thermoregulation behavior of adults[3], duration of exposure[24]), they remain reliant on comparable measures of thermal resistance across species and populations. We found that TFL80 measured seven days after heat-stress produced significantly smaller safety margins than either LT80 or CT_{MAX} (LMER: F = $28.23_{2,84}$, $P < 0.001$, Tukey multiple comparisons P <0.001, Extended Data Figure 3 & Table 3). For populations that already live near the upper edge of their thermal range, safety margins are reduced from a mean of 2.38 ± 3.46°C when based on LT80, to 1.23 ± 3.04°C when based on TFL80 (Figure 2). This predicts that 15 of the 43 species studied here have populations found in environments in which air temperatures exceed safety limits during the hottest part of the year. We illustrate the implications of this with case-study distribution models of *Drosophila flavomontana;* safety margins based on TFL80 predict a 17.9% reduction in habitable landscape compared to an identical LT80-based model under current climate conditions (Figure 3A). The disparity between predictions based on sterility and lethality grew to 48.0% by the year 2080 under moderately optimistic future climate forecasts (ICCP-AR5 RCP 4.5, Figure 3B), and to 58.9% under pessimistic climate change scenarios (ICCP-AR5 RCP 8.5, Figure 3C). TFL-based models also predict that by 2080 the available habitat for *D. flavomontana* will have reduced by 42.3% and 62.9% under RCP4.5 and RCP8.5 respectively.

Figure 3: Potential habitat range of *Drosophila flavomontana* (LT80 = 35.4°C, TFL80 = 31.9°C)

under A) current and B & C) possible future climate scenarios (B = RCP4.5 'moderately

optimistic', C = RCP8.5 'pessimistic', predicted for 2060 - 2080). Colored areas in each panel

represent suitable habitat range predicted by the model that excludes maximum

temperature. Red areas show regions where maximum summer temperatures exceed LT80.

Orange areas show regions where maximum summertime temperatures exceed TFL80. Blue

regions are areas where limits for *D. flavomontana* are not exceeded all year.

If our data for *Drosophila* can be extrapolated to other organisms, then male fertility losses

at high temperatures may be common, occurring at substantially lower temperatures than

the species' upper lethal limit. The limited data on fertility at extreme temperatures supports

this, with losses in male fertility at high temperatures seen across a broad diversity of

organisms[20]. Even species that might be predicted to have high thermal tolerance can show

evidence of thermal fertility losses. For instance, the zebra finch, a high temperature adapted

desert-dwelling endothermic species with naturally high body temperature and good

thermoregulation, shows substantial damage to sperm at temperatures it regularly

experiences in nature[17]. Behavioral thermoregulation could potentially reduce the impact of

high temperatures on fertility in nature. However, while some evidence suggests that

Drosophila are able to behaviorally thermoregulate in the lab[25], evidence for flies choosing

cooler microclimates such as leaf litter, shade, or higher altitudes in nature is mixed[26].

Further, many species are able to survive high temperature periods by aestivating as adults,

eggs or pupae. This may explain why our data predict negative thermal safety margins for

some species. Despite these potential mechanisms, we still find that species' distributions

are predicted by thermal fertility limits.

Our work emphasizes that temperature-driven fertility losses may be a major threat to

biodiversity during climate change. We urgently need to understand the range of organisms

likely to suffer similar fertility losses in nature, and the traits that predict vulnerability.

However, we currently do not understand the physiology underlying variation in TFLs

between species, nor the selective forces that created this variation. Ultimately, we need to

know whether evolution for higher TFLs will allow species to adapt to a warming

environment.

Main Text References

1.   Pecl, G. T. *et al.* Biodiversity redistribution under climate change: Impacts on

ecosystems and human well-being. *Science* **355,** eaai9214 (2017).

2.   Chen, I.-C., Hill, J. K., Ohlemüller, R., Roy, D. B. & Thomas, C. D. Rapid Range Shifts of

Species Associated with High Levels of Climate Warming. *Science* **333,** 1024–1026

(2011).

3.   Sunday, J. M. *et al.* Thermal-safety margins and the necessity of thermoregulatory

behavior across latitude and elevation. *Proc Natl Acad Sci USA* **111,** 5610–5615 (2014).

4.   Kearney, M. & Porter, W. Mechanistic niche modelling: combining physiological and

spatial data to predict species' ranges. *Ecol Lett* **12,** 334–350 (2009).

5.   Elith, J. & Leathwick, J. R. Species Distribution Models: Ecological Explanation and

Prediction Across Space and Time. *Annu. Rev. Ecol. Evol. Syst.* **40,** 677–697 (2009).

6.   Nowakowski, A. J. *et al.* Thermal biology mediates responses of amphibians and reptiles

to habitat modification. *Ecol Lett* **21,** 345–355 (2018).

7.   Metelmann, S. *et al.* The UK's suitability for Aedes albopictus in current and future

climates. *J Roy Soc Interface* **16,** 20180761 (2019).

8.   Kellermann, V. *et al.* Upper thermal limits of Drosophila are linked to species

distributions and strongly constrained phylogenetically. *Proc Natl Acad Sci U.S.A.* **109,**

16228–16233 (2012).

9.   Lancaster, L. T. & Humphreys, A. M. Global variation in the thermal tolerances of plants.

Proc Natl Acad Sci USA **117,** 13580–13587 (2020).

10.   Rezende, E. L., Bozinovic, F., Szilàgyi, A. & Santos, M. Predicting temperature mortality

and selection in natural Drosophila populations. *Science* **369,** 1242–1245 (2020).

11.   Sage, T. L. *et al.* The effect of high temperature stress on male and female reproduction

in plants. *Field Crop Res* **182,** 30–42 (2015).

12.   Sales, K. *et al.* Experimental heatwaves compromise sperm function and cause

transgenerational damage in a model insect. *Nat Commun* **9,** 4771 (2018).

13.   Porcelli, D., Gaston, K. J., Butlin, R. K. & Snook, R. R. Local adaptation of reproductive

performance during thermal stress. *J Evol Biol* **30,** 422–429 (2016).

14.   Saxon, A. D., O'Brien, E. K. & Bridle, J. R. Temperature fluctuations during development

reduce male fitness and may limit adaptive potential in tropical rainforest Drosophila. *J*

Evol Biol **31,** 405–415 (2018).

15.   Breckels, R. D. & Neff, B. D. The effects of elevated temperature on the sexual traits,

immunology and survivorship of a tropical ectotherm. *J Exp Biol* **216,** 2658–2664 (2013).

16. Paxton, C. W., Baria, M. V. B., Weis, V. M. & Harii, S. Effect of elevated temperature on

fecundity and reproductive timing in the coral Acropora digitifera. *Zygote* **24,** 511–516

(2016).

17. Hurley, L. L., McDiarmid, C. S., Friesen, C. R., Griffith, S. C. & Rowe, M. Experimental

heatwaves negatively impact sperm quality in the zebra finch. *Proc R Soc B* **285,**

20172547 (2018).

18. Yogev, L. *et al.* Seasonal variations in pre- and post-thaw donor sperm quality. *Hum*

Repro **19,** 880–885 (2004).

19. Rohmer, C., David, J. R., Moreteau, B. & Joly, D. Heat induced male sterility in

Drosophila melanogaster: Adaptive genetic variations among geographic populations

and role of the Y chromosome. *J Exp Biol* **207,** 2735–2743 (2004).

20. Walsh, B. S. *et al.* The Impact of Climate Change on Fertility. *Trends Ecol. Evol* **34,** 249–

259 (2019).

21. Balanyá, J., Oller, J. M., Huey, R. B., Gilchrist, G. W. & Serra, L. Global Genetic Change

Tracks Global Climate Warming in Drosophila subobscura. *Science* **313,** 1773–1775

(2006).

22. Terblanche, J. S. *et al.* Ecologically relevant measures of tolerance to potentially lethal

temperatures. *J Exp Biol* **214,** 3713–3725 (2011).

23.   van der Linde, K. *et al.* First Records of Zaprionus indianus (Diptera: Drosophilidae), a

pest species on commercial fruits from Panama and the United States of America. *Flo*

Ento **89,** 402–404 (2006).

24.   Overgaard, J., Kearney, M. R. & Hoffmann, A. A. Sensitivity to thermal extremes in

Australian Drosophila implies similar impacts of climate change on the distribution of

widespread and tropical species. *Glob Chan Biol* **20,** 1738–1750 (2014).

25.   Dillon, M. E., Wang, G., Garrity, P. A. & Huey, R. B. Thermal preference in Drosophila. *J*

Therm Biol **34,** 109–119 (2009).

26.   Tratter Kinzner, M. *et al.* Is temperature preference in the laboratory ecologically

relevant for the field? The case of Drosophila nigrosparsa. *Glob Ecol Con* **18,** e00638

(2019).

Materials and Methods:

<u>Animal maintenance:</u>

Drosophila species were obtained from stock centers, other research groups, or field

collections (Supplementary Information Table 1). All species were kept in temperature-

controlled rooms at temperatures selected to avoid effects of heat-hardening or stress

(Supplementary Information Table 1). All rearing temperatures were well below those

observed to be stressful, indeed our warmest rearing temperature (25°C) was several

degrees cooler than the coolest lethal temperature we observe for any species (32°C). All

stocks were kept at moderate density ($20-50$ breeding pairs per vial or $50-100$ pairs per

bottle culture) at 12:12 L:D and ambient humidity. Cultures were tipped to fresh food every 7

days and a new generation was made every $4-6$ weeks depending on the speed of species

development. Species were reared on one of four food types selected to minimize

heterogeneous nutritional stress; A = ASG (10g agar, 85g Sucrose, 20g yeast extract, 60g

maize, 1000ml H_2O, 25ml 10% Nipagin), B = Banana (10g agar, 30g Yeast extract, 150g pulped

banana, 50g Molasses, 30g Malt Extract, 25ml 10% Nipagin, 1000ml H_2O), P = Propionic (10g

Agar, 20g Yeast extract, 70g cornmeal, 10g soya flour, 80g Malt Extract, 22g Molasses, 14ml

10% nipagin, 6ml Propionic acid, 1000ml H_2O), M = Malt (10g agar, 20g Yeast, 60g Semolina,

80g Malt, 14ml 10% Nipagin, 5ml Propionic acid, 1000ml H_2O).

<u>Measuring upper thermal limits</u>

We assayed three metrics of upper thermal limits in sexually mature males from 43 species

of *Drosophila*: Lethal Temperature (LT), Thermal Fertility Limit (TFL) and Maximum Critical

Temperature (CT_{MAX}). We tested these limits in only males because of the substantial

evidence that male fertility is particularly vulnerable to high temperatures[20]. All experiments

were blinded by numbering vials randomly across treatments.

Measuring Lethal Temperature (LT) & Thermal Fertility Limit (TFL):

We measured LT and TFL under static temperature conditions by allocating groups of flies to

separate four-hour temperature pulses and recording their survival and fertility. Using static

temperatures to measure thermal threshold traits has received criticism[22]. However, the use

of ramping temperatures requires an immediate observable response, such as flies losing

their coordinated motor function, which allows the researcher to visually determine the

temperature at which the function was lost. Unfortunately, in _Drosophila_ fertility is internal

and has no visible marker, so there is no directly observable response that would indicate

that a male has become sterile as temperatures increase. Hence, we have to use static

temperatures and assay fertility by subsequently mating the male to females. Measuring LT

under static temperature stress allows us to directly compare measures of fertility loss and

lethality under identical heating conditions.

Newly eclosed virgin adult males were collected into plastic vials in groups of 10 over a 48-

hour period and allowed to sexually mature for 7 – 21 days depending on species (see

Supplementary Information Table 1 for maturation times). Mature males were transferred to

fresh vials with standard ASG food and a cotton-wool stopper and allocated to 3D-printed

floating racks in pre-heated waterbaths (Grant TXF200) set to a range of temperatures

(Supplementary Information Table 1). ASG food was used during heating for all species

because preliminary work had shown that the survival under heat stress was influenced by

food type. We monitored the temperature during the assay in a fresh ASG vial without flies

with a k-type thermocouple attached to a data logger (Pico Technology TC-08). Males were

heated for 4 hours between ~10am - ~2pm and then returned to temperature-controlled

rooms set to the species' benign temperature. We scored survival of males the next morning

to account for any immediate recovery or delayed death. We scored flies as 'dead' if they

were completely immobile or had become stuck on their backs in the fly food – the latter

would indicate that the flies were completely incapacitated which would likely be lethal in

the wild. Surviving males were individually aspirated into separate vials containing their

designated food type and 3-4 sexually mature virgin females. Males remained in these vials

at their benign temperature and were allowed to mate freely for 6 days. After 6 days males

were transferred to a second vial with 1-2 more virgin females and allowed to mate for 24

hours. Giving males two sets of females to mate with allows us to score fertility at two time

points to capture any recovery or delayed sterilizing effect of heat over time. Females in both

sets of vials were left to lay eggs for 3-5 days (depending on fecundity of the species) after

the male had been removed. Vials were scored for the presence of larvae or their distinctive

track marks in the food. We found no significant differences in mortality between heat-

stressed and control males during the 1-week oviposition period, so we do not measure LT

multiple time points.

We used the `drm()` function in the `drc` package in R to fit dose response curves to survival

and fertility data across temperatures for each species. The models generate point estimates

of the temperature required to reduce survival or fertility by a given percentage relative to

the maximum value in the data. We estimate thresholds based on 80% reduction (LT80 and

TFL80 hereafter) because this is likely to represent a substantial threat to population stability

for most species. To test the robustness of our findings we repeated all analyses with more

conservative 50% reduction estimates (LT50 and TFL50 hereafter) and present these results

in the Extended Data (Extended Data Table 3). We used 3-parameter versions of the log-

logistic model which fixes the lower limit of the predicted curve to 0 to reflect that we expect

fertility and survival to eventually reach 0 over an infinite range of temperatures. In these

models we do not consider dead males to be sterile to avoid statistically confounding the

effects of heat stress on survival and fertility. We only allow TFL80 to be lower than or equal

to the species' LT80 because estimating 80% fertility loss in populations that have already

experienced >80% mortality provides little informative data. Furthermore, we only consider a

species' TFL to be statistically lower than its LT if the 95% confidence intervals of these two

point-estimates do not overlap.

In total, 14742 males were exposed to heat treatment, of which 10925 survived and 9064

went on to be tested for heat-induced sterility. The mean sample size at each temperature

per species was 36.5 ± 0.5sem, ranging from $10 - 88$ individuals. Because fertility is

inherently only measurable in males that survive heat stress, we may be measuring TFLs in

more thermally hardy individuals – i.e. those that survive. In the context of our study, this

will make our estimate of any difference between LTs and TFLs more conservative.

Measuring CT$_{MAX}$:

Studies typically estimate upper thermal limits under dynamically ramping heat

conditions[6,8,24,27], rather than under static heat-stress as we have used here for TFL and LT.

To compare our static measures of TFL and LT with patterns of heat tolerance found using

the more widely used dynamic assays, we measured upper critical limits of our 43 *Drosophila*

species under gradually ramping heat conditions – their critical thermal maxima (CT_{MAX}). This

allows us to compare how static and dynamic measures of critical heat tolerance predict

species' distributions.

We measure CT_{MAX} as the mean temperature at which male flies lose coordinated movement

and are unable to right themselves. Sexually mature males were anesthetized with CO_2 and

isolated into individual glass sample tubes sealed with a rubber stopper. Sample tubes were

left at ambient room temperature for at least 30 minutes to allow males to fully recover from

CO_2 anesthetization. Preliminary trials found no difference in CT_{MAX} between anesthetized

and non-anesthetized male *D. melanogaster.* Sample tubes were then submerged in a water-

filled glass aquarium held at the rearing temperature for the species (18, 23 or 25^{o}C) by a

heated circulator (Grant TXF200). The temperature was then increased at a rate of 0.1^{o}C per

minute and the temperature at which flies collapsed for 30 seconds and did not right

themselves after being disturbed (tapping the vial) was recorded. The temperature of the

waterbath was continuously recorded independently of the heating unit with a data logger

(Elitech RC-61). The mean sample size for each species for CT_{MAX} assays was 19.53 ± 1.77.

Genetic and physiological changes due to lab domestication have been reported in a number

of insects[28-30]. Because the 43 species of *Drosophila* we use here varied in the time spent in

lab culture we tested for confounding effects of lab adaptation on physiological thermal

limits (Supplementary Material). We found only minor signals of lab adaptation in measures

of LT80, and no significant signal in either CT_{MAX} or TFL80 (Supplementary Information). We

see no strong evidence that lab adaptation has degraded inter-species variation in thermal

traits. Further, we can replicate our core results with a subset of the 11 most recently caught

species (Supplementary Material). This supports work in other dipteran systems that have

shown long-term lab stocks to be good proxies for thermal traits in natural populations[31].

<u>Repeatability</u>

It was impossible to assay upper limits across all 43 species simultaneously due to the

number of flies that could be processed simultaneously, and the different temperature

ranges required for each species. Further, inter-species variation in development time

prohibited us from synchronizing all 43 species to run the assay in a completely randomized

block design. Instead, we independently assayed CT_{MAX}, TFL and LT for each species between

June 2018 – November 2019.

To test the repeatability of our measure of LT80 we ran a single control block in February

2020 with 22 species that produced enough sexually mature virgin males in synchrony. We

simultaneously exposed these flies to a range of 6 temperatures that covered all species'

estimated LT80. We then tested the strength of the correlation between our original LT80

estimates and those given by this replicate block. We determined if any given species

showed significantly different estimates based on whether the confidence intervals of the

two runs overlapped (Extended Data Figure 4). Sample sizes for this control run ranged from

3 − 10 males per temperature per species, and involved 1152 male flies in total.

To test repeatability for our TFL assay we ran two independent blocks of the full assay on

Drosophila virilis and compared the TFL80 estimate. The blocks were run 6 months apart on

the same equipment and fly stock, and data were collected by two separate researchers. We

compare the estimated TFL80 from each run and consider differences to be significant if the

95% confidence intervals do not overlap (Extended Data Figure 5).

Repeatability of CT_{MAX} was estimated by randomly allocating species from the same rearing

temperatures into blocks of 3 − 5 species and measuring their mean CT_{MAX} as described

above. We compared mean CT_{MAX} values with those obtained when species were run

individually (Extended Data Figure 6).

<u>Correlations between upper thermal limits.</u>

We explored the correlation between all three physiological limits to test if TFL captures

additional differences in species' thermal hardiness. We used several parallel approaches to

test the strength of correlations. We accounted for evolutionary relationships between

species by including information from a *Drosophila* phylogeny published by Patrik Röhner

and colleagues[32]. Firstly, we calculated phylogenetically independent contrasts (PICs) for

each trait and correlated these using standard linear models forced through the origin at 0.

This approach assumes that each trait completely evolves along a Brownian model of

evolution. Second, we analyzed each pair of traits with phylogenetically controlled least

squares regression (`pgls` in the R package `caper`) in which we estimate the phylogenetic

signal (Pagels λ) in model residuals with a maximum likelihood approach. This approach is

sensitive to the order of x and y variables, so we ran both inversions of each pairwise

correlation. Finally, we used the `corphylo` function in the `ape` package to generate a

correlation matrix of all three traits in which phylogenetic signal for each is estimated

assuming an Ornstein-Uhlenbeck process. We also compared the output of these

phylogenetically controlled approaches to standard linear models. In all analyses we used

TFL80 measured 7-days post heat stress because we found that the sterilizing effect of

temperature is delayed in many species.

<u>Species distributions and environmental variables:</u>

Geographic distribution data for each species were obtained manually from TaxoDros.uzh.ch

as coordinates and location names. Location data was systematically cleaned with functions

in the `CoordinateCleaner` R package to check for appropriate decimalization, remove invalid

coordinates, remove country centroids, remove locations over water, remove locations

centered on biodiversity institutions, remove location with equal longitude and latitude, and

flag outliers. Remaining coordinates were then plotted on maps and outliers were

investigated in the primary literature and removed or corrected when appropriate. To

minimize spatially biased oversampling we then used the `thin` function in `spThin` R

package to reduce the dataset down so that no two points for a given species occur within

10km of each other. We ran this independently for every species except three small island

endemics (*D. santomea*, *D. sechellia* and *D. madierensis*) to preserve variation in their sparse

distribution records.

The cleaned dataset of geolocations was integrated with rasterized bioclimate data from the

WorldClim V2.0 database using functions from the `Raster` package in R[33]. Our intention was

not to fully reassess multiple climate variables as predictors for *Drosophila* distributions,

because this has received extensive attention in the literature[8,24]. Instead, we tested if TFLs

correlate and interact with variables previously identified as important for predicting

physiological limits and distribution in *Drosophila*; the maximum temperature reached in the

hottest summer month (WorldClim Bio5 – T_{max} hereafter) and annual precipitation

(WorldClim Bio12 - P_{ANN} hereafter). Mean and standard deviation of each climate variable for

each species were calculated across all global locations.

Comparing fit of CT_{MAX}, LT and TFL to species' thermal habitats

We used phylogenetically controlled linear models to test the relationship between each

physiological limit and maximum air temperatures found in species' natural environment

(T_{max}). We analysed our data with T_{max} as the response variable because previous work has

found that species' phylogenetically constrained physiology determines where they occur

more readily than local adaptation drives phenotypic change[8]. Because of the strong

correlations between the three measures of upper thermal limits we could not include them

all as predictors in the same model with T_{max} as the dependent variable. Instead, we fit

separate models with each of TFL80, LT80 or CT_{MAX} as independent predictors of species' T_{max}

and compare the goodness of fit. We included annual precipitation (P_{ANN}) as an interacting

covariate with each thermal limit because previous work has found that the relationship

between lethal limits and T_{max} degrades for species that occur in more humid habitats[8]. We

used AICc scores and visualization of residuals to retain or drop the interaction term from

models. We do not retain P_{ANN} as a main term because we are not testing if spatial patterns

of temperature correlate with spatial patterns of rainfall independently of *Drosophila*

biology. All predictors are centered and scaled to the mean. We compared the adjusted R^2 of

each best-fit model for each upper limit.

<u>Calculating and modelling thermal safety margins</u>

Thermal safety margins represent the physiological capacity of organisms to tolerate

temperature increases in their native distribution. As per Kellermann (2012)[8], we calculated

this in two ways: "Central Safety Margins" are the physiological limit minus the mean T_{max} for

each species, "Distribution Safety Margins" are the physiological limit minus the mean T_{max} +

1 standard deviation. "Distribution" safety margins have been argued to account for

populations of species that live at the upper thermal range of species' distribution. Safety

margins were calculated separately using either CT_{MAX}, TFL80 or LT80 as the physiological

limit.

We tested for significant differences between safety margins calculated with each

physiological measure with linear mixed models using the `lme4` package in R. We set

'physiological limit' (CT_{MAX}, TFL80 or LT80) and mean absolute latitude of each species known

distribution locations as interacting predictors. Latitude was fit as a curvilinear (latitude^2)

predictor following visualization of the raw data and because previous studies that have

found curvilinear patterns in safety margins over latitudinal range[3,8,34]. We included species

identity as a random intercept to account for non-independence between safety margin

values calculated for TFL80, LT80 and CT_{MAX} for each species. We used AICc and visualization

of model residuals to reach a minimum best-fit model through stepwise selection. We used

Type II Wald's F test in the `car` package to determine the significance of main effects in final

models. We used post-hoc Tukey contrasts implemented with the `emmeans` package to

detect significant differences between safety margins based on the three physiological limits.

"Central" and "Distribution" safety margins were modelled separately.

<u>Predicting current and putative future distributions of *D. flavomontana*</u>

As a case-study demonstration of the importance of TFLs, we used MaxEnt v3.4.1[35] to model

potential distributions of *D. flavomontana*. Occurrence data from TaxoDros.uzh.ch were

further slimmed so that points within 25km of each other were clustered together (*N* = 24).

First, we chose climatic and other variables to predict a potential distribution of *D.*

flavomontana that is independent of the upper thermal limit. After excluding variables that

strongly correlated or had little to no impact on the model results (<3% contribution), we

used four of the tested 15 variables: mean winter temperature (WC Bio 11, from the

WorldClim2.0 database), precipitation seasonality (WC Bio 15), an agricultural index[36], and

elevation[37]. We then ran 10 cross-validations with maximum iterations of 5000, using 10000

random pseudo-absences. To distinguish suitable from unsuitable regions, we used the

threshold that maximized Youden's index (maximum training sensitivity plus specificity)[38].

We then constrained the suitable area by the upper physiological thermal limits measured

above: i) the lethal temperature measured as LT80 (35.4°C), and ii) the sterilizing

temperature measured as absolute TFL80 (31.9°C). We use the maximum temperature of the

warmest month as reference (T_{max}). Using T_{max} directly as a predictor variable for *D.*

flavomontana had close to no effect on the distribution (0.1% contribution).

For future predictions, we chose two carbon emission scenarios from the IPCC-AR5: RCP4.5

(moderately optimistic) and RCP8.5 (pessimistic). We used precipitation and temperature

predictions of the NorESM1-M model from the NEX-NASA GDDP data set[39], as this model

produces a median trend in terms of temperature increase[40]. The agricultural index and

elevation had to be used as present day data. Predicting and constraining the possible future

distribution followed the same procedure as for the current distribution.

Data and Code Availability:

All data and analysis code required to recreate this study will be made freely available on

Dryad upon acceptance of the paper.

Methods References:

27.   Terblanche, J. S., Klok, C. J., Krafsur, E. S. & Chown, S. L. Phenotypic plasticity and

geographic variation in thermal tolerance and water loss of the tsetse Glossina

pallidipes (Diptera: Glossinidae): Implications for distribution modelling. *Am J Trop Med*

Hyg **74,** 786–794 (2006).

28.   Alston, M. A., Lee, J., Moore, M. E., Kingsolver, J. G. & Willett, C. S. The ghost of

temperature past: interactive effects of previous and current thermal conditions on

gene expression in Manduca sexta. *J Exp Biol* **223,** jeb213975 (2020).

29.   Santos, J. *et al.* Tracking changes in chromosomal arrangements and their genetic

content during adaptation. *J Evol Biol* **29,** 1151–1167 (2016).

30.   Krebs, R. A., Roberts, S. P., Bettencourt, B. R. & Feder, M. E. Changes in thermotolerance

and Hsp70 expression with domestication in Drosophila melanogaster. *J Evol Biol* **14,**

75–82 (2001).

31.   Lyons, C. L., Coetzee, M., Terblanche, J. S. & Chown, S. L. Thermal limits of wild and

laboratory strains of two African malaria vector species, Anopheles arabiensis and

Anopheles funestus. *Malar. J.* **11,** 226–14 (2012).

32.   Rohner, P. T. *et al.* Interrelations of global macroecological patterns in wing and thorax

size, sexual size dimorphism, and range size of the Drosophilidae. *Ecography* **41,** 1707–

1717 (2018).

33.   Fick, S. E. & Hijmans, R. J. WorldClim 2: new 1-km spatial resolution climate surfaces for

global land areas. *Intl J Climatol* **37,** 4302–4315 (2017).

34.   Deutsch, C. A. *et al.* Impacts of climate warming on terrestrial ectotherms across

latitude. *Proc Natl Acad Sci USA* **105,** 6668–6672 (2008).

35.   Phillips, S. J., Anderson, R. P. & Schapire, R. E. Maximum entropy modeling of species

geographic distributions. *Ecol Mod* **190,** 231–259 (2006).

36.   Ramankutty, N., Evan, A. T., Monfreda, C. & Foley, J. A. Farming the planet: 1.

Geographic distribution of global agricultural lands in the year 2000. *Glob Biogeochem*

Cy **22** (2008).

37.   Amante, C. & Eakins, B. W. Etopo1 1 Arc-Minute Global Relief Model: Procedures, Data

Sources and Analysis. *National Geographic Data Center* (2009).

38.   Liu, C., White, M. & Newell, G. Selecting thresholds for the prediction of species

occurrence with presence-only data. *J Biogeog* **40,** 778–789 (2013).

39.   Thrasher, B., Maurer, E. P., McKellar, C. & Duffy, P. B. Technical Note: Bias correcting

climate model simulated daily temperature extremes with quantile mapping. *Hydro.*

Earth Syst Sci **16,** 3309–3314 (2012).

40.   Jones, A. E. *et al.* Bluetongue risk under future climates. *Nat Clim Change* **9,** 153–157

(2019).

Acknowledgements:

Natasha Mannion and Dr Angela Sims for assistance with experiments, Dr Ben Longdon and

Dr Katherine Roberts for flies, Rowan Connell for designing 3D-printed equipment. **Funding:**

NERC grant NE/P002692/1 to TP, AB, AH & RS. SNF P300PA_177830 to AM. NIHR HPRU EZI to

SM.

Author Contributions: Conception: TP, AB, RS, AH, SP. Methodology & data collection: SP,

BW, NW. Data curation: SP & SM. Analysis: SP, AM & SM. Original Draft: SP, TP, AB, RS & AH.

Review & Editing: All authors. **Data and materials availability:** All data and analysis R code

will be deposited on Dryad upon acceptance of this manuscript.

Competing Interest:

The authors have no competing interest to declare.

Supplementary Information is available for this paper, including Supplementary Table 1 –

Details of species used.

Correspondence and requests for materials should be addressed to Steven Parratt

(S.parratt@liverpool.ac.uk).

 Extended Data Legends:

 Extended Data Figure 1: TFL80 and LT80 of Drosophila immediately after heat stress. Several

 species of Drosophila lose 80% fertility at cooler-than-lethal temperatures immediately

 following heat-shock. LT80 (black circles) and TFL80 (pink circles). Errors for both measures are

 95% confidence intervals generated from dose response model estimates. Fertility loss

 measured as ability to sire any offspring between $1 - 6$ days post heat-stress.

 Extended Data Figure 2: Relative ranking of species thermal robustness based on LT80 and

 TFL80. Most to least heat tolerant species ranked from top to bottom. Pink bars indicate species

 with significantly lower TFL80 than LT80. Data based on TFL80 recorded 7-days post-heat shock.

 Extended Data Figure 3: Central Thermal safety margins for 43 Drosophila species. Margins

 calculated using the physiological limit minus the mean maximum temperature across all

 recorded locations for each species. Grey squares and line are safety margins calculated using

 CT_{MAX} values, black circles and lines are based on LT80, and pink points and lines are based on

 TFL80 measured 7-days after heat stress. The three physiological limits produce significantly

 different safety margin estimates (Central safety margins: lmer: $F_2 = 28.23$, $P < 0.001$, estimate =

 -1.17). TFL80 measured at 7 days post-heat stress predicts mean central safety margins of 4.97 $\pm$

 2.42^{o}C. This is significantly smaller than margins based on CT_{MAX} (mean central safety margins:

 8.81 $\pm$ 3.26^{o}C, pairwise Tukey tests P<0.001) and LT80 (central safety margins: 6.13 $\pm$ 3.09^{o}C,

 pairwise Tukey tests P<0.001). Latitude shown here is the mean absolute latitude i.e. the sign of

 negative latitudes has been removed to give relative distance from the equator. Fitted lines are

 predictions from models in which the limit type (TFL, LT or CT_{MAX}) and latitude[2] are fixed effects

and species identity is a random effect to account for repeated measures across species (see

Extended Data Table 3).

Extended Data Figure 4: Repeatability of LT80. Repeatability was high across the 22 *Drosophila*

species tested. Left panel: There was no significant difference in the estimate of LT80 between

full runs and control repeats in any of these species (significance measured as non-overlapping

CIs of point estimates of LT80). Pink points = estimate from single species assay, gold point =

estimate from simultaneous multispecies control assay. Right panel: The correlation between

the two independent measurements of LT80 across these species was strongly positive and

explained a high degree of variation in the data (coefficient = 1.06, R^2 = 0.96).

Extended Data Figure 5: Repeatability of TFL80 for Drosophila virilis. We independently

repeated the TFL80 assay for one species in which we found a significant difference between

LT80 and TFL80. Two independent runs of sexually mature males were conducted by two

researchers 6 months apart. Red = original data used to calculate TFL80 for *D. virilis* in this

manuscript, blue = repeated assay. Dashed lines intersect at 80% threshold. Line fits predicted

by 3-parameter dose-response model. The fly stock and equipment were identical.

652 **Extended Data Figure 6: Repeatability of knockdown CT$_{MAX}$.** Left panel; mean CT$_{MAX}$

653 temperature recorded from species' independent assays (gold points), and in mixed species

654 control blocks (pink points). There was no global significant difference between CT$_{MAX}$

655 estimates across all species (Block: F$_{1,831}$ = 3.246, *P* = 0.072). However a significant

656 experimental block*species interaction was found for 4 of the 41 species tested (interaction

657 term: F$_{40,831}$ = 3.7589, P < 0,001); *Drosophila sechelia, D. nasuta, D. yakuba* & *D. tiesseri* all

658 scored significantly higher CT$_{MAX}$ in our mixed species control block than in their own individual

659 CT$_{MAX}$ assays. Right panel: The correlation between estimated CT$_{MAX}$ values from both blocks

660 was strongly positive (coefficient = 1.07, F$_{1,37}$ = 309.84, P < 0.001), and explained 89% of the

661 variation.

663 **Extended Data Table 1:** Correlations between TFL80, LT80, and CT$_{MAX}$. Multiple methods for

664 phylogenetically controlled correlations were used to test the extent to which these three

665 physiological limits are proxies for one another. In all four methods TFL80 correlates less

666 strongly with CT$_{max}$ and LT80 than the two measures of critical limits correlate with each other.

667 Phylogenetic least squares (pgls) allows for estimation of phylogenetic signal (Pagel's λ) in the

668 residuals of the y-variable, thus correlation coefficients from this method are sensitive to

669 which variable is assigned as the predictor and which as the response. To account for this, we

670 present two pgls() outputs for each pair of traits. Phylogenetic independent contrasts (PIC)

671 essentially assumes a phylogenetic signal of λ = 1 (complete Brownian motion evolution of the

672 trait). Linear models ('lm') make no adjustment for non-independence in trait values between

673 closely related species. `corphylo` from the `ape` R package allows for the three-way

674 correlation matrix to be estimated simultaneously and produces estimates of phylogenetic

signal (d) of each trait under an Ornstein-Uhlenbeck process. TFL80 here measured 7-days post

heat-stress. "-"denotes model estimates are identical to the reciprocal x~y configuration.

Extended Data Table 2: Summaries of phylogenetically controlled models that predict

maximum environmental temperature (T_{max}) by either CT_{MAX}, LT80 or TFL80 independently of

each other. Annual precipitation (P_{ANN}) was included as an interaction term with physiological

limits. adjR^2 and phylogenetic signal in model residuals (Pagel's λ) given for final best-fit

model derived from AICc model selection and inspection of model residuals. Terms retained

after model selection shown in italics. All continuous predictors centred and scaled to the

mean.

Extended Data Table 3: Summary model fits of species' safety margins (SM) to absolute

latitude (Lat^2), the physiological limit used in their calculation ('Limit'), and the interaction

between 'Lat^2' and 'Limit'. 'Species' identity is included as a random intercept term. We show

models of both "Central safety margin" and the "Distribution Safety Margin" *as per*

Kellermann[8]. Central safety margins are calculated as the difference between the physiological

limit and the mean maximum temperature experienced in every know location in the species'

distribution. "Distribution safety margins" capture the environmental conditions at the upper

thermal edge of specie's known distribution by adding 1 standard deviation to the mean

maximum temperature. R^2 given for final best-fit model derived from AICc model selection and

inspection of model residuals. Significance of main terms given by Type II sum of squares F-

tests with Kenward-Rogers degrees of freedom. Terms retained after model selection shown

in italics. Post-hoc Tukey tests were run with the `emmeans` package to identify significant

differences between levels of "Limits". TFL80 measured 7-days post heat shock.

Extended Data Table 4: Using 50% thresholds to predict distributions. Summaries of

phylogenetically controlled models that predict maximum environmental temperature (T_{max})

by either CT_{MAX}, LT50 or TFL50 independently of each other. Annual precipitation (P_{ANN}) was

included as an interaction term with physiological limits. R^2 and phylogenetic signal in model

residuals (Pagel's λ) given for final best-fit model derived from AICc model selection and

inspection of model residuals. Terms retained after model selection shown in italics. All

continuous predictors centred and scaled to the mean.