Title: Temperatures that sterilise males better predict global distributions of species than lethal temperatures

Authors: Steven R Parratt[1*], Benjamin S Walsh[1], Soeren Metelmann[2], Nicola White[1], Andri Manser[1†], Amanda J Bretman[3], Ary A Hoffmann[4], Rhonda R Snook[5], Tom AR Price[1*]

Affiliations:

[1] Department of Evolution, Ecology and Behavior, University of Liverpool, Liverpool, UK.

[2] Health Protection Research Unit in Emerging and Zoonotic Infections, University of Liverpool, Liverpool, UK.

[3] Faculty of Biological Sciences, University of Leeds, Leeds, UK.

[4] School of BioSciences, Bio21 Institute, University of Melbourne, Australia.

[5] Department of Zoology, Stockholm University, Stockholm, Sweden

*Correspondence to:s.parratt@liverpool.ac.uk

†Current Address: Department of Evolutionary Biology and Environmental Studies, University of Zürich, Zürich, Switzerland

Abstract:

Climate change is well understood to be a major threat to biodiversity, but sublethal impacts of high temperatures, such as reduced fertility, have been poorly studied. We examined a panel of 43 *Drosophila* species, finding that 19 experience significant fertility loss at temperatures up to 4.3°C cooler than their lethal temperature limits. We found that upper thermal fertility limits explain global distributions of species better than limits based on lethal temperatures. This suggests that limits to reproduction, rather than limits to survival, can underpin species distributions in nature. Given that high temperatures impair male fertility across a broad range of animals and plants, many species may be at increased risk of extinction due to inability to reproduce at high temperatures.

One Sentence Summary:

Species' distributions and response to climate change are strongly affected by the temperature at which they lose fertility.

Main Text:

We urgently need to understand how rises in temperature will impact biodiversity(*1*). To do this

we must understand the physiological, behavioral and evolutionary factors that underpin

species' current thermal distributions(*2*, *3*). Laboratory-derived estimates of the highest

temperatures at which an organism can function provide measures of species' thermal

tolerances(*4*). These measures of upper thermal limits have improved the accuracy of

functional species distribution models(*5*) which can be extrapolated to climate change

scenarios, allowing ecologists to forecast future species distributions(*6*). Accurate predictions of

species' distributions are invaluable for prioritizing conservation efforts(*7*) and predicting the

invasion of disease vectors(*8*).

Upper thermal tolerance limits are usually based on the temperatures that cause loss of

coordinated movement, coma, respiratory failure, or death: the species' critical thermal limit.

Despite these being measured in artificial laboratory conditions, critical limits correlate

reasonably well with species' distributions(*4*) and have been used to estimate species' capacity

to tolerate temperature increases across their current distribution range; their 'thermal safety

margins'(*3*). However, persistence of populations is not determined solely by survival, but also

by reproduction. There is evidence that sub-lethal temperatures cause losses in fertility in

plants(*9*), insects(*10-12*), fish(*13*), aquatic invertebrates(*14*), birds(*15*) and mammals, including

humans(*16*). These effects include direct impacts on physiological processes (*10*, *15*, *17*) and

indirect influences via changes in behavior and phenology(*18*). Previously, we proposed that

temperatures at which fertility is lost, the thermal fertility limits (TFLs) (*18*), may be a critical

but understudied part of species' true upper thermal limits. If TFLs are lower than critical limits,

then many organisms will be more vulnerable to climate change than currently thought. If TFLs

and critical limits correlate poorly then we may misidentify which species are most at risk from

rising temperatures. Critically, we need to know whether TFLs measured in laboratory

conditions can predict the distribution of natural populations better than measures of critical

limits.

Here, we recorded three measures of upper thermal limits in adult males from 43 species of

Drosophila fruit flies. First, we exposed flies to a 4-hour static heat stress at a range of

temperatures from benign through to lethal(*19*). From these data we estimated the

temperature at which 80% of living males are sterilized (TFL80) and the temperature that is

lethal to 80% of individuals (LT80). Fertility was assayed at two time points: (i) over 1-6 days

post-heat, to capture any immediate sterilizing effect of heat, and (ii) 7-days after heat-stress to

capture any recovery of fertility or delayed sterility. These data allow us to compare fertility and

survival thresholds under identical conditions. In a separate experiment, we measured the

temperature at which males lose coordinated motor function under ramping temperature

conditions (CTmax). This measure is widely used across animal systems to predict species'

sensitivity to climate change (*3, 4*).

We found that 11 of 43 species experience an 80% loss in fertility at cooler-than-lethal

temperatures immediately following heat-stress (Fig S1). Interestingly, rather than seeing a

recovery of fertility over time, the impact of high temperatures on fertility was more

pronounced 7-days post heat stress (Fig 1). Using this delayed measure of fertility, nearly half of

species (19/43) showed significant fertility loss. The difference between lethal and fertility

limits ranged from 0°C to 4.3°C (mean = 1.15 ± 0.22°C), and LT80 and TFL80 predict dramatically

different relative ranking of species' robustness to high temperature (Fig S2).

Figure 1: Incorporating fertility loss into estimates of upper thermal limits dramatically

changes the ranking of species' thermal tolerance. LT80 (black circles) and TFL80 (pink

circles). 95% CI are shown as error bars for both measures. Fertility loss measured seven

days post heat stress to account for delayed sterility effects.

These data show that fertility loss at sub-lethal temperatures is common across *Drosophila.* This

suggests that ignoring TFLs has led to the overestimation of thermal tolerance for many species.

This may be particularly problematic for efforts to preserve rare and endangered species or

control emerging pests and disease vectors. For example, we found that the crop pest species

Zaprionus indianus (African fig fly) has a relatively high upper lethal limit (LT80 = 37.9°C) but a

much cooler upper fertility limit (TFL80 = 34.8°C). This switches the species' thermal hardiness

rank from rank 5th to 17th of species here (Fig S2). This pest is currently expanding throughout

the New World, and this disparity could have marked consequences for forecasting its putative

range and hence where to concentrate control efforts(*20*).

Our work demonstrates that TFLs can be substantially lower than lethal temperatures when

measured under the same conditions in the laboratory. However, the key question is whether

TFLs actually shape organisms' distributions in nature. To test this, we used existing data on the

distribution of each *Drosophila* species and integrated this with global climate data for each

location to estimate the maximum temperatures species are likely to encounter in nature(*19*).

Previous work using 95 *Drosophila* species found that the mean maximum summer

temperature measured in this way correlates with CTmax for species from dry (<1000ml annual

rainfall) but not from wet (>1000ml annual rainfall) environments(*4*). We first verified this

finding in our smaller dataset of 43 species; CTmax significantly interacts with annual

precipitation to predict mean maximum environmental temperature (PGLS: F = $6.02_{1,39}$, P = 0.018, adjR^2 = 0.37). We then tested whether either LT80 or TFL80 measured 7-days after heat stress are better predictors of species' thermal habitats than CTmax (Table S1). Across all species, we found that a model with TFL80 as the sole physiological predictor had the strongest correlation with mean maximum environmental temperature with no significant interaction with precipitation (PGLS: interaction term: F = $0.531_{1,39}$ P = 0.471, TFL80: F = $39.3_{1,39}$ P < 0.001, adjR^2 = 0.55). LT80 also did not significantly interact with precipitation to predict environmental temperature but explained considerably less variation than TFL80 (PGLS: interaction term: $F_{1,39}$ = 0.99, P = 0.324, LT80: $F_{1,39}$ = 0.99, P = 0.324, adjR^2 = 0.31). Hence, TFL results in a 48% and 77% improvement in the accuracy (R^2) of species distribution predictions compared to CTmax and LT80 respectively. This dramatic improvement is largely due to TFL80 more accurately predicting the thermal range for species from high humidity environments (Table S2 & Fig S3). We find qualitatively similar results if both lethal and sterilizing limits are included as predictors in the same model (Table S3). The power of TFL80 to predict the distribution of *Drosophila* in nature strongly suggests that fertility losses due to high temperature are an important, but previously ignored, determinant of where species can persist.

Predicting species' vulnerability to current and future temperature extremes is critical to protect biodiversity[1]. "Thermal safety margins" measure species' vulnerability to climate change by taking the difference between an organism's physiological thermal limit and the maximum air temperature it is likely to experience[21]. Although safety margins often use complex measures of environmental temperature to improve accuracy (e.g. microhabitat

availability, thermoregulation behavior of adults(*3*)), they remain reliant on relatively simple

measures of thermal vulnerability. We find that TFL80 measured seven days after heat-stress

produces significantly smaller safety margins than either LT80 or CTmax (LMER: χ^2 = 40.36, df =

2, *P* < 0.001, Fig S4). For populations that already live near the upper edge of their thermal

range, TFL-based safety margins are reduced from 2.59 ± 3.08°C under assumptions of LT80, to

1.43 ± 2.80°C. This predicts that 20 of the 43 species studied here are found in environments in

which air temperatures exceed safety limits during the hottest part of the year. We illustrate

the implications of this with case-study distribution models of *Drosophila flavomontana;* safety

margins based on TFL80 predicts a 17.9% reduction in habitable landscape compared to an

identical LT80-based model under current climate conditions (Fig 2A). This disparity between

predictions based on sterility and lethality grew to 48.0% by the year 2070 under moderately

optimistic future climate forecasts (RCP4.5, Fig 2B), and to 58.9% under pessimistic climate

change scenarios (RCP8.5, Fig 2C). TFL-based models also predict that by 2070 the available

habitat for *D. flavomontana* will have reduced by 42.3% and 62.9% under RCP4.5 and RCP8.5

respectively.

Figure 2: Potential habitat range of *Drosophila flavomontana* (LT80 = 35.8°C, TFL80 =

31.4°C) under A) current and B & C) possible future climate scenarios (B = RCP4.5

'moderately optimistic', C = RCP8.5 'pessimistic', predicted for 2060 - 2080). Colored areas

in each panel represent suitable habitat range predicted by the model that excludes

maximum temperature (Fig S5). Red areas show regions where maximum summer

temperatures exceed LT80. Orange areas show regions where maximum summertime

temperatures exceed TFL80. Blue regions are habitable for *D. flavomontana* all year.

If our data for *Drosophila* can be extrapolated to other organisms, it is likely that male fertility

losses at high temperatures are common and can occur at substantially lower temperatures

than the species upper lethal limit. The limited data on fertility at extreme temperatures

supports this, with losses in male fertility at high temperatures seen across a broad diversity of

organisms[18]. It is possible that behavioral thermoregulation will reduce the impact of high

temperatures on fertility in nature. *Drosophila* are able to behaviorally thermoregulate, moving

to leaf litter, shade, or higher altitudes, with many species able to survive high temperature

periods by aestivating as adults, eggs or pupae[22]. Despite this, we still find that the

distribution of species is predicted by thermal fertility limits. Even species that might be

predicted to have high thermal tolerance can show evidence of thermal fertility losses. For

instance, the zebra finch, a high temperature adapted desert- dwelling endothermic species

with naturally high body temperature and good thermoregulation, shows substantial damage

to sperm at temperatures it regularly experiences in nature[15].

Our work suggests that temperature-driven fertility losses may be a major threat to biodiversity

during climate change. We urgently need to understand the range of organisms likely to suffer

similar fertility losses in nature, and the traits that predict vulnerability. However, we currently

do not understand the physiology underlying variation in TFLs between species, nor the

selective forces that created this variation. Ultimately, we need to know whether evolution for

higher TFLs will allow species to adapt to a warming environment.

References and Notes:

174   1.   G. T. Pecl *et al.*, Biodiversity redistribution under climate change: Impacts on

175   ecosystems and human well-being. *Science*. **355**, eaai9214 (2017).

176   2.   I.-C. Chen, J. K. Hill, R. Ohlemüller, D. B. Roy, C. D. Thomas, Rapid Range Shifts of

177   Species Associated with High Levels of Climate Warming. *Science*. **333**, 1024–1026

178   (2011).

179   3.   J. M. Sunday *et al.*, Thermal-safety margins and the necessity of thermoregulatory

180   behavior across latitude and elevation. *Proceedings of the National Academy of*

181   *Sciences*. **111**, 5610–5615 (2014).

182   4.   V. Kellermann *et al.*, Upper thermal limits of Drosophila are linked to species

183   distributions and strongly constrained phylogenetically. *Proc. Natl. Acad. Sci. U.S.A.*

184   **109**, 16228–16233 (2012).

185   5.   M. Kearney, W. Porter, Mechanistic niche modelling: combining physiological and

186   spatial data to predict species' ranges. *Ecol Lett*. **12**, 334–350 (2009).

187   6.   J. Elith, J. R. Leathwick, Species Distribution Models: Ecological Explanation and

188   Prediction Across Space and Time. *Annu. Rev. Ecol. Evol. Syst.* **40**, 677–697 (2009).

189   7.   A. J. Nowakowski *et al.*, Thermal biology mediates responses of amphibians and

190   reptiles to habitat modification. *Ecol Lett*. **21**, 345–355 (2018).

191   8.   S. Metelmann *et al.*, The UK's suitability for Aedes albopictus in current and future

192   climates. *Journal of The Royal Society Interface*. **16**, 20180761 (2019).

193   9.   T. L. Sage *et al.*, The effect of high temperature stress on male and female

194   reproduction in plants. *Field Crops Research*. **182**, 30–42 (2015).

195   10.   K. Sales *et al.*, Experimental heatwaves compromise sperm function and cause

196   transgenerational damage in a model insect. *Nat Commun*. **9**, 4771 (2018).

197   11.   D. Porcelli, K. J. Gaston, R. K. Butlin, R. R. Snook, Local adaptation of reproductive

198   performance during thermal stress. *J Evol Biol*. **30**, 422–429 (2016).

199   12.   A. D. Saxon, E. K. O'Brien, J. R. Bridle, Temperature fluctuations during

200   development reduce male fitness and may limit adaptive potential in tropical

201   rainforest Drosophila. *J Evol Biol*. **31**, 405–415 (2018).

202   13.   R. D. Breckels, B. D. Neff, The effects of elevated temperature on the sexual traits,

203   immunology and survivorship of a tropical ectotherm. *Journal of Experimental*

204   *Biology*. **216**, 2658–2664 (2013).

205   14.   C. W. Paxton, M. V. B. Baria, V. M. Weis, S. Harii, Effect of elevated temperature on

206   fecundity and reproductive timing in the coral Acropora digitifera. *Zygote*. **24**, 511–

207   516 (2016).

208   15.   L. L. Hurley, C. S. McDiarmid, C. R. Friesen, S. C. Griffith, M. Rowe, Experimental

209   heatwaves negatively impact sperm quality in the zebra finch. *Proc. R. Soc. B.* **285**,

210   20172547 (2018).

211   16.   L. Yogev *et al.*, Seasonal variations in pre- and post-thaw donor sperm quality. *Hum*

212   *Repro*. **19**, 880–885 (2004).

213   17.   C. Rohmer, J. R. David, B. Moreteau, D. Joly, Heat induced male sterility in

214   Drosophila melanogaster: Adaptive genetic variations among geographic

215   populations and role of the Y chromosome. *Journal of Experimental Biology*. **207**,

216   2735–2743 (2004).

217   18.   B. S. Walsh *et al.*, The Impact of Climate Change on Fertility. *Trends Ecol. Evol*. **34**,

218   249–259 (2019).

219   19.   Materials and Methods

220   20.   K. van der Linde, G. J. Steck, K. Hibbard, J. B. Florida, First records of Zaprionus

221   indianus (Diptera: Drosophilidae), a pest species on commercial fruits from Panama

222   and the United States of America. *BioOne* **89,** 402 - 404 (2006).

223   21.   C. A. Deutsch *et al.*, Impacts of climate warming on terrestrial ectotherms across

224   latitude. *Proceedings of the National Academy of Sciences*. **105**, 6668–6672 (2008).

225   22.   M. E. Dillon, G. Wang, P. A. Garrity, R. B. Huey, Thermal preference in Drosophila.

226   *Journal of Thermal Biology*. **34**, 109–119 (2009).

227   23.   J. Overgaard, M. R. Kearney, A. A. Hoffmann, Sensitivity to thermal extremes in

228   Australian Drosophila implies similar impacts of climate change on the distribution

229   of widespread and tropical species. *Global Change Biology*. **20**, 1738–1750 (2014).

230   24.   J. S. Terblanche, J. A. Deere, S. Clusella Trullas, C. Janion, S. L. Chown, Critical

231   thermal limits depend on methodological context. *Proc. R. Soc. B.* **274**, 2935–2942

232   (2007).

233   25.   P. T. Rohner *et al.*, Interrelations of global macroecological patterns in wing and

234   thorax size, sexual size dimorphism, and range size of the Drosophilidae.

235   *Ecography*. **41**, 1707–1717 (2018).

236   26.   S. E. Fick, R. J. Hijmans, WorldClim 2: new 1-km spatial resolution climate surfaces

237   for global land areas. *International Journal of Climatology*. **37**, 4302–4315 (2017).

238   27.   S. J. Phillips, R. P. Anderson, R. E. Schapire, Maximum entropy modeling of species

239   geographic distributions. *Ecological Modelling*. **190**, 231–259 (2006).

240   28.   N. Ramankutty, A. T. Evan, C. Monfreda, J. A. Foley, Geographic distribution of

241   global agricultural lands in the year 2000, *Global Biogeochemical Cycles*, **22**

242   doi:10.1029/2007GB002952, (2008).

243   29.   C. Amante, B. W. Eakins, ETOPO1 1 Arc-Minute Global Relief Model: Procedures,

244   Data Source and Analysis. *National Geographic Data Center* (2009).

30. C. Liu, M. White, G. Newell, Selecting thresholds for the prediction of species

occurrence with presence-only data. *Journal of Biogeography*. **40**, 778–789 (2013).

31. B. Thrasher, E. P. Maurer, C. McKellar, P. B. Duffy, Technical Note: Bias correcting

climate model simulated daily temperature extremes with quantile mapping.

Hydrol. Earth Syst. Sci. **16**, 3309–3314 (2012).

32. A. E. Jones *et al.*, Bluetongue risk under future climates. *Nature Climate Change*. **9**,

153–157 (2019).

33. S. Pitnick, T. A. Markow, G. S. Spicer, Delayed male maturity is a cost of producing

large sperm in Drosophila. *Proceedings of the National Academy of Sciences*. **92**,

10614–10618 (1995).

34. K. E. Roberts, J. D. Hadfield, M. D. Sharma, B. Longdon, Changes in temperature

alter the potential outcomes of virus host shifts. *PLoS pathogens*. **14**, e1007185

(2018).

35. M. D. Garlovsky, R. R. Snook, Persistent postmating, prezygotic reproductive

isolation between populations. *Ecol Evol*. **8**, 9062–9073 (2018).

36. A. A. Hoffmann, S. L. Chown, S. Clusella Trullas, Upper thermal limits in terrestrial

ectotherms: how constrained are they? *Funct Ecol*. **27**, 934–949 (2013).

Acknowledgements: Natasha Mannion and Dr Angela Sims for assistance with experiments, Dr

Ben Longdon and Dr Katherine Roberts for flies, Rowan Connell for designing 3D-printed

equipment. **Funding:** NERC grant NE/P002692/1 to TP, AB, AH & RS. SNF P300PA_177830 to

AM. NIHR HPRU EZI to SM. **Author Contributions:** Conception: TP, AB, RS, AH, SP. Methodology

& data collection: SP, BW, NW. Data curation: SP & SM. Analysis: SP, AM & SM. Original Draft:

SP, TP, AB, RS & AH. Review & Editing: All authors. **Data and materials availability:** All data and

analysis R code will be deposited on Dryad upon acceptance of this manuscript.

List of Supplementary Materials:

Materials and Methods:

<u>Animal maintenance:</u>

All species were kept in temperature-controlled rooms at their noted "rearing temperature"

(see table S4) selected based on observations of when the laboratory populations were most

stable. All stocks were kept at 12:12 L:D and ambient humidity. Stocks were maintained at

moderate density (20 – 50 flies per 23ml vial or 50 – 100 flies per 300ml bottle culture).

Cultures were tipped to fresh food every 7 days and a new generation was made every 4 – 6

weeks depending on the speed of a species development. Species were reared on one of four

food types; A = ASG (10g agar, 85g Sucrose, 20g yeast extract, 60g maize, 1000ml H_2O, 25ml

10% Nipagin), B = Banana (10g agar, 30g Yeast extract, 150g pulped banana, 50g Molasses, 30g

Malt Extract, 25ml 10% Nipagin, 1000ml H_2O), P = Propionic (10g Agar, 20g Yeast extract, 70g

cornmeal, 10g soya flour, 80g Malt Extract, 22g Molasses, 14ml 10% nipagin, 6ml Propionic

acid, 1000ml H_2O), M = Malt (10g agar, 20g Yeast, 60g Semolina, 80g Malt, 14ml 10% Nipagin,

5ml Propionic acid, 1000ml H_2O). To standardize across species, we only included males in our

TFL assay that were sexually mature. This inevitably led to variation in the age of males (days

since eclosion) between species based on published age-to-maturity (e.g. *34*) and our own

observations (Table S4 for details).

<u>Species verification:</u>

Most *Drosophila* species were gained from stock centers, other research groups, or through

field collections. Where species were not distinguishable through evident morphological traits,

we verified position within the phylogeny by nucleotide sequence identity.

We extracted DNA from 2-3 adult male flies with DNeasy kits (Qiagen) following the

manufacturer's invertebrate protocol. We PCR amplified a portion of the mitochondrial

universal barcode gene cytochrome oxidase subunit 1 using the primers C1-J-1718 (5' –

GGAGGATTTGGAAATTGATTAGT – 3') and C1-N-2191 (5' – CCCGGTAAAATTAAAATATAAACTTC –

3') using HotStart Taq (Promega) with (5-minute initial heating, 30 cycles at 95°C for 30s, 56 for

30s, and 72°C for 30, with an final elongation step of 72°C for 120s). PCR products were

visualized by SYBRSafe-stained gel electrophoresis and cleaned up using Exonuclease I and

Shrimp Alkaline Phosphatase incubation per supplier protocol (BioLine). We then used BigDye

based sequence reactions with both forward and reverse primers, followed by NaOH & ethanol

precipitation clean-up and precipitation before sequences were analyzed with an ABi 3500XL

Genetic Analyzer. Forward and reverse sequences for each species were aligned to derive a

consensus sequence. For species with a putative ID based on stock center or collaborator

expertise we assessed the clustering of these sequences with publicly available CO1 sequences

from the same species available on the BOLD database (boldsystems.org). Clustering was

performed with the MAFFT server tool, and clustering via a neighbor-joining tree was

visualized, all based on default parameters. For species that were unknown (two *repleta*-like

species collected from the field in Madeira and Morocco), we performed the same process as

above, but included multiple sequences from multiple species in the *repleta* species group –

initial BLAST of these sequences on the NCBI database had given an approximate idea of the

species ID.

<u>Measuring upper thermal limits</u>

We assayed three metrics of upper thermal limits in males from 43 species of *Drosophila*: Lethal

Temperature 80% (LT80), Thermal Fertility Limit 80% (TFL80) and Maximum Critical

Temperature (CTmax). We chose to test for thermal fertility limits in only males because of the

substantial evidence that male fertility is particularly vulnerable to high temperatures(*18*).

LT80 & TFL80: Newly eclosed virgin adult males were collected over a 48-hour period into

Drosophila vials and allowed to sexually mature for 7 – 21 days depending on species (see Table

S4). Males were kept in groups of maximum 10 individuals to minimize intrasexual aggression.

Once mature, males were transferred to fresh 23ml fly vials with standard maize-sugar-yeast

agar "ASG" medium' food and allocated to 3D-printed floating plastic racks in pre-heated

waterbaths (Grant TXF200) set to a range of temperatures (see Table S4 for temperature

ranges). ASG food was used during heating for all species because preliminary work had shown

that survival under heat stress was influenced by food type. Floating racks were weighted with

ball-bearings at each corner to keep vials containing flies submerged so that the waterline was

above the cotton-wool stopper. This ensured that flies were exposed to homogenous

temperature conditions inside the vials. In each run of the experiment we monitored the

temperature inside a fresh ASG vial without flies with a k-type thermocouple attached to a data

logger (Pico Technology TC-08). Males were heated for 4 hours between ~10am - ~2pm and

then returned to temperature-controlled rooms set to the species' benign temperature. We

scored the survival of males the next morning to account for any immediate recovery or

delayed lethality of the temperature treatment. We scored flies as dead if they were

completely immobile or had become stuck on their backs in the fly food – the latter would

indicate that the flies were completely incapacitated by the treatment temperature which

would likely be lethal in the wild.

Surviving males were then individually aspirated into separate, fresh vials containing their

species' designated food type and 3-4 sexually mature virgin females. Males remained in these

vials at their benign temperature and allowed to mate freely for 6 days. This allows us to score

any immediate lasting effect of temperature on fertility. After 6 days males were transferred to

a second fresh vial with 1-2 more virgin females and allowed to mate for 24 hours. This allowed

us to test for recovery of fertility or any delayed sterilizing effect of temperature. Females in

both sets of vials were left to lay eggs for 3-5 days (depending on fecundity of the species) after

the male had been removed. Fertility was scored qualitatively as the presence or absence of

any offspring: vials were routinely scored for the presence of larval flies by checking for the

larvae themselves or their distinctive track marks in the food. Juveniles flies were allowed to

develop through to adult eclosion before being frozen and the number of adult offspring

counted.

In total, 14742 males were exposed to heat treatment, of which 10925 survived and 9064 went

on to be tested for heat-induced sterility. The average sample size at each temperature per

species was 36.5 ± 0.5sem, ranging from 10 – 88 individuals.

CTmax: The measures of upper tolerance described above are performed under static heat-

stress conditions – this is necessary for estimating TFLs because the phenotype "fertility" can

only be observed after the heat-stress treatment, so matching a male's fertility to a stress

temperature requires that he only experiences a single stressful environment. LT80 is therefore

a directly comparable measure to TFL80. However, several studies in *Drosophila* and other

animals have used estimates of temperature tolerance measured under gradually ramping heat

conditions to estimate species response to climate change (*4*, *7*, *23*, *24*). Because of this, and

because previous work has shown that estimate of upper critical thermal limits can depend on

methodological context(*25*), we also measured upper critical limits of our 43 *Drosophila* species

under ramping heat conditions – critical thermal maxima (CTmax). This allows us to compare

how static and dynamic measures of critical heat tolerance predict species' distributions. It also

allows us to compare upper TFLs to two widely used measures of upper critical limits.

We measure CTmax as the mean temperature at which male flies lose coordinated movement

and are unable to right themselves. Individual males were collected as virgin adults from fly

stocks and allowed to sexually mature (see Table S4). Flies were then anesthetized with CO_2

and isolated into individual glass sample tubes sealed with a rubber stopper. Sample tubes

were left at ambient room temperature for at least 30 minutes to allow males to fully recover

from CO_2 anesthetization – preliminary trials found no difference in CTmax between

anesthetized and non-anesthetized male *D. melanogaster.* Sample tubes then were attached to

a custom-built plastic rack and held at a 45° angle with elastic banding to facilitate the

experimenter identifying when flies collapsed. The rack and vials were submerged in a glass

aquarium held at the rearing temperature for the given species by a water circulator (Grant

TXF200) (18, 23 or 25°C – Table S4). The temperature was then increased at a rate of 0.1°C per

minute until all flies were incapacitated. The temperature at which flies collapsed for 30

seconds and did not right themselves after being physically disturbed by the experimenter

(tapping the vial) was scored. The temperature of the waterbath was continuously recorded

throughout the assay with a data logger (Elitech RC-61). The mean sample size for each species

for CTmax assays was 19.53 ± 1.77.

<u>Repeatability</u>

Due to the logistical constraints in the number of flies that could be processed at a time and the

differences in temperature range required to stress each species it was impossible to assay

upper limits across all 43 species simultaneously. Further, inter-species variation in

development time prohibited us from synchronizing all 43 species to run the assay in a

completely randomized block design. Instead, we assayed each species separately to

independently estimate CTmax, TFL80 and LT80. These assays were run between June 2018 –

November 2019.

To test the repeatability of our measure of Lethal Temperature (LT80) we ran a single control

block in February 2020 with 22 species that we could obtain enough virgin males that would

reach sexual maturity in synchrony. We simultaneously exposed these species to a range of 6

temperatures each that covered their estimated LT80 identified when we had previously run

them individually. We then tested the strength of the correlation between our original LT80

estimates and those given by this replicate block and determined if any given species showed

significantly different estimates based on whether the confidence intervals of the two runs

overlapped. Sample sizes for this control run ranged from 3 – 10 males per temperature per

species, and involved 1152 male flies in total.

As a test of repeatability for our TFL80 estimates we ran two independent TFL assays on

Drosophila virilis. This species was selected because it had demonstrated a clear difference

between TFL80 and LT80 limits. The two repeat blocks were run 6 months apart and data were

collected by two separate researchers. We compare the estimated TFL80 from these

independent runs and consider differences to be significant if the 95% confidence intervals of

our estimates do not overlap.

Repeatability of CTmax was estimated by randomly allocating species from the same rearing

temperatures into blocks of 3 - 5 species and measuring their mean CTmax as described above.

We compared these mean Ctmax values with those obtained when species were run

individually when we collected the CTmax data used in analyses.

<u>Point estimates of TFL80 & LT80</u>

We chose 80% thresholds as our physiological limits for both LT and TFL because i) we wanted

to compare viability and fertility measures on comparable scales, and ii) a loss of 80% of males

is likely to represent a substantial threat to population stability for most species.

We found that general linear models fit our qualitative (1/0 data) fertility and survival data

poorly, and overestimated 80% fertility loss thresholds considerably. Instead, we generated

point estimates of 80% thresholds with dose response models (`drm()`), implemented in the

`drc` package in R. We used 3-parameter versions of the log-logistic model which fixes the

lower limit of the predicted curve to 0 to reflect that we expect fertility and survival to

eventually reach 0 over an infinite range of temperatures. The models then estimate the point

at which we see an 80% reduction in survival or fertility relative to the upper limit present in

the data. In this way, these models allow for a given species to inherently have some degree of

sterility even at benign temperatures, and to estimate the temperature required to reduce this

by 80%. Percentage fertility was analyzed for surviving flies only, i.e males killed by the heat

treatment were not considered to be sterile.

In analyses in which we predict geographic range by physiological limits we use TFL80 estimates

that are 'significantly' lower than the species' LT80 estimate – for the remaining species TFL80

point estimates are defaulted to the LT80 to be conservative. We considered the difference

between TFL80 and LT80 to be 'significant' if the 95% confidence intervals of the two point-

estimates do not overlap.

<u>Correlations between upper thermal limits.</u>

We used phylogenetically controlled linear models to test how much variation in TFL80 is

explained by variation in LT80. These measures are taken from the same assays and are often

very similar, so this correlation gives a quantitative assessment of the mismatch between these

two thermal tolerance traits.

CTmax and LT80 inherently use different heating regimes to capture upper critical limits. Most

notably, CTmax includes an element of temperature ramping, and so may allow species to

plastically harden to high temperature stress. To test how well these two measures capture the

same component of thermal interspecific variation, and are therefore proxies for each other,

we used phylogenetically controlled linear models to explore the correlation between TFL80

and CTmax. Whilst we did not expect these methods to produce identical point estimates of

lethal temperatures for a given species, we did hypothesize that these correlations will have a

high R^2 and a slope estimate close to 1.

If there is a universal upper limit to fertility across *Drosophila* then we would expect the

difference between TFL and critical temperatures to be greater in species with higher

temperature LT80 and CTmax. To test this, we used phylogenetically controlled linear models to

test for significant correlations between ΔLimits (the difference between TFL80 and LT80) and

both LT80 and CTmax.

Accounting for phylogeny:

All of the species we used in this study are present in a phylogeny published by Patrik Röhner

and colleagues(*26*): We use this phylogenetic tree as the basis of all our phylogenetically

controlled analyses.

We tested for phylogenetic signal in all three upper thermal limits (LT80, TFL80 and CTmax). We

used `pgls()` in `caper` to estimate if Pagel's lambda was significantly different from 0 (no

phylogenetic signal) or 1 (complete Brownian motion). The maximum likelihood method was

used to estimate lambda.

When analyzing correlations amoung traits, and between traits and geographic distributions we

corrected for phylogenetic signal in model residuals with `pgls()` in the R package `caper`. Even

where phylogenetic signal was estimated to be not significantly different to 0, we retained the

correction in the model because these models give qualitatively identical results to uncorrected

linear models using `lm()`.

<u>Environmental variables:</u>

Geographic distribution data for each species were obtained manually from TaxoDros.uzh.ch as

coordinates and location names. These were systematically cleaned by deleting any entry set to

zero decimal places in both coordinates. Then, they were globally cleaned by removing

locations with vague names such as entire countries or US states. Plots of each species location

on a map were then visually inspected, and outliers were investigated and removed or

corrected when appropriate.

The cleaned dataset of geolocations was then integrated with rasterized bioclimate data from

the WordlClim V2.0 database (*27*). Our intention was not to fully reassess multiple climate

variables as predictors for *Drosophila* distributions, because this has received extensive

attention in the existing literature(*4, 24*). Instead, we tested if TFLs correlate with variables

previously identified as important for predicting physiological limits and distribution in

Drosophila; the maximum temperature reached in the hottest summer month (WorldClim Bio5)

and annual precipitation (WorldClim Bio12). Means of each climate variable for each species

were calculated across all global locations. We also calculated the mean of each variable

independently for locations in the North and South hemispheres for species whose range

straddles the equator.

Comparing upper thermal limits as predictors of species' thermal environment

We used three distinct modelling approaches to test which measure of upper thermal limits

best predicts species' thermal range:

Firstly, we fitted separate phylogenetically controlled linear models with either TFL80, LT80 or

CTmax as independent predictors, and species' Tmax (WorldClim Bio5) as a response variable.

In all three models we included mean annual precipitation (WorldClim Bio12 or "Pann") as an

interacting covariate, because previous work has found that the power of CTmax to predict

species' thermal environments degrades in more humid habitats(*4*). In these models all

predictors are centered and scaled. We then used AIC-based model selection to reduce models

to minimum-adequate models, and compared the slope estimate, model structure and R^2 of the

competing models.

Secondly, we directly replicated the method of (*4*) in which the authors split species into those

that experience greater-than and less-than 1000mm of rainfall ("wet" and "dry" hereafter). The

authors of (*4*) found that CTmax correlated well for "dry" species but not for "wet" species. We

ran phylogenetically corrected linear models with Tmax as a response and either CTmax, LT80

or TFL80 and predictors for both 'dry' and 'wet' species. We qualitatively compared the R^2 of

these models as indicators of model fit.

Finally, we included both CTmax and TFL80 and their interaction with `Pann` into a single

phylogenetically controlled model of Tmax. We did not include LT80 in this analysis because *a*

priori Variance Inflation Factor tests (VIF) on non-phylogenetically controlled versions of this

model indicated that LT80 correlated very strongly with both other measures. We reduced the

maximal model through AIC-based stepwise simplification starting with higher order interaction

terms to reach a minimum adequate model.

<u>Calculating and modelling thermal safety margins</u>

Thermal safety margins were calculated as the difference between a species' physiological

upper thermal limit and the mean maximum summertime temperature they are likely to

experience across all of their known global distribution locations. As per(*4*), we calculated this

in two ways: firstly by taking the mean Tmax for each species, to estimate safety margins for

the 'central' populations of a species (termed "Central Safety Margin"). Secondly by taking the

mean Tmax + 1 SD to account for populations of species that live at the upper thermal range of

species' distribution (the "distribution safety margin). Safety margins were calculated

separately using either CTmax, TFL80 or LT80 as the physiological limit.

We also treated populations in the North and Southern hemispheres as equivalent to separate

species, because previous work has found minor differences in safety margins in the North and

South hemisphere, likely because of biases in sampling efforts(*3*).

Safety margins were modelled as phylogenetically controlled linear mixed models with

'hemisphere' (N or S) and 'physiological limit' (CTmax, TFL80 or LT80) as linear predictors,

absolute latitude (degrees^2) as a quadratic predictor, and species identity as a random

intercept.

<u>Predicting current and putative future distributions of *D. flavomontana*</u>

As a case-study demonstration of the importance of TFLs, we use MaxEnt v3.4.1 (*28*) to model

potential distributions of *D. flavomontana*. Occurrence data were obtained from

TaxoDros.uzh.ch and points within 25km of each other were clustered together to avoid

oversampling (*N* = 24). First, we chose climatic and other variables to predict a potential

distribution of *D. flavomontana* that is independent of the upper thermal limit. After excluding

variables that strongly correlated or had little to no impact on the model results (<3%

contribution), we used four of the tested 15 variables: mean winter temperature (WC Bio 11,

from the WorldClim2.0 database, precipitation seasonality (WC Bio 15), an agricultural index

(*29*), and elevation (*30*). We then ran 10 cross-validations with maximum iterations of 5000,

using 10000 random pseudo-absences. To distinguish suitable from unsuitable regions, we used

the threshold that maximized Youden's index (maximum training sensitivity plus specificity)

(*31*). The MaxEnt mean output grid is shown in Figure S5.

We then constrained the suitable area by the upper physiological thermal limits measured

above: i) the lethal temperature measured as LT80 (35.8°C), and ii) the TFL80 (31.4°C). We use

the maximum weekly temperature of the warmest month as reference (WC Bio5). Using WC Bio

05 directly as a predictor variable for *D. flavomontana* had close to no effect on the distribution

(0.1% contribution).

For future predictions, we chose two carbon emission scenarios: RCP4.5 (moderately optimistic)

and RCP8.5 (pessimistic). We use precipitation and temperature predictions of the NorESM1-M

model from the NEX-NASA GDDP data set (*32*), as this model produces a median trend in terms

of temperature increase(*33*). The agricultural index and elevation had to be used as present day

data. Predicting and constraining the possible future distribution followed the same procedure

as for the current distribution.

Figure S1: TFL80 and LT80 of *Drosophila* immediately after heat stress. Several species of

Drosophila lose 80% fertility at cooler-than-lethal temperatures immediately following

heat-shock. LT80 (black circles) and TFL80 (pink circles). Errors for both measures are 95%

confidence intervals generated from dose response model estimates. Fertility loss

measured as ability to sire any offspring between 1 – 6 days post heat-stress.

Fig S2: Relative ranking of species thermal robustness based on LT80 and TFL80. Most to least

heat tolerant species ranked from top to bottom. Pink bars indicate species with significantly

lower TFL than LT. Data based on TFL80 recorded 7-days post-heat shock.

Fig S3. Correlations of maximum environmental temperature with CTmax, LT80 and TFL80.

Upper thermal limits to fertility better predicts species' natural thermal habitat than either

measure of critical limits; LT80 includes the same heat stress regime as TFL80, CTmax is widely

used. To aid visualization here, data are split into 'wet' (>1000ml rainfall year^{-1}) and 'dry' (< 1000

ml rainfall year^{-1}) habitats as in(*4*). Model estimates from fitted lines here given in Table S2.

Measures based solely on upper critical limits show no significant correlation for species that live

in naturally wet habitat. This recapitulates the findings of Kellermann *et al*(*4*).

Fig S4: Thermal safety margins. Central (top) and Distribution (bottom) safety margins for species calculated using the physiological limit minus the mean maximum temperature across all recorded locations for each species. For distribution margins Tmax + 1sd is used to calculate safety margins, which has been argued captures populations at the boundary of a species' thermal range (*4*). Grey points and fit are safety margins calculated using knockdown CTmax

values. Black points and lines are safety margins based on LT80, and pink points are margins

based on TFL80. Fitted lines are predictions from models in which the limit type (TFL, LT or

CTmax) and latitude^2 are fixed effects and species identity is a random effect to account for

repeated measures across species. Models and fits for North and South populations are

calculated independently. Based on CTmax we predict that safety margins for central populations

are 9.06 ± 3.24°C and populations at the thermal extremes have margins of 5.23 ± 3.24°C. This

suggests that 3 species have populations at the edge of their range that are currently exposed to

lethal temperatures in the hottest part of the year. LT80 is a more severe heat shock than CTmax

and so inherently predicts smaller margins; 6.36 ± 2.92°C for central populations and 2.59 ±

3.08°C for populations at thermal extremes of species' ranges. This physiological measure

predicts that 11 species have populations at the edge of their range that are currently exposed to

lethal temperatures in the hottest part of the year. Including fertility data into LT80 to get TFL80

predicts that populations at the center of species' ranges have a 5.26 ± 2.55°C safety margin,

whilst populations at the edge of species' thermal ranges have a 1.43 ± 2.8°C safety margin. TFL-

based margins predict that 20 species have populations at the edge of their thermal range that

are currently exposed to sterilizing temperatures in the hottest part of the year. Safety margins

calculated with TFL80 are significantly smaller than those calculated from LT80 (Central safety

margins: lmer: F_1 = 28.04, P < 0.001, estimate = -1.13, distribution safety margins; lmer: F_1 =

17.33, P < 0.001, estimate = -1.13). We did not statistically compare safety margins calculated

from CTmax with those calculated from TFL80 and LT80 because CTmax inherently uses a

different method of heating individuals.

Fig S5: Base putative distribution model of *D. flavomontana*. Created with MaxEnt v3.4 with a

10-fold cross-validation run. The average test AUC for the replicate runs is 0.94, and the standard

deviation is 0.058. We used the maximum training sensitivity plus specificity threshold to

distinguish between suitable and unsuitable areas.

Table S1: Summaries of model that fit maximum environmental temperature (Tmax) to either CTmax, LT80 or TFL80 independently of each other. All models fit with upper physiological limits interacting with annual precipitation (Pann). R^2 given for maximal model. Terms retained after AIC model selection shown in italics. All models are phylogenetically controlled, with no significant phylogenetic signal found in any case (Pagel's lambda not significantly different to 0). All predictors centred and scaled.

Full Model	Terms	DF	F	P	adjR²
Tmax~CTmax*Pann	*CTmax*Pann*	*1,39*	*6.092*	*0.018*	*0.371*
	CTmax		9.351	0.004	
	Pann		12.282	0.001	
Tmax~LT80*Pann	LT80*Pann	1,39	0.995	0.325	0.306
	LT80		*14.311*	*<0.001*	
	Pann		*6.247*	*0.017*	
Tmax~TFL80*Pann	TFL80*Pann	1,39	0.531	0.471	0.544
	TFL80		*4.706*	*<0.001*	
	Pann		*3.551*	*0.001*	

Table S2: Summaries of model that fit either TFL80, LT80 or CTmax to Maximum Environmental temperature (Tmax). Species from 'wet' environments (Pann > 1000 ml) and 'dry' (Pann < 1000ml) modelled separately. All models are phylogenetically controlled, with no significant phylogenetic signal found in any case (Pagel's lambda not significantly different to 0).

Species group	Full Model	Terms	DF	F	P	$^{adj}R^2$
Dry	Tmax~CTmax	CTmax	1,18	19	<0.001	0.487
	Tmax~TFL80	TFL80	1,18	39.858	<0.001	0.672
	Tmax~LT80	LT80	1,18	13.47	0.002	0.396
Wet	Tmax~CTmax	CTmax	1,21	0.456	0.505	0.025
	Tmax~TFL80	TFL80	1,21	6.40	0.019	0.197
	Tmax~LT80	LT80	1,21	1.56	0.225	0.025

Table S3: Summary of model in which both TFL80 and CTmax are included as predictors of maximum environmental temperature (Tmax). R^2 shown for full model. Model was phylogenetically controlled, but no significant phylogenetic signal in residuals was detected. LT80 is excluded as a predictor based on high VIF scores indicating variance-inflating autocorrelation with both other predictors.

Full Model	Terms	DF	F	P	adjR2
Tmax~CTmax*Pann+TFL80*Pann	CTmax*Pann	1,37	0.164	0.688	0.531
	TFL80*Pann		0.329	0.569	
	CTmax		0.985	0.327	
	Pann		12.814	<0.001	
	TFL80		38.254	<0.001	

Table S4: Husbandry details for species used in this study. All species either *D = Drosophila* or *Z = Zaprionus*

Species	Rearing temp (ºC)	StrainID & source	Multiple lines	Food	Mean age at heat (days)	Temperature stress range (ºC)
D. affinis	18	DSSC Stock #: 14012-0141.14	N	M	7	29 – 36
D. albomicans	23	DSSC Stock #: 15112-1751.03	N	A	7	31 – 36
D. aldrichi	23	Wild caught single line, Texas 2019	N	A	7	33 – 40
D. americana	23	Single-line from (*35*)	N	M	7	33 – 38
D. ananassae	25	Single-line from (*35*)	N	A	7	30 – 38
D. arizonae	23	DSSC Stock #: 14012-0181.02	N	B	7	35 – 41
D. biarmipes	23	DSSC Stock #: 14023-0361.11	N	A	7	30 – 35
D. bifasciata	18	DSSC Stock #: 14012-0181.02	N	A	7	25 – 34
D. bipectinata	23	DSSC Stock #: 14024-0381.20	N	A	7	30 – 36
D. borealis	23	DSSC Stock #: 15010-0961.10	N	A	14	31 – 39
D. buzzatii	23	Single-line from (*35*)	N	M	7	32 – 39
D. eohydei	23	DSSC Stock #: 15085-1631.00	N	B	14	31 – 38
D. erecta	23	Single-line from (*35*)	N	A	7	29 – 39
D. flavomontana	23	Single-line from (*35*)	N	M	14	30 – 37
D. hydei	25	Mixed, long-term lab population at University of Liverpool	Y	B	14	29 – 37
D. lacicola	23	Single-line from (*35*)	N	M	7	31 – 37
D. littoralis	23	DSSC Stock #: 15010-1001.03	N	B	14	32 – 37
D. lummei	23	DSSC Stock #:	N	M	7	25 – 37

		15010-1011.07				
D. madeirensis	18	Mix of 3 lines, wild caught March 2019	Y	A	7	27 – 34
D. mayaguana	23	DSSC Stock #: 15081-1397.00	N	M	7	35 – 40
D. melanogaster	25	'Dahommey' lab population	Y	A	7	30 – 38
D. mercatorum	23	Wild-caught Madeira 2018, single line	N	A	7	31 – 40
D. micromelanica	23	DSSC Stock #: 15030-1151.01	N	A	7	30 – 38
D. melanica	23	DSSC Stock #: 15030-1141.03	N	A	7	32 – 38
D. mojavensis	23	Multi-line mixed population from (35)	Y	B	7	34 – 41
D. montana	18	Mixed population of IsoLines in (36)	Y	M*	21	31 – 37
D. nasuta	23	DSSC Stock #: 15112-1781.13	N	A	7	30 – 37
D. novamexicana	23	Multi-line mixed population from (35)	Y	B	7	28 – 38
D. obscura	23	Multi-line mixed population from (35)	Y	P	7	28 – 33
D. paranaensis	23	DSSC Stock #: 15082-1541.09	N	A	7	31 – 37
D. pseudoobscura	18	Mixed population if IsoLines in	Y	A*	7	29 – 35
D. repleta	23	DSSC Stock #: 15084-1611.01	N	A	7	30 – 38
D. santomea	23	Single-line rom (35)	N	A	7	29 – 37
D. sechellia	23	Multi-line mixed population from (35)	Y	P	7	29 – 37
D. simulans	25	Mixed of laboratory lines collected across Europe 2016 - 2018	Y	A	7	30 – 37
D. subobscura	18	Mixed population from Lines	Y	A	7	27 – 34

	23	Mixed populations of 3 IsoLines caught in Madeira (March 2018) and single lines each from the USA and Japan	Y	A	7	27 – 37
D. suzukii						
D. takahashii	23	Single-line from (*35*)	N	A	7	30 – 37
D. teissieri	23	Single-line from (*35*)	N	A	7	28 – 37
D. virilis	23	Multi-line mixed population from (*35*)	Y	P	7	26 – 38
D. yakuba	23	Multi-line mixed population from (*35*)	Y	A	7	3 – 38
Zaprionus indianus	23	Single line, wild caught 2019	N	B	7	33 – 39
Zaprionus tuberculatus	23	DSSC Stock #: 50001-0001.07	Y	B	7	33– 38

Supplementary text

Repeatability of 4-hour heating assay:

Fig ST1.1: Repeatability of LT80. Repeatability was high across the 22 *Drosophila* species

tested. Left panel: There was no significant difference in the estimate of LT80 between full

TFL runs and control repeats in any of these species (significance measured as non-

overlapping CIs of point estimates of LT80). Pink points = estimate from full TFL assay, gold

point = estimate from simultaneous multispecies LT80 assay. Right panel: The correlation

between the two independent measurements of LT80 across these species was strongly

positive and explained a high degree of variation in the data (coefficient = 1.06, R^2 = 0.96).

Figure ST1.2: Repeatability of TFL80 for *Drosophila virilis*. We fully independently repeated

the TFL80 assay for one species in which we found a significant difference between LT80

and TFL80. Two independent runs of sexually mature males were conducted by two

researchers 6 months apart. Red = original data used to calculate TFL80 for *D. virilis* in this

manuscript, blue = repeated assay. Dashed lines intersect at 80% threshold. Line fits

predicted by 3-parameter dose-response model. The fly stock and equipment were

identical.

Repeatability of knockdown CTmax assay

Fig S2.1: Knockdown CTmax assays were generally highly repeatable. Left panel; mean CTmax

temperature recorded from species' independent assays (gold points), and in mixed species

control blocks (pink points). There was no global significant difference between CTmax

estimates across all species (Block: $F_{1,831}$ = 3.246, P = 0.072). However a significant experimental

block*species interaction was found for 4 of the 41 species tested (interaction term: $F_{40,831}$ =

3.7589, P < 0,001); *Drosophila sechelia, D. nasuta, D. yakuba* & *D. tiesseri* all scored significantly

higher CTmax in our mixed species control block than in their own individual CTmax assays.

Right panel: The correlation between estimated CTmax values from both blocks was strongly

positive (coefficient = 1.07, $F_{1,37}$ = 309.84, P < 0.001), and explained 89% of the variation.

Correlations between TFL, LT and CTmax

The correlation between TFL80 and LT80 is significant (PGLS: $F_{1,41}$ = 47.86, P <0.001, coefficient

= 1.631) but LT80 only explains just over half of the variation in TFL80 (PGLS: adjR^2 = 0.53,).

The correlation between absolute TFL80 and CTmax assays is significant (PGLS: $F_{1,41}$ = 19.40, P

<0.001) but considerably weaker than a perfect proxy (PGLS: R^2 = 0.30, coefficient = 0.631).

These analyses were phylogenetically controlled, with no significant phylogenetic signal found

among model residuals (λ = 0.543^{ns}).

The correlation between the two measures of upper lethal limits (LT80 and knockdown CTmax)

is significant and positive (estimate = 0.668, $F_{1,41}$ = 89.176, P <0.001) but also shallower than

would be expected of exact proxies. However, this relationship explains 67.74% of the variation

in LT80 across species. LT80 and knockdown CTmax are different measures of upper thermal

limits and so deviation from parity is not unexpected. Notably, knockdown CTmax inherently

contains an element of thermal ramping, whilst LT80 employs a single static temperature.

These analyses were phylogenetically controlled, with a significant phylogenetic signal found

among model residuals (λ = 0.570**).

The difference between TFL80 and LT80 (ΔLimits) shows no significant relationship with either

CTmax or LT80 (PGLS; knockdown CTmax: coefficient = 0.042, $F_{1,41}$ = 0.127, P = 0.72, R^2 = -0.02;

LT80: coefficient = -0.015, $F_{1,41}$ = 0.011, P = 0.92, R^2 = 0.02). This demonstrates that critical limits

cannot be used as a direct proxy for TFLs, and so TFLs are capturing biological variation that

measurements of critical limits do not. This also suggests that there is not simply an

unavoidable upper limit to fertility shared among species with differing upper critical limits.

713
714
715

 Phylogenetic signal in upper thermal limits

Previous empirical work has found that absolute upper critical limits show strong phylogenetic

signal(*37*). This suggests that these limits may be difficult to rapidly evolve in response to

changes in the environment and suggests that species are sorted into their environment by

their evolutionary history, rather than locally adapted to it(*4*). We find similar degrees of

phylogenetic signal in our measures of critical upper limits (LT80 λ = 0.94 P_0 <0.001, P_1 = 0.108;

κ = 0.866, P = 0.001), (CT_{max} λ = 0.853, P_0 <0.001, P_1 < 0.001; κ = 0.51, P < 0.001). We also find

phylogenetic signal in our estimate of TFL80, however it is less strong than the signal for either

upper lethal limit (λ = 0.722, P_0 = 0.01, P_1 = 0.004; κ = 0.32, P = 0.005). The difference between

LT80 and TFL80 (ΔLimits hereafter) are explained by evolutionary relatedness between species

(λ = 0.823, P_0 = < 0.001, P_1 = 0.024, κ = 0.857, P = 0.001), but this is primarily because the

majority of species in the *Sophophora* subgenus have a ΔLimit of 0°C. Whether this means that

species are able to rapidly evolve TFLs within the range of their ΔLimits is an open question.