Title: Modelling the spatial-temporal distributions and associated determining factors of a keystone pelagic fish

Authors:

Samantha Andrews*
Affiliation: Department of Biology, Memorial University, St. John's, Newfoundland and Labrador, A1B 3X9, Canada

Shawn J. Leroux
Affiliation: Department of Biology, Memorial University, St. John's, Newfoundland and Labrador, A1B 3X9, Canada

Marie-Josée Fortin
Affiliation: Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, Ontario, M5S 3B2, Canada

*Corresponding Author
Email: s-andrews@live.ca
Telephone: +1 709 864 7504

Abstract

Mobile pelagic species habitat is structured around dynamic oceanographic and ecological processes which operate and interact horizontally and vertically throughout the water column and change over time. However, pelagic species movements and distributions are often poorly understood. We use the Maxent species distribution model to assess how changes in the relative importance of modelled oceanographic (e.g., temperature) and climatic variables (e.g., the North Atlantic Oscillation) over 17-years affect the monthly average horizontal and vertical distribution of a keystone pelagic forage species, Atlantic Canadian capelin (*Mallotus villosus*). We show the range and distribution of capelin occurrence probabilities vary across horizontal and vertical axes over time, with binary presence/absence predictions indicating capelin occupy between 0.72% (April) and 3.45% (November) of the total modelled space. Furthermore, our analysis reveals that the importance of modelled oceanographic variables, such as temperature, varying between months (e.g. 44% permutation importance in August to 2% in May). By capturing the spatial dynamics of capelin over horizontal, vertical, and temporal axes, our analysis builds on work that improves our understanding and predictive modelling ability of pelagic species distributions under current and future conditions for pro-active ecosystem-based management.

Key words

Spatial-temporal dynamics; distribution models; marine pelagic fish; capelin (*Mallotus villosus*)

Introduction

Mobile pelagic species have critical functions in marine systems, occupying multiple trophic levels (Sarà and Sarà, 2007; van der Lingen *et al.*, 2010), acting as agents for resource flows (Gounand *et al.*, 2018), and as ecosystem engineers (Breitburg *et al.*, 2010). Despite this, their extensive movements which can cross-jurisdictions means they are typically poorly surveyed. Consequently, the abundance, distribution, and population structures of mobile pelagic species are not well understood.

Species distribution models (SDMs), which combine abiotic (e.g. temperature) and biotic (e.g. prey) variables with species presence data, are commonly used to predict potential species distributions (Guisan *et al.*, 2017). SDMs have been applied to a wide range of marine species including invertebrates (Eger *et al.*, 2017), reef fish (Young and Carr, 2015), and sponges (Knudby *et al.*, 2013). Modelling pelagic species distribution remains challenging, however, because pelagic species habitat is structured around dynamic oceanographic and ecological processes, operating and interacting horizontally and vertically throughout the water column, and changing over time (Angel, 1993). Oceanographic conditions that influence pelagic species distributions include temperature (Rose, 2005), salinity (Vilhjálmsson, 2002), dissolved oxygen content (Bertrand *et al.*, 2011), sea surface height (Zainuddin *et al.*, 2017), and mixed layer depth (Williams *et al.*, 2015). Prey abundance and distribution also plays a pivotal role in pelagic predator distributions (Garrison *et al.*, 2002), with chlorophyll concentration often acting as a proxy for zooplanktivorous prey (Bailey *et al.*, 2012). Furthermore, large-scale climate oscillations such as the North Atlantic Oscillation and Atlantic Multidecadal Oscillation influence regional oceanographic conditions and have been implicated in changing distributions of pelagic species (Gregory *et al.*, 2009).

While distribution models are typically applied to two-dimensional surfaces which

lack temporal variability, the dynamic horizontal, and vertical fluidity of the ocean

necessitates models – particularly those generated for mobile pelagic species which can be

distributed throughout the water column – to account for conditions in four-dimensions –

across longitude and latitude, at depth, and over time. Whereas oceanographic data averaged

over long temporal windows (e.g. decadal, annual) may provide sufficient information to

model the distributions of relatively sedentary species, using such data can lead to loss of

information about how pelagic species are responding spatially to variations in conditions

than more contemporaneous resolutions (e.g. monthly or daily) (Mannocci *et al.*, 2017).

Distribution models that consider changes through time are increasing in number (e.g. Brodie

et al., 2018). However, although the availability in-situ and remotely sensed oceanographic

data and ocean models that provide data at multiple depths is increasing (Kavanaugh *et al.*,

2016), distribution models that incorporate conditions at depth are rare (Duffy and Chown,

2017, but see Hobday and Hartmann, 2006). Such oceanographic information is not truly

three-dimensional, but rather offers snapshots at specific depths creating a "2.5-dimensional"

environment (Duffy and Chown, 2017). Nevertheless, incorporating both depth and time into

SDMs may improve our ability to predict current and future mobile pelagic species

distributions and explore biological responses to global change.

Capelin (*Mallotus villosus*) is a migratory zooplanktivorous pelagic fish species with

a circumpolar distribution in the Pacific, Arctic, and North Atlantic Oceans. Four genetically

distinct populations exist – two in the Pacific, one around West Greenland, and one in the

Northeast Atlantic around Atlantic Canada (Præbel *et al.*, 2008). Within Atlantic Canadian

waters, capelin, whose abundance and distribution is closely tied to environmental conditions,

is regarded as a keystone species (Rose, 2005; Davoren *et al.*, 2006). During the 1990s, the

North Atlantic experienced a substantial change in ocean climate (Greene *et al.*, 2008).

Within Atlantic Canadian waters, alongside numerous groundfish species capelin declined in

abundance, shifted their distributions, began maturing at younger ages and smaller sizes, and

spawned later in the year (Carscadden and Nakashima, 1997). Capelin have not recovered to

their former state, and these changes have been implicated in the lack of recovery of

Canada's Atlantic cod (*Gadus morhua*), which previously had significant economic

importance in the region (Mullowney and Rose, 2014). Despite the ecological and cultural

importance of capelin in Atlantic Canada, their year-round distributions and spatial responses

to changing conditions throughout the region remain poorly understood.

In this study, we address the following research objectives: (1) quantify variation in

the average monthly distribution of capelin across longitude, latitude, and depth to capture

their horizontal and vertical movements, and (2) assess changes in the relative importance of

modelled oceanographic and climatic variables to estimated capelin distributional changes.

We construct monthly "2.5-dimensional" distribution models with Maxent (Phillips *et al.*,

2006), predict the average monthly estimated probability of occurrence across the 2.5-

dimensional geographic space, and derive the relative importance of modelled variables using

Maxent's permutation importance metrics. Results of our analysis will broadly contribute to

the application of distributions models for mobile pelagic species, and more specifically to

the understanding of capelin in Atlantic Canada, a forage species with substantial ecological

and cultural importance in the region.

Methods

Study region

Our study region includes parts of the Atlantic Canadian ocean domain - the ocean and

benthos lying within and adjacent to the Canadian territorial sea (0 – 12 nautical miles from

the low-water line along the coast) and Exclusive Economic Zone (EEZ) (12 – 24 nautical

miles from the low-water line along the coast), broadly extending from 40° latitude to 70°

latitude. To avoid imposing unnatural breaks in capelin distributions, we include France's

EEZ surrounding Saint Pierre and Miquelon, disputed territories with the USA, and areas of

the Grand Banks, Flemish Cap, and Georges Bank and Basin adjacent to Canada's EEZ as

part of the study area (Figure 1).

105
Data – Environmental variables

We obtained oceanographic data (chlorophyll α concentration, density mixed layer depth,

dissolved oxygen, salinity, sea surface height, and temperature), as well as bottom depth,

from the numerical modelling products GLORYS V4.1 (E.U. Copernicus Marine Service

Information, 2018a) and BIOMER V3.2 (E.U. Copernicus Marine Service Information,

2018b). Both products offer global gridded global monthly average measurements at a

horizontal resolution of 0.25˚ across 75 depth levels (0.5 meters to 5,902 meters at varying

intervals) for the years 1998 to 2015 inclusive.

We also obtained monthly averages for two climate oscillations; the Atlantic

Multidecadal Oscillation (AMO) index (ESRL, 2019) calculated from detrended spatially

averaged sea surface temperature anomalies between 87.5° - 87.5°N, 2.5° - 357.5°E and the

North Atlantic Oscillation (NAO) index (NCAR, 2019) calculated from the leading empirical

orthogonal function sea level pressure anomalies between 20°-80°N, 90°W-40°E. In both

cases, index values vary temporally but not spatially across the study region (i.e. characterises

broad temporal variability in ocean conditions within and between months).

Data – Capelin presences

We obtained georeferenced capelin presence data from the Ocean Biogeographic Information

System (OBIS) database (OBIS, 2018). We subset the dataset to the years 1998 to 2014 to

match the oceanographic data availability (we excluded 2015 as there were only ten

presences recorded in the OBIS dataset for the entire year) and only included data points with

sampling depth information. To reduce over-representation of conditions that may arise from

over-sampling of cells (Elith *et al.*, 2006), we reduced the number of presences to one per

spatial-temporal grid cell (see below for details on grid). For example, if a cell had twelve

presence points during March 1999, we reduced the number of presences in that cell to one.

For individual species and size classes, sampling gears differ in their ability to catch

individuals, impacting detectability of both presences, and potentially biasing SDMs (Knudby

et al., 2013). We gleaned sampling gear information from the metadata submitted to OBIS

from data contributors to include as a categorical predictor. Based on the metadata, we

identified nine different, albeit broad, gear type categories (Supplementary File 1).

Spatial-temporal grid

To account for the 3-dimensional nature of the ocean in the SDMs, we created a series of

grids representing each depth included in the model, based on the oceanographic data layers.

Because our distribution model (i.e., Maxent, see below) assumes an equal-area surface, we

projected all layers into an Albers Equal Area (25 km^2) grid for modelling. The grid follows

this same projection. The grid was duplicated to create one grid per month-year period, thus

giving each grid cell a unique x, y, z, and t location. Throughout the remainder of this text,

grids which include only a spatial element will be referred to as the 'spatial grid' while the

grid that includes the month-year period will be referred to as the 'spatial-temporal grid'.

147
Modelling Process

We used Maxent (Phillips *et al.*, 2006 Supplementary File 2) to model the average spatial

distribution of capelin presence probabilities on a month-by-month basis (e.g. one model for

March including March data from all years between 1998 and 2014 for which there is

observation data). We carried out all analysis in R version 3.5.3 (R Core Team, 2019), with

the *Raster* (Hijmans, 2019), *Dismo* (Hijmans *et al.*, 2017), *EcoSpat* (Broennimann *et al.*,

2018), and *enmSdm* (Smith, 2019) packages.

Background Point Generation

Maxent uses background points to characterise the study area. To reduce model over-fitting

and to accurately reflect conditions across a species' range, background points should be

constrained to the geographical range of known occurrence (Elith *et al.*, 2011; Merow *et al.*,

2013). We derived the capelin's horizontal geographic range using a 100% minimum convex

polygon (MCP) (Syfert *et al.*, 2014) (Figure 1). To include depth with the MCP, we also

excluded depth layers falling below the maximum depth of capelin presences (maximum

depth layer 1,045.85 meters). We generated 10,000 background points for each monthly

model (Supplementary File 3). Only one background point per spatial-temporal grid cell was

permitted to prevent over-representation of oceanographic conditions (Elith *et al.*, 2006).

To account for within-month variability (e.g. March 1999 vs March 2000) in

oceanographic conditions, we weighted the number of background points to the number of

presences recorded in each month-year period. For example, if 22% of the unique

observations by cell occurred in 2003, 49% in 2004, 7% in 2005, and 21% in 2006, 22% of

the background points for that model came from 2003, 49% from 2004, 7% in 2005, and 21%

in 2006. To ensure we sufficiently captured conditions across the capelin's range, we also

weighted points by the number of spatial grid cells represented in each Northwest Atlantic

Fisheries Organization (NAFO) division and the area of the Hudson Strait represented in the

MCP (NAFO, 2019). For example, we subdivided the 22% of the background points that

came from 2003 so that 8.8% of the cells came from NAFO division 3K, 7.4 from 3L, 4%

from 3Ps, etc. (Supplementary File 4). This double-weighting of background points also acts

to reduce the influence of sampling bias, which may impact the performance of the models

(Elith et al., 2011).

Data extraction for presence and background points

We extracted ten oceanographic, six climatic, and one static (bathymetry) environmental

predictor variables (Table 1) to the presence and background points. We included both

surface values and values at the depth layer closest to capelin presence/background points as

predictors (Supplementary File 5) except for sea surface height, density mixed layer depth,

and seafloor depth as these are single layer values only. To account for the potential lagged

influence of AMO and NAO on distributions, we added three metrics for the AMO and NAO

predictors: the value of the oscillations during the sampling month, the value during the

previous sampling month, and the mean value from the previous winter (calculated from

December to February values).

Variable selection

To minimise the impact of collinearity, we selected variables for each monthly model a-priori

using Spearman correlation coefficient and variance inflation factor (VIF) (Dormann *et al.*,

2013) (Supplementary File 6). Although Maxent performs well even with small sample sizes

(Pearson *et al.*, 2007), to avoid erroneous estimation of predictor variance, we employed a

minimum events per variable (EPV) rule in which we would only model months that had a

minimum of five samples per predictor (Yalcin and Leroux, 2018).

Maxent modelling

We executed individual monthly models using Maxent version 3.4.1. To allow entropy to be

reached, we set the maximum number of iterations to 5000 for each model. We also set

regularization (β) multiplier values for each monthly model to reduce overfitting and

complexity. To determine the regularization values, we constructed a series of models for

each month with regularization values ranging from 0.25 to 4, at 0.25 step intervals. Optimal

regularization multiplier values were determined using Akaike's Information Criterion

adjusted for small sample size (AICc) (Warren and Seifert, 2011). Final models for each

month used the regularization value that resulted in the lowest AICc.

For each monthly model we used temporally-split cross-validation to train and test the

model (Radosavljevic and Anderson, 2014), in which data from one year was used as the

training data, and tested against all other years, and then repeated until we used all years for

training. We further assessed models performance using several metrics - testing and training

area under the curve (AUC) scores (Phillips *et al.*, 2006), true skills statistic (TSS) (Allouche

et al., 2006), and the continuous Boyce index (CBI) (Boyce *et al.*, 2002) (Supplementary File

7). We calculated each metric separately for each temporal fold which we then averaged to

produce mean predictive performances for each monthly model. Finally, we created

permutation plots for each monthly model to measure mean variable importance of each of

the environmental variables used in the model.

Spatial predictions

For each month, we created average probability maps based on Maxent's cloglog

transformation, which provides an estimated probability of presence (Phillips *et al.*, 2017). In

order to predict estimated probability of occurrence across the depth layers, we concatenated

the depth layers of each oceanographic variables into a single stacked variable layer for each

temporal fold (Bentlage *et al.*, 2013). For example, for March 1999, we had one layer for

chlorophyll α concentrations representing all depth layers, one layer for dissolved oxygen,

one for salinity, and one for temperature. We used these stacked layers to produce probability

maps for each temporal fold, which we then averaged to create monthly average probability

maps for each depth-layer (Supplementary File 8).

To explore general patterns in the probability of capelin presence, we also converted

the probability maps into binary presence/absence maps using maximum Kappa as a

threshold. This threshold is used in a wide range of distribution modelling studies (e.g.

(Davidson *et al.*, 2017; Scherrer *et al.*, 2018), demonstrates resiliency to prevalence which

cannot be accurately determined with presence-only data, and shows good agreement with

the AUC (Phillips *et al.*, 2006; Nenzén and Araújo, 2011).

Results

We obtained a total of 11,517 presence points that matched the required spatial and temporal

extent from OBIS. Of these, we removed 5157 presences that lacked depth information, were

duplicate entries or appeared in the same spatial-temporal cell. A total of 6360 presence

points remained. Presence point data availability was highly variable by month, with June

having the highest number of presences ($n = 1263$) and February the lowest number of

presences ($n = 10$) throughout the study period (1998-2014) (Supplementary File 9).

Of the eighteen variables selected a-priori, multicollinearity analysis indicated only

six, five, and nine variables could be included in the final models for January, February, and

March to December respectively. However, following the EVP rule, we did not run a

February model as there were insufficient presence points. We also chose to not to run a

January model due to the smaller number of sampling years compared to the others (four

years vs eleven+ years for March – December). Selected variables for the March to

December models are presented in Table 1.

Maxent model results

Each of the monthly models on average performed very well (metrics Train AUC, Test AUC, TSS, and CBI; Supplementary File 7). Overall, mean performance of the models was highest in December and lowest in October.

The average (across temporal folds) relative contribution of the variables to each monthly model also varied (Figure 2). Temperature contributed the most to predicting capelin distribution in the June (41%), July (27%), August (44%), September (41%), October (46%), and November (31%) models, and the least to the March (11%), April (3%), and May (2%) models. Salinity had the strongest influence in the March model (38%); however, during no month did it contribute the least to a model fit. Chlorophyll α had the strongest influence on predicted distribution in the December model (41%) and was least important in the May (5%), June (9%), August (1%), and September (3%) models. Dissolved oxygen was the most important predictor of capelin distribution in the May model (51%), and the least in the July (10%), October (8%), and December (12%) models. The climate oscillations (NAO and AMO) played little role in any of the models, nor did they vary substantially in their importance from one month to the next (range 0.06 for NAO value from the previous month in June, to 16% for the NAO value from the previous winter in April).

Monthly probability maps

We produced a total of 460 average (across temporal folds) prediction maps based on Maxent's cloglog transformation, depicting the probability of capelin occurrence from ~0.5 meters depth to ~1,045 meters depth between March and December inclusive (Figure 3).

Alongside latitudinal and longitudinal distributional changes, the average vertical

distribution of probabilities of occurrence also varied by month (Figure 4). On average,

depths of ~244 meters and shallower saw the highest probability scores, except for October,

November, and December which saw higher probabilities at greater depths. Across all depths,

the spread of probability values varies from month to month though some general patterns

emerge when binning the probabilities into ten equally sized groups (Table 2). Most

probability values fall below 0.3 (65.3% in May to 94.07% in September). Between 0.3 and

1, values fall steadily to under 1% of the total number of spatial grid cells. There are two

exceptions to this pattern. In both August and October, there is a small increase in

probabilities falling between 0.7 and 0.79, though values continue to decline thereafter.

When converting the probability maps into binary predicted presence /absence, we see

differences in the horizontal (*x*, *y*) and vertical (*z*) distribution of capelin predicted presences

across the study area (Figures 5 and 6). From March to June, predicted presence cells shifted

eastwards, extending out from the Gulf of St Lawrence and around Nova Scotia to encompass

eastern Newfoundland and the Grand Banks (western trailing cell centroid in March -67.75°

to -64.5° in June, eastern leading cell centroid in March -55.25° to -46.25° in June), with

some cells appearing further north. The vertical distribution of predicted presence cells also

expanded during this period, with cells occupying depth layers from ~0.5 meters to ~147

meters in March to ~42 meters to ~656 meters in June. From July to September, predicted

presence cells largely shrunk back to the Gulf of St Lawrence (western trailing cell centroid

in July -69.25° to -69° in September, eastern leading cell centroid in July -52.5° to -57.5° in

September), before expanding again in October to encompass the eastern side of Nova Scotia

and Newfoundland. During August, predicted presence cells occupied the greatest number of

depth layers (~27 meters to ~1046 meters). In November predicted cells show a shift

eastwards (western trailing cell centroid in October -62.0° to -59.5° in November, eastern

leading cell centroid in October -48° to -46.75° in November) and northwards (southern

trailing cell centroid in October 43° to 44.75° in November, northern leading cell centroid in

October - to -57.5° in September) along Labrador before contracting to eastern

Newfoundland in December. The number of depth layers with predicted presences were also

reduced during this period (~69 meters to ~509 meters in November, and ~69 meters to ~458

meters in December). Following these distribution patterns, the percentage spatial grid cells

predicted to contain capelin in all months remains relatively small – from 0.72% of the total

spatial grid in April up to a maximum of 3.45% in August (Figure 5).

Discussion

Marine-based SDMs are typically applied to 'flat' surfaces, lacking temporal and vertical

information. However, for marine pelagic species which can exhibit extensive movements

horizontally (x, y) and vertically (z) throughout the water column, models should explicitly

consider space in four dimension – across depth and time as well as longitude and latitude.

By coupling long time-series (17 years) of species presence data with "2.5-dimensional"

oceanographic models, we quantified the spatial-temporal dynamics of the estimated

probability of capelin occurrence throughout the water column across Atlantic Canada, and

relative importance of modelled variables on a month-by-month basis.

Our analysis indicates the range and distribution of capelin occurrence probabilities

vary month by month, and over depth, with the highest probability scores for March –

September occurring in shallower depths than for October - December. When binarizing

probability values into presence/absence, we found capelin occupy between 0.72% (April)

and 3.45% (November) of the total modelled space. We also observed depth changes with,

for example, presences primarily located further west in shallower waters in March (~ 26 to

~133 meters) than in December (~78 to ~371 meters). This pattern is consistent with

observations of capelin moving eastwards and into deeper waters towards the end of the year

(Carscadden and Nakashima, 1997; Davoren *et al.*, 2006). While alternative thresholds may

adopt a lower value for binarizing predictions into presence/absence (Nenzén and Araújo,

2011), we expect the general distribution patterns to remain.

By producing monthly models, we were able to assess the relative importance of

modelled variables for monthly capelin distribution. Our analysis revealed that chlorophyll α

concentrations, dissolved oxygen, salinity, and temperature varied in importance as predictors

from one month to the next, with each becoming the most important variable in at least one of

the months modelled. Notably temperature contributed the most to the June – November

models. Temperature is widely considered a key determinate of marine fish distribution,

including for capelin (Rose, 2005), though has also been found to a small role in interannual

variations in the distribution of some species (Thorson *et al.*, 2017). Along similar lines, we

found temperature contributed the least to the March - May and December models. Alongside

temperature, oxygen places substantial metabolic constraints on marine species distributions

(Deutsch *et al.*, 2015), a pattern which emerged in the April and May models in which

dissolved oxygen became the most important predictor of capelin occurrence.

The changes in relative variable importance revealed in our models may indicate

capelin's realized niche changes relative to variations in conditions experienced in each

month and/or with phenology. Indeed, the distribution and frequency of modelled variable

values vary across the modelled space. For example, predictably water temperatures are on

average warmer in August than in March, and chlorophyll α concentrations higher in May

than in August (Supplementary Files 3 and 10). Furthermore, the physiology of any given

species determines which suite of conditions are suitable for survival. While species have

upper and lower limits to physiological tolerances, species may experience conditions that

vary within their tolerance range over time (Stuart-Smith *et al.*, 2017). By capturing changes

in variable importance for pelagic species, models have the potential more closely reflect

temporal dynamics in species realized niche, improving our understanding and predictive

ability of changing distributions under current and future conditions. Additionally, given prey

location plays a role in predator distributions (Garrison et al., 2002), incorporating the spatial-

temporal dynamics of capelin distributions may serve to better inform models for capelin

predator distributions and multispecies assessments in ecosystem-based fishery management.

Distribution models require consideration of the relevance of the spatial and temporal

scales at which species and oceanographic processes operate on (Mannocci *et al.*, 2017). The

oceanographic models chosen to model distributions dictate the resolution of our models

(monthly averaged 25° grid cell). While the oceanographic models allowed us to encompass

a large time-period of capelin data, the resolution may prove too coarse particularly during

biologically important periods, such as during the spawning period when capelin physiology

and behaviour changes (Rose, 2005; Davoren *et al.*, 2006). Furthermore, the models

presented in this study provide monthly average distributions, potentially masking

interannual variation in conditions that determine distributions (Boyce *et al.*, 2002).

Nevertheless, the continued development of finer spatial and temporal scale models may not

only allow for more refined distribution models, but their predictive output can allow for

increased efficiency in the implementation of management measures (Lewison *et al.*, 2015;

Dunn *et al.*, 2016), including nowcasting and forecasting of species distributions for pro-

active management.

Our study adds to the small albeit growing body of distribution models which

embrace a "2.5 dimension" approach, and models that attempt to capture distribution changes

on timescales more relevant to the dynamic nature of pelagic species. With survey effort

spatially and temporally patchily distributed throughout our modelled space, our predicted

occurrences offer new insights into the potential distribution of capelin - a forage species with

substantial ecological importance in Atlantic Canada, and whose life-histories are sensitive to

changing environmental conditions. By capturing species spatial-temporal dynamics over

horizontal and vertical axes, and changes in the relative importance of modelled variables,

such studies can enhance our understanding and predictive modelling ability of pelagic

species distributions under current and future conditions for pro-active ecosystem-based

conservation and fisheries management.

Acknowledgements

This research is sponsored by the Natural Sciences and Engineering Research Council of Canada's Canadian Healthy Oceans Network and its Partners: Department of Fisheries and Oceans Canada and the Institut Nordique de Recherche en Environnement et en Santé au Travail (representing the Port of Sept-Îles and City of Sept-Îles). The authors wish to thank Fran Mowbray (DFO-Newfoundland), the Ecosystem Ecology Lab (Memorial University), Paul Snelgrove (Memorial University), Mariano Kohen-Alonso (DFO-Newfoundland), and Tom Therriault (DFO-Pacific) for their feedback throughout the process. We also thank a number of people who have helped with the underlying code used to create the analysis in this study: David Bazin (Copernicus Marine Environment Monitoring Service); Michael D. Sumner (Australian Antarctic Division); Rémi Daigle (DFO-Maritimes); Simon Beaton (Independent), and the wider Stack Exchange Community, with particular thanks to: TinglTanglBob (Stack Overflow); Nadizan (Stack Overflow); Spacedman (GIS Stack Exchange).

Data Availability Statement

All datasets used in this study were derived from sources in the public domain and are cited
in the text. In order of citation these datasets are:

GLORYS V4.1: E.U. Copernicus Marine Service Information. 2018a. Global Ocean
Reanalysis Products (GLOBAL-REANALYSIS-PHY-001-025) Version 4.1. Available from
https://marine.copernicus.eu/services-portfolio/access-to-products/.

BIOMER V3.2: E.U. Copernicus Marine Service Information. 2018b. Global
Biogeochemical Non assimilative Hindcast Product
(GLOBAL_REANALYSIS_BIO_001_018) Version 3.2. Available from
https://marine.copernicus.eu/services-portfolio/access-to-products/.

The Atlantic Multidecadal Oscillation (AMO) index: ESRL. 2019. Atlantic Multidecadal
Oscillation (AMO) Index (Kaplan SST V2).
https://www.esrl.noaa.gov/psd/data/timeseries/AMO/ (Accessed 2 January 2019).

North Atlantic Oscillation (NAO) index: NCAR. 2019. Hurrell North Atlantic Oscillation
(NAO) Index (PC-Based). https://climatedataguide.ucar.edu/climate-data/hurrell-north-
atlantic-oscillation-nao-index-pc-based (Accessed 2 January 2019).

Capelin presence: OBIS. 2018. Distribution records of Mallotus villosus (Lamarck, 1798).
Available: Ocean Biogeographic Information System. Intergovernmental Oceanographic
Commission of UNESCO. www.iobis.org (Accessed 31 October 2018).

References

Allouche, O., Tsoar, A., and Kadmon, R. 2006. Assessing the accuracy of species distribution models: prevalence, kappa and the true skill statistic (TSS). Journal of Applied Ecology, 43: 1223–1232.

Angel, M. V. 1993. Biodiversity of the pelagic ocean. Conservation Biology, 7: 760–772.

Bailey, H., Fossette, S., Bograd, S. J., Shillinger, G. L., Swithenbank, A. M., Georges, J.-Y., Gaspar, P., *et al.* 2012. Movement patterns for a critically endangered species, the leatherback turtle (*Dermochelys coriacea*), linked to foraging success and population status. PLOS ONE, 7: e36401.

Bentlage, B., Peterson, A. T., Barve, N., and Cartwright, P. 2013. Plumbing the depths: extending ecological niche modelling and species distribution modelling in three dimensions. Global Ecology and Biogeography, 22: 952–961.

Bertrand, A., Chaigneau, A., Peraltilla, S., Ledesma, J., Graco, M., Monetti, F., and Chavez, F. P. 2011. Oxygen: a fundamental property regulating pelagic ecosystem structure in the coastal southeastern tropical Pacific. PLOS ONE, 6: e29558.

Boyce, M. S., Vernier, P. R., Nielsen, S. E., and Schmiegelow, F. K. A. 2002. Evaluating resource selection functions. Ecological Modelling, 157: 281–300.

Breitburg, D. L., Crump, B. C., Dabiri, J. O., and Gallegos, C. L. 2010. Ecosystem engineers in the pelagic realm: alteration of habitat by species ranging from microbes to jellyfish. Integrative and Comparative Biology, 50: 188–200.

Brodie, S., Jacox, M. G., Bograd, S. J., Welch, H., Dewar, H., Scales, K. L., Maxwell, S. M., *et al.* 2018. Integrating dynamic subsurface habitat metrics into species distribution models. Frontiers in Marine Science, 5.

Broennimann, O., Di Cola, V., and Guisan, A. 2018. ecospat: Spatial Ecology Miscellaneous Methods.

Carscadden, J. E., and Nakashima, B. S. 1997. Abundance and changes in distribution, biology and behaviour of capelin in response to cooler waters of the 1990s. *In* Proceedings of Forage Fish in Marine Ecosystems. Alaska Sea Grant Program, pp. 457–468.

Davidson, A. D., Shoemaker, K. T., Weinstein, B., Costa, G. C., Brooks, T. M., Ceballos, G., Radeloff, V. C., *et al.* 2017. Geography of current and future global mammal extinction risk. PLOS ONE, 12: e0186934.

Davoren, G. K., Anderson, J. T., and Montevecchi, W. A. 2006. Shoal behaviour and maturity relations of spawning capelin (*Mallotus villosus*) off Newfoundland: demersal spawning and diel vertical movement patterns. Canadian Journal of Fisheries and Aquatic Sciences, 63: 268–284.

Deutsch, C., Ferrel, A., Seibel, B., Pörtner, H.-O., and Huey, R. B. 2015. Climate change tightens a metabolic constraint on marine habitats. Science, 348: 1132–1135.

Dormann, C. F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz, J. R. G., *et*
al. 2013. Collinearity: a review of methods to deal with it and a simulation study evaluating
their performance. Ecography, 36: 27–46.

Duffy, G. A., and Chown, S. L. 2017. Explicitly integrating a third dimension in marine
species distribution modelling. Marine Ecology Progress Series, 564: 1–8.

Dunn, D. C., Maxwell, S. M., Boustany, A. M., and Halpin, P. N. 2016. Dynamic ocean
management increases the efficiency and efficacy of fisheries management. PNAS, 113: 668–
673.

Eger, A. M., Curtis, J. M. R., Fortin, M.-J., Côté, I. M., and Guichard, F. 2017.
Transferability and scalability of species distribution models: a test with sedentary marine
invertebrates. Canadian Journal of Fisheries and Aquatic Sciences, 74: 766–778.

Elith, J, Graham, C. H., Anderson, R. P., Dudík, M., Ferrier, S., Guisan, A., Hijmans, R. J.,
et al. 2006. Novel methods improve prediction of species' distributions from occurrence data.
Ecography, 29: 129–151.

Elith, J., Phillips, S. J., Hastie, T., Dudík, M., Chee, Y. E., and Yates, C. J. 2011. A statistical
explanation of MaxEnt for ecologists. Diversity and Distributions, 17: 43–57.

ESRL. 2019. Atlantic Multidecadal Oscillation (AMO) Index (Kaplan SST V2).
https://www.esrl.noaa.gov/psd/data/timeseries/AMO/ (Accessed 2 January 2019).

E.U. Copernicus Marine Service Information. 2018a. Global Ocean Reanalysis Products
(GLOBAL-REANALYSIS-PHY-001-025) Version 4.1.
https://marine.copernicus.eu/services-portfolio/access-to-products/ (Accessed 2 January
2019).

E.U. Copernicus Marine Service Information. 2018b. Global Biogeochemical Non
assimilative Hindcast Product (GLOBAL_REANALYSIS_BIO_001_018) Version 3.2.
https://marine.copernicus.eu/services-portfolio/access-to-products/ (Accessed 2 January
2019).

Garrison, L. P., Michaels, W., Link, J. S., and Fogarty, M. J. 2002. Spatial distribution and
overlap between ichthyoplankton and pelagic fish and squids on the southern flank of
Georges Bank. Fisheries Oceanography, 11: 267–285.

Gounand, I., Harvey, E., Little, C. J., and Altermatt, F. 2018. Meta-ecosystems 2.0: rooting
the theory into the field. Trends in Ecology & Evolution, 33: 36–46.

Greene, C. H., Pershing, A. J., Cronin, T. M., and Ceci, N. 2008. Arctic climate change and
its impacts on the ecology of the North Atlantic. Ecology, 89: S24–S38.

Gregory, B., Christophe, L., and Martin, E. 2009. Rapid biogeographical plankton shifts in
the North Atlantic Ocean. Global Change Biology, 15: 1790–1803.

Guisan, A., Thuiller, W., and Zimmermann, N. E. 2017. Habitat Suitability and Distribution
Models: With Applications in R. Cambridge University Press, Cambridge.

Hijmans, R. J., Phillips, S., Leathwick, J. R., and Elith, J. 2017. dismo: Species Distribution
Modeling. https://CRAN.R-project.org/package=dismo.

Hijmans, R. J. 2019. raster: Geographic Data Analysis and Modeling. https://CRAN.R-
project.org/package=raster.

Hobday, A. J., and Hartmann, K. 2006. Near real-time spatial management based on habitat
predictions for a longline bycatch species. Fisheries Management and Ecology, 13: 365–380.

Kavanaugh, M. T., Oliver, M. J., Chavez, F. P., Letelier, R. M., Muller-Karger, F. E., and
Doney, S. C. 2016. Seascapes as a new vernacular for pelagic ocean monitoring, management
and conservation. ICES Journal of Marine Science, 73: 1839–1850.

Knudby, A., Kenchington, E., and Murillo, F. J. 2013. Modeling the distribution of geodia
sponges and sponge grounds in the northwest Atlantic. PLOS ONE, 8: e82306.

Lewison, R., Hobday, A. J., Maxwell, S., Hazen, E., Hartog, J. R., Dunn, D. C., Briscoe, D.,
et al. 2015. Dynamic ocean management: identifying the critical ingredients of dynamic
approaches to ocean resource management. BioScience, 65: 486–498.

Mannocci, L., Boustany, A. M., Roberts, J. J., Palacios, D. M., Dunn, D. C., Halpin, P. N.,
Viehman, S., *et al.* 2017. Temporal resolutions in species distribution models of highly
mobile marine animals: recommendations for ecologists and managers. Diversity and
Distributions, 23: 1098–1109.

Merow, C., Smith, M. J., and Silander, J. A., Jr. 2013. A practical guide to MaxEnt for
modeling species' distributions: what it does, and why inputs and settings matter. Ecography,
36: 1058–1069.

Mullowney, D. R. J., and Rose, G. A. 2014. Is recovery of northern cod limited by poor
feeding? The capelin hypothesis revisited. ICES Journal of Marine Science, 71: 784–793.

NAFO. 2019. Northwest Atlantic Fisheries Organization Geographic Information.
https://www.nafo.int/Data/GIS.

NCAR. 2019. Hurrell North Atlantic Oscillation (NAO) Index (PC-Based).
https://climatedataguide.ucar.edu/climate-data/hurrell-north-atlantic-oscillation-nao-index-pc-
based (Accessed 2 January 2019).

Nenzén, H. K., and Araújo, M. B. 2011. Choice of threshold alters projections of species
range shifts under climate change. Ecological Modelling, 222: 3346–3354.

OBIS. 2018. Distribution records of *Mallotus villosus* (Lamarck, 1798). Available: Ocean
Biogeographic Information System. Intergovernmental Oceanographic Commission of
UNESCO. www.iobis.org (Accessed 31 October 2018).

Pearson, R. G., Raxworthy, C. J., Nakamura, M., and Peterson, A. T. 2007. Predicting species
distributions from small numbers of occurrence records: a test case using cryptic geckos in
Madagascar. Journal of Biogeography, 34: 102–117.

Phillips, S. J., Anderson, R. P., and Schapire, R. E. 2006. Maximum entropy modeling of
species geographic distributions. Ecological Modelling, 190: 231–259.

Phillips, S. J., Anderson, R. P., Dudík, M., Schapire, R. E., and Blair, M. E. 2017. Opening
the black box: an open-source release of Maxent. Ecography, 40: 887–893.

Præbel, K., Westgaard, J. I., Fevolden, S. E., and Christiansen, J. S. 2008. Circumpolar
genetic population structure of capelin *Mallotus villosus*. Marine Ecology Progress Series,
360: 189–199.

R Core Team. 2019. R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. https://www.R-project.org.

Radosavljevic, A., and Anderson, R. P. 2014. Making better Maxent models of species
distributions: complexity, overfitting and evaluation. Journal of Biogeography, 41: 629–643.

Rose, G. A. 2005. Capelin (*Mallotus villosus*) distribution and climate: a sea "canary" for
marine ecosystem change. ICES Journal of Marine Science, 62: 1524–1530.

Sarà, G., and Sarà, R. 2007. Feeding habits and trophic levels of bluefin tuna *Thunnus*
thynnus of different size classes in the Mediterranean Sea. Journal of Applied Ichthyology,
23: 122–127.

Scherrer, D., D'Amen, M., Fernandes, R. F., Mateo, R. G., and Guisan, A. 2018. How to best
threshold and validate stacked species assemblages? Community optimisation might hold the
answer. Methods in Ecology and Evolution, 9: 2155–2166.

Smith, A. B. 2019. enmSdm: Tools for modeling niches and distributions of species. English.
http://www.earthSkySea.org.

Stuart-Smith, R. D., Edgar, G. J., and Bates, A. E. 2017. Thermal limits to the geographic
distributions of shallow-water marine species. Nature Ecology & Evolution, 1: 1846–1852.

Syfert, M. M., Joppa, L., Smith, M. J., Coomes, D. A., Bachman, S. P., and Brummitt, N. A.
2014. Using species distribution models to inform IUCN Red List assessments. Biological
Conservation, 177: 174–184.

Thorson, J. T., Ianelli, J. N., and Kotwicki, S. 2017. The relative influence of temperature and
size-structure on fish distribution shifts: A case-study on Walleye pollock in the Bering Sea.
Fish and Fisheries, 18: 1073–1084.

van der Lingen, C. D., Bertrand, A., Bode, A., Brodeur, R., Cubillos, L., Espinoza, P.,
Friedland, K., *et al.* 2010. Trophic dynamics. *In* Climate change and small pelagic fish, pp.
112–157. Ed. by D. M. Checkley, J. Alheit, Y. Oozeki, and C. Roy. Cambridge University
Press, Cambridge.

Vilhjálmsson, H. 2002. Capelin (Mallotus villosus) in the Iceland–East Greenland–Jan
Mayen ecosystem. ICES Journal of Marine Science, 59: 870–883.

Warren, D. L., and Seifert, S. N. 2011. Ecological niche modeling in Maxent: the importance
of model complexity and the performance of model selection criteria. Ecological
Applications, 21: 335–342.

Williams, A. J., Allain, V., Nicol, S. J., Evans, K. J., Hoyle, S. D., Dupoux, C., Vourey, E., *et*
al. 2015. Vertical behavior and diet of albacore tuna (*Thunnus alalunga*) vary with latitude in

563 the South Pacific Ocean. Deep Sea Research Part II: Topical Studies in Oceanography, 113:
564 154–169.

565 Yalcin, S., and Leroux, S. J. 2018. An empirical test of the relative and combined effects of
566 land-cover and climate change on local colonization and extinction. Global Change Biology,
567 24: 3849–3861.

568 Young, M., and Carr, M. H. 2015. Application of species distribution models to explain and
569 predict the distribution, abundance and assemblage structure of nearshore temperate reef
570 fishes. Diversity and Distributions, 21: 1428–1440.

571 Zainuddin, M., Farhum, A., Safruddin, S., Selamat, M. B., Sudirman, S., Nurdin, N.,
572 Syamsuddin, M., *et al.* 2017. Detection of pelagic habitat hotspots for skipjack tuna in the
573 Gulf of Bone-Flores Sea, southwestern Coral Triangle tuna, Indonesia. PLOS ONE, 12:
574 e0185601.

Tables and Figures

Table 1: Variables used before and after a-priori variable selection based on Spearman
Correlation and VIF for the March – December models. In addition to incompatible VIF and
Spearman scores, gear type information was also too incomplete to be reliably included.

Variable abbreviations: depth – value of the variable at presence depth; surface – value of the
variable at the sea surface; Chl – Chlorophyll concentration (mmol.m $^{-}$3); Depth – Sea floor
depth; MLP - Mixed layer thickness (meters); O2 – Dissolved Oxygen (mmol.m $^{-}$3); Salinity
– Salinity (PSU); SSH- Sea Surface Height (meters) Temperature – Temperature (kelvin);
NAO – North Atlantic Oscillation; AMO – Atlantic Multidecadal Oscillation; sample – value
during the presence month; prev – value from the month prior to presence; winter – value
during the winter previous to presence; Gear – Gear Type (Table SF1 in supplementary
materials).

Data	Variables
All Variables prior to selection	Chl (surface); Chl (depth); Depth; MLP; O2 (surface); O2 (depth); Salinity (surface); Salinity (depth); SSH; Temperature (surface); Temperature (depth); NAO (prev); NAO (sample); NAO (winter); AMO (prev); AMO (sample); AMO (winter); Gear
Variables selected for the March – December models	Chl (surface); O2 (depth); Salinity (depth); Temperature (depth); NAO (prev); NAO (sample); NAO (winter); AMO (sample); AMO (winter)

591   Table 2: Percentage of spatial grid cells falling into Maxent predicted probability of capelin
592   occurrence bins for each of the modelled months. For example, in March 71.88% of cells had
593   a predicted probability of occurrence between 0 and 0.09.

	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0 - 0.09	71.88	64.49	36.52	67.70	71.70	60.29	88.21	47.87	75.13	84.48
0.1 - 0.19	0	0.01	0.18	0.01	0	0	0	0.03	0	0
0.2 - 0.29	11.96	22.03	28.61	16.87	17.18	17.91	5.86	22.58	10.56	4.72
0.3 - 0.39	4.90	6.54	16.89	5.94	4.32	14.43	2.47	11.70	5.48	3.45
0.4 - 0.49	4.46	2.95	6.64	3.20	1.82	5.15	1.36	4.27	2.85	2.29
0.5 - 0.59	2.62	1.69	4.97	2.34	1.42	0.72	0.79	3.34	2.26	1.83
0.6 - 0.69	2.08	1.05	3.08	1.94	1.23	0.46	0.55	3.78	1.83	1.18
0.7 - 0.79	1.34	0.58	1.73	1.32	1.02	0.56	0.47	4.02	1.18	0.95
0.8 - 0.89	0.62	0.45	0.81	0.49	0.85	0.47	0.21	1.73	0.52	0.90
>0.9	0.14	0.22	0.58	0.18	0.45	0.03	0.08	0.69	0.18	0.21

Figure 1: The study region includes waters inside the Canadian EEZ (purple line), the French
EEZ contained within the Canadian EEZ, and adjacent areas of the Flemish Cap, Grand
Banks, Georges Bank, and Georges Basin. Analysis was restricted to the Minimum Convex
Polygon (MCP) boundaries, which was derived from capelin presence data.

Figure 2: Average permutation importance of environmental variables used in the Maxent
models. Values represent averages (across temporal folds) for each month. See Table 1 for a
definition and description of the variables.

Figure 3: Sample of predicted probability of occurrence presence maps generated by Maxent
for March, June, September, and December over four different depth layers. White indicates
the seafloor and grey indicates landmass. Lighter colours indicate lower probabilities and
darker colours higher.

Figure 1: Vertical distribution of Maxent model predicted probability of capelin occurrence from March to December relative to the amount of ocean available at each depth. Note that there is a greater area of ocean at higher depths than lower depths due to increasing intrusion of the sea floor, and that depth layers are not evenly spaced (see Supplementary File 5). Lighter colours indicate lower probability and darker colours higher.

Figure 5: 2-Dimensional Overview of the horizontal (*x,* y) average location of predicted
capelin presence (black) and absence (yellow) for each modelled month, based on maximum
kappa thresholds applied to Maxent model outputs. Areas of occurrence indicates the location
of a presence cell, regardless of depth. The value on the top right indicated the percentage of
cells in the spatial grid with predicted presence.

Figure 6: The vertical average distribution of predicted capelin presence (black) and absence (yellow) relative to the amount of ocean available at each depth layer for each modelled month, based on maximum kappa thresholds. Sea surface is represented at the top of the y-axis and deepest depth at the bottom. Note that there is a greater area of ocean at higher depths than lower depths due to increasing intrusion of the sea floor, and that depth layers are not evenly spaced (see Supplementary File 5).