Expression of a *Malassezia* codon optimized mCherry fluorescent protein in a bicistronic vector

Joleen P.Z. Goh[1], Giuseppe Ianiri[2,3], Joseph Heitman[3],

and Thomas L. Dawson, Jr[1,4]*

[1]Skin Research Institute of Singapore, Agency of Science, Technology and Research, Singapore, Singapore

[2]Department of Agricultural, Environmental and Food Sciences, University of Molise, 86100, Campobasso, Italy

[3]Department of Molecular Genetics and Microbiology, Duke University Medical Centre, Durham, North Carolina, USA

[4]Department of Drug Discovery, School of Pharmacy, Medical University of South Carolina, Charleston, SC, USA

*Correspondence:
Dr Thomas Dawson
thomas.dawson@sris.a-star.edu.sg

Email addresses of authors:

Joseph Heitman
heitm001@duke.edu

Giuseppe Ianiri
giuseppe.ianiri@unimol.it

Joleen P.Z. Goh
joleen.goh@sris.a-star.edu.sg

Keywords: *Malassezia*, fluorescent, multicistronic, bicistronic, transformation, tagging, 2A

Expression of a *Malassezia* codon optimized mCherry fluorescent protein in a bicistronic vector

Abstract

The use of fluorescent proteins allows a multitude of approaches from live imaging and fixed cells to

labelling of whole organisms, making it a foundation of diverse experiments. Tagging a protein of

interest or specific cell type allows visualization and studies of cell localization, cellular dynamics,

physiology, and structural characteristics. In specific instances fluorescent fusion proteins may not be

properly functional as a result of structural changes that hinder protein function, or when overexpressed

may be cytotoxic and disrupt normal biological processes. In our study, we describe application of a

bicistronic vector incorporating a Picornavirus 2A peptide sequence between a NAT antibiotic

selection marker and mCherry. This allows expression of multiple genes from a single open reading

frame and production of discrete protein products through a cleavage event within the 2A peptide. We

demonstrate integration of this bicistronic vector into a model *Malassezia* species, the haploid strain

M. furfur CBS 14141, with both active selection, high fluorescence, and proven proteolytic cleavage.

Potential applications of this technology can include protein functional studies, *Malassezia* cellular

localization, and co-expression of genes required for targeted mutagenesis.

Introduction

Malassezia, comprising 18 currently recognized species, are a unique group of lipophilic

basidiomycetes that evolved independently from related plant pathogen lineages (Xu et al. 2007), and

represent a ubiquitous and dominant eukaryotic microbial community on human skin (Grice and Segre

2011; Findley et al. 2013; Grice and Dawson 2017). In adaption to life on mammalian skin, *Malassezia*

genomes have been reshaped to secrete an armory of proteases, lipases, and phospholipases, amongst

other enzymes to support their growth. These skin-dwelling yeasts usually maintain a symbiotic

relationship with their human host but can quickly shift into opportunistic pathogens, usually induced

by environmental alterations such as breaches in skin barrier integrity, dysfunctional immune response,

or age related changes that affect skin, such as aging, puberty, or menopause (Grice and Segre 2011).

Malassezia are the causative agents of dandruff and seborrheic dermatitis, and are associated with

myriad clinical conditions such as atopic dermatitis, pityriasis versicolor, and folliculitis (Gaitanis et

al. 2012; Theelen et al. 2018). Beyond superficial cutaneous disorders, *Malassezia* are also responsible

for catheter-associated infections and invasive septic fungemia (Barber et al. 1993; Gaitanis et al. 2012;

Kaneko et al. 2012; Iatta et al. 2014). Recent reports illustrate pathogenic roles for *Malassezia* in

Crohn's Disease and pancreatic ductal adenocarcinoma through an elevated inflammatory response

linked to CARD-9 and the complement cascade (Limon et al. 2019; Aykut et al. 2019). Relatively little

is known about the specific mechanisms of *Malassezia* pathogenesis, despite decades of investigation

and their broad significance, meaning much remains to be explored about this important group of

yeasts. Research efforts on *Malassezia* are gaining traction as more species are being identified in

human gut microflora, animal skin, and even in Antarctic and marine environments, making them

amongst the most ubiquitous of fungi (Amend 2014; Theelen et al. 2018). These new findings have

rapidly increased interest, and are driving advances in diverse species identification, definition of

axenic culture conditions, and development and application of tools to dissect *Malassezia* genomic

complexity (Dawson 2019).

Malassezia were widely accepted as highly recalcitrant to conventional transformation techniques

including biolistic, electroporation, and lithium acetate methods. Recent developments in genetic

modifications with the use of *Agrobacterium* have not only established gene transfer, but also provided

tremendous headway in studies of previously uncharacterized *Malassezia* gene function (Ianiri et al.

2016; 2019; Sankaranarayanan et al. 2020). Earlier studies have also applied the use of fluorescence

protein tagging (Celis et al. 2017; Sankaranarayanan et al. 2020). Additionally, the simultaneous co-

expression of multiple genes and fluorescent proteins has applications ranging from monitoring gene

expression (Rasala et al. 2012; Lewis et al. 2015), protein tagging, to live cell or whole organism

labelling and imaging (Provost, Rhee, and Leach 2007; Kim et al. 2011; Ahier and Jarriault 2014),

enabling this technique as a cornerstone in biomedical research.

One of the most common approaches in collective gene expression exploits the incorporation of the

2A oligopeptide, first identified in viral genome of foot-and-mouth disease virus (F2A). Subsequently,

other 2A sequences were discovered in porcine teschovirus-1 (P2A), equine rhinitis A (E2A), and

Thosea asigna virus (F2A). 2A peptides are usually between 18-22 residues and reports have suggested

2A sequences encode a single open reading frame (ORF), and impedes the formation of a peptide bond

between glycine and proline residues, allowing the generation of discrete protein products (Trichas,

Begbie, and Srinivas 2008; Lewis et al. 2015). The use of 2A sequence overcomes the need of

bidirectional or multiple promoters and skips use of numerous co-transfection plasmids, offering

greater efficiency and simplicity in construct designs and transformation.

In this report, we demonstrated use of a bicistronic expression system to simultaneously produce active

fluorescent mCherry and dominant nourseothricin resistance (NAT) non-fusion proteins, delivered via

Agrobacterium tumefaciens-mediated transformation (ATMT) in *Malassezia furfur*. This is achieved

through incorporating viral P2A sequence between gene ORFs, with mCherry upstream of NAT,

designed to ensure expression of mCherry if NAT protein is expressed. *Agrobacterium* Transfer DNA

(T-DNA) expression vector was designed and constructed with unique restriction enzyme sites to allow

straightforward modification to any of the genetic elements to support broad experimental use and

complement alternative experimental needs.

Results

Random Insertional Mutagenesis in *M. furfur*. The expression vector was first constructed in *E. coli*

pUC57 cloning vector by insertion between the actin *ACT1* promoter and terminator of *Malassezia*

sympodialis ATCC 42132, a codon optimized mCherry gene and a P2A viral sequence (Supplementary

Table 1 and Supplementary Figure 1) followed by the *NAT* selection marker gene, forming a single

ORF. The expression cassette was digested from purified pUC57 and cloned into an *Agrobacterium*

tumor-inducing (Ti) backbone binary vector, generating the resulting plasmid, pJG201702 (Figure 1).

The bicistronic expression plasmid pJG201702 was verified by PCR and Sanger sequencing (data not

shown) prior to electroporation into competent *Agrobacterium tumefaciens*.

Transformations were carried out as previously described (Ianiri et al. 2016; Celis et al. 2017), and

randomly selected NAT-resistant colonies were analyzed by PCR to detect the presence of *NAT* and

mCherry ORFs. Predicted amplicons of 576 bp and 708 bp for *NAT* and mCherry respectively, were

observed in 4 out of 23 transformants but absent in wild type (Figure 2A). Transformants and wild type

were serially diluted in PBS and spotted on mDixon and selective medium-supplemented with NAT.

Wild type CBS14141 was not able to proliferate in the presence of NAT while engineered *M. furfur*

strains displayed growth similar to wild type on mDixon (Figure 2B). These results suggest in the

selected transformants, the *NAT* resistance gene was incorporated into the genome, expressed, and

translated into a functional protein without affecting fitness.

P2A cleavage efficiency. To determine whether P2A peptide allows proper and efficient cleavage and

release of the mCherry and NAT proteins, total protein lysates were prepared and analyzed from wild

type and transformants. Immunoblot analysis using a red fluorescent protein antibody (RFP, mCherry)

revealed the expected single band of 29 kDa with no observable band detected in wild type lysate

(Figure 2C).

***In vivo* fluorescence assessment**. *M. furfur* CBS 14141 wild type and three selected genetically

engineered strains, mf::mc-12, -25, and -27 were fluorescently imaged. Live cell imaging revealed

transformants exhibit a higher fluorescence signal compared to wild type (Figure 3A). Wild type

displayed classical autofluorescence observed in *Malassezia* species that localizes mainly in the yeast

cell wall, while all selected transformants displayed a strong mCherry cytoplasmic signal, with mf::mc-

27 demonstrating the strongest fluorescence. In an effort to reduce the autofluorescence background,

mf::mc-27 and wildtype were cultured in liquid media and cells were collected at mid-log phase to

avoid the accumulation of fluorescent metabolites, dead or stationary cells. Further, cells were washed

in PBS to remove residual culture media before imaging but detectable autofluorescence was still

observed in wildtype. Regardless, mf::mc-27 transformant demonstrated higher fluorescence, an

indicator that the mCherry protein was expressed and functional (Figure 3B).

Location of the inserted genetic cassette. Transformant mf::mc-27 was subjected to Illumina

sequencing (Novogene) to determine the site of the T-DNA insertion. Sequenced DNA fragments were

mapped to *M. furfur* CBS 14141 reference genome assembly (Sankaranarayanan et al. 2020) and

RNAseq data (TLD lab, unpublished data). Bioinformatic analysis located the insertion of the

exogenous DNA cassette in an intergenic region on chromosome 2 (Figure 4). Sanger sequencing of

T-DNA through to upstream chromosomal DNA confirmed the presence of a complete ORF of a

putative CDC25-related phosphatase gene. PCR amplifications of the downstream chromosomal DNA

from the insertional point were unsuccessful. However, Illumina sequencing reads were able to detect

both intact transformation DNA cassette and predicted ORF of the adjoining adiponectin receptor.

Discussion

Malassezia are indispensable members of a healthy human skin microbiome, found on almost all

warm-blooded animals, and can even be traced in a marine ecosystem (Amend 2014; Theelen et al.

2018). This diverse group of lipophilic yeasts is often implicated in various cutaneous diseases and

recent studies have identified *Malassezia* as playing pivotal roles in the progression of Crohn's Disease

and exocrine pancreatic cancer (Ashbee and Evans 2002; Limon et al. 2019; Aykut et al. 2019). Yet

much of the specific mechanisms and disease pathogenesis remain elusive due to the lack of capability

to perform gene studies, hampering the advance of research developments. Today, two research groups

have demonstrated the use of a soil bacterium, *Agrobacterium tumefaciens* to genetically modify

Malassezia, paving a new avenue to investigate functional genomics (Ianiri et al. 2016; Celis et al.

2017). In this study, we leveraged the application of *Agrobacterium*-mediated transformation to

introduce a bicistronic expression vector to co-express red fluorescence protein, mCherry, and NAT

resistance marker in *Malassezia furfur*.

The usage of inter-kingdom *Agrobacterium*-mediated transformation in delivering exogenous DNA is

a convenient and versatile tool applied in diverse engineered eukaryotic species. This pathogenic

bacterium effectively housebreaks into the host genome and influences host cellular processes to its

benefit. It also provides an inexpensive and flexible approach in designing and utilizing expression

vectors. In our study, *Agrobacterium tumefaciens* Transfer DNA (T-DNA) does not contain

homologous DNA sequences to *Malassezia furfur* genome, in a deliberate attempt to assess random

integration and gene disruption.

The construction of the bicistronic vector, pJG201702 included the use of actin encoding promoter and

terminator to regulate the expression of mCherry and NAT proteins. The arrangement of the transcribed

proteins was specifically designed to direct the obligatory expression of mCherry when NAT protein

is simultaneously expressed. It was engineered to avoid the possibility of selective gene expression as

Malassezia acquired the potential to undergo genomic rearrangements, excising non-essential genes in

maintaining a condensed genome (Xu et al. 2007; Sankaranarayanan et al. 2020).

Generation of non-fusion proteins was facilitated through the insertion of P2A viral sequence between

mCherry and NAT. Ribosomal skipping of the peptide bond formation between glycine and proline

residues within P2A sequence results in a pseudo-cleavage event to take place, separating the two

protein products (Kim et al. 2011; Szymczak-Workman, Vignali, and Vignali 2012; Liu et al. 2017).

In pJG201702, each genetic component is flanked by restriction enzyme sites, designed for the ease of

future modification including the development of multi-cistronic vectors through incorporating

multiple 2A sequences to generate varied gene products. Additionally, the mCherry and P2A sequences

were codon optimized to improve protein expression in *Malassezia*.

To assess the functionality of P2A sequence in the generation of discrete mCherry and NAT protein

products, a monoclonal antibody specific for red fluorescent protein was probed against whole cell

lysates extracted from wild type and transformants. We detected the predicted protein band at 29 kDa

in all transformants. The pseudo autolytic-cleavage occurs near the end of P2A sequence resulting in

the retention of a 2A tag (21 amino acid residues) at the end of mCherry C-terminus, explaining the

significant shift of the cleaved mCherry protein from its native weight of 26.7 kDa. This suggests

adequate self-cleaving efficiency with the inclusion of a Gly-Ser-Gly (GSG) linker in P2A residues

which promotes cleavage efficiency (Wang et al. 2015; Kim et al. 2011). Correct functionality of the

generated vector was demonstrated by strong mCherry expression of NAT resistant transformants

compared to wild type, which displayed autofluorescence background probably due to the presence of

lipids within the growth media (Croce and Bottiroli 2014) and the production of naturally occurring

intracellular metabolites (Mayser et al. 2002; Maslanka, Kwolek-Mirek, and Zadrag-Tecza 2018).

We sequenced the genome of one fluorescent transformant and confirmed correct T-DNA integration

in *M. furfur* genome. The T-DNA was inserted between two genes, the gene upstream of T-DNA

encodes a putative CDC25-related phosphatase, an yeast ortholog of a Ras guanyl-nucleotide exchange

factor. This gene is thought to be involved in Ras protein signal transduction and cell cycle regulation,

traversing the start control point of the mitotic cell cycle (Chen et al. 2000). Downstream of the T-

DNA insertion encodes for an adiponectin receptor gene that is predicted to be associated with zinc ion

homeostasis (Lyons et al. 2004). Based on the orientation of the genes, it is likely that the T-DNA

integrated in the terminator region of CDC25-related phosphatase and the promoter region of the

predicted adjoining adiponectin-encoding gene. Intergenic insertions of T-DNA are very common and

represent a disadvantage of ATMT in fungi, and they are probably due to a preference for the T-DNA

to insert in low-transcribed regions (Michielse et al. 2005; Idnurm et al. 2017). Intergenic insertions

were also reported in two previous studies in *Malassezia*, suggesting that additional mechanisms that

favor insertion in a ORF-free regions may exist (Ianiri et al. 2016; 2019); this is surprising if we

consider that *Malassezia* genomes are consistently small and compact (Wu et al. 2015). This intergenic

insertion may explain the lack of phenotypic abnormalities or variations in fitness in mc::mf-17

transformant. Intergenic insertions can also be turned into an advantage by using a conditional

promoter that drives the expression of the gene downstream the T-DNA insertion (Kilaru et al. 2015;

Ianiri, Boyce, and Idnurm 2017). Another advantage could be the use of non-protein coding region as

a safe haven for future genetic engineering in *Malassezia*, especially useful for the reintroduction of

genes as complementation without compromising cell viability or disruption of neighboring genes as

reported in *C. neoformans* (Arras et al. 2015; Upadhya et al. 2017).

We demonstrated the effective use of a bicistronic vector in *Malassezia*, to generate a codon optimized

RFP in a NAT-resistant strain. The application of a bicistronic system can potentially be expanded to

a multi-cistronic construct with the aid of multiple cloning sites, to include the expression of assorted

genes according to the users' preference. Most of current research models in understanding skin health

include the use of transgenic mice expressing green fluorescence protein (GFP), mammalian cell

culture, reconstructed skin epidermis (RHE) or even *ex-vivo* skin models, which can be complemented

with these fluorescently labeled *Malassezia*. In addition, the competence to fluorescently label yeast

cells with a separate selectable marker can be applied to deepen our understanding of *Malassezia* gene

functions and host-microbes interactions.

Methods

Strains and culture conditions. Haploid strain *M. furfur* CBS 14141 was obtained from Westerdijk

Fungal Biodiversity Institute. Cells were cultured at 32°C, with mDixon medium (36 g/L malt extract,

20 g/L desiccated oxbile, 6 g/L peptone, 2 ml/L glycerol, 2 ml/L oleic acid, 10 ml/L Tween 40, pH 6).

Genetically transformed cells were maintained on mDixon supplemented with 100 µg/ml

nourseothricin sulfate (NAT).

Plasmid and constructs. pJG201701 was first constructed with *M. sympodialis* ATCC 42132 actin

promoter and terminator sequences, encoding for codon-optimized fluorescent mCherry and porcine

teschovirus-1 2A (P2A) sequences (as detailed in Supplementary Table 1), followed by NAT antibiotic

resistance marker (GenScript) in pUC57 cloning vector. To enable the translation of both proteins, the

stop codon of mCherry was removed and retaining a single stop codon in the second protein, NAT.

Both actin regulatory and NAT sequences were taken from pAIM2 (Ianiri et al. 2016). Each genetic

elements are adjoined with restriction cut sites to allow efficient modification of the construct.

pJG201701 was digested with EcoRI and KpnI, and gene fragments were cloned in pPZP201-BK

(Covert et al. 2001), which contains *Agrobacterium tumefaciens* backbone vector using T4 DNA ligase

(New England Biolabs). Resultant plasmid, pJG201702 was transferred to *A. tumefaciens* EHA105

strain via electroporation and verified by digestion pattern and PCR.

A. tumefaciens-**mediated transformation.** Transformation of *M. furfur* were performed using

previously established protocol (Celis et al. 2017; Ianiri et al. 2016). *A. tumefaciens* harboring

pJG201702 was grown overnight in Luria-Bertani medium supplemented with 50 µg/ml kanamycin at

30°C and 250 rpm in a shaking incubator. Aliquot of cells were used as inoculum, resuspended in

induction medium (IM) containing 100 µM acetosyringone (Sigma) and cultured for additional 6 hours

to OD_{600} of 1. *M. furfur* CBS 14141 cells were collected at mid-log phase and used at OD_{600} of 1.

Proportional volumes of bacterial and yeast cells were mixed and filtered through 0.45 µm mixed

cellulose membrane (Merck, Millipore) before transferring onto IM agar supplemented with 200 µM

acetosyringone. Co-culture plates were incubated at 25°C for 5 days, after which cells were washed in

20 mL sterile PBS and plated on selection medium – mDixon agar containing 100 µg/ml NAT to select

for transformants, and 200 µg/ml cefotaxime with 10 µg/ml tetracycline to halt the growth of *A.*

tumefaciens.

Molecular analysis. Genomic DNA of wild-type and engineered strains of *M. furfur* CBS 14141 were

isolated with MasterPure yeast DNA kit (Epicentre) as per manufacturer's protocol with additional

step of homogenizing at 6 m/s for 50 sec (MP Biomedicals). PCR amplification of mCherry and NAT

genes with mCherry-forward (5'-ATGGTGTCGAAGGGCGAG-3') and mCherry-reverse (5'-

CTTGTAGAGCTCGTCCATGC-3'), and NAT-F (5'-ATGGCGGCCGCCACTCTTGAC-3') and

NAT-R (5'-TTATGGACAAGGCATACTCATATAAAG-3') primers respectively to screen for

positively transformed *Malassezia*.

Total protein extraction and immunoblot. Washed transformant and wild type yeast cells were

collected by centrifugation and resuspended in lysis buffer (50mM Tris-HCl, pH 7.5) with added

protease inhibitor cocktail (Nacalai tesque). Cells were mechanically lysed with mixture of ceramic

and glass beads (Lysing Matrix E, MP Biomedicals) for 50 seconds and repeated twice to achieve

complete cell breakage. Cell debris were separated by centrifugation at 12,000 rpm at 4°C for 5 min

and supernatants were transferred to new tubes for repeated centrifugations. Prepared protein samples

were separated using 4-20% Tris-glycine SDS-PAGE gradient gel and blotted to PVDF membrane.

The blot was blocked in Intercept PBS blocking buffer (Li-cor) and subsequently incubated with RFP

monoclonal antibody (ThermoFisher, MA5-15257) and probed with fluorescent anti-mouse secondary

antibody. Protein bands were visualized with Odyssey CLx digital imaging system (Li-cor).

Genome sequencing. Mf::mc-27 isolate was selected for whole genome analysis and nucleic acid was

extracted using method described previously. Whole genome sequencing was performed with Illumina

Novoseq 6000 at 100x coverage, from 150 bp short insert paired-end reads (NovogeneAIT). CDS

prediction using Augustus gene prediction tool identified putative CDC25 Phosphatase and

Adiponectin Protein Receptor proteins (Supplementary Table 2).

Fluorescence microscopy. Live cell imaging was obtained on inverted wide-field microscope

(Olympus IX-83). Fluorescence and differential interference contrast (DIC) images were achieved with

60x 1.2 oil objective (plan-Apochromat). Exposure time was kept at 700 ms, and cells were held at 37

°C with applied CO_2 incubator chamber during image acquisition. Images were processed and analyzed

with FIJI (ImageJ).

Acknowledgements

This study was supported by Agency for Science, Technology and Research IAF-PP (HBMS) grant

Asian Skin Microbiome Project H18/01a0/016 (to TLD). Work in the Heitman lab was supported by

NIH/NIAID R37 grant AI39115-22 and NIH/NIAID R01 grant AI50113-15. Prof Joseph Heitman is

the co-director and fellow of the CIFAR program Fungal Kingdom: Threats & Opportunities. We

would like to thank Dr Andrea Camattari for providing the *Malassezia sympodialis* codon usage table,

Dr Teun Boekhout at the Westerdijk Fungal Biodiversity Institute for providing the *Malassezia* strains,

and Dr Simon L.I.J. Denil for his help with bioinformatic analysis.

Author Contributions Statement

JPZG, GI, JH and TLD designed the experiments. JPZG performed the experiments and data

analysis. JPZG and TLD wrote the manuscript.

Conflict of interest statement

The authors declare that the research was conducted in the absence of any commercial or financial

relationships that could be construed as a potential conflict of interest.

References

Ahier, Arnaud, and Sophie Jarriault. 2014. "Simultaneous Expression of Multiple Proteins Under a
295   Single Promoter in *Caenorhabditis Elegans* via a Versatile 2A-Based Toolkit." *Genetics* 196
296   (3): 605–13. https://doi.org/10.1534/genetics.113.160846.

297   Amend, Anthony. 2014. "From Dandruff to Deep-Sea Vents: Malassezia-like Fungi Are Ecologically
298   Hyper-Diverse." *PLOS Pathogens* 10 (8): e1004277.
299   https://doi.org/10.1371/journal.ppat.1004277.

300   Arras, Samantha D. M., Jessica L. Chitty, Kirsten L. Blake, Benjamin L. Schulz, and James A.
301   Fraser. 2015. "A Genomic Safe Haven for Mutant Complementation in Cryptococcus
302   Neoformans." Edited by Kirsten Nielsen. *PLOS ONE* 10 (4): e0122916.
303   https://doi.org/10.1371/journal.pone.0122916.

304   Ashbee, H. R., and E. G. V. Evans. 2002. "Immunology of Diseases Associated with Malassezia
305   Species." *Clinical Microbiology Reviews* 15 (1): 21–57.
306   https://doi.org/10.1128/CMR.15.1.21-57.2002.

307   Aykut, Berk, Smruti Pushalkar, Ruonan Chen, Qianhao Li, Raquel Abengozar, Jacqueline I. Kim,
308   Sorin A. Shadaloey, et al. 2019. "The Fungal Mycobiome Promotes Pancreatic Oncogenesis
309   via Activation of MBL." *Nature* 574 (7777): 264–67. https://doi.org/10.1038/s41586-019-
310   1608-2.

311 Barber, Gerard R., Arthur E. Brown, Timothy E. Kiehn, Fitzroy F. Edwards, and Donald Armstrong.
312    1993. "Catheter-Related Malassezia Furfur Fungemia in Immunocompromised Patients." *The*
313    *American Journal of Medicine* 95 (4): 365–70. https://doi.org/10.1016/0002-9343(93)90304-
314    8.

315 Celis, A.M., A.M. Vos, S. Triana, C.A. Medina, N. Escobar, S. Restrepo, H.A.B. Wösten, and H. de
316    Cock. 2017. "Highly Efficient Transformation System for Malassezia Furfur and Malassezia
317    Pachydermatis Using Agrobacterium Tumefaciens-Mediated Transformation." *Journal of*
318    *Microbiological Methods* 134 (March): 1–6. https://doi.org/10.1016/j.mimet.2017.01.001.

319 Chen, R A, T Michaeli, L Van Aelst, and R Ballester. 2000. "A Role for the Noncatalytic N
320    Terminus in the Function of Cdc25, a Saccharomyces Cerevisiae Ras-Guanine Nucleotide
321    Exchange Factor." *Genetics* 154 (4): 1473–84.

322 Covert, Sarah F., Pratibha Kapoor, Mei-ho Lee, Angela Briley, and C. Joseph Nairn. 2001.
323    "Agrobacterium Tumefaciens-Mediated Transformation of Fusarium Circinatum."
324    *Mycological Research* 105 (3): 259–64. https://doi.org/10.1017/S0953756201003872.

325 Croce, A.C., and G. Bottiroli. 2014. "Autofluorescence Spectroscopy and Imaging: A Tool for
326    Biomedical Research and Diagnosis." *European Journal of Histochemistry : EJH* 58 (4).
327    https://doi.org/10.4081/ejh.2014.2461.

328 Dawson, Thomas L. 2019. "Malassezia: The Forbidden Kingdom Opens." *Cell Host & Microbe* 25
329    (3): 345–47. https://doi.org/10.1016/j.chom.2019.02.010.

330 Findley, Keisha, Joy Yang, Sean Conlan, Clayton Deming, Jennifer A. Meyer, Deborah Schoenfeld,
331    Effie Nomicos, Morgan Park, Heidi H. Kong, and Julia A. Segre. 2013. "Topographic
332    Diversity of Fungal and Bacterial Communities in Human Skin." *Nature* 498 (7454): 367–70.
333    https://doi.org/doi:10.1038/nature12171.

334 Gaitanis, G., P. Magiatis, M. Hantschke, I. D. Bassukas, and A. Velegraki. 2012. "The Malassezia
335    Genus in Skin and Systemic Diseases." *Clinical Microbiology Reviews* 25 (1): 106–41.
336    https://doi.org/10.1128/CMR.00021-11.

337 Grice, Elizabeth A, and Thomas L Dawson. 2017. "Host–Microbe Interactions: Malassezia and
338    Human Skin." *Current Opinion in Microbiology* 40 (December): 81–87.
339    https://doi.org/10.1016/j.mib.2017.10.024.

340 Grice, Elizabeth A., and Julia A. Segre. 2011. "The Skin Microbiome." *Nature Reviews*
341    *Microbiology* 9 (4): 244–53. https://doi.org/10.1038/nrmicro2537.

342 Ianiri, Giuseppe, Anna F. Averette, Joanne M. Kingsbury, Joseph Heitman, and Alexander Idnurm.
343    2016. "Gene Function Analysis in the Ubiquitous Human Commensal and Pathogen
344    *Malassezia* Genus." *MBio* 7 (6): e01853-16, /mbio/7/6/e01853-16.atom.
345    https://doi.org/10.1128/mBio.01853-16.

346 Ianiri, Giuseppe, Kylie J. Boyce, and Alexander Idnurm. 2017. "Isolation of Conditional Mutations
347    in Genes Essential for Viability of Cryptococcus Neoformans." *Current Genetics* 63 (3): 519–
348    30. https://doi.org/10.1007/s00294-016-0659-2.

349 Ianiri, Giuseppe, Gabriel Dagotto, Sheng Sun, and Joseph Heitman. 2019. "Advancing Functional
350    Genetics Through Agrobacterium-Mediated Insertional Mutagenesis and CRISPR/Cas9 in the
351    Commensal and Pathogenic Yeast Malassezia." *Genetics* 212 (4): 1163–79.
352    https://doi.org/10.1534/genetics.119.302329.

Iatta, Roberta, Claudia Cafarchia, Teresa Cuna, Osvaldo Montagna, Nicola Laforgia, Ottavio Gentile,
Antonino Rizzo, Teun Boekhout, Domenico Otranto, and Maria Teresa Montagna. 2014.
"Bloodstream Infections by Malassezia and Candida Species in Critical Care Patients."
Medical Mycology 52 (3): 264–69. https://doi.org/10.1093/mmy/myt004.

Idnurm, Alexander, Andy M. Bailey, Timothy C. Cairns, Candace E. Elliott, Gary D. Foster,
Giuseppe Ianiri, and Junhyun Jeon. 2017. "A Silver Bullet in a Golden Age of Functional
Genomics: The Impact of Agrobacterium-Mediated Transformation of Fungi." *Fungal*
Biology and Biotechnology 4 (1): 6. https://doi.org/10.1186/s40694-017-0035-0.

Kaneko, Takamasa, Makiko Murotani, Kiyofumi Ohkusu, Takashi Sugita, and Koichi Makimura.
2012. "Genetic and Biological Features of Catheter-Associated Malassezia Furfur from
Hospitalized Adults." *Medical Mycology* 50 (1): 74–80.
https://doi.org/10.3109/13693786.2011.584913.

Kilaru, S., W. Ma, M. Schuster, M. Courbot, and G. Steinberg. 2015. "Conditional Promoters for
Analysis of Essential Genes in Zymoseptoria Tritici." *Fungal Genetics and Biology* 79
(June): 166–73. https://doi.org/10.1016/j.fgb.2015.03.024.

Kim, Jin Hee, Sang-Rok Lee, Li-Hua Li, Hye-Jeong Park, Jeong-Hoh Park, Kwang Youl Lee,
Myeong-Kyu Kim, Boo Ahn Shin, and Seok-Yong Choi. 2011. "High Cleavage Efficiency of
a 2A Peptide Derived from Porcine Teschovirus-1 in Human Cell Lines, Zebrafish and
Mice." Edited by Volker Thiel. *PLoS ONE* 6 (4): e18556.
https://doi.org/10.1371/journal.pone.0018556.

Lewis, Jo E., John M. Brameld, Phil Hill, Perry Barrett, Francis J.P. Ebling, and Preeti H. Jethwa.
2015. "The Use of a Viral 2A Sequence for the Simultaneous Over-Expression of Both the
Vgf Gene and Enhanced Green Fluorescent Protein (EGFP) in Vitro and in Vivo." *Journal of*
Neuroscience Methods 256 (December): 22–29.
https://doi.org/10.1016/j.jneumeth.2015.08.013.

Limon, Jose J., Jie Tang, Dalin Li, Andrea J. Wolf, Kathrin S. Michelsen, Vince Funari, Matthew
Gargus, et al. 2019. "Malassezia Is Associated with Crohn's Disease and Exacerbates Colitis
in Mouse Models." *Cell Host & Microbe* 25 (3): 377-388.e6.
https://doi.org/10.1016/j.chom.2019.01.007.

Liu, Ziqing, Olivia Chen, J. Blake Joseph Wall, Michael Zheng, Yang Zhou, Li Wang, Haley Ruth
Vaseghi, Li Qian, and Jiandong Liu. 2017. "Systematic Comparison of 2A Peptides for
Cloning Multi-Genes in a Polycistronic Vector." *Scientific Reports* 7 (1): 2193.
https://doi.org/10.1038/s41598-017-02460-2.

Lyons, Thomas J., Nancy Y. Villa, Lisa M. Regalla, Brian R. Kupchak, Anna Vagstad, and David J.
Eide. 2004. "Metalloregulation of Yeast Membrane Steroid Receptor Homologs."
Proceedings of the National Academy of Sciences 101 (15): 5506–11.
https://doi.org/10.1073/pnas.0306324101.

Maslanka, Roman, Magdalena Kwolek-Mirek, and Renata Zadrag-Tecza. 2018. "Autofluorescence
of Yeast Saccharomyces Cerevisiae Cells Caused by Glucose Metabolism Products and Its
Methodological Implications." *Journal of Microbiological Methods* 146 (March): 55–60.
https://doi.org/10.1016/j.mimet.2018.01.017.

Mayser, Peter, Ute Schäfer, Hans-Joachim Krämer, Bernhard Irlinger, and Wolfgang Steglich. 2002.
"Pityriacitrin – an Ultraviolet-Absorbing Indole Alkaloid from the Yeast Malassezia Furfur."

Archives of Dermatological Research 294 (3): 131–34. https://doi.org/10.1007/s00403-002-
0294-2.

Michielse, Caroline B., Paul J. J. Hooykaas, Cees A. M. J. J. van den Hondel, and Arthur F. J. Ram.
2005. "Agrobacterium -Mediated Transformation as a Tool for Functional Genomics in
Fungi." *Current Genetics* 48 (1): 1–17. https://doi.org/10.1007/s00294-005-0578-0.

Provost, Elayne, Jerry Rhee, and Steven D. Leach. 2007. "Viral 2A Peptides Allow Expression of
Multiple Proteins from a Single ORF in Transgenic Zebrafish Embryos." *Genesis* 45 (10):
625–29. https://doi.org/10.1002/dvg.20338.

Rasala, Beth A., Philip A. Lee, Zhouxin Shen, Steven P. Briggs, Michael Mendez, and Stephen P.
Mayfield. 2012. "Robust Expression and Secretion of Xylanase1 in Chlamydomonas
Reinhardtii by Fusion to a Selection Gene and Processing with the FMDV 2A Peptide."
Edited by John W. Stiller. *PLoS ONE* 7 (8): e43349.
https://doi.org/10.1371/journal.pone.0043349.

Sankaranarayanan, Sundar Ram, Giuseppe Ianiri, Marco A Coelho, Md Hashim Reza, Bhagya C
Thimmappa, Promit Ganguly, Rakesh Netha Vadnala, et al. 2020. "Loss of Centromere
Function Drives Karyotype Evolution in Closely Related Malassezia Species." *ELife* 9
(January): e53944. https://doi.org/10.7554/eLife.53944.

Szymczak-Workman, Andrea L., Kate M. Vignali, and Dario A. A. Vignali. 2012. "Design and
Construction of 2A Peptide-Linked Multicistronic Vectors." *Cold Spring Harbor Protocols*
2012 (2): 199–204. https://doi.org/10.1101/pdb.ip067876.

Theelen, Bart, Claudia Cafarchia, Georgios Gaitanis, Ioannis Dimitrios Bassukas, Teun Boekhout,
and Thomas L Dawson. 2018. "Malassezia Ecology, Pathophysiology, and Treatment."
Medical Mycology 56 (suppl_1): S10–25. https://doi.org/10.1093/mmy/myx134.

Trichas, Georgios, Jo Begbie, and Shankar Srinivas. 2008. "Use of the Viral 2A Peptide for
Bicistronic Expression in Transgenic Mice." *BMC Biology* 6 (1): 40.
https://doi.org/10.1186/1741-7007-6-40.

Upadhya, Rajendra, Woei C. Lam, Brian T. Maybruck, Maureen J. Donlin, Andrew L. Chang, Sarah
Kayode, Kate L. Ormerod, James A. Fraser, Tamara L. Doering, and Jennifer K. Lodge.
2017. "A Fluorogenic C. Neoformans Reporter Strain with a Robust Expression of m-Cherry
Expressed from a Safe Haven Site in the Genome." *Fungal Genetics and Biology* 108
(November): 13–25. https://doi.org/10.1016/j.fgb.2017.08.008.

Wang, Yuancheng, Feng Wang, Riyuan Wang, Ping Zhao, and Qingyou Xia. 2015. "2A Self-
Cleaving Peptide-Based Multi-Gene Expression System in the Silkworm Bombyx Mori."
Scientific Reports 5 (1): 16273. https://doi.org/10.1038/srep16273.

Wu, Guangxi, He Zhao, Chenhao Li, Menaka Priyadarsani Rajapakse, Wing Cheong Wong, Jun Xu,
Charles W. Saunders, et al. 2015. "Genus-Wide Comparative Genomics of Malassezia
Delineates Its Phylogeny, Physiology, and Niche Adaptation on Human Skin." Edited by
Gregory S. Barsh. *PLOS Genetics* 11 (11): e1005614.
https://doi.org/10.1371/journal.pgen.1005614.

Xu, J., C. W. Saunders, P. Hu, R. A. Grant, T. Boekhout, E. E. Kuramae, J. W. Kronstad, et al. 2007.
"Dandruff-Associated Malassezia Genomes Reveal Convergent and Divergent Virulence
Traits Shared with Plant and Human Fungal Pathogens." *Proceedings of the National*
Academy of Sciences 104 (47): 18730–35. https://doi.org/10.1073/pnas.0706756104.

Figure 1. Random integration bicistronic vector. Generation of bicistronic construct comprised use of promoter and terminator sequences from *Malassezia sympodialis* ATCC 42132 in *Agrobacterium tumefaciens* backbone vector. Actin encoding regulatory elements govern the sequential expression of mCherry and nourseothricin sulfate (NAT) antibiotic resistance marker, separated by porcine teschovirus-1 2A (P2A) pseudo-autolytic cleavage sequence. pJG201702 vector was created with multiple restriction cut sites between each genetic element for future modification.

Expression of a *Malassezia* codon optimized mCherry fluorescent protein in a bicistronic vector

Figure 2. Characterization and analysis of transformants. (A) PCR screening of transformants and wild-type cells were analyzed with mCherry and NAT ORF spanning primers, indicating the appropriate sized product in transformants and not wild-type cells. **(B)** Transformants and *M. furfur* CBS 14141 wild-type clones were ten-fold serially diluted in PBS and 3uL of cell suspensions were spotted on mDixon growth medium and selection media-supplemented with 100ug/mL NAT, indicating NAT resistance in selected clones. **(C)** Immunoblot was performed with whole cell lysates, probed with anti-RFP primary antibodies. Transformants expressing mCherry contained additional amino acid residues from viral P2A protein tag, produced expected protein bands with an upward shift of molecular weight at 29 kDa. In comparison, 26.7 kDa native molecular weight band was detected in positive control, TurboRFP expressing keratinocyte cell lysate.

Figure 3. Live fluorescent Imaging. (A) Transformants and wild-type cells were collected from mDixon agar and imaged under TRITC channel and differential interference contrast (DIC). Wild-type cells displayed localized autofluorescence in cell wall, in contrast to transformants exhibiting stronger fluorescence distributed uniformly throughout the cells. **(B)** Cells in exponential growth phase were collected to minimize accretion of dead or stationery cells and aggregation of fluorescence metabolites. Transformant mf::mc-27 presented higher fluorescence intensity in comparison to *M. furfur* CBS 14141 wild type cells which displayed persistent low levels of fluorescence background.

Expression of a *Malassezia* codon optimized mCherry fluorescent protein in a bicistronic vector

Figure 4. Random insertional mutagenesis. Using Illumina and Sanger sequencing, exogenous T-

DNA was identified in chromosome 2 of mf::mc-27 mutant. Sanger sequencing further validates the

presence of CDC25 phosphatase gene, adjacent to exogenous DNA of actin terminator and NAT.

Illumina reads were able to detect both complete flanking ORFs of the hypothetical adiponectin rector

protein, and exogenous T-DNA containing *Malassezia* actin promoter, terminator, and mCherry-P2A-

NAT cassette.