Nucleotide excision repair hotspots and coldspots of UV-induced DNA damage in the human genome

Yuchao Jiang[1,2,3,*,#], Wentao Li[4,*], Laura A Lindsey-Boltz[4,*], Yuchen Yang[2], Yun Li[1,2,5], Aziz Sancar[3,4,#]

[1] Department of Biostatistics, Gillings School of Global Public Health, University of North Carolina, Chapel Hill, NC 27599, USA.
[2] Department of Genetics, School of Medicine, University of North Carolina, Chapel Hill, NC 27599, USA.
[3] Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, NC 27599, USA.
[4] Department of Biochemistry and Biophysics, School of Medicine, University of North Carolina, Chapel Hill, NC 27599, USA.
[5] Department of Computer Science, College of Arts and Sciences, University of North Carolina, Chapel Hill, NC 27599, USA.

[*] These authors contributed equally.
[#] To whom correspondence should be addressed. Email: yuchaoj@email.unc.edu, aziz_sancar@med.unc.edu.

ABSTRACT

We recently developed high-throughput sequencing approaches, eXcision Repair sequencing (XR-seq) and Damage-seq, to generate genome-wide mapping of DNA excision repair and damage formation, respectively, with single-nucleotide resolution. Here, we used time-course XR-seq data to profile UV-induced excision repair dynamics, paired with Damage-seq data to quantify the overall induced DNA damage. We identified genome-wide repair hotspots exhibiting high-level nucleotide excision repair immediately after UV irradiation. We show that such repair hotspots do not result from hypersensitivity to DNA damage, and are thus not damage hotspots. We find that the earliest repair occurs preferentially in promoters and enhancers from open-chromatin regions. The repair hotspots are also significantly enriched for frequently interacting regions and super-enhancers, both of which are themselves hotspots for local chromatin interactions. Further interrogation of chromatin organization to include DNA replication timing allows us to conclude that early-repair hotspots are enriched for early-replication domains. Collectively, we report genome-wide early-repair hotspots of UV-induced damage, in association with chromatin states and epigenetic compartmentalization of the human genome.

INTRODUCTION

UV in sunlight is a known mutagen and causative agent of skin cancer that induces DNA lesions such as cyclobutane pyrimidine dimers (CPDs) and pyrimidine-pyrimidone (6-4) photoproducts [(6-4)PPs] (1,2). In humans, both damage types are solely repaired by nucleotide excision repair (excision repair), which removes DNA lesions by dual incisions bracketing modified bases and fills and seals the resulting gap by DNA synthesis and ligation (3). Excision repair consists of two pathways, global repair and transcription-coupled repair (4,5), that differ primarily in the damage recognition step (3-6). For UV-induced DNA damage, CPD repair is highly associated with transcription, specifically with the transcribed strand, while (6-4)PP repair is uniformly distributed throughout the genome (7).

Recently, we developed a high-throughput approach, excision repair sequencing (XR-seq), to isolate the oligonucleotides excised by excision repair and subject them to next-generation sequencing (7). This method has allowed for genome-wide mapping of UV-induced excision repair with single-nucleotide resolution in human (7), Lemur (8), mouse (9), *Drosophila melanogaster* (10), *Saccharomyces cerevisiae* (11), *Escherichia coli* (12), and *Arabidopsis thaliana* (13). We further developed another next-generation sequencing method, Damage-seq (14), to generate genome-wide mapping of UV damage formation with single-nucleotide resolution (15).

Using the combined UV damage maps and repair maps, it has been shown that the induced DNA damage is uniformly distributed throughout the genome and that the overall effect of damage in the genome is primarily driven, not by damage formation, but by repair efficiency (14,15). Indeed, different repair efficiencies across the genome have been previously reported, which are impacted by multiple factors, including transcription, chromatin state and structure, regulatory protein binding to DNA, and posttranscriptional modification of histones (7,15-22). Existing studies, however, have been focused on profiling repair dynamics over a time course – 4 h for (6-4)PP and 48 h for CPD. Genomic regions that harbor high-level early repair shortly after damage formation have not been studied, nor have their associated genomic and epigenomic characteristics been systematically explored.

Here, we performed XR-seq at times as early as 1 min for (6-4)PP and 12 min for
CPD following UV irradiation to identify such repair hotspots. We systematically
characterized the identified hotspots using additional high-throughput sequencing data
that measures DNA damage formation, DNase I hypersensitivity, histone modifications,
3D chromatin interactions, and DNA replication timing. These extensive links between
chromatin states and organization, cell cycle, DNA damage, and excision repair at
specific genomic sites facilitate a better understanding of mutagenesis and
carcinogenesis.
56

RESULTS

Genome-wide profiling of DNA excision repair kinetics through ordered high-
throughput experiments

In this work, we present an experimental and analytical framework where we
systematically assay DNA excision repair of UV-induced DNA damage over a time course.
Figure 1A gives an outline of the experimental design. Specifically, we adopted XR-seq
(7) to measure repair of (6-4)PPs at 1 min, 2 min, 5 min, 20 min, 1 h, 2 h, and 4 h and
CPDs at 12 min in normal human skin fibroblasts (NHF1) after 20J/m$_2$ UV treatment. The
(6-4)PP XR-seq experiments from 1 min to 4 h were all performed with two biological
replicates, with correlation coefficients greater than 0.99 between each pair of replicates
(Figure S1). Quality control procedures show that all samples have good data quality with
expected read length distributions and enrichments of TT and/or TC dinucleotides at the
damage sites (Figure S2). To investigate the interplay between DNA damage and repair,
we additionally adopted Damage-seq data (15) to profile the two types of UV-induced
damage at 0 timepoint after UV irradiation. Refer to the Methods section for details on
library preparation, sequencing, and bioinformatic analysis and Table S1 for detailed
sample information.

The bulk repair pathways and kinetics of these two types of DNA damage are
known to be quite different (23,24). The global repair efficiently removes (6-4)PPs, with
most of the damage excised within 4 h of UV treatment. We provide additional empirical
evidence that the repair levels of (6-4)PP from the transcribed strand (TS) and the non-
transcribed strand (NTS) are on par across all timepoints (Figure S3A-B). In contrast,

CPDs are recognized primarily in a transcription-coupled manner, and their complete
repair requires 12 to 48 h, depending on the UV dose. In our experiment, however, given
the sampling timepoint as early as 12 min for CPDs, transcription-coupled repair has not
contributed much, if at all, to the repair of CPDs. Specifically, as shown in Figure S3C,
we do not observe a difference between TS and NTS repair at 12 min, which is likely too
early for enough polymerases to have stalled and recruited the repair factors (25). This is
of key importance so that our analysis is unbiasedly focused on the global repair of (6-
4)PP at any timepoints and of CPD at the early 12 min timepoint, without transcription as
a confounder. Refer to the Discussion section for more details.

In this study, we focus on characterizing global repair kinetics and identifying early-
repair hotspots and late-repair coldspots. *In vivo* excision assay (24) by gel
autoradiography at 0 min and 2 min reveals that the primary excision repair products can
be detected as early as 2 min after UV treatment (Figure 1B). Using genome-wide repair
data by XR-seq, we performed principal component analysis (PCA) (26) on the top 2,000
highly variable genes to generate a low-dimensional representation of the (6-4)PP repair
data (Figure 1C). Since transcription-coupled repair does not contribute to the repair of
(6-4)PPs, the PCA plots do not differ between the TS and NTS repair, except for reverted
signs of the eigenvectors. Importantly, a repair trajectory on the principal component
space can be reconstructed, which lines up well with the timepoints, suggesting
differential repair kinetics over the time course (Figure 1C).

Identification of early-repair hotspots and late-repair coldspots in normal human fibroblasts

We developed a computational framework to identify genome-wide early-repair hotspots
using time-course XR-seq data. Briefly, we segmented the genome into consecutive bins
of 50 bp long and identified bins that showed significantly enriched repair at earlier or later
timepoints using a thresholding approach on the downsampled reads (Figure S4). We
define such genomic bins as repair hotspots and coldspots, respectively. While such a
method is efficient and effective in identifying the top few hundred repair hotspots and
coldspots, we additionally adopted a more rigorous Poisson log linear model (27,28) on
the read count data for normalization and testing of repair enrichment. The identified

repair hotspots and coldspots show enriched repair levels compared to those as expected under the null (Figure S5). Refer to the Methods section for details.

Figure 1D shows the distributions of read counts per genomic bin across all (6-4)PP samples, and we note enrichment of both early repair at 1 min and late repair at 4 h, corresponding to repair hotspots and coldspots, respectively. For repair of (6-4)PPs, we identified on the genome-wide scale 175 and 156 repair hotspots from the plus and minus strand (Table S2) and 48 and 57 repair coldspots from the plus and minus strand (Table S3), respectively. For repair of CPDs, we focused on repair hotspots at an early timepoint to negate the global impact of transcription, and identified 99 and 93 repair hotspots from the plus and minus strands, respectively (Table S4). We could not search for intrinsic CPD coldspots, because 30 min after UV irradiation transcription-coupled repair becomes an important contributor to the repair profile.

The repair hotspots and coldspots reported are scattered across the entire human genome (Figure S6). XR-seq signals from examples of repair hotspots and coldspots are separated by strand and plotted across all timepoints in Figure 2. We also include epigenomic signals by DNase-seq, ChIP-seq from ENCODE (29), and Damage-seq signals at 0 min after UV treatment (15). Specifically, the XR-seq signals from an example (6-4)PP repair hotspot decrease dramatically from 1 min to 20 min and can be barely seen at 1 h (Figure 2A). In contrast, the XR-seq signals at a late-repair coldspot shown in Figure 2B increases over the time course and peaks at 4 h. Another representative CPD repair hotspot at 12 min is shown in Figure 2C.

We next performed sequence context analysis using all reads that are mapped to the repair hotspots and coldspots, respectively. We trimmed the reads to be 15 bp long centering at the damage sites and calculated strand-specific nucleotide frequencies in repair hotspots, coldspots, and randomly picked spots. Interestingly, we identified an enrichment of cytosine in the flanking regions of the damage/repair sites for both repair hotspots and coldspots (Figure S7). Motif analysis by the MEME suite (30) confirmed the enrichment of cytosine adjacent to the repair sites, which are themselves enriched with canonical sequences of CTCA for (6-4)PP and TT for CPD (Table S5). While it has been previously shown that active transcription factor binding sites have decreased levels of excision repair (21), here we report a preference for cytosine at both early and late

141 timepoints, which have increased levels of repair compared to null regions. Notably, such
142 cytosine enrichment in the flanking regions of the repair sites is not due to sequence
143 context bias, as we show in Figure S8 that not all genomic regions enriched for cytosines
144 are enriched for excision repair. While an interesting observation, the underlying
145 biological mechanisms need further investigation.
146
Early-repair hotspots are not hypersensitive to DNA damage
A recent study assayed DNA damage by next-generation sequencing and reported a total
of 153 hyper-hotspots acquiring CPDs much more frequently than the genomic average
in primary human fibroblasts (31). Of these damage hotspots, 83 are from the plus strand
and 74 are from the minus strand, each having at least five recurrent sequence reads
(31). To investigate whether the identified repair hotspots simply result from increased
levels of DNA damage, we first intersected the reported damage hotspots from Premi et
al. (31) with the 99 and 93 CPD repair hotspots from the plus and minus strand that we
identified. We found that none of our identified repair hotspots overlapped with the
reported damage hotspots.
To further confirm and replicate this seemingly striking result, we analyzed
genome-wide CPD DNA damage data generated by our previously developed Damage-
seq protocol (15), to quantify damage levels at 0 timepoint after UV irradiation with single-
nucleotide resolution. After stringent quality control procedures (refer to the Methods
section for details), we identified 91 and 78 CPD damage hotspots from the plus and
minus strand, respectively, each having at least ten mapped reads (Table S6). Notably,
these Damage-seq hotspots are shown to be enriched for heterochromatin and repressed
regions (Figure S9), which is concordant with previous reports (32,33). Importantly, none
of the CPD damage hotspots, identified from this parallel Damage-seq platform, overlap
with the repair hotspots.
In addition, we compared the DNA damage levels for (6-4)PP and CPD from three
independent sequencing technologies – Damage-seq (15), adductSeq (31), and CPD-
seq (19) – at our identified hotspots and coldspots against those from randomly sampled
regions along the genome. To account for the sparse sampling when measuring DNA
damage by next-generation sequencing, we also extend the regions corresponding to the

repair hotspots, coldspots, and random spots at both ends for 20bp and 500bp,
respectively. Our results, shown in Figure 3, suggest that there is no significant difference
in the damage levels between the three repair categories (hotspot, coldspot, and random
spot). The zoom-in and zoom-out views of three example repair hotspots and coldspots
in Figure 2 also suggest that the Damage-seq reads are uniformly distributed in the
flanking regions. Previous results have demonstrated that the UV-induced DNA damage
is indeed virtually uniform across the entire human genome, while repair is affected by
chromatin states, transcription factor binding, etc., in a manner dependent on the type of
DNA damage (15). While we note that the shallow depth of coverage of Damage-seq can
be a limiting factor (refer to the Discussion section for details), our results validate our
conclusion that the identified repair hotspots are not damage hotspots.
183
Early-repair hotspots are enriched for promoters and enhancers in open chromatin
regions
The packaging of DNA into chromatin can hinder the access of repair proteins and affect
the repair efficiency (16,17) and specific histone modifications are associated with
different functional and cytological chromatin states. To investigate the relationship
between the identified repair hotspots/coldspots and the epigenomic markers, we used
publicly available DNase-seq data and histone modification ChIP-seq data for an adult
human fibroblast cell line (NHLF) from ENCODE (29). Our results suggest that the earliest
repair occurs preferentially in active and open-chromatin regions. Specifically, chromatin
accessibility by DNase-seq is significantly higher in repair hotspots than in random
genomic regions, and it is significantly lower in repair coldspots (Figure 4). For histone
modifications, repair hotspots have significantly higher ChIP-seq signals compared to
random spots for activation markers, including H3K4me1, H3K4me3, and H3K27ac
(Figure 4).
We adopted the segmented chromatin states by chromHMM (34) to annotate the
repair hotspots for (6-4)PP and CPD and found a significant enrichment of active
promoters and enhancers (Figure 5A), which are characterized by nucleosome loss and
open chromatin. Genome-wide repair kinetics inferred across timepoints confirm that
enhancers are repaired at earlier timepoints and that repressed/heterochromatin regions

are repaired at later timepoints (Figure S10). Additionally, we performed genome-wide

annotations for CpG islands and genic/intergenic regions. Our results suggest that the

repair hotspots are enriched for inter CpG regions (i.e., depleted for CpG islands) (Figure

5B) and intronic regions (Figure 5C), both of which are enriched for AT and presumably

result in more repair. Notably, the number of identified repair coldspots is too low to

generate reproducible annotation results between the replicates.

210 Early-repair hotspots are enriched for frequently interacting regions (FIREs) and

211 super-enhancers

With the genome-wide enrichment of active promoters and enhancers in open chromatin

regions of the detected repair hotspots, we hypothesize that the 3D genome structure

contributes to the observed differential repair kinetics between different regions of the

genome. To test this hypothesis, we sought to interrogate the publicly available Hi-C data

of human fibroblast cell line IMR90 (35,36). Specifically, after quality control procedures

and data normalization, we profiled frequently interacting regions (FIREs) using

FIREcaller (37). After overlapping the repair hotspots and coldspots with the called FIREs

(Table S7A), we found that a significantly higher proportion of repair hotspots overlap with

FIREs – 23.16% and 11.76% for (6-4)PP and CPD, respectively – compared to a genome

average of 6.93% based on the profiled FIREs (Figure 6A). Conversely, the overlapping

proportion of (6-4)PP repair coldspots is 3.23%, significantly lower than the genome

average (Figure 6A).

FIREs have been previously reported to be enriched for super-enhancers (38). We

have demonstrated that the repair hotspots are enriched for both FIREs and enhancers.

We also observed that, across many cases, multiple enhancers that overlapped with the

repair hotspots were from the same genomic regions (Figure S11). As such, we

hypothesize that the repair hotspots are also enriched for super-enhancers and thus

additionally adopted a list of previously annotated super-enhancers in the human

fibroblasts (39) (Table S7B). We found that, compared to a genome-wide average of

2.05%, the early-repair hotspots are indeed enriched for super-enhancers (5.14% and

4.69% for (6-4)PP and CPD repair hotspots, respectively), while none of the repair

coldspots overlap with super-enhancers (0% for (6-4)PP repair coldspot) (Figure 6B).

In addition, we called significant interactions based on the Hi-C contact matrix

using the Fit-Hi-C method (40) (Table S7C) and showed that early-repair hotspots also

overlap with a significantly higher number of significant interactions (Figure 6C). Refer to

the Methods section for details on data analysis. The overlapping information of the called

repair hotspots and coldspots with the profiled FIREs, super-enhancers, and significant

chromatin interactions are included in Table S8. Figure 6D illustrates the loop interactions

of two identified repair hotspots. Notably, these two hotspots also overlap with both FIREs

and super-enhancers. These results collectively provide a global picture of genetic

regulation of repair kinetics via 3D genome structures.

Early-repair hotspots are enriched for early-replication domains

We next set out to test the relationship between the repair kinetics and replication timing.

We used the genome-wide Repli-Seq data of the human fibroblast cell line IMR90, which

maps high-resolution DNA replication patterns with respect to both cell-cycle time and

genomic position (41). A recent re-analysis of the data using a deep-learning model (42)

segmented the genome into different replication domains, including the early-replication

domain (ERD), late-replication domain (LRD), up-transition zone (UTZ), and down-

transition zone (DTZ). Refer to Table S9 for genome-wide segmentation results. When

overlapping the identified repair hotspots with the segmented replication domains, we find

that early-repair hotspots are also significantly enriched for early-replication domains

(Figure 7). While replication time has been shown to be correlated with chromatin

accessibility (41), here we have demonstrated a potential cell-cycle effect on DNA

excision repair. That is, regions that are duplicated early also tend to be repaired early,

and this has been shown to result in mutagenesis asymmetries (17) – we further elaborate

such implications in the Discussion section.

Profiling gene-level repair kinetics in normal human fibroblasts

So far, our analysis has been conducted on the whole-genome level. Identifying dynamic

changes at individual gene levels can provide important insights about genes and

pathways that exhibit significant changes in repair dynamics (43). We calculated strand-

specific reads per kilobase per million reads (RPKM) for all the genes in the human

genome. RPKMs across nine genes associated with circadian rhythm (e.g., *CLOCK*, *BMAL1*, *PER1*, *CRY1*) and excision repair (e.g., *XPA*, *XPC*, *CSA*, *CSB*) present different repair dynamics over the time course (Figure S12). We adopted Trendy (44) to perform segmented regression analysis (Table S10) on the ordered XR-seq data. We focused on significant genes with adjusted R^2 greater than 0.8 from the regression analysis and identified 3,017 significant genes with fitted trend as "down," 3,496 significant genes with fitted trend as "up-down," 1,024 significant genes with fitted trend as "up," and 1,452 significant genes with fitted trend as "down-up" (Figure S13). The number of genes that show the "down" trend is two-fold higher than the number of genes that show the "up" trend, which is consistent with our previous observation – there are more repair hotspots than coldspots. For these four groups of significant genes, we further performed gene ontology (GO) enrichment analysis using PANTHER (45). For genes with the "down" and "up-down" trend, GO biological processes are enriched for a large number of biological regulations, including regulation of the metabolic process, gene expression, developmental process, cell cycle, and cell death, etc. (Table S11A-B). On the other hand, there are no obvious GO enrichment terms for the "up" or "down-up" genes (Table S11C-D).

DISCUSSION

We adopted time-course XR-seq data to detect genome-wide repair hotspots and coldspots and integrated additional omics data from various next-generation sequencing platforms to investigate the relationship between DNA damage, excision repair, epigenomic markers, 3D genome organization, and replication timing. To our best knowledge, this, for the first time, demonstrates the connection between repair kinetics and chromatin organization (46) with high resolution in human cell lines. The identified repair hotspots can serve as additional functional genomic features in studies of the human genome at the multidimensional level. We believe the framework and multi-omics data we present here can be useful to scientists both in the fields of chromatin dynamics and those interested in determinants of repair rates in the human genome. Our approach is applicable to studying the response to other types of DNA damage, including that

induced by drugs used in chemotherapy and thus has the potential for informing new
therapeutic strategies.

Using paired XR-seq and Damage-seq data, we provided additional empirical
evidence that the damage levels in the regions of the repair hotspots were not enriched,
in concordance with previous reports (14,15). Therefore, we attributed the observed
differences in repair patterns to heterogeneous repair efficiencies and not to damage
formation. It is, however, noteworthy that the depth of coverage by Damage-seq is an
order of magnitude lower than that by XR-seq. Specifically, the genome-wide average
total number of reads between the two replicates for (6-4)PP Damage-seq is 20.5 million,
while that average for (6-4)PP XR-seq is 181.8 million (Table S1). As such, it is possible
that we do not have enough sequencing depth to detect Damage-seq hotspots with high
sensitivity. The fact that a good proportion of Damage-seq reads get mapped to
heterochromatin regions further complicates analysis and lowers detection power. This
issue also persists for the recently developed adductSeq and FreqSeq, where a Poisson
model was adopted with a genomic average rate of 0.07 reads per pyrimidine dinucleotide
in fibroblasts (31). A Poisson distribution with a mean parameter under the null much less
than one could potentially suffer from overdispersion and inflated false positive rates.
Targeted Damage-seq offers a solution for both discovery and validation. However, as
with targeted DNA sequencing, the data can be extremely noisy due to targeting,
amplification, and sequencing biases and artifacts (47).

Using the genome-wide profile of DNA replication timing, we demonstrate that the
early-repair hotspots also tend to reside in early-replication domains. The inevitable effect
of replication on gene dosage and copy number could also have regulatory consequences
(48). Genes/regions that are duplicated early will be present at twice the copy number of
late replicating domains for most of the duration of S phase, increasing both the DNA
amount (as sister chromatids) and the transcriptional output. The ability to perform
excision repair for a greater fraction of the cell cycle has been shown to result in lower
mutation rates in the early-replication domains (49,50).

UV-induced DNA damage and excision repair have been linked with mutagenesis
and carcinogenesis. Recent studies have shown that mutation hotspots exhibit strong
increases in CPD formation efficacy (51-53) and that excision repair is attenuated in

transcription factor binding sites, leading to elevated somatic mutation rates (19,21). In
terms of repair kinetics, Adar et al. (17) showed that globally late-repaired regions are
associated with a higher level of cancer-linked mutations. To further investigate this in our
study context, we quantified the rates of somatic point mutations associated with
melanoma (1) in the repair hotspots and coldspots. However, due to the low number of
hotspots and coldspots, as well as their small genomic lengths, we found very few
overlapped with mutations and were thus underpowered to test for the differences in
mutation rates between different repair categories.

While the repair of (6-4)PP is predominantly carried out by global repair,
transcription-coupled repair plays an essential role in removing the more abundant and
less helix-distorting CPD adducts. In this study, we have intentionally focused on profiling
global repair hotspots for both (6-4)PP and CPD at early timepoints without the effect of
transcription-coupled repair. Genes exhibit both transcriptional dynamics over a time
course and biological fluctuations due to transcriptional bursting (54) at the same
timepoint. This confounds and complicates the analysis. To disentangle the effects of
global repair and transcriptional-coupled repair in identifying repair hotspots is an
unsolved yet challenging problem.

METHODS
Experimental methods
Cell culture and UV irradiation
Human NHF1 cells were obtained from W.K. Kaufmann (University of North Carolina,
Chapel Hill) (55) and cultured in Dulbecco's Modified Eagle Medium (DMEM) with 10%
FBS at 37 °C in a 5% CO_2 humidified chamber. For (6-4)PP XR-seq at 1 min and 2 min
timepoints, UV irradiation was performed as previously described (7,56). Briefly, the 80%
confluent NHF1 cells in one petri dish were irradiated for 20 s under a 250 nm UV lamp
(1 $J/m_2/s$) after removing the culture medium. 37 °C DMEM with 10% FBS medium was
immediately added into the petri dish, then the medium was poured off and the petri dish
was put on ice promptly at the end of 1 min or 2 min after UV irradiation. The time count
starts from the end of 20 s UV irradiation and ends at the timepoint when the petri dish is
put on ice. The cells were washed one time with ice-cold PBS before being harvested by

a cell scraper in 10 ml ice-cold PBS. In each replicate of (6-4)PP XR-seq experiment, 50

and 30 petri dishes (150mm x 15mm) containing NHF1 cells were treated one by one at

1 min and 2 min timepoints, respectively. Cell culture, UV treatment, and library

preparation for (6-4)PP XR-seq at 5 min, 20 min, 1 h, 2 h, and 4 h, and CPD XR-seq at

12 min were performed in previous studies (7,10,17). For *in vivo* excision assay, UV

irradiation was performed as aforementioned, and 10 and 5 petri dishes (150mm x 15mm)

containing NHF1 cells were used at 0 min and 2 min timepoints respectively.

Excision assay

The *in vivo* excision assay was performed as described (24,57). Following UV irradiation,

the excision products were isolated by gentle cell lysis and nonchromatin fraction

separation and purified by TFIIH immunoprecipitation. The purified excision products

were then 3' radiolabeled by terminal deoxynucleotidyl transferase and [α-$_{32}$P]-3'-dATP,

and resolved in a 10% denaturing acrylamide gel. Ten and five petri dishes (150mm x

15mm) of NHF1 cells were used at 0 min and 2 min, respectively.

XR-seq library preparation and sequencing

XR-seq libraries were prepared as described in the previous protocol (57). Briefly, the

excision products were isolated by TFIIH immunoprecipitation following gentle cell lysis

and non-chromatin fraction separation, and ligated with adaptors. The ligated excision

products were then further purified by immunoprecipitation with anti-(6-4)PP antibody and

repaired by (6-4)PP photolyase before the library amplification by PCR. Libraries were

sequenced on an Illumina HiSeq 4000 platform.

Data collection

(6-4)PP XR-seq data at 5 min, 20 min, 1 h, 2 h, and 4 h were downloaded from the Gene

Expression Omnibus (GEO) with accession numbers GSE67941 (7) and GSE76391 (17).

CPD XR-seq data at 12 min were downloaded from GEO with accession number

GSE138846 (10). CPD and (6-4)PP damage data of NHF1 by Damage-seq were

downloaded from GEO with accession number GSE98025 (15); CPD damage data of

NHF1 by CPD-seq were downloaded from GEO with accession number GSM2772322

and GSM2772323 (19); CPD damage data of human primary fibroblast by adductSeq were downloaded from GEO with accession number GSM4073616 and GSM4073634 (31). Hyper-hotspots for UV-induced CPD damage in primary human fibroblasts were downloaded from Premi et al. (31). NHDF H3K4me1 (ENCODE Data Coordination Center accession number ENCSR000ARV), H3K4me3 (accession number ENCSR000DPR), H3K27ac (accession number ENCSR000APN), H3K27me3 (accession number ENCSR000APO), H3K9me3 (accession number ENCSR000ARX), and DNase-seq (accession number ENCSR000EMP) data were downloaded from the ENCODE portal (29). NHLF chromatin state segmentation results by chromHMM were downloaded from UCSC accession number wgEncodeEH000792 (34). Hi-C data of IMR90 were downloaded from GEO with accession number GSE43070 (35) and from https://bioconductor.org/packages/HiCDataHumanIMR90/ (36). The list of annotated super-enhancers in IMR90 was downloaded from the Roadmap Epigenomics Consortium (39). Genomic categories of replication timing from Repli-Seq data of IMR90 (41) were downloaded from GSE53984 (42).

Bioinformatic and statistical analysis

XR-seq bioinformatic pre-processing

For XR-seq, cutadapt (58) was used to trim reads with adaptor sequence TGGAATTCTCGGGTGCCAAGGAACTCCAGTNNNNNNNACGATCTCGTATGCCGTCT TCTGCTTG at the 3' end and discard untrimmed reads. BWA (59) was used for alignment of single-end short reads. Unmapped reads and reads that map to multiple locations with the same alignment quality were removed using Samtools (60). Post-alignment filtering steps were adopted using Rsamtools (http://bioconductor.org/packages/Rsamtools/). Specifically, if multiple reads share the same 5' and 3' end coordinates, we keep only one to perform deduplication. We also only keep reads that have mapping quality greater than 20 and are of lengths 21 bp to 31 bp.

Gene-level quantification of excision repair

Reads from the TS and NTS strands were separated using known gene annotations for hg19 by ENSEMBL. We use RPKM for within-sample normalization for the XR-seq data.

To perform gene-level quantification and downstream analysis including segmented
regression, we adopted a stringent quality control procedure and only retained genes that:
(i) had at least ten TT or TC dinucleotides from either TS or NTS; (ii) were less than 300Kb;
and (iii) had at least ten reads in total across all XR-seq samples. In addition, we took the
ratio of the reads from the TS and the NTS [TS/(TS+NTS)] to remove biases and artifact
that are shared between the two DNA strands, i.e., library size, gene length, and other
gene-specific biases, such as sequencing bias and antibody pulldown efficiency, etc. The
ratio is bound between 0 and 1 and sheds light upon how transcription-coupled repair and
global repair interplay (Figure S3).

Identification of repair hotspots and coldspots
We started by segmenting the human reference genome into consecutive bins of 50 bp
long. We then calculated the observed depth of coverage per bin by XR-seq, separating
the plus-strand reads (+) and the minus-strand reads (-). To mitigate the effect of library
size/sequencing depth, we downsampled the reads in each sample to 7.7 million without
replacement. To identify repair hotspots and coldspot, we set a threshold on the number
of read counts per genomic bin in the 1 min and 4 h samples. Specifically, to identify (6-
4)PP repair hotspots, we require at least 15 reads mapped in both replicates at 1min and
at most 5 reads mapped in both replicates at 4 h. The read count threshold is relaxed for
the identification of coldspots, which have a smaller number compared to the hotspots.
For CPD repair, to avoid complicatedness due to transcription-coupled repair at later
timepoints, we focused on CPD repair hotspots only.
In addition to the thresholding approach, we adopted a more rigorous cross-
sample Poisson log linear model (27,28) for data normalization. Specifically, we denote
Y as the observed repair matrix, with row i corresponding to the ith genomic bin and
column j corresponding to the jth sample. The "null" model, which reflects the expected
coverage when there is no biologically relevant repair enrichment, is

$$Y_{ij} \sim \text{Poisson}(\lambda_{ij}), \ \ \lambda_{ij} = N_j \beta_i f_j(TC_i),$$

where N_j is the total number of mapped reads for sample j (fixed for downsampled data),
β_i reflects the bin-specific bias due to library preparation and sequencing bias, and
$f_j(TC_i)$ is the sample-specific bias due to TC (thymine and cytosine) content for

damage/repair. The goal of fitting the null model to the data is to estimate the various

sources of biases, which can then be used for normalization. We adopt a robust iterative

maximum-likelihood algorithm (28) for estimating the parameters of the null model. Plus

and minus strands are analyzed separately.

Given a first-pass of the calling algorithms, we identified strong repair hotspots in

pericentromeric regions, which were collapsed repeats annotated as unique sequences

in the reference genome (e.g., ribosomal DNA (9)). It is important to exclude artifacts as

stringently as possible, and thus we undertook an additional quality control step. "Blacklist"

bins,   including   segmental   duplication   regions

(http://humanparalogy.gs.washington.edu/build37/data/GRCh37GenomicSuperDup.tab),

gaps in reference assembly from telomere, centromere, and/or heterochromatin regions

(https://gist.github.com/leipzig/6123703), and repeating elements by RepeatMasker

(https://genome.ucsc.edu/cgi-bin/hgTrackUi?g=rmsk) are masked in downstream

analysis.

Hi-C data analysis

We adopted the Hi-C data of human fibroblast cell line IMR90 (35,36) to investigate the

relationship between identified repair hotspots and the 3D genome structure. We took the

raw contact matrix with 40 kb resolution as input and detected FIREs, which play

important roles in transcriptional regulations, across the entire genome using FIREcaller

(37). To further investigate whether these repair hotspots are involved in functional

chromatin looping between regulatory elements and their target genes, we adopted the

Fit-Hi-C approach (40) to identify long-range chromatin interactions on all 40 kb bin pairs

within a maximal 3 MB region. The interactions with p-value < 2.31e-11 were considered

as statistically significant (61).

DATA AND CODE AVAILABILITY

The data reported in this paper have been deposited in GEO with accession number

GSE148303.   Scripts   used   in   this   paper   are   available   at

https://github.com/yuchaojiang/damage_repair.

COMPETING INTEREST STATEMENT

The authors declare no competing interests.

ACKNOWLEDGEMENTS

This work was supported by NIH Grants R35 GM118102 (to A.S.), R01 ES027255 (to A.S.), R35 GM138342 (to Y.J.), K99 ES030015 (to W.L.), and a pilot award from the UNC Computational Medicine Program (to Y.J.). We thank the Sancar Lab members and Drs. Sheera Adar, Ming Hu, and Jeremy Simon for useful comments and feedback.

AUTHOR CONTRIBUTIONS

AS envisioned and initiated the study, while LW and LAL-B performed the experiments. YJ devised the analytical framework, and all authors executed the data analysis. YJ wrote the manuscript, with contributions from WL on experimental methods. The manuscript was further edited and approved by all authors.

REFERENCES

1. Hodis, E., Watson, I.R., Kryukov, G.V., Arold, S.T., Imielinski, M., Theurillat, J.P., Nickerson, E., Auclair, D., Li, L., Place, C. *et al.* (2012) A landscape of driver mutations in melanoma. *Cell*, **150**, 251-263.

2. Martincorena, I., Roshan, A., Gerstung, M., Ellis, P., Van Loo, P., McLaren, S., Wedge, D.C., Fullam, A., Alexandrov, L.B., Tubio, J.M. *et al.* (2015) Tumor evolution. High burden and pervasive positive selection of somatic mutations in normal human skin. *Science*, **348**, 880-886.

3. Sancar, A. (2016) Mechanisms of DNA repair by photolyase and excision nuclease (Nobel Lecture). *Angewandte Chemie International Edition*, **55**, 8502-8527.

4. Hanawalt, P.C. and Spivak, G. (2008) Transcription-coupled DNA repair: two decades of progress and surprises. *Nat Rev Mol Cell Biol*, **9**, 958-970.

5. Mellon, I., Spivak, G. and Hanawalt, P.C. (1987) Selective removal of transcription-blocking DNA damage from the transcribed strand of the mammalian DHFR gene. *Cell*, **51**, 241-249.

6. Kusakabe, M., Onishi, Y., Tada, H., Kurihara, F., Kusao, K., Furukawa, M., Iwai, S., Yokoi, M., Sakai, W. and Sugasawa, K. (2019) Mechanism and regulation of DNA damage recognition in nucleotide excision repair. *Genes Environ*, **41**, 2.

7. Hu, J., Adar, S., Selby, C.P., Lieb, J.D. and Sancar, A. (2015) Genome-wide analysis of human global and transcription-coupled excision repair of UV damage at single-nucleotide resolution. *Genes Dev*, **29**, 948-960.

8.   Akkose, U., Kaya, O., Lindsey-Boltz, L., Karagoz, Z., Brown, A.D., Larsen, P.A., Yoder, A.D.,
Sancar, A. and Adebali, O. (2020) Comparative analyses of two primate species diverged
by more than 60 million years show different rates but similar distribution of genome-
wide UV repair events. *bioRxiv*.

9.   Yang, Y., Hu, J., Selby, C.P., Li, W., Yimit, A., Jiang, Y. and Sancar, A. (2019) Single-
nucleotide resolution analysis of nucleotide excision repair of ribosomal DNA in humans
and mice. *J Biol Chem*, **294**, 210-217.

10.   Deger, N., Yang, Y., Lindsey-Boltz, L.A., Sancar, A. and Selby, C.P. (2019) Drosophila, which
lacks canonical transcription-coupled repair proteins, performs transcription-coupled
repair. *J Biol Chem*, **294**, 18092-18098.

11.   Li, W., Adebali, O., Yang, Y., Selby, C.P. and Sancar, A. (2018) Single-nucleotide resolution
dynamic repair maps of UV damage in Saccharomyces cerevisiae genome. *Proc Natl Acad*
Sci U S A, **115**, E3408-E3415.

12.   Adebali, O., Chiou, Y.Y., Hu, J., Sancar, A. and Selby, C.P. (2017) Genome-wide
transcription-coupled repair in Escherichia coli is mediated by the Mfd translocase. *Proc*
Natl Acad Sci U S A, **114**, E2116-E2125.

13.   Oztas, O., Selby, C.P., Sancar, A. and Adebali, O. (2018) Genome-wide excision repair in
Arabidopsis is coupled to transcription and reflects circadian gene expression patterns.
Nat Commun, **9**, 1503.

14.   Hu, J., Lieb, J.D., Sancar, A. and Adar, S. (2016) Cisplatin DNA damage and repair maps of
the human genome at single-nucleotide resolution. *Proc Natl Acad Sci U S A*, **113**, 11507-
11512.

15.   Hu, J., Adebali, O., Adar, S. and Sancar, A. (2017) Dynamic maps of UV damage formation
and repair for the human genome. *Proc Natl Acad Sci U S A*, **114**, 6758-6763.

16.   Yimit, A., Adebali, O., Sancar, A. and Jiang, Y. (2019) Differential damage and repair of
DNA-adducts induced by anti-cancer drug cisplatin across mouse organs. *Nat Commun*,
10, 309.

17.   Adar, S., Hu, J., Lieb, J.D. and Sancar, A. (2016) Genome-wide kinetics of DNA excision
repair in relation to chromatin state and mutagenesis. *Proc Natl Acad Sci U S A*, **113**,
E2124-2133.

18.   Mao, P., Smerdon, M.J., Roberts, S.A. and Wyrick, J.J. (2016) Chromosomal landscape of
UV damage formation and repair at single-nucleotide resolution. *Proc Natl Acad Sci U S A*,
113, 9057-9062.

19.   Mao, P., Brown, A.J., Esaki, S., Lockwood, S., Poon, G.M.K., Smerdon, M.J., Roberts, S.A.
and Wyrick, J.J. (2018) ETS transcription factors induce a unique UV damage signature
that drives recurrent mutagenesis in melanoma. *Nat Commun*, **9**, 2626.

20.   Garcia-Nieto, P.E., Schwartz, E.K., King, D.A., Paulsen, J., Collas, P., Herrera, R.E. and
Morrison, A.J. (2017) Carcinogen susceptibility is regulated by genome architecture and
predicts cancer mutagenesis. *EMBO J*, **36**, 2829-2843.

21.   Sabarinathan, R., Mularoni, L., Deu-Pons, J., Gonzalez-Perez, A. and Lopez-Bigas, N. (2016)
Nucleotide excision repair is impaired by binding of transcription factors to DNA. *Nature*,
532, 264-267.

22.   Poulos, R.C., Thoms, J.A.I., Guan, Y.F., Unnikrishnan, A., Pimanda, J.E. and Wong, J.W.H.
(2016) Functional Mutations Form at CTCF-Cohesin Binding Sites in Melanoma Due to
Uneven Nucleotide Excision Repair across the Motif. *Cell reports*, **17**, 2865-2872.

23.   Reardon, J.T. and Sancar, A. (2003) Recognition and repair of the cyclobutane thymine
dimer, a major cause of skin cancers, by the human excision nuclease. *Genes Dev*, **17**,
2539-2551.

24.   Hu, J., Choi, J.H., Gaddameedhi, S., Kemp, M.G., Reardon, J.T. and Sancar, A. (2013)
Nucleotide excision repair in human cells: fate of the excised oligonucleotide carrying
DNA damage in vivo. *J Biol Chem*, **288**, 20918-20926.

25.   Deger, N., Yang, Y., Lindsey-Boltz, L.A., Sancar, A. and Selby, C.P. (2019) Drosophila, which
lacks canonical transcription-coupled repair proteins, performs transcription-coupled
repair. *Journal of Biological Chemistry*, **294**, 18092-18098.

26.   Jolliffe, I.T. and Cadima, J. (2016) Principal component analysis: a review and recent
developments. *Philos Trans A Math Phys Eng Sci*, **374**, 20150202.

27.   Witten, D.M. (2011) Classification and clustering of sequencing data using a Poisson
model. *The Annals of Applied Statistics*, **5**, 2493-2518.

28.   Jiang, Y., Oldridge, D.A., Diskin, S.J. and Zhang, N.R. (2015) CODEX: a normalization and
copy number variation detection method for whole exome sequencing. *Nucleic Acids Res*,
43, e39.

29.   Consortium, E.P. (2012) An integrated encyclopedia of DNA elements in the human
genome. *Nature*, **489**, 57-74.

30.   Bailey, T.L., Boden, M., Buske, F.A., Frith, M., Grant, C.E., Clementi, L., Ren, J., Li, W.W.
and Noble, W.S. (2009) MEME SUITE: tools for motif discovery and searching. *Nucleic
Acids Res*, **37**, W202-208.

31.   Premi, S., Han, L., Mehta, S., Knight, J., Zhao, D., Palmatier, M.A., Kornacker, K. and Brash,
D.E. (2019) Genomic sites hypersensitive to ultraviolet radiation. *Proc Natl Acad Sci U S A*,
116, 24196-24205.

32.   Han, C., Srivastava, A.K., Cui, T., Wang, Q.E. and Wani, A.A. (2016) Differential DNA lesion
formation and repair in heterochromatin and euchromatin. *Carcinogenesis*, **37**, 129-138.

33.   Hauer, M.H. and Gasser, S.M. (2017) Chromatin and nucleosome dynamics in DNA
damage and repair. *Genes Dev*, **31**, 2204-2221.

34.   Ernst, J. and Kellis, M. (2012) ChromHMM: automating chromatin-state discovery and
characterization. *Nat Methods*, **9**, 215-216.

35.   Jin, F., Li, Y., Dixon, J.R., Selvaraj, S., Ye, Z., Lee, A.Y., Yen, C.A., Schmitt, A.D., Espinoza, C.A.
and Ren, B. (2013) A high-resolution map of the three-dimensional chromatin
interactome in human cells. *Nature*, **503**, 290-294.

36.   Dixon, J.R., Selvaraj, S., Yue, F., Kim, A., Li, Y., Shen, Y., Hu, M., Liu, J.S. and Ren, B. (2012)
Topological domains in mammalian genomes identified by analysis of chromatin
interactions. *Nature*, **485**, 376-380.

37.   Crowley, C., Yang, Y., Qiu, Y., Hu, B., Won, H., Ren, B., Hu, M. and Li, Y. (2019) FIREcaller:
an R package for detecting frequently interacting regions from Hi-C data. *bioRxiv*, 619288.

38.   Schmitt, A.D., Hu, M., Jung, I., Xu, Z., Qiu, Y., Tan, C.L., Li, Y., Lin, S., Lin, Y., Barr, C.L. *et al.*
(2016) A Compendium of Chromatin Contact Maps Reveals Spatially Active Regions in the
Human Genome. *Cell Rep*, **17**, 2042-2059.

39.   Consortium, R.E., Kundaje, A., Meuleman, W., Ernst, J., Bilenky, M., Yen, A., Heravi-
Moussavi, A., Kheradpour, P., Zhang, Z., Wang, J. *et al.* (2015) Integrative analysis of 111
reference human epigenomes. *Nature*, **518**, 317-330.

40.   Ay, F., Bailey, T.L. and Noble, W.S. (2014) Statistical confidence estimation for Hi-C data
reveals regulatory chromatin contacts. *Genome Res*, **24**, 999-1011.

41.   Hansen, R.S., Thomas, S., Sandstrom, R., Canfield, T.K., Thurman, R.E., Weaver, M.,
Dorschner, M.O., Gartler, S.M. and Stamatoyannopoulos, J.A. (2010) Sequencing newly
replicated DNA reveals widespread plasticity in human replication timing. *Proc Natl Acad*
Sci U S A, **107**, 139-144.

42.   Liu, F., Ren, C., Li, H., Zhou, P., Bo, X. and Shu, W. (2016) De novo identification of
replication-timing domains in the human genome by deep learning. *Bioinformatics*, **32**,
641-649.

43.   Li, W., Liu, W., Kakoki, A., Wang, R., Adebali, O., Jiang, Y. and Sancar, A. (2019) Nucleotide
excision repair capacity increases during differentiation of human embryonic carcinoma
cells into neurons and muscle cells. *J Biol Chem*, **294**, 5914-5922.

44.   Bacher, R., Leng, N., Chu, L.F., Ni, Z., Thomson, J.A., Kendziorski, C. and Stewart, R. (2018)
Trendy: segmented regression analysis of expression dynamics in high-throughput
ordered profiling experiments. *BMC Bioinformatics*, **19**, 380.

45.   Thomas, P.D., Campbell, M.J., Kejariwal, A., Mi, H., Karlak, B., Daverman, R., Diemer, K.,
Muruganujan, A. and Narechania, A. (2003) PANTHER: a library of protein families and
subfamilies indexed by function. *Genome Res*, **13**, 2129-2141.

46.   van Eijk, P., Nandi, S.P., Yu, S., Bennett, M., Leadbitter, M., Teng, Y. and Reed, S.H. (2019)
Nucleosome remodeling at origins of global genome-nucleotide excision repair occurs at
the boundaries of higher-order chromatin structure. *Genome Res*, **29**, 74-84.

47.   Jiang, Y., Wang, R., Urrutia, E., Anastopoulos, I.N., Nathanson, K.L. and Zhang, N.R. (2018)
CODEX2: full-spectrum copy number variation detection by high-throughput DNA
sequencing. *Genome Biol*, **19**, 202.

48.   Rhind, N. and Gilbert, D.M. (2013) DNA replication timing. *Cold Spring Harb Perspect Biol*,
630   **5**, a010132.

631   49.   Stamatoyannopoulos, J.A., Adzhubei, I., Thurman, R.E., Kryukov, G.V., Mirkin, S.M. and
632   Sunyaev, S.R. (2009) Human mutation rate associated with DNA replication timing. *Nat*
633   *Genet*, **41**, 393-395.

634   50.   Lang, G.I. and Murray, A.W. (2011) Mutation rates across budding yeast chromosome VI
635   are correlated with replication timing. *Genome Biol Evol*, **3**, 799-811.

636   51.   Elliott, K., Bostrom, M., Filges, S., Lindberg, M., Van den Eynden, J., Stahlberg, A., Clausen,
637   A.R. and Larsson, E. (2018) Elevated pyrimidine dimer formation at distinct genomic bases
638   underlies promoter mutation hotspots in UV-exposed cancers. *PLoS Genet*, **14**, e1007849.

639   52.   Pich, O., Muinos, F., Sabarinathan, R., Reyes-Salazar, I., Gonzalez-Perez, A. and Lopez-
640   Bigas, N. (2018) Somatic and Germline Mutation Periodicity Follow the Orientation of the
641   DNA Minor Groove around Nucleosomes. *Cell*, **175**, 1074-1087 e1018.

642   53.   Brown, A.J., Mao, P., Smerdon, M.J., Wyrick, J.J. and Roberts, S.A. (2018) Nucleosome
643   positions establish an extended mutation signature in melanoma. *PLoS Genet*, **14**,
644   e1007823.

645   54.   Jiang, Y., Zhang, N.R. and Li, M. (2017) SCALE: modeling allele-specific gene expression by
646   single-cell RNA sequencing. *Genome Biol*, **18**, 74.

647   55.   Heffernan, T.P., Simpson, D.A., Frank, A.R., Heinloth, A.N., Paules, R.S., Cordeiro-Stone, M.
648   and Kaufmann, W.K. (2002) An ATR- and Chk1-dependent S checkpoint inhibits replicon
649   initiation following UVC-induced DNA damage. *Mol Cell Biol*, **22**, 8552-8561.

650   56.   Li, W., Hu, J., Adebali, O., Adar, S., Yang, Y., Chiou, Y.Y. and Sancar, A. (2017) Human
651   genome-wide repair map of DNA damage caused by the cigarette smoke carcinogen
652   benzo[a]pyrene. *Proc Natl Acad Sci U S A*, **114**, 6752-6757.

653   57.   Hu, J., Li, W., Adebali, O., Yang, Y., Oztas, O., Selby, C.P. and Sancar, A. (2019) Genome-
654   wide mapping of nucleotide excision repair with XR-seq. *Nat Protoc*, **14**, 248-282.

655   58.   Martin, M. (2011) Cutadapt removes adapter sequences from high-throughput
656   sequencing reads. *EMBnet. journal*, **17**, 10-12.

657   59.   Li, H. and Durbin, R. (2009) Fast and accurate short read alignment with Burrows-Wheeler
658   transform. *Bioinformatics*, **25**, 1754-1760.

659   60.   Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
660   Durbin, R. and Genome Project Data Processing, S. (2009) The Sequence Alignment/Map
661   format and SAMtools. *Bioinformatics*, **25**, 2078-2079.

662   61.   Giusti-Rodríguez, P., Lu, L., Yang, Y., Crowley, C.A., Liu, X., Juric, I., Martin, J.S., Abnousi,
663   A., Allred, S.C. and Ancalade, N. (2019) Using three-dimensional regulatory chromatin
664   interactions from adult and fetal cortex to interpret genetic results for psychiatric
665   disorders and cognitive traits. *bioRxiv*, 406330.

 FIGURE & TABLE LEGENDS

Figure 1. Excision repair kinetics and early-repair hotspots in normal human

fibroblasts. (A) Experimental design to measure DNA damage and excision repair of the

UV-induced (6-4)PP and CPD across different timepoints. (B) Detection of excision

products at 0 min and 2 min timepoints *in vivo*. Following UV irradiation, the excised

oligonucleotides were purified by TFIIH immunoprecipitation, radiolabeled, and resolved

in a 10% sequencing gel. DNA excision products of (6-4)PP can be detected as early as

2 min upon damage induction. (C) Principal component analysis of genome-wide excision

repair as measured by XR-seq shows repair kinetics across different timepoints. Between

1 min and 5 min, excised oligonucleotides were not degraded, and thus XR-seq measured

cumulative repair. (D) Each row is a sample, and each column is a specific total number

of reads per genomic bin. The color in the heatmap corresponds to the log counts of the

number of bins with specific read depths. Early-repair hotspots exist in samples collected

at early timepoints.

Figure 2. Distribution of DNA damage, repair, and epigenomic markers at identified

repair hotspots and coldspots. XR-seq and Damage-seq data are shown for both

strands, marked with + and -. Epigenetic data from ChIP-seq of histone modifications and

DNase-seq are plotted on the same scale for cross comparison. Read count data

normalized by sequencing depth are visualized in the Integrative Genomics Viewer. (A)

A (6-4)PP repair hotspot from chr2. (B) A (6-4)PP repair coldspot from chr1. (C) A CPD

repair hotspot from chr1. All examples are from intronic gene regions overlapping

annotated enhancers. Zoomed-in view of canonical sequences is overlaid in the bottom,

with the damage/repair sites shown in dashed boxes.

Figure 3. Repair hotspots are not hypersensitive to ultraviolet radiation. There is no

enriched DNA damage, as measured by Damage-seq, adductSeq, and CPD-seq, at the

identified repair hotspots for (6-4)PP and CPD. (A) Read counts for DNA damage are

computed in repair hotspots, coldspots, and random spots. The regions corresponding to

the different repair categories are extended at both ends for 20bp and 500bp, respectively,

to account for the shallow sequencing depth by quantifying DNA damage.

Figure 4. Genome-wide repair hotspots are associated with epigenomic markers.
Chromatin accessibility (DNase I hypersensitivity) is higher for repair hotspots and lower
for coldspots. Repair hotspots are also characterized by higher ChIP-seq signals for
H3K4me1, H3K4me3, and H3K27ac, markers for gene activations.

Figure 5. Genome-wide repair hotspots are enriched for enhancers. Annotations for
(A) chromatin states, (B) CpG islands, and (C) genic/intergenic regions are shown for
repair hotspots for (6-4)PP (top) and CPD (bottom), separated by strands. Repair
hotspots are enriched for enhancers and promoters, which are in open-chromatin regions.
Repair hotspots are also enriched in inter-CpG and intronic regions, both of which are
AT-rich.

Figure 6. Genome-wide repair hotspots overlap with FIREs and super-enhancers
identified by Hi-C. FIREs and super-enhancers were identified and annotated using Hi-
C data from human fibroblasts. Repair hotspots overlap with (A) FIREs and (B) super-
enhancers with significantly higher proportions compared to the genome-wide averages.
The solid density curves are generated from bootstrapping different regions along the
genome as the null case; the dashed vertical lines are the observed proportions for the
repair hotspots and coldspots. (C) Repair hotspots have a significantly higher number of
significant interactions, identified by Hi-C. (D) Two examples of (6-4)PP repair hotspots
(chr8:13224201-13224300 and chr14:61994601-61994700) that overlap with both FIREs
and super-enhancers, and loop to different regions of the genome. Hi-C data has low
resolution, and the significant interactions are drawn from the center of each bin, which
does not exactly overlap with the identified hotspot shown in red.

Figure 7. Genome-wide repair hotspots are enriched for early-replication domains.
Replication timing domains were identified using Repli-Seq data. Compared to genome-
wide average (random), there is a significantly higher proportion of the early-repair
hotspots located in the early-replication domains.

Figure 1

Figure 2

Figure 3

Figure 4

(A)

(B)

(C)

Figure 5

Figure 6

Figure 7