Nucleotide excision repair hotspots of UV-induced DNA damage in the human genome

Yuchao Jiang[1,2,3,*,#], Wentao Li[4,*], Laura A Lindsey-Boltz[4,*], Yuchen Yang[2], Yun Li[1,2,5], Aziz Sancar[3,4,#]

1   Department of Biostatistics, Gillings School of Global Public Health, University of North Carolina, Chapel Hill, NC 27599, USA.
2   Department of Genetics, School of Medicine, University of North Carolina, Chapel Hill, NC 27599, USA.
3   Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, NC 27599, USA.
4   Department of Biochemistry and Biophysics, School of Medicine, University of North Carolina, Chapel Hill, NC 27599, USA.
5   Department of Computer Science, College of Arts and Sciences, University of North Carolina, Chapel Hill, NC 27599, USA.

∗   These authors contributed equally.
To whom correspondence should be addressed. Email: yuchaoj@email.unc.edu, aziz_sancar@med.unc.edu.

ABSTRACT

We recently developed high-throughput sequencing approaches, eXcision Repair sequencing (XR-seq) and Damage-seq, to generate genome-wide mapping of DNA excision repair and damage formation, respectively, with single-nucleotide resolution. Here, we used time-course XR-seq data to profile UV-induced excision repair dynamics, paired with Damage-seq data to quantify the overall induced DNA damage. We identified genome-wide repair hotspots exhibiting high-level nucleotide excision repair immediately after UV irradiation. We show that such repair hotspots do not result from hypersensitivity to DNA damage, and are thus not damage hotspots. We find that the earliest repair occurs preferentially in promoters and enhancers from open-chromatin regions. The repair hotspots are also significantly enriched for frequently interacting regions and super-enhancers, both of which are themselves hotspots for local chromatin interactions. Further interrogation of chromatin organization to include DNA replication timing allows us to conclude that early-repair hotspots are enriched for early-replication domains. Collectively, we report genome-wide early-repair hotspots of UV-induced damage, in association with chromatin states and epigenetic compartmentalization of the human genome.

INTRODUCTION

UV in sunlight is a known mutagen and causative agent of skin cancer that induces DNA lesions as cyclobutane pyrimidine dimers (CPDs) and pyrimidine-pyrimidone (6-4) photoproducts [(6-4)PPs] (Hodis et al. 2012; Martincorena et al. 2015). In humans, both damage types are solely repaired by nucleotide excision repair (excision repair), which removes DNA lesions by dual incisions bracketing modified bases and fills and seals the resulting gap by DNA synthesis and ligation (Reardon and Sancar 2005). Excision repair consists of two pathways, global repair and transcription-coupled repair (Mellon et al. 1987; Hanawalt and Spivak 2008), that differ primarily in the damage recognition step (Kusakabe et al. 2019). For UV-induced DNA damage, CPD repair is highly associated with transcription, specifically with the transcribed strand, while (6-4)PP repair is distributed uniformly distributed throughput the genome (Hu et al. 2015).

Recently, we developed a high-throughput approach, excision repair sequencing (XR-seq), to isolate the oligonucleotides excised by excision repair and subject them to next-generation sequencing (Hu et al. 2015). This method has allowed for genome-wide mapping of UV-induced excision repair with single-nucleotide resolution in human (Hu et al. 2015), Lemur (Akkose et al. 2020), mouse (Yang et al. 2019), *Drosophila melanogaster* (Deger et al. 2019), *Saccharomyces cerevisiae* (Li et al. 2018), *Escherichia coli* (Adebali et al. 2017), and *Arabidopsis thaliana* (Oztas et al. 2018). We further developed another next-generation sequencing method, Damage-seq (Hu et al. 2016), to generate genome-wide mapping of UV damage formation with single-nucleotide resolution (Hu et al. 2017).

Using the combined UV damage maps and repair maps, it has been shown that the induced DNA damage is uniformly distributed throughout the genome and that the overall effect of damage in the genome is primarily driven, not by damage formation, but by repair efficiency (Hu et al. 2016; Hu et al. 2017). Indeed, different repair efficiencies across the genome have been previously reported, which are impacted by multiple factors, including transcription, chromatin state and structure, regulatory protein binding to DNA, and posttranscriptional modification of histones (Hu et al. 2015; Adar et al. 2016; Mao et al. 2016; Poulos et al. 2016; Sabarinathan et al. 2016; Garcia-Nieto et al. 2017; Hu et al. 2017; Mao et al. 2018; Yimit et al. 2019). Existing studies, however, have been focused

on profiling repair dynamics over a time course – 4 h for (6-4)PP and 48 h for CPD.

Genomic regions that harbor high-level early repair shortly after damage formation have

not been studied, nor have their associated genomic and epigenomic characteristics been

systematically explored.

Here, we performed XR-seq at as early as 1 min for (6-4)PP and 12 min for CPD

following UV irradiation to identify such repair hotspots. We systematically characterized

the identified hotspots using additional high-throughput sequencing data that measures

DNA damage formation, DNase I hypersensitivity, histone modifications, 3D chromatin

interactions, and DNA replication timing. These extensive links between chromatin states

and organization, cell cycle, DNA damage, and excision repair at specific genomic sites

facilitate better understanding of mutagenesis and carcinogenesis and have the potential

to improve prevention strategies and develop novel therapeutics for human diseases such

as cancer.

RESULTS

Genome-wide profiling of DNA excision pair kinetics through ordered high-

throughput experiments

In this work, we present an experimental and analytical framework where we

systematically assay DNA excision repair of UV-induced DNA damage over a time course.

Figure 1A gives an outline of the experimental design. Specifically, we adopted XR-seq

(Hu et al. 2015) to measure repair of (6-4)PPs at 1 min, 2 min, 5 min, 20 min, 1 h, 2 h,

and 4 h and CPDs at 12 min in normal human skin fibroblasts (NHF1) after 20J/m^2 UV

treatment. The (6-4)PP XR-seq experiments from 1 min to 4 h were all performed with

two biological replicates, with greater than 0.99 correlation coefficients between each pair

of replicates (Figure S1). Quality control procedures show that all samples have good

data quality with expected read length distributions and enrichments of TT and/or TC

dinucleotides at the damage sites (Figure S2). To investigate the interplay between DNA

damage and repair, we additionally adopted Damage-seq data (Hu et al. 2017) to profile

the two types of UV-induced damage at 0 timepoint after UV irradiation. Refer to the

Methods section for details on library preparation, sequencing, and bioinformatic analysis

and Table S1 for detailed sample information.

The bulk repair pathways and kinetics of these two types of DNA damage are
known to be quite different (Reardon and Sancar 2003; Hu et al. 2013). The global repair
efficiently removes (6-4)PPs, with most of the damage excised within 4 h of UV treatment.
We provide additional empirical evidence that the repair levels of (6-4)PP from the
transcribed strand (TS) and the non-transcribed strand (NTS) are on par across all
timepoints (Figure S3A-B). In contrast, CPDs are recognized primarily in a transcription-
coupled manner and their complete repair requires 12 to 48 h, depending on the UV dose.
In our experiment, however, given the sampling timepoint as early as 12 min for CPDs,
transcription-coupled repair has not contributed much, if any at all, to the repair of CPDs
(Figure S3C). This is of key importance so that our analysis is unbiasedly focused on
global repair of (6-4)PP at any timepoints and of CPD at the early 12 min timepoint,
without transcription as a confounder. Refer to the Discussion section for more details.
In this study, we focus on characterizing global repair kinetics and identifying early-
repair hotspots. *In vivo* excision assay by gel autoradiography at 0 min and 2 min reveals
that the primary excision products repair can be detected as early as 2 min after UV
treatment (Figure 1B). Using genome-wide repair data by XR-seq, we performed principal
component analysis (PCA) on the top 2,000 highly variable genes to generate a low-
dimensional representation of the (6-4)PP repair data (Figure 1C). Since transcription-
coupled repair does not contribute to the repair of (6-4)PPs, the PCA plots do not differ
between the TS and NTS repair, except for reverted signs of the eigenvectors. Importantly,
a repair trajectory on the principal component space can be reconstructed, which lines up
well with the timepoints, suggesting differential repair kinetics over the time course (Figure
1C).
103

Identification of early-repair hotspots in normal human fibroblasts

Identification of early-repair hotspots in normal human fibroblasts
We developed a computational framework to identify genome-wide early-repair hotspots
using time-course XR-seq data. Refer to the Methods section for details. Briefly, we
segmented the genome into consecutive bins of 50 bp long and identified bins that
showed significantly higher number of reads at earlier or later timepoints (Figure S4). We
define such genomic bins as repair hotspots and coldspots, respectively. Figure 1D
shows the distributions of read counts per genomic bin across all (6-4)PP samples and

111 we note enrichment of both early repair at 1 min and late repair at 4 h, corresponding to
112 repair hotspots and coldspots, respectively. For repair of (6-4)PPs, we identified on the
113 genome-wide scale 175 and 156 repair hotspots from the plus and minus strand (Table
114 S2) and 48 and 57 repair coldspots from the plus and minus strand (Table S3),
115 respectively. For repair of CPDs, we focused on repair hotspots at an early timepoint to
116 negate the global impact of transcription, and identified 99 and 93 repair hotspots from
117 the plus and minus strands, respectively (Table S4). We could not search for intrinsic
118 CPD coldspots, because 30 min after UV irradiation transcription-coupled repair becomes
119 an important contributor to the repair profile.

120  The repair hotspots and coldspots reported are scattered across the entire human
121 genome (Figure S5). XR-seq signals from examples of repair hotspots and coldspots are
122 separated by strand and plotted across all timepoints in Figure 2. We also include
123 epigenomic signals by DNase-seq, ChIP-seq from ENCODE (ENCODE Project
124 Consortium 2012), and Damage-seq signals at 0 min after UV treatment (Hu et al. 2017).
125 Specifically, the XR-seq signals from an example (6-4)PP repair hotspot decrease
126 dramatically from 1 min to 20 min and can be barely seen at 1 h (Figure 2A). In contrast,
127 the XR-seq signals at a late-repair coldspot shown in Figure 2B increases over the time
128 course and peaks at 4 h. Another representative CPD repair hotspot at 12 min is shown
129 in Figure 2C.

130  We next performed sequence context analysis using all reads that are mapped to
131 the repair hotspots and coldspots, respectively. We trimmed the reads to be 15 bp long
132 centering at the damage sites and calculated strand-specific nucleotide frequencies in
133 repair hotspots, coldspots, and randomly picked spots. Interestingly, we identified an
134 enrichment of cytosine in the flanking regions of the damage/repair sites for both repair
135 hotspots and coldspots (Figure S6). Motif analysis by the MEME suite (Bailey et al. 2009)
136 confirmed such enrichment of cytosine adjacent to the repair sites, which are themselves
137 enriched with canonical sequences of CTCA for (6-4)PP and TT for CPD (Table S5).
138 While it has been previously shown that active transcription factor binding sites have
139 decreased levels of excision repair (Sabarinathan et al. 2016), here we report a
140 preference for cytosine at both early and late timepoints, which have increased levels of

141 repair compared to null regions. While an interesting observation, the underlying
142 biological mechanisms need further investigation.
143
Early-repair hotspots are not hypersensitive to DNA damage
A recent study assayed DNA damage by next-generation sequencing and reported a total
of 153 hyper-hotspots acquiring CPDs much more frequently than the genomic average
in primary human fibroblasts (Premi et al. 2019). Of these damage hotspots, 83 are from
the plus strand and 74 are from the minus strand, each having at least five recurrent
sequence reads (Premi et al. 2019). To investigate whether the identified repair hotspots
simply result from increased levels of DNA damage, we intersected the reported damage
hotspots from Premi et al. (2019) with the 99 and 93 CPD repair hotspots from the plus
and minus strand that we identified. We found that none of our identified repair hotspots
overlapped with the reported damage hotspots.
To further confirm this seemingly striking result, we analyzed genome-wide CPD
DNA damage data generated by our previously developed Damage-seq protocol (Hu et
al. 2017), to quantify damage levels at 0 timepoint after UV irradiation with single-
nucleotide resolution. After stringent quality control procedures (refer to the Methods
section for details), we identified 91 and 78 CPD damage hotspots from the plus and
minus strand, respectively, each having at least ten mapped reads (Table S6). Notably,
these Damage-seq hotspots are shown to be enriched for heterochromatin and repressed
regions (Figure S7), which is concordant with previous reports (Han et al. 2016; Hauer
and Gasser 2017). Importantly, none of the CPD damage hotspots, identified from this
parallel Damage-seq platform, overlap with the repair hotspots.
In addition, we compared the Damage-seq signals for (6-4)PP and CPD at our
identified hotspots and coldspots against those from randomly sampled regions along the
genome. Regions corresponding to the repair hotspots, coldspots, and random spots are
extended at both ends for 20bp and 500bp, respectively, to account for the different
sampling densities between Damage-seq and XR-seq. Our results, shown in Figure 3,
suggest that there is no significant difference in the damage levels between the three
repair categories (hotspot, coldspot, and random spot). The zoom-in and zoom-out views
of three example repair hotspots and coldspots in Figure 2 also suggest that the Damage-

seq reads are uniformly distributed in the flanking regions. Previous results have
demonstrated that the UV-induced DNA damage is indeed virtually uniform across the
entire human genome, while repair is affected by chromatin states, transcription factor
binding, etc., in a manner dependent on the type of DNA damage (Hu et al. 2017). While
we note that the shallow depth of coverage of Damage-seq can be a limiting factor (refer
to the Discussion section for details), our results validate our conclusion that the identified
repair hotspots are not damage hotspots.
179
Early-repair hotspots are enriched for promoters and enhancers in open chromatin
regions
The packaging of DNA into chromatin can hinder the access of repair proteins and affect
the repair efficiency (Adar et al. 2016; Yimit et al. 2019) and specific histone modifications
are associated with different functional and cytological chromatin states. To investigate
the relationship between the identified repair hotspots/coldspots and the epigenomic
markers, we used publicly available DNase-seq data and histone modification ChIP-seq
data for an adult human fibroblast cell line (NHLF) from ENCODE (ENCODE Project
Consortium 2012). Our results suggest that the earliest repair occurs preferentially in
active and open-chromatin regions. Specifically, chromatin accessibility by DNase-seq is
significantly higher in repair hotspots than in random genomic regions, and it is
significantly lower in repair coldspots (Figure 4). For histone modifications, repair hotspots
have significantly higher ChIP-seq signals compared to random spots for activation
markers, including H3K4me1, H3K4me3, and H3K27ac (Figure 4).
We adopted the segmented chromatin states by chromHMM (Ernst and Kellis 2012)
to annotate the repair hotspots for (6-4)PP and CPD and found a significant enrichment
of active promoters and enhancers (Figure 5A), which are characterized by nucleosome
loss and open chromatin. Genome-wide repair kinetics inferred across timepoints confirm
that enhancers are repaired at earlier timepoints and that repressed/heterochromatin
regions are repaired at later timepoints (Figure S8). Additionally, we performed genome-
wide annotations for CpG islands and genic/intergenic regions. Our results suggest that
the repair hotspots are enriched for inter CpG regions (i.e., depleted for CpG islands)
(Figure 5B) and intronic regions (Figure 5C), both of which are enriched for AT and

presumably result in more repair. Notably, the number of identified repair coldspots is too
low to generate reproducible annotation results between the replicates.
205

Early-repair hotspots are enriched for frequently interacting regions (FIREs) and
super-enhancers
With the genome-wide enrichment of active promoters and enhancers in open chromatin
regions of the detected repair hotspots, we hypothesize that the 3D genome structure
contributes to the observed differential repair kinetics between different regions of the
genome. To test this hypothesis, we sought to interrogate the publicly available Hi-C data
of human fibroblast cell line IMR90 (Dixon et al. 2012; Jin et al. 2013). Specifically, after
quality control procedures and data normalization, we profiled frequently interacting
regions (FIREs) using FIREcaller (Crowley et al. 2019). After overlapping the repair
hotspots and coldspots with the called FIREs (Table S7A), we found that a significantly
higher proportion of repair hotspots overlap with FIREs – 23.16% and 11.76% for (6-4)PP
and CPD, respectively – compared to a genome average of 6.93% based on the profiled
FIREs (Figure 6A). Conversely, the overlapping proportion of (6-4)PP repair coldspots is
3.23%, significantly lower than the genome average (Figure 6A).
FIREs have been previously reported to be enriched for super-enhancers (Schmitt
et al. 2016). We have demonstrated that the repair hotspots are enriched for both FIREs
and enhancers. We also observed that, across many cases, multiple enhancers that
overlapped with the repair hotspots were from the same genomic regions (Figure S9). As
such, we hypothesize that the repair hotspots are also enriched for super-enhancers and
thus additionally adopted a list of previously annotated super-enhancers in the human
fibroblasts (Roadmap Epigenomics Consortium, 2015) (Table S7B). We found that,
compared to a genome-wide average of 2.05%, the early-repair hotspots are indeed
enriched for super-enhancers (5.14% and 4.69% for (6-4)PP and CPD repair hotspots,
respectively), while none of the repair coldspots overlap with super-enhancers (0% for (6-
4)PP repair coldspot) (Figure 6B).
In addition, we called significant interactions based on the Hi-C contact matrix
using the Fit-Hi-C method (Ay et al. 2014) (Table S7C) and showed that early-repair
hotspots also overlap with a significantly higher number of significant interactions (Figure

6C). Refer to the Methods section for details on data analysis. The overlapping

information of the called repair hotspots and coldspots with the profiled FIREs, super-

enhancers, and significant chromatin interactions are included in Table S8. Figure 6D

illustrates the loop interactions of two identified repair hotspots. Notably, these two

hotspots also overlap with both FIREs and super-enhancers. These results collectively

provide a global picture of genetic regulation of repair kinetics via 3D genome structures.

Early-repair hotspots are enriched for early-replication domains

We next set out to test the relationship between the repair kinetics and replication timing.

We used the genome-wide Repli-Seq data of the human fibroblast cell line IMR90, which

maps high-resolution DNA replication patterns with respect to both cell-cycle time and

genomic position (Hansen et al. 2010). A recent re-analysis of the data using a deep-

learning model (Liu et al. 2016) segmented the genome into different replication domains

including the early-replication domain (ERD), late-replication domain (LRD), up-transition

zone (UTZ), and down-transition zone (DTZ). Refer to Table S9 for genome-wide

segmentation results. When overlapping the identified repair hotspots with the segmented

replication domains, we find that early-repair hotspots are also significantly enriched for

early-replication domains (Figure 7). While replication time has been shown to be

correlated with chromatin accessibility (Hansen et al. 2010), here we have demonstrated

a potential cell-cycle effect on DNA excision repair. That is, regions that are duplicated

early also tend to be repaired early and this has been shown to result in mutagenesis

asymmetries (Adar et al. 2016) – we further elaborate such implications in the Discussion

section.

Profiling gene-level repair kinetics in normal human fibroblasts

So far, our analysis has been conducted on the whole-genome level. Identifying dynamic

changes at individual gene levels can provide important insights about genes and

pathways that exhibit significant changes in repair dynamics (Li et al. 2019). We

calculated strand-specific reads per kilobase per million reads (RPKM) for all the genes

in the human genome. RPKMs across nine genes associated with circadian rhythm (e.g.,

CLOCK, *BMAL1*, *PER1*, *CRY1*) and excision repair (e.g., *XPA*, *XPC, CSA, CSB*) present

different repair dynamics over the time course (Figure S10). We adopted Trendy (Bacher et al. 2018) to perform segmented regression analysis (Table S10) on the ordered XR-seq data. We focused on significant genes with adjusted R^2 greater than 0.8 from the regression analysis and identified 3,017 significant genes with fitted trend as "down," 3,496 significant genes with fitted trend as "up-down," 1,024 significant genes with fitted trend as "up," and 1,452 significant genes with fitted trend as "down-up" (Figure S11). The number of genes that show the "down" trend is two-fold higher than the number of genes that show the "up" trend, which is consistent with our previous observation – there are more repair hotspots than coldspots. For these four groups of significant genes, we further performed gene ontology (GO) enrichment analysis using PANTHER (Thomas et al. 2003). For genes with the "down" and "up-down" trend, GO biological processes are enriched for a large number of biological regulations, including regulation of metabolic process, gene expression, developmental process, cell cycle, and cell death, etc. (Table S11A-B). On the other hand, there are no obvious GO enrichment terms for the "up" or "down-up" genes (Table S11C-D).

DISCUSSION

We adopted time-course XR-seq data to detect genome-wide repair hotspots and coldspots and integrated additional omics data from various next-generation sequencing platforms to investigate the relationship between DNA damage, excision repair, epigenomic markers, 3D genome organization, and replication timing. To our best knowledge, this is the first demonstration at high-resolution of the connection between repair kinetics and chromatin organization. The identified repair hotspots can serve as additional functional genomic features in studies of the human genome at multidimensional level. We believe the framework and multi-omics data we present here can be useful to scientists both in the fields of chromatin dynamics and those interested in determinants of repair rates in human genome. Our approach is applicable to studying the response to other types of DNA damage including that induced by drugs used in chemotherapy and thus has the potential for informing new therapeutic strategies.

Using paired XR-seq and Damage-seq data, we provided additional empirical evidence that the damage levels in the regions of the repair hotspots were not enriched,

in concordance with previous reports (Hu et al. 2016; Hu et al. 2017). Therefore, we
attributed the observed differences in repair patterns to heterogeneous repair efficiencies
and not to damage formation. It is, however, noteworthy that the depth of coverage by
Damage-seq is an order of magnitude lower than that by XR-seq. Specifically, the
genome-wide average total number of reads between the two replicates for (6-4)PP
Damage-seq is 20.5 million, while that average for (6-4)PP XR-seq is 181.8 million (Table
S1). As such, it is possible that we do not have enough sequencing depth to detect
Damage-seq hotspots with high sensitivity. The fact that a good proportion of Damage-
seq reads get mapped to heterochromatin regions further complicates analysis and
lowers detection power. This issue also persists for the recently developed adductSeq
and FreqSeq, where a Poisson model was adopted with a genomic average rate of 0.07
reads per pyrimidine dinucleotide in fibroblasts (Premi et al. 2019). A Poisson distribution
with a mean parameter under the null much less than one could potentially suffer from
overdispersion and inflated false positive rates. Targeted Damage-seq offers a solution
for both discovery and validation. However, as with targeted DNA sequencing, the data
can be extremely noisy due to targeting, amplification, and sequencing biases and
artifacts (Jiang et al. 2018).

Using the genome-wide profile of DNA replication timing, we demonstrate that the
early-repair hotspots also tend to reside in early-replication domains. The inevitable effect
of replication on gene dosage and copy number could also have regulatory consequences
(Rhind and Gilbert 2013). Genes/regions that are duplicated early will be present at twice
the copy number of late replicating domains for most of the duration of S phase,
increasing both the DNA amount (as sister chromatids) and the transcriptional output.
The ability to perform excision repair for a greater fraction of the cell cycle has been shown
to result in lower mutation rates in the early-replication domains (Stamatoyannopoulos et
al. 2009; Lang and Murray 2011).

Granted, UV-induced DNA damage and excision repair have been linked with
mutagenesis and carcinogenesis. Recent studies have shown that mutation hotspots
exhibit strong increases in CPD formation efficacy (Elliott et al. 2018) and that excision
repair is attenuated in transcription factor binding sites, leading to elevated somatic
mutation rates (Sabarinathan et al. 2016). In terms of repair kinetics, Adar et al. (2016)

showed that globally late-repaired regions are associated with a higher level of cancer-
linked mutations. To further investigate this in our study context, we quantified the rates
of somatic point mutations associated with melanoma (Hodis et al. 2012) in the repair
hotspots and coldspots. However, due to the low number of hotspots and coldspots, as
well as their small sizes, we found very few overlapped with mutations and were thus
extremely underpowered to test for the differences in mutation rates between different
repair categories.

While the repair of (6-4)PP is predominantly carried out by global repair,
transcription-coupled repair plays an essential role in removing the more abundant and
less helix-distorting CPD adducts. In this study, we have intentionally focused on profiling
global repair hotspots for both (6-4)PP and CPD at early timepoints without the effect of
transcription-coupled repair. Genes exhibit both transcriptional dynamics over a time
course and biological fluctuations due to transcriptional bursting (Jiang et al. 2017) at the
same timepoint. This confounds and complicates the analysis. To disentangle the effects
of global repair and transcriptional-coupled repair in identifying repair hotspots is an
unsolved yet challenging problem.

METHODS
Experimental methods
Cell culture and UV irradiation
Human NHF1 cells were obtained from W.K. Kaufmann (University of North Carolina,
Chapel Hill) (Heffernan et al. 2002) and cultured in Dulbecco's Modified Eagle Medium
(DMEM) with 10% FBS at 37 °C in a 5% CO_2 humidified chamber. For (6-4)PP XR-seq
at 1 min and 2 min timepoints, UV irradiation was performed as previously described (Hu
et al. 2015; Li et al. 2017). Briefly, the 80% confluent NHF1 cells were irradiated for 20 s
under a 250 nm UV lamp (1 $J/m^2/s$) after removing the culture medium and incubated at
37 °C in a 5% CO_2 humidified chamber for 1 min or 2 min with 37 °C culture medium.
Then, the cells were put on ice immediately and washed one time with ice-cold PBS
before being harvested by a cell scarper in 10 ml ice-cold PBS. In each replicate of (6-
4)PP XR-seq experiment, 50 and 30 petri dishes (150mm x 15mm) containing NHF1 cells
were used at 1 min and 2 min timepoints respectively. Cell culture, UV treatment, and

library preparation for (6-4)PP XR-seq at 5 min, 20 min, 1 h, 2 h, and 4 h, and CPD XR-

seq at 12 min were performed in previous studies (Hu et al. 2015; Adar et al. 2016; Deger

et al. 2019). For *in vivo* excision assay, UV irradiation was performed as aforementioned,

and 10 and 5 petri dishes (150mm x 15mm) containing NHF1 cells were used at 0 min

and 2 min timepoints respectively.

Excision assay

The *in vivo* excision assay was performed as described (Hu et al. 2013; Hu et al. 2019).

Following UV irradiation, the excision products were isolated by gentle cell lysis and

nonchromatin fraction separation, and purified by TFIIH immunoprecipitation. The purified

excision products were then 3' radiolabeled by terminal deoxynucleotidyl transferase and

[α-^{32}P]-3'-dATP, and resolved in a 10% denaturing acrylamide gel. Ten and five petri

dishes (150mm x 15mm) of NHF1 cells were used at 0 min and 2 min, respectively.

XR-seq library preparation and sequencing

XR-seq libraries were prepared as described in the previous protocol (Hu et al. 2019).

Briefly, the excision products were isolated by TFIIH immunoprecipitation following gentle

cell lysis and non-chromatin fraction separation, and ligated with adaptors. The ligated

excision products were then further purified by immunoprecipitation with anti-(6-4)PP

antibody and repaired by (6-4)PP photolyase before the library amplification by PCR.

Libraries were sequenced on an Illumina HiSeq 4000 platform.

Data collection

(6-4)PP XR-seq data at 5 min, 20 min, 1 h, 2 h, and 4 h were downloaded from the Gene

Expression Omnibus (GEO) with accession numbers GSE67941 (Hu et al. 2015) and

GSE76391 (Adar et al. 2016). CPD XR-seq data at 12 min were downloaded from GEO

with accession number GSE138846 (Deger et al. 2019). CPD and (6-4)PP Damage-seq

data of NHF1 were downloaded from with accession number GSE98025 (Hu et al. 2017).

Hyper-hotspots for UV-induced CPD damage in primary human fibroblasts were

downloaded from Premi et al. (2019). NHDF H3K4me1 (ENCODE Data Coordination

Center accession number ENCSR000ARV), H3K4me3 (accession number

ENCSR000DPR), H3K27ac (accession number ENCSR000APN), H3K27me3 (accession number ENCSR000APO), H3K9me3 (accession number ENCSR000ARX), and DNase-seq (accession number ENCSR000EMP) data were downloaded from the ENCODE portal (ENCODE Project Consortium 2012). NHLF chromatin state segmentation results by chromHMM were downloaded from UCSC accession number wgEncodeEH000792 (Ernst and Kellis 2012). Hi-C data of IMR90 were downloaded from GEO with accession number GSE43070 (Jin et al. 2013) and from https://bioconductor.org/packages/HiCDataHumanIMR90/ (Dixon et al. 2012). The list of annotated super-enhancers in IMR90 was downloaded from the Roadmap Epigenomics Consortium (Roadmap Epigenomics Consortium, 2015). Genomic categories of replication timing from Repli-Seq data of IMR90 (Hansen et al. 2010) were downloaded from GSE53984 (Liu et al. 2016).

Bioinformatic and statistical analysis

XR-seq bioinformatic pre-processing

For XR-seq, cutadapt (Martin 2011) was used to trim reads with adaptor sequence TGGAATTCTCGGGTGCCAAGGAACTCCAGTNNNNNNACGATCTCGTATGCCGTCT TCTGCTTG at the 3' end and discard untrimmed reads. BWA (Li and Durbin 2009) was used for alignment of single-end short reads. Unmapped reads and reads that map to multiple locations with the same alignment quality were removed using Samtools (Li et al. 2009). Post-alignment filtering steps were adopted using Rsamtools (http://bioconductor.org/packages/Rsamtools/). Specifically, if multiple reads share the same 5' and 3' end coordinates, we keep only one to perform deduplication. We also only keep reads that have mapping quality greater than 20 and are of lengths 21 bp to 31 bp.

Gene-level quantification of excision repair

Reads from the TS and NTS strands were separated using known gene annotations for hg19 by ENSEMBL. We use RPKM for within-sample normalization for the XR-seq data. To perform gene-level quantification and downstream analysis including segmented regression, we adopted a stringent quality control procedure and only retained genes that: (i) had at least 10 TT or TC dinucleotides from either TS or NTS; (ii) were less than 300Kb;

and (iii) had at least 10 reads in total across all XR-seq samples. In addition, we took the
ratio of the reads from the TS and the NTS [TS/(TS+NTS)] to remove biases and artifact
that are shared between the two DNA stands, i.e., library size, gene length, and other
gene-specific biases, such as sequencing bias and antibody pulldown efficiency, etc. The
ratio is bound between 0 and 1 and sheds light upon how transcription-coupled repair and
global repair interplay (Figure S3).

Computational approach to identify repair hotspots and coldspots

We started by segmenting the human reference genome into consecutive bins of 50 bp
long. We then calculated the observed depth of coverage per bin by XR-seq, separating
the plus-strand reads (+) and the minus-strand reads (-). To mitigate the effect of library
size/sequencing depth, we downsampled the reads in each sample to 7.7 million without
replacement. Given such data input, we identified strong repair hotspots in
pericentromeric regions, which were collapsed repeats annotated as unique sequences
in the reference genome (e.g., ribosomal DNA (Yang et al. 2019)). It is important to
exclude artifacts as stringently as possible and thus we undertook an additional quality
control step. "Blacklist" bins, including segmental duplication regions
(http://humanparalogy.gs.washington.edu/build37/data/GRCh37GenomicSuperDup.tab),
gaps in reference assembly from telomere, centromere, and/or heterochromatin regions
(https://gist.github.com/leipzig/6123703), and repeating elements by RepeatMasker
(https://genome.ucsc.edu/cgi-bin/hgTrackUi?g=rmsk) are masked in downstream
analysis. To identify repair hotspots and coldspot, we set a threshold on the number of
read counts per genomic bin in the 1 min and 4 h samples. Specifically, to identify (6-
4)PP repair hotspots, we require at least 15 reads mapped in both replicates at 1min and
at most 5 reads mapped in both replicates at 4 h. The read count threshold is relaxed for
the identification of coldspots, which have a smaller number compared to the hotspots.
For CPD repair, to avoid complicatedness due to transcription-coupled repair at later
timepoints, we focused on CPD repair hotspots only.

Hi-C data analysis

We adopted the Hi-C data of human fibroblast cell line IMR90 (Dixon et al. 2012; Jin et al. 2013) to investigate the relationship between identified repair hotspots and the 3D genome structure. We took the raw contact matrix with 40 kb resolution as input and detected FIREs, which play important roles in transcriptional regulations, across the entire genome using FIREcaller (Crowley et al. 2019). To further investigate whether these repair hotspots are involved in functional chromatin looping between regulatory elements and their target genes, we adopted the Fit-Hi-C approach (Ay et al. 2014) to identify long-range chromatin interactions on all 40 kb bin pairs within a maximal 3 MB region. The interactions with p-value < 2.31e-11 were considered as statistically significant (Giusti-Rodríguez et al. 2019).

DATA AND CODE AVAILABILITY

The data reported in this paper have been deposited in GEO with accession number GSE148303. Scripts used in this paper are available at https://github.com/yuchaojiang/damage_repair.

COMPETING INTEREST STATEMENT

The authors declare no competing interests.

ACKNOWLEDGEMENTS

This work was supported by NIH Grants R35 GM118102 (to A.S.), R01 ES027255 (to A.S.), P01 CA142538 (to Y.J.), UL1 TR002489 (to Y.J.), K99 ES030015 (to W.L.), and a pilot award from the UNC Computational Medicine Program (to Y.J.). We thank the Sancar Lab members and Drs. Sheera Adar, Ming Hu, and Jeremy Simon for useful comments and feedback.

AUTHOR CONTRIBUTIONS

AS envisioned and initiated the study, while LW and LAL-B performed the experiments. YJ devised the analytical framework and all authors executed the data analysis. YJ wrote

479 the manuscript, with contributions from WL on experimental methods. The manuscript
480 was further edited and approved by all authors.
481

<u>REFERENCES</u>

Adar S, Hu J, Lieb JD, Sancar A. 2016. Genome-wide kinetics of DNA excision repair in relation to
chromatin state and mutagenesis. *Proc Natl Acad Sci U S A* **113**: E2124-2133.

Adebali O, Chiou YY, Hu J, Sancar A, Selby CP. 2017. Genome-wide transcription-coupled repair
in Escherichia coli is mediated by the Mfd translocase. *Proc Natl Acad Sci U S A* **114**: E2116-
E2125.

Akkose U, Kaya O, Lindsey-Boltz L, Karagoz Z, Brown AD, Larsen PA, Yoder AD, Sancar A, Adebali
O. 2020. Comparative analyses of two primate species diverged by more than 60 million
years show different rates but similar distribution of genome-wide UV repair events.
bioRxiv.

Ay F, Bailey TL, Noble WS. 2014. Statistical confidence estimation for Hi-C data reveals regulatory
chromatin contacts. *Genome Res* **24**: 999-1011.

Bacher R, Leng N, Chu LF, Ni Z, Thomson JA, Kendziorski C, Stewart R. 2018. Trendy: segmented
regression analysis of expression dynamics in high-throughput ordered profiling
experiments. *BMC Bioinformatics* **19**: 380.

Bailey TL, Boden M, Buske FA, Frith M, Grant CE, Clementi L, Ren J, Li WW, Noble WS. 2009.
MEME SUITE: tools for motif discovery and searching. *Nucleic Acids Res* **37**: W202-208.

Consortium EP. 2012. An integrated encyclopedia of DNA elements in the human genome. *Nature*
489: 57-74.

Consortium RE, Kundaje A, Meuleman W, Ernst J, Bilenky M, Yen A, Heravi-Moussavi A,
Kheradpour P, Zhang Z, Wang J et al. 2015. Integrative analysis of 111 reference human
epigenomes. *Nature* **518**: 317-330.

Crowley C, Yang Y, Qiu Y, Hu B, Won H, Ren B, Hu M, Li Y. 2019. FIREcaller: an R package for
detecting frequently interacting regions from Hi-C data. *bioRxiv*: 619288.

Deger N, Yang Y, Lindsey-Boltz LA, Sancar A, Selby CP. 2019. Drosophila, which lacks canonical
transcription-coupled repair proteins, performs transcription-coupled repair. *J Biol Chem*
294: 18092-18098.

Dixon JR, Selvaraj S, Yue F, Kim A, Li Y, Shen Y, Hu M, Liu JS, Ren B. 2012. Topological domains in
mammalian genomes identified by analysis of chromatin interactions. *Nature* **485**: 376-
380.

Elliott K, Bostrom M, Filges S, Lindberg M, Van den Eynden J, Stahlberg A, Clausen AR, Larsson E.
2018. Elevated pyrimidine dimer formation at distinct genomic bases underlies promoter
mutation hotspots in UV-exposed cancers. *PLoS Genet* **14**: e1007849.

Ernst J, Kellis M. 2012. ChromHMM: automating chromatin-state discovery and characterization.
Nat Methods **9**: 215-216.

Garcia-Nieto PE, Schwartz EK, King DA, Paulsen J, Collas P, Herrera RE, Morrison AJ. 2017.
Carcinogen susceptibility is regulated by genome architecture and predicts cancer
mutagenesis. *EMBO J* **36**: 2829-2843.

Giusti-Rodríguez P, Lu L, Yang Y, Crowley CA, Liu X, Juric I, Martin JS, Abnousi A, Allred SC,
Ancalade N. 2019. Using three-dimensional regulatory chromatin interactions from adult
and fetal cortex to interpret genetic results for psychiatric disorders and cognitive traits.
bioRxiv: 406330.

Han C, Srivastava AK, Cui T, Wang QE, Wani AA. 2016. Differential DNA lesion formation and
repair in heterochromatin and euchromatin. *Carcinogenesis* **37**: 129-138.

Hanawalt PC, Spivak G. 2008. Transcription-coupled DNA repair: two decades of progress and
surprises. *Nat Rev Mol Cell Biol* **9**: 958-970.

Hansen RS, Thomas S, Sandstrom R, Canfield TK, Thurman RE, Weaver M, Dorschner MO, Gartler
SM, Stamatoyannopoulos JA. 2010. Sequencing newly replicated DNA reveals widespread
plasticity in human replication timing. *Proc Natl Acad Sci U S A* **107**: 139-144.

Hauer MH, Gasser SM. 2017. Chromatin and nucleosome dynamics in DNA damage and repair.
Genes Dev **31**: 2204-2221.

Heffernan TP, Simpson DA, Frank AR, Heinloth AN, Paules RS, Cordeiro-Stone M, Kaufmann WK.
2002. An ATR- and Chk1-dependent S checkpoint inhibits replicon initiation following
UVC-induced DNA damage. *Mol Cell Biol* **22**: 8552-8561.

Hodis E, Watson IR, Kryukov GV, Arold ST, Imielinski M, Theurillat JP, Nickerson E, Auclair D, Li L,
Place C et al. 2012. A landscape of driver mutations in melanoma. *Cell* **150**: 251-263.

Hu J, Adar S, Selby CP, Lieb JD, Sancar A. 2015. Genome-wide analysis of human global and
transcription-coupled excision repair of UV damage at single-nucleotide resolution. *Genes*
Dev **29**: 948-960.

Hu J, Adebali O, Adar S, Sancar A. 2017. Dynamic maps of UV damage formation and repair for
the human genome. *Proc Natl Acad Sci U S A* **114**: 6758-6763.

Hu J, Choi JH, Gaddameedhi S, Kemp MG, Reardon JT, Sancar A. 2013. Nucleotide excision repair
in human cells: fate of the excised oligonucleotide carrying DNA damage in vivo. *J Biol*
Chem **288**: 20918-20926.

Hu J, Li W, Adebali O, Yang Y, Oztas O, Selby CP, Sancar A. 2019. Genome-wide mapping of
nucleotide excision repair with XR-seq. *Nat Protoc* **14**: 248-282.

Hu J, Lieb JD, Sancar A, Adar S. 2016. Cisplatin DNA damage and repair maps of the human
genome at single-nucleotide resolution. *Proc Natl Acad Sci U S A* **113**: 11507-11512.

Jiang Y, Wang R, Urrutia E, Anastopoulos IN, Nathanson KL, Zhang NR. 2018. CODEX2: full-
spectrum copy number variation detection by high-throughput DNA sequencing. *Genome*
Biol **19**: 202.

Jiang Y, Zhang NR, Li M. 2017. SCALE: modeling allele-specific gene expression by single-cell RNA
sequencing. *Genome Biol* **18**: 74.

Jin F, Li Y, Dixon JR, Selvaraj S, Ye Z, Lee AY, Yen CA, Schmitt AD, Espinoza CA, Ren B. 2013. A high-
resolution map of the three-dimensional chromatin interactome in human cells. *Nature*
503: 290-294.

Kusakabe M, Onishi Y, Tada H, Kurihara F, Kusao K, Furukawa M, Iwai S, Yokoi M, Sakai W,
Sugasawa K. 2019. Mechanism and regulation of DNA damage recognition in nucleotide
excision repair. *Genes Environ* **41**: 2.

Lang GI, Murray AW. 2011. Mutation rates across budding yeast chromosome VI are correlated
with replication timing. *Genome Biol Evol* **3**: 799-811.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics **25**: 1754-1760.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, Genome
Project Data Processing S. 2009. The Sequence Alignment/Map format and SAMtools.
Bioinformatics **25**: 2078-2079.

Li W, Adebali O, Yang Y, Selby CP, Sancar A. 2018. Single-nucleotide resolution dynamic repair
maps of UV damage in Saccharomyces cerevisiae genome. *Proc Natl Acad Sci U S A* **115**:
E3408-E3415.

Li W, Hu J, Adebali O, Adar S, Yang Y, Chiou YY, Sancar A. 2017. Human genome-wide repair map
of DNA damage caused by the cigarette smoke carcinogen benzo[a]pyrene. *Proc Natl*
Acad Sci U S A **114**: 6752-6757.

Li W, Liu W, Kakoki A, Wang R, Adebali O, Jiang Y, Sancar A. 2019. Nucleotide excision repair
capacity increases during differentiation of human embryonic carcinoma cells into
neurons and muscle cells. *J Biol Chem* **294**: 5914-5922.

Liu F, Ren C, Li H, Zhou P, Bo X, Shu W. 2016. De novo identification of replication-timing domains
in the human genome by deep learning. *Bioinformatics* **32**: 641-649.

Mao P, Brown AJ, Esaki S, Lockwood S, Poon GMK, Smerdon MJ, Roberts SA, Wyrick JJ. 2018. ETS
transcription factors induce a unique UV damage signature that drives recurrent
mutagenesis in melanoma. *Nature communications* **9**: 2626.

Mao P, Smerdon MJ, Roberts SA, Wyrick JJ. 2016. Chromosomal landscape of UV damage
formation and repair at single-nucleotide resolution. *Proc Natl Acad Sci U S A* **113**: 9057-
9062.

Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet journal **17**: 10-12.

Martincorena I, Roshan A, Gerstung M, Ellis P, Van Loo P, McLaren S, Wedge DC, Fullam A,
Alexandrov LB, Tubio JM et al. 2015. Tumor evolution. High burden and pervasive positive
selection of somatic mutations in normal human skin. *Science* **348**: 880-886.

Mellon I, Spivak G, Hanawalt PC. 1987. Selective removal of transcription-blocking DNA damage
from the transcribed strand of the mammalian DHFR gene. *Cell* **51**: 241-249.

Oztas O, Selby CP, Sancar A, Adebali O. 2018. Genome-wide excision repair in Arabidopsis is
coupled to transcription and reflects circadian gene expression patterns. *Nat Commun* **9**:
1503.

Poulos RC, Thoms JAI, Guan YF, Unnikrishnan A, Pimanda JE, Wong JWH. 2016. Functional
Mutations Form at CTCF-Cohesin Binding Sites in Melanoma Due to Uneven Nucleotide
Excision Repair across the Motif. *Cell reports* **17**: 2865-2872.

Premi S, Han L, Mehta S, Knight J, Zhao D, Palmatier MA, Kornacker K, Brash DE. 2019. Genomic
sites hypersensitive to ultraviolet radiation. *Proc Natl Acad Sci U S A* **116**: 24196-24205.

Reardon JT, Sancar A. 2003. Recognition and repair of the cyclobutane thymine dimer, a major
cause of skin cancers, by the human excision nuclease. *Genes Dev* **17**: 2539-2551.

Reardon JT, Sancar A. 2005. Nucleotide excision repair. *Prog Nucleic Acid Res Mol Biol* **79**: 183-
235.

Rhind N, Gilbert DM. 2013. DNA replication timing. *Cold Spring Harb Perspect Biol* **5**: a010132.

Sabarinathan R, Mularoni L, Deu-Pons J, Gonzalez-Perez A, Lopez-Bigas N. 2016. Nucleotide
excision repair is impaired by binding of transcription factors to DNA. *Nature* **532**: 264-
267.

Schmitt AD, Hu M, Jung I, Xu Z, Qiu Y, Tan CL, Li Y, Lin S, Lin Y, Barr CL et al. 2016. A Compendium
of Chromatin Contact Maps Reveals Spatially Active Regions in the Human Genome. *Cell*
Rep **17**: 2042-2059.

Stamatoyannopoulos JA, Adzhubei I, Thurman RE, Kryukov GV, Mirkin SM, Sunyaev SR. 2009.
Human mutation rate associated with DNA replication timing. *Nat Genet* **41**: 393-395.

Thomas PD, Campbell MJ, Kejariwal A, Mi H, Karlak B, Daverman R, Diemer K, Muruganujan A,
Narechania A. 2003. PANTHER: a library of protein families and subfamilies indexed by
function. *Genome Res* **13**: 2129-2141.

Yang Y, Hu J, Selby CP, Li W, Yimit A, Jiang Y, Sancar A. 2019. Single-nucleotide resolution analysis
of nucleotide excision repair of ribosomal DNA in humans and mice. *J Biol Chem* **294**: 210-
217.

Yimit A, Adebali O, Sancar A, Jiang Y. 2019. Differential damage and repair of DNA-adducts
induced by anti-cancer drug cisplatin across mouse organs. *Nat Commun* **10**: 309.

FIGURE & TABLE LEGENDS

Figure 1. Excision repair kinetics and early-repair hotspots in normal human

fibroblasts. (A) Experimental design to measure DNA damage and excision repair of the

UV-induced (6-4)PP and CPD across different timepoints. (B) Detection of excision

products at 0 min and 2 min timepoints *in vivo*. Following UV irradiation, the excised

oligonucleotides were purified by TFIIH immunoprecipitation, radiolabeled, and resolved

in a 10% sequencing gel. DNA excision products of (6-4)PP can be detected as early as

2 min upon damage induction. (C) Principal component analysis of genome-wide excision

repair as measured by XR-seq shows repair kinetics across different timepoints. Between

1 min and 5 min, excised oligonucleotides were not degraded and thus XR-seq measured

cumulative repair. (D) Each row is a sample and each column is a specific total number

of reads per genomic bin. The color in the heatmap corresponds to the log counts of the

number of bins with specific read depths. Early-repair hotspots exist in samples collected

at early timepoints.

Figure 2. Distribution of DNA damage, repair, and epigenomic markers at identified

repair hotspots and coldspots. XR-seq and Damage-seq data are shown for both

strands, marked with + and -. Epigenetic data from ChIP-seq of histone modifications and

DNase-seq are plotted on the same scale for cross comparison. (A) A (6-4)PP repair

hotspot from chr2. (B) A (6-4)PP repair coldspot from chr1. (C) A CPD repair hotspot from

chr1. All examples are from intronic gene regions overlapping annotated enhancers.

Zoomed-in view of canonical sequences are overlaid in the bottom, with the

damage/repair sites shown in dashed boxes.

Figure 3. Repair hotspots are not damage hotspots. There is no enriched DNA

damage, as measured by Damage-seq, at the identified repair hotspots for (A) (6-4)PP

and (B) CPD. Regions corresponding to the different repair categories are extended at

both ends for 20bp and 500bp, respectively, to account for the different sampling densities

between Damage-seq and XR-seq.

Figure 4. Genome-wide repair hotspots are associated with epigenomic markers.

Chromatin accessibility (DNase I hypersensitivity) is higher for repair hotspots and lower

for coldspots. Repair hotspots are also characterized by higher ChIP-seq signals for

H3K4me1, H3K4me3, and H3K27ac, markers for gene activations.

Figure 5. Genome-wide repair hotspots are enriched for enhancers. Annotations for

(A) chromatin states, (B) CpG islands, and (C) genic/intergenic regions are shown for

repair hotspots for (6-4)PP (top) and CPD (bottom), separated by strands. Repair

hotspots are enriched for enhancers and promoters, which are in open-chromatin regions.

Repair hotspots are also enriched in inter-CpG and intronic regions, both of which are AT

rich.

Figure 6. Genome-wide repair hotspots overlap with FIREs and super-enhancers

identified by Hi-C. FIREs and super-enhancers were identified and annotated using Hi-

C data from human fibroblasts. Repair hotspots overlap with (A) FIREs and (B) super-

enhancers with significantly higher proportions compared to the genome-wide averages.

The solid density curves are generated from bootstrapping different regions along the

genome as the null case; the dashed vertical lines are the observed proportions for the

repair hotspots and coldspots. (C) Repair hotspots have significantly higher number of

significant interactions, identified by Hi-C. (D) Two examples of (6-4)PP repair hotspots

(chr8:13224201-13224300 and chr14:61994601-61994700) that overlap with both FIREs

and super-enhancers, and loop to different regions of the genome. Hi-C data has low

resolution and the significant interactions are drawn from the center of each bin, which

does not exactly overlap with the identified hotspot shown in red.

Figure 7. Genome-wide repair hotspots are enriched for early-replication domains.

Replication timing domains were identified using Repli-Seq data. Compared to genome-

wide average (random), there is a significantly higher proportion of the early-repair

hotspots located in the early-replication domains.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7