1 **Species-specific quorum sensing represses the chitobiose utilization locus**
2 **in *Vibrio cholerae***
3

4 Catherine A. Klancher, Jane D. Newman, Alyssa S. Ball, Julia C. van Kessel, and
5 Ankur B. Dalia*
6
7 Department of Biology, Indiana University, Bloomington, IN 47405
8 *Author for correspondence: ankdalia@indiana.edu

9
10

Abstract

11
The marine facultative pathogen *Vibrio cholerae* forms complex multicellular
communities on the chitinous shells of crustacean zooplankton in its aquatic
reservoir. *V. cholerae*-chitin interactions are critical for the growth, evolution, and
waterborne transmission of cholera. This is due, in part, to chitin-induced
changes in gene expression in this pathogen. Here, we sought to identify factors
that influence chitin-induced expression of one locus, the chitobiose utilization
operon (*chb*), which is required for the uptake and catabolism of the chitin
disaccharide. Through a series of genetic screens, we identified that the master
regulator of quorum sensing, HapR, is a direct repressor of the *chb* operon. We
also found that the levels of HapR in *V. cholerae* are regulated by the ClpAP
protease. Furthermore, we show that the canonical quorum sensing cascade in
V. cholerae regulates *chb* expression in a HapR-dependent manner. Through
this analysis we found that signaling via the species-specific autoinducer CAI-1,
but not the inter-species autoinducer AI-2, influences *chb* expression. This
phenomenon of species-specific regulation may enhance the fitness of this
pathogen in its environmental niche.

Importance
29
In nature, bacteria live in multicellular and multispecies communities. Microbial
species can sense the density and composition of their community through
chemical cues using a process called quorum sensing (QS). The marine
pathogen *Vibrio cholerae* is found in communities on the chitinous shells of
crustaceans in its aquatic reservoir. *V. cholerae* interactions with chitin are critical
for the survival, evolution, and waterborne transmission of this pathogen. Here,
we show that *V. cholerae* uses QS to regulate the expression of one locus
required for *V. cholerae*-chitin interactions.
38

Introduction
39
The facultative bacterial pathogen *Vibrio cholerae*, the causative agent of the
diarrheal disease cholera, natively resides in the aquatic environment. In this
niche, *V. cholerae* forms multicellular communities on biotic and abiotic chitinous
surfaces, like the shells of crustaceans or marine snow (1, 2). Chitin is a
polysaccharide made up of β-1,4 linked *N*-acetylglucosamine (GlcNAc) and
serves as a major nutrient source for *V. cholerae* in the marine environment (1, 3,
4). The ability of *V. cholerae* to form chitin biofilms is critical for the waterborne

transmission of cholera (5, 6). As chitin is the most abundant biopolymer in the
ocean, the ability of *Vibrio* species to break down and utilize this highly insoluble
polysaccharide also serves an important role in global nitrogen and carbon
recycling (1, 4).
51
When *V. cholerae* is associated with a chitinous surface, chitin induces the
expression of a subset of genes in *V. cholerae*. The genes induced by chitin
include those required for chitin degradation, uptake, and catabolism (termed the
chitin utilization program), as well as the genes required for natural
transformation (7, 8). Transcriptional responses resulting from *Vibrio*-chitin
interactions are highly regulated. One major chitin-responsive regulator is the
orphan hybrid sensor kinase ChiS (7, 9). ChiS senses chitin indirectly through the
periplasmic <u>c</u>hitin <u>b</u>inding <u>p</u>rotein (CBP) (9, 10). In the absence of chitin, CBP
represses ChiS through interactions with its periplasmic domain (9, 10). In the
presence of chitin, the CBP-chitin complex stimulates ChiS activity (9, 10). Thus,
in the presence of chitin, ChiS is active and can facilitate expression of the chitin
utilization program. Alternatively, ChiS can be genetically activated in the
absence of chitin by deleting *cbp* (10, 11).
65
In the marine environment, *V. cholerae* not only senses chitin to modulate gene
expression, but also the presence of other bacteria through a process termed
"quorum sensing" (QS) (12). This is a process by which bacteria indirectly sense
other microbes in their community via small diffusible molecules called
autoinducers (AIs). AIs are sensed by cognate sensor proteins. *V. cholerae*
encodes four AI sensors, although the autoinducer molecules that serve as
inducing cues are only known for two of them (13). AI sensing allows for cell-
density specific gene expression programs, which regulate "group" or "individual"
behaviors (14). *V. cholerae* senses both chitin and AIs to regulate natural
transformation on chitinous surfaces (15). Though a link between chitin utilization
and quorum sensing has previously been suggested, it has not been directly
studied (16).
78
To investigate regulation of the chitin utilization program in *V. cholerae*, most
studies employ the chitobiose utilization operon (*chb*) (10, 11, 17, 18). The *chb*
operon encodes the genes required for uptake and catabolism of the chitin
disaccharide chitobiose, and is highly induced in the presence of chitin
oligosaccharides (7). Several mechanisms of *chb* regulation have already been
identified. ChiS is the master regulator of the chitin utilization program in *V.*
cholerae, and we have recently shown this protein is a direct transcriptional
activator required for induction of the *chb* locus (9, 10). Previous work has shown
that carbon catabolite repression (CCR) can also play a role in regulating chitin
responsive phenotypes including ChiS-dependent induction of *chb* and natural
transformation (18, 19). In addition, our group has previously found that the cell
division licensing factor SlmA plays an essential role in activating *chb* expression
(11). Tight regulation via these diverse signaling systems may act to ensure that

the chitin utilization program is only expressed under conditions in which it will
provide a competitive advantage.
94
95   Here, we sought to identify additional regulators of *chb*. Through a number of
96   genetic screens and complementary molecular methods, we show that quorum
97   sensing is an additional regulatory system that tunes expression of a chitin
98   utilization locus in *V. cholerae*.
99
Results
ClpA is identified in an unbiased screen for activators of P$_{chb}$
To identify additional genes required for activation of the *chb* locus, we
conducted a transposon mutant screen. This was carried out in a strain
containing a chromosomally-integrated P$_{chb}$-*lacZ* transcriptional reporter. As
shown previously, induction of P$_{chb}$ is dependent on the activity of the master
regulator ChiS (7, 10, 11). In the absence of chitin, ChiS activity is repressed by
CBP. In the presence of chitin, CBP repression of ChiS is relieved, which allows
for ChiS-dependent activation of P$_{chb}$. In addition to being induced by chitin, ChiS
can be activated genetically in the absence of chitin by deleting *cbp* (10, 11). As
chitin oligomers are prohibitively expensive, a Δ*cbp* mutation was used to induce
ChiS-dependent P$_{chb}$-*lacZ* expression in our genetic screen exactly as previously
described (11). So, the starting genotype for our screen was a strain containing
P$_{chb}$-*lacZ* and a Δ*cbp* mutation. This strain formed blue colonies on X-gal
containing plates and we screened for white colonies to identify putative
activators that contribute to P$_{chb}$ induction.
116
Of approximately 60,000 transposon mutants visually screened for loss of P$_{chb}$-
lacZ expression, one gene identified was *clpA* (2 unique transposon insertions).
Other hits identified in this screen are listed in **Table S1**. To study the effect of
clpA on P$_{chb}$ activity moving forward, we utilized a previously described
chromosomally-integrated P$_{chb}$-GFP reporter (10, 11). Using this reporter, we
found that a Δ*clpA* mutation resulted in a ~3-fold decrease in P$_{chb}$ expression
relative to the parent (**Fig. 1**). Importantly, complementation of this strain with an
ectopic copy of *clpA in trans* restored P$_{chb}$ expression to parent levels (**Fig. S1**).
ClpA is a AAA+ unfoldase that recognizes protein substrates, unfolds them, and
feeds them into the ClpP protease where they are degraded (20). If ClpA was
exhibiting its effect on P$_{chb}$ expression as a part of the ClpAP machine, we
hypothesized that a Δ*clpP* mutation should phenocopy a Δ*clpA* mutation. Indeed,
Δ*clpP* and Δ*clpP* Δ*clpA* strains phenocopied a Δ*clpA* mutant for P$_{chb}$ expression
(**Fig. 1**). These results suggest that loss of the ClpAP protease decreases P$_{chb}$
expression. As ClpAP degrades proteins, we hypothesized that ClpAP may
indirectly promote activation of P$_{chb}$ by degrading a repressor of the *chb* locus.
133
HapR is a repressor of P$_{chb}$ that is degraded by ClpAP
To identify a putative repressor of P$_{chb}$ that is targeted by ClpAP for degradation,
we conducted a counter-screen using the P$_{chb}$-*lacZ* reporter. For the Δ*clpA*
counter-screen, the parent strain had both Δ*cbp* and Δ*clpA* mutations. This

mutant is white on X-gal plates because P$_{chb}$-lacZ is poorly expressed;
inactivation of the putative repressor in this strain should result in restoration of
P$_{chb}$-lacZ expression and yield a blue colony phenotype. In the ΔclpA counter-
screen, we visually screened approximately 30,000 transposon mutants for re-
activation of P$_{chb}$-lacZ expression and identified *hapR* (9 unique transposon
insertions). Other hits identified in this screen are listed in **Table S1**. A ΔhapR
mutation restored P$_{chb}$ expression in the ΔclpA mutant background (**Fig. 1**). In
fact, a ΔhapR mutation allowed for higher P$_{chb}$ expression than the parent strain.
This suggests that HapR represses P$_{chb}$ expression when ClpAP is intact and
that ClpAP does not degrade the entire pool of HapR in the cell (**Fig. 1**).
Importantly, the level of P$_{chb}$ expression observed in the ΔhapR ΔclpA mutant
phenocopied the ΔhapR strain (**Fig. 1**). This epistasis between *clpA* and *hapR*
suggests that they are involved in the same pathway for regulating P$_{chb}$
expression. In addition, complementation of the ΔhapR strain with an ectopic
copy of *hapR in trans* decreased P$_{chb}$ expression almost to the level in the parent
(**Fig. S1**). Further, complementation of the ΔhapR ΔclpA strain with *hapR in trans*
brought P$_{chb}$ expression down to the level observed in the ΔclpA parent (**Fig. S1**).
155
We hypothesized that the reason P$_{chb}$ expression was decreased in ΔclpAP
mutants was due to increased HapR protein levels. Western blotting in these
backgrounds revealed that HapR levels were, indeed, increased in strains
containing mutations to *clpA* and/or *clpP* (**Fig. 1**). ClpAP dependent degradation
of HapR is not unique to *V. cholerae*, but has previously been observed in *Vibrio*
vulnificus where ClpAP degrades the HapR homolog SmcR (21). In addition to
the ClpAP machine, it was shown that the Lon protease also plays a role in
SmcR degradation. So, we next sought to investigate the role of other protease
machines on induction of P$_{chb}$ and HapR protein levels. Mutations to other Clp
components (the ClpS adaptor protein or the ClpX unfoldase*)* did not have a
marked impact on HapR protein levels. Also, a ΔclpS mutation did not affect P$_{chb}$
expression levels, while a ΔclpX mutation slightly decreased P$_{chb}$ expression
(**Fig. 1**). Because HapR expression was not affected by the ΔclpX mutation, the
observed decrease in P$_{chb}$ expression may be attributed to a pleiotropic effect
(*i.e.*, an effect of *clpX* that is independent of HapR-dependent P$_{chb}$ repression). In
contrast to the effect of the Lon protease on SmcR levels in *V. vulnificus*, we did
not observe an impact of ΔlonA on HapR levels in *V. cholerae*; and ΔlonA,
correspondingly, did not affect P$_{chb}$ expression (**Fig. 1**). Together, these results
establish that HapR is a repressor of P$_{chb}$ and that HapR levels are controlled
specifically by the ClpAP protease in *V. cholerae*.
176
HapR-mediated repression of P$_{chb}$ occurs on chitinous surfaces
Thus far, we have studied P$_{chb}$ regulation using a Δcbp mutation to induce ChiS
activity. Natural induction of this locus, however, occurs in the presence of chitin
oligosaccharides. So, we next wanted to test whether the repression of P$_{chb}$ by
HapR was observed in a more physiologically relevant setting. To test this, we
cultured *V. cholerae* strains with CBP intact, a P$_{chb}$-mCherry reporter, and a
construct that constitutively expresses GFP (P$_{const2}$-GFP) on chitin beads (**Fig.**

2A). The P_{const2}-GFP construct (derived from the insulated *proD* promoter (22))
served as an internal control for noise in gene expression and was used to
normalize P_{chb}-mCherry expression in single cells (see Methods for details).
187
When cells were cultured on chitin beads, both the parent strain and the $\Delta hapR$
strain exhibited a bimodal distribution of P_{chb} expression (**Fig. 2A-C**). There were
at least two possible explanations for the observed bimodality in P_{chb} gene
expression in this experiment. One, the signaling pathway that leads to activation
of *chb* has switch-like behavior that results in a population that exhibits bimodality
in *chb* expression. Or two, the environment during growth on chitin beads is
heterogeneous and only some cells within the population have access to the
chitin oligosaccharides necessary for activation of *chb*. Those cells that do not
have access to chitin do not activate P_{chb} expression (P_{chb} Off), while those that
do have access to chitin do induce P_{chb} expression (P_{chb} On). To differentiate
between these two possibilities, we induced expression of P_{chb} either using the
native inducer, chitin oligosaccharides, or genetically by deleting *cbp*. If the
signaling circuit responsible for P_{chb} activation has switch-like behavior, we would
predict that bimodality in gene expression would be maintained in both of these
conditions; however, if bimodality is the result of heterogeneous access to chitin
oligosaccharides when cells are cultured on chitin beads, we would expect
bimodality to be lost. When P_{chb} was induced by chitin oligosaccharides or *via*
deletion of *cbp*, all cells in the population are uniformly "P_{chb} On" (**Fig. 2D, 2F**)
and the population becomes unimodal (**Fig. 2E, 2G**). Thus, these results showed
that only the "P_{chb} On" cells have access to chitin oligosaccharides when grown
on chitin beads. These results are consistent with chitin-induction of natural
transformation on chitin beads, where cells display bimodality in gene expression
that is also likely due to heterogeneous access to inducing chitin
oligosaccharides (23).
212
Next, we tested whether HapR influenced activation of P_{chb} on chitin beads by
just assessing the expression level among "P_{chb} On" cells. As observed in bulk
populations, the $\Delta hapR$ strain exhibited a ~1.7 fold increase in P_{chb} expression
when cultured on chitin beads (**Fig. 2C**) and a ~2.5 fold increase when cultured
with chitin oligosaccharides (**Fig. 2E**). These values were consistent with the ~2-
fold increase in P_{chb} expression observed in single cells when the population was
induced via deletion of *cbp* (**Fig. 2G**). Finally, native *chb* transcripts (measured
via qRT-PCR) were also induced ~2-fold higher in a $\Delta hapR$ mutant compared to
the parent when strains were induced with chitin oligosaccharides (**Fig. S2**).
Together, these results indicate that HapR is a *bona fide* repressor of P_{chb} under
physiologically relevant inducing conditions.
224
HapR repression of P_{chb} is conserved in other V. cholerae El Tor isolates
Previously, it was suggested that HapR was an activator of P_{chb} expression in the
V. cholerae El Tor isolate A1552 (16). Specifically, the authors showed that
deletion of *hapR* resulted in a decrease in P_{chb} expression when cells were
cultured on chitin flakes. In the present study, we use the *V. cholerae* El Tor

isolate E7946 to study P_{chb} regulation. To assess if HapR exhibits different
effects on P_{chb} expression depending on the strain background, we assessed
expression of a P_{chb}-mCherry reporter in $hapR^+$ and $\Delta hapR$ derivatives of both
E7946 and A1552. Consistent with our previous results, we observed that P_{chb}
expression is elevated ~2-fold in the $\Delta hapR$ derivative of both strain backgrounds
when induced with chitin oligosaccharides or via deletion of cbp, which is
consistent with HapR acting as a repressor of this locus (**Fig. S3**). It is unclear
what explains the discrepancy between our findings and those that were
previously reported; however, these data suggest that they cannot be attributed
to differences between strain backgrounds.
240
Deletion of HapR does not confer a growth advantage during growth on
chitobiose
Our data indicates that in the absence of HapR, P_{chb} expression is elevated. The
chb locus encodes the genes required for uptake and degradation of the chitin
disaccharide, chitobiose. Thus, we wanted to assess if the increased expression
of P_{chb} confers a fitness advantage to $\Delta hapR$ cells during growth on chitobiose.
To test this, we conducted competitive growth assays with a 1:1 mixture of a
parent and a $\Delta hapR$ mutant strain on minimal medium with chitobiose as the sole
carbon source. We hypothesized that if the $\Delta hapR$ mutant had a competitive
growth advantage due to increased expression of *chb*, then it should outcompete
the parent strain in this assay. Even after ~48 generations of growth on
chitobiose, however, we did not observe a competitive advantage for the $\Delta hapR$
mutant (**Fig. S4**). HapR is a global regulator that controls the expression of
dozens of genes (24). Thus, even though a $\Delta hapR$ mutant has increased P_{chb}
expression, there may be negative pleiotropic effects associated with the $\Delta hapR$
mutation that masks any competitive advantage of derepression of P_{chb} in this
mutant during growth on chitobiose.
258
HapR binds P_{chb} in vitro and in vivo
Thus far, our data suggest that HapR is a repressor of P_{chb}, but it does not
distinguish whether HapR is a direct or indirect regulator of this locus. To assess
if HapR could be regulating P_{chb} directly, we used an *in silico* approach to identify
putative HapR binding sites in P_{chb} based on consensus binding sequences
generated for the HapR homologs LuxR (from *Vibrio harveyi*) and SmcR (from *V.*
vulnificus) via chromatin immunoprecipitation sequencing (ChIP-seq) (25, 26).
Using the Motif Alignment and Search Tool (MAST) in the Multiple Em for Motif
Elicitiation (MEME) Suite (27), we identified two potential HapR binding sites in
P_{chb} (**Fig. 3A**). Interestingly, these binding sites (BSs) overlap with other elements
in the *chb* promoter required for transcriptional activation. HapR BS 1 overlaps
with the SlmA binding site, which is a critical activator of P_{chb} expression (11).
HapR BS 2 overlaps with the -35 signal, which is required for RNA polymerase to
bind to the promoter and initiate transcription. This sequence analysis suggests
that the repressive effect of HapR may be due to HapR binding directly
antagonizing SlmA and/or RNAP at this locus.
275

We next sought to determine whether HapR could directly bind to P$_{chb}$ using both
in vitro and *in vivo* approaches. First, we tested whether HapR could bind to P$_{chb}$
in vitro using EMSA assays. Consistent with the presence of two HapR binding
sites in P$_{chb}$, we observed two shifts by EMSA when using a DNA probe of the
chb promoter (**Fig. 3B**). HapR does not regulate the ClpP promoter and thus did
not bind P$_{clpP}$ (**Fig. 3B**), which is consistent with previous studies (28). EMSAs
were also done using 32 bp probes that encompassed each putative HapR BS
(probe sequences in **Fig. 3A**). We observed that HapR was able to bind to both
probes, suggesting that HapR binds to both HapR BS 1 and HapR BS 2 (**Fig.**
3C). Next, we wanted to assess if HapR bound to P$_{chb}$ *in vivo via* ChIP assays
under physiologically relevant conditions. To that end, we first generated a
FLAG-HapR strain that was functional for regulating P$_{chb}$ expression (**Fig. S5**).
Using this strain in ChIP-qPCR assays, we found that P$_{chb}$ was, indeed, bound by
HapR *in vivo*, while the negative control P$_{clpP}$ locus was not bound by HapR (**Fig.**
3D). Together, these data demonstrate that HapR binds to P$_{chb}$, which suggests
that it is a direct repressor of this locus.
292
Quorum sensing regulates expression of P$_{chb}$ through the cholera-specific
autoinducer CAI-1
HapR is the master regulator of quorum sensing (QS) in *V. cholerae*, and is
highly expressed at high cell density (HCD) (29). Thus far, we have established
that HapR is a repressor of P$_{chb}$ expression. Next, we wanted to probe the role of
QS in regulating P$_{chb}$. To that end, we sought to test the effect of mutations in
genes upstream of HapR in the *V. cholerae* QS cascade on P$_{chb}$ induction and
HapR protein levels.
301
QS in *Vibrio* species is controlled by autoinducer-responsive sensor proteins that
indirectly modulate phosphorylation of the response regulator LuxO, which in turn
indirectly regulates production of HapR (14). When autoinducer concentrations
are low (*i.e.*, at low cell density (LCD)), multiple histidine kinase sensors acts as
kinases (30-32). This results in high levels of phosphorylated LuxO, which
prevents HapR production (**Fig. S6A**) (33). By contrast, at high autoinducer
concentrations (*i.e.*, at HCD), the sensors act as phosphatases, which ultimately
leads to dephosphorylation of LuxO and allows for HapR production (**Fig. S6B**)
(14, 34).
311
First, we assessed the impact of HCD on P$_{chb}$ induction and HapR levels by
deleting *luxO*, which genetically locks cells in a HCD state. Induction of the
parent strain is tested under HCD conditions, thus, as expected, HapR levels
were similar between the parent and the Δ*luxO* mutant; accordingly, expression
of P$_{chb}$ was also similar between the parent and Δ*luxO* (**Fig. 4**). Next, we tested
the effect of LCD on P$_{chb}$ and HapR expression by generating a *luxO^{D47E}* mutant,
which mimics phosphorylated LuxO and genetically locks cells in a LCD state. In
this strain, we saw that P$_{chb}$ expression increased and was correlated with a
decrease in HapR protein levels (**Fig. 4**). Together, these data suggest that
HapR repression of P$_{chb}$ is mediated through the canonical QS circuit.

322
Next, we wanted to move further upstream in the QS regulatory cascade to
address if distinct autoinducers differentially affected expression of P$_{chb}$ and
HapR. There are four parallel histidine kinase sensors that coordinate QS-
dependent control of HapR expression (13); however, the autoinducer signal is
only known for two of these systems. The sensor LuxPQ is responsive to the
inter-species autoinducer AI-2, and the sensor CqsS is responsive to the *V.*
cholerae-specific autoinducer CAI-1 (**Fig. S6**) (35). To assess the role of each
autoinducer in regulation of P$_{chb}$, we made mutations to the synthase genes
responsible for production of each autoinducer. LuxS makes AI-2 (36) and CqsA
makes CAI-1 (35) (**Fig. S6**). In a strain that no longer produces AI-2 (Δ*luxS*),
expression of P$_{chb}$ was similar to that observed in the parent and HapR levels
remained similar in these two strains (**Fig. 4**). By contrast, a strain that is unable
to produce CAI-1 (Δ*cqsA*) had increased expression of P$_{chb}$, likely due to the low
level of HapR produced (**Fig. 4**). The observed decrease in P$_{chb}$ expression in the
Δ*cqsA* background was, indeed, due to a lack of CAI-1 production as
exogenously adding back synthetic CAI-1 restored HapR levels and repression of
P$_{chb}$ to the parent level (**Fig. 4**). In addition, a strain that does not make CAI-1
induced P$_{chb}$ to the same level as a strain that does not produce both CAI-1 or
HapR (**Fig. 4**). This epistasis indicates that CAI-1 production and HapR are
involved in the same regulatory pathway for modulating expression of P$_{chb}$.
These data support previous results, which indicate that CqsS signaling plays a
dominant role in regulating HapR (23, 35, 37, 38). Together, these results
indicate that P$_{chb}$ expression is strongly influenced by QS signaling mediated by
the *V. cholerae*-specific autoinducer CAI-1 and less via the inter-species
autoinducer AI-2.
348
Discussion
Here, we show that HapR acts as a repressor of chitobiose utilization genes in *V.*
cholerae. When this organism forms communities on a chitinous surface, chitin
induces expression of P$_{chb}$ through activation of the chitin sensor ChiS (**Fig. 5**).
V. cholerae then modulates P$_{chb}$ expression depending on the presence of other
V. cholerae cells, which it senses via the *V. cholerae*-specific autoinducer CAI-1.
At low CAI-1 concentrations, P$_{chb}$ expression is high because the repressor HapR
is produced at a low level (**Fig. 5A**). At high CAI-1 concentrations, HapR is
produced at higher levels and can repress P$_{chb}$ expression (**Fig. 5B**). As the
HapR binding sites in P$_{chb}$ overlap with the SlmA binding site and the -35 signal,
it is possible that HapR competes with these activators for binding at P$_{chb}$. Thus,
HapR-mediated repression may occur through antagonism of the binding activity
of SlmA and/or RNAP at P$_{chb}$.
362
When HapR is produced, it gets proteolyzed by the ClpAP machine (**Fig. 5B**).
While the dynamic range of HapR-dependent P$_{chb}$ repression is only ~2-fold
when ClpAP is intact, the dynamic range of repression increases to ~7-fold in the
absence of ClpAP. Thus, it is tempting to speculate that regulation of ClpAP may
modulate P$_{chb}$ expression under some conditions. Currently, however, little is

known about the regulation of ClpAP in *V. cholerae*. ClpAP is induced during
heat shock in *V. vulnificus* (21). Also, in *E. coli*, ClpP expression is induced
during heat shock (39-41). Thus, it is believed that the ClpP protease plays a role
in the heat shock response. In *Bacillus subtilis*, ClpP is upregulated under
various stress conditions, suggesting that this protease may also be induced by a
general stress response (42). Thus, one possibility is that stress-dependent
regulation of ClpAP indirectly regulates P$_{chb}$. It has also been hypothesized that
V. cholerae makes use of proteolysis machines to rapidly respond to changes in
their environment; namely, the transition from the human gut to the aquatic
environment after infection (43). ClpAP acting as an activator of P$_{chb}$ may be a
way for cells to rapidly induce expression of a locus that is not important in the
human gut, but is useful for survival in its aquatic reservoir. Consistent with this
idea, *chb* is induced late in infection, suggesting this is a critical locus for
preparing to re-enter the aquatic environment after infection of a host (44).
382
The mechanisms underlying distinct responses to AI-2 or CAI-1 remain a topic of
interest in QS. It has been shown that the CAI-1 sensor CqsS plays a dominant
role over LuxPQ in modulating HapR levels. Thus, it remains possible that the
presence of CAI-1 allows for more robust regulation of HapR-regulated genes.
CAI-1 has been shown to be critical for expression of the virulence factor *hapA*
(38), natural transformation (15), and for repressing chitobiose utilization as
shown in this study. HapA is a protease that has been implicated in mediating *V.*
cholerae detachment from host epithelial cells, thereby promoting dissemination
of cells back into the aquatic environment (45, 46); natural transformation and
chitin utilization aid in *V. cholerae* fitness in the marine environment. By contrast,
expression of *tcpA*, a protein which contributes to *V. cholerae* pathogenesis in
the human gut, was shown to be primarily regulated by AI-2 (37). It is tempting to
speculate that CAI-1 controls *V. cholerae* behaviors important for marine survival,
whereas AI-2 controls behaviors involved in infection. It has been hypothesized
that sensing both of these autoinducers plays a critical role in biofilm dispersal,
and that only when both are sensed, *V. cholerae* cells are induced to leave a
surface (*i.e.* QS works as a coincidence detector for both signals) (37). Thus,
signaling *via* distinct autoinducers may allow *V. cholerae* to modulate responses
depending on the context of the environment they inhabit.
402
The data we present here suggests that at HCD, *V. cholerae* dampens its
expression of a chitin utilization locus. Below, we speculate on a few reasons
why this regulation may be advantageous. Chitin polymers in the shells of
crustacean zooplankton are long chain polysaccharides in a crystalline insoluble
lattice. In order to be used as a nutrient source, the long chain chitin must be
broken down into smaller, soluble chitin oligosaccharides. *V. cholerae* secretes
chitinases that enzymatically degrade insoluble chitin into soluble chitin
oligosaccharides for uptake and catabolism. The production and secretion of
chitinases is an energetically costly process; thus, liberated chitin is a valuable
"public good" in the context of a chitin biofilm (47). So, it is possible that QS
regulation of chitin utilization allows *V. cholerae* to modulate chitin uptake based

on the composition of its microbial community. When the level of *V. cholerae*
cells (and the corresponding concentration of cholera specific autoinducer CAI-1)
in the community is high, chitin uptake may decrease among individual cells
within the population in an effort to "share" liberated chitin oligosaccharides. By
contrast, when the level of *V. cholerae* in the community is low, sensing of only
the inter-species signal AI-2 (which is produced by many bacterial species) does
not suppress chitin uptake and utilization, thus, allowing the *V. cholerae* within
this population to maximally compete for liberated chitin oligosaccharides.
422
Another possibility is that this regulation allows *V. cholerae* to control its
production of toxic metabolites. In a previous study, it was shown that cells in a
HCD state alter their metabolic flux to produce neutral byproducts as opposed to
organic acids when grown in LB supplemented with a fermentable sugar like
glucose (48). This regulation allows for a more stable community; by contrast,
cells locked in a LCD state will excrete harmful metabolic byproducts, which
leads to the demise of the community (48). It has been shown that *V. cholerae*
excretes ammonium as a potentially toxic byproduct when grown on chitin (16).
Also, because chitin oligosaccharides likely feed into glycolysis, it is possible that
catabolism of chitin results in the production of potentially toxic organic acids.
Thus, repressing chitin uptake and utilization may also help slow the rate of
metabolism to prevent the accumulation of toxic intermediates in a dense
community setting.
436
Materials & Methods
Bacterial strains and culture conditions
V. cholerae strains were routinely grown at 30°C in LB medium and on LB agar
supplemented when necessary with carbenicillin (20 μg/mL or 50 μg/mL),
kanamycin (50 μg/mL), spectinomycin (200 μg/mL), trimethoprim (10 μg/mL),
and/or chloramphenicol (2 μg/mL). Strains were grown in DASW medium for
microscopy (see below for details), Instant Ocean for generating mutant strains
(see below for details), or M9 minimal medium for competition experiments (see
below for details).
446
Transposon mutagenesis
Transposon mutant libraries were generated with a Carb^R mini-Tn10 transposon
exactly as previously described (49). Briefly, the transposon mutagenesis
plasmid pDL1086 was first mated into parent strains containing chromosomally-
integrated P$_{chb}$-*lacZ* and a Δ*cbp* mutation (activator screen) or Δ*clpA* Δ*cbp*
mutations (Δ*clpA* counter-screen). The parent strains also carried a
chromosomally-integrated P$_{chb}$-mCherry reporter at an ectopic site to ensure that
candidate transposon mutants affected expression of P$_{chb}$ and did not simply
disrupt the P$_{chb}$-*lacZ* reporter. The activator screen parent also carried an
additional copy of ChiS at an ectopic site, which prevented transposon hits in this
known regulator of P$_{chb}$. Transposition was induced by plating cultures on LB
agar supplemented with 50 μg/mL Carb at 42°C. To screen colonies, plates also

contained 40 µg/mL X-gal and 5 mM IPTG. IPTG was added to competitively
inhibit the basal activity of the P$_{chb}$-*lacZ* reporter.
461
The sequence of transposon-genomic junctions in transposon mutants were
determined by inverse PCR followed by Sanger sequencing. Briefly, genomic
DNA was purified from mutants of interest and digested with the FatI restriction
enzyme per manufacturer's instructions (NEB). Digested genomic DNAs were
then incubated with T4 DNA ligase per manufacturer's instructions (NEB) to
generate self-ligated circles. The transposon-genomic junction was then
amplified by PCR using the primers specified in **Table 2** and subsequently
Sanger sequenced (Eurofins Genomics).
470
Generating mutant strains
V. cholerae E7946 served as the parent for all strains generated in this study,
except for those used in **Fig. S3** where we compare E7946 to A1552 (50, 51).
Mutant constructs were generated by splicing-by-overlap extension (SOE) PCR
exactly as previously described with the primers indicated in **Table 2**. (52). PCRs
were performed to generate Up (F1/R1), Middle, and Down (F2/R2) arms. Up
and Down arms were designed to have 3kb arms of homology to the genome at
the site targeted for mutagenesis. All three arms were then mixed and used as
template for SOE PCR reactions with the F1 and R2 primers to generate the full
length mutant construct SOE product. Mutant constructs were introduced into *V.*
cholerae cells by chitin-induced natural transformation and/or cotransformation
exactly as previously described (53, 54) or by chitin-independent transformations
using a plasmid that ectopically expresses *tfoX* and *qstR* as previously described
(55). For chitin-dependent natural transformation, *V. cholerae* was grown to mid-
log in LB medium, washed with instant ocean medium (7 g/L; Aquarium
Systems), and incubated with chitin flakes (Alfa Aesar) at final OD$_{600}$ of 0.1
overnight at 30°C. The next day, SOE PCR products were added to chitin
induced cells, incubated at 30°C for 5 hrs, and then outgrown and plated on
selective media. For chitin-independent transformation, cells harboring
pMMB67EH-*tfoX-qstR* were grown overnight in LB medium supplemented with
100 µM IPTG and 1 µg/mL chloramphenicol. The next day, 7 µL of the overnight
culture was diluted into 350 µL of instant ocean medium. Then, SOE products
were added as described above for chitin-dependent transformations. Mutant
strains were confirmed by colony PCR, MAMA PCR (56), and/or sequencing. A
complete list of strains and primers used in this study are outlined in **Table 1** and
Table 2, respectively.
497
Measuring GFP and mCherry fluorescence in bulk populations
GFP and mCherry fluorescence in reporter strains was determined exactly as
previously described (57). Briefly, single colonies were grown in LB overnight at
30°C. Where indicated, media was supplemented with 10 µM CAI-1. CAI-1 was
synthesized exactly as previously described (58). The next day, cells were
washed and resuspended to an OD$_{600}$ of 1.0 in instant ocean medium (7 g/L;

Aquarium systems). Fluorescence was determined on a BioTek H1M plate
reader. For GFP measurements, excitation was set to 500 nm and emission was
set to 540 nm; for mCherry measurements, excitation was set to 580 nm and
emission was set to 610 nm.
508
509 *Antibody generation*
510 Purified *Vibrio harveyi* LuxR protein (300 µg; purified as previously described
511 (59)) was sent to Cocalico Biologicals Inc. for serial injection into a rabbit host for
512 antibody generation. Serum obtained from the third bleed has been used for
513 Western analyses.
514
Western blotting
From overnight cultures, cells were concentrated to an OD_{600} of 20 in instant
ocean medium. Cells were lysed on a FastPrep-24 Classic Instrument at 4°C,
then lysates were clarified by centrifugation. Lysates were then boiled with an
equal volume of 2x SDS PAGE sample buffer (220 mM Tris pH 6.8, 25%
glycerol, 1.2% SDS, 0.02% Bromophenol Blue, and 5% β-mercaptoethanol).
Proteins were separated on a 15% SDS polyacrylamide gel by SDS
electrophoresis, electropheretically transferred to a PVDF membrane, and
probed with rabbit polyconal α-LuxR serum or mouse monoclonal α-RpoA
(Biolegend) primary antibodies. LuxR is the HapR homolog in *Vibrio harveyi* and
has 72% identity and 86% similarity to *V. cholerae* HapR. The LuxR antibody
was empirically found to be cross-reactive with *V. cholerae* HapR, and so it was
used to detect HapR protein levels. Blots were then incubated with α-rabbit or α-
mouse HRP conjugated secondary antibodies, developed using Pierce ECL
Western Blotting Substrate (ThermoFisher), and exposed to film.
530
Chitin bead culturing
Chitin beads (200 µL of a 50% slurry; NEB) and overnight cultures of the
indicated strains were washed with defined artificial salt water medium (DASW),
which was prepared exactly as previously described (8). Chitin beads were
inoculated with *V. cholerae* cells to an OD_{600} of 0.1 in a final volume of 5 mL in a
Costar 6-well plate (Corning). Chitin mixtures were incubated statically at 30°C
for 7 days before imaging.
538
Microscopy data collection and analysis
To image chitin beads, cultured beads were gently transferred to a coverslip
using wide-bore pipette tips. To image chitin grown cells, chitin bead reactions
were vortexed then centrifuged at 250 x g for 1 minute. Cells found in the
supernatant were transferred to a coverslip. To image Δ*cbp* strains, overnight
cultures grown in LB were washed and resuspended to an OD_{600} of 0.2 in DASW
and then transferred to a coverslip. Samples on the coverslip were placed under
a 0.2% gelzan pad and imaged on an inverted Nikon Ti-2 microscope with a Plan
Apo 60x objective, YFP & mCherry filter cubes, a Hamamatsu ORCAFlash 4.0
camera, and Nikon NIS Elements imaging software.
549

The strains used to examine P$_{chb}$ expression across a population contained 1) a
P$_{chb}$-mCherry reporter and 2) a reporter that drove constitutive expression of
GFP, P$_{const2}$-GFP. By using these reporters, the expression of the P$_{chb}$-mCherry
could be normalized to GFP expression in each cell. Images were analyzed on
Fiji using the MicrobeJ plugin (60) to determine mean intensity of cells in the YFP
and mCherry channels. GFP was assessed using a YFP filter set to avoid
background fluorescence from chitin beads, which was stronger in the GFP
channel. Background fluorescence was subtracted from each channel and the
mCherry / GFP fluorescence was determined for each individual cell.
559
HapR protein purification
A plasmid expressing a hexahistidine-tagged *hapR* wild-type allele was
generated using Gibson assembly. The *hapR* insert was amplified from *V.*
cholerae E7946 and inserted into a pET28b vector (Novagen) using the primers
listed in **Table 2**. The plasmid was then electroporated into *E. coli* BL21(DE3) for
protein overexpression. This strain was grown overnight in LB medium with
kanamycin, back-diluted 1:100 into 1 L of LB medium with kanamycin, and grown
to an OD$_{600}$ of 0.4-0.6 at 30°C. Expression of HapR was induced by IPTG to a
final concentration of 1 mM and cultures grown for 4 h shaking at 30°C. The cells
were pelleted and frozen at -80°C. The pellet was resuspended in 25 mL Buffer A
(25 mM Tris pH 8, 500 mM NaCl), and an Avestin EmulfiFlex-C3 emulsifier was
used to lyse cells. The soluble lysate was applied to a HisTrap HP Ni-NTA
column using an Äkta Pure FPLC in Buffer A and eluted from the column with a
gradient of Buffer B (25 mM Tris pH 8, 500 mM NaCl, 1 M imidazole). The
purified protein was concentrated to approximately 5 ml using Sartorius Vivaspin
Turbo 10,000 MWCO centrifugal concentrators. The sample was manually
injected into the Äkta Pure and separated via size exclusion chromatography on
a HiLoad™ 16/600 Superdex™ 75 pg column equilibrated with gel filtration buffer
(25 mM Tris pH 7.5, 200 mM NaCl). Eluted fractions were analyzed by SDS-
PAGE (15% gel), pooled, and concentrated using the same centrifugal
concentrators previously mentioned. The samples were then immediately frozen
in liquid nitrogen with a final concentration of 10% glycerol and stored at -80°C.
582
Electrophoretic mobility shift assay (EMSA)
Binding reactions contained 10 mM Tris HCl pH 7.5, 1 mM EDTA, 10 mM KCl, 1
mM DTT, 50 µg/mL BSA, 0.1 mg/mL salmon sperm DNA, 5% glycerol, 1 nM of a
Cy5 labeled DNA probe, and HapR at the indicated concentrations (diluted in 10
mM Tris pH 7.5, 10 mM KCl, 1 mM DTT, and 5% glycerol). Reactions were
incubated at room temperature for 20 minutes in the dark, then
electrophoretically separated on polyacrylamide gels in 0.5x Tris Borate EDTA
(TBE) buffer. Gels were imaged for Cy5 fluorescence on a Typhoon-9210
instrument. Short DNA probes (30bp) were made by end-labeling one primer of a
complementary pair (primers listed in **Table 2**) using 20 µM Cy5-dCTP and
Terminal deoxynucleotidyl Transferase (TdT; Promega). Complementary primers
(one labeled with Cy5 and the other unlabeled) were annealed by slow cooling at
equimolar concentrations in annealing buffer (10 mM Tris pH 7.5 and 50 mM

NaCl). P$_{chb}$ and P$_{clpP}$ probes were made by Phusion PCR, where Cy5-dCTP was
included in the reaction at a level that would result in incorporation of 1–2 Cy5
labeled nucleotides in the final probe as previously described.
599
ChIP-qPCR assays
Assays were carried out exactly as previously described (10). Briefly, overnight
cultures were diluted to an OD$_{600}$ of 0.08 and then grown for 6 hours at 30°C.
Cultures were crosslinked using 1% paraformaldehyde, then quenched with a 1.2
molar excess of Tris. Cells were washed with PBS and stored at -80°C overnight.
The next day, cells were resuspended in lysis buffer (1x FastBreak cell lysis
reagent (Promega), 50 µg/mL lysozyme, 1% Triton X-100, 1 mM PMSF, and 1x
protease inhibitor cocktail; 100x inhibitor cocktail contained the following: 0.07
mg/mL phosphoramidon (Santa Cruz), 0.006 mg/mL bestatin
(MPbiomedicals/Fisher Scientific), 1.67 mg/mL AEBSF (DOT Scientific), 0.07
mg/mL pepstatin A (Gold Bio), 0.07 mg/mL E64 (Gold Bio)) and then lysed by
sonication, resulting in a DNA shear size of ~500 bp. Lysates were incubated
with Anti-FLAG M2 Magnetic Beads (Sigma), washed to remove unbound
proteins, and then bound protein-DNA complexes were eluted off with SDS.
Samples were digested with Proteinase K, then crosslinks were reversed. DNA
samples were cleaned up and used as template for quantitative PCR (qPCR)
using iTaq Universal SYBR Green Supermix (Bio-Rad) and primers specific for
the genes indicated (primers are listed in **Table 2**) on a Step-One qPCR system.
Standard curves of genomic DNA were included in each experiment and were
used to determine the abundance of each amplicon in the input (derived from the
lysate prior to ChIP) and output (derived from the samples after ChIP). Primers to
amplify *rpoB* served as a baseline control in this assay because HapR does not
bind this locus. Data are reported as 'Fold Enrichment', which is defined as the
ratio of the test promoter (P$_{chb}$ or P$_{clpP}$) / *rpoB* found in the output divided by the
same ratio found in the input.
625
Chitobiose competition
Overnight cultures grown in LB were washed with M9 medium and mixed 1:1.
For each growth reaction, 10^2 cells of this mixture was added to M9 medium
supplemented with 0.2% chitobiose and 10 µM synthetic CAI-1 and grown
shaking at 30°C for 24 hours. After 24 hours, ~10^2 cells from this mixture was
used to inoculate fresh growth reactions to achieve additional generations of
growth on chitobiose. This was repeated a third time after another 24 hours. CAI-
1 was supplemented throughout these experiments to ensure consistently high
levels of HapR expression throughout transfer steps. After each 24 hour growth
period, the CFU/mL was determined for each strain in the mixture by dilution
plating on LB agar supplemented with X-gal. Competing strains were discerned
by blue/white screening as one strain was *lacZ*$^+$ and the other was Δ*lacZ*. A *lacZ*$^+$
strain was competed against a Δ*lacZ* strain (parent : parent competition) or
against a Δ*lacZ* Δ*hapR* strain (Δ*hapR* : parent competition). *V. cholerae* can grow
on chitobiose through the activity of two sugar transporters: the GlcNAc PTS
transporter (VC0995) or the chitobiose ABC transporter (VC0616-0620).

Because we wanted to study the regulation of the transporter encoded within the
chb locus, all strains for this assay had a deletion in VC0995, which renders
growth on chitobiose dependent on the *chb* locus as previously described (11).
Competitive indices were calculated as the CFU ratio of $\Delta lacZ$ / $lacZ^+$ after
growth for the indicated number of generations divided by the CFU ratio of the
$\Delta lacZ$ / $lacZ+$ in the initial inoculum.
648
649 *Statistics*
650 Statistical comparisons were determined using Student's t-test or One-way
651 ANOVA with Tukey's post-test using GraphPad Prism software.
652

653 **Acknowledgements**
654 We thank Ryan Chaparian for offering experimental advice, David Grainger for
655 helpful discussions, and Wai-Leung Ng for providing synthetic CAI-1. This work
656 was supported by grant R35GM128674 from the National Institutes of Health to
657 ABD and R35GM124698 from the National Institutes of Health to JVK.

658
659

References

1. Pruzzo C, Vezzulli L, & Colwell RR (2008) Global impact of *Vibrio cholerae* interactions with chitin. *Environ Microbiol* 10(6):1400-1410.

2. Huq A*, et al.* (1983) Ecological relationships between *Vibrio cholerae* and planktonic crustacean copepods. *Appl Environ Microbiol* 45(1):275-283.

3. Nahar S*, et al.* (2011) Role of Shrimp Chitin in the Ecology of Toxigenic *Vibrio cholerae* and Cholera Transmission. *Front Microbiol* 2:260.

4. Hunt DE, Gevers D, Vahora NM, & Polz MF (2008) Conservation of the chitin utilization pathway in the *Vibrionaceae*. *Appl Environ Microbiol* 74(1):44-51.

5. Colwell RR*, et al.* (2003) Reduction of cholera in Bangladeshi villages by simple filtration. *Proc Natl Acad Sci U S A* 100(3):1051-1055.

6. Huq A*, et al.* (2010) Simple sari cloth filtration of water is sustainable and continues to protect villagers from cholera in Matlab, Bangladesh. *mBio* 1(1).

7. Meibom KL*, et al.* (2004) The *Vibrio cholerae* chitin utilization program. *Proc Natl Acad Sci U S A* 101(8):2524-2529.

8. Meibom KL, Blokesch M, Dolganov NA, Wu CY, & Schoolnik GK (2005) Chitin induces natural competence in *Vibrio cholerae*. *Science* 310(5755):1824-1827.

9. Li X & Roseman S (2004) The chitinolytic cascade in *Vibrios* is regulated by chitin oligosaccharides and a two-component chitin catabolic sensor/kinase. *Proc Natl Acad Sci U S A* 101(2):627-631.

10. Klancher CA, Yamamoto S, Dalia TN, & Dalia AB (2020) ChiS is a noncanonical DNA-binding hybrid sensor kinase that directly regulates the chitin utilization program in *Vibrio cholerae*. *Proc Natl Acad Sci U S A* In press.

11. Klancher CA, Hayes CA, & Dalia AB (2017) The nucleoid occlusion protein SlmA is a direct transcriptional activator of chitobiose utilization in *Vibrio cholerae*. *PLoS Genet* 13(7):e1006877.

12. Fuqua WC, Winans SC, & Greenberg EP (1994) Quorum sensing in bacteria: the LuxR-LuxI family of cell density-responsive transcriptional regulators. *J Bacteriol* 176(2):269-275.

13. Jung SA, Chapman CA, & Ng WL (2015) Quadruple quorum-sensing inputs control *Vibrio cholerae* virulence and maintain system robustness. *PLoS Pathog* 11(4):e1004837.

14. Ball AS, Chaparian RR, & van Kessel JC (2017) Quorum Sensing Gene Regulation by LuxR/HapR Master Regulators in *Vibrios*. *J Bacteriol* 199(19).

15. Suckow G, Seitz P, & Blokesch M (2011) Quorum sensing contributes to natural transformation of *Vibrio cholerae* in a species-specific manner. *J Bacteriol* 193(18):4914-4924.

16. Sun S, Tay QX, Kjelleberg S, Rice SA, & McDougald D (2015) Quorum sensing-regulated chitin metabolism provides grazing resistance to *Vibrio cholerae* biofilms. *ISME J* 9(8):1812-1820.

17. Yamamoto S, *et al.* (2014) Regulation of natural competence by the
orphan two-component system sensor kinase ChiS involves a non-
canonical transmembrane regulator in *Vibrio cholerae*. *Molecular*
microbiology 91(2):326-347.
18. Yamamoto S & Ohnishi M (2017) Glucose-Specific Enzyme IIA of the
Phosphoenolpyruvate:Carbohydrate Phosphotransferase System
Modulates Chitin Signaling Pathways in *Vibrio cholerae*. *J Bacteriol*
199(18).
19. Blokesch M (2012) Chitin colonization, chitin degradation and chitin-
induced natural competence of Vibrio cholerae are subject to catabolite
repression. *Environ Microbiol* 14(8):1898-1912.
20. Sauer RT & Baker TA (2011) AAA+ proteases: ATP-fueled machines of
protein destruction. *Annu Rev Biochem* 80:587-612.
21. Lee KJ, Jung YC, Park SJ, & Lee KH (2018) Role of Heat Shock
Proteases in Quorum-Sensing-Mediated Regulation of Biofilm Formation
by *Vibrio* Species. *mBio* 9(1).
22. Davis JH, Rubin AJ, & Sauer RT (2011) Design, construction and
characterization of a set of insulated bacterial promoters. *Nucleic Acids*
Res 39(3):1131-1141.
23. Lo Scrudato M & Blokesch M (2012) The regulatory network of natural
competence and transformation of *Vibrio cholerae*. *PLoS Genet*
8(6):e1002778.
24. Zhu J & Mekalanos JJ (2003) Quorum sensing-dependent biofilms
enhance colonization in *Vibrio cholerae*. *Dev Cell* 5(4):647-656.
25. van Kessel JC, Ulrich LE, Zhulin IB, & Bassler BL (2013) Analysis of
activator and repressor functions reveals the requirements for
transcriptional control by LuxR, the master regulator of quorum sensing in
Vibrio harveyi. *mBio* 4(4).
26. Lee DH, *et al.* (2008) A consensus sequence for binding of SmcR, a *Vibrio*
vulnificus LuxR homologue, and genome-wide identification of the SmcR
regulon. *J Biol Chem* 283(35):23610-23618.
27. Bailey TL, *et al.* (2009) MEME SUITE: tools for motif discovery and
searching. *Nucleic Acids Res* 37(Web Server issue):W202-208.
28. Chaparian RR, Tran MLN, Miller Conrad LC, Rusch DB, & van Kessel JC
(2020) Global H-NS counter-silencing by LuxR activates quorum sensing
gene expression. *Nucleic Acids Res* 48(1):171-183.
29. Jobling MG & Holmes RK (1997) Characterization of *hapR*, a positive
regulator of the *Vibrio cholerae* HA/protease gene hap, and its
identification as a functional homologue of the *Vibrio harveyi* luxR gene.
Molecular microbiology 26(5):1023-1034.
30. Freeman JA, Lilley BN, & Bassler BL (2000) A genetic analysis of the
functions of LuxN: a two-component hybrid sensor kinase that regulates
quorum sensing in *Vibrio harveyi*. *Molecular microbiology* 35(1):139-149.
31. Freeman JA & Bassler BL (1999) Sequence and function of LuxU: a two-
component phosphorelay protein that regulates quorum sensing in *Vibrio*
harveyi. *J Bacteriol* 181(3):899-906.

32. Neiditch MB, Federle MJ, Miller ST, Bassler BL, & Hughson FM (2005) Regulation of LuxPQ receptor activity by the quorum-sensing signal autoinducer-2. *Mol Cell* 18(5):507-518.

33. Tu KC & Bassler BL (2007) Multiple small RNAs act additively to integrate sensory information and control quorum sensing in *Vibrio harveyi*. *Genes Dev* 21(2):221-233.

34. Freeman JA & Bassler BL (1999) A genetic analysis of the function of LuxO, a two-component response regulator involved in quorum sensing in *Vibrio harveyi*. *Molecular microbiology* 31(2):665-677.

35. Miller MB, Skorupski K, Lenz DH, Taylor RK, & Bassler BL (2002) Parallel quorum sensing systems converge to regulate virulence in *Vibrio cholerae*. *Cell* 110(3):303-314.

36. Surette MG, Miller MB, & Bassler BL (1999) Quorum sensing in *Escherichia coli*, *Salmonella typhimurium*, and *Vibrio harveyi*: a new family of genes responsible for autoinducer production. *Proc Natl Acad Sci U S A* 96(4):1639-1644.

37. Bridges AA & Bassler BL (2019) The intragenus and interspecies quorum-sensing autoinducers exert distinct control over *Vibrio cholerae* biofilm formation and dispersal. *PLoS Biol* 17(11):e3000429.

38. Hurley A & Bassler BL (2017) Asymmetric regulation of quorum-sensing receptors drives autoinducer-specific gene expression programs in *Vibrio cholerae*. *PLoS Genet* 13(5):e1006826.

39. Kroh HE & Simon LD (1990) The ClpP component of Clp protease is the sigma 32-dependent heat shock protein F21.5. *J Bacteriol* 172(10):6026-6034.

40. Grossman AD, Erickson JW, & Gross CA (1984) The *htpR* gene product of *E. coli* is a sigma factor for heat-shock promoters. *Cell* 38(2):383-390.

41. Katayama Y, *et al.* (1988) The two-component, ATP-dependent Clp protease of *Escherichia coli*. Purification, cloning, and mutational analysis of the ATP-binding component. *J Biol Chem* 263(29):15226-15236.

42. Gerth U, Kruger E, Derre I, Msadek T, & Hecker M (1998) Stress induction of the *Bacillus subtilis clpP* gene encoding a homologue of the proteolytic component of the Clp protease and the involvement of ClpP and ClpX in stress tolerance. *Molecular microbiology* 28(4):787-802.

43. Pennetzdorfer N, *et al.* (2019) Regulated proteolysis in *Vibrio cholerae* allowing rapid adaptation to stress conditions. *Front Cell Infect Microbiol* 9:214.

44. Schild S, *et al.* (2007) Genes induced late in infection increase fitness of *Vibrio cholerae* after release into the environment. *Cell Host Microbe* 2(4):264-277.

45. Benitez JA, *et al.* (1997) Adherence of *Vibrio cholerae* to cultured differentiated human intestinal cells: an in vitro colonization model. *Infect Immun* 65(8):3474-3477.

46. Benitez JA, Silva AJ, & Finkelstein RA (2001) Environmental signals controlling production of hemagglutinin/protease in *Vibrio cholerae*. *Infect Immun* 69(10):6549-6553.

798   47.   Drescher K, Nadell CD, Stone HA, Wingreen NS, & Bassler BL (2014)
799   Solutions to the public goods dilemma in bacterial biofilms. *Curr Biol*
800   24(1):50-55.
801   48.   Hawver LA, Giulietti JM, Baleja JD, & Ng WL (2016) Quorum sensing
802   coordinates cooperative expression of pyruvate metabolism genes To
803   maintain a sustainable environment for population stability. *mBio* 7(6).
804   49.   McDonough E, Kamp H, & Camilli A (2016) *Vibrio cholerae* phosphatases
805   required for the utilization of nucleotides and extracellular DNA as
806   phosphate sources. *Molecular microbiology* 99(3):453-469.
807   50.   Miller VL, DiRita VJ, & Mekalanos JJ (1989) Identification of *toxS*, a
808   regulatory gene whose product enhances *toxR*-mediated activation of the
809   cholera toxin promoter. *J Bacteriol* 171(3):1288-1293.
810   51.   Yildiz FH, Liu XS, Heydorn A, & Schoolnik GK (2004) Molecular analysis
811   of rugosity in a *Vibrio cholerae* O1 El Tor phase variant. *Molecular*
812   *microbiology* 53(2):497-515.
813   52.   Dalia AB, Lazinski DW, & Camilli A (2013) Characterization of
814   undermethylated sites in *Vibrio cholerae*. *J Bacteriol* 195(10):2389-2399.
815   53.   Dalia AB, McDonough E, & Camilli A (2014) Multiplex genome editing by
816   natural transformation. *Proc Natl Acad Sci U S A* 111(24):8937-8942.
817   54.   Dalia AB (2018) Natural Cotransformation and Multiplex Genome Editing
818   by Natural Transformation (MuGENT) of *Vibrio cholerae*. *Methods Mol Biol*
819   1839:53-64.
820   55.   Simpson CA, Podicheti R, Rusch DB, Dalia AB, & van Kessel JC (2019)
821   Diversity in natural transformation frequencies and regulation across
822   *Vibrio* species. *mBio* 10(6).
823   56.   Cha RS, Zarbl H, Keohavong P, & Thilly WG (1992) Mismatch
824   amplification mutation assay (MAMA): application to the *c-H-ras* gene.
825   *PCR Methods Appl* 2(1):14-20.
826   57.   Dalia AB (2016) RpoS is required for natural transformation of *Vibrio*
827   *cholerae* through regulation of chitinases. *Environ Microbiol* 18(11):3758-
828   3767.
829   58.   Ng WL*, et al.* (2011) Signal production and detection specificity in *Vibrio*
830   CqsA/CqsS quorum-sensing systems. *Molecular microbiology* 79(6):1407-
831   1417.
832   59.   Chaparian RR, Olney SG, Hustmyer CM, Rowe-Magnus DA, & van Kessel
833   JC (2016) Integration host factor and LuxR synergistically bind DNA to
834   coactivate quorum-sensing genes in *Vibrio harveyi*. *Molecular*
835   *microbiology* 101(5):823-840.
836   60.   Ducret A, Quardokus EM, & Brun YV (2016) MicrobeJ, a tool for high
837   throughput bacterial cell detection and quantitative analysis. *Nat Microbiol*
838   1(7):16077.
839
840

Figure 1. *HapR is a repressor of P$_{chb}$ that is degraded by the ClpAP protease.*
Expression of a P$_{chb}$-GFP reporter and HapR protein levels were determined in
the indicated mutant strains. The parent strain contained a P$_{chb}$-GFP reporter and
a Δcbp mutation. A representative western blot is shown below bars to indicate
the protein levels for HapR and RpoA (a loading control) in the corresponding
strains. An antibody against LuxR, which has 72% identity and 86% similarity to
HapR, is cross-reactive with HapR and so was used to detect HapR protein
levels. Fluorescence of cultures was determined on a plate reader from at least
six independent biological replicates and is shown as the mean ± SD. Statistical
comparisons were made by one-way ANOVA with Tukey's post-test. NS, not
significant. ***, p < 0.001.

 Figure 2. *HapR-mediated repression of P_chb occurs on chitinous surfaces.* **A**)
 Representative images of the indicated *V. cholerae* strains grown on chitin

beads. The parent strain background contains both a P$_{chb}$-mCherry reporter and
a P$_{const2}$-GFP construct, which exhibits constitutive GFP expression. Scale bar =
10 µm. **B-C**) P$_{chb}$ expression in the indicated *V. cholerae cbp^{+}* strains grown on
chitin beads. **B**) Representative images of cells that were vortexed off chitin
beads. Arrows demarcate cells where P$_{chb}$ expression is induced ("P$_{chb}$ On"
cells). Scale bar = 2 µm. **C**) Scatter plot and bar graph showing the relative
expression of the P$_{chb}$-mCherry reporter in cells cultured on chitin beads. Scatter
plots (left Y-axis) represent the entire population, whereas the bars (right Y-axis)
represent only the "P$_{chb}$ On" cells (bracketed in black on scatter plots). The
percentage of cells in the "P$_{chb}$ On" population is indicated. n = 2120 for Parent; n
= 2191 for *ΔhapR;* n = 333 for Parent P$_{chb}$ On; n = 1139 for *ΔhapR* P$_{chb}$ On. Data
shown are from two independent biological replicates. **D-E**) P$_{chb}$ expression of
the indicated *V. cholerae* strains where P$_{chb}$ expression is induced by chitin
oligosaccharides. **D**) Representative images of cells grown with chitin
oligosaccharides. Scale bar = 2 µm. **E**) Scatter plot showing the relative
expression of a P$_{chb}$-mCherry reporter in the indicated *V. cholerae* strains. n =
2735 for Parent; n = 2384 for *ΔhapR*. Data shown are from two independent
biological replicates. **F-G**) P$_{chb}$ expression of the indicated *V. cholerae* strains
where P$_{chb}$ expression is induced via deletion of *cbp*. **F**) Representative images
of cells grown in the absence of chitin. Scale bar = 2 µm. **G**) Scatter plot showing
the relative expression of a P$_{chb}$-mCherry reporter in the indicated *V. cholerae*
strains. n = 2407 for *Δcbp*; n = 2313 for *ΔhapR Δcbp*. Data shown are from two
independent biological replicates. Statistical comparisons in **D**, **G**, and **J** were
made using Student's t-test. ***, p < 0.001.
897

898
899
900
Figure 3. *HapR binds P$_{chb}$ in vitro and in vivo.* **A**) Promoter map of P$_{chb}$
highlighting putative HapR binding sites (BSs). Other sites required for P$_{chb}$
activation (the SlmA BS and the -35 and -10 signals) are highlighted. The exact
sequences of the region containing the HapR BSs (bolded), the SlmA BS, and
the -35 signal are shown. The transcriptional start site (+1) and the translational
start site (+137) are also shown. **B**) A representative EMSA using HapR and
Cy5-labeled DNA probes of the indicated promoters. The P$_{chb}$ probe was
incubated with (from left to right) 0 nM (-), 12.5 nM, 25 nM, 50 nM, 100 nM, 200
nM, 400 nM, or 800 nM purified HapR. The P$_{clpP}$ probe was incubated with 0 nM
(-) or 800 nM HapR (+). **C**) A representative EMSA using HapR and 32 bp Cy5-
labeled probes encompassing each of the putative HapR binding sites within the
chb promoter (exact probe sequences are shown in **A**). The 32 bp probes were
incubated with (from left to right) 0 nM (-), 100 nM, 200 nM, 400 nM, 800 nM, 1.6
µM, or 3.2 µM purified HapR. **D**) ChIP-qPCR assays showing enrichment of the
indicated promoters relative to *rpoB*, a reference locus that HapR does not bind
to. Data are from five independent biological replicates and shown as the mean ±
SD. Statistical comparisons were made by Student's t-test. *, p = 0.0240.
918

Figure 4. *Quorum sensing regulates expression of P$_{chb}$ through the cholera-*
specific autoinducer CAI-1. **A**) Model of QS regulation in *V. cholerae*. There are
four sensor kinases that contribute to canonical QS in *V. cholerae* to regulate
HapR production: LuxQ, CqsS, VpsS, and CqsR. The signals for VpsS and CqsR
are unknown. LuxPQ senses the interspecies autoinducer AI-2 and CqsS senses
the *V. cholerae*-specific autoinducer CAI-1. AI-2 is produced by the LuxS
synthase and CAI-1 is produced by the CqsA synthase. (Left) At low cell density,
the LuxQ and CqsS sensors act as kinases, resulting in phosphorylation of LuxO,
which subsequently decreases expression of HapR. (Right) At high cell density
when concentrations of AI-2 and CAI-1 are high, LuxQ and CqsS act as
phosphatases, which results in dephosphorylation of LuxO, and subsequent
activation of HapR expression. **B**) Expression of a P$_{chb}$-GFP reporter and HapR
protein levels were determined in the indicated strains. The parent strain contains
a P$_{chb}$-GFP reporter and a Δ*cbp* mutation. A representative western blot is shown
below the bars to indicate the protein levels for HapR and RpoA (a loading
control) in the corresponding strains. An antibody against LuxR, which has 72%
identity and 86% similarity to HapR, is cross-reactive with HapR and so was used
to detect HapR protein levels. Statistical comparisons were made by one-way
ANOVA with Tukey's post-test. NS, not significant. ***, $p < 0.001$. The P$_{chb}$-GFP
fluorescence data for "Parent" and "Δ*clpA*" are identical to the data presented in
Fig. 1 and were included here to ease comparisons.
953

Figure 5. *Model of quorum sensing regulation of chitobiose utilization genes in V.*
cholerae. When *V. cholerae* grows on a chitinous surface, the periplasmic chitin
binding protein (CBP) binds to chitin oligosaccharides. This de-represses ChiS,
which can then activate expression of the chitobiose utilization operon (*chb*)
through recruitment of RNA Polymerase (RNAP) (10). Binding of SlmA to P$_{chb}$ is
also required for transcriptional activation (11). **A**) At low CAI-1 concentrations,
the CqsS sensor indirectly (dashed line) promotes LuxO phosphorylation, which
indirectly blocks HapR expression. In this state, P$_{chb}$ is maximally expressed. **B**)
At high CAI-1 concentrations, the CqsS sensor indirectly dephosphorylates
LuxO, which results in indirect activation of HapR expression. HapR then exerts
a repressive effect on P$_{chb}$ by decreasing expression of this locus ~2-fold in the
presence of the ClpAP protease machine. If HapR is not degraded by ClpAP, its
repressive effect can result in a ~7-fold decrease in P$_{chb}$ expression. The
mechanism of HapR repression may be through occluding binding of SlmA or
RNAP to P$_{chb}$.

974 **Table 1.** *Strains used in this study.*

Strain Designation	Reference in Manuscript	Genotype	Reference
SAD 030	*V. cholerae* E7946 WT; parent for all other E7946 strains; Fig. S2 Parent	*V. cholerae* E7946 O1 El Tor	(50)
SAD 2825	Activator screen strain	*V. cholerae* E7946 ΔVC1807::P$_{chb}$-mCherry, Kan^R; P$_{chb}$-*lacZ*, Spec^R; ΔVCA0692::*chiS*, Tm^R; Δ3' *cbp*, pDL1086 Cm^R Carb^R	This study
SAD 2826	Δ*clpA* counter-screen strain	*V. cholerae* E7946 ΔVC1807::P$_{chb}$-mCherry, Kan^R; P$_{chb}$-*lacZ*, Spec^R; Δ*clpA*; Δ*cbp*::Tm^R; pDL1086, Cm^R Carb^R	This study
SAD 1309	Fig. 1 & 4 Parent; Fig. S5 HapR WT	*V. cholerae* E7946 Δ*lacZ*:: P$_{chb}$-GFP, Kan^R; Δ*cbp*::Spec^R	(11)
SAD 2827	Fig. 1 & 4 Δ*clpA*	*V. cholerae* E7946 Δ*lacZ*:: P$_{chb}$-GFP, Kan^R; Δ*clpA*::Carb^R; Δ*cbp*::Spec^R	This study
SAD 2828	Fig. 1 Δ*clpP*	*V. cholerae* E7946 Δ*lacZ*:: P$_{chb}$-GFP, Kan^R; Δ*clpP*::Carb^R; Δ*cbp*::Tm^R	This study
SAD 2829	Fig. 1 Δ*clpAP*	*V. cholerae* E7946 Δ*lacZ*:: P$_{chb}$-GFP, Kan^R; Δ*clpA*::Carb^R; Δ*clpP*::Tm^R; Δ*cbp*::Spec^R	This study
SAD 2830	Fig. 1 Δ*clpS*	*V. cholerae* E7946 Δ*lacZ*:: P$_{chb}$-GFP, Kan^R; Δ*clpS*::Carb^R; Δ*cbp*::Tm^R	This study
SAD 2831	Fig. 1 Δ*clpX*	*V. cholerae* E7946 Δ*lacZ*:: P$_{chb}$-GFP, Kan^R; Δ*clpX*::Carb^R; Δ*cbp*::Tm^R	This study
SAD 2832	Fig. 1 Δ*lonA*	*V. cholerae* E7946 Δ*lacZ*:: P$_{chb}$-GFP, Kan^R; Δ*lonA*::Tm^R; Δ*cbp*::Carb^R	This study
SAD 2833	Fig. 1 Δ*hapR* Δ*clpA*	*V. cholerae* E7946 Δ*lacZ*:: P$_{chb}$-GFP, Kan^R; Δ*hapR*::Spec^R; Δ*clpA*::Carb^R; Δ*cbp*::Tm^R	This study
SAD 2834	Fig. 1 Δ*hapR*	*V. cholerae* E7946 Δ*lacZ*:: P$_{chb}$-GFP, Kan^R; Δ*hapR*::Spec^R; Δ*cbp*::Carb^R	This study
SAD 2835	Fig. 2 Parent	*V. cholerae* E7946 Δ*lacZ*::P$_{chb}$-mCherry, Kan^R; ΔVCA0692::P$_{const2}$-GFP, Spec^R	This study
SAD 2836	Fig. 2 Δ*hapR*	*V. cholerae* E7946 Δ*lacZ*::P$_{chb}$-mCherry, Kan^R; ΔVCA0692::P$_{const2}$-GFP, Spec^R; Δ*hapR*::Cm^R	This study
SAD 2837	Fig. 2 Δ*cbp* & Fig. S3 E7946 Parent	*V. cholerae* E7946 Δ*lacZ*::P$_{chb}$-mCherry, Kan^R; ΔVCA0692::P$_{const2}$-GFP, Spec^R; Δ*cbp*::Tm^R	This study
SAD 2838	Fig. 2 Δ*hapR* Δ*cbp* & Fig. S3 E7946 Δ*hapR*	*V. cholerae* E7946 Δ*lacZ*::P$_{chb}$-mCherry, Kan^R; ΔVCA0692::P$_{const2}$-GFP, Spec^R; Δ*hapR*::Cm^R; Δ*cbp*::Tm^R	This study
SAD 2839	Fig. 3 ChIP strain and Fig. S5 1x FLAG-HapR	*V. cholerae* E7946 Δ*lacZ*::P$_{chb}$-GFP, Kan^R; 1x FLAG-*hapR*; Δ*cbp*::Tm^R	This study
SAD 2840	Fig. 4 Δ*luxO*	*V. cholerae* E7946 Δ*lacZ*::P$_{chb}$-GFP, Kan^R; Δ*luxO*::Spec^R; Δ*cbp*::Carb^R	This study
SAD 2841	Fig. 4 *luxO*^{D47E}	*V. cholerae* E7946 Δ*lacZ*::P$_{chb}$-GFP, Kan^R; *luxO*^{D47E}; Δ*cbp*::Spec^R	This study
SAD 2842	Fig. 4 Δ*luxS*	*V. cholerae* E7946 Δ*lacZ*::P$_{chb}$-GFP, Kan^R; Δ*luxS*::Cm^R; Δ*cbp*::Spec^R	This study
SAD 2843	Fig. 4 Δ*cqsA*	*V. cholerae* E7946 Δ*lacZ*::P$_{chb}$-GFP, Kan^R; Δ*cqsA*::Tm^R; Δ*cbp*::Spec^R	This study
SAD 2844	Fig. 4 Δ*cqsA* Δ*hapR*	*V. cholerae* E7946 Δ*lacZ*::P$_{chb}$-GFP, Kan^R; Δ*cqsA*::Tm^R; Δ*hapR*::Cm^R; Δ*cbp*::Spec^R	This study

SAD 306	*V. cholerae* A1552 WT; parent for all other A1552 strains	*V. cholerae* A1552 O1 El Tor	(51)
SAD 2908	*E. coli* strain used to mate pMMB tfoX-qstR into complementation strains used in Fig. S1	*E. coli* S17 harboring pMMB67EH *tfoX qstR*, Cm^R	This study
SAD 2909	Fig. S1 Δ*hapR* P$_{hapR}$-*hapR*	*V. cholerae* E7946 Δ*lacZ*::P$_{chb}$-GFP Kan^R; Δ*hapR*::Spec^R; Δ*cbp*::Carb^R; ΔVCA0692::P$_{hapR}$-*hapR*, Tm^R	This study
SAD 2910	Fig. S1 Δ*hapR* Δ*clpA* P$_{hapR}$-*hapR*	*V. cholerae* E7946 Δ*lacZ*::P$_{chb}$-GFP Kan^R; Δ*hapR*::Spec^R; Δ*clpA*::Carb^R; Δ*cbp*::Tm^R; ΔVCA0692::P$_{hapR}$-*hapR*, Zeo^R	This study
SAD 2911	Fig. S1 Δ*clpA* P$_{clpSA}$-*clpSA*	*V. cholerae* E7946 Δ*lacZ*::P$_{chb}$-GFP Kan^R; Δ*clpA*::Carb^R; Δ*cbp*::Spec^R; ΔVCA0692::P$_{clpSA}$-*clpSA*, Tm^R	This study
SAD 2912	Fig. S2 Δ*hapR*	*V. cholerae* E7946 Δ*hapR*::Cm^R	This study
SAD 2845	Fig. S3 A1552 Parent	*V. cholerae* A1552 ΔVCA0692::P$_{const2}$-GFP, Spec^R; Δ*lacZ*::P$_{chb}$-mCherry, Kan^R; Δ*cbp*::Tm^R	This study
SAD 2846	Fig. S3 A1552 Δ*hapR*	*V. cholerae* A1552 ΔVCA0692::P$_{const2}$-GFP, Spec^R; Δ*lacZ*::P$_{chb}$-mCherry, Kan^R; Δ*hapR*::Cm^R; Δ*cbp*::Tm^R	This study
SAD 2847	Fig. S4 WT *lacZ*$^+$	*V. cholerae* E7946 ΔVC1807::Cm^R; ΔVC0995	This study
SAD 2848	Fig. S4 WT Δ*lacZ*	*V. cholerae* E7946 Δ*lacZ*::Spec^R; ΔVC1807::Cm^R; ΔVC0995	This study
SAD 2849	Fig. S4 Δ*hapR* Δ*lacZ*	*V. cholerae* E7946 Δ*lacZ*::Spec^R; ΔVC1807::Cm^R; ΔVC0995; Δ*hapR*::Tm^R	This study
JDN 71	Fig. 3B, 3C purified HapR	*E. coli* BL21(DE3) harboring pET28b-*6XHis-hapR*, Kan^R	This study

975
976
977
978

979 **Table 2.** *Primers used in this study.*

Primers for Inverse PCR		
Primer	**Sequence**	**Description**
BBC 244	CCCGGGATCCTGTGTGAAATTGTTATCCGC	Tn10-specific primer for Inverse PCR
BBC 434	GTGTGGGCACTCGACATATGACAAG	Tn10-specific primer for Inverse PCR
Primers for Reporter Constructs		
Primer	**Sequence**	**Description**
ABD 332	GGCTGAACGTGGTTGTCGAAAATGAC	$\Delta lacZ$ F1 (Up Arm)
BBC 219	GTTTATTTTTGTCGACTGTACAGCGTTTAAATAGAGGTCGATATTGACCC	$\Delta lacZ$ R1 (Up Arm)
ABD 342	ATTTTTCAGTTGGCCTACAATGCTTTCC	ΔVC1807 F1 (Up Arm)
BBC 719	CACCATACCCACGCCGAAACAAGGATTTTGAATTAAACGTTTCATTAGTC	ΔVC1807 R1 (Up Arm)
BBC 218	CGCTGTACAGTCGACAAAAATAAAC	Kan^R F (Middle Arm)
BBC 262	TACCGAGGACGCGAAGCTG	Kan^R R (Middle Arm)
BBC 266	CAGCTTCGCGTCCTCGGTAGAATAAAGCAATCCGCAAGCG	P$_{chb}$ F (Middle Arm)
BBC 267	CCCGGGATCCTGTGTGAAATTGAGTTGCTTTCATTTCACTAATGG	P$_{chb}$ R (Middle Arm)
BBC 732	ctcaagccgaggagtaaagaagAGTTGCTTTCATTTCACTAATGG	P$_{chb}$ fuse to *lacZ* R (Middle Arm)
BBC 252	CAATTTCACACAGGATCCCGGGAGGAGGTAACGTAATGCGTAAAGGAGAAGAAC	GFP F (Middle Arm)
BBC 254	tgtaggctggagctgcttcTTAGTTGTATAGTTCATCCATGCC	GFP R (Middle Arm)
BBC 1356	TTGTTTCGGCGTGGGTATGGTGCGCTGTACAGTCGACAAAAATAAAC	mCherry F (Middle Arm)
BBC 206	tgtaggctggagctgcttcttacttgtacagctcgtccatg	mCherry R (Middle Arm)
ABD 785	CTTCTTTACTCCTCGGCTTGAG	P$_{chb}$ stitch to *lacZ* F (Down Arm)
ABD 255	gaagcagctccagcctacaCCACAATAAGCCAGAGAGCCTTAAG	$\Delta lacZ$ F2 (Down Arm)
ABD 256	CCCAAATACGGCAACTTGGCG	$\Delta lacZ$ R2 (Down Arm)
ABD 341	gaagcagctccagcctacaTAGTCGAAAATAAAAAAAAGAGGCTCGCCTC	ΔVC1807 F2 (Down Arm)
ABD 345	CTTGCTAACCGTTGGTGTTACCAGC	ΔVC1807 R2 (Down Arm)
BBC 3083	tggataactttacgggcatgcataaggctcgtataatatattcagggagaccacaacggtttccctctacaaataattttgtttaactttCAATTTCACACAGGATCCCGGG	P$_{const2}$ F
BBC 3082	ttatacgagccttatgcatgcccgtaaagttatccagcaaccactcatagacctagggcagcagatagggacgacgtggtgttagctgtgCTCATTAGGCACCCCAGGC	P$_{const2}$ R
Primers for SOE deletion, FLAG fusions, and point mutants		
Primer	**Sequence**	**Description**
BBC 742	attccgggggatccgtcgacCTGCAGTTCagaagcagctccagcctaca	MiniFRT F
BBC 743	tgtaggctggagctgcttctGAACTGCAGgtcgacggatccccggaat	MiniFRT R

ABD 123	ATTCCGGGGATCCGTCGAC	Kan^R, $Spec^R$, $Carb^R$, Cm^R or Tm^R cassette F
ABD 124	TGTAGGCTGGAGCTGCTTC	Kan^R, $Spec^R$, $Carb^R$, Cm^R or Tm^R cassette R
ABD 796	TTAGAATCTGCGCCAGAAGCG	Δ*cbp* F1 (Up Arm)
ABD 797	gtcgacggatccccggaatCATAGCTGTTCCTTACTAGTTGC	Δ*cbp* R1 (Up Arm)
BBC 920	gtcgacggatccccggaatGCTCATCAGGTCGTCAGCC	Δ3' *cbp* R1 (Up Arm)
ABD 798	gaagcagctccagcctacaGTACTGGATCTGAAACCAGTTAAG	Δ*cbp* F2 (Down Arm)
ABD 799	GTATTGCGGAATGACCAGCATG	Δ*cbp* R2 (Down Arm)
BBC 2755	TTACCCCTAAGTCGGCGAGC	ΔVC0995 F1 (Up Arm)
BBC 2756	gtcgacggatccccggaatAATATTCACCTTAAGTTCCCCC	ΔVC0995 R1 (Up Arm)
BBC 2757	gaagcagctccagcctacaCTGCCTTAATCGAGTTTAAACCC	ΔVC0995 F2 (Down Arm)
BBC 2758	GCACCACGATAGCAATAAGC	ΔVC0995 R2 (Down Arm)
CAK 407	GCTACGCACTGCCAAATTACC	Δ*clpA* or Δ*clpS* F1 (Up Arm)
CAK 408	gtcgacggatccccggaatAAGCATAAGGCCTCCTTAAGGAAC	Δ*clpA* R1 (Up Arm)
CAK 409	gaagcagctccagcctacaCACTAGACATCTCACAATACGC	Δ*clpA* F2 (Down Arm)
CAK 410	CCGCTAACATCTCAGGACTG	Δ*clpA* or Δ*clpS* R2 (Down Arm)
CKP 449	TGCTCGGTTTTGATCCGTTC	Δ*clpP* or Δ*clpX* F1 (Up Arm)
CKP 450	gtcgacggatccccggaatGGGCGACATTGCATTTTTTTC	Δ*clpP* R1 (Up Arm)
CKP 451	gaagcagctccagcctacaGGCGAGTAAGCTCGTAATTG	Δ*clpP* F2 (Down Arm)
CKP 454	GTCAAACTAGAAACCAGCGC	Δ*clpP* or Δ*clpX* R2 (Down Arm)
CKP 447	gtcgacggatccccggaatTTTACTCATGACACTTCAATATTTG	Δ*clpS* R1 (Up Arm)
CKP 448	gaagcagctccagcctacaCAAGCTTAAACTTGCCGGGC	Δ*clpS* F2 (Down Arm)
CKP 452	gtcgacggatccccggaatGTCTGTCATTCGCTAACCTC	Δ*clpX* R1 (Up Arm)
CKP 453	gaagcagctccagcctacaGCGGAGTAATCTCAAGCAAC	Δ*clpX* F2 (Down Arm)
CKP 489	TTGACGCTCTGAAAGCAGAAG	Δ*lonA* F1 (Up Arm)
CKP 490	gtcgacggatccccggaatCAAGTTCATATTTTTCTCTCTTCCG	Δ*lonA* R1 (Up Arm)
CKP 491	gaagcagctccagcctacaGATGCATAGCAAAAATAAGTAAATC	Δ*lonA* F2 (Down Arm)
CKP 492	CGAAGAATTATAAGTGCAAAGGC	Δ*lonA* R2 (Down Arm)
BBC 422	ACGTTCAATAACCAGAATGTTGG	Δ*hapR* F1 (Up Arm)
BBC 423	gtcgacggatccccggaatTTTTCGATTGATGCGTCCATAG	Δ*hapR* R1 (Up Arm)
CKP 511	GTCtttgtcatcgtcatccttataatcCATAGGGGTATATCCTTGCC	*1x FLAG-hapR* R1 (Up Arm)

BBC 408	gaagcagctccagcctacaAACTAGTTTCTTGGGCAGCACAA	Δ*hapR* F2 (Down Arm)
CKP 510	ATGgattataaggatgacgatgacaaaGACGCATCAATCGAAAAACG	*1x FLAG-hapR* F2 (Down Arm)
BBC 409	GTTCATAATGATTTCCTTGGTGCC	Δ*hapR* R2 (Down Arm)
BBC 374	TGGCAAAAAGCGAGAGAAGAAG	Δ*luxO* F1 (Up Arm)
BBC 375	gtcgacggatccccggaatCATGAGGACATATTTTGTTTTCTGC	Δ*luxO* R1 (Up Arm)
CKP 519	TCATATCTGGCAAACGTAAcTCCAGCAGGATTAAGTCAGG	*luxO*^{D47E} R1 (Up Arm)
BBC 376	gaagcagctccagcctacaTAAGCGATGAGAGAATGGATCAAC	Δ*luxO* F2 (Down Arm)
CKP 518	TGACTTAATCCTGCTGGAgTTACGTTTGCCAGATATGACG	*luxO*^{D47E} F2 (Down Arm)
BBC 377	TCACACCCGAATTTCCATCATGC	Δ*luxO* R2 (Down Arm)
CKP 554	GTCTCTTAGCCGAGGTACTG	Δ*cqsA* F1 (Up Arm)
CKP 555	gtcgacggatccccggaatCTTGTTCATCGCAATATATCCTAG	Δ*cqsA* R1 (Up Arm)
CKP 556	gaagcagctccagcctacaTTTCGTTAAATGCATAAATAACAAAAAC	Δ*cqsA* F2 (Down Arm)
CKP 557	AGTTGGAACCACTTCTTGTC	Δ*cqsA* R2 (Down Arm)
CKP 549	TACAACTGCTTGGCACGC	Δ*luxS* F1 (Up Arm)
CKP 550	gtcgacggatccccggaatTGGCATTTCCTTTCTCCC	Δ*luxS* R1 (Up Arm)
CKP 551	gaagcagctccagcctacaCACTAAGTCGGTTCTGTAAACG	Δ*luxS* F2 (Down Arm)
CKP 552	TTCTTAGCGTGATCAATTGC	Δ*luxS* R2 (Down Arm)
ABD 334	AGTGCTCCGGACTCTTTGCTCTG	Δ*lacZ* F1 (Up Arm)
ABD 254	gtcgacggatccccggaatCATCCCTCAAGCCGAGGAGTAAAG	Δ*lacZ* R1 (Up Arm)
ABD 255	gaagcagctccagcctacaCCACAATAAGCCAGAGAGCCTTAAG	Δ*lacZ* F2 (Down Arm)
ABD 256	CCCAAATACGGCAACTTGGCG	Δ*lacZ* R2 (Down Arm)
ABD 725	GAAGCAGCTCCAGCCTACA	Detect F for all deletions
BBC 082	gtcgacggatccccggaatCATAACTTACACCTTACTCACCCAG	Δ*cbp* detect R
CAK 411	TTGTTTGGTGCGATTATTGG	Δ*clpA* detect R
CKP 464	TTGGCAGAACATCTTTGATC	Δ*clpP* detect R
CKP 463	AAAGTATCCAGCTCACGGCG	Δ*clpS* detect R
CKP 465	CCATGTGTGGATAGACAACC	Δ*clpX* detect R
CKP 493	CAGTGCCACTTGGTCACCTG	Δ*lonA* detect R
BBC 410	TAAATGGGGCTTGGAGAATTTAG	Δ*hapR* detect R
BBC 1911	CGTAATCAAACTGCGAAAGTG	Δ*luxO* detect R
CKP 558	AATTCGTAACTCTGAGCATG	Δ*cqsA* detect R
CKP 553	TGGACCACGAACCTTAAACG	Δ*luxS* detect R
BBC 2759	TTGTCAGAAAGCGTTTCTGC	ΔVC0995 detect R
ABD 399	AACTGATGGCAGAAAAAGCCACTCAG	Δ*lacZ* detect R
BBC 993	ttgattataaggatgacgatgac	*1x FLAG* detect F
ABD 846	CATAAACATGTTTCTGATCAGCAG	*1x FLAG-hapR* detect R
BBC 380	GCCAATAGAATGAGTCTATTGGCTG	*luxO*^{D47E} detect F
CKP 520	TCATATCTGGCAAACGTAgc	*luxO*^{D47E} detect R
Primers for EMSA probes		

Primer	Sequence	Description
BBC 744	cagcttcgcgtcctcggtaCGCAAATATAACTCAGGCAAAG	P*chb* F
CKP 072	cccgggatcctgtgtgaaattgCTTTGGCAGGAGTAAGAAAACACCTAG	P*chb* R
CKP 865	TAAGCAAACTGTAGCGTAGAAG	P*clpP* F
CKP 866	TGCATTTTTTTCTTGGTAGC	P*clpP* R
CKP 504	GCCAAATAAGTAAGTAAACACTAACAAACCTA	HapR BS 1 F
CKP 505	TAGGTTTGTTAGTGTTTACTTACTTATTTGGC	HapR BS 1 R
CKP 506	AATTGCAATTGATAAATTTCTTGGCTAGGTGT	HapR BS 2 F
CKP 507	ACACCTAGCCAAGAAATTTATCAATTGCAATT	HapR BS 2 R
Primers for qPCR and qRT-PCR		
Primer	Sequence	Description
BBC 989	GCATCTAGGTTTTGACGTTTTTAACG	P*chb* amplify F
BBC 990	AACACTCTCCAAGACCTACCTC	P*chb* amplify R
ABD 132	CTGTCTCAAGCCGGTTACAA	*rpoB* amplify F
ABD 133	TTTCTACCAGTGCAGAGATGC	*rpoB* amplify R
CKP 865	TAAGCAAACTGTAGCGTAGAAG	P*clpP* amplify F
CKP 867	TCGTATGAACGCTCACCACG	P*clpP* amplify R
Primers for purification vector		
Primer	Sequence	Description
JDN 92	agcagcggcctggtgccgcgcggcagcCatatggacgcatcaatcgaaaaa	Cloning *6X-His-hapR* into pET28b vector via IDA, F
JDN 93	tcagtggtggtggtggtggtgCTCGAGctagttcttatagatacacagcatattgagg	Cloning *6X-His-hapR* into pET28b vector via IDA, R
JDN 22	CTCGAGcaccaccaccacca	Amplify pET28b vector backbone for IDA, F
JDN 23	atGgctgccgcgcggcacca	Amplify pET28b vector backbone for IDA, R
Primers for complementation constructs		
BBC 832	GCTTTTTGCTACAACGACCG	ΔVCA0692 F1 for Up arm
BBC 828	CACCATACCCACGCCGAAACAACAGTGATGTAGCGAATCGGAC	ΔVCA0692 R1 for Up arm
BBC 243	TTGTTTCGGCGTGGGTATGGTG	Tm^R or Zeo^R F for Middle arm
BBC 647	tttttctatttctgaatcgattcatacgaCTCATTAGGCACCCCAGGC	Tm^R F or Zeo^R for Middle arm
CKP 948	tcgtatgaatcgattcagaaatagaaaaaACCATTCTCGTTGTGTTGGG	P*hapR*-*hapR* F for Middle arm
BBC 729	tgtaggctggagctgcttcCTAGTTCTTATAGATACACAGCATATTG	P*hapR*-*hapR* R for Middle arm
CKP 949	tcgtatgaatcgattcagaaatagaaaaaCCCTCATGCATTTTATAACTG	P*clpSA*-*clpSA* F for Middle arm
CKP 950	tgtaggctggagctgcttcCTAGTGGACCACCTCTTCGC	P*clpSA*-*clpSA* R for Middle arm
BBC 830	gaagcagctccagcctacaGTTGAGTTGGATGCAGCACC	ΔVCA0692 F2 for Down arm
BBC 834	CACAATTTCTCGCTTAAAATGTCC	ΔVCA0692 R2 for Down arm
CKP 509	gcaggtggagcaggtggaGACGCATCAATCGAAAAACG	ΔVCA0692::P*hapR*-*hapR* detect F

BBC 1101	CAGACGTACTATTAACAGGACTGAC	ΔVCA0692::P$_{hapR}$-*hapR* detect R
CKP 235	gtcgacggatccccggaatCAAATATATCCTCCTCACTATTTTGATTAG	ΔVCA0692::P$_{clpSA}$-*clpSA* detect F
CKP 447	gtcgacggatccccggaatTTTACTCATGACACTTCAATATTTG	ΔVCA0692::P$_{clpSA}$-*clpSA* detect R

Supplemental Information for

Species-specific quorum sensing represses the chitobiose utilization locus in *Vibrio cholerae*

Catherine A. Klancher, Jane D. Newman, Alyssa S. Ball, Julia C. van Kessel, and Ankur B. Dalia*

Figure S1. *Complementation of ΔhapR and ΔclpA mutant strains.* Expression of a P*chb*-GFP transcriptional reporter was determined in the indicated strains. The parent strain contained a chromosomally-encoded P*chb*-GFP reporter and a Δ*cbp* mutation. Strains were complemented with a chromosomally-encoded single copy of the indicated gene driven by its native promoter at an ectopic site (*hapR* or *clpA* Complement). Fluorescence of cultures was determined on a plate reader from at least three independent biological replicates and is shown as the mean ± SD. Statistical comparisons were made by one-way ANOVA with Tukey's post-test. NS, not significant. *, p = 0.0104. ***, p < 0.001. The P*chb*-GFP fluorescence data for "Δ*clpA*", "Δ*hapR* Δ*clpA*", and "Δ*hapR*" are identical to the data presented in **Fig. 1** and were included here for ease of comparison.

Figure S2. *HapR is a repressor of native chb expression.* The indicated strains were induced with sterile water (no inducer) or chitin oligosaccharides (+ $(GlcNAc)_6$), then assessed for *chb* transcript abundance by qRT PCR. The parent strain used was *V. cholerae* E7946 WT. Data are shown on **A**) a log scale or **B**) a linear scale. The latter plot helps accentuate that deletion of *hapR* results in a two-fold increase in native *chb* transcripts in response to chitin oligosaccharides, which is similar to what is observed using transcriptional reporters. Data are from at least three independent biological replicates and is shown as the mean ± SD. Statistical comparisons were made by one-way ANOVA with Tukey's post-test. **, p = 0.0064.

Figure S3. *HapR is a repressor of P$_{chb}$ in V. cholerae E7946 and V. cholerae A1552.* **A**) Expression of a P$_{chb}$-mCherry reporter was determined in the indicated mutant strains. Each respective parent strain contained a P$_{chb}$-mCherry reporter and a Δ*cbp* mutation to activate ChiS. **B**) Scatter plot showing the relative expression of a P$_{chb}$-mCherry reporter in the indicated *V. cholerae* A1552 strains that have *cbp* intact when using chitin oligosaccharides to induce cells. The parent strain background contains both a P$_{chb}$-mCherry reporter, a P$_{const2}$-GFP construct (which exhibits constitutive GFP expression), and *cbp* intact. n = 2189 for Parent; n = 2164 for Δ*hapR.* Data shown are from two independent biological replicates. **C**) Representative images of *V. cholerae* A1552 strains analyzed in **B**. Scale bar = 2 μm. Statistical comparisons in **A** were made using a one way ANOVA with Tukey's post test and in **B** were made using Student's t-test. ***, p < 0.001.

Figure S4. A *ΔhapR mutant does not have a fitness advantage for growth on chitobiose.* The indicated *V. cholerae* strains were mixed and co-cultured in M9 minimal media supplemented with 0.2% chitobiose + 10 µM CAI-1 and grown for the indicated number of generations. Each data point represents an independent biological replicate.

Figure S5. *N-terminally FLAG-tagged HapR is functional for regulation of P$_{chb}$.* Strains expressing the indicated HapR allele at the native locus were assessed for expression of a P$_{chb}$-GFP reporter. Both strains also harbored a Δ*cbp* deletion to activate ChiS. Statistical comparisons were made using Student's t-test. NS, not significant.

Figure S6. Model of quorum sensing regulation in *V. cholerae*. There are four parallel sensor kinases that contribute to QS in *V. cholerae*: LuxQ, CqsS, VpsS, and CqsR. The signals for VpsS and CqsR are unknown. LuxPQ senses the interspecies autoinducer AI-2 and CqsS senses the *V. cholerae*-specific autoinducer CAI-1. AI-2 is produced by the LuxS synthase and CAI-1 is produced by the CqsA synthase. (**A**) At low cell density, the LuxQ and CqsS sensors act as kinases, resulting in phosphorylation of LuxO, which subsequently decreases expression of HapR. (**B**) At high cell density, LuxQ and CqsS act as phosphatases, which results in dephosphorylation of LuxO, and subsequent activation of HapR expression. Dashed lines indicate indirect effects of one protein on another.

Table S1. *Other hits from P_{chb} activator screen and ∆clpA counter-screen.*

Activator or ∆*clpA* counter-screen?	Gene	VC gene designation	Putative function	Number of unique insertions
Activator	*treC*	VC0911	Trehalose-6-phosphate hydrolase	5
Activator	*cyaA*	VC0122	Adenylate cyclase	1
∆*clpA* counter-screen	*nagC*	VC0993	GlcNAc transcriptional repressor protein	4
∆*clpA* counter-screen	*cytR*	VC2677	Cytidine repressor protein	1