1 **Species-specific quorum sensing represses the chitobiose utilization locus**
2 **in *Vibrio cholerae***
3

4 Catherine A. Klancher, Jane D. Newman, Alyssa S. Ball, Julia C. van Kessel, and
5 Ankur B. Dalia*
6
7 Department of Biology, Indiana University, Bloomington, IN 47405
8 *Author for correspondence: ankdalia@indiana.edu

9
10

Abstract

The marine facultative pathogen *Vibrio cholerae* forms complex multicellular
communities called biofilms on the shells of crustacean zooplankton in its aquatic
reservoir. These shells are primarily composed of the polysaccharide chitin. *V.*
cholerae-chitin interactions are critical for the growth, evolution, and waterborne
transmission of cholera. This is due, in part, to chitin inducing changes in gene
expression in this pathogen. Here, we sought to identify factors that influence
chitin-induced expression of one locus, the chitobiose utilization operon (*chb*),
which is required for the uptake and catabolism of the chitin disaccharide.
Through a series of genetic screens, we identified that the master regulator of
quorum sensing, HapR, is likely a direct repressor of the *chb* operon. We also
found that the levels of HapR in *V. cholerae* are regulated by the ClpAP
protease. Furthermore, we show that the canonical quorum sensing cascade in
V. cholerae regulates *chb* expression in a HapR-dependent manner. Through
this analysis we found that signaling via the species-specific autoinducer CAI-1,
but not the inter-species autoinducer AI-2, influences *chb* expression. This
phenomenon of species-specific regulation may enhance the fitness of this
pathogen in its environmental niche.

Importance
In nature, bacteria live in multicellular and multispecies communities. Microbial
species can sense the density and composition of their community through
chemical cues using a process called quorum sensing (QS). The marine
pathogen *Vibrio cholerae* forms communities on the chitinous shells of
crustaceans in its aquatic reservoir, which is critical for the survival, evolution,
and waterborne transmission of this pathogen. Here, we show that *V.*
cholerae uses QS to regulate the expression of one locus required for *V.*
cholerae-chitin interactions.
39

Introduction
The facultative bacterial pathogen *Vibrio cholerae* is the causative agent of the
diarrheal disease cholera and natively resides in the aquatic environment. In this
niche, *V. cholerae* forms multicellular communities on biotic and abiotic chitinous
surfaces, like the shells of crustaceans or marine snow (1). Chitin is a
polysaccharide made up of β-1,4 linked *N*-acetylglucosamine (GlcNAc) and
serves as a major nutrient source for *V. cholerae* in the marine environment. The

ability of *V. cholerae* to form chitin biofilms is critical for the waterborne
transmission of *V.* cholerae (2, 3). As chitin is the most abundant biopolymer in
the ocean, the ability of *Vibrio* species to break down and utilize this highly
insoluble polysaccharide also serves an important role in global nitrogen and
carbon recycling (1). Chitin induces genetic programs in many *Vibrio* species that
may increase their fitness in the environment, such as the chitin utilization
program and natural competence (4, 5).
54
The chitin utilization program in *V. cholerae* is robust and highly regulated. To
study this program, most studies employ the chitobiose utilization operon (*chb*),
which is required for the uptake and catabolism of the chitin disaccharide and is
highly induced in the presence of chitin oligosaccharides (4). Presently, there are
several regulators of this program that have been identified. ChiS is the master
regulator of the chitin utilization program in *V. cholerae*, and we have recently
shown this protein is a direct transcriptional activator required for induction of the
chb locus (6, 7). Previous work has shown that carbon catabolite repression
(CCR) can also play a role in regulating chitin responsive phenotypes including
ChiS-dependent induction of *chb* and natural competence (8, 9). In addition, our
group has previously found that the cell division licensing factor SlmA plays an
essential role in activating *chb* expression (10). Tight regulation via these diverse
signaling systems may act to ensure that the chitin utilization program is only
expressed under conditions in which it will provide a competitive advantage.
69
70　Here, through a number of genetic screens and complementary molecular
71　methods we show that quorum sensing is an additional regulatory system that
72　tunes expression of a chitin utilization locus in *V. cholerae*.
73
Results
ClpA is identified in an unbiased screen for activators of P_{chb}
To identify additional genes required for activation of the *chb* locus, we
conducted a transposon mutant screen. This was carried out in a strain
containing a P_{chb}-*lacZ* transcriptional reporter. As shown previously, induction of
P_{chb} is dependent on the activity of the master regulator ChiS (4, 7). In the
absence of chitin, ChiS activity is repressed by CBP. In the presence of chitin,
CBP repression of ChiS is relieved, which allows for ChiS-dependent activation
of P_{chb}. ChiS can also be activated genetically in the absence of chitin by deleting
cbp (7, 10). As chitin oligomers are prohibitively expensive, a Δcbp mutation was
used to induce ChiS-dependent P_{chb}-*lacZ* expression in our genetic screen
exactly as previously described (10). So, the starting genotype for our screen
was a strain containing P_{chb}-*lacZ* and a Δcbp mutation. This strain formed blue
colonies on X-gal containing plates and we screened for white colonies to identify
putative activators that act downstream of ChiS to contribute to P_{chb} induction.
89
One gene identified in this genetic screen was *clpA*. To study the effect of ClpA
on P_{chb} activity moving forward, we utilized a previously described P_{chb}-GFP
reporter (7, 10). Using this reporter, we found that a $\Delta clpA$ mutation resulted in a

~3-fold decrease in P$_{chb}$ expression relative to the parent (**Fig. 1**). ClpA is a
AAA+ unfoldase that recognizes protein substrates, unfolds them, and feeds
them into the ClpP protease where they are degraded (11). If ClpA was exhibiting
its effect on P$_{chb}$ expression as a part of the ClpAP machine, we hypothesized
that a ΔclpP mutation should phenocopy a ΔclpA mutation. Indeed, ΔclpP and
ΔclpP ΔclpA strains phenocopied a ΔclpA mutant for P$_{chb}$ expression (**Fig. 1**).
These results suggest that loss of the ClpAP protease decreases P$_{chb}$
expression. As ClpAP normally degrades proteins, we hypothesized that ClpAP
may indirectly promote activation of P$_{chb}$ by degrading a repressor of the *chb*
locus.
103
HapR is a repressor of P$_{chb}$ that is degraded by ClpAP
To identify a putative repressor of P$_{chb}$ that is targeted by ClpAP for degradation,
we conducted a counter-screen using the P$_{chb}$-*lacZ* reporter. For the counter-
screen, the parent strain had both Δcbp and ΔclpA mutations. This mutant is
white on X-gal plates because P$_{chb}$-*lacZ* is poorly expressed; inactivation of the
putative repressor in this strain should result in restoration of P$_{chb}$-*lacZ*
expression and yield a blue colony phenotype. In this counter-screen, we
identified *hapR*. A ΔhapR mutation restored P$_{chb}$ expression in the ΔclpA mutant
background (**Fig. 1**). In fact, a ΔhapR mutation allowed for higher P$_{chb}$ expression
than the parent strain. This suggests that HapR also represses P$_{chb}$ expression
when ClpAP is intact; and that ClpAP does not degrade the entire pool of HapR
in the cell (**Fig. 1**). Importantly, the level of P$_{chb}$ expression observed in the
ΔhapR ΔclpA mutant phenocopied the ΔhapR strain (**Fig. 1**). This epistasis
between *clpA* and *hapR* suggests that they are involved in the same pathway for
regulating P$_{chb}$ expression.
119
We hypothesized that the reason P$_{chb}$ expression is decreased in *clpAP* mutants
was due to increased HapR protein levels. Western blotting in these
backgrounds revealed that HapR levels were, indeed, increased in strains
containing mutations to *clpA* and/or *clpP* (**Fig. 1**). ClpAP dependent degradation
of HapR is not unique to *V. cholerae*, but has previously been observed in *Vibrio*
vulnificus where ClpAP degrades the HapR homolog SmcR (12). In addition to
the ClpAP machine, it was shown that the Lon protease also plays a role in
SmcR degradation. So, we next sought to investigate the role of other protease
machines on induction of P$_{chb}$ and HapR protein levels. Mutations to other Clp
components (the ClpS adaptor protein or the ClpX unfoldase*)* did not have a
marked impact on HapR protein levels. Also, a ΔclpS mutation did not affect P$_{chb}$
expression levels, while a ΔclpX mutation slightly decreased P$_{chb}$ expression
(**Fig. 1**). Because HapR levels were not elevated in the ΔclpX mutant
background, however, the observed decrease in P$_{chb}$ expression may be
attributed to a pleiotropic effect. In contrast to the effect of the Lon protease on
SmcR levels in *V. vulnificus*, we did not observe an impact of ΔlonA on HapR
levels in *V. cholerae*; and ΔlonA, correspondingly, did not affect P$_{chb}$ expression
(**Fig. 1**). Together, these results establish that HapR is a repressor of P$_{chb}$ and
that HapR levels are controlled specifically by the ClpAP protease in *V. cholerae*.

HapR-mediated repression of P$_{chb}$ occurs on chitinous surfaces

Thus far, we have demonstrated that HapR is a repressor of P$_{chb}$ that acts downstream of ChiS activation using a Δ*cbp* mutation to induce ChiS activity. Natural induction of this locus, however, occurs in the presence of chitin oligosaccharides. So, we next wanted to test whether the repression of P$_{chb}$ by HapR was observed in a more physiologically relevant setting. To test this, we cultured *V. cholerae* strains with CBP intact, a P$_{chb}$-mCherry reporter, and a construct that constitutively expresses GFP (P$_{const2}$-GFP) on chitin beads (**Fig. 2A**). The P$_{const2}$-GFP construct [derived from the insulated proD promoter (13)] served as an internal control for noise in gene expression and was used to normalize P$_{chb}$-mCherry expression in single cells (see Methods for details).

When cells were cultured on chitin beads, both the parent strain and the Δ*hapR* strain exhibited a bimodal distribution of P$_{chb}$ expression (**Fig. 2A-D**). There were at least two possible explanations for the observed bimodality in P$_{chb}$ gene expression in this experiment. One, the signaling pathway that leads to activation of *chb* has switch-like behavior that results in a population that exhibits bimodality in *chb* expression. Or two, the environment during growth on chitin beads is heterogenous and only some cells within the population have access to the chitin oligosaccharides necessary for activation of *chb*. Those cells that do not have access to chitin do not activate P$_{chb}$ expression (P$_{chb}$ Off), while those that do have access to chitin do induce P$_{chb}$ expression (P$_{chb}$ On). To differentiate between these two possibilities, we genetically activated P$_{chb}$ in the absence of chitin by deleting *cbp*. If the signaling circuit responsible for P$_{chb}$ activation has switch like behavior, we would predict that bimodality in gene expression would be maintained; however, if bimodality is the result of heterogenous access to chitin oligosaccharides when cells are cultured on chitin beads, we would expect bimodality to be lost. When P$_{chb}$ is activated via a Δ*cbp* mutation, all cells in the population are uniformly "P$_{chb}$ On" (**Fig. 2E**) and the population becomes unimodal (**Fig. 2F**). Thus, these results showed that only the "P$_{chb}$ On" cells have access to chitin oligosaccharides when grown on chitin beads. These results are consistent with chitin-induction of natural competence genes on chitin beads, where cells display bimodality in gene expression that is also likely due to heterogeneous access to inducing chitin oligosaccharides (14).

Next, we tested whether HapR influenced activation of P$_{chb}$ on chitin beads by just assessing the expression level among "P$_{chb}$ On" cells. As observed in bulk populations, the Δ*hapR* strain exhibited a ~1.7 fold increase in P$_{chb}$ expression (**Fig. 2D**). This was also consistent with the ~2-fold increase in P$_{chb}$ expression observed in single cells when the population was induced via deletion of *cbp* (**Fig. 2G**). Together, these results indicate that HapR is a *bona fide* repressor of P$_{chb}$ even under physiologically relevant inducing conditions.

HapR repression of P$_{chb}$ is conserved in other V. cholerae El Tor isolates

Previously, it was suggested that HapR was actually an activator of P*chb*
expression in the *V. cholerae* El Tor isolate A1552 (15). Specifically, the authors
showed that deletion of *hapR* resulted in a decrease in P*chb* expression when
cells were cultured on chitin flakes. In the present study, we use the *V. cholerae*
El Tor isolate E7946 to study P*chb* regulation. To assess if HapR exhibits different
effects on P*chb* expression depending on the strain background, we assessed
expression of a P*chb*-mCherry reporter in *hapR*⁺ and Δ*hapR* derivatives of both
E7946 and A1552. Consistent with our previous results, we observed that P*chb*
expression is elevated ~2-fold in the Δ*hapR* derivative of both strain
backgrounds, which is consistent with HapR acting as a repressor of this locus
(**Fig. S1**). It is unclear what explains the discrepancy between our findings and
those that were previously reported; however, these data suggest that they
cannot be attributed to differences between strain backgrounds.
197
Deletion of HapR does not confer a growth advantage during growth on
chitobiose
Our data indicate that in the absence of HapR, P*chb* expression is elevated. The
chb locus encodes the genes required for uptake and degradation of the chitin
disaccharide, chitobiose. So next, we wanted to assess if the increased
expression of P*chb* confers a fitness advantage to Δ*hapR* cells during growth on
chitobiose. To test this, we conducted competitive growth assays with a 1:1
mixture of a parent and a Δ*hapR* mutant strain on minimal medium with
chitobiose as the sole carbon source. We hypothesized that if the Δ*hapR* mutant
had a competitive growth advantage due to increased expression of *chb*, then it
should outcompete the parent strain in this assay. Even after ~48 generations of
growth on chitobiose, however, we did not observe a competitive advantage for
the Δ*hapR* mutant (**Fig. S2**). HapR is a global regulator that controls the
expression of dozens of genes. Thus, even though a Δ*hapR* mutant has
increased P*chb* expression, there may be negative pleiotrophic effects associated
with the Δ*hapR* mutation that masks any competitive advantage of derepression
of P*chb* in this mutant during growth on chitobiose.
215
*HapR binds P*chb* in vitro and in vivo*
Thus far, our data suggest that HapR is a repressor of P*chb*, but it does not
distinguish whether HapR is a direct or indirect regulator of this locus. To assess
if HapR could be regulating P*chb* directly, we used an *in silico* approach to identify
putative HapR binding sites in P*chb* based on consensus binding sequences
generated for the HapR homologs LuxR (from *Vibrio harveyi*) and SmcR (from *V.*
vulnificus) via chromatin immunoprecipitation sequencing (ChIP-seq) (16, 17).
Through this analysis, we identified two potential HapR binding sites in P*chb* (**Fig.**
3A; exact sequences in **Fig. S3**). Interestingly, these binding sites (BSs) overlap
with other elements in the *chb* promoter required for transcriptional activation.
HapR BS 1 overlaps with the SlmA binding site, which is a critical activator of
P*chb* expression (10). HapR BS 2 overlaps with the -35 signal, which is required
for RNA polymerase to bind to the promoter and initiate transcription. This

sequence analysis suggests that the repressive effect of HapR may be due to
HapR binding directly antagonizing SlmA and/or RNAP at this locus.
231
We next sought to determine whether HapR could directly bind to P_{chb} using both
in vitro and *in vivo* approaches. First, we tested whether HapR could bind to P_{chb}
in vitro using EMSA assays. Consistent with the presence of two HapR binding
sites in P_{chb}, we observed two shifts by EMSA when using a DNA probe of the
chb promoter (**Fig. 3B**). HapR does not regulate the ClpP promoter and thus did
not bind P_{clpP} (**Fig. 3B**), which is consistent with previous studies (18). EMSAs
were also done using 32 bp probes that encompassed each putative HapR BS
(probe sequences in **Fig. S3**). We observed that HapR was able to bind to both
probes, suggesting that HapR binds to both HapR BS 1 and HapR BS 2 (**Fig.
3C**). Next, we wanted to assess if HapR bound to P_{chb} *in vivo* via ChIP assays
under physiologically relevant conditions. To that end, we first generated a
FLAG-HapR strain that was functional for regulating P_{chb} expression (**Fig. S4**).
Using this strain in ChIP-qPCR assays, we found that P_{chb} was, indeed, bound by
HapR *in vivo*, while the negative control P_{clpP} locus was not bound by HapR (**Fig.
3D**). Together, these data demonstrate that HapR binds to P_{chb}, which suggests
that it may be a direct repressor of this locus.
248
Quorum sensing regulates expression of P_{chb} through the cholera-specific
autoinducer CAI-1
HapR is the master regulator of quorum sensing (QS) in *V. cholerae*, and is
highly expressed at high cell density (HCD). Thus far, we have only established
that HapR is a repressor of P_{chb} expression. Next, we wanted to probe the role of
QS in regulating P_{chb}. To that end, we sought to test the effect of mutations to
genes upstream of HapR in the *V. cholerae* QS cascade on P_{chb} induction and
HapR protein levels.
257
QS in *Vibrio* species is controlled by autoinducer-responsive sensor proteins that
indirectly modulate phosphorylation of the response regulator LuxO, which in turn
indirectly regulates production of HapR. When autoinducer concentrations are
low (*i.e.*, at low cell density (LCD)), multiple histidine kinase sensors acts as
kinases. This results in high levels of phosphorylated LuxO, which prevents
HapR production (**Fig. 4A**, left). By contrast, at high autoinducer concentrations
(*i.e.*, at HCD), the sensors act as phosphatases, which ultimately leads to
dephosphorylation of LuxO and allows for HapR production (**Fig. 4A**, right).
266
First, we assessed the impact of HCD on P_{chb} induction and HapR levels by
deleting *luxO*, which genetically locks cells in a HCD state. Induction of the
parent strain is tested under HCD conditions, thus, as expected, HapR levels
were similar between the parent and the Δ*luxO* mutant; accordingly, expression
of P_{chb} was also similar between the parent and Δ*luxO* (**Fig. 4B**). Next, we tested
the effect of LCD on P_{chb} and HapR expression by generating a *luxO*^{D47E} mutant,
which mimics phosphorylated LuxO and genetically locks cells in a LCD state. In
this strain, we saw that P_{chb} expression increased and was correlated with a

decrease in HapR protein levels (**Fig. 4B**). Together, these data suggest that
HapR repression of P_{chb} is mediated through the canonical QS circuit.
277
Next, we wanted to move further upstream in the QS regulatory cascade to
address if distinct autoinducers differentially affected expression of P_{chb} and
HapR. There are potentially four histidine kinase sensors that coordinate QS-
dependent control of HapR expression (19); however, the autoinducer signal is
only known for two of these systems. The sensor LuxPQ is responsive to the
inter-species autoinducer AI-2, and the sensor CqsS is responsive to the *V.*
cholerae-specific autoinducer CAI-1 (**Fig. 4A**) (20). To assess the role of each
autoinducer in regulation of P_{chb}, we made mutations to the synthase genes
responsible for production of each autoinducer. LuxS makes AI-2 (21) and CqsA
makes CAI-1 (20) (**Fig. 4A**). In a strain that no longer produces AI-2 (Δ*luxS*),
expression of P_{chb} was similar to that observed in the parent and HapR levels
remained similar in these two strains (**Fig. 4B**). By contrast, a strain that is
unable to produce CAI-1 (Δ*cqsA*) had increased expression of P_{chb}, likely due to
the low level of HapR produced (**Fig. 4B**). The observed decrease in P_{chb}
expression in the Δ*cqsA* background was, indeed, due to a lack of CAI-1
production as exogenously adding back synthetic CAI-1 restored HapR levels
and repression of P_{chb} to the parent level (**Fig. 4B**). In addition, a strain that does
not make CAI-1 induced P_{chb} to the same level as a strain that does not produce
CAI-1 or HapR (**Fig. 4B**). This epistasis indicates that CAI-1 production and
HapR are involved in the same regulatory pathway for modulating expression of
P_{chb}. These data support previous results, which indicate that CqsS signaling
plays a dominant role in regulating HapR (14, 20, 22, 23). Together, these results
indicate that P_{chb} expression is strongly influenced by QS signaling mediated by
the *V. cholerae*-specific autoinducer CAI-1 and less via the inter-species
autoinducer AI-2.
303
Discussion
Here, we show that HapR acts as a repressor of chitobiose utilization genes in *V.*
cholerae. When this organism forms communities on a chitinous surface, chitin
induces expression of P_{chb} through activation of the chitin sensor ChiS (**Fig. 5**).
V. cholerae then modulates P_{chb} expression depending on the presence of other
V. cholerae cells, which it senses via the *V. cholerae*-specific autoinducer CAI-1.
At low CAI-1 concentrations, P_{chb} expression is high because the repressor HapR
is produced at a low level (**Fig. 5A**). At high CAI-1 concentrations, HapR is
produced at higher levels and can repress P_{chb} expression (**Fig. 5B**). Because
HapR directly binds to P_{chb} both *in vitro* and *in vivo*, we speculate that this
repressive effect is direct.
315
When HapR is produced, it gets proteolyzed by the ClpAP machine (**Fig. 5B**).
While the dynamic range of HapR-dependent P_{chb} repression is only ~2-fold
when ClpAP is intact, the dynamic range of repression increases to ~7-fold in the
absence of ClpAP. Thus, it is tempting to speculate that regulation of ClpAP may
modulate P_{chb} expression under some conditions. Currently, however, little is

known about the regulation of ClpAP in *V. cholerae*. ClpAP is induced during
heat shock in *V. vulnificus* (12). Also, in *E. coli*, ClpP expression is induced
during heat shock (24-26). Thus, it is believed that the ClpP protease plays a role
in the heat shock response. In *Bacillus subtilis*, ClpP is upregulated under
various stress conditions, suggesting that this protease may also be induced by a
general stress response (27). Thus, one possibility is that stress-dependent
regulation of ClpAP indirectly regulates P_{chb}. It has also been hypothesized that
V. cholerae makes use of proteolysis machines to rapidly respond to changes in
their environment; namely, the transition from the human gut to the aquatic
environment after infection (28). ClpAP acting as an activator of P_{chb} may be a
way for cells to rapidly induce expression of a locus that is not important in the
human gut, but is useful for survival in its aquatic reservoir. Consistent with this
idea, *chb* is induced late in infection, suggesting this is a critical locus for
preparing to re-enter the aquatic environment after infection of a host (29).
335
The mechanisms underlying distinct responses to AI-2 or CAI-1 remain a topic of
interest in QS. It has been shown that the CAI-1 sensor CqsS plays a dominant
role over LuxPQ in modulating HapR levels. Thus, it remains possible that the
presence of CAI-1 allows for more robust regulation of HapR-regulated genes.
Indeed, CAI-1 has been shown to be critical for expression of the virulence factor
hapA (23), natural transformation (30), and for repressing chitobiose utilization as
shown in this study. HapA has been implicated in mediating *V. cholerae*
detachment from host epithelial cells, thereby promoting dissemination of cells
back into the aquatic environment (31, 32); natural transformation and chitin
utilization aid in *V. cholerae* fitness in the marine environment. By contrast,
expression of the major subunit of the toxin co-regulated pilus *tcpA* was shown to
be primarily regulated by AI-2 (22). It is tempting to speculate that CAI-1 controls
V. cholerae behaviors important for marine survival, whereas AI-2 controls
behaviors involved in infection (where the community is complex and has many
bacteria producing AI-2). It has been hypothesized that sensing both of these
autoinducers plays a critical role in biofilm dispersal, and that only when both are
sensed, *V. cholerae* cells are induced to leave a surface (i.e. QS works as a
coincidence detector for both signals) (22). Thus, signaling via distinct
autoinducers may allow *V. cholerae* to modulate responses depending on the
context of the environment they are inhabiting.
356
The data we present here suggests that at HCD, *V. cholerae* dampens its
expression of a chitin utilization locus. Below, we speculate on a few reasons
why this regulation may be advantageous. *V. cholerae* must secrete chitinases
that enzymatically degrade long chain insoluble chitin polymers into soluble chitin
oligosacchardies that can then be taken up and catabolized. This is an
energetically costly process, and as a result, liberated chitin is a valuable "public
good" in the context of a chitin biofilm (33). Thus, one possible explanation is that
QS regulation of chitin utilization allows *V. cholerae* to modulate chitin uptake
based on the composition of its microbial community. When other *V. cholerae*
cells are nearby, which will signal via the *V. cholerae*-specific autoinducer CAI-1,

chitin uptake may decrease among individual cells within the population in an
effort to "share" liberated chitin. By contrast, sensing the inter-species AI-2 does
not suppress chitin uptake and utilization.
370
Another possibility is that this regulation allows *V. cholerae* to control its
production of toxic metabolites. In a previous study, it was shown that cells in a
HCD state alter their metabolic flux to produce neutral byproducts as opposed to
organic acids (34). This regulation allows for a more stable community; by
contrast, cells locked in a LCD state will excrete harmful metabolic byproducts,
which leads to the demise of the community. Thus, it is possible that repressing
chitin uptake and utilization also helps slow the rate of metabolism to prevent the
accumulation of toxic intermediates in a dense community setting.
379
Materials & Methods
Bacterial strains and culture conditions
V. cholerae strains were routinely grown in LB medium and on LB agar
supplemented when necessary with Carbenicillin (20 µg/mL or 50 µg/mL),
Kanamycin (50 µg/mL), Spectinomycin (200 µg/mL), Trimethoprim (10 µg/mL),
and/or Chloramphenicol (2 µg/mL).
386
Transposon mutagenesis
Transposon mutant libraries were generated with a Carb^R mini-Tn10 transposon
exactly as previously described (35). Briefly, the transposon mutagenesis
plasmid pDL1086 was first mated into P$_{chb}$-*lacZ* Δcbp or P$_{chb}$-*lacZ* $\Delta clpA$ Δcbp
parent strains. The parent strains also carried a P$_{chb}$-mCherry reporter at an
ectopic site to ensure that candidate transposon mutants affected expression of
P$_{chb}$ and did not simply disrupt the P$_{chb}$-*lacZ* reporter. The activator screen parent
carried an additional copy of ChiS, as this is a known activator of this locus.
Transposition was induced by plating cultures on LB agar supplemented with 50
µg/mL Carb at 42°C. To screen colonies, plates also contained 40 µg/mL X-gal
and 5 mM IPTG. IPTG was added to competitively inhibit the basal activity of the
P$_{chb}$-*lacZ* reporter.
399
Generating mutant strains
V. cholerae E7946 served as the parent for all strains generated in this study,
except for those used in **Fig. S1** where we compare E7946 to A1552 (36, 37).
Mutant constructs were generated using splicing-by-overlap extension PCR
exactly as previously described (38). Transformations and cotransformations
were carried out by chitin-induced natural transformation exactly as previously
described (39, 40). Mutant strains were confirmed by PCR and/or sequencing. A
complete list of strains and primers used in this study are outlined in **Table S1**
and **Table S2**, respectively.
409
Measuring GFP and mCherry fluorescence in bulk populations
GFP and mCherry fluorescence in reporter strains was determined exactly as
previously described (41). Briefly, single colonies were grown in LB

(supplemented with 10 µM CAI-1 where noted) overnight at 30°C. The next day,
cells were washed and resuspended to an OD of 1.0 in instant ocean medium
(7g/L; Aquarium systems). Fluorescence was determined on a BioTek H1M plate
reader. For GFP measurements, excitation was set to 500 nm and emission was
set to 540 nm; for mCherry measurements, excitation was set to 580 nm and
emission was set to 610 nm.
419
420   *Antibody generation*
421   Purified LuxR protein (300 µg; purified as previously described (42)) was sent to
422   Cocalico Biologicals Inc. for serial injection into a rabbit host for antibody
423   generation. Serum obtained from the third bleed has been used for Western
424   analyses.
425
Western blotting
From overnight cultures, cells were concentrated to an OD_{600} of 20 in instant
ocean medium. Cells were lysed on a FastPrep-24 Classic Instrument at 4°C,
then lysates were clarified by centrifugation. Lysates were then boiled with an
equal volume of 2x SDS PAGE sample buffer (220 mM Tris pH 6.8, 25%
glycerol, 1.2% SDS, 0.02% Bromophenol Blue, and 5% β-mercaptoethanol).
Proteins were separated by SDS polyacrylamide gel electrophoresis,
electropheretically transferred to a PVDF membrane, and probed with rabbit
polyconal α-LuxR serum or mouse monoclonal α-RpoA (Biolegend) primary
antibodies. LuxR is the HapR homolog in *Vibrio harveyi* and has 72% identity and
86% similarity to *V. cholerae* HapR. Blots were then incubated with α-rabbit or α-
mouse HRP conjugated secondary antibodies, developed using Pierce ECL
Western Blotting Substrate (ThermoFisher), and exposed to film.
439
Chitin bead culturing
Chitin beads (200 µL of a 50% slurry; NEB) and overnight cultures of the
indicated strains were washed with defined artificial salt water medium (DASW),
which was prepared exactly as previously described (5). Chitin beads were
inoculated with *V. cholerae* cells to an OD of 0.1 in a final volume of 5 mL in a
Costar 6-well plate (Corning). Chitin mixtures were incubated statically at 30°C
for 7 days before imaging.
447
Microscopy data collection and analysis
To image chitin beads, cultured beads were gently transferred to a coverslip
using wide-bore pipette tips. To image chitin grown cells, chitin bead reactions
were vortexed then centrifuged at 250 x g for 1 minute. Cells found in the
supernatant were transferred to a coverslip. To image Δ*cbp* strains, overnight
cultures grown in LB were resuspended to an OD_{600} of 0.2 in DASW and then
transferred to a coverslip. Samples on the coverslip were placed under a 0.2%
gelzan pad and imaged on an inverted Nikon Ti-2 microscope with a Plan Apo
60x objective, YFP & mCherry filter cubes, a Hamamatsu ORCAFlash 4.0
camera, and Nikon NIS Elements imaging software.
458

The strains used to examine P_{chb} expression across a population contained 1) a
P_{chb}-mCherry reporter and 2) a reporter that drove constitutive expression of
GFP, P_{const2}-GFP. By using these reporters, the expression of the P_{chb}-mCherry
could be normalized to GFP expression in each cell. Images were analyzed on
Fiji using the MicrobeJ plugin (43) to determine mean intensity of cells in the YFP
and mCherry channels. GFP was assessed using the YFP channel to avoid
background fluorescence from chitin beads, which was stronger in the GFP
channel. Background fluorescence was subtracted from each channel and the
mCherry / GFP fluorescence was determined for each individual cell.
468
HapR protein purification
A plasmid expressing a hexahistidine-tagged *hapR* wild-type allele was
generated using Gibson assembly. The hapR insert was amplified from *V.*
cholerae E7946 and inserted into a pET28b vector (Novagen) using the primers
listed in **Table S2**. The plasmid was then electroporated into *E. coli* BL21(DE3)
for protein overexpression. This strain was grown overnight in LB medium with
kanamycin, back-diluted 1:100 into 1 L of LB medium with kanamycin, and grown
to an OD_{600} of 0.4-0.6 at 30°C. Expression of HapR was induced by IPTG to a
final concentration of 1 mM and cultures grown for 4 h shaking at 30°C. The cells
were pelleted and frozen at -80°C. The pellet was resuspended in 25 mL buffer A
(25 mM Tris pH 8, 500 mM NaCl), and an Avestin EmulfiFlex-C3 emulsifier was
used to lyse cells. The soluble lysate was applied to a HisTrap HP Ni-NTA
column using an Äkta Pure FPLC in buffer A and eluted from the column with a
gradient of buffer B (25 mM Tris pH 8, 500 mM NaCl, 1 M imidazole). The
purified protein was concentrated to approximately 5 ml using Sartorius Vivaspin
Turbo 10,000 MWCO centrifugal concentrators. The sample was manually
injected into the Äkta Pure and separated via size exclusion chromatography on
a HiLoad™ 16/600 Superdex™ 75 pg column equilibrated with gel filtration buffer
(25 mM Tris pH 7.5, 200 mM NaCl). Eluted fractions were analyzed by SDS-
PAGE, pooled, and concentrated using the same centrifugal concentrators
previously mentioned. The samples were then immediately frozen in liquid
nitrogen with a final concentration of 10% glycerol and stored at -80°C.
491
Electrophoretic mobility shift assay (EMSA)
Binding reactions contained 10 mM tris HCl pH 7.5, 1 mM EDTA, 10 mM KCl, 1
mM DTT, 50 µg/mL BSA, 0.1 mg/mL salmon sperm DNA, 5% glycerol, 1 nM of a
Cy5 labeled DNA probe, and HapR at the indicated concentrations (diluted in 10
mM tris pH 7.5, 10 mM KCl, 1 mM DTT, and 5% glycerol). Reactions were
incubated at room temperature for 20 minutes in the dark, then
electrophoretically separated on polyacrylamide gels in 0.5x Tris Borate EDTA
(TBE) buffer. Gels were imaged for Cy5 fluorescence on a Typhoon-9210
instrument. Short DNA probes (30bp) were made by end-labeling one primer of a
complementary pair (primers listed in **Table S2**) using 20 µM Cy5-dCTP and
Terminal deoxynucleotidyl Transferase (TdT; Promega). Complementary primers
(one labeled with Cy5 and the other unlabeled) were annealed by slow cooling at
equimolar concentrations in annealing buffer (10 mM Tris pH 7.5 and 50 mM

NaCl). P_{chb} and P_{clpP} probes were made by Phusion PCR, where Cy5-dCTP was
included in the reaction at a level that would result in incorporation of 1–2 Cy5
labeled nucleotides in the final probe as previously described.
508
ChIP-qPCR assays
Assays were carried out exactly as previously described (7). Briefly, overnight
cultures were diluted to an OD_{600} of 0.08 and then grown for 6 hours at 30°C.
Cultures were crosslinked using 1% paraformaldehyde, then quenched with a 1.2
molar excess of Tris. Cells were washed with PBS and stored at -80°C overnight.
The next day, cells were resuspended in lysis buffer (1x FastBreak cell lysis
reagent (Promega), 50 µg/mL lysozyme, 1% Triton X-100, 1 mM PMSF, and 1x
protease inhibitor cocktail; 100x inhibitor cocktail contained the following: 0.07
mg/mL phosphoramidon (Santa Cruz), 0.006 mg/mL bestatin
(MPbiomedicals/Fisher Scientific), 1.67 mg/mL AEBSF (DOT Scientific), 0.07
mg/mL pepstatin A (Gold Bio), 0.07 mg/mL E64 (Gold Bio)) and then lysed by
sonication, resulting in a DNA shear size of ~500 bp. Lysates were incubated
with Anti-FLAG M2 Magnetic Beads (Sigma), washed to remove unbound
proteins, and then bound protein-DNA complexes were eluted off with SDS.
Samples were digested with Proteinase K, then crosslinks were reversed. DNA
samples were cleaned up and used as template for quantitative PCR (qPCR)
using iTaq Universal SYBR Green Supermix (Bio-Rad) and primers specific for
the genes indicated (primers are listed in **Table S2**) on a Step-One qPCR
system. Standard curves of genomic DNA were included in each experiment and
were used to determine the abundance of each amplicon in the input (derived
from the lysate prior to ChIP) and output (derived from the samples after ChIP).
Primers to amplify *rpoB* served as a baseline control in this assay because HapR
does not bind this locus. Data are reported as 'Fold Enrichment', which is defined
as the ratio of the test promoter (P_{chb} or P_{clpP}) / *rpoB* found in the output divided
by the same ratio found in the input.
534
Chitobiose competition
Overnight cultures grown in LB were washed with M9 medium and mixed 1:1.
For each growth reaction, 10^2 cells of this mixture was added to M9 medium
supplemented with 0.2% chitobiose and 10 µM synthetic CAI-1 and grown
shaking at 30°C for 24 hours. After 24 hours, ~10^2 cells from this mixture was
used to inoculate fresh growth reactions to achieve additional generations of
growth on chitobiose. This was repeated a third time after another 24 hours. CAI-
1 was supplemented throughout these experiments to ensure consistently high
levels of HapR expression throughout transfer steps. After each 24 hour growth
period, the CFU/mL was determined for each strain in the mixture by dilution
plating on LB agar supplemented with X-gal. Competing strains were discerned
by blue/white screening as one strain was *lacZ*⁺ and the other was Δ*lacZ*. *V.*
cholerae can grow on chitobiose through the activity of two sugar transporters:
the GlcNAc PTS transporter (VC0995) or the chitobiose ABC transporter
(VC0616-0620). Because we wanted to study the regulation of the transporter
encoded within the *chb* locus, strains for this assay had a deletion in VC0995,

which renders growth on chitobiose dependent on the *chb* locus as previously
described (10).
553
554 *Statistics*
555 Statistical comparisons were determined using Student's t-test or One-way
556 ANOVA with Tukey's post-test using GraphPad Prism software.
557

558 **Acknowledgements**
559 We thank Ryan Chaparian for offering experimental advice, David Grainger for
560 helpful discussions, and Wai-Leung Ng for providing synthetic CAI-1. This work
561 was supported by grant R35GM128674 from the National Institutes of Health to
562 ABD and R35GM124698 from the National Institutes of Health to JVK.

563
564

References

1. Pruzzo C, Vezzulli L, & Colwell RR (2008) Global impact of Vibrio cholerae interactions with chitin. *Environ Microbiol* 10(6):1400-1410.
2. Colwell RR*, et al.* (2003) Reduction of cholera in Bangladeshi villages by simple filtration. *Proc Natl Acad Sci U S A* 100(3):1051-1055.
3. Huq A*, et al.* (2010) Simple sari cloth filtration of water is sustainable and continues to protect villagers from cholera in Matlab, Bangladesh. *mBio* 1(1).
4. Meibom KL*, et al.* (2004) The Vibrio cholerae chitin utilization program. *Proc Natl Acad Sci U S A* 101(8):2524-2529.
5. Meibom KL, Blokesch M, Dolganov NA, Wu CY, & Schoolnik GK (2005) Chitin induces natural competence in Vibrio cholerae. *Science* 310(5755):1824-1827.
6. Li X & Roseman S (2004) The chitinolytic cascade in Vibrios is regulated by chitin oligosaccharides and a two-component chitin catabolic sensor/kinase. *Proc Natl Acad Sci U S A* 101(2):627-631.
7. Klancher CAY, S.; Dalia, T. N.; Dalia, A. B. (2020) ChiS is a noncanonical DNA-binding hybrid sensor kinase that directly regulates the chitin utilization program in Vibrio cholerae. *In review*.
8. Yamamoto S & Ohnishi M (2017) Glucose-Specific Enzyme IIA of the Phosphoenolpyruvate:Carbohydrate Phosphotransferase System Modulates Chitin Signaling Pathways in Vibrio cholerae. *J Bacteriol* 199(18).
9. Blokesch M (2012) Chitin colonization, chitin degradation and chitin-induced natural competence of Vibrio cholerae are subject to catabolite repression. *Environ Microbiol* 14(8):1898-1912.
10. Klancher CA, Hayes CA, & Dalia AB (2017) The nucleoid occlusion protein SlmA is a direct transcriptional activator of chitobiose utilization in Vibrio cholerae. *PLoS Genet* 13(7):e1006877.
11. Sauer RT & Baker TA (2011) AAA+ proteases: ATP-fueled machines of protein destruction. *Annu Rev Biochem* 80:587-612.
12. Lee KJ, Jung YC, Park SJ, & Lee KH (2018) Role of Heat Shock Proteases in Quorum-Sensing-Mediated Regulation of Biofilm Formation by Vibrio Species. *mBio* 9(1).
13. Davis JH, Rubin AJ, & Sauer RT (2011) Design, construction and characterization of a set of insulated bacterial promoters. *Nucleic Acids Res* 39(3):1131-1141.
14. Lo Scrudato M & Blokesch M (2012) The regulatory network of natural competence and transformation of Vibrio cholerae. *PLoS Genet* 8(6):e1002778.
15. Sun S, Tay QX, Kjelleberg S, Rice SA, & McDougald D (2015) Quorum sensing-regulated chitin metabolism provides grazing resistance to Vibrio cholerae biofilms. *ISME J* 9(8):1812-1820.
16. van Kessel JC, Ulrich LE, Zhulin IB, & Bassler BL (2013) Analysis of activator and repressor functions reveals the requirements for

611     transcriptional control by LuxR, the master regulator of quorum sensing in
612     Vibrio harveyi. *mBio* 4(4).

613  17. Lee DH*, et al.* (2008) A consensus sequence for binding of SmcR, a Vibrio
614     vulnificus LuxR homologue, and genome-wide identification of the SmcR
615     regulon. *J Biol Chem* 283(35):23610-23618.

616  18. Chaparian RR, Tran MLN, Miller Conrad LC, Rusch DB, & van Kessel JC
617     (2020) Global H-NS counter-silencing by LuxR activates quorum sensing
618     gene expression. *Nucleic Acids Res* 48(1):171-183.

619  19. Jung SA, Chapman CA, & Ng WL (2015) Quadruple quorum-sensing
620     inputs control Vibrio cholerae virulence and maintain system robustness.
621     *PLoS Pathog* 11(4):e1004837.

622  20. Miller MB, Skorupski K, Lenz DH, Taylor RK, & Bassler BL (2002) Parallel
623     quorum sensing systems converge to regulate virulence in Vibrio
624     cholerae. *Cell* 110(3):303-314.

625  21. Surette MG, Miller MB, & Bassler BL (1999) Quorum sensing in
626     Escherichia coli, Salmonella typhimurium, and Vibrio harveyi: a new family
627     of genes responsible for autoinducer production. *Proc Natl Acad Sci U S A*
628     96(4):1639-1644.

629  22. Bridges AA & Bassler BL (2019) The intragenus and interspecies quorum-
630     sensing autoinducers exert distinct control over Vibrio cholerae biofilm
631     formation and dispersal. *PLoS Biol* 17(11):e3000429.

632  23. Hurley A & Bassler BL (2017) Asymmetric regulation of quorum-sensing
633     receptors drives autoinducer-specific gene expression programs in Vibrio
634     cholerae. *PLoS Genet* 13(5):e1006826.

635  24. Kroh HE & Simon LD (1990) The ClpP component of Clp protease is the
636     sigma 32-dependent heat shock protein F21.5. *J Bacteriol* 172(10):6026-
637     6034.

638  25. Grossman AD, Erickson JW, & Gross CA (1984) The htpR gene product
639     of E. coli is a sigma factor for heat-shock promoters. *Cell* 38(2):383-390.

640  26. Katayama Y*, et al.* (1988) The two-component, ATP-dependent Clp
641     protease of Escherichia coli. Purification, cloning, and mutational analysis
642     of the ATP-binding component. *J Biol Chem* 263(29):15226-15236.

643  27. Gerth U, Kruger E, Derre I, Msadek T, & Hecker M (1998) Stress induction
644     of the Bacillus subtilis clpP gene encoding a homologue of the proteolytic
645     component of the Clp protease and the involvement of ClpP and ClpX in
646     stress tolerance. *Molecular microbiology* 28(4):787-802.

647  28. Pennetzdorfer N*, et al.* (2019) Regulated Proteolysis in Vibrio cholerae
648     Allowing Rapid Adaptation to Stress Conditions. *Front Cell Infect Microbiol*
649     9:214.

650  29. Schild S*, et al.* (2007) Genes induced late in infection increase fitness of
651     Vibrio cholerae after release into the environment. *Cell Host Microbe*
652     2(4):264-277.

653  30. Suckow G, Seitz P, & Blokesch M (2011) Quorum sensing contributes to
654     natural transformation of Vibrio cholerae in a species-specific manner. *J*
655     *Bacteriol* 193(18):4914-4924.

31. Benitez JA*, et al.* (1997) Adherence of Vibrio cholerae to cultured
differentiated human intestinal cells: an in vitro colonization model. *Infect*
Immun 65(8):3474-3477.
32. Benitez JA, Silva AJ, & Finkelstein RA (2001) Environmental signals
controlling production of hemagglutinin/protease in Vibrio cholerae. *Infect*
Immun 69(10):6549-6553.
33. Drescher K, Nadell CD, Stone HA, Wingreen NS, & Bassler BL (2014)
Solutions to the public goods dilemma in bacterial biofilms. *Curr Biol*
24(1):50-55.
34. Hawver LA, Giulietti JM, Baleja JD, & Ng WL (2016) Quorum Sensing
Coordinates Cooperative Expression of Pyruvate Metabolism Genes To
Maintain a Sustainable Environment for Population Stability. *mBio* 7(6).
35. McDonough E, Kamp H, & Camilli A (2016) Vibrio cholerae phosphatases
required for the utilization of nucleotides and extracellular DNA as
phosphate sources. *Molecular microbiology* 99(3):453-469.
36. Miller VL, DiRita VJ, & Mekalanos JJ (1989) Identification of toxS, a
regulatory gene whose product enhances toxR-mediated activation of the
cholera toxin promoter. *J Bacteriol* 171(3):1288-1293.
37. Yildiz FH, Liu XS, Heydorn A, & Schoolnik GK (2004) Molecular analysis
of rugosity in a Vibrio cholerae O1 El Tor phase variant. *Molecular*
microbiology 53(2):497-515.
38. Dalia AB, Lazinski DW, & Camilli A (2013) Characterization of
undermethylated sites in Vibrio cholerae. *J Bacteriol* 195(10):2389-2399.
39. Dalia AB, McDonough E, & Camilli A (2014) Multiplex genome editing by
natural transformation. *Proc Natl Acad Sci U S A* 111(24):8937-8942.
40. Dalia AB (2018) Natural Cotransformation and Multiplex Genome Editing
by Natural Transformation (MuGENT) of Vibrio cholerae. *Methods Mol Biol*
1839:53-64.
41. Dalia AB (2016) RpoS is required for natural transformation of Vibrio
cholerae through regulation of chitinases. *Environ Microbiol* 18(11):3758-
3767.
42. Chaparian RR, Olney SG, Hustmyer CM, Rowe-Magnus DA, & van Kessel
JC (2016) Integration host factor and LuxR synergistically bind DNA to
coactivate quorum-sensing genes in Vibrio harveyi. *Molecular*
microbiology 101(5):823-840.
43. Ducret A, Quardokus EM, & Brun YV (2016) MicrobeJ, a tool for high
throughput bacterial cell detection and quantitative analysis. *Nat Microbiol*
1(7):16077.
694
695

Figure 1. *HapR is a repressor of P$_{chb}$ that is degraded by the ClpAP protease.* Expression of a P$_{chb}$-GFP reporter and HapR protein levels were determined in the indicated mutant strains. The parent strain contained a P$_{chb}$-GFP reporter and a Δcbp mutation. A representative western blot is shown below bars to indicate the protein levels for HapR and RpoA (a loading control) in the corresponding strains. Fluorescence of cultures was determined on a plate reader from at least six independent biological replicates and is shown as the mean ± SD. Statistical comparisons were made by one-way ANOVA with Tukey's post-test. NS, not significant. ***, $p < 0.001$.

722
Figure 2. *HapR-mediated repression of P_{chb} occurs on chitinous surfaces.* **A**)
Representative images of the indicated *V. cholerae* strains grown on chitin
beads. The parent strain background contains both a P_{chb}-mCherry reporter and
a P_{const2}-GFP construct, which exhibits constitutive GFP expression. Scale bar =
10 μm. **B-D**) P_{chb} expression in the indicated *V. cholerae cbp*^+ strains grown on
chitin beads. **B**) Representative images of cells that were vortexed off chitin
beads. Arrows demarcate cells where P_{chb} expression is induced ("P_{chb} On"
cells). Scale bar = 4 μm. **C**) Histogram showing the relative expression of the
P_{chb}-mCherry reporter. n = 2120 for Parent; n = 2191 for ΔhapR. Data shown are
from two independent biological replicates. **D**) Box and whisker plot showing

average mCherry intensity of the "P$_{chb}$ On" populations shown in **C**. Box
represents the first quartile, median, and third quartile while whiskers represent
the minimum and maximum values; n = 293 for Parent; n = 1092 for Δ*hapR.* **E-G**)
P$_{chb}$ expression of the indicated *V. cholerae* strains where P$_{chb}$ expression is
induced via deletion of *cbp.* **E**) Representative images of cells grown in the
absence of chitin. Scale bar = 4 µm. **F**) Histogram showing the relative
expression of a P$_{chb}$-mCherry reporter in the indicated *V. cholerae* strains. n =
2407 for Δ*cbp*; n = 2313 for Δ*hapR* Δ*cbp.* Data shown are from two independent
biological replicates. **G**) Box and whisker plot showing average mCherry intensity
of the populations depicted in **F**. Box represents the first quartile, median, and
third quartile while whiskers represent the minimum and maximum values; n =
2407 for WT; n = 2313 for Δ*hapR.* Statistical comparisons in **D** and **G** were made
using Student's t-test. ***, p < 0.001.
746

747
Figure 3. *HapR binds P$_{chb}$ in vitro and in vivo.* **A**) Promoter map of P$_{chb}$
highlighting putative HapR binding sites. Other sites required for P$_{chb}$ activation
(the SlmA binding site and the -35 and -10 signals) are highlighted. Also shown is
the transcriptional start site (+1) and the translational start site (+137). **B**) A
representative EMSA using HapR and Cy5-labeled DNA probes of the indicated
promoters. The P$_{chb}$ probe was incubated with (from left to right) 0 nM (-), 12.5
nM, 25 nM, 50 nM, 100 nM, 200 nM, 400 nM, or 800 nM purified HapR. The P$_{clpP}$
probe was incubated with 0 nM (-) or 800 nM HapR (+). **C**) A representative
EMSA using HapR and 32 bp Cy5-labeled probes encompassing each of the
putative HapR binding sites within the *chb* promoter. The 32 bp probes were
incubated with (from left to right) 0 nM (-), 100 nM, 200 nM, 400 nM, 800 nM, 1.6
µM, or 3.2 µM purified HapR. **D**) ChIP-qPCR assays showing enrichment of the
indicated promoters relative to *rpoB*, a reference locus that HapR does not bind
to. Data are from five independent biological replicates and shown as the mean ±
SD. Statistical comparisons were made by Student's t-test. *, p = 0.0240.
763

764
Figure 4. *Quorum sensing regulates expression of P$_{chb}$ through the cholera-*
specific autoinducer CAI-1. **A**) Model of QS regulation in *V. cholerae*. There are
four sensor kinases that contribute to canonical QS in *V. cholerae* to regulate
HapR production: LuxQ, CqsS, VpsS, and CqsR. The signals for VpsS and CqsR
are unknown. LuxPQ senses the interspecies autoinducer AI-2 and CqsS senses
the *V. cholerae*-specific autoinducer CAI-1. AI-2 is produced by the LuxS
synthase and CAI-1 is produced by the CqsA synthase. (Left) At low cell density,
the LuxQ and CqsS sensors act as kinases, resulting in phosphorylation of LuxO,
which subsequently decreases expression of HapR. (Right) At high cell density
when concentrations of AI-2 and CAI-1 are high, LuxQ and CqsS act as
phosphatases, which results in dephosphorylation of LuxO, and subsequent
activation of HapR expression. **B**) Expression of a P$_{chb}$-GFP reporter and HapR
protein levels were determined in the indicated strains. The parent strain contains
a P$_{chb}$-GFP reporter and a Δ*cbp* mutation. A representative western blot is shown
below the bars to indicate the protein levels for HapR and RpoA (a loading
control) in the corresponding strains. Statistical comparisons were made by one-
way ANOVA with Tukey's post-test. NS, not significant. ***, $p < 0.001$. The P$_{chb}$-
GFP fluorescence data for "Parent" and "ΔclpA" are identical to the data
presented in **Fig. 1** and were included here to ease comparisons.
784

Figure 5. *Model of quorum sensing regulation of chitobiose utilization genes in V.*
cholerae. When *V. cholerae* grows on a chitinous surface, the periplasmic chitin
binding protein (CBP) binds to chitin oligomers. This de-represses ChiS, which
can then activate expression of the chitobiose utilization operon (*chb*). **A**) At low
CAI-1 concentrations, the CqsS sensor indirectly promotes LuxO phosphoylation,
which indirectly blocks HapR expression. In this state, P_{chb} is maximally
expressed. **B**) At high CAI-1 concentrations, the CqsS sensor indirectly
dephosphorylates LuxO, which results in activation of HapR expression. HapR
then exerts a repressive effect on P_{chb} by decreasing expression of this locus ~2-
fold in the presence of the ClpAP protease machine. If HapR is not degraded by
ClpAP, its repressive effect can result in a ~7-fold decrease in P_{chb} expression.
797

Figure S1. *HapR is a repressor of P$_{chb}$ in V. cholerae E7946 and V. cholerae A1552.* Expression of a P$_{chb}$-mCherry reporter was determined in the indicated mutant strains. Each respective parent strain contained a P$_{chb}$-mCherry reporter and a Δ*cbp* mutation to activate ChiS.

837 **Figure S2.** A Δ*hapR mutant does not have a fitness advantage for growth on*
838 *chitobiose.* The indicated *V. cholerae* strains were mixed and co-cultured in M9
839 minimal media supplemented with 0.2% chitobiose + 10 µM CAI-1 and grown for
840 the indicated number of generations.
841

Figure S3. *Sequence of predicted HapR binding sites.* Sequences of the two 32

bp probes used in EMSAs in **Fig. 3C** (for HapR BS 1 and HapR BS 2) are

indicated. Predicted HapR binding sites are bolded and bracketed in green;

conserved nucleotide residues are underlined. HapR BS 1 was predicted from

the SmcR consensus site and HapR BS 2 was predicted from the LuxR

consensus site. The SlmA binding site is indicated by the blue bracket and the -

35 signal is indicated by the orange bracket.

867 **Figure S4.** *N-terminally FLAG-tagged HapR is functional for regulation of P$_{chb}$.*
868 Strains expressing the indicated HapR allele at the native locus were assessed
869 for expression of a P$_{chb}$-GFP reporter. Both strains also harbored a Δcbp deletion
870 to activate ChiS. Statistical comparisons were made using Student's t-test. NS,
871 not significant.
872
873

Table S1. *Strains used in this study.*

Strain Designation (Reference)	Reference in Manuscript	Genotype
SAD 2825/CAK 550 (This Study)	Activator screen strain	Pchb-mCherry @ VC1807, KanR; Pchb-lacZ @ lacZ, SpecR; ChiS @ VCA0692, TmR; Δ3' CBP; pDL1086, CmR CarbR
SAD 2826/CAK 762 (This Study)	ΔclpA counter-screen strain	Pchb-mCherry @ VC1807, KanR; Pchb-lacZ @ lacZ, SpecR; ΔclpA; Δcbp::TmR; pDL1086, CmR CarbR
SAD 1309/CAK 009 (10)	Fig. 1 & 4 Parent; Fig. S4 HapR WT	Pchb-GFP @ lacZ, KanR; Δcbp::SpecR
SAD 2827/CAK 692 (This Study)	Fig. 1 & 4 ΔclpA	Pchb-GFP @ lacZ, KanR; ΔclpA::CarbR; Δcbp::SpecR
SAD 2828/CAK 710 (This Study)	Fig. 1 ΔclpP	Pchb-GFP @ lacZ, KanR; ΔclpP::CarbR; Δcbp::TmR
SAD 2829/CAK 734 (This Study)	Fig. 1 ΔclpAP	Pchb-GFP @ lacZ, KanR; ΔclpA::CarbR; ΔclpP::TmR; Δcbp::SpecR
SAD 2830/CAK 708 (This Study)	Fig. 1 ΔclpS	Pchb-GFP @ lacZ, KanR; ΔclpS::CarbR; Δcbp::TmR
SAD 2831/CAK 712 (This Study)	Fig. 1 ΔclpX	Pchb-GFP @ lacZ, KanR; ΔclpX::CarbR; Δcbp::TmR
SAD 2832/CAK 843 (This Study)	Fig. 1 ΔlonA	Pchb-GFP @ lacZ, KanR; Δlon::TmR; Δcbp::CarbR
SAD 2833/CAK 786 (This Study)	Fig. 1 ΔhapR ΔclpA	Pchb-GFP @ lacZ, KanR; ΔhapR::SpecR; ΔclpA::CarbR; Δcbp::TmR
SAD 2834/CAK 665 (This Study)	Fig. 1 ΔhapR	Pchb-GFP @ lacZ, KanR; ΔhapR::SpecR; Δcbp::CarbR
SAD 2835/CAK 1492 (This Study)	Fig. 2 Parent	Pchb-mCherry @ lacZ, KanR; Pconst2-GFP @ VCA0692, SpecR
SAD 2836/CAK 1505 (This Study)	Fig. 2 ΔhapR	Pchb-mCherry @ lacZ, KanR; Pconst2-GFP @ VCA0692, SpecR; ΔhapR::CmR
SAD 2837/CAK 1498 (This Study)	Fig. 2 Δcbp & Fig. S1 E7946 Parent	Pchb-mCherry @ lacZ, KanR; Pconst2-GFP @ VCA0692, SpecR; Δcbp::TmR
SAD 2838/CAK 1502 (This Study)	Fig. 2 ΔhapR Δcbp & Fig. S1 E7946 ΔhapR	Pchb-mCherry @ lacZ, KanR; Pconst2-GFP @ VCA0692, SpecR; ΔhapR::CmR; Δcbp::TmR
SAD 2839/CAK 869 (This Study)	Fig. 3 ChIP strain and Fig. S4 1x FLAG-HapR	Pchb-GFP @ lacZ, KanR; 1x FLAG-HapR; Δcbp::TmR
SAD 2840/CAK 663 (This Study)	Fig. 4 ΔluxO	Pchb-GFP @ lacZ, KanR; ΔluxO::SpecR; Δcbp::CarbR
SAD 2841/CAK 886 (This Study)	Fig. 4 luxO^{D47E}	Pchb-GFP @ lacZ, KanR; LuxO D47E; Δcbp::SpecR
SAD 2842/CAK 917 (This Study)	Fig. 4 ΔluxS	Pchb-GFP @ lacZ, KanR; ΔluxS::CmR; Δcbp::SpecR
SAD 2843/CAK 915 (This Study)	Fig. 4 ΔcqsA	Pchb-GFP @ lacZ, KanR; ΔcqsA::TmR; Δcbp::SpecR
SAD 2844/CAK 1504 (This Study)	Fig. 4 ΔcqsA ΔhapR	Pchb-GFP @ lacZ, KanR; ΔcqsA::TmR; ΔhapR::CmR; Δcbp::SpecR
SAD 2845/CAK 1506 (This Study)	Fig. S1 A1552 Parent	V. cholerae A1552 Pconst2-GFP @ VCA0692, SpecR; Pchb-mCherry @ lacZ, KanR; Δcbp::TmR
SAD 2846/CAK 1510 (This Study)	Fig. S1 A1552 ΔhapR	V. cholerae A1552 Pconst2-GFP @ VCA0692, SpecR; Pchb-mCherry @ lacZ, KanR; ΔhapR::CmR; Δcbp::TmR
SAD 2847/CAK 942 (This Study)	Fig. S2 WT lacZ+	ΔVC1807::CmR; ΔVC0995
SAD 2848/CAK 944 (This Study)	Fig. S2 WT ΔlacZ	ΔlacZ::SpecR; ΔVC1807::CmR; ΔVC0995

SAD 2849/CAK 1026 (This Study)	Fig. S2 ΔhapR ΔlacZ	ΔlacZ::SpecR; ΔVC1807::CmR; ΔVC0995; ΔhapR::TmR
JDN 71 (This Study)	Fig. 3B, 3C purified HapR	*E. coli* BL21(DE3) harboring pET28b-6XHis-HapR, KanR

Table S2. *Primers used in this study.*

Pimers for Reporter Constructs		
Primer	**Sequence**	**Description**
ABD 332	GGCTGAACGTGGTTGTCGAAAATGAC	ΔlacZ F1 (Up Arm)
BBC 219	GTTTATTTTTGTCGACTGTACAGCGTTTAAATAGAGGTCGATATTGACCC	ΔlacZ R1 (Up Arm)
ABD 342	ATTTTTCAGTTGGCCTACAATGCTTTCC	ΔVC1807 F1 (Up Arm)
BBC 719	CACCATACCCACGCCGAAACAAGGATTTTGAATTAAACGTTTCATTAGTC	ΔVC1807 R1 (Up Arm)
BBC 218	CGCTGTACAGTCGACAAAAATAAAC	Kan^R F (Middle Arm)
BBC 262	TACCGAGGACGCGAAGCTG	Kan^R R (Middle Arm)
BBC 266	CAGCTTCGCGTCCTCGGTAGAATAAAGCAATCCGCAAGCG	P_{chb} F (Middle Arm)
BBC 267	CCCGGGATCCTGTGTGAAATTGAGTTGCTTTCATTTCACTAATGG	P_{chb} R (Middle Arm)
BBC 732	ctcaagccgaggagtaaagaagAGTTGCTTTCATTTCACTAATGG	P_{chb} fuse to lacZ R (Middle Arm)
BBC 252	CAATTTCACACAGGATCCCGGGAGGAGGTAACGTAATGCGTAAAGGAGAAGAAC	GFP F (Middle Arm)
BBC 254	tgtaggctggagctgcttcTTAGTTGTATAGTTCATCCATGCC	GFP R (Middle Arm)
BBC 1356	TTGTTTCGGCGTGGGTATGGTGCGCTGTACAGTCGACAAAAATAAAC	mCherry F (Middle Arm)
BBC 206	tgtaggctggagctgcttcttacttgtacagctcgtccatg	mCherry R (Middle Arm)
ABD 785	CTTCTTTACTCCTCGGCTTGAG	P_{chb} stitch to lacZ F (Down Arm)
ABD 255	gaagcagctccagcctacaCCACAATAAGCCAGAGAGCCTTAAG	ΔlacZ F2 (Down Arm)
ABD 256	CCCAAATACGGCAACTTGGCG	ΔlacZ R2 (Down Arm)
ABD 341	gaagcagctccagcctacaTAGTCGAAAATAAAAAAAAGAGGCTCGCCTC	ΔVC1807 F2 (Down Arm)
ABD 345	CTTGCTAACCGTTGGTGTTACCAGC	ΔVC1807 R2 (Down Arm)
BBC 3083	tggataactttacgggcatgcataaggctcgtataatatattcagggagaccacaacggtttccctctacaaataattttgtttaactttCAATTTCACACAGGATCCCGGG	P_{const2} F
BBC 3082	ttatacgagccttatgcatgcccgtaaagttatccagcaaccactcatagacctagggcagcagatagggacgacgtggtgttagctgtgCTCATTAGGCACCCCAGGC	P_{const2} R
Primers for SOE deletion, FLAG fusions, and point mutants		
Primer	**Sequence**	**Description**
BBC 742	attccggggatccgtcgacCTGCAGTTCagaagcagctccagcctaca	MiniFRT F
BBC 743	tgtaggctggagctgcttctGAACTGCAGgtcgacggatccccggaat	MiniFRT R
ABD 123	ATTCCGGGGATCCGTCGAC	Kan^R, $Spec^R$, $Carb^R$, Cm^R or Tm^R cassette F
ABD 124	TGTAGGCTGGAGCTGCTTC	Kan^R, $Spec^R$, $Carb^R$, Cm^R or Tm^R cassette R
ABD 796	TTAGAATCTGCGCCAGAAGCG	Δcbp F1 (Up Arm)
ABD 797	gtcgacggatccccggaatCATAGCTGTTCCTTACTAGTTGC	Δcbp R1 (Up Arm)
BBC 920	gtcgacggatccccggaatGCTCATCAGGTCGTCAGCC	Δ3'cbp R1 (Up Arm)

ABD 798	gaagcagctccagcctacaGTACTGGATCTGAAACCAGTTAAG	Δcbp F2 (Down Arm)
ABD 799	GTATTGCGGAATGACCAGCATG	Δcbp R2 (Down Arm)
BBC 2755	TTACCCCTAAGTCGGCGAGC	ΔVC0995 F1 (Up Arm)
BBC 2756	gtcgacggatccccggaatAATATTCACCTTAAGTTCCCCC	ΔVC0995 R1 (Up Arm)
BBC 2757	gaagcagctccagcctacaCTGCCTTAATCGAGTTTAAACCC	ΔVC0995 F2 (Down Arm)
BBC 2758	GCACCACGATAGCAATAAGC	ΔVC0995 R2 (Down Arm)
CAK 407	GCTACGCACTGCCAAATTACC	ΔclpA or ΔclpS F1 (Up Arm)
CAK 408	gtcgacggatccccggaatAAGCATAAGGCCTCCTTAAGGAAC	ΔclpA R1 (Up Arm)
CAK 409	gaagcagctccagcctacaCACTAGACATCTCACAATACGC	ΔclpA F2 (Down Arm)
CAK 410	CCGCTAACATCTCAGGACTG	ΔclpA or ΔclpS R2 (Down Arm)
CKP 449	TGCTCGGTTTTGATCCGTTC	ΔclpP or ΔclpX F1 (Up Arm)
CKP 450	gtcgacggatccccggaatGGGCGACATTGCATTTTTTTC	ΔclpP R1 (Up Arm)
CKP 451	gaagcagctccagcctacaGGCGAGTAAGCTCGTAATTG	ΔclpP F2 (Down Arm)
CKP 454	GTCAAACTAGAAACCAGCGC	ΔclpP or ΔclpX R2 (Down Arm)
CKP 447	gtcgacggatccccggaatTTTACTCATGACACTTCAATATTTG	ΔclpS R1 (Up Arm)
CKP 448	gaagcagctccagcctacaCAAGCTTAAACTTGCCGGGC	ΔclpS F2 (Down Arm)
CKP 452	gtcgacggatccccggaatGTCTGTCATTCGCTAACCTC	ΔclpX R1 (Up Arm)
CKP 453	gaagcagctccagcctacaGCGGAGTAATCTCAAGCAAC	ΔclpX F2 (Down Arm)
CKP 489	TTGACGCTCTGAAAGCAGAAG	ΔlonA F1 (Up Arm)
CKP 490	gtcgacggatccccggaatCAAGTTCATATTTTTCTCTCTTCCG	ΔlonA R1 (Up Arm)
CKP 491	gaagcagctccagcctacaGATGCATAGCAAAAATAAGTAAATC	ΔlonA F2 (Down Arm)
CKP 492	CGAAGAATTATAAGTGCAAAGGC	ΔlonA R2 (Down Arm)
BBC 422	ACGTTCAATAACCAGAATGTTGG	ΔhapR F1 (Up Arm)
BBC 423	gtcgacggatccccggaatTTTTCGATTGATGCGTCCATAG	ΔhapR R1 (Up Arm)
CKP 511	GTCtttgtcatcgtcatccttataatcCATAGGGGTATATCCTTGCC	1x FLAG-HapR R1 (Up Arm)
BBC 408	gaagcagctccagcctacaAACTAGTTTCTTGGGCAGCACAA	ΔhapR F2 (Down Arm)
CKP 510	ATGgattataaggatgacgatgacaaaGACGCATCAATCGAAAAACG	1x FLAG-HapR F2 (Down Arm)
BBC 409	GTTCATAATGATTTCCTTGGTGCC	ΔhapR R2 (Down Arm)
BBC 374	TGGCAAAAAGCGAGAGAAGAAG	ΔluxO F1 (Up Arm)
BBC 375	gtcgacggatccccggaatCATGAGGACATATTTTGTTTTCTGC	ΔluxO R1 (Up Arm)
CKP 519	TCATATCTGGCAAACGTAAcTCCAGCAGGATTAAGTCAGG	luxO^{D47E} R1 (Up Arm)
BBC 376	gaagcagctccagcctacaTAAGCGATGAGAGAATGGATCAAC	ΔluxO F2 (Down

		Arm)
CKP 518	TGACTTAATCCTGCTGGAgTTACGTTTGCCAGATATGACG	luxO^{D47E} F2 (Down Arm)
BBC 377	TCACACCCGAATTTCCATCATGC	ΔluxO R2 (Down Arm)
CKP 554	GTCTCTTAGCCGAGGTACTG	ΔcqsA F1 (Up Arm)
CKP 555	gtcgacggatccccggaatCTTGTTCATCGCAATATATCCTAG	ΔcqsA R1 (Up Arm)
CKP 556	gaagcagctccagcctacaTTTCGTTAAATGCATAAATAACAAAAAC	ΔcqsA F2 (Down Arm)
CKP 557	AGTTGGAACCACTTCTTGTC	ΔcqsA R2 (Down Arm)
CKP 549	TACAACTGCTTGGCACGC	ΔluxS F1 (Up Arm)
CKP 550	gtcgacggatccccggaatTGGCATTTCCTTTCTCCC	ΔluxS R1 (Up Arm)
CKP 551	gaagcagctccagcctacaCACTAAGTCGGTTCTGTAAACG	ΔluxS F2 (Down Arm)
CKP 552	TTCTTAGCGTGATCAATTGC	ΔluxS R2 (Down Arm)
ABD 334	AGTGCTCCGGACTCTTTGCTCTG	ΔlacZ F1 (Up Arm)
ABD 254	gtcgacggatccccggaatCATCCCTCAAGCCGAGGAGTAAAG	ΔlacZ R1 (Up Arm)
ABD 255	gaagcagctccagcctacaCCACAATAAGCCAGAGAGCCTTAAG	ΔlacZ F2 (Down Arm)
ABD 256	CCCAAATACGGCAACTTGGCG	ΔlacZ R2 (Down Arm)
ABD 725	GAAGCAGCTCCAGCCTACA	Detect F for all deletions
BBC 082	gtcgacggatccccggaatCATAACTTACACCTTACTCACCCAG	Δcbp detect R
CAK 411	TTGTTTGGTGCGATTATTGG	ΔclpA detect R
CKP 464	TTGGCAGAACATCTTTGATC	ΔclpP detect R
CKP 463	AAAGTATCCAGCTCACGGCG	ΔclpS detect R
CKP 465	CCATGTGTGGATAGACAACC	ΔclpX detect R
CKP 493	CAGTGCCACTTGGTCACCTG	ΔlonA detect R
BBC 410	TAAATGGGGCTTGGAGAATTTAG	ΔhapR detect R
BBC 1911	CGTAATCAAACTGCGAAAGTG	ΔluxO detect R
CKP 558	AATTCGTAACTCTGAGCATG	ΔcqsA detect R
CKP 553	TGGACCACGAACCTTAAACG	ΔluxS detect R
BBC 2759	TTGTCAGAAAGCGTTTCTGC	ΔVC0995 detect R
ABD 399	AACTGATGGCAGAAAAAGCCACTCAG	ΔlacZ detect R
BBC 993	ttgattataaggatgacgatgac	1x FLAG detect F
ABD 846	CATAAACATGTTTCTGATCAGCAG	1x FLAG HapR detect R
BBC 380	GCCAATAGAATGAGTCTATTGGCTG	luxO^{D47E} detect F
CKP 520	TCATATCTGGCAAACGTAgc	luxO^{D47E} detect R
Primers for EMSA probes		
Primer	**Sequence**	**Description**
BBC 744	cagcttcgcgtcctcggtaCGCAAATATAACTCAGGCAAAG	P$_{chb}$ F
CKP 072	cccgggatcctgtgtgaaattgCTTTGGCAGGAGTAAGAAAACACCTAG	P$_{chb}$ R
CKP 865	TAAGCAAACTGTAGCGTAGAAG	P$_{clpP}$ F
CKP 866	TGCATTTTTTTCTTGGTAGC	P$_{clpP}$ R
CKP 504	GCCAAATAAGTAAGTAAACACTAACAAACCTA	HapR BS 1 F
CKP 505	TAGGTTTGTTAGTGTTTACTTACTTATTTGGC	HapR BS 1 R
CKP 506	AATTGCAATTGATAAATTTCTTGGCTAGGTGT	HapR BS 2 F
CKP 507	ACACCTAGCCAAGAAATTTATCAATTGCAATT	HapR BS 2 R
Primers for qPCR		
Primer	**Sequence**	**Description**

BBC 989	GCATCTAGGTTTTGACGTTTTTAACG	P$_{chb}$ amplify F
BBC 990	AACACTCTCCAAGACCTACCTC	P$_{chb}$ amplify R
ABD 132	CTGTCTCAAGCCGGTTACAA	rpoB amplify F
ABD 133	TTTCTACCAGTGCAGAGATGC	rpoB amplify R
CKP 865	TAAGCAAACTGTAGCGTAGAAG	P$_{clpP}$ amplify F
CKP 867	TCGTATGAACGCTCACCACG	P$_{clpP}$ amplify R
Primers for purification vector		
Primer	**Sequence**	**Description**
JDN 92	agcagcggcctggtgccgcgcggcagcCatatggacgcatcaatcgaaaaa	Cloning 6X-His HapR into pET28b vector via IDA, F
JDN 93	tcagtggtggtggtggtggtgCTCGAGctagttcttatagatacacagcatattgagg	Cloning 6X-His HapR into pET28b vector via IDA, R
JDN 22	CTCGAGcaccaccaccacca	Amplify pET28b vector backbone for IDA, F
JDN 23	atGgctgccgcgcggcacca	Amplify pET28b vector backbone for IDA, R

880