Structure of the bacterial cellulose ribbon and its assembly-guiding cytoskeleton by electron cryotomography

William J. Nicolas[1,2], Debnath Ghosal[1,3], Elitza I. Tocheva[1,4], Elliot M. Meyerowitz[1,2] and Grant J. Jensen[1,2]

[1] Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, California 91125

[2] Howard Hughes Medical Institute, Pasadena, California 91125

[3]Present address: Division of Medicine, Dentistry and Health Sciences, University of Melbourne Parkville VIC 3010, Australia

[4]Present address: Department of Microbiology and Immunology, University of British Colombia, Vancouver, British Columbia V6T 1Z3

Abstract

Cellulose is a widespread component of bacterial biofilms, where its properties of exceptional water retention, high tensile strength and stiffness prevents dehydration and mechanical disruption of the biofilm. Bacteria in the *Gluconacetobacter* genus secrete crystalline cellulose, with a structure very similar to that found in plant cell walls. How this higher-order structure is produced is poorly understood. We used cryo-electron tomography and focused ion beam milling of native bacterial biofilms to image cellulose-synthesizing *G. hansenii* and *G. xylinus* bacteria in a frozen-hydrated, near-native state. We confirm previous results suggesting that cellulose crystallization occurs serially following its secretion along one side of the cell, leading to a cellulose ribbon that can reach several microns in length and combine with ribbons from other cells to form a robust biofilm matrix. We were able to take direct measurements in a near-native state of the cellulose sheets. Our results also reveal a novel cytoskeletal structure, that

we name the cortical belt, adjacent to the inner membrane and underlying the sites where

cellulose is seen emerging from the cell. We find that this structure is not present in other

cellulose-synthesizing bacterial species, *Agrobacterium tumefaciens* and *Escherichia coli* 1094,

which do not produce organized cellulose ribbons. We therefore propose that the cortical belt

holds the cellulose synthase complexes in a line, to form higher-order cellulose structures such

as sheets and ribbons.

Introduction

Humans rely on cellulose for building material, clothing and fuel[1–3]. More recently the polymer

has sparked interest in the biotechnology field as a potential source of biofuel feedstock[4], and

in the biomedical industry as a biologically neutral scaffold to promote tissue regeneration[5,6] .

Cellulose is a linear polymer of glucose molecules connected with β1-4 linkages by a

glucosyltransferase. Individual linear glucan chains can pack via hydrogen bonding and Van

Der Waals interactions in various ways to form different types of celluloses, with different

properties[3,7,8]. The most common way glucan chains organize in nature is to form hydrogen-

bonded planes which then stack into parallel layers via Van Der Waals interactions. These

stacked layers give rise to cellulose I microfibrils, or "native cellulose", that can then coalesce

to form larger arrays. Because glucan chains pack in a regular lattice, cellulose I is considered

crystalline. This form of cellulose is mainly found in plants, where it is a major structural

element of the cell wall[9].

In the prokaryotic world, cellulose is an important component of bacterial biofilms[10,11], which

increase cells' tolerance for a range of biotic and abiotic stresses and enhance surface adhesion,

cell cooperation and resource capture[10]. Cellulose-containing biofilms have also been

implicated in pathogenicity, enabling bacteria to resist antibiotics and disinfection[12,13]. Most

cellulose-synthesizing bacteria produce amorphous (non-crystalline) cellulose, but a few genera, including *Gluconacetobacter*, can produce cellulose I microfibrils like those found in plants. In *Gluconacetobacter*, these crystalline cellulose microfibrils can further aggregate into wide ribbon structures and larger arrays, giving rise to thick biofilms of pure cellulose I[14] .

Bacterial cellulose is synthesized by an envelope-spanning machinery called the Bacterial Cellulose Synthase (BCS) complex, encoded by the BCS operon and first identified in *Gluconacetobacter*[11]. While the components vary, most of the species encode BcsA, a component in the inner membrane that, with BcsB, catalyzes transfer of UDP-glucose to the nascent glucan chain[11,15,16]. BcsD forms a periplasmic ring that gathers glucan chains from several BcsA/B units[17]. BcsA and B are essential for cellulose synthesis *in vivo*, and BcsD is essential for the crystallization of nascent glucan chains[18]. BcsC forms a pore in the OM and very recent work has solved its crystallographic structure[19]. Consistent with previous data relying on sequence homology with the exopolysaccharide secretin components AlgE and AlgK from *P. aeruginosa*, BcsC is found to form an outer-membrane β-barrel pore at its C-terminal end, secreting the nascent elementary cellulose fibrils into the environment[18–22]. It is hypothesized that the elementary cellulose fibrils can aggregate with neighboring elementary fibrils upon secretion to form microfibrils[23,24]. Genes flanking the operon, *cmcax, ccpAX* and *bglxA*, are essential for cellulose crystallization and despite knowledge of their enzymatic functions, how they take part in this process is unclear[24–27].

In the following report, the terms used to describe the cellulose assembly process are adapted from the ones defined in[24], elaborating on the cell-directed hierarchical model for cellulose crystallization[7,28]. Glucan chains are linear polymers of β-1,4 linked glucose residues synthesized by a single catalytic site of a cellulose synthase. An elementary fibril (also termed mini-crystal in previous work[28–30]) is the product of the periplasmic aggregation of multiple glucan chains which is then extruded through a single BcsC subunit into the environment.

Microfibrils result from the aggregation of several elementary fibrils, at least three according

to earlier work[30], outside the cell. These microfibrils can then crystallize into sheets that stack

on each other to form ribbons.

Much work has already been done to understand the synthesis of crystalline cellulose[16,17,34–

37,18,25–27,29,31–33] . In particular freeze-fracture electron microscopy (EM) studies have found that

the *G. hansenii* BCS complexes are arrayed linearly along the side of the cell[29,36,37], and this

arrangement seems to determine the extracellular organization of cellulose I into ribbons[29,37].

How this linear arrangement is achieved is not known.

Here we used cryo-electron tomography (cryo-ET) of isolated cells and cryo-Focused Ion Beam

(FIB)-milled biofilms to visualize native cellulose production in *G. hansenii* and *G. xylinus*,

allowing the morphological characterization of the cellulose ribbons in a near-native state. We

identified a novel cytoplasmic structure, which we call the cortical belt. We found that this

cortical belt is absent from *Escherichia coli* 1094, which produces amorphous cellulose, and

Agrobacterium tumefaciens, which produces crystalline microfibrils but not higher-order

sheets, suggesting that the cortical belt functions to align BCS complexes to produce cellulose

sheets.

Results

<u>Cellulose is laid out in stacked sheets on one side of the cells.</u>

To visualize bacterial cellulose production, we used cryo-ET to image intact frozen-hydrated

G. hansenii cells separated from their cellulose biofilm according to the original method from

Brown et al. 1976. Previous work showed that newly synthesized cellulose ribbons are visible

under the electron microscope at one hour post-separation[36]. To assure that the cells would have

enough time to synthesize cellulose ribbons we imaged cells 5 hours (300 minutes) after

separation. To confirm cellulose production, we stained cells with mitoTracker Deep Red FM to visualize membranes and Calcofluor-White to visualize cellulose. By confocal imaging, we observed cellulose filamentous structures tens of microns long (Fig. 1A and B, cyan arrowheads). We next plunge-froze cells at the same timepoint and imaged them by cryo-ET. The rod-shaped cells always lay flat on the grids, but their long axis was oriented randomly in the grid plane. Of 33 cells imaged, we found putative cellulose ribbons associated with 29 (88%), always on one side of the cell, including the top and bottom, and always aligned with the cell's long axis (Fig 1C-E, yellow arrows). To confirm that the ribbon was in fact cellulose, we treated cells with cellulase and observed a large reduction in the occurrence of ribbons in cryo-EM and negative stained images (Supplemental figure 1, yellow arrowheads). Instead, we observed aggregated material we think is likely partially digested cellulose (Supplemental figure 1F, orange arrowheads).

The spatial relation between the cellulose ribbons and the OM was examined. In 3 out of the 29 tomograms, the cellulose ribbon was observed running beneath or on top of the cell, causing it to be normal to the electron beam thus inherently not well resolved and difficult to assess its spatial relation with the OM. Therefore, data from these 3 tomograms was excluded for these measurements. In the remaining tomograms two distinct configurations were observed: a "tight" configuration in 23 out of 26 tomograms (88%), where the average outer membrane (OM)-to-ribbon distance was 16 ± 5 nm (n = 23) (Fig. 1C-H, supplemental video 1), and a "loose" configuration in 3 out of 26 tomograms (12%), where the average OM-to-closest sheet distance was 99 ± 49 nm (n = 3) (Fig. 2). Among the tomograms showing a "tight" configuration, 17 out of 23 (65%) displayed multiple clear direct contacts between the OM and the ribbon (Fig. 1F-H, white arrows). Tomograms in the "loose" configuration exhibited ribbons that seemed detached from the OM, with an increased OM-to-closest sheet distance compared to the "tight" configuration (Fig. 2E). All three tomograms presented disorganized

aggregates bearing a mesh-like appearance between the OM and the ribbon (Fig. 2A-D, orange asterisks and dashed bracket). These aggregates always connected to the ribbon (Fig. 2A, black lined orange arrows). Similar cellulose aggregates have been seen previously by negative staining[23].

These cells and their cellulose structures (the ribbons) were imaged in a near-native (frozen hydrated) state, allowing measurement of their native dimensions. In our description of the cellulose ribbons below, by length we mean the dimension parallel to the long axis of the cell (Fig. 3A). By thickness we refer to the dimension normal to the cell surface (Fig. 3A, black inset). By width we refer to the dimension tangential to the cell surface (Fig. 3B). The cellulose ribbons we observed were very similar to what has been seen previously by negative stain EM[23,36]. Ribbons comprised long flexible sheets, too long to be measured by cryo-ET because they are never entirely in the field of view. All structures in an electron tomogram suffer from the missing wedge artefact, inherent to electron tomography, which causes elongation in the direction parallel to the electron beam[38,39]. For this reason, measurements of the width of the cellulose sheets are systematically overestimated. Sheet width was estimated at 38 ± 14 nm (n = 45) (Fig. 3C). These sheets then stack into a ribbon (2.3 ± 0.9 sheets on average; n = 24), with a variable inter-sheet distance (16 ± 7 nm; n = 23). Inter-sheet distance was accurately measured peak-to-peak (Fig. 3D), which encompasses 2 halves of the two neighboring sheets' density and the space between them (Fig. 3A, black inset). Because the apparent thickness of single densities in cryo-ET is strongly affected by the defocus applied, individual cellulose sheet thickness measurements will be overestimated. Therefore, we can only say confidently that they are thinner than the inter-sheet distance. Despite careful inspection, although densities could be seen in the periplasmic space, we did not recognize a consistent shape which we could confidently attribute to the BCS machinery. This is likely due to the large cell diameter (~800nm), and the small size and/or flexibility of the BCS complexes.

Sheets arise from the stacking of microfibrils

To visualize earlier stages of cellulose synthesis, we plunge-froze cells at earlier timepoints

after separation from the biofilm. A total of 6 and 15 tomograms were acquired at 13- and 20-

minutes post-separation, respectively. At 13 minutes (the most quickly we could complete

plunge freezing), no cells exhibited a cellulose ribbon, however, disorganized aggregates were

observed in the vicinity of 1 out of the 6 tomograms. At 20 minutes post-separation, cellulose

ribbons were observed adjacent to the cell in 9 out of 15 tomograms (64% versus 88% (n = 33)

at 300 minutes post-separation) (Fig. 4A). Out of these 9 cells harboring an adjacent cellulose

ribbon, 3 had it on the top or bottom of the cell and were excluded from the analysis for the

same reason explained above. Therefore, the analysis of the OM-ribbon interface was

conducted on the remaining 6. The cellulose ribbons observed at 20min post-separation

comprised only one cellulose sheet (n = 6) (Fig. 4B). Four out of these 6 tomograms (67%)

exhibited a "tight" configuration. The average OM-to-closest sheet distance of 14 ± 3 nm (n =

4) was not significantly different from the 300 minutes post-separation "tight" configuration

average OM-to-closest sheet distance (Fig. 4C-D). The two other tomograms bore ribbons in

the "loose" configuration, i.e. at an OM-to-closest sheet distance >40 nm with disorganized

aggregates in-between. These "loose" ribbons had an OM-to-closest sheet distance of 43 and

59 nm, respectively. The disorganized aggregates visible at 20 minutes post-separation

emanated perpendicularly from the OM to connect to the nascent cellulose sheet. They were

thinner than the ones observed at 300 minutes post-separation and rod-shaped (Fig. 4E-F, red

arrowheads). Average density profiles normal to the direction of the cylindrical-shaped

densities were traced to estimate their diameter (Fig. 4G). We again emphasize the inherent

overestimation of such measurements due to defocus. The average estimates on the two cells,

11 ± 2 nm (n = 12) and 6.5 ± 1 nm (n = 4), respectively (Fig. 4G), therefore establish upper

limits of the true diameter. These estimates are also less than the above-measured inter-sheet distances (Fig. 4H). Because elementary fibrils are thought to be between 1.5 and 6 nm in thickness[14,36,37], we interpret these structures as microfibrils composed of several elementary fibrils. The variability of the microfibril diameter measurements between cells (Fig. 4G, Cells #1 and #2) suggests these structures can contain a varying number of elementary fibrils more-or-less tightly packed together. This configuration is reminiscent of what was seen in previous studies of microfibrils coming out of clusters of pores[23,36] and likely represents an early stage of cellulose sheet formation that has been mechanically disturbed. Sheets at 20 minutes post-separation had an estimated width of 25 ± 8 nm (n = 6) (Fig. 4I), smaller than those at 300 minutes.

These results show that 1) the microfibrils emanating from the OM have roughly the same thickness as the cellulose sheet, 2) sheet width seems to increase over time and 3) the number of cellulose sheets comprising a ribbon increases over time.

A novel cytoplasmic structure is associated with cellulose production

We next examined the interior of *G. hansenii* cells during cellulose synthesis. These cells had extensive cytoplasmic vesicles in the center and at the periphery of the cell (Supplemental figure 2). The most notable feature we observed was another ribbon-like structure closely associated with the inner membrane (24 ± 4 nm from it; n = 19, for an example peak-to-peak measurement see Fig. 3D) and several hundred nanometers in length (Fig. 5A, purple arrows). We found it in 90% of cells with a cellulose ribbon (n = 29), always on the same side as, and underlying, the nascent cellulose sheet (Fig. 5B-C, supplemental video 2). This cytoplasmic structure is not a tube but rather a stack of sheet-like structures, 47 ± 23 nm wide (n = 10), parallel to the inner membrane and spaced (peak-to-peak) by 15 ± 5 nm (n = 7) (Fig. 5D-F). We refer to it here as the "cortical belt".

<u>Structural hallmarks of crystalline cellulose synthesis are also present in intact biofilms</u>

It is possible that separating bacteria from the cellulose mat for whole cell cryo-ET imaging

could have altered structures associated with cellulose synthesis. We therefore imaged *G.*

hansenii cells *in situ* in young cellulose biofilms grown on gold grids. We imaged biofilms after

3 or 6 hours before plunge-freezing in hope of visualizing any change in the ordering of the

fibers or the aspect of the cells over the course of biofilm growth. To access cells within the 5-

to 10-micron thick biofilm, we used cryo-FIB milling to generate thin (~200 nm) lamellae

suitable for imaging by cryo-ET (Fig. 6A-C). In a total of 19 analyzed tomograms (9 and 10

tomograms for 6h and 3h biofilms, respectively, Table 1), we observed fields of living and dead

bacteria encased in a matrix of bundled cellulose ribbons at both time points (Fig. 6D-E and

supplemental video 3). Overview tomograms (low magnification with low total dose) and high-

resolution composite images of the lamellae allowed extraction of positional information of the

cells in relation to the biofilm. There were 0.10 ± 0.02 and 0.27 ± 0.04 cells/um^2 and 15% and

28% of the volume of the lamellae was occupied by cells at 3 and 6h time points, respectively

(Fig. 6F) (n = 6 and 4 lamellae, respectively). This approximate 2-fold increase in cell density

from a 3-hour to a 6-hour biofilm suggests that cell division is occurring during biofilm growth.

As dead cells can be easily identified by their appearance (Fig. 6D, red asterisks), the live-to-

dead cell ratio was calculated at 0.9 ± 0.1 in both 3- and 6-hour biofilms, revealing no increase

in the proportion of dead cells between these two timepoints (Fig. 6G). Because lamellae give

access to the native organization and layering of the cells within the biofilm, the depth of

dead/living cells within the biofilm was assessed by measuring their distance from the leading

edge of the lamella (see methods). No trend between cell depth within the biofilm and state of

the cells was detected (Fig. 6H).

In all 19 tomograms (combining 3h and 6h lamellae), we observed numerous cellulose ribbons

surrounding the cells (Fig. 7A, yellow arrowheads). In 5 out of the 19 tomograms (26%), a

cellulose ribbon was closely appended to the cell's OM, as we previously had seen in separated

cells (Fig. 7B-C, dark-lined yellow arrowhead). Among those 5 tomograms, 4 showed a cortical

belt adjacent to the cellulose ribbon (Fig. 7B-D and supplemental video 3). The OM-to-

cellulose ribbon distance (19.2 ± 8 nm, n = 4) and inner membrane to cortical belt distance (22

± 2 nm, n = 4) were very similar to those measured before in separated cells. In 5 out of the 10

tomograms in 3h biofilm lamellae, disorganized cellulose aggregates were observed connected

to well-ordered ribbons just as in the separated cells, whereas this was never observed in the 6h

biofilms. This suggests that crystallization is disrupted more often in early biofilm growth (Fig.

7E-G, orange dashed lining). Because *Gluconacetobacter* cells are thick, electron transmittance

in the central region of the cytoplasm is very low when imaging whole cells, making it difficult

to visualize this area. Reducing sample thickness to approximately 200 nm by cryo-FIB-milling

allowed us to observe these central regions with greater contrast and visualize the extensive

vesicle network deep inside the cell (Fig. 7E, white arrowheads). Overall, the morphology of

cells and cellulose structures in a biofilm was the same as in isolated cells.

<u>The cortical belt is specific to bacterial species that produce crystalline cellulose ribbons</u>

To see whether the cortical belt is specific to *G. hansenii*, we imaged another strain of

Gluconacetobacter, *G. xylinus*, by cryo-ET at 300 minutes post-separation. Four out of 8 cells

(50%) exhibited an extracellular cellulose ribbon along the cells' long axis (Supplemental

figure 3A). The cellulose ribbons observed had 2 sheets of cellulose, with an estimated average

width of 27 ± 16 nm (n = 5). All four cells also possessed a cortical belt (Supplemental figure

3A-B, purple arrows), with similar dimensions to those in *G. hansenii.* The average distance

from the cortical belt to the inner membrane was 24 ± 4 nm (n = 4). In one instance, the cortical

belt also contained three stacked layers spaced (peak-to-peak) by 9 nm (Supplemental figure 3C). Aside from *Gluconacetobacter*, other bacterial species produce different types of cellulose. For instance, *Escherichia coli* 1094 can make amorphous cellulose[40] and *Agrobacterium tumefaciens* makes crystalline cellulose microfibrils during plant infection[41]. Neither of these species are known to make cellulose ribbons. We asked whether structures similar to the cortical belt observed in *Gluconacetobacter* were present in these species. Our lab had previously imaged *A. tumefaciens* for other studies, and therefore cryo-tomograms of *A. tumefaciens* were already available. We confirmed by confocal microscopy that *A. tumefaciens* produces cellulose in the same growth conditions as had been used for the earlier experiments (Fig. 8A), and then screened the available tomograms for the presence of cellulose. As the purpose of the previous studies had not been cellulose synthesis observation, relatively few (65 out of 1,854 tomograms) showed distinct cellulose fibers in the vicinity of the cells (Fig. 8B-C, yellow arrowheads, supplemental video 4). These fibers did not adopt any preferential orientation and ran in all directions around the cell. They also had a decreased width (14 ± 5 nm, n = 52 fibers measured in 5 tomograms) compared to *G. hansenii* cellulose sheets, confirming that *A. tumefaciens* does not elaborate wide cellulose sheets nor ribbons but rather simpler structures of crystalline cellulose, presumably bundles of microfibrils. In the 65 cellulose-producing cells, we never observed a cortical belt structure. Two notable features were however observed: 1) a polar outer-membrane flattening in 28 cells with a thickening of the OM (43% out of the 65 cells presenting cellulose, Fig. 8B, cyan arrow) and 2) polar amorphous aggregates in 24 cells (37% out of the 65 cells presenting cellulose), (Fig. 8B, orange dashed lining). 19 cells exhibited all three described features, the polar flattening, the amorphous aggregates and the cellulose fibers. We suspect these polar amorphous aggregates to be the unipolar polysaccharides (UPP) described in previous work and shown to allow the attachment of *A. tumefaciens* to biotic and abiotic surfaces in the early stages of biofilm

formation[42]. The very close proximity of the putative UPP to the polar flattening suggests the

latter could hold the UPP-secreting complexes.

We confirmed that *Escherichia coli* 1094 grown in minimal medium produces cellulose (Fig.

8D). The cells aggregated, making it difficult to image single cells by cryo-ET, so instead we

FIB milled through bacterial mats, producing approximately 200 nm-thick lamellae. To identify

cellulose structures, we also imaged lamellae from cultures grown in minimal medium

supplemented with cellulase. In 3 of the 5 tomograms of untreated cells, we observed

amorphous fibrous material (Fig. 8E, orange asterisk), that was not visible in 2 tomograms of a

cellulase-treated culture (Fig. 8F). None of the cells imaged in either condition contained a

cortical belt (n = 13 untreated and 5 cellulase-treated cells), suggesting that it is unique to

bacteria producing higher-order crystalline cellulose structures, *i.e.* sheets.

Discussion

Here we characterized bacterial cellulose synthesis in two *Gluconacetobacter* spp. and

compared it to two other species by cryo-ET. We identified a novel cytoplasmic structure

associated with the production of cellulose I ribbons in *Gluconacetobacter* spp. We also

performed cryo-FIB milling followed by cryo-ET on a native biofilm.

<u>Cryo-ET confirms the need of a tight interaction between the nascent sheet and the OM</u>

The cell-directed hierarchical model proposes linearly arranged 3.5-nm diameter pores on the

surface of the cell[37], each extruding an elementary fibril[23,29]. The arrangement of these pores in

lines allows the crystallization of the elementary fibrils upon secretion and integration into a

cellulose sheet parallel to the long axis of the cell[7,43,44]. Our results agree with this model.

Indeed, we observed that when the gap between the nascent sheet and the OM exceeds

approximately 40 nm, disorganized aggregates occur (Fig. 2). Along with previous work that

observed similar events[23], we hypothesize that these aggregates are microfibrils failing to

integrate into an ordered sheet. Furthermore, it has been shown that the addition of compounds

which bind directly to cellulose drastically alters the assembly of the sheets and leads to the

formation of similar aggregates[23,44]. It appears as though preventing the nascent microfibrils

from interacting with each other upon secretion prevents them from forming one organized

sheet. Conversely, a confined spacing between the nascent sheet and the OM promotes proper

crystallization of the nascent microfibrils. This proximity could be maintained either by a

previously synthesized sheet preventing the nascent one from separating too far from the OM,

or by specialized cellulose binding enzymes situated in the outer-leaflet of the OM, such as

CmcAx, which has the ability to bind cellulose[45].

Cryo-ET sheds light on the buildup of a microfibril

Many values have been reported for the elementary fibrils' dimension, mainly through direct

observation by negative staining electron microscopy[14,29,36]. The most favored hypothesis is an

approximately 1.5-nm thick elementary fibril (thoroughly discussed in [23]). Very recently, the

characterization of the structure of the BcsC subunit (the OM pore) strengthens this hypothesis,

since the pore was seen to have an internal diameter of 1.5 nm at maximum[19]. Therefore, the

simplest hypothesis is that each extrusion pore comprises a single BcsC sub-unit secreting an

elementary fibril not more than 1.5 nm thick.

While negative staining has provided high-resolution views of cellulose ribbons [23,36], observing

them in a frozen-hydrated state enables more accurate measurements of their dimensions and

observation of their interaction with the OM. This is particularly important for extracellular

polysaccharides, which have been shown to collapse and undergo drastic conformational

changes upon dehydration, staining and embedding[46].

We were able to image in two tomograms, microfibrils extruded perpendicularly to the OM and

integrating to form a thin parallel sheet (Fig. 4E-F). A possible interpretation of why these

events are rare is that they result from an accidental mechanical separation of the nascent sheet

from the OM, revealing intermediate forms of cellulose bundling such as thin microfibrils. As

explained earlier, precise measurement of the thickness of densities is difficult in cryo-ET since

it is influenced by the defocus applied during imaging (causing overestimation of the true

thickness). Despite this uncertainty, our measurements are done in a near native state. We

estimated these microfibrils to be less than 11 nm in diameter (Fig. 4G-H). Previous work

measured microfibril thicknesses from 3 to 12 nm in cellulose sheets splayed apart by cellulase

treatments[47]. If we assume an elementary fibril is 1.5 nm in diameter since this is the maximal

opening of a BcsC pore[19], a 10 nm cylindrical microfibril would comprise approximately 35

elementary fibrils. This would require a cluster of 35 extrusion pores. Previous reports have

stated the extrusion pores cluster in bunches from 2 to 4 pores[7,29], substantially less than our

calculation. Our data therefore suggest that there is more than one BcsC subunit per extrusion

pore. For example, if each 3.5 nm diameter extrusion pore[37] held 4 BcsC subunits, a cluster of

9 extrusion pores could produce a 9 nm diameter microfibril (Fig. 9). In this particular case,

each extrusion pore holding multiple BcsC subunits would produce a crystalline aggregate of

elementary fibrils which would pack with its neighboring aggregates to form a microfibril.

<u>Cryo-ET sheds light on the assembly of a cellulose Iα sheet</u>

We found that ribbons were stacks of sheets that likely interact loosely with one another since

the inter-sheet distance varied from 7- to 31-nm. This loose stacking corroborates previous

observations[30]. Previous measurements done by negative staining had estimated cellulose sheet

width to range from 40 to 600 nm[23,36,47], wider than our measurements ranging from 11 to 69

nm (Fig. 4I). These variations have been attributed to the cell strain, growth conditions and

intercellular variation[23,36,37]. We found that the thickness of cellulose sheets is similar to the

diameter of the microfibrils. Therefore, our data suggest that microfibrils lie down in rows to

create the width of the sheet. This was also suggested in[7].

While the number of sheets produced by a single cell increased with time, the main dimension

of growth appears to be ribbon length, as suggested by previous work and our fluorescence data

showing cellulose ribbons several cell lengths long (Fig. 1A-B)[24]. Wider sheets occur in later

time points (Fig. 4I), suggesting that sheet width also grows with time. However, in the current

model, sheet width is correlated with the number of extrusion pores, hence to cell length[7,37]. It

is possible that at 300min post-separation, cells are longer and possess more extrusion pores,

therefore producing wider sheets.

<u>Cryo-ET on *G. hansenii* cells allowed the visualization of a novel cytoskeletal element, the</u>

<u>cortical belt</u>

Negative stain, cryo-fracture and immuno-EM studies have shown that cellulose extrusion

pores in *Gluconacetobacter* align in a line on one side of the cell [23,37,48], but what causes this

alignment is unknown. Here, we identify a novel cytoplasmic structure in two species of

Gluconacetobacter that spatially correlates with the nascent cellulose ribbon (Fig. 1C-E and

Fig. 5). This structure, which we term the cortical belt, is found at a fixed distance from the

inner membrane (24 ± 4 nm) and remains intact upon cellulase treatment (Supplemental figure

1F, purple arrow), suggesting that it is stable even in the absence of the cellulose ribbon.

We observed the cortical belt in both *Gluconacetobacter* spp. imaged but not in other bacteria

that produce less-ordered forms of cellulose, including *Escherichia coli* 1094, which

synthesizes amorphous cellulose[40], and *Agrobacterium tumefaciens,* which synthesizes

cellulose I microfibrils[49] (Fig. 8). This suggests that the cortical belt functions in the formation

of cellulose I ribbons. BcsD and CcpAx, as well as two cell wall-related enzymes, have been

shown to be involved in the crystallization process of the ribbons [24,26,50]. It is possible that the

cortical belt interacts with one or more of these components to guide the positioning of the BCS

complexes, but the interaction may also be indirect. If the cortical belt is responsible for

scaffolding the BCS complexes, it represents a novel prokaryotic cytoskeletal element, i.e. "a

cytoplasmic protein filament and its associated superstructures that move or scaffold material

within the cell"[51]. Other bacterial cytoskeletal elements have been observed to form belt-like

structures, including bactofilins[52], or to stack, like the CTP synthase[53], although with different

dimensions. *G. hansenii* has homologs of both bactofilin and CTP synthase, but neither shows

co-evolution with the *bcs* gene cluster. We hope that future work will identify the component(s)

that form the cortical belt, shedding more light on the molecular processes involved in the

organization and clustering of the BCS complexes in *G. hansenii*.

The cortical belt reveals another similarity between cellulose synthesis in *Gluconacetobacter*

and land plants.

Historically, the first plant cellulose synthase genes were identified by cDNA homology with

the *G. xylinum* acsA (bcsA) gene[54]. Later on, phylogenetic studies highlighted an early

divergence between cyanobacterial and plant cellulose synthases[55,56]. A large number of

cellulose I synthesizing organisms have in common that the synthase complexes arrange in

specific patterns, determining the final architecture of the cellulose structures[7]. A simple row

in systems like *Gluconacetobacter* spp. or certain charophytes and chlorophytes[57] and

hexameric rosette structures called Cellulose Synthase Complexes (CSC) in land plants. In

both, the extrusion of a crystalline form of cellulose exerts a force believed to be able to propel

the CSCs in plants[58,59] and the whole cell in *Gluconacetobacter*[24,36]. Our work uncovers an

additional similarity, the involvement of a cytoskeletal element, the cortical belt, to guide the

synthase complexes. In land plants CSCs have been shown to interact indirectly with underlying

cortical microtubules, mediating trans-membrane cross-talk[60–62], guiding and regulating CSC

velocity[63–65]. While CSCs were shown to be motile in land plants, they are believed to be static

in *Gluconacetobacter*[23], perhaps held in place by the cortical belt, in order to transfer the

propelling force to the whole cell.

<u>Insights from FIB-milling native biofilms</u>

Cryo-FIB milling through native biofilms offers the possibility of observing bacteria in the

context of their original biofilm environment and retrieving high resolution morphological and

positional information about the cells relative to one another and relative to the biofilm layers.

Visualization of the density and organization of the extracellular matrix and its interaction with

the cells is also rendered possible by cryo-FIB milling. This is especially important since in

nature most bacterial species are found in complex interacting communities, in the form of

homogeneous or heterogeneous communities that organize in biofilms[10].

Milling the *Gluconacetobacter* biofilms to 200 nm revealed numerous cytosolic vesicles of

variable shapes and sizes. The cortical belt was also visible, as in the isolated cells. The

cellulose ribbons aligned with each other to form larger arrays 2-3 µm wide (Fig. 6D, yellow

arrowheads and supplemental video 3), showing the propensity of these structures to interact

with each other. This propensity was previously characterized by live imaging of the cellulose

biosynthesis and crystallization process in *Gluconacetobacter,* which showed that the bacterial

cells preferentially follow already established tracks, *i.e.* previously synthesized cellulose

ribbons[24]. The occurrence of disorganized cellulose clusters in biofilms grown for 3h but not

6h, suggests that such aggregates are either 1) digested by enzymes, likely CmcAx, reported to

have an endoglucanase capable of digesting amorphous cellulose[66] and to be present on the

surface of *G. hansenii* or released in the environment[25,45] or 2) diluted by a gradual increase in

well-ordered ribbons over time.

Cell death in biofilms, with the fraction of dead cells measured at 10% in our biofilms, is a

well-known phenomenon[10], caused by programmed cell death mechanisms, cannibalistic

behaviors such as already described in *B. subtilis*[67] or nutrient/oxygen depletion[68,69]. We did

not observe a preferential location of dead cells at the bottom of the biofilm, ruling out anoxic

conditions being the primary cause of cell death. This could be because the thickness of the

biofilm, between 1.5- and 3-um according to the cell depth distribution (Fig. 6H), is too small

to have a significant oxygen gradient, as suggested by studies that measured total anoxia being

reached generally between 70- and 80-um depth[69–71]. Processing thicker biofilms in the range

of tens of microns would allow visualization of the effects of nutrient/oxygen gradients on cell

distribution and physiology. For now, plunge freezing such as performed in this study can only

properly vitrify samples less than ~10 microns thick[72]. Moreover, milling thicknesses above 8-

10 microns becomes labor intensive and technically difficult. A possible course of action for

further studies would be to perform high-pressure freezing on thicker biofilms and then produce

thin sections either by cryosectioning, hybrid cryosectioning/FIB-milling methods such as

described in[73–75] or following a cryo-lift out procedure[75].

Conflicts of interest

The authors declare that there are no conflicts of interest.

Acknowledgments

This work was supported by NIH grant R35-GM122588 to GJJ, the Howard Hughes Medical

Institute (HHMI) and the Center for Environmental Microbial Interactions (CEMI) pilot grant

program. Cryo-electron microscopy was performed in the Beckman Institute Resource Center

for Transmission Electron Microscopy at Caltech. We thank Jean Marc Ghigo for kindly

providing us the E. Coli 1094 strain. Special acknowledgments to Catherine Oikonomou for all

the help and scientific advice given during this study and also to Candace Haigler for sharing

her thoughts and her precious experience on the not so common *Gluconacetobacter* spp.

Methods

<u>Cell culture</u>

Gluconacetobacter hansenii (ATCC 23769) was cultured as previously described [35] in SH

medium: 2% glucose, 0.5% bactopeptone, 0.5% yeast extract, pH 6. For solid medium, 2.5%

bacto-agar was added. Cells were separated from the cellulose biofilm by mechanical disruption

as previously described[36]. Briefly, the bacterial cellulose biofilm developing at the air-media

interface was picked up with a single-use sterile inoculating loop and transferred to fresh

medium, where it was vigorously shaken and then removed. In preparation for freezing, cells

were pelleted by centrifugation for 10 minutes at 2500rcf at 20C and resuspended in 0.5mL of

SH media. The culture was incubated for the desired length of time at 30ºC without shaking

before plunge freezing. For cellulose digestion, 0.2g/L cellulase (Worthington, purified exo-

and endo-glucanases, #LS002598) was added.

Gluconacetobacter xylinus (ATCC 700178/BPR2001) was cultured as described above in

Fructose–Peptone–Yeast Extract (FPY) media: 2% fructose, 1% bactopeptone, 0.5% yeast

extract and 0.25% K_2HPO_4.

Escherichia coli 1094 was cultured in Lysogeny Broth (LB) and induced for cellulose

production in minimal medium: 0.2% $(NH_4)_2SO_4$, 1.4% KH_2PO_4, 0.1% $MgSO_4$, 0.5%

$FeSO_4.7H_2O$, 0.4% glucose, 0.01% thiamine, pH 7. A saturated overnight LB culture was

diluted 1:50 into 3mL of minimal medium with or without 0.2g/L cellulase (Worthington,

purified exo- and endo-glucanases, #LS002598). Cultures were incubated at 37ºC with shaking at 220rpm. When the medium transitioned from turbid to clear and white flakes appeared (cellulose and bacteria), the induction of cellulose synthesis is considered successful.

Agrobacterium tumefaciens was cultured as described in previous work[76]. Briefly, *A. tumefaciens* C58 was cultivated in liquid AB medium (glucose 0.2%, NH4Cl 18.7mM, MgSO4 2.5uM, KCl 2mM, CaCl2 0.07mM, FeSO4 0.01mM, K2HPO4 8.4mM, NaH2PO4.7H2O 4.16mM, pH 7) at 30C overnight. Induction was done by pipetting 100uL of overnight culture and spreading onto AB induction plates (glucose 0.2%, NH4Cl 18.7mM, MgSO4 2.5uM, KCl 2mM, CaCl2 0.07mM, FeSO4 0.01mM, K2HPO4 8.4mM, NaH2PO4.7H2O 4.16mM, Bactagar 1.7%, Acetosyringone 100uM, pH 5.8). Plates were then incubated for 3 days at 20C. Cells were resolubilized by scraping a small amount from the plate with an inoculation loop and resuspending it in 100uL of liquid induction AB medium.

The following strains are the ones included in the tomogram analysis: **NT1** is a C58 strain without the pTiC58 (tumor inducing) plasmid; **A139** strain is NT1REB(pJK270) + pJZ041. NT1REB is a "bald strain", no flagellin mutant, derived from NT1. The pJK270 is pTiC58 with the transposed NPTII gene for kanamycin resistance. The pJZ041 plasmid carries a GFP tagged VirB8 gene, a component of the T4SS (Aguilar et al. 2011); **JX148** strain is a C58 derived mutant of the rem gene. The strain is non motile; **AD348** is a GV3101(pMP90) strain with its whole VirB system deleted. GV3101 is a pTiC58 free, rifampicin resistant C58 strain and pMP90 is a helper pTiC58 without the T-DNA; **AD1484** is a AD348 variant, transformed with pAD2079 containing the whole VirB system.

  Confocal microscopy

  Cellulose was stained with Calcofluor-white (Sigma-Aldrich, #18909) at a concentration of

  0.001% and cell membranes were stained with MitoTracker Deep Red FM (Thermo-Fisher,

  #M22426) at a concentration of 0.5ug/uL. Stack acquisition was done on a Zeiss LSM880 Airy

  Scan microscope. Airy scan acquisitions were performed in super-resolution mode with Z-step

  set at the optimal optical sectioning. The Mito-Tracker Deep Red FM channel was set as the

  following: excitation at 633 nm, use of the 488/561/633 main beam splitter and a band-pass

  570-620 + long-pass 645 filter. The Calcofluor White channel was set as the following:

  excitation at 405 nm, use of the 405 main beam splitter and a band-pass 420-480 + band-pass

  495-550 filter. Airy scan processing was performed on the fly by the in-built algorithm of Zeiss

  Black.

  Sample preparation for cryo-EM

  For isolated cells, Quantifoil Cu R2/2 Finder grids (*Quantifoil Micro Tools GmbH*) were glow-

  discharged at 15mA for 1min. The grids were pre-incubated with fiducial marker solution

  prepared as follows: 50μL of 10nm colloidal gold (*Ted Pella, Inc*) mixed with 50uL of 5%

  BSA, vortexed 1 min and centrifuged at 15,000rcf for 15 min, supernatant discarded, and pellet

  resuspended in 40μL of PBS buffer. 3μL were deposited on each grid, left for 1 minute then

  back-blotted with Whatman paper. Cells were plunge frozen with a Vitrobot Mark IV (*Thermo*

  Fisher Scientific) with 100% humidity at 30ºC and back-blotted for 3 to 5s.

  For native biofilms, Quantifoil gold R2/2 Finder grids were placed in 35mm glass bottom petri

  dishes (*MatTek Corporation* #P35G-1.0-2.0C) containing 1mL of SH media inoculated with a

  2-day old biofilm. The dishes were sealed with Micropore tape (*3M*) and incubated without

  shaking at 30ºC for 3 to 6 hours. Plunge-freezing was done at 22C, 50% humidity, either with

manual blotting on both sides of the grids (first back-blotted then front-blotted) or using the

automatic blotting function of the Vitrobot with a blot time of 5-6s, blot force of 15 and drain

time of 2s.

For *E. coli* 1094, after 4 hours of incubation in minimal media, the medium should turn from

turbid to clear with white flakes. OD_{600} of the cultures was monitored using the culture (always

turbid) where cellulose induction was performed in the presence of cellulase to keep the cells

from aggregating. It was then used as a reference to concentrate the cells to high OD_{600} (10-

20), in order to form bacterial mats on the EM grids, for control and cellulase conditions.

Plunge-frozen was done at 20C, 100%, either with manual back-blotting for 5-7s and a drain

time of 1s or using the automatic blotting function of the Vitrobot with a wait time of 10s, blot

time of 5-6s, blot force of 3 and drain time of 1s.

<u>FIB milling</u>

Grids were clipped in Autogrid holders (*Thermo Fisher*) machined with a notch to allow FIB

milling closer to the edge of the grid. Autogrids were placed in a custom-built shuttle and

inserted into a Versa 3D dual-beam FIB/SEM microscope with FEG (*FEI*) equipped with a

PP3000T cryo-transfer apparatus (*Quorum Technologies*). They were maintained at -175ºC at

all times by a custom-built cryo-stage[77]. To reduce sample charging and protect the sample

from curtaining during milling, the grids were sputter-coated with platinum at 15 mA for 60

seconds. Thin lamellae were generated with the Ga^+ ion beam at 30 kV at angles ranging from

10 to 17 degrees. Rough milling was done at high currents, ranging from 0.3 nA to 100 pA until

the lamellae measured 1 micron in thickness under the FIB view. Current was then

progressively brought down to 10 pA for the final milling steps until the measured thickness

was between 100-200 nm. Final polishing of the back end of the lamella is also done at 10pA

where the sample is tilted +0.5 to 1° to homogenize the lamella thickness. During the whole

procedure, imaging with the SEM beam was done at 5 kV and 13 pA.

<u>Electron cryo-tomography</u>

Tomography of whole cells and FIB-milled lamellae was performed on either a Titan Krios or

Tecnai G2 Polara transmission electron microscope (*Thermo Fisher*) equipped with 300 kV

field emission gun, energy filter (*Gatan*) and K2 or K3 Summit direct electron detector (*Gatan*).

The Krios is equipped with a Volta phase plate (*Thermo Fisher*) [78]. Tilt-series acquisition was

done with SerialEM [79] with a 2-3˚ tilt increment for a total range of ±60˚ or ±50˚, defocus of -

4, -6 or -8 μm, and total dose up to 180 e$^-$/Å^2. Volta phase plate imaging was performed in

Figures 1, 2, 5 and 7A-B with a defocus of -2μm and a measured phase shift of 0.5 π/rad before

tilt series acquisitions.

Low magnification tomography on the biofilm lamellae was performed at 6500 magnification

(14 Å^2 pixel size) with a -10 or -15 μm defocus and a total dose between 5 and 10 e$^-$/Å^2.

Tomography of FIB-milled lamellae was done exclusively on the Titan Krios. Because samples

were thinner, the total dose was limited to ~80 e$^-$/Å^2.

<u>Data processing</u>

Tomograms were reconstructed using the IMOD software (http://bio3d.colorado.edu/imod/) [80].

Alignment was done on 1k x 1k binned tilt-series with fiducial-based alignment. Aligned stacks

were low-pass filtered (0.35, σ = 0.05) to eliminate high-frequency noise. Weighted back

projection reconstruction was performed and the "SIRT-like filter" was used with 20 iterations.

Segmentation was also done using IMOD and drawing tools developed by Andrew Noske

(http://www.andrewnoske.com/student/imod.php). To better distinguish features during the

segmentation steps, tomograms were filtered with the 3D non-linear anisotropic diffusion filter

in IMOD. The cell contours and cortical belt were segmented manually on a Cintiq 21uX tablet

(*Wacom*) and cellulose was segmented using a semi-automated thresholded method. 1) A

denoising Non-linear Anisotropic Diffusion filter was applied (included in the *etomo* package,

http://bio3d.colorado.edu/imod/) on the tomogram; 2) Precise boundary models are drawn

around the structures to be thresholded; 3) Thresholding segmentation is performed with

3Dmod using the isosurface function and the previously drawn contours are used as a mask.

When the contours are precisely following the contours, this technic allows to raise the

isosurface threshold without picking up background noise.

Measurements for all distances between elements (cellulose sheet – outer-membrane, width of

the cellulose ribbon, cortical belt – inner-membrane) were taken by generating normalized

density profile plots and measuring the distances between the density peaks of the

corresponding sub-cellular features (Fig. 3). This was automated with a custom script, *sideview-*

profile-average, written by Davi Ortega (https://www.npmjs.com/package/sideview-profile-

average).

Estimation of the cell depth in the native biofilm lamellae was calculated as follows: 1) using

the two parallel walls of the milled trench, a perpendicular line is traced at the leading edge of

the lamella (where the platinum meets the frozen material); 2) Lines are drawn from the center

of the cells to the leading edge perpendicular line (Fig. 6H, red line in top view of lamella); 3)

The distance from the cell center to the limit of the platinum on the leading edge, which is the

surface of the sample, is measured. The real depth is then calculated using the following

equation: opposite side (real depth) = tan (*a*) x adjacent side (distance measured, d in Fig. 6H).

The angle *a* is the angle between the grid surface and the FIB gun during the milling process,

which can be accurately measured during reconstruction with 3dmod.

All statistics were performed with GraphPad Prism software (https://www.graphpad.com/scientific-software/prism/). All datasets were first analyzed for normality using the Shapiro-Wilk test and homoscedasticity (equal standard deviations). If dataset is normal, appropriate parametric tests were performed and if not, appropriate non-parametric tests were performed.

Figure 2E: n = 3 and 24 for the "loose" and "tight" configuration respectively. Two tailed P-value = 0.0008, Mann-Whitney test.

Figure 4A: n = 6, 15, 33 for 13-, 20- and 300-minutes, respectively.

Figure 4B: n = 6 and n = 21 tomograms for 20- and 300-minutes post-separation, respectively. Two tailed P-value < 0.0001, One sample Wilcoxon signed rank test against a theoretical value of 1 (number of sheets observed at 20-min post-separation).

Figure 4H: n = 12 and 4 microfibril thickness measurements performed on two separate tomograms (Cell #1 and #2, left side of the graph). N = 47 measurements for inter-sheet distances performed on 23 tomograms. ANOVA followed by Tukey's multiple comparison test was performed. Cell #1 vs Cell #2, Cell #1 vs 300-min inter-sheet distances and Cell #2 vs 300-min inter-sheet distances showed adjusted P-values of 0.073, 0.15 and 0.0015, respectively.

Figure 4I: n = 6 and 45 sheets measured at 20- and 300-minutes post-separation. Welch's t test (parametric t-test without equal SD assumption) showed a P-value of 0.23.

Figure 6F: n = 6 and 4 for biofilms let to grow for 3h and 6h, respectively. Unpaired T-test showed a two-tailed P-value of 0.0011.

Figure 6G: n = 6 and 4 for biofilms let to grow for 3h and 6h, respectively. Unpaired T-test showed a two-tailed P-value of 0.2720.

Figure 6H: n = 49, 46, 4 and 11 for live and dead cells in 3h and 6h biofilms, respectively.

Mann-Whitney tests were performed on live vs dead cells in 3h and 6h biofilms conditions,

showing two-tailed P-values of 0.82 and 0.54, respectively.

Bibliography

1.   Pauly, Markus & Keegstra, Kenneth. Cell-wall carbohydrates and their modification as

a resource for biofuels. *Plant Journal* vol. 54 559–568 (2008).

2.   Pauly, Markus & Keegstra, Kenneth. Plant cell wall polymers as precursors for

biofuels. *Current Opinion in Plant Biology* vol. 13 305–312 (2010).

3.   Hon, David N. S. Cellulose: a random walk along its historical path. *Cellulose* **1**, 1–25

(1994).

4.   Xu, Youjie, Zhang, Meng, Roozeboom, Kraig & Wang, Donghai. Integrated bioethanol

production to boost low-concentrated cellulosic ethanol without sacrificing ethanol

yield. *Bioresour. Technol.* **250**, 299–305 (2018).

5.   Gatenholm, Paul & Klemm, Dieter. Bacterial Nanocellulose as a Renewable Material

for Biomedical Applications. *MRS Bull.* **35**, 208–213 (2010).

6.   Cavalcante, Aline Ribeiro Teixeira, Lima, Rodrigo Pontes de, Souza, Veridiana Sales

Barbosa de, Pinto, Flávia Cristina Morone, Campos Júnior, Olavio, Silva, Jaiurte

Gomes Martins da, Albuquerque, Amanda Vasconcelos de & Aguiar, José Lamartine

de Andrade. Effects of bacterial cellulose gel on the anorectal resting pressures in rats

submitted to anal sphincter injury. *Heliyon* **4**, e01058 (2018).

7.   Brown Jr., R. M. The biosynthesis of cellulose. *J. Macromol. Sci. Part A Pure Appl.*

Chem. **33**, 1345–1373 (1996).

8.   McNamara, Joshua T., Morgan, Jacob L. W. & Zimmer, Jochen. A molecular

description of cellulose biosynthesis. *Annu. Rev. Biochem.* **84**, 895–921 (2015).

9.   Cosgrove, Daniel J. Growth of the plant cell wall. *Nat. Rev. Mol. Cell Biol.* **6**, 850–861
(2005).

10.   Flemming, Hans Curt, Wingender, Jost, Szewzyk, Ulrich, Steinberg, Peter, Rice, Scott
A. & Kjelleberg, Staffan. Biofilms: An emergent form of bacterial life. *Nature Reviews*
Microbiology vol. 14 563–575 (2016).

11.   Römling, Ute & Galperin, Michael Y. Bacterial cellulose biosynthesis: Diversity of
operons, subunits, products, and functions. *Trends in Microbiology* vol. 23 545–557
(2015).

12.   De Vos, Willem M. Microbial biofilms and the human intestinal microbiome. *npj*
Biofilms and Microbiomes vol. 1 15005 (2015).

13.   Costerton, J. W., Cheng, K. J., Geesey, G. G., Ladd, T. I., Nickel, J. C., Dasgupta, M.
& Marrie, T. J. Bacterial Biofilms in Nature and Disease. *Annu. Rev. Microbiol.* **41**,
435–464 (1987).

14.   Haigler, Candace Hope, Brown, R. Malcolm & Benziman, Moshe. The fine structure of
cellulose microfibrils. *Science* **119**, 80–82 (1980).

15.   Morgan, Jacob L. W., McNamara, Joshua T., Fischer, Michael, Rich, Jamie, Chen,
Hong Ming, Withers, Stephen G. & Zimmer, Jochen. Observing cellulose biosynthesis
and membrane translocation in crystallo. *Nature* **531**, 329–334 (2016).

16.   Du, Juan, Vepachedu, Venkata, Cho, Sung Hyun, Kumar, Manish & Nixon, B. Tracy.
Structure of the cellulose synthase complex of Gluconacetobacter hansenii at 23.4 Å
resolution. *PLoS One* **11**, e0155886 (2016).

17.   Hu, S. Q., Gao, Y. G., Tajima, Kenji, Sunagawa, Naoki, Zhou, Yong, Kawano, Shin,
Fujiwara, Takaaki, Yoda, Takanori, Shimura, Daisuke, Satoh, Yasuharu, Munekata,
Masanobu, Tanaka, Isao & Yao, Min. Structure of bacterial cellulose synthase subunit
D octamer with four inner passageways. *Proc. Natl. Acad. Sci.* **107**, 17957–17961

(2010).

18. Saxena, I. M., Kudlicka, K., Okuda, K. & Brown, R. M. Characterization of genes in

the cellulose-synthesizing operon (acs operon) of Acetobacter xylinum: Implications

for cellulose crystallization. *J. Bacteriol.* **176**, 5735–5752 (1994).

19. Acheson, Justin F., Derewenda, Zygmunt S. & Zimmer, Jochen. Architecture of the

Cellulose Synthase Outer Membrane Channel and Its Association with the Periplasmic

TPR Domain. *Structure* (2019).

20. Whitney, John C., Hay, Iain D., Li, Canhui, Eckford, Paul D. W., Robinson, Howard,

Amaya, Maria F., Wood, Lynn F., Ohman, Dennis E., Bear, Christine E., Rehm, Bernd

H. & Lynne Howell, P. Structural basis for alginate secretion across the bacterial outer

membrane. *Proc. Natl. Acad. Sci.* **108**, 13083–13088 (2011).

21. Rehman, Zahid U., Wang, Yajie, Moradali, M. Fata, Hay, Iain D. & Rehm, Bernd H.

A. Insights into the assembly of the alginate biosynthesis machinery in Pseudomonas

aeruginosa. *Appl. Environ. Microbiol.* **79**, 3264–3272 (2013).

22. Keiski, Carrie-Lynn, Harwich, Michael, Jain, Sumita, Neculai, Ana Mirela, Yip,

Patrick, Robinson, Howard, Whitney, John C., Riley, Laura, Burrows, Lori L., Ohman,

Dennis E. & Howell, P. Lynne. AlgK Is a TPR-Containing Protein and the Periplasmic

Component of a Novel Exopolysaccharide Secretin. *Structure* **18**, 265–273 (2010).

23. Haigler, Candace Hope. Alteration of cellulose assembly in Acetobacter xylinum by

fluorescenet brightening agents, direct dyes and cellulose derivatives. (University of

North Carolina, 1982).

24. Mehta, Kalpa, Pfeffer, Sarah & Brown, R. Malcolm. Characterization of an acsD

disruption mutant provides additional evidence for the hierarchical cell-directed self-

assembly of cellulose in Gluconacetobacter xylinus. *Cellulose* **22**, 119–137 (2015).

25. Nakai, Tomonori, Sugano, Yasushi, Shoda, Makoto, Sakakibara, Hitoshi, Oiwa,

Kazuhiro, Tuzi, Satoru, Imai, Tomoya, Sugiyama, Junji, Takeuchi, Miyuki, Yamauchi, Daisuke & Mineyukia, Yoshinobu. Formation of highly twisted ribbons in a carboxymethylcellulase gene-disrupted strain of a cellulose-producing bacterium. *J. Bacteriol.* **195**, 958–964 (2013).

26. Sunagawa, Naoki, Fujiwara, Takaaki, Yoda, Takanori, Kawano, Shin, Satoh, Yasuharu, Yao, Min, Tajima, Kenji & Dairi, Tohru. Cellulose complementing factor (Ccp) is a new member of the cellulose synthase complex (terminal complex) in Acetobacter xylinum. *J. Biosci. Bioeng.* **115**, 607–612 (2013).

27. Deng, Ying, Nagachar, Nivedita, Xiao, Chaowen, Tien, Ming & Kao, Teh Hui. Identification and characterization of non-cellulose-producing mutants of Gluconacetobacter hansenii generated by Tn5 transposon mutagenesis. *J. Bacteriol.* **195**, 5072–5083 (2013).

28. Cousins, Susan K. & Brown, R. Malcolm. Cellulose I microfibril assembly: computational molecular mechanics energy analysis favours bonding by van der Waals forces as the initial step in crystallization. *Polymer* **36**, 3885–3888 (1995).

29. Haigler, Candace H. & Benziman, Moshe. Biogenesis of Cellulose I Microfibrils Occurs by Cell-Directed Self-Assembly in Acetobacter xylinum. in *Cellulose and Other Natural Polymer Systems* 273–297 (Springer US, 1982).

30. Cousins, Susan K. & Brown, R. Malcolm. Photoisomerization of a dye-altered β-1,4 glucan sheet induces the crystallization of a cellulose-composite. *Polymer* **38**, 903–912 (1997).

31. Saxena, I. M. & Brown, R. M. Identification of a second cellulose synthase gene (acsAII) in Acetobacter xylinum. *J. Bacteriol.* **177**, 5276–83 (1995).

32. Florea, Michael, Reeve, Benjamin, Abbott, James, Freemont, Paul S. & Ellis, Tom. Genome sequence and plasmid transformation of the model high-yield bacterial

cellulose producer Gluconacetobacter hansenii ATCC 53582. *Sci. Rep.* **6**, 23635

(2016).

33.   Toyosaki, Hiroshi, Kojima, Yukiko, Tsuchida, Takayasu, Hoshino, Ken-Ichiro,

Yamada, Yuzo & Yoshinaga, Fumihiro. The characterization of an acetic acid

bacterium useful for producingbacterial cellulose in agitation cultures: The proposal of

Acetobacter xylinum subsp. sucrofermentans subsp. nov. *J. Gen. Appl. Microbiol.* **41**,

307–314 (1995).

34.   Park, Joong Kon, Jung, Jae Yong & Park, Youn Hee. Cellulose production by

Gluconacetobacter hansenii in a medium containing ethanol. *Biotechnol. Lett.* **25**,

2055–2059 (2003).

35.   Schramm, M. & Hestrin, S. Factors affecting Production of Cellulose at the Air/ Liquid

Interface of a Culture of Acetobacter xylinum. *J. Gen. Microbiol.* **11**, 123–129 (1954).

36.   Brown, R. M., Willison, J. H., Richardson, C. L. & Richardson, C. L. Cellulose

biosynthesis in Acetobacter xylinum: visualization of the site of synthesis and direct

measurement of the in vivo process. *Proc. Natl. Acad. Sci. U. S. A.* **73**, 4565–9 (1976).

37.   Zaar, K. Visualization of pores (export sites) correlated with cellulose production in the

envelope of the gram-negative bacterium Acetobacter xylinum. *J. Cell Biol.* **80**, 773–

777 (1979).

38.   Hawkes, Peter W. The electron microscope as a structure projector. in *Electron*

Tomography: Methods for Three-Dimensional Visualization of Structures in the Cell

vol. 9780387690087 83–111 (Springer New York, 2006).

39.   Radermacher, Michael. Weighted back-projection methods. in *Electron Tomography:*

Methods for Three-Dimensional Visualization of Structures in the Cell vol.

9780387690087 245–273 (Springer New York, 2006).

40.   Le Quéré, Benjamin & Ghigo, Jean Marc. BcsQ is an essential component of the

Escherichia coli cellulose biosynthesis apparatus that localizes at the bacterial cell pole.

Mol. Microbiol. **72**, 724–740 (2009).

41.   Matthysse, A. G., Holmes, K. V & Gurlitz, R. H. G. Elaboration of cellulose fibrils by

Agrobacterium tumefaciens during attachment to carrot cells. *J. Bacteriol.* **145**, 583–

595 (1981).

42.   Xu, Jing, Kim, Jinwoo, Koestler, Benjamin J., Choi, Jeong Hyeon, Waters, Christopher

M. & Fuqua, Clay. Genetic analysis of agrobacterium tumefaciens unipolar

polysaccharide production reveals complex integrated control of the motile-to-sessile

switch. *Mol. Microbiol.* **89**, 929–948 (2013).

43.   Ross, Peter, Mayer, Raphael, Benziman, A. N. D. Moshe & Benziman, M. Cellulose

Biosynthesis and Function in Bacteria. *Microbiology* **55**, 35–58 (1991).

44.   Benziman, Moshe, Haigler, Candace H., Brown, R. Malcolm, White, Alan R. &

Cooper, Kay M. Cellulose biogenesis: Polymerization and crystallization are coupled

processes in Acetobacter xylinum. *Proc. Natl. Acad. Sci. U. S. A.* **77**, 6678–6682

(1980).

45.   Yasutake, Yoshiaki, Kawano, Shin, Tajima, Kenji, Yao, Min, Satoh, Yasuharu,

Munekata, Masanobu & Tanaka, Isao. Structural characterization of the Acetobacter

xylinum endo-β-1,4-glucanase CMCax required for cellulose biosynthesis. *Proteins*

Struct. Funct. Bioinforma. **64**, 1069–1077 (2006).

46.   Dohnalkova, Alice C., Marshall, Matthew J., Arey, Bruce W., Williams, Kenneth H.,

Buck, Edgar C. & Fredrickson, James K. Imaging hydrated microbial extracellular

polymers: Comparative analysis by electron microscopy. *Appl. Environ. Microbiol.* **77**,

1254–1262 (2011).

47.   White, Alan R. & Brown, R. M. Enzymatic hydrolysis of cellulose: Visual

characterization of the process. *Proc. Natl. Acad. Sci. U. S. A.* **78**, 1047–1051 (1981).

48.   Kimura, S., Chen, H. P., Saxena, I. M., Brown, Jr & Itoh, T. Localization of c-di-GMP-

777   binding protein with the linear terminal complexes of Acetobacter xylinum. *J.*

778   *Bacteriol.* **183**, 5668–5674 (2001).

779   49.   Matthysse, Ann G., White, Sally & Lightfoot, Richard. Genes required for cellulose

780   synthesis in Agrobacterium tumefaciens. *J. Bacteriol.* **177**, 1069–1075 (1995).

781   50.   Deng, Ying, Nagachar, Nivedita, Fang, Lin, Luan, Xin, Catchmark, Jeffrey M., Tien,

782   Ming & Kao, Teh Hui. Isolation and characterization of two cellulose morphology

783   mutants of Gluconacetobacter hansenii ATCC23769 producing cellulose with lower

784   crystallinity. *PLoS One* **10**, e0119504 (2015).

785   51.   Pilhofer, Martin & Jensen, Grant J. *The bacterial cytoskeleton: More than twisted*

786   *filaments*. *Current Opinion in Cell Biology* vol. 25 1–9 (2013).

787   52.   Kühn, Juliane, Briegel, Ariane, Mörschel, Erhard, Kahnt, Jörg, Leser, Katja, Wick,

788   Stephanie, Jensen, Grant J. & Thanbichler, Martin. Bactofilins, a ubiquitous class of

789   cytoskeletal proteins mediating polar localization of a cell wall synthase in Caulobacter

790   crescentus. *EMBO J.* **29**, 327–339 (2010).

791   53.   Ingerson-Mahar, Michael, Briegel, Ariane, Werner, John N., Jensen, Grant J. & Gitai,

792   Zemer. The metabolic enzyme CTP synthase forms cytoskeletal filaments. *Nat. Cell*

793   *Biol.* **12**, 739–746 (2010).

794   54.   Pear, J. R., Kawagoe, Y., Schreckengost, W. E., Delmer, D. P. & Stalker, D. M. Higher

795   plants contain homologs of the bacterial celA genes encoding the catalytic subunit of

796   cellulose synthase. *Proc. Natl. Acad. Sci. U. S. A.* **93**, 12637–42 (1996).

797   55.   Nobles, David R. & Brown, R. Malcolm. The pivotal role of cyanobacteria in the

798   evolution of cellulose synthases and cellulose synthase-like proteins. *Cellulose* **11**,

799   437–448 (2004).

56.   Nobles, D. R., Romanovicz, D. K. & Brown, Jr. Cellulose in cyanobacteria. Origin of

vascular plant cellulose synthase? *Plant Physiol.* **127**, 529–542 (2001).

57.   Lampugnani, Edwin R., Flores-Sandoval, Eduardo, Tan, Qiao Wen, Mutwil, Marek,

Bowman, John L. & Persson, Staffan. Cellulose Synthesis – Central Components and

Their Evolutionary Relationships. *Trends in Plant Science* vol. 24 402–412 (2019).

58.   Diotallevi, Fabiana & Mulder, Bela. The cellulose synthase complex: A polymerization

driven supramolecular motor. *Biophys. J.* **92**, 2666–2673 (2007).

59.   Chan, Jordi, Coen, Enrico, Chan, Jordi & Coen, Enrico. Interaction between

Autonomous and Microtubule Guidance Systems Controls Cellulose Synthase Report

Interaction between Autonomous and Microtubule Guidance Systems Controls

Cellulose Synthase Trajectories. *Curr. Biol.* 1–7 (2020).

60.   Paredez, Alexander R., Somerville, Christopher R. & Ehrhardt, David W. Visualization

of cellulose synthase demonstrates functional association with microtubules. *Science*

312, 1491–1495 (2006).

61.   Li, Shundai, Lei, Lei, Somerville, Christopher R. & Gu, Ying. Cellulose synthase

interactive protein 1 (CSI1) mediates the intimate relationship between cellulose

microfibrils and cortical microtubules. *Plant Signal. Behav.* **7**, 1–5 (2012).

62.   Sampathkumar, Arun, Peaucelle, Alexis, Fujita, Miki, Schuster, Christoph, Persson,

Staffan, Wasteneys, Geoffrey O. & Meyerowitz, Elliot M. Primary wall cellulose

synthase regulates shoot apical meristem mechanics and growth. *Development* **146**,

(2019).

63.   Fujita, Miki, Himmelspach, Regina, Ward, Juliet, Whittington, Angela, Hasenbein,

Nortrud, Liu, Christine, Truong, Thy T., Galway, Moira E., Mansfield, Shawn D.,

Hocart, Charles H. & Wasteneys, Geoffrey O. The anisotropy1 D604N mutation in the

Arabidopsis cellulose synthase1 catalytic domain reduces cell wall crystallinity and the

velocity of cellulose synthase complexes. *Plant Physiol.* **162**, 74–85 (2013).

64.   Fujita, Miki, Himmelspach, Regina, Hocart, Charles H., Williamson, Richard E.,

827   Mansfield, Shawn D. & Wasteneys, Geoffrey O. Cortical microtubules optimize cell-

828   wall crystallinity to drive unidirectional growth in Arabidopsis. *Plant J.* **66**, 915–928

829   (2011).

830   65.   Liu, Zengyu, Schneider, Rene, Kesten, Christopher, Zhang, Youjun Yi, Somssich,

831   Marc, Zhang, Youjun Yi, Fernie, Alisdair R. & Persson, Staffan. Cellulose-

832   Microtubule Uncoupling Proteins Prevent Lateral Displacement of Microtubules during

833   Cellulose Synthesis in Arabidopsis. *Dev. Cell* **38**, 305–315 (2016).

834   66.   Standal, R., Iversen, T. G., Coucheron, D. H., Fjaervik, E., Blatny, J. M. & Valla, S. A

835   new gene required for cellulose production and a gene encoding cellulolytic activity in

836   Acetobacter xylinum are colocalized with the bcs operon. *J. Bacteriol.* **176**, 665–672

837   (1994).

838   67.   López, Daniel, Vlamakis, Hera, Losick, Richard & Kolter, Roberto. Cannibalism

839   enhances biofilm development in bacillus subtilis. *Mol. Microbiol.* **74**, 609–618 (2009).

840   68.   Billings, Nicole, Birjiniuk, Alona, Samad, Tahoura S., Doyle, Patrick S. & Ribbeck,

841   Katharina. Material properties of biofilms - A review of methods for understanding

842   permeability and mechanics. *Reports Prog. Phys.* **78**, (2015).

843   69.   Stewart, Philip S. Diffusion in biofilms. *Journal of Bacteriology* vol. 185 1485–1491

844   (2003).

845   70.   Xu, Karen D., Stewart, Philip S., Xia, Fuhu, Huang, Ching Tsan & McFeters, Gordon

846   A. Spatial physiological heterogeneity in Pseudomonas aeruginosa biofilm is

847   determined by oxygen availability. *Appl. Environ. Microbiol.* **64**, 4035–4039 (1998).

848   71.   Jo, Jeanyoung, Cortez, Krista L., Cornell, William Cole, Price-Whelan, Alexa &

849   Dietrich, Lars E. P. An orphan cbb3-type cytochrome oxidase subunit supports

850   Pseudomonas aeruginosa biofilm growth and virulence. *Elife* **6**, (2017).

72.   Sartori, N., Richter, Karsten & Dubochet, Jacques. Vitrification depth can be increased

more than 10-fold by high-pressure freezing. *J. Microsc.* **172**, 55–61 (1993).

73.   Harapin, Jan, Börmel, Mandy, Sapra, K. Tanuj, Brunner, Damian, Kaech, Andres &

Medalia, Ohad. Structural analysis of multicellular organisms with cryo-electron

tomography. *Nat. Methods* (2015).

74.   Hsieh, Chyongere, Schmelzer, Thomas, Kishchenko, Gregory, Wagenknecht, Terence

& Marko, Michael. Practical workflow for cryo focused-ion-beam milling of tissues

and cells for cryo-TEM tomography. *J. Struct. Biol.* **185**, 32–41 (2014).

75.   Schaffer, Miroslava, Pfeffer, Stefan, Mahamid, Julia, Kleindiek, Stephan, Laugks, Tim,

Albert, Sahradha, Engel, Benjamin D., Rummel, Andreas, Smith, Andrew J.,

Baumeister, Wolfgang & Plitzko, Juergen M. A cryo-FIB lift-out technique enables

molecular-resolution cryo-ET within native Caenorhabditis elegans tissue. *Nat.*

Methods **16**, 757–762 (2019).

76.   Das, Aditi & Das, Anath. Delineation of polar localization domains of *Agrobacterium*

tumefaciens type IV secretion apparatus proteins VirB4 and VirB11. *Microbiologyopen*

3, 793–802 (2014).

77.   Rigort, Alexander, Bäuerlein, Felix J. B., Leis, Andrew, Gruska, Manuela, Hoffmann,

Christian, Laugks, Tim, Böhm, Ulrike, Eibauer, Matthias, Gnaegi, Helmut, Baumeister,

Wolfgang & Plitzko, Jürgen M. Micromachining tools and correlative approaches for

cellular cryo-electron tomography. *J. Struct. Biol.* **172**, 169–179 (2010).

78.   Danev, Radostin, Buijsse, Bart, Khoshouei, Maryam, Plitzko, Jürgen M. & Baumeister,

Wolfgang. Volta potential phase plate for in-focus phase contrast transmission electron

microscopy. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 15635–40 (2014).

79.   Mastronarde, David N. Automated electron microscope tomography using robust

prediction of specimen movements. *J. Struct. Biol.* **152**, 36–51 (2005).

80.   Kremer, J. R., Mastronarde, D. N. & McIntosh, J. R. Computer visualization of three-dimensional image data using IMOD. *J. Struct. Biol.* **116**, 71–6 (1996).

Supplemental movie legends

Supplemental video 1 | Movie showing multiple views of the tomographic volume shown in Figure 1C-E. The disposition of the cellulose ribbon relative to the cell and its very close contact with the bacterial envelope is demonstrated.

Supplemental video 2 | Movie showing multiple views of the tomographic volume shown in Figure 5A-C. The close association between the cortical belt and the cellulose ribbon are shown. The second part of the animation shows the tomographic volume shown in Figure 2 and Figure 5D-F. The multilayered structure of the cortical belt is visible.

Supplemental video 3 | Movie of the FIB-milling of a lamella through the *G. hansenii* biofilms shown in Figure 6 and the 3D organization of the cells and the cellulose within the biofilm shown in Figure 6. The second part of the animation shows the tomographic volume shown in Figure 7A-D. Numerous cytoplasmic vesicles and the cortical belt underneath a cellulose ribbon are visible.

Supplemental video 4 | Movie showing multiple views of the tomographic volume shown in Figure 8B-C. The cellulose fibers are seen around the cell. No cortical belt is visible. A polar flattening is visible at the top of the cell with a "thickening" of the OM and a large periplasmic density underneath. The putative UPP (amorphous aggregate) is seen on the side of the polar flattening.

Figures

A — MT-Deep red FM, Calcofluor-W, 5 μm

B — 5 μm

C — OM, IM, CB, S, 100 nm

Legend:
- Outer membrane (dark green)
- Inner membrane (light green)
- Crystalline cellulose ribbon (yellow)
- Storage granule (blue)
- Cortical belt (purple)

D — 100 nm

E — 90°, 100 nm

F — OM, IM, CB, Cytosol, Exterior, 50 nm

G — OM, IM, CB, 50 nm

H — OM, IM, CB, 50 nm

Figure 1 | Interactions between the bacterial envelope and the cellulose ribbon: the "tight" configuration

(**A**) Confocal-Airy scan optical slices show representative examples of *G. hansenii* cells in red (MitoTracker Deep Red FM) displaying the cellulose ribbon on their side in cyan (Calcofluor-white). (**B**) Enlarged view indicated by white dashed rectangle in (A). The cellulose structure is clearly seen closely appended to one side of the cell (cyan arrowheads). (**C**) 9-nm thick tomographic slice showing the typical *G. hansenii* cell harboring the cellulose ribbon on its right side (yellow arrows). White arrowheads point to ribosomes and red arrows point to cytosolic vesicles. Here and below, IM: Inner-membrane; OM: Outer-membrane; S: Storage granule; CB: Cortical belt. (**D**) Manual segmentation of the cell shown in (C). (**E**) Rotated segmented volume shown in (D) showing the very close contact between the cellulose ribbon (yellow) and the outer membrane (green). (**F-H**) Transverse 9-nm thick tomographic slices through the bacterial envelope of the cell shown in (C) at the levels indicated by the blue, black and red dashed lines, respectively. Two cellulose sheets (yellow arrows) are seen. One interacts with the OM all along (white arrow). Our interpretation is that integration of the cellulose fibers into the sheet occurs immediately upon secretion.

Outer membrane ▮ (green)
Inner membrane ▮ (light green)
Crystalline cellulose ribbon ▮ (yellow)
Cortical belt ▮ (purple)
Disorganized cellulose aggregates ▮ (red)

Figure 2 | Interactions between the bacterial envelope and the cellulose ribbon: the "loose"

configuration

(**A**) 9-nm thick tomographic slice showing a cell where aggregates of disorganized cellulose

(orange asterisks) occur between the ribbon (yellow arrows) and the OM. Note the cortical belt

(CB) cannot be seen in this slice. Black line orange arrows indicate points of contact between

the cellulose sheet and the disorganized aggregates. Red arrows point to vesicles. (**B**) Manual

segmentation of the tomogram in (A) showing these disorganized aggregates in 3-D. (**C-D**)

Transverse 9-nm thick tomographic slices through the envelope of the cell shown in (A) at the

levels indicated by the blue and pink dashed lines highlighting the distance between the two

cellulose sheets (yellow arrows) and the OM and the presence of the disorganized clusters

(orange dashed brackets). (**E**) Plot showing the OM-to-closest sheet distance in the two types

of configuration.

Figure 3 | Cellulose sheet dimensions

(**A-B**) Longitudinal and transverse schematic depiction defining the different dimensions

measured, namely OM-to-sheet distance, sheet width and inter-sheet distance. Identical

terminology is used for the measurements of the cortical belt. (**C**) Transverse 12-nm thick slice

of the bacterial envelope of the cell shown in (D) at the level indicated by the blue dashed line.

The yellow arrows highlight the two stacked sheets. On the right, the average density profile

along the red line demonstrates how the cellulose sheet widths were estimated. Vertical axis is

length in nm along the red line and horizontal axis is the normalized electron density. (**D**) 12-

nm thick tomographic slice showing the typical organization of the bacterial envelope on the

side where cellulose sheets (yellow arrows) are being synthesized. The average density profile

on the right taken along the red line shows the CB-IM, IM-OM OM-sheet and inter-sheet

distances (green dashed lines).

A

Cellulose ribbons
Disorganized aggregates

Fraction of cells (%)

13min 20min 300min

B

cellulose sheets / ribbon

20min 300min

C

OM
IM
CB

100 nm

+ 20 min

D

OM
IM

100 nm

+ 300 min

E

100 nm

F

Bacterial envelope
Cellulose

100 nm

G

50 nm

11 nm

density (arbitrary unit)

0 2 4 6 8 10 12 14 16 18 20 22 24 26 28 30 32 34 36 38 40

H

Microfibril diameter (nm)

Intersheet distance (nm)

Cell #1   Cell #2   t-300min

I

Sheet width (nm)

20min 300min

Figure 4 | The cellulose ribbon is a composite structure made of stacked sheets

(**A**) Percentages of cells exhibiting disorganized aggregates (blue) and cellulose ribbons (red) at 13-, 20- and 300minutes post-separation. While disorganized aggregate occurrence is steady, there is an increase in the occurrence of cellulose ribbons over time. (**B**) Number of cellulose sheets composing the ribbons as a function of time after cell separation. (**C**) Composite image composed of 10-nm thick tomographic slices spaced by 24 nm in Z, of a cell 20 minutes post-separation in the "tight" configuration. The cellulose ribbon is thin (yellow arrows), composed of one sheet immediately adjacent to the OM. Limits of the two original images are indicated by the red dashed line. (**D**) 11 nm thick tomographic slice of a cell 300 minutes post-separation. The cellulose ribbon (yellow arrows) is large and composed of multiple sheets. (**E**) Nascent cellulose sheet 20 minutes post-separation (yellow arrow). Putative microfibrils can be seen coming out perpendicularly from the outer membrane (red arrowheads). (**F**) Corresponding manual segmentation of (E). (**G**) Enlarged view of the blue boxed region in (E). Below is the average density profile showing the estimation of the diameter of one putative microfibril (red line). (**H**) Estimated diameters of microfibrils observed at 20-minutes post-separation in the two cells where they are visible (left vertical axis) as in (E) and the inter-sheet distances measured in the 300-minutes post-separation cellulose ribbons (right vertical axis). (**I**) Sheet width estimations at 20- and 300-minutes post separation.

Outer membrane
Inner membrane
Cellulose ribbon
Cortical belt

Figure 5 | The cortical belt lies below the cellulose ribbon in the cytoplasm

(**A**) 9-nm thick tomographic slice showing a representative cortical belt (purple arrows) just inside the IM and proximal to the cellulose ribbon on the outside of the cell (yellow arrows). (**B**) Manual segmentation of the tomogram shown in (A) highlighting the cellulose ribbon and the cortical belt. (**C**) Same segmentation rotated 90° about the long axis of the cell shows how the cortical belt and the cellulose ribbon follow the same trajectory. (**D**) 9-nm thick tomographic slice taken from the same tomogram as in figure 2, showing one out of several cases where the cortical belt presented stacked layers (red dashed box). (**E**) Enlarged view of the red dashed boxed region in (D) showing the arrangement of the stacked layers. On the right is a density profile displayed normal to the cortical belt to measure the inter-layer distance (15 nm). (**F**) Transverse 9-nm thick tomographic slice of the cell region shown in (D), at the level indicated by the blue dashed line, highlighting stacked layers of the cortical belt. The cellulose ribbon can be seen at a distance (yellow arrowheads) with disorganized aggregates in between (orange dashed brackets and asterisk).

A 1 mm

B 50 um

C 40 um

D
- Live cells
- Dead cells
- Cellulose

1 um

E 1 um

F

Volume of the lamella occupied by cells (%)

3h   6h

G

Live cell ratio

1.0

0.5

0.0

3h   6h

H

Sputtered platinum

Distance from the cell to leading front of the lamella "d"

d

Sputtered platinum

d

a

lamella

depth

cell

original biofilm

depth = d x tan(a)

Top view of lamella   Longitudinal side view of lamella

Live cells   Dead cells

Cell depth in lamella (um)

• 3h biofilms
× 6h biofilms

Figure 6 | FIB-milling through native *G. hansenii* biofilms

(**A**) cryo-SEM overview of a 6-hour biofilm (outlined in red) grown on a gold quantifoil grid. (**B**) cryo-SEM view of a thick biofilm area (boxed in blue in (A)). Wrinkles in the biofilm are typical of a biofilm a few microns thick. (**C**) Milled lamella (boxed in yellow) from the green boxed region shown in (A). (**D**) 23-nm thick tomographic slice of a low mag tomogram taken on the lamella shown in (C). Living (when frozen) and dead cells are visible (green and red asterisks, respectively) and large cellulose arrays can be seen filling the gaps between the cells (yellow arrowheads). (**E**) Manual segmentation of the tomogram shown in (D). (**F**) Fraction of the lamella volume occupied by the cells was assessed for each lamella (**G**) Live cell ratio in 3h and 6h biofilms. (**H**) Violin boxplots reporting the absolute depth of the live and dead cells within the biofilms grown for 3 and 6 hours. The dashed red lines indicate the first and third quartiles and solid red line represents median. This shows that while the biofilms get thicker with time, the ratio of live-to-dead cells appears constant through depth and time. Method of calculation is detailed on the left of the panel and in the methods section. Lamella is drawn in blue, with the platinum coated leading edge represented in gray.

Figure 7 | Lamellae of native biofilms also reveal numerous vesicles and the cortical belt

(**A-B**) Two tomographic slices of a *G. hansenii* cell from a biofilm grown for 6h surrounded by cellulose ribbons (yellow arrowheads). The cortical belt is visible in (B) (purple arrow) and seems to follow the trajectory of the cellulose sheet proximal to the OM (dark lined yellow arrowhead). (**C**) Manual segmentation of the tomogram displayed in (A) and (B) showing the juxtaposition of the cortical belt (purple to red) and the nascent cellulose ribbon (yellow). (**D**) Enlargement of the boxed region in (B) showing the layered cortical belt. (**E**) Tomographic slice of a cell surrounded by cellulose ribbons (yellow arrowheads) from a biofilm grown for 3h and harboring numerous vesicles in its cytosol (white arrowheads). Disorganized aggregates (orange dashed lines) are visible at this timepoint. (**F-G**) Tomographic slices showing additional examples of disorganized cellulose aggregates (orange dashed lines) surrounded by cellulose ribbons (yellow arrowheads) visible in 3h biofilms. Scale bars = 100 nm. All tomographic slices are 11-nm thick.

Agrobacterium tumefaciens

A
- C. White (cellulose)
- MitoTracker (cells)
- 2 um

B
- OM
- IM
- S
- 200 nm

C
- Storage granule
- Inner/outer membrane
- Putative UPP
- Cellulose microfibrils
- 200 nm

Escherichia coli 1094

D
- C. White (cellulose)
- MitoTracker (cells)
- 5um

E
- 200 nm
- Untreated

F
- 200 nm
- + Cellulase

Figure 8 | The cortical belt is not found in other cellulose-synthesizing species

(**A**) Maximum projection of *A. tumefaciens* cells synthesizing cellulose. Cells are stained with

Mito Tracker Deep Red (red) and cellulose with Calcofluor-white (cyan). (**B**) 10-nm thick

tomographic slice of a typical *A. tumefaciens* cell with cellulose microfibrils around (yellow

arrowheads). No cortical belt can be seen in the cells. A polar flattening can be seen at the lower

pole (cyan arrow) with an amorphous aggregate (orange dashed lines). These aggregates are

most probably the UniPolar-Polysaccharide (UPP) synthesized specifically by *A. tumefaciens*.

(**C**) Manual segmentation of the tomogram in (B) showing the organization of the cellulose

microfibrils around the cell, the absence of the cortical belt and the putative UPP.   (**D**) 50-nm

optical slice of an induced *E. coli 1094* cellulose biofilm. Cells are stained with mitoTracker

Deep Red (red) and cellulose with Calcofluor-white (cyan). (**E**) 6-nm tomographic slice of a

lamella tomogram of bacterial mat showing three *E. coli 1094* cells and an amorphous cellulose

aggregate between them (orange asterisk). (**F**) 6-nm tomographic slice of a lamella through a

bacterial mat treated with cellulase, showing multiple cells. No cellulose was visible in this

condition. No cortical belt can be seen in the cells in either condition.

1086

1087

1088

1089

1090

1091

1092

1093

1094

1095

Top view of the extrusion pores

BcsC subunits
Cortical belt
Elementary fibrils aggregates
Extrusion pore

IM   OM

~10 nm diameter microfibrils

Cellulose sheet

Lengthening over time

Width

90°

Side view of the extrusion pores

IM

OM

Cellulose sheet

Lengthening over time

Thickness

Elementary fibril aggregates

Microfibril

Cellulose sheet

IM   OM

Figure 9 | Updated cell-directed hierarchical model

Top (left) and side (right) view of a *G. hansenii* cell showing the different aggregation steps leading to a cellulose sheet, how microfibrils contribute to sheet width and the role of the cortical belt. In this model, clusters of 9 extrusion pores are depicted (green circles), the real numbers and distribution are unknown. Each extrusion pore is presented as comprising 4 BcsC subunits each (red circles), the actual number is not known. Inset in blue is a magnified view of the line of 9 extrusion pores, each hypothesized to extrude an aggregate of multiple elementary fibrils. All aggregates then coalesce to form a microfibril. These microfibrils then stack together, contributing to the width of the cellulose sheet. Adapted from the cell in figure 4E-G.

Whole cell tomography

Species	Condition/strain	Lamellae	Tomograms	VPP	Source
G. hansenii (ATCC 23769)	Untreated - 13min post sep	N/A	6	-	ATCC
	Untreated - 20min post sep	N/A	14	-	
	Untreated - 300min post sep	N/A	33	24	
	Cellulase 0.2g/L	N/A	4	-	
G. xylinus (ATCC 700178)	Untreated - 300min post sep	N/A	8	-	ATCC
A. tumefaciens C58[1, 3]	WT[2]	N/A	47	-	gift from Patricia Zambrisky (UC Berkeley) to Elitza Tocheva
	A139[2]	N/A	10	-	gift from Patricia Zambrisky (UC Berkeley) to Elitza Tocheva
	AD348[2]	N/A	1	-	gift from Anath Das (University of Minnesota) to Debnath Ghosal
	AD1484[2]	N/A	1	-	gift from Anath Das (University of Minnesota) to Debnath Ghosal
	JX148[2]	N/A	4	-	gift from Patricia Zambrisky (UC Berkeley) to Elitza Tocheva
	NT1[2]	N/A	2	-	gift from Patricia Zambrisky (UC Berkeley) to Elitza Tocheva

Tomography on milled lamellae

Species	Condition/strain	Lamellae	Tomograms		Source
G. hansenii (ATCC 23769)	Native biofilm - untreated	12	33	3	ATCC
E. coli 1094 induced for cellulose synthesis	Untreated	2	6	-	gift from Jean Marc Ghigo (Institut Pasteur)
	Cellulase 0.2g/L	1	2	-	

[1] C58 or ATCC 33970 is the wild type *A. tumefaciens*

[2] All these strains have a C58 background.

[3] These strains were imaged for other purposes, but used here as well