Expanding the diversity of bacterioplankton isolates and modeling isolation efficacy with large scale dilution-to-extinction cultivation

Michael W. Henson[1,#], V. Celeste Lanclos[1], David M. Pitre[2], Jessica Lee Weckhorst[2,†], Anna M. Lucchesi[2], Chuankai Cheng[1], Ben Temperton[3*], and J. Cameron Thrash[1*]

[1]Department of Biological Sciences, University of Southern California, Los Angeles, CA 90089, U.S.A.
[2]Department of Biological Sciences, Louisiana State University, Baton Rouge, LA 70803, U.S.A.
[3]School of Biosciences, University of Exeter, Stocker Road, Exeter, EX4 4QD, U.K.
[#]Current affiliation: Department of Geophysical Sciences, University of Chicago, Chicago, IL 60637, U.S.A.
[†]Quantitative and Computational Biosciences Program, Baylor College of Medicine, Houston, TX 77030, U.S.A.

*Correspondence:

J. Cameron Thrash
thrash@usc.edu

Ben Temperton
b.temperton@exeter.ac.uk

Running title: Evaluation of large-scale DTE cultivation

Keywords: dilution-to-extinction, cultivation, bacterioplankton, LSUCC, microbial ecology, coastal microbiology

47
Abstract

Cultivated bacterioplankton representatives from diverse lineages and locations are
essential for microbiology, but the large majority of taxa either remain uncultivated or lack
isolates from diverse geographic locales. We paired large scale dilution-to-extinction (DTE)
cultivation with microbial community analysis and modeling to expand the phylogenetic and
geographic diversity of cultivated bacterioplankton and to evaluate DTE cultivation success.
Here, we report results from 17 DTE experiments totaling 7,820 individual incubations over
three years, yielding 328 repeatably transferable isolates. Comparison of isolates to microbial
community data of source waters indicated that we successfully isolated 5% of the observed
bacterioplankton community throughout the study. 43% and 26% of our isolates matched
operational taxonomic units (OTUs) and amplicon single nucleotide variants (ASVs),
respectively, within the top 50 most abundant taxa. Isolates included those from previously
uncultivated clades such as SAR11 LD12 and *Actinobacteria* acIV, as well as geographically
novel members from other ecologically important groups like SAR11 subclade IIIa, SAR116,
and others; providing the first isolates in eight genera and seven species. Using a newly
developed DTE cultivation model, we evaluated taxon viability by comparing relative abundance
with cultivation success. The model i) revealed the minimum attempts required for successful
isolation of taxa amenable to growth on our media, and ii) identified important and abundant taxa
such as SAR11 with low viability that likely impacts cultivation success. By incorporating
viability in experimental design, we can now statistically constrain the number of attempts
necessary for successful cultivation of a given taxon on a defined medium.
69
70

Importance

Even before the coining of the term "great plate count anomaly" in the 1980s, scientists
had noted the discrepancy between the number of microorganisms observed under the
microscope and the number of colonies that grew on traditional agar media. New cultivation
approaches have reduced this disparity, resulting in the isolation of some of the "most wanted"
bacterial lineages. Nevertheless, the vast majority of microorganisms remain uncultured,
hampering progress towards answering fundamental biological questions about many important
microorganisms. Furthermore, few studies have evaluated the underlying factors influencing
cultivation success, limiting our ability to improve cultivation efficacy. Our work details the use
of dilution-to-extinction (DTE) cultivation to expand the phylogenetic and geographic diversity
of available axenic cultures. We also provide a new model of the DTE approach that uses
cultivation results and natural abundance information to predict taxon-specific viability and
iteratively constrain DTE experimental design to improve cultivation success.

Introduction

Axenic cultures of environmentally important microorganisms are critical for fundamental microbiological investigation, including generating physiological information about environmental tolerances, determining organismal-specific metabolic and growth rates, testing hypotheses generated from *in situ* 'omics observations, and experimentally examining microbial interactions. Research using important microbial isolates has been critical to a number of discoveries such as defining microorganisms involved in surface ocean methane saturation (1–3), the role of proteorhodopsin in maintaining cellular functions during states of carbon starvation (4, 5), the complete nitrification of ammonia within a single organism (6), and identifying novel metabolites and antibiotics (7–10). However, the vast majority of taxa remain uncultivated (11–13), restricting valuable experimentation on such topics as genes of unknown function, the role of analogous gene substitutions in overcoming auxotrophy, and the multifaceted interactions occurring in the environment inferred from sequence data (11, 14–16).

The quest to bring new microorganisms into culture, and the recognition that traditional agar-plate based approaches have limited success (17–19), have compelled numerous methodological advances spanning a wide variety of techniques like diffusion chambers, microdroplet encapsulation, and slow acclimatization of cells to artificial media (20–25). Dilution-to-extinction (DTE) cultivation using sterile seawater as the medium has also proven highly successful for isolating bacterioplankton (26–32). Pioneered by Don Button and colleagues for the cultivation of oligotrophic bacteria, this method essentially pre-isolates organisms after serial dilution by separating individual or small groups of cells into their own incubation vessel (32, 33). This prevents slow-growing, obligately oligotrophic bacterioplankton from being outcompeted by faster-growing organisms, as would occur in enrichment-based isolation methods, particularly those that would target aerobic heterotrophs. It is also a practical method for taxa that cannot grow on solid media. Natural seawater media provide these taxa with the same chemical surroundings from which they are collected, reducing the burden of anticipating all the relevant compounds required for growth (33).

Improvements to DTE cultivation in multiple labs have increased the number of inoculated wells and decreased the time needed to detect growth (26, 28, 34), thereby earning the moniker "high-throughput culturing" (26, 28). We (35) and others (30) have also adapted DTE culturing by incorporating artificial media in place of natural seawater media to successfully isolate abundant bacterioplankton. Thus far, DTE culturing has lead to isolation of many numerically abundant marine and freshwater groups such as marine SAR11 Alphaproteobacteria (28, 29, 34–36), the freshwater SAR11 LD12 clade (29), SUP05/Arctic96BD-19 *Gammaproteobacteria* (37–39), OM43 *Betaproteobacteria* (26, 27, 31, 40, 41), HIMB11-Type Roseobacter (35, 42), numerous so-called Oligotrophic Marine *Gammaproteobacteria* (43), and acI *Actinobacteria* (44).

Despite the success of DTE cultivation, many taxa continue to elude domestication (11–13, 16). Explanations include a lack of required nutrients or growth factors in media (20, 45–49) and biological phenomena such as dormancy and/or phenotypic heterogeneity within populations (47, 48, 50–56). However, there have been few studies empirically examining the factors underlying isolation success in DTE cultivation experiments (34, 57, 58), restricting our ability to determine the relative importance of methodological vs. biological influences on cultivation reliability for any given organism. Moreover, even for those taxa that we have successfully cultivated, in many cases we lack geographically diverse strains, restricting comparisons of the phenotypic and genomic diversity that may influence taxon-specific cultivability.

We undertook a three-year cultivation effort in the coastal northern Gulf of Mexico
(nGOM), from which we lack representatives of many common bacterioplankton groups, to
provide new model organisms for investigating microbial function, ecology, biogeography, and
evolution. Simultaneously, we paired our cultivation efforts with 16S rRNA gene amplicon
analyses to compare cultivation results with the microbial communities in the source waters. We
have previously reported on the success of our artificial media in obtaining abundant taxa over
the course of seven experiments from this campaign (35). Here, we expand our report to include
the cultivation results from a total of seventeen experiments, and update the classic viability
calculations of Button *et al.* (33) with a new model to estimate the viability of individual taxa
using relative abundance information. Individuals from eight and seven cultivated groups
belonged to putatively novel genera and species, respectively, and our isolates expand the
geographic representation for many important clades like SAR11. Additionally, using model-
based predictions, we identify possible taxon-specific viability variation that can influence
cultivation success. By incorporating these new viability estimates into the model, our method
facilitates statistically-informed experimental design for targeting individual taxa, thereby
reducing uncertainty for future culturing work (59).
155
156   **Material and Methods**
157
Sampling
Surface water samples were collected at six different sites once a year for three years, except for
Terrebonne Bay, which was collected twice. The sites sampled were Lake Borgne (LKB; Shell
Beach, LA), Bay Pomme d'Or (JLB; Buras, LA), Terrebonne Bay (TBON; Cocodrie, LA),
Atchafalaya River Delta (ARD; Franklin, LA), Freshwater City (FWC; Kaplan, LA), and
Calcasieu Jetties (CJ; Cameron, LA) (lat/long coordinates provided in Table S1). Water
collection for biogeochemical and biological analysis followed the protocol in (35). Briefly, we
collected surface water in a sterile, acid-washed polycarbonate bottle. Duplicate 120 ml water
samples were filtered serially through 2.7 µm Whatman GF/D (GE Healthcare, Little Chalfort,
UK) and 0.22 µm Sterivex (Millipore, Massachusetts, USA) filters and placed on ice until
transferred to -20˚C in the laboratory (maximum 3 hours on ice). The University of Washington
Marine Chemistry Laboratory analyzed duplicate subsamples of 50 ml 0.22 µm-filtered water
collected in sterile 50 ml falcon tubes (VWR, Pennsylvania, USA) for concentrations of SiO_4,
PO_4^{3-}, NH_4^+, NO_3^-, and NO_2^-. Samples for cell counts were filtered through a 2.7 µm GF/D filter,
fixed with 10% formaldehyde, and stored on ice until enumeration (maximum 3 hours).
Temperature, salinity, pH, and dissolved oxygen were measured using a handheld YSI 556
multiprobe system (YSI Inc., Ohio, USA). All metadata is available in Table S1.
175
Dilution-to-extinction culturing and propagation
Isolation, propagation, and identification of isolates were completed as previously reported (29,
35). A subsample of 2.7 µm filtered surface water was stained with 1X SYBR Green (Lonza,
Basal, Switzerland), and enumerated using a flow cytometer as described (60). After serial
dilution to a predicted 1-3 cells·µl^{-1}, 2 µl water was inoculated into five, 96-well PTFE plates
(Radleys, Essex, UK) containing 1.7 ml artificial seawater medium (Table S1) to achieve an
estimated 1-3 cells·well^{-1} (Table 1). The salinity of the medium was chosen to match *in situ*
salinity after experiment JLB (January 2015) (Table S1). After year two (TBON3-LKB3), a
second generation of media, designated MWH, was designed to incorporate additional important

osmolytes, reduced sulfur compounds, and other constituents (Table S1) potentially necessary
for *in vitro* growth of uncultivated clades (49, 61–67). Cultures were incubated at *in situ*
temperatures (Table S1) in the dark for three to six weeks and evaluated for positive growth (>
10^4 cells·ml^{-1}) by flow cytometry. 200 µl from positive wells was transferred to duplicate 125 ml
polycarbonate flasks (Corning, New York, USA) containing 50 ml of medium. At FWC, FWC2,
JLB2c, and JLB3, not all positive wells were transferred because of the large number of positive
wells. At each site, 48/301, 60/403, 60/103, and 60/146 of the positive wells were transferred,
respectively, selecting for flow cytometric signals that maximized our chances of isolating small
microorganisms that encompass many of the most abundant, and most wanted taxa, like SAR11.
194
Culture identification
Cultures reaching $\geq$ 1 x 10^5 cells·ml^{-1} had 35 ml of the 50 ml volume filtered for identification
via 16S rRNA gene PCR onto 25 mm 0.22-µm polycarbonate filters (Millipore, Massachusetts,
USA). DNA extractions were performed using the MoBio PowerWater DNA kit (QIAGEN,
Massachusetts, USA) following the manufacturer's instructions and eluted in sterile water. The
16S rRNA gene was amplified as previously reported in Henson et al. 2016 (35) and sequenced
at Michigan State University Research Technology Support Facility Genomics Core. Evaluation
of Sanger sequence quality was performed with 4Peaks (v. 1.7.1)
(http://nucleobytes.com/4peaks/) and forward and reverse complement sequences (converted via
http://www.bioinformatics.org/sms/rev_comp.html) were assembled where overlap was
sufficient using the CAP3 web server (http://doua.prabi.fr/software/cap3).
206
Community iTag sequencing, operational taxonomic units, and single nucleotide variants
Sequentially (2.7µm, 0.22µm) filtered duplicate samples were extracted and analyzed using our
previously reported protocols and settings (35, 68). We sequenced the 2.7-0.22 µm fraction for
this study because this fraction corresponded with the < 2.7 µm communities that were used for
the DTE experiments. To avoid batch sequencing effects, DNA from the first seven collections
reported in (35) was resequenced with the additional samples from this study (FWC2 and after-
Table 1). We targeted the 16S rRNA gene V4 region with the 515F, 806RB primer set (that
corrects for poor amplification of taxa like SAR11) (69, 70) using Illumina MiSeq 2 x 250bp
paired-end sequencing at Argonne National Laboratories, resulting in 2,343,106 raw reads for
the 2.7-0.22 µm fraction. Using Mothur v1.33.3 (71), we clustered 16S rRNA gene amplicons
into distinctive OTUs with a 0.03 dissimilarity threshold (OTU$_{0.03}$) and classified them according
to the Silva v119 database (72, 73). After these steps, 55,256 distinct OTUs$_{0.03}$ remained. We
also used minimum entropy decomposition (MED) to partition reads into fine-scale amplicon
single nucleotide variants (ASVs) (74). Reads were first analyzed using Mothur as described
above up to the *screen.seqs()* command. The cleaned reads fasta file was converted to MED-
compatible headers with the 'mothur2oligo' tool renamer.pl from the functions in MicrobeMiseq
(https://github.com/DenefLab/MicrobeMiseq) (75) using the fasta output from *screen.seqs()* and
the Mothur group file. These curated reads were analyzed using MED (v. 2.1) with the flags –M
60, and -d 1. MED resulted in 2,813 refined ASVs. ASVs were classified in Mothur using
classify.seqs(), the Silva v119 database, and a cutoff bootstrap value of 80% (76). After
classification, we removed ASVs identified as "chloroplast", "mitochondria", or "unknown"
from the dataset.
229
Community analyses

OTU (OTU$_{0.03}$) and ASV abundances were analyzed within the R statistical environment v.3.2.1
(77) following previously published protocols (29, 35, 68). Using the package PhyloSeq (78),
OTUs and ASVs were rarefied using the command *rarefy_even_depth()* and OTUs/ASVs
without at least two reads in four of the 34 samples (2 sites; ~11%) were removed. This cutoff
was used to remove any rarely occurring or spurious OTUs/ASVs. Our modified PhyloSeq
scripts are available on our GitHub repository https://github.com/thrash-lab/Modified-Phyloseq.
After filtering, the datasets contained 777 unique OTUs and 1,323 unique ASVs (Table S1). For
site-specific community comparisons, beta diversity between sites was examined using Bray-
Curtis distances via ordination with non-metric multidimensional scaling (NMDS) (Table S1).
The nutrient data were normalized using the R function *scale* which subtracts the mean and
divides by the standard deviation for each column. The influence of the transformed
environmental parameters on beta diversity was calculated in R with the *envfit* function. Relative
abundances of an OTU or ASV from each sample were calculated as the number of reads over
the sum of all the reads in that sample. The relative abundance was then averaged between
biological duplicates for a given OTU or ASV. To determine the best matching OTU or ASV for
a given LSUCC isolate, the OTU representative fasta file, provided by Mothur using
get.oturep(), and the ASV fasta file were used to create a BLAST database (*makeblastdb*) against
which the LSUCC isolate 16S rRNA genes could be searched via blastn (BLAST v 2.2.26)
("OTU_ASVrep_db" - Available as Supplemental Information at
https://doi.org/10.6084/m9.figshare.12142098). We designated a LSUCC isolate 16S rRNA gene
match with an OTU or ASV sequence based on ≥ 97% or ≥ 99% sequence identity, respectively,
as well as a ≥ 247 bp alignment.
253
16S rRNA gene phylogeny
Taxa in the *Alpha-*, *Beta-*, and *Gammaproteobacteria* phylogenies from (35) served as the
backbone for the trees in the current work. For places in these trees with poor representation near
isolate sequences, additional taxa were selected by searching the 16S rRNA genes of LSUCC
isolates against the NCBI nt database online with BLASTn (79) and selecting a variable number
of best hits. The *Bacteroidetes* and *Actinobacteria* trees were composed entirely of non-
redundant top 100-300 MegaBLAST hits to a local version of the NCBI nt database, accessed
August 2018. Sequences were aligned with MUSCLE v3.6 (80) using default settings, culled
with TrimAl v1.4.rev22 (81) using the -automated1 flag, and the final alignment was inferred
with IQ-TREE v1.6.11 (82) with default settings and -bb 1000 for ultrafast bootstrapping (83).
Tips were edited with the nw_rename script within Newick Utilities v1.6 (84) and trees were
visualized with Archaeopteryx (85). Fasta files for these trees and the naming keys are available
as Supplemental Information at https://doi.org/10.6084/m9.figshare.12142098.
267
Assessment of isolate novelty
We quantified taxonomic novelty using BLASTn of our isolate 16S rRNA genes to those of
other known isolates collected in three databases: 1) The NCBI nt database (accessed August
2018) - "NCBIdb"; 2) a custom database comprised of sequences from other DTE experiments -
"DTEdb"; and 3) a database containing all our isolate 16S rRNA genes - "LSUCCdb". The
DTEdb and LSUCCdb fasta files are available as Supplemental Information at
https://doi.org/10.6084/m9.figshare.12142098. We compared our isolate sequences to these
databases as follows:

1) All representative sequences were searched against the nt database using BLASTn
(BLAST+v. 2.7.1) with the flags -perc_identity 84, -evalue 1E-6, -task blastn, -outfmt "6
qseqid sseqid pident length slen qlen mismatch evalue bitscore sscinames sblastnames
stitle", and -negative_gilist to remove uncultured and environmental sequences. The
negative GI list was obtained by searching `"environmental samples"[organism] OR
metagenomes[orgn]"` in the NCBI Nucleotide database (accessed September 12th, 2018)
and hits were downloaded in GI list format. This negative GI list is available as
Supplemental Information at https://doi.org/10.6084/m9.figshare.12142098. The resultant
hits from the nt database search were further manually curated to remove sequences
classified as single cell genomes, clones, duplicates, and previously deposited LSUCC
isolates.
2) We observed that many known HTCC, IMCC, and HIMB isolates that had previously
been described as matching our clades (Figs. S1-5) were missing from the resultant lists
of nt hits, so we extracted isolate accession numbers from numerous DTE experiments
(26–28, 31, 34, 37, 44, 86, 87) from the nt database via *blastdbcmd* and generated a
separate DTEdb using *makeblastdb*. Duplicate accession numbers found in the NCBIdb
were removed. The same BLASTn settings as in 1) were used to search our isolate
sequences against DTEdb. Any match that fell below the lowest percent identity hit to the
NCBIdb was removed from the DTEdb search since the match would not have been
present in the first NCBIdb search.
3) Finally, using the same BLASTn settings, we compared all pairwise identities of our 328
LSUCC isolate 16S rRNA gene sequences via the LSUCCdb.
The output from these searches is available in Table S1 under the "taxonomic novelty" tab.
We placed our LSUCC isolates into 55 taxonomic groups based on sharing ≥ 94%
identity and/or their occurrence in monophyletic groups within our 16S rRNA gene trees (Figs.
S1-5, see above). For visualization purposes, in groups with multiple isolates we used our
chronologically first cultivated isolate as the representative sequence for blastn searches, and
these are the top point (100% identity to itself) in each group column of Figure 1. Sequences
from the other DTE culture collections were labeled with the corresponding collection name,
while all other hits were labeled as "Other".
Geographic novelty was assessed by manually screening the accession numbers from hits
to LSUCC isolates with ≥ 99% 16S rRNA gene sequence identity for the latitude and longitude
from a connected publication or location name (e.g. source, country, site) in the NCBI
description. LSUCC isolates in the *Janibacter* sp., *Micrococcus* sp., *Altererythrobacter* sp.,
Pseudomonas sp., and *Phycicoccus* sp. groups (16 total isolates) were not assessed because of
missing isolation source information and no traceable publication.
312
Modeling DTE cultivation via Monte Carlo simulations
We developed a model using Monte Carlo simulation to estimate the median number of positive
and pure wells (and associated 95% confidence intervals (CI)) expected from a DTE experiment
for a given taxon at different inoculum sizes (λ), relative abundances (r), and viability (V) (Fig.
5). For each bootstrap, the number of cells added to each well was simulated using a Poisson
distribution at a mean inoculum size of λ cells per well across n wells. The number of cells added
to each well that belonged to a specific taxon was then estimated using a binomial distribution
where the number of trials was set as the number of cells in a well and the probability of a cell
belonging to a specific taxon, r, was the relative abundance of its representative ASV in the

community analysis. Wells that contained at least one cell of a specific taxon were designated
'positive'. Wells in which all the cells belonged to a specific taxon were designated as 'pure'.
Finally, the influence of taxon-specific viability on recovery of 'pure' wells was simulated using
a second binomial distribution, where the number of cells within a 'pure' well was used as the
number of trials and the probability of growth was a viability score ranging from 0 to 1. For each
simulation, 9,999 bootstraps were performed. Code for the model and all simulations is available
in the 'viability_test.py' at our GitHub repository https://github.com/thrash-lab/viability_test.
329
Actual versus expected number of isolates
For each taxon in each DTE experiment, the Monte Carlo simulation was used to evaluate
whether the number of recovered pure wells for each taxon was within 95% CI of simulated
estimates, assuming optimum growth conditions (i.e. $V = 100\%$). For each of 9,999 bootstraps,
460 wells were simulated with the inoculum size used for the experiment and the relative
abundance of the ASV. For taxa where the number of expected wells fell outside the 95% CI of
the model, a deviance score was calculated as the difference between the actual number of wells
observed and median of the simulated dataset. The results of this output are presented in Table
S1 under the "Expected vs actual" tab, and the R script for visualizing this output as Figure 6 is
available at our GitHub repository https://github.com/thrash-lab/EvsA-visualization.
340
Estimating viability in under-represented taxa
For taxa where the observed number of positive wells was lower than the 95% CI lower limit
within a given experiment, and because our analysis was restricted to only those organisms for
which our media was sufficient for growth at least once, the deviance was assumed to be a
function of a viability term, $V,$ (ranging from 0 to 1) associated with suboptimal growth
conditions, dormancy, persister cells, etc. To estimate a value of viability for a given taxon
within a particular experiment, the Monte Carlo simulation was run using an experiment-
appropriate inoculum size, relative abundance, and number of wells (460 for each experiment).
Taxon-specific viability was tested across a range of decreasing values from 99% to 1% until
such time as the observed number of pure wells for a given taxon fell between the 95% CI
bounds of the simulated data. At this point, the viability value is the maximum viability of the
taxon that enables the observed number of pure wells for a given taxon to be explained by the
model. The results of this output are presented in Table S1 under the "Expected vs actual" tab.
354
Likelihood of recovering taxa at different relative abundances
To estimate the number of wells required in a DTE experiment to have a significant chance of
recovering a taxon with a relative abundance of *r,* assuming optimum growth conditions ($V =$
100%), the Monte Carlo model was used to simulate experiments from 92 wells to 9,200 wells
per experiment across a range of relative abundances from 0 to 100% in 0.1% increments, and a
range of inoculum sizes (cells per well of 1, 1.27, 1.96, 2, 3, 4 and 5). Each experiment was
bootstrapped 999 times and the number of bootstraps in which the lower-bound of the 95% CI
was ≥ 1 was recorded.
363
Data accessibility
All iTag sequences are available at the Short Read Archive with accession numbers
SRR6235382-SRR6235415 (29). PCR-generated 16S rRNA gene sequences from this study are
accessible on NCBI GenBank under the accession numbers MK603525-MK603769. Previously

generated 16S rRNA genes sequences are accessible on NCBI GenBank under the accession
numbers KU382357-KU38243 (35). Table S1 is available at
https://doi.org/10.6084/m9.figshare.12142113.
371
Results
General cultivation campaign results
We conducted a total of seventeen DTE cultivation experiments to isolate bacterioplankton (sub
2.7 µm fraction), with paired microbial community characterization of source waters (0.22 µm -
2.7 µm fraction), from six coastal Louisiana sites over a three year period (Table S1). We
inoculated 7,820 distinct cultivation wells (all experiments) with an estimated 1-3 cells·well^{-1}
using overlapping suites of artificial seawater media, JW (years 1 and 2, (35)) and MWH (year
3), designed to match the natural environment (Table 1). The MWH suite of media was modified
from the JW media to additionally include choline, glycerol, glycine betaine, cyanate, DMSO,
DMSP, thiosulfate, and orthophosphate. These compounds have been identified as important
metabolites and osmolytes for marine and freshwater prokaryotes and were absent in the first
iteration (JW) of our media (Table S1) (88–94). A total of 1,463 wells were positive ($> 10^4$
cells·ml^{-1}), and 738 of these were transferred to 125 mL polycarbonate flasks. For four
experiments (FWC, FWC2, JLB2, and JLB3) we only transferred a subset of positives (48/301,
60/403, 60/103, and 60/146) because the number of isolates exceeded our ability to maintain and
identify them at that time (Table 1). The subset of positive wells for these four experiments was
selected using flow cytometry signatures with $< 10^2$ green fluorescence and $< 10^2$ side scatter-
usually indicative of smaller oligotrophic cells like SAR11 strain HTCC1062 (49) using our
settings. Of the 738 wells from which we transferred cells across all experiments, 328 of these
yielded repeatably transferable isolates that we deemed as pure cultures based on 16S rRNA
gene PCR and Sanger sequencing.
393
394 **Table 1. Cultivation statistics, including whole community viability estimates**

Site	Date	n	z	p	λ	V* (ASE)	CV	our model: estimated # wells with 1 cell (bootstrapped median: (xx-xx) 95% CI) if $V==1$ **	our model: V_{est}: min-max 95% CI ***	in situ salinity	Medium salinity	Medium	Study
CJ	Sep 2014	460	15	0.033	1.27	2.6 (0.67)	0.259	164 (144-185)	1.5-4.2	24.6	34.8	JWAMPFe	(35)
ARD	Nov 2014	460	1	0.002	1.5	0.1 (0.15)	1.451	154 (134-174)	0.1-0.7	1.72	34.8	JW1	(35)
JLB	Jan 2015	460	61	0.133	1.96	7.3 (0.93)	0.127	127 (109-146)	5.6-9.2	26.0	34.8	JW1	(35)
FWC†	Mar 2015	460	301	0.654	2	53.1 (3.2)	0.06	125 (106-143)	47.1-59.7	5.39	5.79	JW4	(35)
LKB	June 2015	460	15	0.033	1.8	1.8 (0.48)	0.266	137 (118-156)	1.1-3.0	2.87	5.79	JW4	(35)
Tbon2	Aug 2015	460	41	0.089	1.56	6.0 (0.93)	0.156	151 (132-171)	4.3-8.1	14.2	11.6	JW3	(35)
CJ2	Oct 2015	460	61	0.133	2	7.1 (0.91)	0.128	125 (106-143)	5.6-9.1	22.2	23.2	JW2	(35)
FWC2†	Apr 2016	460	403	0.876	2	104.4 (6.2)	0.059	125 (106-143)	>92.3	20.9	23.2	JW2	This study
ARD2c	Jun 2016	460	7	0.015	2	0.8 (0.29)	0.362	125 (106-143)	0.3-1.5	0.18	1.45	JW5	This study
JLB2c†	May 2016	460	103	0.224	2	12.7 (1.25)	0.099	125 (106-143)	10.3-15.4	6.89	5.79	JW4	This study
LKB2	Jul 2016	460	39	0.085	2	4.4 (0.71)	0.161	125 (106-143)	3.2-6.0	2.39	1.45	JW5	This study
Tbon3	Jul 2016	460	78	0.17	2	9.3 (1.05)	0.113	125 (106-143)	7.4-11.5	17.7	34.8	MWH1	This study
CJ3	Sep 2016	460	69	0.15	2	8.1 (0.98)	0.121	125 (106-143)	6.4-10.2	23.7	23.2	MWH2	This study
FWC3	Nov 2016	460	27	0.059	2	3.0 (0.58)	0.194	125 (106-143)	2.0-4.4	18.0	23.2	MWH2	This study
ARD3	Dec 2016	460	58	0.126	2	6.7 (0.89)	0.132	125 (106-143)	5.2-8.6	3.72	1.45	MWH5	This study
JLB3†	Jan 2017	460	146	0.317	2	19.1 (1.59)	0.083	125 (106-143)	16.1-22.4	12.4	11.6	MWH3	This study
LKB3	Feb 2017	460	38	0.083	2	4.3 (0.70)	0.163	125 (106-143)	3.1-5.8	3.55	1.45	MWH5	This study

395 *Viability according to equation 1, above. Asymptotic Standard Error (ASE) is presented in parentheses.

 **Based on 9,999 bootstraps.
397 ***Based on 9,999 bootstraps tested at viability increments of 0.1%.
398 †Experiments where a subset of positive wells were transferred.
399 FWC2 shows the advantage of our method over equation 1 for extreme values.
400
Phylogenetic and geographic novelty of our isolates
The 328 isolates belonged to three Phyla: *Proteobacteria* (n = 319), *Actinobacteria* (n = 8), and
Bacteroidetes (n = 1) (Figs. S1-S5). We placed these isolates into 55 groups based on their
positions within 16S rRNA gene phylogenetic trees (Figs. S1-S5) and as a result of having ≥
94% 16S rRNA gene sequence identity to other isolates. We applied a nomenclature to each
group based on previous 16S rRNA gene database designations and/or other cultured
representatives (Fig. 1, Table S1). Isolates represented eight putatively novel genera with ≤
94.5% 16S rRNA gene identity to a previously cultured representative: the *Actinobacteria* acIV
subclades A and B, and one other unnamed *Actinobacteria* group; an undescribed
Acetobacteraceae clade (*Alphaproteobacteria*); the freshwater SAR11 LD12 (*Candidatus*
Fonsibacter ubiquis (29)); the MWH-UniPo and an unnamed *Burkholderaceae* clade
(*Betaproteobacteria*); and the OM241 *Gammaproteobacteria* (Fig. 1, Table S1). Seven
additional putatively novel species were also cultured (between 94.6 and 96.9% 16S rRNA gene
sequence identity) in unnamed *Commamonadaceae* and *Burkholderiales* clades
(*Betaproteobacteria*); the SAR92 clade and *Pseudohonigella* genus (*Gammaproteobacteria*); and
unnamed *Rhodobacteraceae* and *Bradyrhizobiaceae* clades, as well as *Maricaulis* spp.
(*Alphaproteobacteria*) (Fig. 1). LSUCC isolates belonging to the groups BAL58
Betaproteobacteria (Fig. S4), OM252 *Gammaproteobacteria*, HIMB59 *Alphaproteobacteria*,
and what we designated the LSUCC0101-type *Gammaproteobacteria* (Fig. S5) had close 16S
rRNA gene matches to other cultivated taxa at the species level, however, none of those
previously cultivated organisms have been formally described (Fig. 1). The OM252, BAL58, and
MWH-UniPo clades were the most frequently cultivated, with 124 of our 328 isolates belonging
to these three groups (Table S1). In total, 73 and 10 of the 328 isolates belonged in putatively
novel genera and species, respectively. We estimated that at least 310 of these isolates were
geographically novel, being the first of their type cultivated from the nGOM (Fig. 2). This
included isolates from cosmopolitan groups like SAR11 subclade IIIa, OM43
Betaproteobacteria, SAR116, and HIMB11-type "Roseobacter" spp.. Cultivars from Vibrio sp.
and Alteromonas sp. were the only two groups that contained hits to organisms isolated from the
GoM.
430
Natural abundance of isolates
We matched LSUCC isolate 16S rRNA gene sequences with both operational taxonomic units
(OTUs) and amplicon single nucleotide variants (ASVs) from bacterioplankton communities to
assess the relative abundances of our isolates in the coastal nGOM waters that served as inocula.
While OTUs provide a broad group-level designation (97% sequence identity), this approach can
artificially combine multiple ecologically distinct taxa (95). Due to higher stringency for
defining a matching 16S rRNA gene, ASVs can increase the confidence that our isolates
represent environmentally relevant organisms (74, 96). However, while many abundant
oligotrophic bacterioplankton clades such as SAR11 (29, 97), OM43 (40, 41), SAR116 (98), and
Sphingomonas spp. (99) have a single copy of the rRNA gene operon, other taxa can have
multiple rRNA gene copies (97, 100), complicating ASV analyses. Since we could not *a priori*

rule out multiple rRNA gene operons for novel groups with no genome sequenced
representatives, we used both OTU and ASV approaches.
In total, we obtained at least one isolate from 40 of the 777 OTUs and 71 of the 1,323
ASVs observed throughout the three-year dataset. 43% and 26% of LSUCC isolates matched the
top 50 most abundant OTUs (median relative abundances, all sites, from 0.11-8.1%; Fig. S6A)
and ASVs (mean relative abundances, all sites, from 0.11-3.8%; Fig. S6B), respectively, across
all sites and samples. Microbial communities from all collected samples clustered into two
groups corresponding to those inhabiting salinities below 7 and above 12, and salinity was the
primary environmental driver distinguishing community beta diversity (OTU: R^2=0.88, P=0.001,
ASV: R^2=0.89, P =0.001). As part of the cultivation strategy after the first five experiments, we
used a suite of five media differing by salinity and matched the experiment with the medium that
most closely resembled the salinity at the sample site. Consequently, our isolates matched
abundant environmental groups from both high and low salinity regimes. At salinities above
twelve, LSUCC isolates matched 13 and 14 of the 50 most abundant OTUs and ASVs,
respectively (Figs. 3A, 3B; Table S1). These taxa included the abundant SAR11 subclade IIIa.1,
HIMB59, HIMB11-type "Roseobacter", and SAR116 *Alphaproteobacteria*; the OM43
Betaproteobacteria; and the OM182 and LSUCC0101-type *Gammaproteobacteria*. At salinities
below seven, 10 and 9 of the 50 most abundant OTUs and ASVs, respectively, were represented
by LSUCC isolates, including one of most abundant taxa in both cluster sets, SAR11 LD12
(Figs. 3C, 3D). Some taxa, such as SAR11 IIIa.1 and OM43, were among the top 15 most
abundant taxa in both salinity regimes (Fig. 3, Table S1), suggesting a euryhaline lifestyle. In
fact, our cultured SAR11 IIIa.1 ASV7471 was the most abundant ASV in the aggregate dataset
(Fig. S6).
Overall, this effort isolated taxa representing 18 and 12 of the top 50 most abundant
OTUs and ASVs, respectively (Table 2, Fig. S6). When looking at different median relative
abundance categories of > 1%, 0.1% - 1%, and < 0.1%, isolate OTUs were distributed across
those categories in the following percentages: 15%, 20%, and 27%; isolate ASVs were
distributed accordingly: 4%, 26%, and 37% (Table 2). Isolates with median relative abundances
of < 0.1%, such as *Pseudohongiella* spp., *Rhodobacter* spp., and *Bordetella* spp., would
canonically fall within the rare biosphere (101) (Table S1). A number of isolates did not match
any identified OTUs or ASVs (38% and 33% of LSUCC isolates when compared to available
OTUs and ASVs, respectively), either because their matching OTUs/ASVs were below our
thresholds for inclusion (at least two reads from at least two sites), or because they were below
the detection limit from our sequencing effort (Table 2). Thus, 43% and 30% of our isolates
belonged to OTUs and ASVs, respectively, with median relative abundances > 0.1%.
477
478 **Table 2. Median relative abundances (*r*) of cultured OTUs and ASVs across all samples**

	In top 50 ranks	*r* > 1%	1% - 0.1% *r*	*r* < 0.1%	Not detected
OTUs	18 (140 isolates)	50 isolates (15%)	90 isolates (27%)	65 isolates (20%)	123 isolates (38%)
ASVs	12 (84 isolates)	13 isolates (4%)	84 isolates (26%)	122 isolates (37%)	109 isolates (33%)

479
Modeling DTE cultivation
An enigma that became immediately apparent through a review of our data was the absence of an
obvious relationship between the abundance of a given taxon in the inoculum and the frequency
of obtaining an isolate of the same type from a DTE cultivation experiment (Figs. S7, S8). For
example, although we could culture SAR11 LD12 over a range of media conditions (29), and the
matching ASV had relative abundances of > 5% in six of our seventeen experiments (Fig. 4), we

only isolated one representative (LSUCC0530). In an ideal DTE cultivation experiment where
cells are randomly subsampled from a Poisson-distributed population, if the medium were
sufficient for a given microorganism's growth, then the number of isolates should correlate with
that microorganism's abundance in the inoculum. However, a qualitative examination of several
abundant taxa that grew in our media, some of which we cultured on multiple occasions,
revealed no clear pattern between abundance and isolation success (Fig. 4). Considering that
medium composition was sufficient for cultivation of these organisms on at least some
occasions, we hypothesized that cultivation frequency may reflect viability differences in the
populations of a given taxon. Thus, we decided to model cultivation frequency in relationship to
estimated abundances in a way that could generate estimates of cellular viability. We also hoped
that this information might help us inform experimental design and make DTE cultivation efforts
more predictable (59).
Previously, Don Button and colleagues developed a statistical model for viability (V) of
cells in the entire sample for a DTE experiment (33):
500
501   (1) $V = -(ln(1-p)/\lambda)$
502
Where p is the proportion of wells or tubes, n, with growth, z, ($p=z/n$) and λ is the estimated
number of cells inoculated per well (the authors used X originally). The equation uses a Poisson
distribution to account for the variability in cell distribution within the inoculum and therefore
the variability in the number of wells or tubes receiving the expected number of cells. We and
others have used this equation in the past (26, 28, 35) to evaluate the efficacy of our cultivation
experiments in the context of commonly cited numbers for cultivability using agar-plate based
methods (13, 17, 102).
While Equation 1 is effective for its intended purpose, it has a number of drawbacks that
limit its utility for taxon-specific application: 1) If $p=1$, i.e. all wells are positive, then the
equation is invalid; 2) At high values of p and low values of λ, estimates of V can exceed 100%
(Table 2); 3) Accuracy of viability, calculated by the asymptotic standard error, $ASEV$, or the
coefficient of variation, CVV, was shown to be non-uniform across a range of λ, with greatest
accuracy when true viability was ~10% (33). Thus, low viability, low values of λ, and small
values of n were found to produce unreliable results; 4) If $p=0$, i.e. no positive wells are
observed, estimates of viability that could produce 0 positive wells cannot be calculated. In
addition, 5) Button's original model assumes that a well will only produce a pure culture if the
inoculated well contains one cell. By contrast, in low diversity samples, samples dominated by a
single taxon, or experiments evaluating viability from axenic cultures across different media, a
limitation that only wells with single cells are axenic will underestimate the expected number of
pure wells. To overcome these limitations, we developed a Monte Carlo simulation model that
facilitates the incorporation of relative abundance data from complementary community profiling
data (e.g. 16S rRNA gene amplicons) to calculate the likelihood of positive wells, pure wells,
and viability at a taxon-specific level, based on the observed number of wells for we obtained an
isolate of a particular taxon (Fig. 5). By employing a Monte Carlo approach, our model is robust
across all values of p and n with uniform prediction accuracy, and we can estimate the accuracy
of our prediction within 95% confidence intervals (CI). Furthermore, the width of 95% CI
boundaries of viability, as well as the expected number of positive and pure wells, are entirely
controllable and dependent only on available computational capacity for bootstrapping (i.e.,
these can be improved with more bootstrapping, but at greater computational cost). When zero

positive wells are observed experimentally, our approach enables estimation of a maximum
viability that could explain such an observation by identifying the range of variability values for
which zero resides within the bootstrapped 95% CI. Finally, the ability to calculate the viability
of the entire community, as in Equation 1, is retained simply by estimating viability using a
relative abundance of one.
We compared our model to that of Button et al. for evaluating viability from whole
community experimental results, similarly to previous reports (26, 28, 35) (Table 1). Our
viability estimates (V_{est}) generally agreed with those using the Button et al. calculation, but we
have now provided 95% CI to depict the maximum and minimum viability that would match the
returned positive well distribution, as well as maximum and minimum values for the number of
wells that ought to have contained a single cell. Maximum V_{est} ranged from 1.1% to > 92.3%
depending on the experiment, with a median V_{est} across all our experiments of 8.6% (Table 1). In
one case, the extremely high value (FWC2) was better handled by our model compared to
equation 1, because it did not lead to a viability estimation greater than 100%. FWC and FWC2
represent V_{est} outliers compared with the entire dataset (maximums of 59.7% and > 92.3%,
respectively; Table 1). We believe these high numbers most likely resulted from underestimating
the number of cells inoculated into each well because of the use of microscopy, clumped cells, or
pipet error, thus increasing the estimated viability (35).
550
Isolate-specific viability estimates
Our new model also facilitates taxon-specific viability estimates. Cultivation efficacy was
evaluated for 71 cultured ASVs (219 isolates) across 17 sites (1,207 pairwise combinations) by
comparing the number of observed pure wells to those predicted by the Monte Carlo simulation
using 9,999 bootstraps, 460 wells per experiment, and an assumption that all cells were viable
(i.e. $V = 100\%$). In total, for 1,158 out of 1,207 pairwise combinations (95.9%) the observed
number of pure wells fell within the 95% CI of data simulated at matching relative abundance
and inoculum size, suggesting that these two parameters alone could explain the observed
cultivation success for most taxa (Table S1). 1,059 out of these 1,158 combinations (91%)
recorded zero observed wells, but with a maximum relative abundance of 2.8% within these
combinations, a score of zero fell within predicted 95% CI of simulations with 460 wells.
Sensitivity analysis showed that with 460 wells per experiment, an observation of zero pure
wells falls below the 95% confidence intervals lower-bound (and is thus significantly depleted to
enable viability to be estimated) for taxa with relative abundances of 2.3%, 2.9% and 4.5% for
inoculum sizes of one, two and three cells per well, respectively (Fig. S9). In fact, modeling DTE
experiments from 92 wells to 9,200 wells per experiment showed that for taxa comprising ~1%
of a microbial community 1,104 wells (or 12 plates at 92 wells per plate), 1,380 wells (15 plates)
and 2,576 wells (28 plates) were required to be statistically likely to recover at least one positive,
pure well using inocula of one, two or three cells per well, respectively, with $V = 100\%$ (Fig. S9).
A small, but taxonomically relevant minority (49 out of 1,207) of pairwise combinations
had a number of observed pure wells that fell outside of the simulated 95% CI with $V = 100\%$
(Fig. 6). Of these, 28 had either one, two, or three more observed pure wells than the upper 95%
CI (Table S1), suggesting cultivability *higher* than expected based purely a model capturing the
interaction between a Poisson-distributed inoculum and a binomially-distributed relative
abundance, with $V = 100\%$. However, the deviance from the expected number of positive wells
for those above the 95% CI was limited to three or fewer cells, meaning that we only obtained 1-
3 more isolates than expected (Table S1). On the other hand, those organisms that we isolated

less frequently than expected showed greater deviance. 21 out of the 49 outliers had lower than expected cultivability (Fig. 6). These taxa had relative abundances ranging from 2.7% to 14.5%, but recorded only 0, 1, or 2 isolates. In the most extreme case, ASV7629 (SAR11 LD12) at Site ARD2c comprised 14.5% of the community but recorded no observed pure wells, compared to expected number of 13-30 isolates (95% CI) predicted by the Monte Carlo simulation. All the examples of taxa that were isolated less frequently than expected given the assumption of V = 100% belonged to either SAR11 LD12, SAR11 IIIa.1, or one particular OM43 ASV (7241) (Fig. 6).

We used our model to calculate estimated viability (V_{est}) for these organisms based on their cultivation frequency at sites where the assumption of V = 100% appeared violated (Table 3). Using the extreme example of SAR11 LD12 ASV7629 at Site ARD2c, simulations across a range of V indicated that a result of zero positive wells fell within 95% of simulated values when the associated taxon $V_{est} \leq 15\%$. When considering all anomalous cultivation results, LD12 had estimated maximum viabilities that ranged up to 55% (Table 3). OM43 (ASV7241) estimated maximum viabilities ranged from 52-80%, depending on the site, and similarly, SAR11 IIIa.1 ranged between 22-82% viability (Table 3).

Table 3. Estimated viabilities for taxa cultivated less frequently than expected

ASV	Group	Site	r*	n	z	λ	Estimated # wells with 1 cell (bootstrapped median: (xx-xx) 95% CI) if V==1 **	V_{est}: min-max 95% CI based on cultivation results***
7241	OM43	ARD3	0.03	460	0	2	4 (1-9)	0.1-80
7241	OM43	FWC†	0.04	460	0	2	5 (1-9)	0.1-77
7241	OM43	JLB	0.05	460	0	1.96	7 (2-12)	0.1-52
7471	SAR11 IIIa.1	ARD3	0.11	460	0	2	15 (8-23)	0.1-22
7471	SAR11 IIIa.1	CJ	0.03	460	0	1.27	4 (1-9)	0.1-82
7471	SAR11 IIIa.1	FWC3	0.07	460	2	2	9 (4-15)	2.5-80
7471	SAR11 IIIa.1	JLB	0.05	460	0	1.96	6 (2-11)	0.1-59
7471	SAR11 IIIa.1	JLB2c†	0.08	460	0	2	11 (5-18)	0.1-31
7471	SAR11 IIIa.1	JLB3†	0.04	460	0	2	5 (1-9)	0.1-74
7471	SAR11 IIIa.1	LKB	0.05	460	0	1.8	6 (2-12)	0.1-55
7471	SAR11 IIIa.1	LKB2	0.04	460	0	2	5 (1-9)	0.1-77
7471	SAR11 IIIa.1	LKB3	0.09	460	0	2	11 (6-19)	0.1-30
7471	SAR11 IIIa.1	TBON2	0.04	460	0	1.56	6 (2-12)	0.1-56
7471	SAR11 IIIa.1	TBON3	0.04	460	0	2	5 (1-10)	0.1-73
7629	SAR11 LD12	ARD	0.11	460	0	1.5	18 (10-27)	0.1-20
7629	SAR11 LD12	ARD2c	0.15	460	0	2	21 (13-30)	0.1-15
7629	SAR11 LD12	ARD3	0.05	460	0	2	7 (2-12)	0.1-53
7629	SAR11 LD12	FWC†	0.09	460	0	2	12 (6-19)	0.1-28
7629	SAR11 LD12	LKB	0.05	460	0	1.8	7 (2-13)	0.1-51
7629	SAR11 LD12	LKB2	0.08	460	1	2	11 (5-18)	0.3-49
7629	SAR11 LD12	LKB3	0.06	460	0	2	7 (3-13)	0.1-48

*Fractional relative abundance
**Based on 9,999 bootstraps.
***Based on 9,999 bootstraps tested at viability increments of 0.1%.
†Experiments where a subset of positive wells were transferred.

Discussion

This work paired 17 DTE cultivation experiments with cultivation-independent assessments of microbial community structure in source waters to evaluate cultivation efficacy. We generated 328 new bacterial isolates representing 40 of the 777 OTUs and 71 of the 1,323 ASVs observed across all samples from which we inoculated DTE experiments. Stated another way, we successfully cultivated 5% of the total three year bacterioplankton community observed via either OTU or ASV analyses. A large fraction of our isolates (43% of cultured OTUs, 30% of

cultured ASVs) represented taxa present at median relative abundances > 0.1%, with 15% and
4% of cultured OTUs and ASVs, respectively, at median abundances > 1%. 140 of our isolates
matched the top 50 most abundant OTUs, and 84 isolates matched the top 50 most abundant
ASVs.
This campaign led to the first isolations of the abundant SAR11 LD12 and *Actinobacteria*
acIV; the second isolate of the HIMB59 *Alphaproteobacteria*; and new genera within the
Acetobacteraceae, *Burkholderiaceae*, OM241 and LSUCC0101-type *Gammaproteobacteria*, and
MWH-UniPo *Betaproteobacteria*; thereby demonstrating again that continued DTE
experimentation leads to isolation of previously uncultured organisms with value for aquatic
microbiology. We have also added a considerable collection of isolates to previously cultured
groups like OM252 *Gammaproteobacteria*, BAL58 *Betaproteobacteria*, and HIMB11-type
"Roseobacter" spp., and the majority of our isolates represent the first versions of these types of
taxa from the Gulf of Mexico, adding comparative biogeographic value to these cultures.
Our viability model improved upon the statistical equation developed by Button and
colleagues (33) to extend viability estimates to individual taxa within a mixed community and
provide 95% CI constraining those viability estimates. We cultured several groups of organisms
abundant enough to evaluate viability with 460 wells (Figs. 6, S9). The fact that these organisms
were successfully cultured at least once meant that we could reasonably assume that the medium
was sufficient for growth. Some taxa were cultivated more frequently than expected (Fig. 6). We
explore two possible explanations for this phenomenon- errors in quantification and variation in
microbial cell organization. Any systematic error that led to underestimating the abundance of an
organism would have correspondingly resulted in our underestimating the number of wells in
which we would expect to find a pure culture of that organism. Such underestimations could
come from primer biases associated with amplicon sequencing (69, 70, 103), but we do not know
if those protocols specifically underestimate the OM252, MWH-UniPo, and HIMB11-type taxa
cultured more frequently than expected (Fig. 6). However, due the low number of expected
isolates in these groups, and the small deviances in actual isolates from those expected numbers
(within 1-3 isolates compared to expected values), the biases inherent in the relative abundance
estimations for these taxa were probably small. Furthermore, one of the microorganisms isolated
more frequently than expected matched the OM43 ASV1389 (Fig. S6), whereas another OM43
ASV (7241) was cultivated less frequently than expected (see below), meaning that if primer
bias were the cause of this discrepancy, it would have to be operating differently on very closely
related organisms.
There exists a biological explanation for why some isolates might have been cultured
more frequently than expected: clumped cells. If cells of any given taxon in nature grew in small
clusters, then the number of cells we added to a well was greater than expected based on a
Poisson distribution. Furthermore, the model assumes that each cell is independent, and that the
composition of a subset of cells is only a function of the relative abundance of the taxon in the
community. Within a cluster of cells, this assumption is violated as the probability of cells being
from the same taxon is higher. Thus, the model will underestimate the probability of a well being
pure and therefore underestimate the number of pure wells likely to be observed within an
experiment, leading to a greater number of isolates than expected. Future microscopy work could
examine whether microorganisms such as OM252 and MWH-UniPo form small clusters *in situ*
and/or in pure culture, and whether this phenomenon may be different for different ASVs of
OM43, or if clumping may be a transient phenotype.

We also identified three taxa- SAR11 LD12, SAR11 subclade IIIa.1, and the
aforementioned OM43 ASV7241- that were isolated much less frequently than expected based
on their abundances (Fig. 6, Table 3). This could mean that our assumption of $V = 100\%$ was
incorrect, or that, in contrast to the taxa that were cultured more frequently than expected
(above), our methods had biases that *over*estimated the abundance of these organisms, thereby
over-inflating the expected number of isolates. We used the modified 515/806RB primers that
have been shown to be much more accurate in quantifying SAR11 compared to FISH than the
original 515/806 primers (within $6\% \pm 4\%$ SD), and this protocol almost always underestimates
SAR11 abundances (69). This suggests that our expected number of isolates may have actually
been underestimated, our cultivation success was poorer than we measured, and therefore we
may be overestimating viability for the SAR11 taxa. Other sources of systematic error that might
impinge on successful transfers, and thereby reduce our recovery, include sensitivity to pipette
tip and/or flask material. However, the fact that these taxa were sometimes successfully isolated
means that if these mechanisms were impacting successful transfers, then their activity was less
than 100% efficient, which implies variations in subpopulation vulnerability that would be very
similar conceptually to variations in subpopulation viability.
Another possible source of error that could have resulted in lower than expected numbers
of isolates was the subset of experiments for which we did not transfer all positive wells due to
limitations in available personnel time (Tables 1 & 3). However, our selection criteria for the
subset of wells to transfer was based on flow cytometric signatures that would have encompassed
small cells like SAR11 (see Results), and in any case, there were many examples of lower than
expected recovery from other experiments where we transferred all positive wells (Table 3).
Therefore, these four experiments were unlikely to contribute major errors biasing our estimates
of viability for SAR11 LD12, SAR11 IIIa.1, and other small cells like OM43.
If we instead explore biological reasons for the lower than expected numbers of positive
wells in DTE experiments, a plausible explanation supported by the literature is simply that a
large fraction of the population is in some state of inactivity or at least not actively dividing
(104). Studies using uptake of a variety of radiolabelled carbon and sulfur sources have
demonstrated substantial fractions of SAR11 cells may be inactive depending on the population
(105–108). SAR11 cells in the northwest Atlantic and Mediterranean showed variable uptake of
labelled leucine (30-50% -(105, 106); ~25-55% (108, 109) and amino acids (34-61% (105, 106);
34-66% (105, 106). Taken in reverse, this means that up to 75% of the SAR11 population may
be inactive at any given time. In another study focused on brackish communities, less than 10%
of SAR11 LD12 cells took up labelled leucine and/or thymidine (107). While this was likely not
the ideal habitat for LD12 based on salinities above 6 (29, 107), this study supports the others
above that show substantial proportions of inactive SAR11 cells, the fraction of which may
depend on environmental conditions and other unknown factors. Bi-orthagonal non-canonical
amino acid tagging (BONCAT) shows similar trends for SAR11 (110). These data also match
general data indicating prevalent inactivity among aquatic bacterioplankton (104, 111–113).
Although labelled uptake methods do not directly measure rates of cell division, the
incorporation of these compounds requires active DNA replication or translation, which
represent an even more fundamental level of activity than cell division (114).
Why might selection favor low viability? One possibility is as an effective defense
mechanism against abundant viruses. Viruses infecting SAR11 have been shown to be extremely
abundant in both marine (115, 116) and freshwater (117) systems. Indeed, the paradox of high
viral abundances and high host abundances in SAR11 has led to a refining of negative density

dependent selection through Lokta-Volterra predator-prey dynamics (118) to include
heterogeneous susceptibility at the strain level (119, 120) and positive density dependent
selection through intraspecific proliferation of defense mechanisms (121). Activity of lytic
viruses infecting SAR11 *in situ* demonstrated that phages infecting SAR11 have lower ratios of
viral transcripts to host cells compared to other abundant taxa, and that observed abrupt changes
in these ratios suggest co-existence of several SAR11 strains with different life strategies and
phage susceptibility (122). Phenotypic stochasticity of phage receptor expression has been
shown to maintain a small proportion of phage-insensitive hosts within a population, enabling
coexistence of predator and prey without extinction (123). Phages adsorb to a vast array of
receptor proteins on their hosts, with many well-characterised receptors (e.g. OmpC, TonB,
BtuB, LamB) associated with nutrient uptake or osmoregulation (124). Selection therefore
favours phenotypes that limit receptor expression, with an associated fitness cost, particularly in
nutrient-limited environments. However, an alternative mechanism is possible if a population of
cells comprised a small number of viable cells, and a large number of non-viable or dormant
cells where metabolism is restricted but presentation of receptor proteins is retained. The
majority of host-virus encounters would occur with non-viable cells, and constrain viral
propagation through inefficient or failed infection, effectively acting as a sink for infectious
particles. Thus, the population as a whole can maintain expression of receptor proteins, while
reducing viral predation, with low-viability cells retaining the possibility of a return to normal
phenotypic growth if and when viral selection pressure abates. Similar systems of bacterial
phenotypic bistability associated with antibiotic tolerance and survival in harsh environments are
well documented (125, 126) and have been suggested as a possible mechanism for maintaining
long-term co-stability between the abundant phage φCrAss001 and its host in the human gut
(127).
Detailed measurements of dormancy in SAR11 and what types of cellular functions
become inactivated are part of our ongoing work. In the meantime, it is worth it to think about
the implications of a substantial proportion of non-dividing cells for our understanding of basic
growth dynamics. Studies attempting to measure SAR11 growth rates in nature have yielded a
wide range of results, ranging from 0.03-1.8 day^{-1} (97, 105, 108, 128–130). These span wider
growth rates than observed for axenic cultures of SAR11 (0.4-1.2 day^{-1}), but isolate-specific
growth ranges within that spread are much more constrained (29, 36, 49, 131, 132)). Conversion
factors for determining production from ^{3}H-leucine incorporation appear to be accurate for at
least Ia subclade members of SAR11 (133), so variations in growth rate estimates from
microradiography experiments likely have other explanations. It is possible that different strains
of SAR11 simply have variations in growth rate not captured by existing isolates. Another, not
mutually exclusive, possibility is that the differences in *in situ* growth rate estimates also reflect
variations in the proportion of actively dividing cells within the population. A simple model of
cell division with binary fission where only a subset of cells divide and non-dividing cells
persist, rather than die, can still yield logarithmic growth curves (Fig. S10) like those observed
for SAR11 in pure culture (29, 49, 134). However, this means that the division rate for the subset
of cells that are actively dividing is much higher than calculated when assuming 100% viability
in the population. Based on our estimated viability for SAR11 LD12 of 15-55%, to obtain our
previously calculated maximum division rate (0.5 day^{-1}) for the whole culture (29), the per-cell
division rate for only a subpopulation would span 2.48-0.79 day^{-1} (Fig. S10, Supplemental Text).
Verifying the proportion of SAR11 cells actively dividing in a culture may be challenging. Time-

lapse microscopy (135) offers an elegant solution if SAR11 can be maintained for the requisite time periods for accurate measurements in a microfluidic device.

In addition to identifying taxa whose isolation success suggested deviations from biological assumptions of single planktonic cells with 100% viability, the model also revealed the limitations of DTE cultivation in assessing viability depending on relative abundance (Fig. S9). We cannot ascertain whether any given taxon may violate an assumption of $V = 100\%$ unless we have enough wells to demonstrate that it grew in fewer wells than expected. For example, taxa at 1% of the microbial community require more than 1,000 wells before the lack of a cultured organism represents a statistically important negative event, rather than a taxon simply lacking sufficient abundance to ensure inclusion in a well within 95% CI. In our 460 well experiments, we could not resolve whether taxa may have had viabilities below 100% if they were less than 3% of the community for any given experiment (Fig. S9). Modeling DTE experiments showed that for experiments targeting rare taxa, lower inoculum sizes are favoured where selective media for enrichment is either unknown or undesirable. The exponential increase in the number of required wells with respect to the inoculum size is a function of a pure well requiring *all* cells within it to belong to the same taxon, assuming all cells are equally and optimally viable.

By providing taxon-specific predictions of viability from cultivation data, our model now facilitates an iterative process to improve experimental design and make cultivation more reliable. First, we use the cultivation success rates to determine for which taxa the assumption of 100% viability was violated. Second, we use the model to estimate viability for those organisms. Third, we use the viability and relative abundance data to determine, within 95% CI, the appropriate number of inoculation attempts required to isolate a new version of that organism. Using SAR11 LD12 as an example, given a relative abundance of 10%, and a viability of 15%, 800 DTE wells should yield four pure, positive wells (1-8 95% CI). This means that, for microorganisms that we know successfully grow in our media, we can now statistically constrain the appropriate number of wells required to culture a given taxon again. For organisms that were not abundant enough to estimate viability using the model, we can use a conservative viability assumption (e.g., 50% (111)) with which to base our cultivation strategy, thereby still reducing uncertainty about the experimental effort necessary to re-isolate one of these microorganisms.

Conclusions
This work has provided hundreds of new cultures for microbiological research, many among the most abundant members of the nGOM coastal bacterioplankton community. It also provides another demonstration of the effectiveness of sustained cultivation efforts for bringing previously uncultivated strains into culture. Our modeled cultivation results have generated compelling evidence for low viability within subpopulations of SAR11 LD12 and IIIa.1, as well as OM43 *Betaproteobacteria*. The prevalence of, and controls on, dormancy in these clades deserves further study. We anticipate that future work with larger DTE experiments will yield similar viability data about other groups of taxa with lower abundance, highlighting a valuable diagnostic application of DTE cultivation/modeling beyond the primary role in isolating new microorganisms. The integration of cultivation results, natural abundance data from inoculum communities, and DTE modeling represents an important step forward in quantifying the risk associated with DTE efforts to isolate valuable taxa from new sources, or repeating isolation from the same locations. We hope variations of this approach will be incorporated into wider community efforts to invest in culturing the uncultured.

790
791
792
793
794 **Acknowledgments**
795 We would like to thank Dr. Nancy Rabalais for her comments and edits on a very early draft of
796 this manuscript. Portions of this research were conducted with high-performance computing
797 resources provided by Louisiana State University (http://www.hpc.lsu.edu) and the University of
798 Southern California (http://hpc.usc.edu).
799
Funding Information
This work was supported by the Department of Biological Sciences at Louisiana State
University, a Louisiana Board of Regents grant to JCT (LEQSF(2014-17)-RD-A-06), a National
Academies of Science, Engineering, and Medicine Gulf Research Program Early Career
Fellowship to JCT, and the Dornsife College of Letters, Arts and Sciences at the University of
Southern California. BT was partially funded by NERC award NE/R010935/1 and by the Simons
Foundation BIOS-SCOPE program. The funders had no role in study design, data collection, and
interpretation, or the decision to submit the work for publication.
808
Author Contributions
MWH led sample collections, cultivation experiments, nucleic acid extraction, amplicon
sequencing, and analyses; VCL, DMP, JLW, and AML supported sample collections, cultivation
experiments, and nucleic acid extractions; MWH and JCT conducted cultivation comparisons
and phylogenetic analyses; BT developed the viability model and led the statistical analyses; CC
derived the cell-specific growth rate equations incorporating viability; JCT designed the study
and assisted with sample collections and model refinement; MWH, BT, and JCT led manuscript
writing; and all authors contributed to and reviewed the text.

822   **Figures**
823
824
825

826
827
Figure 1. Percent identity of LSUCC isolate 16S rRNA genes compared with those from other
isolates in NCBI ("Other", gray dots) or from the DTE culture collections IMCC (gold dots),
HTCC (blue dots), and HIMB (green dots). Each dot represents a pairwise 16S rRNA gene
comparison (via BLASTn). X-axis categories are groups designated according to ≥ 94%
sequence identity and phylogenetic placement (see Figs S1-S4). Above the graph is the 16S
rRNA gene sequence percent identity to the closest non-LSUCC isolate within a column. Groups
colored in red indicate those where LSUCC isolates represent putatively novel genera, whereas
orange indicates putatively novel species.
836
837
838
839

846
847
848 **Figure 2.** A global map of the isolation location of isolates from selected important aquatic
849 bacterioplankton clades. Circles represent LSUCC isolates, while triangles are non-LSUCC
850 isolates. Inset: a zoomed view of the coastal Louisiana region where LSUCC bacterioplankton
851 originated.

878
879
880

Figure 3. Rank abundances of the 50 most abundant taxa based on median relative abundance at salinities less than seven (top) and greater than twelve (bottom) for OTUs (A, C) and ASVs (B, D) across all sites. The boxes indicate the interquartile range (IQR) of the data, with vertical lines indicating the upper and lower extremes according to 1.5 x IQR. Horizontal lines within each box indicate the median. The data points comprising the distribution are plotted on top of the boxplots. The shade of the dot represents the salinity at the sample site (red-blue :: lower-higher), while the color of the box indicates broad taxonomic identity. LSUCC labels indicate OTUs and ASVs with at least one cultivated representative.

Figure 4. Relative abundance of ASVs within key taxonomic groups compared with salinity. ASV types are colored independently, and triangle points indicate experiments for which at least one isolate was obtained. Non-linear regression lines are provided as a visual aid for abundance trends.

923
924
925

Step 1: Simulate *n* wells

Step 2: Simulate inoculation of wells from Poisson distribution (λ=inoculum)

Step 3: Simulate likelihood of taxon from Binomial distribution (n = # cells in well, p = rel. abund)

Step 4: Count positive wells, taxon positive wells, pure wells and taxon pure wells

Step 5: For taxon pure wells, simulate likelihood of viability from Binomial distribution(n = # cells in well, p = viability)

Step 6: Count wells where viable cells >=1

Step 7: Bootstrap steps 1-6 *k* times at different levels of viability, *0 ≤ v ≤ 1*

Step 8: Identify *min, max* values of *v* where # observed wells falls within bootstrapped 95% CI

926
927
928   **Figure 5.** Graphical depiction of the viability model.
929
930

Figure 6. Actual vs. expected numbers of isolates. Each point represents the actual number of
isolates for every ASV/experiment pair compared to the expected number of isolates based on
our model assuming 100% viability. Colors represent the relationship to the model predictions:
green- relative abundance low enough whereby 0 isolates fell within the 95% CI; pink- isolation
results within the 95% CI for taxa with relative abundance ≥ 2.9%; orange- actual isolates >
maximum 95% CI for expected isolates; blue- actual isolates < minimum 95% CI for expected
isolates. Circle size is proportional to the deviation of the number of actual isolates from the
maximum (for orange) or minimum (for blue) 95% CI for expected isolates. The dotted line is
the 1:1 ratio. Notable taxa on the extremities of the actual and expected values are labeled.

954 **Supplemental Information**
955
956 **Supplemental Text**
957 Math for back calculating the per-cell division rates in Henson et al. 2018 assuming 15%--55%
958 viability.
959
960 Denote the initial cell count as X_0, and the cell count in n^{th} generation as X_n. If all the cells are
961 viable, per each generation, all the cells are doubling:
962

$$X_n = X_{n-1} \cdot 2 \tag{eq. S1.1}$$

963
964
965 And therefore, after n generations, the cell count X_n would be:

$$X_n = X_0 \cdot 2^n \tag{eq. S1.2}$$

966
967
968 When not all the cells are viable, for each generation, instead of simply doubling the whole
969 population, only a portion of the cells (denoted as $V < 1$) replicates. In this case:
970

$$X_n = X_{n-1} \cdot (1 + V) \tag{eq. S1.3}$$

971
972
973 Here we can see that (eq. 3.1) is actually a special case of (eq. 3.3), when $V = 1$,
974

$$X_n = X_{n-1} \cdot (1 + 1) = X_{n-1} \cdot 2$$

975
976
977 When $V < 1$, after n generations:

$$X_n = X_0 \cdot (1 + V)^n \tag{eq. S1.4}$$

978
979
980 Denoting the total time to get n generations as t, which means the number of generations per unit
981 time, also known as the per-cell division rate, is $\mu = n/t$. Rewriting (eq. 3.4), we have:
982

$$X_n = X_0 \cdot (1 + V)^{\mu \cdot t} = X_0 \cdot 2^{\mu \cdot log_2(1+V) \cdot t} \tag{eq. S1.5}$$

983
984
985 According to (eq. 3.5), a cell population of V viability and μ per cell division rate, the population
986 division rate is:

$$\mu_{population} = \mu \cdot log_2(1 + V) \tag{eq. S1.6}$$

987
988
989 Henson et al. 2018 show that the optimal population division rate of SAR11 LD12 is
990 $\mu_{population} = 0.5 \, day^{-1}$, assuming the viability of LD12 is $V = 0.15$, by plugging in these two
991 numbers to (eq. S1.6), we can get the per-cell division rate of LD12 is:

$$\mu = \mu_{population}/log_2(1 + V)$$
$$= 0.5 \, day^{-1}/log_2(1 + 0.15)$$
$$= 0.5 \, day^{-1}/0.20163$$
$$= 2.48 \, day^{-1}$$

992
993
994
995
996 If assuming $V = 0.55$, $\mu = 0.79 \, day^{-1}$. Therefore, given the range of the viability of LD12 is
997 between 15%--55%, the corresponding per cell division rate is between 0.79 divisions to 2.48
998 divisions per day.

Supplemental Table
Supplemental Table S1 is a spreadsheet, Table_S1.xlsx. This includes MWH media recipes, ASV and OTU tables, taxonomic and relative abundance information for all isolates, NMDS information, biogeochemical data for the samples, and the BLAST results of isolate hits. Available at https://doi.org/10.6084/m9.figshare.12142113.

Supplemental Figures
Figure S1. 16S rRNA gene phylogeny of LSUCC isolates in the Phylum *Actinobacteria*. Scale bar represents 0.01 changes per position. Values at internal nodes indicate bootstrap values (n = 1000).

Figure S2. 16S rRNA gene phylogeny of LSUCC isolates in the Class *Alphaproteobacteria*. Scale bar represents 0.01 changes per position. Values at internal nodes indicate bootstrap values (n = 1000).

Figure S3. 16S rRNA gene phylogeny of LSUCC isolates in the Phylum *Bacteroidetes*. Scale bar represents 0.01 changes per position. Values at internal nodes indicate bootstrap values (n = 1000).

Figure S4. 16S rRNA gene phylogeny of LSUCC isolates in the Class *Betaproteobacteria*. Scale bar represents 0.01 changes per position. Values at internal nodes indicate bootstrap values (n = 1000).

Figure S5. 16S rRNA gene phylogeny of LSUCC isolates in the Class *Gammaproteobacteria*. Scale bar represents 0.01 changes per position. Values at internal nodes indicate bootstrap values (n = 1000).

These trees have been attached at the end of the manuscript for greater visibility

1045
1046
1047
1048

1049
1050
1051
1052
Figure S6. Rank abundance plot of the top 50 most abundant (A) OTUs and (B) ASVs across all
sites. The boxes indicate the interquartile range (IQR) of the data, with vertical lines indicating
the upper and lower extremes according to 1.5 x IQR. Horizontal lines within each box indicate
the median. The underlying data points are each individual OTU's sample relative abundances.
The shade of the dot represents the salinity, while the color of the box is the Phylum of the OTU.
OTUs with cultured representatives from the LSU culture collection are labeled with one
LSUCC isolate.

1087
1088
1089
1090 **Figure S7.** Relative abundance of cultivated OTUs according to site salinity. The color of the
1091 line represents the different OTUs classified within the LSUCC group. Non-linear regressions
1092 have been added for reference.

1104
1105
1106
1107

1108
1109
1110
1111 **Figure S8.** Relative abundance of ASVs with a cultured LSUCC representative according to site
1112 salinity. The color of the line represents the different ASVs classified within the LSUCC group.
1113 Non-linear regressions have been added for reference. Arrows represent sites where isolates were
1114 cultivated.

1132
1133
1134
1135
1136 **Figure S9.** Sensitivity testing of the viability model to determine minimum numbers of wells
1137 required to overcome false negatives associated with differing levels of taxon relative
1138 abundance. Different inoculum densities (λ) are depicted with colors. The dotted line indicates
1139 460 wells which corresponded to the experiments in this study.

Figure S10. Logarithmic growth with different percentages of cell viability. Under different cell viabilities, such as 100%, 55% and 15% shown in (A), the culture can have logarithmic growth in the population size if we assume no cell death. It takes around 25 generations to have 10^7 fold expansion of population size when all cells are active (100% viable), whereas when 15% of the population is actively duplicating, it takes around 115 generations to reach the same population size expansion. B) For the SAR11 LD12 strain LSUCC0530, the previously measured population doubling rate was 0.5/day (29). If all the cells are active (100% viability), each cell within the population will double 0.5 times per day (Fit 1 in B). However, with 15% viability, the actual active cell doubling rate is 2.48 doublings day^{-1} (Fit 3 in B).

References

1. Metcalf WW, Griffin BM, Cicchillo RM, Gao J, Janga SC, Cooke HA, Circello BT, Evans BS, Martens-Habbena W, Stahl DA, van der Donk WA. 2012. Synthesis of methylphosphonic acid by marine microbes: a source for methane in the aerobic ocean. Science 337:1104–1107.

2. Carini P, White AE, Campbell EO, Giovannoni SJ. 2014. Methane production by phosphate-starved SAR11 chemoheterotrophic marine bacteria. Nat Commun 5:4346.

3. Karl DM, Beversdorf L, Björkman KM, Church MJ, Martinez A, Delong EF. 2008. Aerobic production of methane in the sea. Nat Geosci 1:473–478.

4. Steindler L, Schwalbach MS, Smith DP, Chan F, Giovannoni SJ. 2011. Energy starved Candidatus Pelagibacter ubique substitutes light-mediated ATP production for endogenous carbon respiration. PLoS One 6:e19725.

5. Gómez-Consarnau L, González JM, Coll-Lladó M, Gourdon P, Pascher T, Neutze R, Pedrós-Alió C, Pinhassi J. 2007. Light stimulates growth of proteorhodopsin-containing marine Flavobacteria. Nature 445:210–213.

6. Daims H, Lebedeva EV, Pjevac P, Han P, Herbold C, Albertsen M, Jehmlich N, Palatinszky M, Vierheilig J, Bulaev A, Kirkegaard RH, von Bergen M, Rattei T, Bendinger B, Nielsen PH, Wagner M. 2015. Complete nitrification by Nitrospira bacteria. Nature 528:504–509.

7. Lam KS. 2006. Discovery of novel metabolites from marine actinomycetes. Curr Opin Microbiol 9:245–251.

8. Jensen PR, Williams PG, Oh D-C, Zeigler L, Fenical W. 2007. Species-specific secondary metabolite production in marine actinomycetes of the genus Salinispora. Appl Environ Microbiol 73:1146–1152.

9. Nett M, Erol Ö, Kehraus S, Köck M, Krick A, Eguereva E, Neu E, König GM. 2006. Siphonazole, an Unusual Metabolite from Herpetosiphon sp. Angew Chem Int Ed 45:3863–3867.

10. Ling LL, Schneider T, Peoples AJ, Spoering AL, Engels I, Conlon BP, Mueller A, Schäberle TF, Hughes DE, Epstein S, Jones M, Lazarides L, Steadman VA, Cohen DR, Felix CR, Fetterman KA, Millett WP, Nitti AG, Zullo AM, Chen C, Lewis K. 2015. A new antibiotic kills pathogens without detectable resistance. Nature 517:455–459.

11. Lloyd KG, Steen AD, Ladau J, Yin J, Crosby L. 2018. Phylogenetically Novel Uncultured Microbial Cells Dominate Earth Microbiomes. mSystems 3:e00055–18.

12. Hug LA. 2018. Sizing Up the Uncultured Microbial Majority. mSystems 3:e00185–18.

13. Steen AD, Crits-Christoph A, Carini P, DeAngelis KM, Fierer N, Lloyd KG, Thrash JC. 2019. High proportions of bacteria and archaea across most biomes remain uncultured.

ISME J 13:3126–3130.

14. Overmann J, Abt B, Sikorski J. 2017. Present and Future of Culturing Bacteria. Annu Rev
Microbiol 71:711–730.

15. Rappé MS. 2013. Stabilizing the foundation of the house that 'omics builds: the evolving
value of cultured isolates to marine microbiology. Curr Opin Microbiol 16:618–624.

16. Carini P. 2019. A "Cultural" Renaissance: Genomics Breathes New Life into an Old Craft.
mSystems 4:e00092–19.

17. Staley JT, Konopka A. 1985. Measurement of in situ activities of nonphotosynthetic
microorganisms in aquatic and terrestrial habitats. Annu Rev Microbiol 39:321–346.

18. Amann RI, Ludwig W, Schleifer KH. 1995. Phylogenetic identification and in situ detection
of individual microbial cells without cultivation. Microbiol Rev 59:143–169.

19. Hahn MW, Koll U, Schmidt J. 2019. Isolation and Cultivation of Bacteria, p. 313–351. *In*
Hurst, CJ (ed.), The Structure and Function of Aquatic Microbial Communities. Springer
International Publishing, Cham.

20. Kaeberlein T, Lewis K, Epstein SS. 2002. Isolating "uncultivable" microorganisms in pure
culture in a simulated natural environment. Science 296:1127–1129.

21. Zengler K, Toledo G, Rappe M, Elkins J, Mathur EJ, Short JM, Keller M. 2002. Cultivating
the uncultured. Proc Natl Acad Sci U S A 99:15681–15686.

22. Steinert G, Whitfield S, Taylor MW, Thoms C, Schupp PJ. 2014. Application of Diffusion
Growth Chambers for the Cultivation of Marine Sponge-Associated Bacteria. Mar
Biotechnol 16:594–603.

23. Nichols D, Cahoon N, Trakhtenberg EM, Pham L, Mehta A, Belanger A, Kanigan T, Lewis
K, Epstein SS. 2010. Use of Ichip for High-Throughput In Situ Cultivation of
"Uncultivable" Microbial Species. Appl Environ Microbiol 76:2445–2450.

24. Tandogan N, Abadian PN, Epstein S, Aoi Y, Goluch ED. 2014. Isolation of microorganisms
using sub-micrometer constrictions. PLoS One 9:e101429.

25. Hahn MW, Stadler P, Wu QL, Pöckl M. 2004. The filtration–acclimatization method for
isolation of an important fraction of the not readily cultivable bacteria. J Microbiol Methods
57:379–390.

26. Connon SA, Giovannoni SJ. 2002. High-throughput methods for culturing microorganisms
in very-low-nutrient media yield diverse new marine isolates. Appl Environ Microbiol
68:3878–3885.

27. Yang SJ, Kang I, Cho JC. 2016. Expansion of Cultured Bacterial Diversity by Large-Scale
Dilution-to-Extinction Culturing from a Single Seawater Sample. Microb Ecol 71:29–43.

28. Stingl U, Tripp HJ, Giovannoni SJ. 2007. Improvements of high-throughput culturing
yielded novel SAR11 strains and other abundant marine bacteria from the Oregon coast and
the Bermuda Atlantic Time Series study site. ISME J 1:361–371.

29. Henson MW, Lanclos VC, Faircloth BC, Thrash JC. 2018. Cultivation and genomics of the
first freshwater SAR11 (LD12) isolate. ISME J 12:1846–1860.

30. Hahnke RL, Bennke CM, Fuchs BM, Mann AJ, Rhiel E, Teeling H, Amann R, Harder J.
2015. Dilution cultivation of marine heterotrophic bacteria abundant after a spring
phytoplankton bloom in the North Sea. Environ Microbiol 17:3515–3526.

31. Sosa OA, Gifford SM, Repeta DJ, DeLong EF. 2015. High molecular weight dissolved
organic matter enrichment selects for methylotrophs in dilution to extinction cultures. ISME
J 9:2725–2739.

32. Schut F, de Vries EJ, Gottschal JC, Robertson BR, Harder W, Prins RA, Button DK. 1993.
Isolation of Typical Marine Bacteria by Dilution Culture: Growth, Maintenance, and
Characteristics of Isolates under Laboratory Conditions. Appl Environ Microbiol 59:2150–
2160.

33. Button DK, Schut F, Quang P, Martin R, Robertson BR. 1993. Viability and isolation of
marine bacteria by dilution culture: theory, procedures, and initial results. Appl Environ
Microbiol 59:881–891.

34. Song J, Oh HM, Cho JC. 2009. Improved culturability of SAR11 strains in dilution-to-
extinction culturing from the East Sea, West Pacific Ocean. FEMS Microbiol Lett 295:141–
147.

35. Henson MW, Pitre DM, Weckhorst JL, Lanclos VC, Webber AT, Thrash JC. 2016.
Artificial Seawater Media Facilitate Cultivating Members of the Microbial Majority from
the Gulf of Mexico. mSphere 1:e00028–16.

36. Rappé MS, Connon SA, Vergin KL, Giovannoni SJ. 2002. Cultivation of the ubiquitous
SAR11 marine bacterioplankton clade. Nature 418:630–633.

37. Marshall KT, Morris RM. 2013. Isolation of an aerobic sulfur oxidizer from the
SUP05/Arctic96BD-19 clade. ISME J 7:452–455.

38. Shah V, Chang BX, Morris RM. 2017. Cultivation of a chemoautotroph from the SUP05
clade of marine bacteria that produces nitrite and consumes ammonium. ISME J 11:263–
271.

39. Spietz RL, Lundeen RA, Zhao X, Nicastro D, Ingalls AE, Morris RM. 2019. Heterotrophic
carbon metabolism and energy acquisition in Candidatus Thioglobus singularis strain PS1, a
member of the SUP05 clade of marine Gammaproteobacteria. Environ Microbiol 21:2391–
2401.

40. Huggett MJ, Hayakawa DH, Rappé MS. 2012. Genome sequence of strain HIMB624, a

cultured representative from the OM43 clade of marine Betaproteobacteria. Stand Genomic
Sci 6:11–20.

41. Giovannoni SJ, Hayakawa DH, Tripp HJ, Stingl U, Givan SA, Cho J-C, Oh H-M, Kitner
JB, Vergin KL, Rappé MS. 2008. The small genome of an abundant coastal ocean
methylotroph. Environ Microbiol 10:1771–1782.

42. Durham BP, Grote J, Whittaker KA, Bender SJ, Luo H, Grim SL, Brown JM, Casey JR,
Dron A, Florez-Leiva L, Others. 2014. Draft genome sequence of marine
alphaproteobacterial strain HIMB11, the first cultivated representative of a unique lineage
within the Roseobacter clade possessing an unusually small genome. Stand Genomic Sci
9:632–645.

43. Cho JC, Giovannoni SJ. 2004. Cultivation and Growth Characteristics of a Diverse Group
of Oligotrophic Marine Gammaproteobacteria. Appl Environ Microbiol 70:432–440.

44. Kim S, Kang I, Seo J-H, Cho J-C. 2019. Culturing the ubiquitous freshwater actinobacterial
acI lineage by supplying a biochemical "helper"catalase. ISME J 13:2252–2263.

45. Tripp HJ. 2013. The unique metabolism of SAR11 aquatic bacteria. J Microbiol 51:147–
153.

46. Nichols D, Lewis K, Orjala J, Mo S, Ortenberg R, O'Connor P, Zhao C, Vouros P,
Kaeberlein T, Epstein SS. 2008. Short Peptide Induces an "Uncultivable" Microorganism
To Grow In Vitro. Appl Environ Microbiol 74:4889–4897.

47. Stewart EJ. 2012. Growing unculturable bacteria. J Bacteriol 194:4151–4160.

48. Kell DB, Young M. 2000. Bacterial dormancy and culturability: the role of autocrine
growth factors. Curr Opin Microbiol 3:238–243.

49. Carini P, Steindler L, Beszteri S, Giovannoni SJ. 2013. Nutrient requirements for growth of
the extreme oligotroph "Candidatus Pelagibacter ubique" HTCC1062 on a defined medium.
ISME J 7:592–602.

50. Epstein SS. 2013. The phenomenon of microbial uncultivability. Curr Opin Microbiol
16:636–642.

51. Shah D, Zhang Z, Khodursky A, Kaldalu N, Kurg K, Lewis K. 2006. Persisters: a distinct
physiological state of E. coli. BMC Microbiol 6:53.

52. Kell D, Potgieter M, Pretorius E. 2015. Individuality, phenotypic differentiation, dormancy
and "persistence" in culturable bacterial systems: commonalities shared by environmental,
laboratory, and clinical microbiology. F1000Res 4:179.

53. Lennon JT, Jones SE. 2011. Microbial seed banks: the ecological and evolutionary
implications of dormancy. Nat Rev Microbiol 9:119–130.

54. Grassi L, Di Luca M, Maisetta G, Rinaldi AC, Esin S, Trampuz A, Batoni G. 2017.
Generation of persister cells of Pseudomonas aeruginosa and Staphylococcus aureus by
chemical treatment and evaluation of their susceptibility to membrane-targeting agents.
Front Microbiol 8:1917.

55. Bergkessel M, Basta DW, Newman DK. 2016. The physiology of growth arrest: uniting
molecular and environmental microbiology. Nat Rev Microbiol 14:549–562.

56. Kussell E, Leibler S. 2005. Phenotypic diversity, population growth, and information in
fluctuating environments. Science 309:2075–2078.

57. Kurm V, van der Putten WH, Gera Hol WH. 2019. Cultivation-success of rare soil bacteria
is not influenced by incubation time and growth medium. PLoS One 14:e0210073.

58. D'Onofrio A, Crawford JM, Stewart EJ, Witt K, Gavrish E, Epstein S, Clardy J, Lewis K.
2010. Siderophores from neighboring organisms promote the growth of uncultured bacteria.
Chem Biol 17:254–264.

59. Thrash JC. 2019. Culturing the Uncultured: Risk versus Reward. mSystems 4:e00130–19.

60. Thrash JC, Weckhorst JL, Pitre DM. 2017. Cultivating Fastidious Microbes, p. 57–78. *In*
McGenity, TJ, Timmis, KN, Nogales, B (eds.), Hydrocarbon and Lipid Microbiology
Protocols. Springer Berlin Heidelberg, Berlin, Heidelberg.

61. Vila-Costa M, Simó R, Harada H, Gasol JM, Slezak D, Kiene RP. 2006.
Dimethylsulfoniopropionate uptake by marine phytoplankton. Science 314:652–654.

62. Mou X, Hodson RE, Moran MA. 2007. Bacterioplankton assemblages transforming
dissolved organic compounds in coastal seawater. Environ Microbiol 9:2025–2037.

63. Stein LY. 2015. Microbiology: Cyanate fuels the nitrogen cycle. Nature 524:43–44.

64. Repeta DJ, Ferrón S, Sosa OA, Johnson CG, Repeta LD, Acker M, Delong EF, Karl DM.
2016. Marine methane paradox explained by bacterial degradation of dissolved organic
matter. Nat Geosci 9:884–887.

65. Curson ARJ, Liu J, Bermejo Martínez A, Green RT, Chan Y, Carrión O, Williams BT,
Zhang S-H, Yang G-P, Bulman Page PC, Zhang X-H, Todd JD. 2017.
Dimethylsulfoniopropionate biosynthesis in marine bacteria and identification of the key
gene in this process. Nat Microbiol 2:17009.

66. Widner B, Mulholland MR. 2017. Cyanate distribution and uptake in North Atlantic coastal
waters. Limnol Oceanogr 62:2538–2549.

67. Widner B, Mulholland MR, Mopper K. 2016. Distribution, Sources, and Sinks of Cyanate
in the Coastal North Atlantic Ocean. Environ Sci Technol Lett 3:297–302.

68. Henson MW, Hanssen J, Spooner G, Fleming P, Pukonen M, Stahr F, Thrash JC. 2018.

Nutrient dynamics and stream order influence microbial community patterns along a 2914
kilometer transect of the Mississippi River. Limnol Oceanogr 63:1837–1855.

69. Apprill A, McNally S, Parsons R, Weber L. 2015. Minor revision to V4 region SSU rRNA
806R gene primer greatly increases detection of SAR11 bacterioplankton. Aquat Microb
Ecol 75:129–137.

70. Walters W, Hyde ER, Berg-Lyons D, Ackermann G, Humphrey G, Parada A, Gilbert JA,
Jansson JK, Caporaso JG, Fuhrman JA, Apprill A, Knight R. 2016. Improved Bacterial 16S
rRNA Gene (V4 and V4-5) and Fungal Internal Transcribed Spacer Marker Gene Primers
for Microbial Community Surveys. mSystems 1:e00009–15.

71. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski R a.,
Oakley BB, Parks DH, Robinson CJ, Sahl JW, Stres B, Thallinger GG, Van Horn DJ,
Weber CF. 2009. Introducing mothur: Open-source, platform-independent, community-
supported software for describing and comparing microbial communities. Appl Environ
Microbiol 75:7537–7541.

72. Pruesse E, Quast C, Knittel K, Fuchs BM, Ludwig W, Peplies J, Glockner FO, Glöckner
FO. 2007. SILVA: a comprehensive online resource for quality checked and aligned
ribosomal RNA sequence data compatible with ARB. Nucleic Acids Res 35:7188–7196.

73. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO.
2013. The SILVA ribosomal RNA gene database project: Improved data processing and
web-based tools. Nucleic Acids Res 41:590–596.

74. Eren AM, Borisy GG, Huse SM, Mark Welch JL. 2014. Oligotyping analysis of the human
oral microbiome. Proc Natl Acad Sci U S A 111:E2875–84.

75. Fujimoto M, Cavaletto J, Liebig JR, McCarthy A, Vanderploeg HA, Denef VJ. 2016.
Spatiotemporal distribution of bacterioplankton functional groups along a freshwater
estuary to pelagic gradient in Lake Michigan. J Great Lakes Res 42:1036–1048.

76. Cole JR, Wang Q, Fish JA, Chai B, McGarrell DM, Sun Y, Brown CT, Porras-Alfaro A,
Kuske CR, Tiedje JM. 2014. Ribosomal Database Project: data and tools for high
throughput rRNA analysis. Nucleic Acids Res 42:D633–42.

77. Team RC. 2017. R Core Team (2017). R: A language and environment for statistical
computing. R Found Stat Comput Vienna, Austria URL http://www R-project org/ , page R
Foundation for Statistical Computing.

78. McMurdie PJ, Holmes S. 2013. phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. PLoS One 8:e61217.

79. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009.
BLAST+: architecture and applications. BMC Bioinformatics 10:421.

80. Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Res 32:1792–1797.

81. Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. Bioinformatics 25:1972–1973.

82. Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol
32:268–274.

83. Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: Improving
the Ultrafast Bootstrap Approximation. Mol Biol Evol 35:518–522.

84. Junier T, Zdobnov EM. 2010. The Newick utilities: high-throughput phylogenetic tree
processing in the UNIX shell. Bioinformatics 26:1669–1670.

85. Han MV, Zmasek CM. 2009. phyloXML: XML for evolutionary biology and comparative
genomics. BMC Bioinformatics 10:356.

86. Stingl U, Cho JC, Foo W, Vergin KL, Lanoil B, Giovannoni SJ. 2008. Dilution-to-
extinction culturing of psychrotolerant planktonic bacteria from permanently ice-covered
lakes in the McMurdo Dry Valleys, Antarctica. Microb Ecol 55:395–405.

87. Page KA, Connon SA, Giovannoni SJ. 2004. Representative freshwater bacterioplankton
isolated from Crater Lake, Oregon. Appl Environ Microbiol 70:6542–6550.

88. Noell SE, Giovannoni SJ. 2019. SAR11 bacteria have a high affinity and multifunctional
glycine betaine transporter. Environ Microbiol 21:2559–2575.

89. Thume K, Gebser B, Chen L, Meyer N, Kieber DJ, Pohnert G. 2018. The metabolite
dimethylsulfoxonium propionate extends the marine organosulfur cycle. Nature 563:412–
415.

90. Asher EC, Dacey JWH, Stukel M, Long MC, Tortell PD. 2017. Processes driving seasonal
variability in DMS, DMSP, and DMSO concentrations and turnover in coastal Antarctic
waters. Limnol Oceanogr 62:104–124.

91. Mizuno CM, Rodriguez-Valera F, Ghai R. 2015. Genomes of planktonic Acidimicrobiales:
widening horizons for marine Actinobacteria by metagenomics. MBio 6:e02083–14.

92. Lee J, Kwon KK, Lim S-I, Song J, Choi AR, Yang S-H, Jung K-H, Lee J-H, Kang SG, Oh
H-M, Cho JC. 2019. Isolation, cultivation, and genome analysis of proteorhodopsin-
containing SAR116-clade strain Candidatus Puniceispirillum marinum IMCC1322. J
Microbiol 57:676–687.

93. Kitzinger K, Padilla CC, Marchant HK, Hach PF, Herbold CW, Kidane AT, Könneke M,
Littmann S, Mooshammer M, Niggemann J, Petrov S, Richter A, Stewart FJ, Wagner M,
Kuypers MMM, Bristow LA. 2019. Cyanate and urea are substrates for nitrification by
Thaumarchaeota in the marine environment. Nat Microbiol 4:234–243.

94. Kamennaya NA, Post AF. 2011. Characterization of cyanate metabolism in marine
Synechococcus and Prochlorococcus spp. Appl Environ Microbiol 77:291–301.

95. Eren AM, Maignien L, Sul WJ, Murphy LG, Grim SL, Morrison HG, Sogin ML. 2013.
Oligotyping: Differentiating between closely related microbial taxa using 16S rRNA gene
data. Methods Ecol Evol 4:1111–1119.

96. Needham DM, Fuhrman JA. 2016. Pronounced daily succession of phytoplankton, archaea
and bacteria following a spring bloom. Nat Microbiol 1:16005.

97. Campbell BJ, Yu L, Heidelberg JF, Kirchman DL. 2011. Activity of abundant and rare
bacteria in a coastal ocean. Proc Natl Acad Sci U S A 108:12776–12781.

98. Oh HM, Kwon KK, Kang I, Kang SG, Lee JH, Kim SJ, Cho JC. 2010. Complete genome
sequence of "Candidatus puniceispirillum marinum" IMCC1322, a representative of the
SAR116 clade in the Alphaproteobacteria. J Bacteriol 192:3240–3241.

99. Fegatella F, Lim J, Kjelleberg S, Cavicchioli R. 1998. Implications of rRNA operon copy
number and ribosome content in the marine oligotrophic ultramicrobacterium
Sphingomonas sp. strain RB2256. Appl Environ Microbiol 64:4433–4438.

100. Acinas SG, Marcelino LA, Klepac-Ceraj V, Polz MF. 2004. Divergence and redundancy
of 16S rRNA sequences in genomes with multiple rrn operons. J Bacteriol 186:2629–2635.

101. Pedrós-Alió C. 2012. The rare bacterial biosphere. Ann Rev Mar Sci 4:449–466.

102. Martiny AC. 2019. High proportions of bacteria are culturable across major biomes.
ISME J 13:2125–2128.

103. Parada AE, Needham DM, Fuhrman JA. 2015. Every base matters: assessing small
subunit rRNA primers for marine microbiomes with mock communities, time series and
global field samples. Environ Microbiol 18:1403–1414.

104. Del Giorgio PA, Gasol JM. 2008. Physiological structure and single-cell activity in
marine bacterioplankton. Microbial Ecology of the Oceans 2:243–298.

105. Malmstrom RR, Kiene RP, Cottrell MT, Kirchman DL. 2004. Contribution of SAR11
bacteria to dissolved dimethylsulfoniopropionate and amino acid uptake in the North
Atlantic Ocean. Appl Environ Microbiol 70:4129–4135.

106. Malmstrom RR, Cottrell MT, Elifantz H, Kirchman DL. 2005. Biomass production and
assimilation of dissolved organic matter by SAR11 bacteria in the Northwest Atlantic
Ocean. Appl Environ Microbiol 71:2979–2986.

107. Piwosz K, Salcher MM, Zeder M, Ameryk A, Pernthaler J. 2013. Seasonal dynamics and
activity of typical freshwater bacteria in brackish waters of the Gulf of Gdańsk. Limnol
Oceanogr 58:817–826.

108. Laghdass M, Catala P, Caparros J, Oriol L, Lebaron P, Obernosterer I. 2012. High
contribution of SAR11 to microbial activity in the north west Mediterranean Sea. Microb
Ecol 63:324–333.

109. Salter I, Galand PE, Fagervold SK, Lebaron P, Obernosterer I, Oliver MJ, Suzuki MT,
Tricoire C. 2015. Seasonal dynamics of active SAR11 ecotypes in the oligotrophic
Northwest Mediterranean Sea. ISME J 9:347–360.

110. Leizeaga A, Estrany M, Forn I, Sebastián M. 2017. Using Click-Chemistry for
Visualizing in Situ Changes of Translational Activity in Planktonic Marine Bacteria. Front
Microbiol 8:2360.

111. Kirchman DL. 2016. Growth Rates of Microbes in the Oceans. Ann Rev Mar Sci 8:285–
309.

112. Samo TJ, Smriga S, Malfatti F, Sherwood BP, Azam F. 2014. Broad distribution and high
proportion of protein synthesis active marine bacteria revealed by click chemistry at the
single cell level. Frontiers in Marine Science 1:48.

113. Smriga S, Samo TJ, Malfatti F, Villareal J, Azam F. 2014. Individual cell DNA synthesis
within natural marine bacterial assemblages as detected by "click"chemistry. Aquat Microb
Ecol 72:269–280.

114. Bradley JA, Amend JP, LaRowe DE. 2019. Survival of the fewest: Microbial dormancy
and maintenance in marine sediments through deep time. Geobiology 17:43–59.

115. Zhao Y, Temperton B, Thrash JC, Schwalbach MS, Vergin KL, Landry ZC, Ellisman M,
Deerinck T, Sullivan MB, Giovannoni SJ. 2013. Abundant SAR11 viruses in the ocean.
Nature 494:357–360.

116. Martinez-Hernandez F, Fornas Ò, Lluesma Gomez M, Garcia-Heredia I, Maestre-
Carballa L, López-Pérez M, Haro-Moreno JM, Rodriguez-Valera F, Martinez-Garcia M.
2018. Single-cell genomics uncover Pelagibacter as the putative host of the extremely
abundant uncultured 37-F6 viral population in the ocean. ISME J 13:232–236.

117. Chen L-X, Zhao Y, McMahon KD, Mori JF, Jessen GL, Nelson TC, Warren LA,
Banfield JF. 2019. Wide Distribution of Phage That Infect Freshwater SAR11 Bacteria.
mSystems 4:e00410–19.

118. Thingstad TF, Lignell R. 1997. Theoretical models for the control of bacterial growth
rate, abundance, diversity and carbon demand. Aquat Microb Ecol 13:19–27.

119. Våge S, Storesund JE, Thingstad TF. 2013. SAR11 viruses and defensive host strains.
Nature 499:E3–4.

120. Thingstad TF, Vage S, Storesund JE, Sandaa RA, Giske J. 2014. A theoretical analysis of
how strain-specific viruses can control microbial species diversity. Proc Natl Acad Sci U S
A 111:7813–7818.

121.   Giovannoni S, Temperton B, Zhao Y. 2013. Giovannoni et al. reply. Nature 499:E4–E5.

122.   Alonso-Sáez L, Morán XAG, Clokie MR. 2018. Low activity of lytic pelagiphages in
1492   coastal marine waters. ISME J 12:2100–2102.

1493   123.   Chapman-McQuiston E, Wu XL. 2008. Stochastic receptor expression allows sensitive
1494   bacteria to evade phage attack. Part I: experiments. Biophys J 94:4525–4536.

1495   124.   Bertozzi Silva J, Storms Z, Sauvageau D. 2016. Host receptors for bacteriophage
1496   adsorption. FEMS Microbiol Lett 363:fnw002.

1497   125.   Oliver JD. 2005. The viable but nonculturable state in bacteria. J Microbiol 43 Spec
1498   No:93–100.

1499   126.   Dubnau D, Losick R. 2006. Bistability in bacteria. Mol Microbiol 61:564–572.

1500   127.   Shkoporov AN, Khokhlova EV, Fitzgerald CB, Stockdale SR, Draper LA, Ross RP, Hill
1501   C. 2018. ΦCrAss001 represents the most abundant bacteriophage family in the human gut
1502   and infects Bacteroides intestinalis. Nat Commun 9:4781.

1503   128.   Teira E, Martínez-García S, Lønborg C, Alvarez-Salgado XA. 2009. Growth rates of
1504   different phylogenetic bacterioplankton groups in a coastal upwelling system. Environ
1505   Microbiol Rep 1:545–554.

1506   129.   Lankiewicz TS, Cottrell MT, Kirchman DL. 2016. Growth rates and rRNA content of
1507   four marine bacteria in pure cultures and in the Delaware estuary. ISME J 10:823–832.

1508   130.   Ferrera I, Gasol JM, Sebastián M, Hojerová E, Koblížek M. 2011. Comparison of growth
1509   rates of aerobic anoxygenic phototrophic bacteria and other bacterioplankton groups in
1510   coastal Mediterranean waters. Appl Environ Microbiol 77:7451–7458.

1511   131.   Carini P, Campbell EO, Morré J, Sañudo-Wilhelmy SA, Thrash JC, Bennett SE,
1512   Temperton B, Begley T, Giovannoni SJ. 2014. Discovery of a SAR11 growth requirement
1513   for thiamin's pyrimidine precursor and its distribution in the Sargasso Sea. ISME J 8:1727–
1514   1738.

1515   132.   Grant SR, Church MJ, Ferrón S, Laws EA, Rappé MS. 2019. Elemental Composition,
1516   Phosphorous Uptake, and Characteristics of Growth of a SAR11 Strain in Batch and
1517   Continuous Culture. mSystems 4:e00218–18.

1518   133.   White AE, Giovannoni SJ, Zhao Y, Vergin K, Carlson CA. 2019. Elemental content and
1519   stoichiometry of SAR11 chemoheterotrophic marine bacteria. Limnology and
1520   Oceanography Letters 4:44–51.

1521   134.   Smith DP, Cameron Thrash J, Nicora CD, Lipton MS, Burnum-Johnson KE, Carini P,
1522   Smith RD, Giovannoni SJ. 2013. Proteomic and Transcriptomic Analyses of "Candidatus
1523   Pelagibacter ubique" Describe the First PII-Independent Response to Nitrogen Limitation in
1524   a Free-Living Alphaproteobacterium. MBio 4:e00133–12.

1525 135. van Vliet S, Dal Co A, Winkler AR, Spriewald S, Stecher B, Ackermann M. 2018.
1526   Spatially Correlated Gene Expression in Bacterial Groups: The Role of Lineage History,
1527   Spatial Gradients, and Cell-Cell Interactions. Cell Syst 6:496–507.e6.

1528

Figure S1

Figure S2

Figure S4

Figure S5