Sexually dimorphic traits and male-specific differentiation are actively regulated by Doublesex during specific developmental windows in *Nasonia vitripennis*

Yidong Wang [a], Anna Rensink [b], Ute Fricke [a,*], Megan C. Riddle [c,$], Carol Trent [c], Louis van de Zande [b], Eveline C. Verhulst [a,d]

a. Wageningen University, Laboratory of Entomology, Wageningen, The Netherlands
b. Evolutionary Genetics, Development and Behaviour, Groningen Institute for Evolutionary Life Sciences, University of Groningen, Groningen, The Netherlands
c. Biology Department, Western Washington University, Washington, United States of America
d. Wageningen University, Laboratory of Genetics, Wageningen, The Netherlands

* Current address: Departement of Animal Ecology and Tropical Biology, Biocenter, University of Würzburg, Würzburg, Germany

$ Current addresses: Department of Psychiatry and Behavioral Sciences, University of Washington, Seattle, WA, United States of America; Eating Recovery Center
1231 116th Ave NE Suite 800, Bellevue, WA 98004

Corresponding author: eveline.verhulst@wur.nl

Keywords: *Nasonia,* Doublesex, sexually dimorphic traits, sexual differentiation, parasitoid

wasps, sex determination

Abstract

Sexually dimorphic traits in insects are rapidly evolving due to sexual selection which can

ultimately lead to speciation. However, our knowledge of the underlying sex-specific

molecular mechanisms is still scarce. Here we show that the highly conserved gene,

Doublesex (Dsx), regulates rapidly diverging sexually dimorphic traits and differentiation in the

model parasitoid wasp *Nasonia vitripennis* (Hymenoptera: Pteromalidae). We present here

the revised full *Dsx* gene structure with an alternative exon one and two additional male

NvDsx isoforms, which gives important insights into the evolution of the sex-specific

oligomerization domains and C-termini. We show the sex-specific *NvDsx* expression

throughout development, and demonstrate that transient *NvDsx* silencing in different male

developmental stages dramatically shifts the morphology of two sexually dimorphic traits from

male to female, with the effect being dependent on the timing of silencing. In addition,

transient silencing of *NvDsx* in early male larvae affects male genitalia tissue growth but not

morphology. This shows that male *NvDsx* is actively required to suppress female-specific

traits and promote male-specific traits during specific developmental windows. These results

also strongly suggests that in *N. vitripennis* most sex-specific tissues fully differentiate in the

embryonic stage and only need *NvDsx* input for growth afterwards. This provides a first

insight into the regulatory activity of *Dsx* in Hymenoptera and will help us to better understand

the evolutionary and molecular mechanisms involved in sex-specific development, which can

eventually lead to the development of new synthetic genetics-based tools for biological pest

control by parasitoid wasps.

Significance Statement

In insects, male and female differentiation is regulated by the highly conserved transcription

factor *Doublesex* (*Dsx*)*.* The role of *Dsx* in regulating rapidly evolving sexually dimorphic traits

has received less attention, especially in wasps and bees. Here, we mainly focused on *Dsx*

regulation of two sexually dimorphic traits and male genitalia morphology in the parasitoid

wasp, *Nasonia vitripennis.* We demonstrate that *Dsx* actively regulates male-specific tissue

growth and morphology during specific developmental windows. These findings will help us to

better understand the molecular mechanisms underlying the rapid evolution of sexual

differentiation and sexually dimorphic traits in insects, but may also be the starting point for

the development of new tools for biological control of pest insects by parasitoid wasps.

Introduction

Insects are well known for their distinct differences between male and female morphology.

The reproductive organs and sexually dimorphic traits such as wing shapes and colour patterns are one of the fastest evolving traits in insects and are therefore often used to distinguish species, as these traits can differ dramatically between species. Development of these fast changing sexually dimorphic traits usually depends on specific genes that are expressed tissue and sex specifically (1). Many of these genes are under control of the transcription factor, Doublesex (Dsx), that initiates sexual differentiation in insects (2–4). *Dsx* pre-mRNA is alternatively spliced to yield male-specific *Dsx* (*DsxM*) or female-specific *Dsx* (*DsxF*) transcripts, resulting in distinct sex-specific protein isoforms (5). Two functional domains are present in all insect Dsx homologs: 1) the DM domain which consists of a DNA binding domain (DBD) and an oligomerization domain (OD1) with a zinc module and binding sites to facilitate binding downstream targets; 2) a dimerization domain (OD2) with a common N-terminal part and a sex specific C-terminal part. These C-terminal sex-specific differences are assumed to be involved in the regulation of the development of sex-specific traits (6,7). For instance, in *Drosophila*, DsxM represses and DsxF activates the expression of *bric-à-brac (bab)* which results in male-specific abdominal pigmentation (8). A similar action is observed in the formation of sex combs, an important feature for performing mating rituals in males of some *Drosophila* species (9). In *Bactrocera dorsalis,* DsxF knockdown in females triggers the interruption of yolk protein gene expression, leading to ovipositor deformation and a delay in ovary development which confirms the essential role of DsxF to initiate and maintain female development in Diptera (10). In Lepidoptera (butterflies and moths) and Coleoptera (beetles), sex-specific *Dsx* expression in both sexes also affects many sex specific traits. For example, transient expression of *DsxM* in female silkmoths *Bombyx mori* distorts the development of female-specific genital organs by partially introducing male characters in female genitalia (11), and introducing mutations in the male-specific *Dsx* isoform of *B. mori* yields either malfunctioning males or intersexes (12). In *Papilio polytes* butterflies differential expression of *Dsx* isoforms results in sex-specific wing patterns (13). *Dsx* also either promotes or constrains the mandible growth in males and females of *Cyclommatus metallifer* beetles (14), and the formation of head horns in *Trypoxylus dichotomus* beetles is antagonistically regulated by *DsxM* and *DsxF*, while the thoracic horn formation is *DsxM* independent (15,16). Noteworthy, all these experiments were conducted only in a specific developmental stage and only few focused on sexually dimorphic traits. However, the expression of *Dsx* starts in early embryogenesis and continues through maturity which implies its ongoing action (10,17,18).

Thus far, the specific role of *Dsx* in regulating sexually dimorphic traits has not been studied in the Hymenoptera. *Dsx* sex-specific splice variants and expression differences have been identified in honeybees, (*Apis mellifera*), bumble bees (*Bombus ignites),* Japanese ants

(*Vollenhovia emeryi*) and fire ants (*Solenopsis invicta*) (19–23), but the role of *Dsx* in

regulating reproductive organ development in males has only been shown in sawflies, *Athalia*

rosae (24). However, to understand the rapid evolution of sexual dimorphism in other insect

orders and to compare the molecular mechanisms of *Dsx*-mediated regulation, it is

paramount to gain more knowledge on how *Dsx* directs sex-specific and sexually dimorphic

trait development in Hymenoptera. Therefore, we set out to 1) identify the sex-specific traits

that are regulated by *Dsx* and 2) study the timing by which *Dsx* regulates these traits during

development in the parasitoid wasp *Nasonia vitripennis*.

N. vitripennis belongs to the genus *Nasonia* (*Hymenoptera: Pteromalidae*), and has become

the model system for parasitoids as it is highly suitable for evolutionary and developmental

genetic studies. Multiple sexually dimorphic traits within the genus *Nasonia* makes it a perfect

system to study sex differentiation (25,26). For instance, the femur of hind legs and antennae

of female *Nasonia* are pigmented and have a dark-brown color, whereas in males these are

pale yellow (26). Only in *N. vitripennis,* males have dramatically shorter forewings and are

flightless, (26), while the females can fly and have large forewings in all species. Although

Dsx has been identified in *N. vitripennis* (27), its role in regulating these sexually dimorphic

traits has not been shown before. Previous research identified one male *Dsx* (*NvDsxM*) and

one female *Dsx* (*NvDsxF*) splice variant (27) and showed that the conserved splicing factor

transformer (Nvtra) is responsible for female-specific splicing of *Dsx* transcripts, while the

male-specific *Dsx* transcript is spliced by default into the male isoform (28).

To assess the role of *Dsx* in regulating sexual dimorphy in *Nasonia,* we first revisited the

NvDsx genetic structure. We confirmed existing splice-variants and identified an alternative

starting exon and two additional male *NvDsx* splice-variants. We then used RNA interference

to temporally silence *Dsx* in different male developmental stages to study its function in

sexual differentiation and dimorphism. We found that transient silencing of *NvDsx* in males

leads to the emergence of different female-specific morphological traits with the effect being

dependent on the timing of silencing during development. In addition, transient silencing of

NvDsx in early male larvae affects male genitalia tissue growth but not morphology. This

indicates that the appearance of these sexually dimorphic traits depends on the action of

NvDsx in a very specific time window during development. Sex-specific tissue differentiation

appears to occur only during embryonic development and constitutive *NvDsx* expression is

required to further promote the growth of the reproductive systems, which has not been

reported in other insect systems (24,29). Ultimately, our results provide a great start to study

the molecular basis of sexual differentiation and rapidly evolving sexual dimorphism in insects

in general and particularly in Hymenoptera.

Results and discussion

Identification and characterization of two additional male-specific *Dsx* splice variants

Previously, only one female (*NvDsxF*) and one male (*NvDsxM*) *Dsx* splice form were

identified in *Nasonia vitripennis* (27). The *NvDsxF* splice variant shared the same first 4 exons

with the *NvDsxM* splice variant, but 108 bases are spliced out from exon 5 which leads to

female-specific exons 5F and 6F. In *NvDsxM* exon 5M results from retaining this 108bp

intron which has not been observed in other insects (27) (Fig. 1, *NvDsxF* and *NvDsxM1*). This

splice isoform translates into a Dsx protein with a very short OD2 domain of only four amino

acids. This *NvDsx* gene structure does not resemble any previously published *Dsx* structures.

To investigate the possibility of additional splice variants of *Dsx* pre-mRNA in *N. vitripennis*

we performed extensive 5' and 3' RACEs. The results revealed a novel alternative starting

exon that we designated exon 1' (Fig. 1). Note that the start codon of protein translation is in

exon 2. Using RT-PCR to target exon 1' to exon 5 we verified its presence in transcripts of

both sexes but with a higher frequency in female transcripts (Fig. S1). Alternative splicing at

the 5' end has not been shown for *Dsx* before. We assume the alternative exon is used in all

splice variants (Fig. 1) but long-read RNA sequencing would be required to confirm this. From

here on we refer to the alternative variants collectively. The 3' RACEs identified two additional

longer male splice variants (Fig. 1). We designated the previously published male-specific

variant *NvDsxM1* (MT043363); the newly found longer male splice forms were designated

NvDsxM2 (M2: MT043359; M2': MT043360) and *NvDsxM3* (M3: MT043361; M3': MT043362).

All male isoforms share exons 1 (or 1') to 5 with the female isoform *NvDsxF* (MT043364). The

NvDsxM2 open reading frame ends with a stop codon in exon 8 and translates into a protein

of 328 amino acids (aa) which is 105 aa longer than the open reading frame of *NvDsxM1*.

Compared to *NvDsxM2* we find that exon 8 is skipped in *NvDsxM3* which puts the stop codon

in exon 9 yielding a protein of 330 amino acids. These additional male specific splice variants

skip the female-specific exon 6 completely which is a shared feature of male splice forms in

many different insect species including fruit flies (*D. melanogaster*) (30) (Fig. 1), wild

silkmoths (*Antheraea assama*) (31), honeybees (*A. mellifera*) (21) and ants (*S. invicta*) (20).

We did not find any non-sex specific splice variants in *N. vitripennis* as opposed to *A.*

mellifera (Fig. 1) (21). NvDsxM2 and NvDsxM3 protein isoforms are completely identical and

only differ at the C-terminus after the male-specific OD2 domain, while NvDsxM1 has a male-

specific OD2 domain of only 4 aa (Fig. S6A).

NvDsxM2 and NvDsxM3 are spliced by default

To determine whether the additionally identified splice variants are not exclusively expressed in females, we first observed the overall expression of different *Nvdsx* splice forms in both males and females. Across development, *NvDsxF* is only expressed in females while *NvDsxM1, NvDsxM2* and *NvDsxM3* are highly expressed in males (Fig. S2), and expressed at a low level in females. Interestingly, only in males a slight increase of *NvDsxM1, -2 and -3* expression from 36h toward the early pupal stage was observed. Afterwards, the expression remained relatively constant until adult stage (Fig. S2). Overall, *NvDsxM2* is more abundant than *NvDsxM3* (Fig. S2). Females express *NvDsxM1, NvDsxM2* and *NvDsxM3* at a low level probably due to the leaky default splicing of male-specific transcripts (Fig. 2B and Fig. S2). Additionally, we studied the effect of *Nvtra* silencing on *Nvdsx* splicing in females as male-specific splice variants are processed by default and do not depend on NvtraF regulation. This means that all three male splice variants should be higher expressed in females after *Nvtra* silencing as there is less NvtraF to regulate splicing of all *Nvdsx* transcripts into the female-specific form. We have shown previously that *Nvtra* pRNAi knockdown in the female pupal stage leads to a shift in splicing from *NvtraF* to *NvtraM* (28). This indeed induced a shift from *NvDsxF* to *NvDsxM* (now *NvDsxM1*) splicing in both the treated adult females and the offspring, without consequences for adult female fertility (28). To verify the default and NvTra independent splicing of *NvDsxM2* and *NvDsxM3*, we used *Nvtra* RNA interference in the last larval instar (4th instar) of females followed by RT-PCR amplification of the different *NvDsx* and *Nvtra* splice variants from 3 hours post injection (hpi) to 5 days post injection (dpi). After confirming that *Nvtra* RNAi leads to *NvtraM* splicing in females (Fig. 2), we did not observe a clear reduction of female *NvDsxF* expression. Instead, a minor increase of *NvDsxM1* expression from 4dpi onwards was observed. *NvDsxM2* and *NvDsxM3* expression levels did increase substantially from 1dpi onward compared to *gfp* knockdown control.

NvDsxM represses female-specific pigmentation

To assess the sex-specific function of Dsx, we studied the phenotypic effect of *NvDsx* silencing at different developmental time points. We were unable to silence the male-specific splice variants (*NvDsxM*) of *NvDsx* specifically (Fig. S3B), therefore we targeted the common region of *NvDsx* (*NvDsxU*). We observed that the developmental stage at which NvdsxU dsRNA was injected greatly influenced a measurable decrease of *NvDsx* using qRT-PCR. We found no significant decrease of *NvDsx* expression in 2nd instar *NvDsx* silenced males in comparison to non-treated and *gfp* control when sampling after white pupal stage (Fig. S3A). The strong increase of *NvDsx* expression from 2nd larval instar towards pupal stage (Fig. S3)

may be the root cause as the amount of siRNA present may be too limited to silence the sudden increase of transcript amounts. However, this allowed us to transiently suppress *NvDsx* during different developmental time windows. Injecting *NvDsxU* dsRNA in 4th instar males led to a significant reduction of *NvDsx* in newly emerged adults when compared with both non-treated (NT) (Tukey's HSD, P<.001) and *gfp* mock silenced males (Tukey's HSD, P<.001), confirming successful *NvDsx* knockdown starting in the 4th instar male larvae with an effect lasting into adulthood (Fig. S3*B*). *Gfp* mock silenced males also showed significant reduction of *NvDsx* expression compared to NT (Tukey's HSD, P=0.021) suggesting that in 4th instar larvae there might be a non-target effects of *gfp* dsRNA as was shown before in honey bee (32). Similar reduction in *NvDsx* silenced pupal stage males was observed when sampling at adult stage compared to NT (Tukey's HSD, P=0.003) and *gfp* control (Tukey's HSD, P=0.002) (Fig. S3*C*) which indicates that we also obtained a highly significant knockdown of *NvDsx* from pupa stage onwards.

Non-treated and *gfp* dsRNA injected males showed unpigmented antennae and legs (Fig. 3*A*, *B*, *C*), whereas females have pigmented antennae and legs (Fig. 3*D*). Depleting *NvDsx* from the 2nd larval instar onwards increases the amount of pigmentation slightly (Fig. 3*F*), but depleting *NvDsx* from male 4th instar larval stages onwards increases pigmentation in their adult stage to the point that it resembled female-like pigmentation (Fig. 3*E*). No pigmentation is observed in adult males when *NvDsx* is depleted from early white pupal stage onwards (Fig. S4*B*).

Given the fact that we do not find changes in pigmentation intensity by *NvDsx* RNAi experiments in 4th instar female larvae (Fig. S5*A* and *B*), and silencing *NvDsx* led to female-like pigmentation in males, we conclude that female pigmentation is the default developmental state and in males NvDsxM actively interferes with the melanin synthesis pathway. As we find no true intersex phenotype, we hypothesize that NvDsxF plays no role in enhancing the genes required for pigmentation in females. We observed a slight reduction in antenna pigmentation of 4th instar *NvDsx* knockdown male larvae which is likely due to 1) structural difference in the antennae with males having less plate organs on the subsegments than females causing a different color reflection (33), and/or 2) incomplete knockdown of *NvDsx.*

In *D. melanogaster,* expression of sex-specific abdominal pigmentation is regulated by interaction of the HOX protein Abdominal-B (Abd-B) and Dsx. Combined input of Abd-B and DsxF leads to the expression of *bric-a-brac* (*bab*) which suppresses the pigmentation in the posterior abdomen segments while DsxM represses *bab* to maintain the posterior abdomen

pigmentation (8). In *Drosophila*, the deposition of the pigments in the cuticle starts from late

pupal to early adult stages (34), therefore, initiating the RNAi response from 4th instar larva in

N. vitripennis onwards creates enough time to deplete *NvDsx* mRNA and NvDsx protein

production to inhibit the expression of pigmentation genes.

NvDsx silencing leads to increased forewing size of males

The second sex-specific character that we assessed was forewing size and shape in

N.vitripennis adult males. Silencing *NvDsx* in both 2nd and 4th instar resulted in increased

forewing length, and width when compared to *gfp* mock silencing (Tukey's HSD, P<0.001)

and NT males (Tukey's HSD, P<0.001) (Fig. 4*B* and *C*). Length/width (L/W) ratio of *NvDsx*

knockdown treatments also showed severe reduction compared to *gfp* mock (Tukey's HSD,

P<0.001) and NT males (Tukey's HSD, P<0.001) (Fig. 4*A*) indicating that the wing got wider

and more female-like. However, the forewings of the *NvDsx* knockdown (*NvDsx-i*), mock *gfp*

(*gfp-i*) knockdown and non-treated males were still smaller and narrower than those of the

non-treated females (Length: Tukey's HSD, P<0.001; Width: Tukey's HSD, P<0.001; L/W

ratio: Tukey's HSD, P<0.001) (Fig. 4*A*, *B* and *C*). Interestingly, also here we found an effect of

timing as the forewing length and width increased significantly between 2nd and 4th instar

NvDsx treatments of which earlier stage *NvDsx* knockdown led to more severe changes

(Tukey's HSD, P<0.001), steering the L/W ratio towards NT female's dramatically (Tukey's

HSD, P<0.001). Between 2nd instar *gfp* and 4th instar *gfp* knockdown (Tukey's HSD,

P=0.062), 2nd instar *gfp* knockdown and NT (Tukey's HSD, P=0.070) and 4th instar *gfp*

knockdown and NT (Tukey's HSD, P=0.999) no significant difference were observed (Fig. 4*A*,

B and *C*). Depleting *NvDsx* from the white pupal stage onwards had no effect on forewing

size or shape (Fig. S4*F*).

Previous research showed that the non-coding region of *Dsx* in *N. giraulti*, the wing-size1

locus (*ws1*), is correlated with the development of long wings in males, and the *ws1* variant in

N. vitripennis is correlated with the severe reduced forewing size in males (35). Although

higher *NvDsxM1* expression levels was detected in individuals with the *N. vitripennis* ws1

variant (35), the mechanism of *NvDsx* regulation is unclear and the *NvDsxM2* and -*3* splice

variants were unknown at the time. Our research confirms that the regulation of the male

forewing size depends on high *NvDsxM* levels and illustrates the role of *NvDsxM* in

repressing wing growth. Changes in spatial and temporal expression of *unpaired-like* has

been shown to result in changes in wing width in *Nasonia* males caused by multiple

regulatory changes (36). NvDsxM regulation is likely responsible for differentially expressed

upd-like. In contrast to the observed complete female-specific pigmentation after *NvDsxM*

depletion in males, the depletion of *NvDsxM* from 2nd instar onwards is inadequate to restore female forewing size. Wing size and shape development already takes place during embryonal development and our chosen time window for RNAi is too late to induce a full switch. We cannot rule out antagonistic effects of NvDsxM and NvDSXF on wing size development in *Nasonia*, since in females *Aedes aegypti, Dsx* knockdown leads to smaller wing development (37). However, we observed no wing size change after *NvDsx* knockdown in females (Fig. S4*F*).

Growth of male genitalia is controlled by *NvDsx*

Next, we depleted *NvDsx* at either 2nd or 4th instar larva stage to observe the role of *NvDsx* in male genitalia development and growth after embryogenesis is completed. This is in contrast to many studies being done using genome editing in which *Dsx* is knocked out from embryogenesis onwards (10,12). *NvDsx* RNAi in both developmental stages reduced the aedeagus length significantly when compared to *gfp* mock RNAi in both developmental stages and NT males (Tukey's HSD, P<0.001) (Fig. 5A). Both developmental stages had the same extent of reduction (Tukey's HSD, P=0.998). Aside from the shortened aedeagus, the phallomere structure (genital phallic lobes) is also severely reduced in size in both 2nd and 4th instar *NvDsx* depleted males (Fig. 5B-c and e) compared to *gfp* RNAi and NT males (Fig. 5B-a, b and d) but the structure of the paramere including the outer parameres, cuspis and digitus, is still intact (Fig. 5B). Compared to *gfp* RNAi males (Movie S1), the severe length reduction of the aedeagus renders *NvDsx* RNAi males unable to copulate with a female (Movie S2), showing that in this case size matters.

Dsx is shown to be involved in the differentiation of genitalia structures in male and female insects in many cases by interactions with HOX genes during early embryogenesis (39). However, whether Dsx is continuously required for genitalia development has not received much attention. Previous studies using *Dsx* RNAi experiments in other species conducted at late larval stage resulted in either deformed external genitalia or chimeras that contain the structures of both sexes (24,29). Here, even though *NvDsx* RNAi was done at a distinctly earlier stage (2nd instar larva) to other studied insect species, we could not observe any abnormalities in genital organs other than size. This shows that contrary to other insects, in *N. vitripennis* the differentiation of genital organs is completed before 2nd instar larval stage and only growth is required during later development. It suggests that besides the sex-specific tissue differentiation, *NvDsx* is also required for the direct sex-specific trait growth as it has been shown in stag beetles *Cyclommatus metallifer* where it was demonstrated that *Dsx* can manipulate the sensitivity of JH in a sex-specific way to regulate mandible growth in males

275 and females (14).

Evolution of Dsx sex-specific OD2 domains and C-termini

Aligning the OD2 domains of NvDsxM1, NvDsxM2 and NvDsxM3 (Fig. S6A) or NvDsxF (Fig.
S6B) with other DsxM or DsxF OD2 domains, respectively, shows that across insect orders
the male-specific OD2 domain is far less conserved than the female-specific OD2 domain.
Within insect orders, also a strong reduction of the sequence identity in male-specific OD2
domain alignment is seen. This observed rapid evolution of male-specific *dsx* regions is
supported by other research and is indicative of its fast evolution likely due to increased
diversity of secondary sexual traits in male insects (40). In females, the sex-specific region of
OD2 is relatively conserved across different insect orders. However, within Hymenoptera the
female-specific OD2 domain shows a lower similarity with only one out of eight conserved
amino acids being shared with all other tested species in the hymenopteran sequences, when
compared to other insect orders that have half of the conserved amino acids (Fig. S6B). In *D.*
melanogaster the C-terminus including the OD2 domain of DsxF binds the co-factor Intersex
(41). The differences in female-specific OD2 amino acid composition in Hymenoptera could
indicate that either no co-factors or different co-factors play a role in regulating the sex in this
order compared to the other insect orders. In addition, in Hymenoptera the female specific
OD2 domain was found to evolve much faster than the male-specific regions. This is contrary
to Diptera, Lepidoptera and Coleoptera and has been suggested to correlate with the
evolution of female-dominated eusociality and female-specific caste differentiation (40).
However, *Nasonia* is solitary as are many other hymenopteran species. In Hymenoptera,
females arise from fertilized eggs while male development initiates by default from haploid,
unfertilized eggs. We suggest that the rapid evolution of female-specific OD2 domain in
Hymenoptera could be linked to this difference. All in all, the identification of the two additional
male NvDsx isoforms opens the door to better comparative phylogenetic studies to better
understand the evolution of Dsx in insects.

Outlook

In this research we identified two sexually dimorphic traits and one male-specific trait that are
actively regulated during specific developmental windows by *Dsx* in *N. vitripennis.* Our results
suggest that male-specific lack of antennae and femur pigmentation is strictly regulated
during pupal development as silencing *NvDsx* from the larval stage onwards resulted in build-
up of pigmentation in the males, while silencing *NvDsx* from the pupal stage onwards did not

give any effect. In addition, our results suggest that sex-specific wing size and shape is

regulated by *NvDsx* during embryogenesis and early larval development as silencing *NvDsx*

from the 2nd larval instar onwards significantly increases wing size in males, while silencing

from the pupa onwards does not change wing size and shape. However, 2nd larval instar

NvDsx RNAi did not result in complete female-like wing size, suggesting that a big part of the

wing developmental window is during embryogenesis and possibly the 1st larval instar. We

additionally show that *NvDsx* regulates only male reproductive organ growth and not

morphology from 2nd larval instar onwards, suggesting that reproductive organ tissue

differentiation takes place before the 2nd instar which has not been reported before.

Visualizing *Dsx*-expressing cells in *D. melanogaster* has demonstrated before that *Dsx* gene

expression is under elaborate temporal and spatial control to regulate tissue-specific sexual

differentiation (19). Here, we expand this knowledge using a non-drosophilid species by

showing that different morphological characteristics require their own specific timing and

action of Dsx protein and we speculate that this timing and action is also species-specific. Our

findings solidify the view that in insects, sexual development is not only a set and forget

mechanism that only operates during embryogenesis, but one that requires continuous input

from Dsx on a species-specific basis.

Material and methods:

Insects rearing

The *Nasonia vitripennis* lab strain AsymCx were reared on *Calliphora sp.* hosts *at* 16h/8h

light/dark and 25℃. AsymCx strain is in the lab for over 10 years and cured from *Wolbachia*

infection.

5' and 3' RACE-PCR to identify additional *NvDsx* splice variants

For identification of the 3' ends of *NvDsx* mRNA*,* total RNA of 3 adult male- and female

AsymCx wasps were extracted individually using TriZol (Invitrogen, Carlsbad, CA, USA) and

cleared from genomic DNA by using the DNA-*free* DNA removal kit according to

manufacturer's instructions (Invitrogen, Carlsbad, CA, USA). Total RNA was annealed with

the 3' adapter from the FirstChoice RLM-RACE kit (Ambion, Austin, TX, USA) during the

reverse-transcription using RevertAidTM H Minus First Strand cDNA Synthesis Kit

(Fermentas, Hanover, MD, USA) and stored at -80°C. The cDNA was subjected to series of

(semi-) nested PCRs using primers complementary to the adapter and gene-specific primers

targeting NvDSXU, and NvDsx_X7, NvDsx_X8, NvDsx_X9 and NvDSXMR1 at 94°C for 3

minutes, 35 cycles of 94°C for 30 seconds, 55-62°C for 30 seconds and 72°C for 2 minutes, with a final extension of 10 minutes at 72°C. Primers were designed based on the initial *NvDsx* gene structure (27) and sequences are listed in the Table S1.

For identification on the 5' end of *NvDsx*, total RNA of 3 female AsymCx wasps were extracted using TriZol (Invitrogen, Carlsbad, CA, USA). After decapping, a 45-base RNA adapter has been ligated to the RNA population using the FirstChoice RLM-RACE kit (Ambion, Austin, TX, USA) and reverse transcribed using the RevertAidTM H Minus First Strand cDNA Synthesis Kit (Fermentas, Hanover, MD, USA). The cDNA was subjected to series of (semi-) nested PCRs using primers complementary to the adapter and gene-specific primers targeting NvDSXU at 94°C for 3 minutes, 35 cycles of 94°C for 30 seconds, 55-60°C for 30 seconds and 72°C for 2 minutes, with a final extension of 10 minutes at 72°C. Primers were designed based on the initial *NvDsx* gene structure (27) and sequences are listed in the Table S1.

5' and 3' RACE-PCR fragments were run and visualized on ethidium bromide-containing 1.5% agarose gel with 1×TAE buffer. All RACE-PCR products were purified using GeneJET Gel Purification Kit (Fermentas, Hanover, MD, USA) and ligated into pGEM-T vector (Promega, Madison, WI, USA) and transformed into competent JM-109 *Escherichia coli* cells (Promega, Madison, WI, USA). Colony-PCRs were conducted by using primers recommended by the kit at 94°C for 3 minutes, 40 cycles of 94°C for 30 seconds, 55°C for 30 seconds and 72°C for 2 minutes, with a final extension of 7 minutes at 72°C. All isoforms were sequenced on an ABI 3730XL capillary sequencer (Applied Biosystems) and reads were visualized and aligned in MEGA4 (42). Exon-intron structure of the genes was constructed by aligning the mRNA sequences to *N. vitripennis* genome and visualizing with Geneious Prime 2019.1.3 and CLC Workbench 6.

RT-PCR to confirm usage of alternative 1' exon

Total RNA was extracted from four female samples and three male samples containing three adults each using NucleoSpin XS (Brand) with final elution done in 15 µl nuclease-free water. Eleven µl of eluted total RNA was converted to cDNA using RevertAidTM H Minus First Strand cDNA Synthesis Kit (Fermentas, Hanover, MD, USA) with random hexamers according to manufacturer's protocol and stored at -80°C. One female sample was not converted to cDNA but used as gDNA contamination control in RT-PCR. For the RT-PCR reactions with the primersets NvEF-1a (Table S1) and Nv_DsxU_F6 or Nv_DsxU_F7 with Nv_Dsx_qPCR_R, GoTaq® G2 Flexi DNA Polymerase (Promega) was used to prepare the mastermix following manufacturer's instruction and amplified with a standard PCR profile: 3

374 min at 95°C, 35 amplification cycles of 30s at 95°C, 30s at 55°C, 30s at 72°C and a final

375 extension of 7 min at 72 °C in a thermal cycler (T100TM Thermal Cycler, Bio-Rad). Five µl of

376 PCR product was visualized on a non-denaturing 1% agarose gel with a 100bp ladder. PCR

377 fragments were Sanger sequenced at Eurofins Scientific to confirm correct amplification.

Alignment and phylogenetic analysis of Dsx OD2 domain across insect orders

Dsx isoforms from 22 species across insect taxa were downloaded from NCBI protein

database with accession number attached in the end of each sequence name (Fig. S6A and

S6B). OD2 common domain and sex-specific region of OD2 domain selection is based on An

et al. 1996 (2). Both the male and female Dsx OD2 domain sequences were separately

aligned using Guidance2 Online Server with Guidance2 method, settings are MAFFT

algorithm, maxiteration of 10, 100 alternative guide-trees. Geneious 2019.1 (created by

Biomatters. Available from https://www.geneious.com) was used to shade the amino acids

with Blosum62 score matrix applied to display the similarity between each sequence

(threshold at 3).

RNAi of *NvTraF* and *NvDsx* gene

For the *Nvtra* RNAi knockdowns, MEGAscript RNAi Kit (Thermo Fisher) was used to generate

dsRNA to target exon 5-8 of *Nvtra* mRNA in 4th instar larva stage of females which resulted in

a male splice form of *NvDsx* as shown in Verhulst *et al.* (28). To conduct functional analysis of

NvDsx, MEGAscript RNAi Kit (Thermo Fisher) was used to generate dsRNA based on *NvDsx*

common region (exon 2-5) present in both male and female splice-variants or the specific

region (exon 7-9) only presents in two longer male splice-variants. *Gfp* dsRNA was generated

from the vector pOPINEneo-3C-GFP and was a gift from Ray Owens (Addgene plasmid #

53534; http://n2t.net/addgene: 53534; RRID: Addgene_53534). This *gfp* dsRNA was used as

control in all experiments as it has no target in *Nasonia*. Primers designed using Geneious

10.0.9 to construct the dsRNA T7 template are listed in the Table S1. For male-specific

NvDsx RNAi, offspring of AsymCX strain was collected at 2nd and 4th larval instars and white

pupa stages which are described in Table S2. All dsRNAs were diluted with RNase-free

elution buffer (MEGAscript RNAi Kit (Thermo Fisher)) to 4000 ng/ul (NanoDrop™ 2000

Spectrophotometer, ThermoFisher) before proceeding to microinjection.

Microinjection of different developmental stages of wasps was carried out with IM300

Microinjecctor (Narishige) and FemtoJet® 4i (eppendorf) which followed the protocols

described by Werren *et al.* (43) and Lynch & Desplan (44) with minor changes. Red colour

food dye was pre-mixed with the dsRNAs in a ratio of 1:9. Experiments were performed using

AsymCx strain at different developmental stages for different purposes (Table S3).

2nd and 4th instar larvae were collected from parasitoid hosts and placed on a 1X PBS agar
plate before injection. Early pupal stage samples were fixed on the glass slides with double-
sided tape. After injection, 2nd instar larvae were transferred back to host (6-8 per host) and
sealed with the host shell. Afterward, sealed hosts were kept in 8 tube-PCR strips with open
lid. Subsequently, all injected samples either in 8 tube-PCR strip or on slides were placed on
1X PBS plates to prevent dehydration and incubated at standard rearing conditions until adult
emergence.

NvDsx splicing and expression throughout development

Male and female wasps were separated at black pupal stage by the sex-specific traits such as
forewing size and presence/absence of the ovipositor. To generate female offspring, one
selected male and one selected female pupa were collected in a single glass vial clogged with
cotton plug until eclosion and kept for one more day to make sure they had mated. After
fertilization, two hosts per day for the first two days were provided to mated females to initiate
oviposition. Afterwards, one-hour host presentations were repeated twice in a row per day.
Since mated *N. vitripennis* tend to lay extremely female-biased eggs, offspring from mated
females was considered female (45). Male offspring was collected by using virgin females
with the same setup. To facilitate early stage (eggs) sample collection, egg-laying chamber
(1.5ml Eppendorf tube with cut-off tip presenting the head of host pupa plugged in 1000 μL
filter pipet tip) was made to allow the female only access to the head portion of the fly host.

To confirm that male-specific splice variants were not under the regulation of NvTraF but
spliced in default mode independently of NvTraF, female samples were collected from 3h, 1d,
2d, 3d, 4d, 5d and 6d after Nv*traF* knockdown. To study the expression pattern of *NvDsx* sex-
specific splice variants throughout whole developmental stages, samples were collected at
specific time points listed in Table S2. More than 50 embryo eggs were collected at 36h and
five to ten larvae, pupae or adults were collected at 2.5d and onwards. Collected samples
were subsequently frozen in liquid nitrogen and stored in -80°C. *NvDsx* sex-specific primers
were designed using Geneious 10.0.9. Both RP49 and NvEF-1a were assessed for their
expression stability in this semi-quantitative RT-PCR setup, and *NvEF-1a* was finally chosen
as the reference gene. Primer sequences are listed in the Table S1.

Frozen samples were homogenized by pestles which were designed for 1.5 ml
microcentrifuge tubes (Biosphere SafeSeal Tube 1.5ml). Total RNA was extracted using ZR
Tissue & Insect RNA MicroPrep™(Zymo) following manufacturer's instructions. On column

DNase treatment step was added to all samples. Total RNA of each samples was eluted in 16 µl of DNase/RNase free water. After verifying the purity and concentration by NanoDrop™ 2000 Spectrophotometer (ThermoFisher), 1µg of each template was synthesized into cDNA with standard reaction mix (SensiFAST™ cDNA Synthesis Kit, Bioline) in a thermal cycler (Bio-Rad T100TM Thermal Cycler, Bio-Rad) with 5 min priming at 25°C, 30 min reverse transcription at 46°C and 5 min reverse transcriptase inactivation at 85°C.

For all RT-PCR reactions, GoTaq® G2 Flexi DNA Polymerase (Promega) was used to prepare the mastermix following manufacture's instruction and amplified with a standard PCR profile: 3min at 95°C, 34 (male-specific splicing verification) / 29 (expression pattern of *NvDsx*) amplification cycles of 30s at 95°C, 30s at 55°C, 50s at 72°C and a final extension of 5 min at 72 °C in a thermal cycler (T100TM Thermal Cycler, Bio-Rad). In order to standardize the expression pattern of *NvDsx*, the number of PCR cycles needed was determined by comparison of the brightness of the bands in gels of different PCR reactions with cycles ranging from 25 to 34 cycles to achieve non-oversaturated brightness. In the end 29 cycles were used for both reference genes and target genes.

Silencing efficiency of *NvDsx*

qPCR was conducted to verify the silencing efficiency of *Dsx* knockdown. After *dsRNA* injection, samples were collected at either adult stages (injected at 4[th] instar larva and white pupa stages) or white pupa stage (injected at 2[nd] instar larva stage). Four to five pupae or adults were pooled to produce one biological replicate and six replicates per treatment were prepared. All testing samples were used in qPCR according to SensiFAST™ SYBR® No-ROX Kit manual (Bioline). *NvDsx* qPCR primers were designed on the common region (exon 3-5) of male and female splice variants (Table S1). *NvEF-1a* was used as reference genes. qPCR was carried out using the CFX96TM Real-Time System (Bio-Rad) with Bio-Rad CFX Manager 3.1 Software (Bio-Rad). The standard qPCR profiles consist of 95°C for 3min, 45 amplification cycles of 15s at 95,15s of 55°C*,* 30s of 72°C with a final standard dissociation curve step to check for non-specific amplification.

Measurements of male forewing size, external reproduction organs

Forewing size was measured by wing length and width based on the standard methods described by Loehlin *et al.* (46). Tibia length of the hind leg was used to standardize the body size differences among individuals (47). Forewings and external reproduction organ of adult males were dissected on clean slides with a drop of 1%PBS buffer using dissection tweezers and subsequently mounted in a drop of ethanol on slides which was allowed to evaporate.

External reproduction apparatuses were identified based on the description in Liu et al (48)

and E. Snodgrass (38). Photos were taken and measured by Dino-Lite Edge 5MP digital

microscope using DinoCapture 2.0 software.

Data analysis

qPCR data was imported to LinRegPCR software (LinRegPCR, 2017.1.0.0, HFRC,

Amsterdam, The Netherlands) (49). After baseline correction, the initial number of templates

(N0) were calculated based on the average PCR efficiency of each amplicon. Relative

expression levels of *NvDsx* in each sample was obtained by dividing the N0 value of *NvDsx*

by N0 value of *EF-1a*. Statistical analysis of qPCR and measurement data was performed in

R using One-way ANOVA and Tukey's Honest Significant Difference (HSD) for post hoc test.

Acknowledgment

We thank Hans Smid for assisting with the phenotype recording; Rutger Diepeveen for his

contribution to the *NvDsx* RNAi work; Joan Diaz Calafat, Age Muller, Simon Pleiter and Romy

Gielings for their contribution to the *NvtraF* RNAi work; Min Xu for supporting the sample

preparation, collection and measuring; and Weizhao Sun for recording the mating behaviour

of RNAi treated males. We additionally thank all the tested *N. vitripennis* for their great

sacrifice to science. This work is part of the Dutch Research Council (NWO) research

programme Innovational Research Incentives Scheme Veni with project number 863.13.014

granted to ECV.

Reference

1. O. Barmina and A. Kopp, Sex-specific expression of a HOX gene associated with rapid

morphological evolution. *Dev. Biol.* **311**, 277–286 (2007).

2. W. An, S. Cho, H. Ishii and P. C. Wensink, Sex-specific and non-sex-specific

oligomerization domains in both of the *doublesex* transcription factors from *Drosophila*

melanogaster. *Mol. Cell. Biol.* **16**, 3106–11 (1996).

3. J. N. Shukla and S. R. Palli, Doublesex target genes in the red flour beetle, *Tribolium*

castaneum. *Sci. Rep.* **2**, 948 (2012).

4. F. Ohbayashi *et al.*, A homologue of the *Drosophila doublesex* gene is transcribed into sex-

specific mRNA isoforms in the silkworm, *Bombyx mori*, *Comp. Biochem. Physiol. - B Biochem.*

Mol. Biol. **128**, 145-158 (2001).

5. K. C. Burtis and B. S. Baker, *Drosophila doublesex* gene controls somatic sexual

differentiation by producing alternatively spliced mRNAs encoding related sex-specific

polypeptides. *Cell.* **56**, 997–1010 (1989).

6. D. C. Price, A. Egizi and D. M. Fonseca, The ubiquity and ancestry of insect *doublesex*. *Sci.*

Rep. **5**, 13068 (2015).

7. E. C. Verhulst and L. van de Zande, Double nexus-Doublesex is the connecting element in

sex determination. *Brief. Funct. Genomics* **14**, 396–406 (2015).

8. T. M. Williams *et al.*, The Regulation and Evolution of a Genetic Switch Controlling Sexually

Dimorphic Traits in *Drosophila*. *Cell* **134**, 610–623 (2008).

9. A. Kopp, *Drosophila* sex combs as a model of evolutionary innovations. *Evol. Dev.* **13**,

504–22 (2011).

10. X. Chen *et al.*, Disruption of sex-specific *doublesex* exons results in male- and female-

specific defects in the black cutworm, *Agrotis ipsilon*. *Pest Manag. Sci.* **75**, 1697-1706 (2019).

11. M. G. Suzuki *et al.*, Role of the male BmDSX protein in the sexual differentiation of

Bombyx mori. *Evol. Dev.* **7**, 58–68 (2005).

12. J. Xu *et al.*, Sexually dimorphic traits in the silkworm, *Bombyx mori*, are regulated by

doublesex. *Insect Biochem. Mol. Biol.* **80**, 42–51 (2017).

13. K. Kunte *et al.*, *doublesex* is a mimicry supergene. *Nature* **507**, 229–232 (2014).

14. H. Gotoh *et al*., Developmental link between sex and nutrition; *doublesex* regulates sex-
specific mandible growth via juvenile hormone signalling in stag beetles. *PLoS Genet.* **10**,
e1004098 (2014).

15. Y. Ito *et al*., The role of *doublesex* in the evolution of exaggerated horns in the Japanese
rhinoceros beetle. *EMBO Rep.* **14**, 561–567 (2013).

16. C. C. Ledón-Rettig, E. E. Zattara and A. P. Moczek, Asymmetric interactions between
doublesex and tissue- and sex-specific target genes mediate sexual dimorphism in beetles.
Nat. Commun. **8**, 14593 (2017).

17. C. C. Robinett, A. G. Vaughan, J.-M. Knapp and B. S. Baker, Sex and the single cell. II.
There is a time and place for sex. *PLoS Biol.* **8**, e1000365 (2010).

18. J. L. Morrow, M. Riegler, M. Frommer and D. C. A. Shearman, Expression patterns of
sex-determination genes in single male and female embryos of two *Bactrocera* fruit fly
species during early development. *Insect Mol. Biol.* **23**, 754–767 (2014).

19. A. Klein *et al*., Evolution of social insect polyphenism facilitated by the sex differentiation
cascade, *PLOS Genet.* **12**, e1005952 (2016).

20. M. Nipitwattanaphon *et al*., Effects of ploidy and sex-locus genotype on gene expression
patterns in the fire ant *Solenopsis invicta*. *Proc. R. Soc. B Biol. Sci.* **281**, 20141776–20141776
(2014).

21. S. Cho, Z. Y. Huang and J. Zhang, Sex-specific splicing of the honeybee *doublesex* gene
reveals 300 million years of evolution at the bottom of the insect sex-determination pathway.
Genetics **177**, 1733–41 (2007).

22. A. Ugajin *et al*., Expression profile of the sex determination gene *doublesex* in a
gynandromorph of bumblebee, *Bombus ignites. Sci. Nat.* **103**, 17 (2016).

23. M. O. Miyakawa, K. Tsuchida and H. Miyakawa, The *doublesex* gene integrates multi-
locus complementary sex determination signals in the Japanese ant, *Vollenhovia emeryi*.
Insect Biochem. Mol. Biol. **94**, 42–49 (2018).

24. S. Mine *et al*., Identification and functional characterization of the sex-determining gene
doublesex in the sawfly, *Athalia rosae* (Hymenoptera: Tenthredinidae). *Appl. Entomol. Zool.*
52, 497–509 (2017).

25. R. Raychoudhury *et al*., Behavioral and genetic characteristics of a new species of

Nasonia, *Heredity.* **104**, 278–288 (2010).

26. D. C. Darling and J. H. Werren, Biosystematics of *Nasonia* (Hymenoptera: Pteromalidae):

Two new species reared from birds' nests in north America. *Ann. Entomol. Soc. Am.* **83**, 352–

370 (1990).

27. D. C. S. G. Oliveira *et al*., Identification and characterization of the *doublesex* gene of

Nasonia. *Insect Mol. Biol.* **18**, 315–324 (2009).

28. E. C. Verhulst, L. W. Beukeboom and L. van de Zande, Maternal control of haplodiploid

sex determination in the wasp *Nasonia*. *Science.* **328**, 620–623 (2010).

29. J.-C. Zhuo *et al*., Identification and functional analysis of the *doublesex* gene in the sexual

development of a hemimetabolous insect, the brown planthopper. *Insect Biochem. Mol. Biol.*

102, 31–42 (2018).

30. B. S. Baker and M. F. Wolfner, A molecular analysis of *doublesex*, a bifunctional gene that

controls both male and female sexual differentiation in *Drosophila melanogaster*. *Genes Dev.*

2, 477–489 (1988).

31. J. N. Shukla and J. Nagaraju, Two female-specific DSX proteins are encoded by the sex-

specific transcripts of *dsx*, and are required for female sexual differentiation in two wild

silkmoth species, *Antheraea assama* and *Antheraea mylitta* (Lepidoptera, Saturniidae). *Insect*

Biochem. Mol. Biol. **40**, 672–682 (2010).

32. F. Nunes *et al*., Non-target effects of green fluorescent protein (GFP)-derived double-

stranded RNA (dsRNA-GFP) used in honey bee RNA interference (RNAi) assays, *Insects* **4**,

90–103 (2013).

33. E. H. Slifer, Sense organs on the antenna of a parasitic wasp, *Nasonia vitripennis*

(Hymenoptera, Pteromalidae). *Biol. Bull.* **136**, 253–263 (1969).

34. J. H. Massey and P. J. Wittkopp, The genetic basis of pigmentation differences within and

between *Drosophila* species. *Curr. Top. Dev. Biol.* **119**, 27-61.

35. D. W. Loehlin *et al*., Non-coding changes cause sex-specific wing size differences

between closely related species of *Nasonia*, *PLoS Genet.* **6**, e1000821 (2010).

36. D. W. Loehlin and J. H. Werren, Evolution of shape by multiple regulatory changes to a

growth gene. *Science* **335**, 943–7 (2012).

37. K. Mysore et al., siRNA-mediated silencing of *doublesex* during female development of
the dengue vector mosquito *Aedes aegypti*. *PLoS Negl. Trop. Dis.* **9**, e0004213 (2015).

38. R. E. Snodgrass, A revised interpretation of the external reproductive organs of male
insects. *Smithson. Misc. Collect.* **136**, 1–60 (1957).

39. A. E. Christiansen, E. L. Keisman, S. M. Ahmad and B. S. Baker, Sex comes in from the
cold: the integration of sex and pattern. *Trends Genet.* **18**, 510–516 (2002).

40. S. Baral, G. Arumugam, R. Deshmukh and K. Kunte, Genetic architecture and sex-
specific selection govern modular, male-biased evolution of *doublesex*. *Sci. Adv.* **5**, eaau3753
(2019).

41. Y. Yang, W. Zhang, J. R. Bayrer and M. A. Weiss, *Doublesex* and the regulation of sexual
dimorphism in *Drosophila melanogaster*. *J. Biol. Chem.* **283**, 7280–7292 (2008).

42. K. Tamura, J. Dudley, M. Nei and S. Kumar, MEGA4: Molecular evolutionary genetics
analysis (MEGA) software version 4.0. *Mol. Biol. Evol.* **24**, 1596–1599 (2007).

43. J. H. Werren, D. W. Loehlin and J. D. Giebel, Larval RNAi in *Nasonia* (parasitoid wasp).
Cold Spring Harb. Protoc. DOI:10.1101/pdb.prot5311 (2009).

44. J. A. Lynch and C. Desplan, A method for parental RNA interference in the wasp *Nasonia*
vitripennis. *Nat. Protoc.* **1**, 486–494 (2006).

45. W. D. Hamilton, Extraordinary sex ratios. *Science.* **156**, 477–488 (1967).

46. D. W. Loehlin, L. S. Enders and J. H. Werren, Evolution of sex-specific wing shape at the
widerwing locus in four species of *Nasonia*. *Heredity.* **104**, 260–269 (2010).

47. H. C. J. Godfray, Parasitoids: behavioral and evolutionary ecology. *Parasitoids Behav.*
Evol. Ecol. (1994).

48. N.-Y. Liu *et al*., Morphology and ultrastructure of the male reproductive system of the
jewel wasp, *Nasonia vitripennis* (Walker) (Hymenoptera: Pteromalidae). *J. Asia. Pac. Entomol.*
20, 577–582 (2017).

49. C. Ramakers, J. M. Ruijter, R. H. L. Deprez and A. F. Moorman, Assumption-free analysis
of quantitative real-time polymerase chain reaction (PCR) data. *Neurosci. Lett.* **339**, 62–66
(2003).

 Figures

Figure 1. Comparison of insect *Doublesex* (*Dsx*) gene models. The eight *N. vitripennis* male and female *Dsx* splice variants are compared to *Dsx* splice variants of *D. melanogaster* and *A. mellifera*. Figure was modified from Oliveira *et al.* (27) to include the alternative 1st exon (1') and the additional *NvDsx* male splice variants. Male and female specific splice forms are indicated by ♂ and ♀. Exon numbers are noted on each exon. Light grey, black and dark grey blocks indicate coding region, DM domain and OD2 dsx dimer region respectively. Scales are provided behind each species splice structures.

Figure 2. Confirmation of sex-specific *Dsx* splicing after *Nvtra* knockdown in *N.vitripennis.* (A) Sex-specific splice variants of *Nvtra* and *NvDsx* in *N. vitripennis* females that are injected in the 4th instar larval stage with *Nvtra* dsRNA or *gfp* dsRNA. (B) male and female non-treated (NT) control samples for each primer sets. Arrows indicate male (in blue) and female (in pink) splice forms of the different genes. Lanes 1-12: Female *Nvtra* knockdown samples from 3-hour post injection (hpi) to 5-day post injection (dpi). Lane 13: 100bp ladder (Thermo). Lanes 14-25: Female *gfp* knockdown samples from 3dpi to 5dpi. For each stage two samples are used. *EF-1a* is used as expression and loading control.

Figure 3. Pigmentations of antennae and hind legs of adult wasps after RNAi treatment in different developmental stages. (A) NT male (B) 4th instar *gfp* RNAi (*gfp-i*) male (C) 2nd instar *gfp* RNAi (*gfp-i*) male (D) NT female (E) 4th instar *NvDsx* RNAi (*NvDsx-i*) male and (F) 2nd instar *NvDsx* RNAi (*NvDsx-i*) male. Light source and exposure time are equal in all pictures.

Figure 4. Comparision of forewing size of adult males after *NvDsx* (*NvDsx-i*) or mock *gfp* (*gfp-i*) dsRNA treatment in the 2nd or 4th instar larval stage and NT control. Y-axis represent (A) forwing length/width ratio corrected with tibia length; (B) length corrected with tibia length; (C) width corrected with tibia length. X-axis shows 6 different groups: males treated with *NvDsx* (*NvDsx-i*) and *gfp* (*gfp-i*) dsRNA in the 2nd and 4th instar larval stage, NT males and NT females. Different colors were assigned for different treatments. Boxplots were generated in R and displayed with minimum, first quartile (Q1), median, third quartile (Q3), and maximum. Dots outside the boxplots represent outliers from the treatments. Replicates were provided in the figures of each treatment. One-way ANOVA and post-hoc Tukey's HSD were used for data analysis. Letters in the figures denote statistical significance (P<0.05) among treatments.

Figure 5. Comparison of genitalia length of adult males after *NvDsx* (*NvDsx-i*) or mock *gfp*
(*gfp-i*) dsRNA treatment in the 2nd or 4th instar larval stage and NT control. A) Y-axis
represents genitalia length corrected with tibia length. X axis shows 5 experiment groups:
males treated with *NvDsx* (*NvDsx-i*) and *gfp* (*gfp-i*) dsRNA in the 2nd and 4th instar larval
stage and NT control. Different colours were assigned for different treatments. Boxplots were
generated in R and displayed with minimum, first quartile (Q1), median, third quartile (Q3),
and maximum. Dots outside the boxplot represent outliers from each treatment. Replicates
were provided in the figures of each treatment. One-way ANOVA and post hoc Tukey's HSD
were used for data analysis. Letters in the figures denote statistical significance (P<0.05)
among treatments. B) Comparison of paramere length (without aedeagus) of adult males after
NvDsx or mock *gfp* dsRNA treatment in the 2nd or 4th instar larval stage and NT control. (a)
paramere length (without aedeagus) in NT males, (b and d) after treatment with *NvDsx*
(*NvDsx-i*) dsRNA in 4th and 2nd instar male larvae, (c and e) after treament with *gfp* (*gfp-i*)
dsRNA in 4th and 2nd instar larvae. Different treatments are indicated below the figure. Scales
are provided on top of each photos. Each part of paramere is described in *E. Snodgrass*
(1957) (38) and indicated with the following abbreviations: outer parameres (opar), cuspis
(cus) and digitus (dig).